

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Hydroperoxyl radical (HOO•) as a reducing agent: unexpected synergy with antioxidants. A review

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Baschieri, A., Jin, Z., Amorati, R. (2023). Hydroperoxyl radical (HOO•) as a reducing agent: unexpected synergy with antioxidants. A review. FREE RADICAL RESEARCH, 57(2), 115-129 [10.1080/10715762.2023.2212121].

Availability:

This version is available at: <https://hdl.handle.net/11585/943253> since: 2023-09-28

Published:

DOI: <http://doi.org/10.1080/10715762.2023.2212121>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Andrea Baschieri, Zongxin Jin and Riccardo Amorati*

Hydroperoxyl Radical (HOO•) as a reducing agent: unexpected synergy with antioxidants. A review.

Free Radical Research, 57:2, 115-129

The final published version is available online at: **DOI:**
[10.1080/10715762.2023.2212121](https://doi.org/10.1080/10715762.2023.2212121)

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Hydroperoxyl Radical (HOO[•]) as a reducing agent: unexpected synergy with antioxidants. A review.

Andrea Baschieri, Zongxin Jin and Riccardo Amorati*

^a Institute for Organic Synthesis and Photoreactivity (ISOF), National Research Council of Italy (CNR), Via P. Gobetti 101, I-40129 Bologna, Italy.

^b Department of Chemistry "G. Ciamician", University of Bologna, Via S. Giacomo 11, 40126 Bologna, Italy. Tel: +39 051 2095689. Fax: +39 051 2095688. E-mail: riccardo.amorati@unibo.it

Keywords

Superoxide, lipid peroxidation, nitroxide, polydopamine, ferroptosis, γ -terpinene


Abstract

This review highlights the progress made in recent years in harnessing the peculiar chemistry of the hydroperoxyl, or perhydroxyl, radical (HOO^\bullet) during lipid peroxidation, particularly with regard to its interaction with antioxidants. The HOO^\bullet radical, the protonated form of superoxide, plays an important role in the propagation and termination of lipid peroxidation in nonaqueous systems. However, differently from alkylperoxyl (ROO^\bullet) radicals that have only oxidizing ability, HOO^\bullet has a two-faced oxidizing and reducing activity. The HOO^\bullet radical can reduce the radical of the antioxidant (phenols and aromatic amines) by H-atom transfer ($\text{A}^\bullet + \text{HOO}^\bullet \longrightarrow \text{AH} + \text{O}_2$) thus increasing the length of the inhibition period and the effectiveness of the antioxidant. The simultaneous presence of HOO^\bullet and ROO^\bullet radicals triggers the catalytic antioxidant activity of quinones and nitroxides, and explains the antioxidant activity of melanin-like polymers. The HOO^\bullet radical can be formed by fragmentation of ROO^\bullet radicals deriving from amines, alcohols, substituted alkenes and may be present at low concentrations in many oxidizing systems. Pro-aromatic compounds, like the natural essential oil component γ -terpinene, are the most effective sources of HOO^\bullet , and behave as co-antioxidants in the presence of nitroxides or quinones. The future developments and applications of the HOO^\bullet chemistry in the context of the inhibition of autoxidation are also discussed.

1. Introduction

Autoxidation is a reaction of organic compounds under mild conditions with atmospheric O_2 , leading to their deterioration by producing hydroperoxides and other oxygenated compounds. Autoxidation affects not only man-made materials like plastics, cosmetics, pharmaceuticals and food, but also biomolecules in living organism, especially those rich in polyunsaturated lipids; thus, it is of extreme importance in food, polymers and biomedical applications. Strikingly, the base mechanism of the reaction with O_2 of all these materials shares the same mechanistic features, consisting of a radical-chain reaction with the classical initiation, propagation, and termination steps [1-3]. The most effective and convenient way to protect organic substrates from oxygen reactivity, both in nature, as in man-made materials, is to use antioxidants. Depending on the mechanism of inhibition of the autoxidation reaction, antioxidants can be classified between preventive and chain-breaking (or radical-trapping antioxidants, RTA) if they reduce the initiation or the propagation step, respectively [1].

The peroxidation mechanism is well established, and is based on a radical chain mechanism propagated by carbon-centred (alkyl, R^\bullet) and oxygen-centred (alkylperoxyl, ROO^\bullet) radicals, as shown in Scheme 1 [1].



Scheme 1. Co-oxidation of an organic substrate (RH) with a compound known to oxidize via hydroperoxyl radical (EH), initiated by In^\bullet radicals, in the presence of antioxidants (AH), showing the role of hydroperoxyl (HOO^\bullet) radicals in the regeneration mechanism.

In such system, the autoxidation rate depends on the reactivity of the substrate (RH) with ROO^\bullet radicals (propagation step, in Scheme 1 exemplified by a H-atom abstraction), and on the steady state concentration of ROO^\bullet radicals. This, in turn, depends on the amount of ROO^\bullet generated by the initiation process, and on the decay rate of ROO^\bullet radical by self-termination or by reaction with chain-breaking antioxidants (AH). Effective antioxidants must be able to compete with RH for ROO^\bullet radicals, thus must have a high inhibition constant (k_{inh}). Moreover, the radical of the antioxidant (A^\bullet) must not propagate the oxidative chain, and possibly should trap further radicals [4]. This last aspect is very important in real conditions because antioxidants are expected to protect oxidizable materials for a long time. The length of the protection by chain-breaking antioxidants can

be increased by agents able to regenerate the active antioxidant by reducing the A^\bullet radical back to AH. For example, it is well known that the antioxidant activity of α -tocopherol in liposomes is prolonged by the presence of water soluble co-antioxidants (like vitamin C) in the water phase [5].

Besides the well-established chemistry of R^\bullet and ROO^\bullet radicals, the hydroperoxyl radical (HOO^\bullet) has been recently discovered as an important player in biological systems, in atmospheric pollutant degradation [6,7], in photocatalysis [8], and in organic synthesis [9]. Although the formation of HOO^\bullet during the autoxidation of certain organic substrates was recognized long time ago [10,11], the role of HOO^\bullet radical has been appreciated in its full importance only after the pioneering work by Foti and Ingold on the antioxidant effect of γ -terpinene [12].

HOO^\bullet is known as a source of oxidative damage in biological systems and is essentially similar to alkylperoxyl radicals in H-atom abstraction. On the other hand, the knowledge about the effects of the reducing properties of HOO^\bullet and of its conjugate base $O_2^{\bullet-}$, is still very limited [13-15]. The reducing ability of superoxide has been previously recognized as an important pathway for the repairing of oxidative damage of tyrosyl residues in proteins subject to oxidative stress. For example, by using mass spectrometry it was found that superoxide prevented myoglobin dimer formation by reducing the protein tyrosyl radical to the parent phenol. However, this reaction is also accompanied by the addition of superoxide at Tyr151, fully demonstrating the two-faced reducing and oxidizing character of the $HOO^\bullet/O_2^{\bullet-}$ couple [16].

In the context of the autoxidation reaction, HOO^\bullet is formed typically by the fragmentation

of ROO^\bullet radicals of specific oxidizable substrates (EH), which therefore represent a possible source of this radical. As summarized in Scheme 1, and explained in this review, the HOO^\bullet radical can potentially impact on both propagation and termination steps of the autoxidation reaction, as effect of its peculiar chemical characteristics. Furthermore, the HOO^\bullet reactivity with several phenolic and nonphenolic antioxidants (AH) has been somewhat neglected and, only recently, its enhancing effect on their antioxidant activity has been observed.


2. Chemical characteristics of hydroperoxyl radical

HOO^\bullet (hydroperoxyl radical), also known as perhydroxyl radical, is the protonated form of superoxide $\text{O}_2^{\bullet-}$. In an aqueous environment at physiological pH, HOO^\bullet is mostly in the deprotonated form ($\text{p}K_a = 4.8$) [17]. The role of $\text{HOO}^\bullet / \text{O}_2^{\bullet-}$ in autoxidation reactions is dependent on its ability to abstract or donate a H-atom, and this property in turn is a function of the strength of the bonds being formed (or broken). In order to have a quantitative description of the oxidizing and reducing properties and to compare them with other radicals in solution, we use the Bond Dissociation Free Energy (BDFE) and redox potentials (E°) values critically reviewed by Mayer and co-workers, as they represent the most complete source of chemical data for proton-coupled electron transfer (PCET) reagents [17].

2.1 H-atom abstraction by HOO^\bullet

The BDFE of the HOO-H bond is 86.5 kcal/mol, that is superimposable to that of the ROO-H bond in alkylperoxyl radicals, that ranges from 90.2 to 86.6 kcal/mol for MeOO-H and

^tBuOO-H, respectively (see Scheme 2) [17]. Regarding H-atom abstraction, HOO• behaves similarly to ROO• radicals. The hydroperoxyl radical propagates the oxidative chain in the autoxidation of 1,4-cyclohexadienes and other easily oxidizable substrates [10]. Moreover, HOO• abstracts the phenolic H-atom from α -tocopherol with a $k_H = 1.6 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$ in PhCl [18], that is very similar to that reported for ROO• under identical conditions ($k_H = 2.7 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$) [18].



Scheme 2: Aqueous proton-coupled electron transfer thermochemistry of (a) hydrogen peroxide, (b) H₂O and (c) CH₃OOH, with pK_a values above horizontal arrows, redox potentials (in V vs NHE) beside vertical arrows, and BDFE values on diagonal lines. Data from ref. [17].

However, differently from ROO•, the H-atom abstracting ability of HOO• in organic solvents is influenced by the formation H-bonds with the solvent. We have shown that the formation of S---HOO• interactions between HOO• and the solvent (**S**) reduces the rate constant of H-atom abstraction from C-H and ArO-H bonds because the H-bond becomes weaker going

from the reactants ($S\text{---}HOO^\bullet + ArOH$) to the products ($S\text{---}HOOH + ArO^\bullet$) [18]. Superoxide, $O_2^{\bullet-}$, is much less active as H-atom abstracting radical than HOO^\bullet [19], due to the relatively small BDFE of $^{\bullet}OO\text{---}H$ that is reported as 77.1 kcal/mol in water [17].

2.2. H-atom donation by HOO^\bullet

The unusual chemical property of the hydroperoxyl radicals is rooted in its hydrogen-donating ability. The BDFE ($^{\bullet}OO\text{---}H$) has been estimated as 51.2 kcal/mol in water and 45.1 kcal/mol in the gas phase [17]. Likewise, $O_2^{\bullet-}$ is a strong one-electron reducing agent, since the $E^\circ(O_2)$ is -0.35 V (vs NHE). The $H\text{---}OO^\bullet$ BDFE is much smaller than that of phenolic and non-phenolic antioxidants (typically from 74 to 84 kcal/mol). This demonstrates from a thermodynamic point of view the possibility of regenerating antioxidants from their radicals [17].


2.3. Comparison with H_2O/HO^\bullet

Despite the chemical similitude (only H and O atoms!) radicals derived from H_2O have a completely different reactivity. The BDFE of $HO\text{---}H$ is extremely high (115.8 kcal/mol), due to the absence of any resonance or hyperconjugative stabilization in HO^\bullet . The hydroxyl radical is also a strong one-electron oxidizing agent, $E^\circ(OH^\bullet)$ is 1.90 V (Scheme 2b). Moreover, the H-O BDFE of the $H\text{---}O^\bullet$ radical is also very high (102 kcal/mol) due to the instability of the O (atomic oxygen) specie, therefore HO^\bullet has no reducing ability [17].

2.4 Thermodynamic basis of antioxidant regeneration

In Scheme 3, a BDFE and potential ladder showing the comparison between the BDFE($^{\bullet}OO\text{---}H$) and that of relevant species for the chemistry of peroxidation and antioxidant

activity are reported. In this scheme, BDFE of all compounds that can form a bond stronger than 51.2 kcal/mol with a H-atom can potentially be reduced by HOO^\bullet , although the reaction will be more efficient if the strength of the newly formed bond is much higher than this value. An analogous ladder can be built by considering redox potentials, and species with a reduction potential bigger than -0.35 V vs HNE are amenable to be reduced by $\text{O}_2^{\bullet-}$. For example, HOO^\bullet reduces alkylperoxyl radicals ($\text{CH}_3\text{OO}^\bullet$ in Scheme 3) at a nearly diffusion rate, as reported by Ingold and Foti [12], while we estimated that the H-atom donation from HOO^\bullet to the α -tocopheroxyl radical (in PhCl at $30\text{ }^\circ\text{C}$) is as high as $10^9\text{ M}^{-1}\text{s}^{-1}$ [18]. As most autoxidation reactions occur in nonaqueous media, where deprotonation of acids is relatively unimportant, the actual interaction between HOO^\bullet and antioxidants is better described by the BDFE ladder. It should also be kept in mind that BDFE values are expected to change considerably as effect of H-bond with the solvent, so the relative order of the species reported in the ladder may change when considering the reactivity in solvents less able to form H-bonds. Besides PCET reactivity, $\text{HOO}^\bullet / \text{O}_2^{\bullet-}$ radicals can also easily add to phenoxyl radicals forming hydroperoxides which are no longer able to trap radicals or to be reduced to the parent phenols (see section 4.1) [16,18]. This reaction is in competition with ET or PCET from $\text{HOO}^\bullet / \text{O}_2^{\bullet-}$ to the radical of the antioxidant and reduces the regeneration. In summary, the data reported show that the HOO^\bullet radical has a two-faced oxidizing and reducing ability, that render it very different from the alkylperoxyl radical (ROO^\bullet) that is the main chain-carrying specie in autoxidation. While HOO^\bullet can potentially propagate the oxidative chain, indeed it is able to react, in some cases with high-rate constants, with a series of redox active substrates that would be otherwise inert under oxidative-only




Scheme 3: BDFE (upper) and reduction potential (vs NHE) (lower) ladders showing the thermodynamics of the donation of a H-atom from HOO• or the electron donation from O₂^{•-} to various acceptors. Data from [17].


3. Formation of hydroperoxyl radicals during autoxidation

Hydroperoxyl radicals can be present along with alkylperoxyl radicals during autoxidation reaction by fragmentation of ROO•. The oxidizable substrate must have specific structural characteristics to allow an efficient generation of HOO• radicals. The presence of a H-atom in 4 position respect to the terminal oxygen of the peroxy group is essential, in order to allow an intramolecular 1,4-hydrogen atom transfer (intra-1,4-HAT), that causes the

production of HOO[•]. However, 1,4-HAT will be more or less effective depending on structural features of the molecule. Some examples of HOO[•] production by 1,4-HAT, subdivided by type of functional group, are summarized below (Scheme 4).




ALCOHOLS




AMINES

Primary and secondary amines



Tertiary amines



Scheme 4: Box: general mechanism of HOO[•] elimination by 1,4-HAT from alkyl peroxy radicals, and typical reaction pathway in alcohols and amines.


3.1 Alcohols: The α-C-H group in primary and secondary alcohols (**1**), is preferentially attacked by radicals, owing to the adjacent hydroxyl group which stabilizes the alkyl radical [20]. The α-hydroxy alkyl radical, also named ketyl radical (**2**) reacts with O₂ at diffusion rate

to afford the α -hydroxy-peroxyl radical **3** which decomposes to a carbonyl compound (**4**) and HOO^\bullet [21].

3.2 Amines Like the oxygen atom in alcohols, nitrogen in amines stabilizes the α alkyl radical by resonance with the unpaired electron. Thus, the α -position of alkyl-substituted amines (**5**, **9**) is strongly activated toward H-atom abstraction, forming amino alkyl radicals (**6**, **10**) that, in the presence of O_2 , form the corresponding α -amino peroxyl radicals (**7**, **11**). In the case of primary and secondary amines, the aminoperoxyl radical decomposes to HOO^\bullet and an imine **8** [22]. In the case of tertiary alkylamines having at least one $-\text{CH}_2\text{CH}_2-$ substituent, such in the case of compound **9**, HOO^\bullet is generated with great efficiency clearly indicating that the aminoalkene **12** is the product of the reaction (see Scheme 4) [23].

3.3. Alkenes and pro-aromatic compounds


Recently has it been reported that the HOO^\bullet radical is formed during the autoxidation of styrene or cyclooctene at 70 °C and for hexadecene at 100 °C [24]. According to Scheme 5, based on theoretical calculations, HOO^\bullet production could be explained on the basis of a chain reaction mechanism involving alkoxy radicals. In the case of styrene, that undergoes autoxidation only by radical addition, the alkoxy radicals (generated together with epoxides by an addition-fragmentation pathway) combine with the double bond providing the alkoxy-alkyl radical **14**. The alkyl radical then reacts with oxygen to yield a peroxyl radical, which in turn produces HOO^\bullet and the alkoxy-substituted alkene **16** via intra-1,4-HAT. However, at low temperature and at high concentration of oxidizable substrate, the autoxidation of styrene affords a very limited amount of HOO^\bullet [25].



Scheme 5: Mechanism proposed for the formation of HOO• during the autoxidation of alkenes.

Some reports, based on the antioxidant effect of superoxide dismutase (SOD) on autoxidation rates, suggest that during the autoxidation of methyl linoleate in micelles a small fraction of the propagating peroxy radicals is represented by HOO•/O₂^{•-} radicals [26,27]. Considering the biological relevance of peroxidation of polyunsaturated lipids, these preliminary results call for further investigations about the mechanism of HOO• production.

1,4-Cyclohexadiene derivatives such as γ -terpinene **17**, a common terpene found in the essential oils (Scheme 6), represent a unique class of alkenes as they produce high yields of HOO• during their autoxidation [12]. γ -Terpinene reacts with radicals exclusively by H-atom transfer from the bis-allylic CH₂ groups forming a highly stabilized alkyl radical which reacts with O₂ to yield a peroxy radical with conjugated double bonds. Then, HOO• is released by intra-1,4-HAT mechanism, whose driving force is the formation of an aromatic product, *p*-cymene [28]. Other pro-aromatic compounds such as 1,4-cyclohexadiene **18** itself, methyl-1,4-cyclohexadiene **19**, 1,4-dihydronaphthalene **20** quantitatively generate HOO• during their autoxidation [9,10].




Scheme 6: Mechanism proposed for the formation of HOO^\bullet during the autoxidation of alkenes.

Contrary to γ -terpinene, the 1,3-cyclohexadiene isomer α -terpinene **21**, which has conjugated double bonds, only partially propagates the chain by HOO^\bullet [29]. The main products of α -terpinene's autoxidation are epoxides and diketones, with the latter likely resulting from the breakdown of alkyl hydroperoxides. Only about 20% of the products of the autoxidation are *p*-cymene, which is produced by an intra-1,4-HAT from the peroxy radical of α -terpinene, suggesting that HOO^\bullet is only marginally involved in chain propagation [30]. Recent kinetic studies support the view that the different behavior between γ - and α -terpinene is due to the preference of α -terpinene for ROO^\bullet addition rather than HAT [31]. Therefore, while potentially all pro-aromatic hydrocarbons may generate HOO^\bullet during their autoxidation, the efficiency of the process depends on the nature of the alkyl radical formed after their reaction with the chain-propagating peroxy radicals.

3.3. Semiquinones and phenolates

Hydroquinones are an important class of natural and synthetic redox-active organic molecules, able to switch between the oxidized (quinone), the radical (semiquinone) and the reduced (hydroquinone) forms. Many hydroquinones have antioxidant properties, and are used as additive to counteract lipid peroxidation [32]. After the reaction with ROO^\bullet , hydroquinones generate the corresponding semiquinones. The reaction with O_2 to produce HOO^\bullet or $\text{O}_2^{\bullet-}$ is a well-known process that is usually related to quinone toxicity [33]. From a mechanistic point of view, (protonated) semiquinones can produce hydroperoxyl radical by H-atom abstraction, reacting directly with molecular oxygen, or by an addition-fragmentation pathway [21,34]. However, the reaction of semiquinones with O_2 may have also unexpected antioxidant effect. Ingold and co-workers demonstrated that ubiquinol (QH_2) is a much more effective chain-breaking antioxidant than α -tocopherol (α -TOH) in low-density lipoproteins (LDL). Unexpectedly, after the reaction of α -TOH with a ROO^\bullet radical, the α -tocopheroxyl (α - TO^\bullet) radical remains trapped into the LDL. Being unable to decay, α - TO^\bullet eventually propagates the oxidative chain by reacting with polyunsaturated lipid. On the other hand, the semiquinone radical formed after the reaction of QH_2 reacts with oxygen to form the corresponding quinone and HOO^\bullet . The hydrophilic HOO^\bullet radical migrates from LDL to the aqueous phase, where forms after deprotonation the low-oxidizing $\text{O}_2^{\bullet-}$ radical, thus stopping the radical reaction inside the lipid core (Scheme 7) [35,36].



Scheme 7: Formation of HOO• from 1,4-semiquinone and protection of LDL autoxidation by HOO• production by the ubiquinol radical.

Superoxide is also produced by electron transfer to O₂ from deprotonated phenols [37]. This reaction is the basis for the autoxidation of aqueous solutions of polyphenols under O₂, that for example is responsible for degradation of polyphenolic antioxidants in cell culture medium [38] or to the formation of melanine-like polymers [39,40].

3.4. Kinetics aspects of Intra-1,4-HAT


The autoxidation of pro-aromatic 1,4-cyclohexadienes can be considered as an interesting strategy to have HOO• as chain-carrying radicals, in order to study their reactivity. For quantitative chemico-physical determinations, well-characterized oxidizable substrates must be chosen. Some examples of typical oxidizable substrates with known propagation and termination rate constants are given in Table 1, together with their respective k_p , $2k_t$, and oxidizability $Ox = k_p/(2k_t)^{1/2}$ values [15]. Interestingly, 1,4-cyclohexadiene displays a very large propagation constant, due to the bis-allylic nature of the reactive C-H groups, but at the same time it has a high termination constant, due to the

self-decay of HOO• radicals.

Table 1. Rate constants for the autoxidation of selected organic substrates in chlorobenzene at 30 °C.

Substrate	k_p [M ⁻¹ s ⁻¹]	$2k_t$ [M ⁻¹ s ⁻¹]	Ox	Ref
Cumene	0.32	4.6×10^4	0.0015	[41]
<i>p</i> -Cymene	0.83	2.9×10^6	0.0005	[42]
Styrene	41	4.2×10^7	0.0063	[43]
Methyl linoleate	62	8.8×10^6	0.0209	[43]
1,4-Cyclohexadiene	1400	1.3×10^9	0.0388	[10]


The generation of HOO• from the cleavage of ROO• radicals requires that the intra-1,4-HAT is faster than other reaction involving ROO•, like addition to double bonds or H-atom abstraction from the substrate, as shown in Scheme 8.



Scheme 8. Competition between fragmentation and propagation of the cyclohexadienyl peroxy radical. The rate of propagation is calculated for a hypothetical concentration of

methyl linoleate of 1 M from its propagation constant (see Table 1) [28,43].

In Scheme 9, selected experimental and theoretical values of intra-1,4-HAT are reported, including the HOO^\bullet production from the putative alkoxyperoxyl radical (see Scheme 5) formed during the autoxidation of styrene (a), of ethanol (b) and 1,4-cyclohexadiene (c). It is evident that the fragmentation is very effective for peroxy radicals deriving from pro-aromatic derivatives, while it is much slower for those deriving from alcohols and simple alkenes [24,28,44]. This clearly demonstrates the importance of aromatization in determining the production of HOO^\bullet [45].



Scheme 9: Rate constant for the elimination of hydroperoxyl from peroxy radicals by intra-1,4-HAT. ^a Calculated, at 70 °C [24], ^b Experimental, in water at 22 °C [15], ^c experimental, in acetonitrile at 20 °C [28].

From a quantitative point of view, considering that the propagation occurs with a pseudo-first order given by k_p [substrate], the HOO^\bullet generation in the case of styrene autoxidation

will be efficient only when $[\text{styrene}] \times 41 \text{ M}^{-1}\text{s}^{-1} < 4.4 \text{ s}^{-1}$, that is $[\text{styrene}] < 0.1 \text{ M}$. In case of the autoxidation of pure 1,4-cyclohexadiene ($k_p = 1400 \text{ M}^{-1}\text{s}^{-1}$), HOO^\bullet generation occurs at concentrations of the substrate $< 2.8 \text{ M}$. This line of reasoning can also be applied to mixtures of 1,4-cyclohexadiene and other oxidizable lipids. For example, with methyl linoleate in the 1 M concentration range (like in case of an oil rich in polyunsaturated lipid), HOO^\bullet generation by co-oxidizing 1,4-cyclohexadiene will be feasible, as the 1,4-intra-HAT ($4 \times 10^4 \text{ s}^{-1}$) is faster than the hypothetical propagation of the cyclohexadienyl radical ($1 \text{ M} \times 62 \text{ M}^{-1}\text{s}^{-1} = 62 \text{ s}^{-1}$) (see Scheme 8). The fast fragmentation of cyclohexadienyl radical is the basis for the use of 1,4-cyclohexadiene derivatives as additives to generate HOO^\bullet during the autoxidation of different substrates. We used this method to investigate the interaction of HOO^\bullet with antioxidants, as it is shown in the next sections of this review.

4. Synergy of HOO^\bullet with Antioxidants

4.1. Synergy of HOO^\bullet with Phenols and Aromatic Amines

The ability of HOO^\bullet to increase the length of the inhibition period, and thus the duration of the antioxidant effect, of an analogue of α -tocopherol, was demonstrated by us by studying the autoxidation of 1,4-cyclohexadiene [18] and was then confirmed in further studies by employing different phenols and pro-aromatic derivatives. For example, when studying the autoxidation of sunflower oil at $130 \text{ }^\circ\text{C}$, the addition of small amount of γ -terpinene was able to increase the inhibition period of the α -tocopherol naturally present in the oil. Experiments performed with stripped sunflower oil (that is, containing only triglycerides

and no endogenous tocopherols or carotenes) [46], showed that γ -terpinene itself had no retarding effect on the autoxidation, but it only acted in synergy with α -tocopherol [47] (see Figure 1).





Figure 1. A) Inhibition activity on the autoxidation of sunflower oil triglycerides (130 °C) of α -tocopherol (α -TOH) and γ -terpinene (γ -T), alone or in mixture. B) Inhibition period length due to α -TOH 0.1% and variable amounts of γ -T. Reprinted with permission from 47. Copyright 2023 American Chemical Society.

This effect was also visible using γ -terpinene in association with other oxidizable substrates, like styrene and squalene, and with different antioxidants like caffeic acid [48]. The mechanism of antioxidant regeneration is exemplified in Scheme 10.




Scheme 10. Reaction pathways for the reaction of HOO^\bullet and ROO^\bullet with (a) the analog of α -tocopherol PMHC, (b) 3,5-di-*tert*-butylcatechol.

Due to its double-faced oxidizing and reducing ability, HOO^\bullet can reduce the radical of the antioxidant and regenerate the parent phenol. However, especially in the case of phenols, this reaction does not occur indefinitely because HOO^\bullet is also capable to add to the phenoxyl ring, forming oxidized products unable to be regenerated. For catechols this regenerating effect is effectively observed even when the semiquinone is already fully oxidized to the corresponding *ortho*-quinone [48] (Scheme 10b).


Aromatic amines are also important radical-trapping antioxidants, used to protect petroleum-based products from autoxidation and limiting their deterioration. Some of them are able to preserve phospholipid membranes from peroxidation, preventing cell death by ferroptosis (see for instance ferrostatin-1, Fer-1 in Scheme 11) [49]. As reported by Ingold, superoxide thermal source (SOTS), can be used as $\text{O}_2^{\bullet-}$ source [50], see Scheme

11. Recently, Pratt and co-workers used this knowledge, to address whether $\text{HOO}^\bullet/\text{O}_2^{\bullet-}$ could regenerate aromatic amines when used as antioxidants to protect unsaturated phospholipids in liposomes. The kinetic data, including inhibition constant (k_{inh}) and stoichiometric factor (n) for different amines in the absence or presence of SOTS were calculated as shown in Scheme 11. As expected, the aliphatic amine 2,2,6,6-tetramethylpiperidine (TMP) is inactive both in the presence and absence of SOTS. On the contrary, the other aromatic amines show interesting results. For instance, the most active amine phenoxazine (PHOXN), had a rate constant of reaction with ROO^\bullet radicals (k_{inh}) approximately identical in both conditions, but showed an increase in the duration of its antioxidant activity (inhibited period) of about 8-times by adding SOTS into the system. This effect, although less evident, is also present for the other compounds reported in Scheme 11.



Scheme 11. Superoxide generation by SOTS, and kinetic data for the antioxidant activity of amines in the presence and absence of SOTS, autoxidation of liposomes at 37 °C [49].

Interestingly, under identical conditions, 2,2,5,7,8-pentamethyl-6-hydroxychroman (PMHC) and α -TOH, were not regenerated by $\text{HOO}^\bullet/\text{O}_2^{\bullet-}$, and conversely an erosion in the radical trapping stoichiometry was observed in the presence of SOTS [49]. This suggests that the H-atom transfer from HOO^\bullet to the phenoxyl radicals derived from α -TOH or PMHC is slow if compared with the addition of HOO^\bullet to form a stable peroxidic adduct. As already discussed, there are two parallel competing pathways when the phenoxyl radical of the phenolic antioxidant is generated: the regeneration of the starting antioxidant compound or the formation of the addition product which is no longer regenerable. For this reason, phenoxyl and aminyl radicals derived from α -TOH/PMHC and PHOXN respectively, perform differently. By CBS-QB3 calculations it was observed that the H-atom transfer from HOO^\bullet to the aminyl radical derived from PHOXN is similarly exergonic to the H-atom transfer from HOO^\bullet to the phenoxyl radical derived from PMHC (or α -TOH). On the other hand, the radical addition of HOO^\bullet with PHOXN-derived radical is 9.4 kcal/mol less advantageous than with the PMHC-derived radical (shown in Scheme 12), presumably because the C=N bond is much weaker than the C=O bond [49].




Scheme 12. Competing Reactions of HOO^\bullet with phenoxyl and aminyl radicals derived from α -TOH/PMHC and PHOXN [49].

4.2 Synergy of HOO• with Nitroxides and Quinones

When HOO• radicals are present, molecules that are devoid of antioxidant activity can act as autoxidation inhibitors. For example, the nitroxide 2,2,6,6-tetramethylpiperidinoxyl (TEMPO, Scheme 13a) doesn't react with alkylperoxyl radicals in organic solvents, but the HOO• radical assists the reduction of the stable radical TEMPO to the corresponding unstable hydroxylamine, that instead can react with ROO•. TEMPO is a strong antioxidant in water [51] and in organic solvents but only when weak acids are present [52,53]. In these circumstances, nitroxides trap ROO• by giving an electron to the ROO• radical, which is subsequently protonated by the solvent or by the acid added [54]. Instead, in non-protic systems, nitroxides are ineffective against organic substrate autoxidation. The addition of 1,4-cyclohexadiene (CHD) to the oxidizable substrate significantly boosts the antioxidant activity of the nitroxide TEMPO (Scheme 13) in non-protic solvents. The co-oxidation of 1,4-cyclohexadiene with the substrate causes the production of hydroperoxyl radicals, that can be effectively quenched by TEMPO. In this system a catalytic cycle is generated, resulting in a strong antioxidant effect, and an extremely long duration. The end is not visible in common reaction times, because peroxy radicals don't add to nitroxides. The cyclic mechanism can be interrupted only by the recombination of alkyl radicals with the nitroxide, but their concentration is low in the presence of O₂ [55]. This mechanism demonstrates why nitroxides are effective as antioxidants during the autoxidation of alkenes at high temperature where the chain-carrying radicals are both ROO• and HOO• [49]. Moreover, the antioxidant effect of TEMPO is visible in the presence of three more

model compounds, 2,5-di-*tert*-butylhydroquinone (QH₂), trimethylamine, and benzyl alcohol, which are known to generate HOO• upon interaction with free radicals (Scheme 13 b).




Scheme 13. Top: mechanism of the synergistic antioxidant activity of TEMPO with HOO•-

generating molecules. Lower panels: O₂ consumption during the autoxidation of styrene (4.3 M) in chlorobenzene initiated by AIBN at 30° without antioxidants (a in panels A, B, C and D), or in the presence of TEMPO 5 μM (b in panels A, B, C and D), CHD 0.27 M or CHD 0.27 M and TEMPO 5 μM (c and d in panel A), Et₃N 0.018 M or Et₃N 0.018 M and TEMPO 5 μM (c and d in panel B), QH₂ 5 μM or QH₂ 5 μM and TEMPO 5 μM (c and d in panel C), PhCH₂OH 0.24 M or PhCH₂OH 0.24 M and TEMPO 5 μM (c and d in panel D). Adapted with permission from reference 55. Copyright 2023 American Chemical Society.

Lastly, it has been observed that also quinones show antioxidant activity in the presence of hydroperoxyl radicals [56-58]. This observation is very counterintuitive considering that quinones are the oxidized form of catechols and hydroquinones. It is interesting to note that *ortho*-benzoquinones have a greater antioxidant impact than *para*-benzoquinones in organic solvents, since the *ortho* isomer's HAT from HOO• to the quinone is more exothermic [59]. However, unlike nitroxides, the semiquinone radicals also react via addition with ROO•/ HOO• radicals, shortening the duration of inhibition.

These unexpected properties can help us explain the antioxidant effect of polymeric nanoparticles based on melanin and polydopamine. These phenol-based compounds are rich in *ortho*-quinone moieties, and therefore in the presence of ROO•/HOO• systems, shows good radical trapping activity [59] (Scheme 14).



Scheme 14. Antioxidation mechanism of polydopamine nanoparticle [59].

In nonaqueous systems, the *ortho*-quinone moiety is better reduced by HOO• than the *para*-quinone one, and thus the former behaves as better antioxidant than the latter [59], in agreement with recent findings about the lack of superoxide-scavenging activity of thymoquinone in neural oxidative stress models [60]. Recently, we have demonstrated that honokiol-related neolignans, after the reaction with alkylperoxyl radicals, form quinonic species that are able to trap HOO• radicals similarly to polydopamine *ortho*-quinones [61]. This suggests that polyphenols with a catechol group (like quercetin or catechins) may also display a prolongation of their antioxidant effects when a source of HOO• is present in the system.

5. Conclusions

With this review, we have provided some recent examples of the importance of HOO• in the inhibition of autoxidation reactions by chain-breaking antioxidants. Despite the simplicity of this radical, many aspects of its chemistry are still unexplored. For example, the influence of HOO•/O₂^{•-} on the autoxidation of emulsions and LDL, and the capability of natural lipids like

fatty acids, phospholipids and phytosterols, to produce HOO^\bullet has not been investigated in detail. Due to the HOO^\bullet acidity, the relative importance of HOO^\bullet and $\text{O}_2^{\bullet-}$ may change in the different compartments of the cells, as at physiological pH deprotonated $\text{O}_2^{\bullet-}$ would be dominant, while at acidic pH (such as in lysosomes) [62,63] the main specie would be HOO^\bullet . It can be expected that the hydrophilicity of HOO^\bullet and its deprotonation to the not-oxidizing superoxide radical ($\text{O}_2^{\bullet-}$), explaining the reduction of radical chain propagation in LDL, may play a role also in other systems such as micelles, liposomes and nano / micro emulsions. At the same time, $\text{O}_2^{\bullet-}$ and HOO^\bullet are able to regenerate the antioxidants present at the interface, thus providing an increase of the radical trapping stoichiometry. In the case of aromatic amines, the regeneration by HOO^\bullet has been related to the efficacy to inhibit ferroptosis. The catalytic antioxidant behavior of redox-active nanomaterials (also called “nanozymes”) could also be explained by the interplay between HOO^\bullet and ROO^\bullet radicals. In conclusion, a better understanding of the HOO^\bullet role in oxidative radical chain processes is expected to increase our comprehension of the mechanisms of lipid peroxidation, and to foster the rational development of novel antioxidant systems.

Acknowledgements

Z. J. acknowledges a fellowship from China Scholarship Council (CSC: 202209120002).

Author contributions

R.A. planned the content, R. A., Z. J. and A.B. wrote and R.A and A.B. revised the manuscript.

Disclosure statement

The authors report no conflict of interest.

ORCID

Riccardo Amorati: <https://orcid.org/0000-0002-6417-9957>

Andrea Baschieri: <https://orcid.org/0000-0002-2108-8190>

References

- [1] Helberg J, Pratt DA. Autoxidation vs. antioxidants - the fight for forever. *Chem Soc Rev* 2021;50(13):7343-7358.
- [2] Foret MK, Lincoln R, Do Carmo S, Cuello AC, Cosa G. Connecting the "Dots": From Free Radical Lipid Autoxidation to Cell Pathology and Disease. *Chem Rev* 2020;120(23):12757-12787.
- [3] Amorati R, Valgimigli L. Advantages and limitations of common testing methods for antioxidants. *Free Radic Res* 2015;49(5):633-49.
- [4] Amorati R, Baschieri A, Valgimigli L. Measuring Antioxidant Activity in Bioorganic Samples by the Differential Oxygen Uptake Apparatus: Recent Advances. *Journal of Chemistry* 2017;2017:1-12.
- [5] Doba T, Burton GW, Ingold KU. Antioxidant and co-antioxidant activity of vitamin C. The effect of vitamin C, either alone or in the presence of vitamin E or a water-soluble vitamin E analogue, upon the peroxidation of aqueous multilamellar phospholipid liposomes. *Biochim Biophys Acta* 1985;835(2):298-303.
- [6] Kumar M, Francisco JS. Red-Light-Induced Decomposition of an Organic Peroxy Radical: A New Source of the HO₂ Radical. *Angew Chem Int Ed Engl* 2015;54(52):15711-4.
- [7] Anglada JM, Martins-Costa M, Francisco JS, Ruiz-Lopez MF. Interconnection of reactive oxygen species chemistry across the interfaces of atmospheric, environmental, and biological processes. *Acc Chem Res* 2015;48(3):575-83.
- [8] Colmenares JC, Luque R. Heterogeneous photocatalytic nanomaterials: prospects and challenges in selective transformations of biomass-derived compounds. *Chem Soc Rev* 2014;43(3):765-78.
- [9] Baschieri A, Amorati R, Valgimigli L, Sambri L. 1-Methyl-1,4-cyclohexadiene as a Traceless Reducing Agent for the Synthesis of Catechols and Hydroquinones. *J Org Chem* 2019;84(21):13655-13664.
- [10] Howard JA, Ingold KU. Absolute rate constants for hydrocarbon autoxidation. V. The hydroperoxy radical in chain propagation and termination. *Canadian Journal of Chemistry* 1967;45(8):785-792.
- [11] Denisov ET. Cyclic mechanisms of chain termination in the oxidation of organic compounds. *Russian Chemical Reviews* 1996;65(6):505-520.
- [12] Foti MC, Ingold KU. Mechanism of inhibition of lipid peroxidation by gamma-terpinene, an unusual and potentially useful hydrocarbon antioxidant. *J Agric Food Chem* 2003;51(9):2758-65.
- [13] Nolte TM, Peijnenburg W. Use of quantum-chemical descriptors to analyse reaction rate constants between organic chemicals and superoxide/hydroperoxyl (O₂^{•-}/HO[•]). *Free Radic Res* 2018;52(10):1118-1131.
- [14] Sawyer DT, Valentine JS. How super is superoxide? *Accounts of Chemical Research* 2002;14(12):393-400.
- [15] Guo Y, Amorati R. The Underrecognized Role of the Hydroperoxyl (HOO[•]) Radical in Chain

- Propagation of Lipids and its Implication in Antioxidant Activity. *Lipid Oxidation in Food and Biological Systems* 2022. p 115-132.
- [16] Das AB, Nagy P, Abbott HF, Winterbourn CC, Kettle AJ. Reactions of superoxide with the myoglobin tyrosyl radical. *Free Radic Biol Med* 2010;48(11):1540-7.
- [17] Agarwal RG, Coste SC, Groff BD, Heuer AM, Noh H, Parada GA, Wise CF, Nichols EM, Warren JJ, Mayer JM. Free Energies of Proton-Coupled Electron Transfer Reagents and Their Applications. *Chem Rev* 2022;122(1):1-49.
- [18] Cedrowski J, Litwinienko G, Baschieri A, Amorati R. Hydroperoxyl Radicals (HOO[•]): Vitamin E Regeneration and H-Bond Effects on the Hydrogen Atom Transfer. *Chemistry* 2016;22(46):16441-16445.
- [19] Hayyan M, Hashim MA, AlNashef IM. Superoxide Ion: Generation and Chemical Implications. *Chem Rev* 2016;116(5):3029-85.
- [20] Denisov ET, Afanas'ev IB. *Oxidation and Antioxidants in Organic Chemistry and Biology*. 2005.
- [21] Valgimigli L, Amorati R, Fumo MG, DiLabio GA, Pedulli GF, Ingold KU, Pratt DA. The unusual reaction of semiquinone radicals with molecular oxygen. *J Org Chem* 2008;73(5):1830-41.
- [22] Schumperli MT, Hammond C, Hermans I. Reactivity of alpha-amino-peroxyl radicals and consequences for amine oxidation chemistry. *Phys Chem Chem Phys* 2012;14(31):11002-7.
- [23] Howard JA, Yamada T. Absolute rate constants for hydrocarbon autoxidation. 31. Autoxidation of cumene in the presence of tertiary amines. *Journal of the American Chemical Society* 2002;103(24):7102-7106.
- [24] Harrison KA, Haidasz EA, Griesser M, Pratt DA. Inhibition of hydrocarbon autoxidation by nitroxide-catalyzed cross-dismutation of hydroperoxyl and alkylperoxyl radicals. *Chem Sci* 2018;9(28):6068-6079.
- [25] Baschieri A, Pulvirenti L, Muccilli V, Amorati R, Tringali C. Chain-breaking antioxidant activity of hydroxylated and methoxylated magnolol derivatives: the role of H-bonds. *Org Biomol Chem* 2017;15(29):6177-6184.
- [26] Roginsky V, Barsukova T. Superoxide dismutase inhibits lipid peroxidation in micelles. *Chem Phys Lipids* 2001;111(1):87-91.
- [27] Tikhonov IV, Pliss EM, Borodin LI, Sen' VD. Effect of superoxide dismutase on the oxidation of methyl linoleate in micelles inhibited by nitroxyl radicals. *Russian Chemical Bulletin* 2017;65(12):2985-2987.
- [28] Sortino S, Petralia S, Foti MC. Absolute rate constants and transient intermediates in the free-radical-induced peroxidation of γ -terpinene, an unusual hydrocarbon antioxidant. *New J. Chem.* 2003;27(11):1563-1567.
- [29] Vardanyan RL, Denisov ET. Regeneration of inhibitors in oxidized 1,3-cyclohexadiene. *Bulletin of the Academy of Sciences of the USSR Division of Chemical Science* 1971;20(12):2681-2683.
- [30] Rudback J, Bergstrom MA, Borje A, Nilsson U, Karlberg AT. alpha-Terpinene, an antioxidant in tea tree oil, autoxidizes rapidly to skin allergens on air exposure. *Chem Res Toxicol* 2012;25(3):713-21.
- [31] Do Q, Lee DD, Dinh AN, Seguin RP, Zhang R, Xu L. Development and Application of a Peroxyl Radical Clock Approach for Measuring Both Hydrogen-Atom Transfer and Peroxyl Radical Addition Rate Constants. *J Org Chem* 2021;86(1):153-168.
- [32] Mollica F, Bonoldi L, Amorati R. Kinetic Analysis of High-Temperature Sunflower Oil Peroxidation Inhibited by the Major Families of Phenolic Antioxidants Unveils the Extraordinary Activity of 1,4-Hydroquinones. *Antioxidants (Basel)* 2022;11(11).

- [33] Bolton JL, Trush MA, Penning TM, Dryhurst G, Monks TJ. Role of quinones in toxicology. *Chem Res Toxicol* 2000;13(3):135-60.
- [34] Korth HG, Mulder P. Phenolic Hydrogen Transfer by Molecular Oxygen and Hydroperoxyl Radicals. Insights into the Mechanism of the Anthraquinone Process. *J Org Chem* 2020;85(4):2560-2574.
- [35] Ingold KU, Bowry VW, Stocker R, Walling C. Autoxidation of lipids and antioxidation by alpha-tocopherol and ubiquinol in homogeneous solution and in aqueous dispersions of lipids: unrecognized consequences of lipid particle size as exemplified by oxidation of human low density lipoprotein. *Proc Natl Acad Sci U S A* 1993;90(1):45-9.
- [36] Bowry VW, Stocker R. Tocopherol-mediated peroxidation. The prooxidant effect of vitamin E on the radical-initiated oxidation of human low-density lipoprotein. *Journal of the American Chemical Society* 2002;115(14):6029-6044.
- [37] Lee D-H, Son JB, Jung S, Song J, Ham SW. Mechanism of action of base-catalyzed oxygenation of phenol derivatives. *Tetrahedron Letters* 2005;46(45):7721-7723.
- [38] Xiao J, Hogger P. Stability of dietary polyphenols under the cell culture conditions: avoiding erroneous conclusions. *J Agric Food Chem* 2015;63(5):1547-57.
- [39] d'Ischia M, Napolitano A, Ball V, Chen CT, Buehler MJ. Polydopamine and eumelanin: from structure-property relationships to a unified tailoring strategy. *Acc Chem Res* 2014;47(12):3541-50.
- [40] Mollica F, Lucernati R, Amorati R. Expanding the spectrum of polydopamine antioxidant activity by nitroxide conjugation. *J Mater Chem B* 2021;9(48):9980-9988.
- [41] Valgimigli L, Bartolomei D, Amorati R, Haidasz E, Hanthorn JJ, Nara SJ, Brinkhorst J, Pratt DA. 3-Pyridinols and 5-pyrimidinols: Tailor-made for use in synergistic radical-trapping co-antioxidant systems. *Beilstein J Org Chem* 2013;9:2781-92.
- [42] Zielinski ZA, Pratt DA. Lipid Peroxidation: Kinetics, Mechanisms, and Products. *J Org Chem* 2017;82(6):2817-2825.
- [43] Howard JA, Ingold KU. Absolute rate constants for hydrocarbon autoxidation. VI. Alkyl aromatic and olefinic hydrocarbons. *Canadian Journal of Chemistry* 1967;45(8):793-802.
- [44] Bothe E, Behrens G, Schulte-Frohlinde D. Mechanism of the First Order Decay of 2-Hydroxy-propyl-2-peroxyl Radicals and of O₂ Formation in Aqueous Solution. *Zeitschrift für Naturforschung B* 1977;32(8):886-889.
- [45] Walton JC, Studer A. Evolution of functional cyclohexadiene-based synthetic reagents: the importance of becoming aromatic. *Acc Chem Res* 2005;38(10):794-802.
- [46] Mollica F, Gelabert I, Amorati R. Correction to "Synergic Antioxidant Effects of the Essential Oil Component γ -Terpinene on High Temperature Oil Oxidation". *ACS Food Science & Technology* 2022;2(6):1050-1050.
- [47] Mollica F, Gelabert I, Amorati R. Synergic Antioxidant Effects of the Essential Oil Component γ -Terpinene on High-Temperature Oil Oxidation. *ACS Food Science & Technology* 2022;2(1):180-186.
- [48] Guo Y, Baschieri A, Amorati R, Valgimigli L. Synergic antioxidant activity of gamma-terpinene with phenols and polyphenols enabled by hydroperoxyl radicals. *Food Chem* 2021;345:128468.
- [49] Poon JF, Zilka O, Pratt DA. Potent Ferroptosis Inhibitors Can Catalyze the Cross-Dismutation of Phospholipid-Derived Peroxyl Radicals and Hydroperoxyl Radicals. *J Am Chem Soc* 2020;142(33):14331-14342.
- [50] Ingold KU, Paul T, Young MJ, Doiron L. Invention of the First Azo Compound To Serve as a Superoxide Thermal Source under Physiological Conditions: Concept, Synthesis, and Chemical Properties. *Journal of the American Chemical Society* 1997;119(50):12364-12365.

- [51] Griesser M, Shah R, Van Kessel AT, Zilka O, Haidasz EA, Pratt DA. The Catalytic Reaction of Nitroxides with Peroxyl Radicals and Its Relevance to Their Cytoprotective Properties. *J Am Chem Soc* 2018;140(10):3798-3808.
- [52] Amorati R, Pedulli GF, Pratt DA, Valgimigli L. TEMPO reacts with oxygen-centered radicals under acidic conditions. *Chem Commun (Camb)* 2010;46(28):5139-41.
- [53] Haidasz EA, Meng D, Amorati R, Baschieri A, Ingold KU, Valgimigli L, Pratt DA. Acid Is Key to the Radical-Trapping Antioxidant Activity of Nitroxides. *J Am Chem Soc* 2016;138(16):5290-8.
- [54] Genovese D, Baschieri A, Vona D, Baboi RE, Mollica F, Prodi L, Amorati R, Zaccheroni N. Nitroxides as Building Blocks for Nanoantioxidants. *ACS Appl Mater Interfaces* 2021;13(27):31996-32004.
- [55] Baschieri A, Valgimigli L, Gabbanini S, DiLabio GA, Romero-Montalvo E, Amorati R. Extremely Fast Hydrogen Atom Transfer between Nitroxides and HOO[•] Radicals and Implication for Catalytic Coantioxidant Systems. *J Am Chem Soc* 2018;140(32):10354-10362.
- [56] Denisov ET. Mechanism of the inhibiting effect of quinone in the chain reaction of oxidation of isopropyl alcohol. *Bulletin of the Academy of Sciences of the USSR Division of Chemical Science* 1969;18(2):282-285.
- [57] Vardanyan RL, Denisov ET, Zozulya VI. Measurement of the rate constants of the reaction of peroxy radicals with alcohols by the interlinked oxidation method with a selective inhibitor. *Bulletin of the Academy of Sciences of the USSR Division of Chemical Science* 1972;21(3):572-574.
- [58] Degtyareva TG, Denisov ET, Martem'yanov VS, Badretdinova LY. The activity of polyatomic alcohols in reactions with peroxide radicals. *Bulletin of the Academy of Sciences of the USSR Division of Chemical Science* 1979;28(6):1139-1144.
- [59] Guo Y, Baschieri A, Mollica F, Valgimigli L, Cedrowski J, Litwinienko G, Amorati R. Hydrogen Atom Transfer from HOO[•] to ortho-Quinones Explains the Antioxidant Activity of Polydopamine. *Angew Chem Int Ed Engl* 2021;60(28):15220-15224.
- [60] Krewenka, C.; Rizzi, S.; Nguyen, C.H.; Delijewski, M.; Gille, L.; Staniek, K.; Duvigneau, J.C.; Radad, K.; Müllebner, A.; Kranner, B.; Moldzio, R. Radical Scavenging Is Not Involved in Thymoquinone-Induced Cell Protection in Neural Oxidative Stress Models. *Antioxidants* 2023, 12, 858.
- [61] Cardullo, N.; Monti, F.; Muccilli, V.; Amorati, R.; Baschieri, A. Reaction with ROO[•] and HOO[•] Radicals of Honokiol-Related Neolignan Antioxidants. *Molecules* 2023, 28, 735.
- [62] Ojo OO, Leake DS. Vitamins E and C do not effectively inhibit low density lipoprotein oxidation by ferritin at lysosomal pH. *Free Radic Res* 2021;55(5):525-534.
- [63] Ojo OO, Leake DS. Effects of lysosomal low-density lipoprotein oxidation by ferritin on macrophage function. *Free Radic Res*, 2022;56(5-6): 436-446.