



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Detection of *Lotmaria passim* in honeybees from Emilia Romagna (Italy) based on a culture method

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Rudelli C., Isani G., Andreani G., Tedesco P., Galuppi R. (2023). Detection of *Lotmaria passim* in honeybees from Emilia Romagna (Italy) based on a culture method. *JOURNAL OF INVERTEBRATE PATHOLOGY*, 201, 1-4 [10.1016/j.jip.2023.108007].

Availability:

This version is available at: <https://hdl.handle.net/11585/953446> since: 2024-01-18

Published:

DOI: <http://doi.org/10.1016/j.jip.2023.108007>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 Detection of *Lotmaria passim* in honeybees from Emilia Romagna (Italy) by cultural method

2

3 Cecilia Rudelli, Gloria Isani, Giulia Andreani, Perla Tedesco*, Roberta Galuppi

4 Department of Veterinary Medical Sciences, Alma Mater Studiorum University of Bologna, via

5 Tolara di sopra 50, Ozzano dell'Emilia, 40064 Bologna, Italy

6 *Corresponding author: perla.tedesco@unibo.it

7

8 Abstract:

9 *Lotmaria passim* is considered an emerging field of study in honeybee pathology, since it can
10 threaten the health of the colony leading to a higher mortality rate. However, there is a lack of
11 knowledge regarding the diffusion of this trypanosomatid in Italy. In this study, we highlight the
12 presence of *L. passim* in the province of Bologna through its culture isolation from honeybee guts
13 and microscopic observation.

14

15 Keywords: *Apis mellifera*, *Lotmaria passim*, trypanosomatid, cultural method, morphological
16 identification, honeybee pathogens

17

18 1. Introduction

19 The family Trypanosomatidae (Euglenozoa, Kinetoplastea) consists of 25 genera with a wide host
20 range, including vertebrates and insects (Maslov et al., 2013; Kostygov et al., 2021). Until a few
21 years ago, in honeybee, the report of monoxenous trypanosomatids in the digestive system was
22 limited to only two species: *Leptomonas apis* Lotmar 1946, no longer reported, and *Crithidia*
23 *mellificae* Langridge & McGhee, 1967. The presence of trypanosomatids in *Apis mellifera* has been
24 increasingly recognized, in particular after the description of the species *Lotmaria passim* (Schwarz
25 et al., 2015), which nowadays is considered as the most widespread bee trypanosomatid all over the
26 world (Arismendi et al., 2016; Stevanovic et al., 2016; Vargas et al., 2017; Williams et al., 2019
27 Castelli et al., 2019; Ribani et al., 2021; Michalczyk & Sokół, 2022). It has been suggested that
28 trypanosomatids infestation could be related to an impairment of bee immune system, a nutritional
29 deficit and an alteration of bee behavior (Buendía-Abad et al., 2022; Gómez-Moracho et al., 2020;
30 Liu et al., 2020; Lukeš et al., 2018); however, the details of the pathogenic effects are still not fully
31 understood.

32 Recently, studies have begun to deepen the interaction of *L. passim* with other well-known bee
33 pathogen: Arismendi et al. (2020) have explored the possible synergic effect on the survival of
34 honeybees infected at the same time by *Nosema ceranae* and *L. passim*; Quintana et al. (2021) have

35 reported the finding of *L. passim* in the mite *Varroa destructor*, while Nanetti et al. (2021) in
36 *Aethina tumida*. Despite its wide range of distribution, there is a lack of knowledge about the real
37 diffusion in the bee colonies of *L. passim*, and in particular few studies have evaluated its
38 distribution in Italy (Bordin et al., 2022; Cilia et al., 2022; Ribani et al., 2021).
39 At present, methods based on DNA assay are the most used to detect the presence of *L. passim*
40 (Arismendi et al., 2016; Castelli et al., 2019; Cilia et al., 2022; Michalczyk & Sokół, 2022;
41 Stevanovic et al., 2016), and few publications have reported the isolation of the parasite directly
42 from the honeybee's intestine (Schwarz et al., 2015; Buendía-Abad et al., 2021; Ribani et al., 2021).
43 In the present preliminary study, we aimed to detect the presence of trypanosomatids in *A. mellifera*
44 from different apiaries of Emilia-Romagna region (Italy) through its direct isolation from the
45 intestine of single honeybees and microscopic observation.

46

47 2. Material and Methods

48 Five apiaries have been included in this study: four (A, B, C, D) belonging to the same beekeeper
49 are in three municipalities of the province of Bologna, and one belonging to a different beekeeper in
50 the province of Rimini. The apiaries A and B are in the suburban area of Bologna (54 m above sea
51 level), the apiary C is located in Argelato (25 m a.s.l.), a town rich of cropland, while the apiary D
52 is located in Pianoro (200 m a.s.l.), surrounded by semi-intensive vineyards. The last is an organic
53 apiary in the municipality of Montescudo (province of Rimini, 209 m a.s.l.) in an area characterized
54 by semi-intensive cultivations. All the samples were collected in spring and summer 2022.

55 From each apiary of the province of Bologna, 6 colonies were selected, and 15 honeybees were
56 sampled from each colony, while in the apiary of the province of Rimini 3 colonies were selected
57 and 21 honeybees were sampled from each one. The specimens were collected from the external
58 frames in all the colonies investigated. The search for intestinal flagellates was performed on single
59 guts sampled from each honeybee using the culture method previously reported (Ribani et al.,
60 2021). Briefly, the gut of each honeybee was dipped and grinded in 0.5 mL of supplemented DS2
61 medium and incubated at 26 °C. Wet mount slides with 10 µL of each culture were observed with
62 light microscope at 3 and 7 days after the incubation, to verify the presence of free active
63 flagellates. Some cultures were maintained by subculture steps every 4–10 days in fresh medium
64 (ratio 1:5).

65 Morphological observation insights and image acquisition were performed on some positive
66 cultures, in May-Grunwald Giemsa stained slides, at 400 × and 1000 × magnification through Leica
67 DMLS light microscope (Leica, Wetzlar, Germany), equipped with a digital camera Nikon DS-Fi2
68 with imaging software NIS Elements 4.10.01 (Nikon, Tokyo, Japan). The scanning electron

69 microscopy (SEM) analysis was also performed as follows: pelleted cells were fixed with 3%
70 glutaraldehyde in phosphate buffer (for 2 h at room temperature, then overnight at 4 °C), washed
71 three times in PBS, dehydrated in a graded ethanol series and dried with hexamethyldisilazane.
72 Subsequently, samples were mounted on aluminum stubs, sputter coated with gold-palladium using
73 a SC7620 Mini Sputter Coater (Quorum Technologies) and observed using a Phenom XL G2
74 Desktop SEM operating at 10 kV. Axenic cultures of reference strains of *C. mellifica* (ATCC
75 30254) and *L. passim* (Ribani et al., 2021) maintained on the same medium at the same condition of
76 incubation were used for morphological comparisons.

77

78 3. Results and Discussion

79 To microscopical examination, the presence of trypanosomatid flagellates was observed in several
80 cultures of bee gut, mostly after 7 days of incubation. Overall, of the 360 bees analyzed from the
81 Bologna province, 81 (22.5%) were positive for flagellates in culture [CI 95%; 18.19 - 26.81]. The
82 total positivity rate was similar to the ones of each apiary (Table 1). No positivity occurred in the
83 samples from the province of Rimini.

84

85

Apiary	n. examined colonies from each apiary	n. examined honeybees for each colony	average number of positive honeybees for each colony (min-max)	number of honeybees for each apiary	positive honeybees for each apiary (number and %)	confidence interval IC 95%
Bologna A	6	15	3.5 (2-5)	90	21 (23.3%)	14.6 – 32.0
Bologna B	6	15	3.5 (0-7)	90	21 (23.3%)	14.6 – 32.0
Bologna C	6	15	2.6 (0-10)	90	16 (17.7%)	9.8 – 25.3
Bologna D	6	15	3.8 (0-9)	90	23 (25.5%)	16.5 – 34.5
Rimini	3	21	0	63	0	0

86

87 Table 1. Maximum and minimum number of honeybees positive to flagellate morphologically
88 identified as *Lotmaria passim* for each apiary, positivity frequencies and confidence interval in the
89 apiaries.

90

91 Morphological observations performed either in wet slides or in slides stained with May Grunwald-
92 Giemsa showed predominant elongated and tear-drop shaped cells typical of a promastigote
93 morphotype. The cells narrowed posteriorly to a short caudate (tail-like) extension, often with a
94 characteristic “nose” of the posterior end (Figure 1A, B, and C), consistent with the description of

95 *L. passim* (Schwarz et al., 2015; Ribani et al., 2021; Buendía-Abad et al., 2022). These
96 characteristics were confirmed by SEM observation of some specimens. Moreover, in wet mounts,
97 these trypanosomatids were actively moving. The morphology observed in the isolates, also in wet
98 slides, clearly differed from that of a *C. mellificae* strain (ATCC 30254) grown in the same culture
99 medium and incubation temperature (Figure 1- D, E, and F), used as a control. Moreover, *C.*
100 *mellificae* showed much slower movements. Therefore, we can assume that the isolates observed in
101 the present study can be attributable to *L. passim*.

102 To date, prevalence studies available in the literature have been carried out using molecular
103 methods such as PCR (Bartolomé et al., 2018; Bordin et al., 2022; Stevanovic et al., 2016), or real
104 time PCR (Arismendi et al., 2022; Cilia et al., 2022; Vajnovic et al., 2018; Xu et al., 2018) that
105 quantified the parasitic load of *Lotmaria* on pools of bees. To our knowledge, no study evaluated
106 the frequency of positive subjects within an infected colony. This information could be useful to
107 better understand the degree of the parasite's diffusion among cohabiting bees, and to define in
108 more detail the sample size needed to evaluate the presence/absence of the parasite. Moreover,
109 albeit time consuming, isolation methods allow to observe the presence of living flagellates,
110 amplifying their number (observation after 7 days gave us the highest number of positives), thus
111 allowing to verify the actual colonization of the bee intestine and to obtain strains which could be
112 further studied.

113 In the present paper, *L. passim* was absent in the bees of the organic apiary from the province of
114 Rimini, while it was observed in all four apiaries tested in the province of Bologna. This high
115 frequency may be due to the fact that these latter apiaries were held by the same beekeeper, who is
116 used to exchange frames and brood between colonies. A wide variability was observed among the
117 colonies of the same apiary, from negative ones to colonies with up to 66.6% of positive bees. This
118 variability could explain the failure to find positive apiaries in Emilia Romagna by Cilia et al.
119 (2022) by PCR on pools of ten bees from each colony. Differently, Ribani et al. (2021) reported in
120 the same region a percentage of 88% of honey samples from different apiaries positive to *L. passim*
121 DNA, suggesting a high diffusion of the parasite in the region.

122 Due to the limited number of colonies analyzed in the present study, it would not be appropriate to
123 consider the results indicative of the prevalence of *L. passim* in the apiaries of Emilia-Romagna.
124 Nevertheless, the current study provides evidence of the presence of bees actually colonized by this
125 trypanosomatid in this region, and underlines the need for a deeper investigation regarding the
126 epidemiology of this flagellate in Italy.

127

128 Declarations of interest: none.

129 Acknowledgements

130 This study was funded by Regione Emilia-Romagna, BEE-RER-3 project—CUP
131 E37G22000030007—del Regolamento (UE) no. 1308/2013 (OCM Apicoltura).

132

133 Arismendi, N., Bruna, A., Zapata, N., & Vargas, M. (2016). PCR-specific detection of recently
134 described *Lotmaria passim* (Trypanosomatidae) in Chilean apiaries. *Journal of Invertebrate*
135 *Pathology*, *134*, 1–5. <https://doi.org/10.1016/j.jip.2015.12.008>

136 Arismendi, N., Caro, S., Castro, M. P., Vargas, M., Riveros, G., & Venegas, T. (2020). Impact of
137 mixed infections of gut parasites *Lotmaria passim* and *Nosema ceranae* on the lifespan and
138 immune-related biomarkers in *Apis mellifera*. *Insects*, *11*(7), 1–12.
139 <https://doi.org/10.3390/insects11070420>

140 Arismendi, N., Castro, M. P., Vargas, M., Zapata, C., & Riveros, G. (2022). The trypanosome
141 *Lotmaria passim* prevails in honey bees of different ages and stages of development. *Journal*
142 *of Apicultural Research*, *61*(1), 63–69. <https://doi.org/10.1080/00218839.2020.1828239>

143 Bartolomé, C., Buendía, M., Benito, M., De la Rúa, P., Ornos, C., Martín-Hernández, R., Higes,
144 M., & Maside, X. (2018). A new multiplex PCR protocol to detect mixed trypanosomatid
145 infections in species of *Apis* and *Bombus*. *Journal of Invertebrate Pathology*, *154*, 37–41.
146 <https://doi.org/10.1016/j.jip.2018.03.015>

147 Bordin, F., Zulian, L., Granato, A., Caldon, M., Colamonico, R., Toson, M., Trevisan, L., Biasion,
148 L., & Mutinelli, F. (2022). Presence of Known and Emerging Honey Bee Pathogens in
149 Apiaries of Veneto Region (Northeast of Italy) during Spring 2020 and 2021. *Applied*
150 *Sciences*, *12*(4). <https://doi.org/10.3390/app12042134>

151 Buendía-Abad, M., Higes, M., Martín-Hernández, R., Barrios, L., Meana, A., Fernández Fernández,
152 A., Osuna, A., & De Pablos, L. M. (2021). Workflow of *Lotmaria passim* isolation:
153 Experimental infection with a low-passage strain causes higher honeybee mortality rates than
154 the PRA-403 reference strain. *International Journal for Parasitology: Parasites and Wildlife*,
155 *14*(December 2020), 68–74. <https://doi.org/10.1016/j.ijppaw.2020.12.003>

156 Buendía-Abad, M., García-Palencia, P., de Pablos, L. M., Alunda, J. M., Osuna, A., Martín-
157 Hernández, R., & Higes, M. (2022). First description of *Lotmaria passim* and *Crithidia mellificae*
158 haptomonad stages in the honeybee hindgut. *International Journal for Parasitology*, *52*(1), 65–75.
159 <https://doi.org/10.1016/j.ijpara.2021.06.005>

160 Castelli, L., Branchiccela, B., Invernizzi, C., Tomasco, I., Basualdo, M., Rodriguez, M., Zunino, P.,
161 & Antúnez, K. (2019). Detection of *Lotmaria passim* in Africanized and European honey bees
162 from Uruguay, Argentina and Chile. *Journal of Invertebrate Pathology*, *160*(November 2018),

163 95–97. <https://doi.org/10.1016/j.jip.2018.11.004>

164 Cilia, G., Tafi, E., Zavatta, L., Caringi, V., & Nanetti, A. (2022). The Epidemiological Situation of
165 the Managed Honey Bee (*Apis mellifera*) Colonies in the Italian Region Emilia-Romagna.
166 *Veterinary Sciences*, 9(8), 1–15. <https://doi.org/10.3390/vetsci9080437>

167 Gómez-Moracho, T., Buendía-Abad, M., Benito, M., García-Palencia, P., Barrios, L., Bartolomé,
168 C., Maside, X., Meana, A., Jiménez-Antón, M.D., Olías-Molero, A.I., Alunda, J.M., Martín-
169 Hernández, R., & Higes, M. (2020). Experimental evidence of harmful effects of *Crithidia*
170 *mellificae* and *Lotmaria passim* on honey bees. *International Journal for Parasitology*, 50(13),
171 1117-1124. <https://doi.org/10.1016/j.ijpara.2020.06.009>

172 Kostygov, A. Y., Karnkowska, A., Votýpka, J., Tashyreva, D., Maciszewski, K., Yurchenko, V., &
173 Lukeš, J. (2021). Euglenozoa: taxonomy, diversity and ecology, symbioses and viruses. *Open*
174 *Biology*, 11(3), 200407. <https://doi.org/10.1098/rsob.200407>

175 Liu, Q., Lei, J., Darby, A.C., & Kadowaki, T. (2020). Trypanosomatid parasite dynamically
176 changes the transcriptome during infection and modifies honey bee physiology.
177 *Communications Biology*. 3, 51. <https://doi.org/10.1038/s42003-020-0775-x>.

178 Lukeš, J., Butenko, A., Hashimi, H., Maslov, D. A., Votýpka, J., & Yurchenko, V. (2018).
179 Trypanosomatids Are Much More than Just Trypanosomes: Clues from the Expanded Family
180 Tree. *Trends in Parasitology*, 34(6), 466–480. <https://doi.org/10.1016/j.pt.2018.03.002>

181 Maslov, D. A., Votýpka, J., Yurchenko, V., & Lukeš, J. (2013). Diversity and phylogeny of insect
182 trypanosomatids: all that is hidden shall be revealed. *Trends in parasitology*, 29(1), 43-52.
183 <https://doi.org/10.1016/j.pt.2012.11.001>

184 Michalczyk, M., & Sokół, R. (2022). Detection of *Lotmaria passim* and *Crithidia mellificae* in
185 Selected Bumblebee Species. *Pathogens*, 11(9). <https://doi.org/10.3390/pathogens11091053>

186 Nanetti, A., Ellis, J. D., Cardaio, I., & Cilia, G. (2021). Detection of *Lotmaria passim*, *Crithidia*
187 *mellificae* and Replicative Forms of Deformed Wing Virus and Kashmir Bee Virus in the
188 Small Hive Beetle (*Aethina tumida*). *Pathogens (Basel, Switzerland)*, 10(3).
189 <https://doi.org/10.3390/pathogens10030372>

190 Quintana, S., Plischuk, S., Brasesco, C., Revainera, P., Genchi García, M. L., Bravi, M. E.,
191 Reynaldi, F., Eguaras, M., & Maggi, M. (2021). *Lotmaria passim* (Kinetoplastea:
192 Trypanosomatidae) in honey bees from Argentina. *Parasitology International*, 81(May 2020),
193 102244. <https://doi.org/10.1016/j.parint.2020.102244>

194 Ribani, A., Utzeri, V. J., Taurisano, V., Galuppi, R., & Fontanesi, L. (2021). Analysis of honey
195 environmental DNA indicates that the honey bee (*Apis mellifera* L.) trypanosome parasite
196 *Lotmaria passim* is widespread in the apiaries of the North of Italy. *Journal of Invertebrate*

197 *Pathology*, 184(March), 107628. <https://doi.org/10.1016/j.jip.2021.107628>

198 Schwarz, R. S., Bauchan, G. R., Murphy, C. A., Ravoet, J., de Graaf, D. C., & Evans, J. D. (2015).
199 Characterization of Two Species of Trypanosomatidae from the Honey Bee *Apis mellifera*:
200 *Crithidia mellificae* Langridge and McGhee, and *Lotmaria passim* n. gen., n. sp. *The Journal*
201 *of Eukaryotic Microbiology*, 62(5), 567–583. <https://doi.org/10.1111/jeu.12209>

202 Stevanovic, J., Schwarz, R. S., Vejnovic, B., Evans, J. D., Irwin, R. E., Glavinic, U., &
203 Stanimirovic, Z. (2016). Species-specific diagnostics of *Apis mellifera* trypanosomatids: A
204 nine-year survey (2007–2015) for trypanosomatids and microsporidians in Serbian honey bees.
205 *Journal of Invertebrate Pathology*, 139, 6–11.
206 <https://doi.org/https://doi.org/10.1016/j.jip.2016.07.001>

207 Vargas, M., Arismendi, N., Riveros, G., Zapata, N., Bruna, A., Vidal, M., Rodriguez, M., &
208 Gerding, M. (2017). Viral and intestinal diseases detected in *Apis mellifera* in Central and
209 Southern Chile. *Chilean journal of agricultural research*, 77(3), 243-249.
210 <http://dx.doi.org/10.4067/S0718-58392017000300243>

211 Vejnovic, B., Stevanovic, J., Schwarz, R.S., Aleksic, N., Mirilovic, M., Jovanovic, N.M.,
212 Stanimirovic, Z. (2018). Quantitative PCR assessment of *Lotmaria passim* in *Apis mellifera*
213 colonies co-infected naturally with *Nosema ceranae*. *Journal of Invertebrate Pathology*, 151,
214 76-81. <https://doi.org/10.1016/j.jip.2017.11.003>

215 Williams M.K.F., Tripodi A.D., Szalanski A.L. (2019). Molecular survey for the honey bee (*Apis*
216 *mellifera* L.) trypanosome parasites *Crithidia mellificae* and *Lotmaria passim*. *Journal of*
217 *Apicultural Research*, 58(4), 553 - 558. <https://doi.org/10.1080/00218839.2019.1568956>

218 Xu, G., Palmer-Young, E., Skyrn, K., Daly, T., Sylvia, M., Averill, A., & Rich, S. (2018). Triplex
219 real-time PCR for detection of *Crithidia mellificae* and *Lotmaria passim* in honey bees.
220 *Parasitology Research*, 117(2), 623–628. <https://doi.org/10.1007/s00436-017-5733-2>

221

222

223 Figure captions:

224 Figure 1: Wet slides (A, D), May Grunwald-Giemsa-stained slides (B, E), and SEM (C, F) images
225 from cell cultures on Insectagro DS2 medium of a field isolate of *Lotmaria passim* (A, B, C) and
226 *Crithidia mellificae* ATCC 30254 (D, E, F). Bar = 10 micron (A-D and F) and 15 micron (C).

227