



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

A tentative I/O curve with consciousness: Effects of multiple simultaneous ambiguous figures presentation on perceptual reversals and time estimation

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

D'Anselmo A., Pisani A., Brancucci A. (2022). A tentative I/O curve with consciousness: Effects of multiple simultaneous ambiguous figures presentation on perceptual reversals and time estimation. *CONSCIOUSNESS AND COGNITION*, 99, 1-11 [10.1016/j.concog.2022.103300].

Availability:

This version is available at: <https://hdl.handle.net/11585/955191> since: 2024-02-01

Published:

DOI: <http://doi.org/10.1016/j.concog.2022.103300>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 A tentative I/O curve with consciousness: effects of multiple simultaneous
2 ambiguous figures presentation on perceptual reversals and time
3 estimation

4

5 Anita D'Anselmo¹, Angelo Pisani², Alfredo Brancucci³

6 *1) Department of Psychological, Health and Territory Sciences (DiSPuTeR), "G. d'Annunzio" University of Chieti-Pescara, Italy*

7 *2) Department of Psychology and Cognitive Sciences, University of Trento, Italy*

8 *3) Department of Motor, Human and Health Sciences, University of Rome "Foro Italico", Italy*

9

10

11

12 **Abstract**

13 This study was aimed at investigating mechanisms of consciousness using bistable perception. In 4
14 experimental conditions, 1, 2, 4 or 8 Rubin's face-vase ambiguous figures were presented for 3 minutes. In
15 Experiment 1, 40 subjects looked at the center of the screen and pressed a specific key correspondent to the
16 figure where they perceived a reversal. In Experiment 2, 32 subjects controlled with eye-tracker performed a
17 similar task in which they pressed the spacebar whenever they perceived a reversal in any of the figures. At
18 the end of each condition subjects estimated its duration. Results showed that changing the number of figures
19 does not alter the number of reversals, producing a flat I/O curve between the two parameters. Estimated
20 time lapse showed a negative correlation with the number of reversals. These findings are discussed
21 considering the relationships between bistable perception, attention, and consciousness, as well as the time
22 perception literature.

23

24

25

26

27

28 **Keywords:** consciousness, attention, bistable perception, time perception.

29

30 INTRODUCTION

31 In the last two decades, the study of the behavioural and neural bases of consciousness has found a
32 prominent place in the literature of psychology and neuroscience (De Graaf et al., 2012; Koch et al.,
33 2016; Zeman, 2001). Considerable advancements have been made in the field on both the empirical
34 (Overgaard, 2017) and theoretical aspects of research (Dehaene & Naccache, 2001). Some scientists
35 have even begun to measure diverse features of consciousness creating in fact a mathematics of
36 experience (Balduzzi & Tononi, 2009; Seth et al., 2008) and bringing thus consciousness to all effects
37 in the ranks of the empirical sciences.

38 One broadly used paradigm to investigate the behavioural and neural aspects of consciousness is
39 based on ambiguous figures. These figures are physically invariant pictures allowing more than one
40 interpretation. They give often rise to bistable percepts, that is, to perceptual experiences made up
41 by a sequence of different visual images elicited by the objects which are intrinsically part of the
42 figure. Most ambiguous figures allow two object interpretations, whose perception reverses
43 alternately from one to the other. This perceptual bistability has been investigated in its neural bases
44 (Kornmeier & Bach, 2012; Leopold & Logothetis, 1999), in the influence on it of the mechanisms of
45 priming (Goolkasian & Woodberry, 2010), in the effects of spatial context in which the images are
46 presented (Ouhana et al., 2017), in its occurrence in clinical cases (Allen & Chambers, 2011) and in
47 its significance for the neural and behavioural correlates of consciousness (Brancucci & Tommasi,
48 2011; for a general review see: Brascamp et al., 2018). Examples of ambiguous figures are the Rubin's
49 vase-face figure (Rubin, 1915), the Necker cube (Necker, 1832), the duck-rabbit figure (McManus et
50 al., 2010), and the old/young woman (Boring, 1930). A perhaps even wider literature has grown for
51 a companion paradigm of ambiguous figures, i.e. binocular rivalry, another method to obtain bistable
52 perception whereby two different figures are simultaneously presented one to the left and the other
53 to the right eye (Tong et al., 2006; Brascamp et al., 2015).

54 Bistable perception is particularly worthwhile for the research on consciousness (Brancucci et al.,
55 2011, 2016, 2018; Wang et al., 2013). The reason for this is that while the ambiguous figure is one
56 and does not change during its presentation, the associated conscious experience reverses from one
57 percept to the other. This allows researchers to ascribe measurable events which occur
58 simultaneously with the perceptual reversals to behavioural or neural events strictly related to
59 consciousness (Parkkonen et al., 2008; Sterzer et al., 2009). A more usual way to elicit different
60 percepts would be to present different images (e.g., a face followed by a tree, and so on). In this way,
61 however, behavioural or neural changes associated with the consequent perceptual changes cannot

62 be exclusively ascribed to processes related to consciousness as they could have been elicited simply
63 by the differences due to the physical inputs.

64 In a recent paper (Brancucci et al., 2020) we described the outcomes of a study in which we used the
65 divided visual field paradigm with vertical or horizontal division and the simultaneous presentation
66 of two identical ambiguous figures. Results showed that the temporal interdependence of the
67 reversals in the two hemifields was very low, and that, during in average 1/3 of the stimulation time,
68 subjects experienced simultaneously the two different interpretations of the same figure in the two
69 hemifields. The type of visual field division did not influence either frequency or temporal
70 interdependence of the reversals. Moreover, when one single ambiguous figure was presented, the
71 number of reversals was approximately the sum of the reversals observed with two simultaneously
72 presented figures.

73 Now, we move from these two main outcomes to further investigate the mechanisms of
74 consciousness hidden in bistable perception. Established that the perception of two ambiguous
75 figures presented simultaneously can reverse independently for each figure, presenting more
76 ambiguous figures at the same time the present study aims at drawing an input/output (I/O) relation
77 between the number of ambiguous figures simultaneously presented and the observed number of
78 perceptual reversals. The analysis of the relationship between the number of simultaneously
79 presented ambiguous figures and the number of observed perceptual reversal can shed light on the
80 nature of the mechanisms eliciting reversals and on the behavioural and neural correlates of
81 consciousness. An increase of the number of reversals associated to a higher number of figures
82 presented would suggest a strong dependence of perceptual bistability on field effects and possibly
83 the presence of multiple central mechanisms generating bistable conscious perception. Of note, the
84 notion of multiple central mechanisms is related to the topographical organization of the visual
85 system where the presentation of many (ambiguous) figures generates activity maps at different
86 levels, starting from the retina, in which the representation of each figure is spatially segregated from
87 the others. Conversely, a non-increased number of reversals would point to a strong role of top-down
88 attention in the generation of the reversals and on the presence of a limited number (possibly one)
89 of central mechanisms at the basis of bistable perception and possibly of consciousness. In fact, given
90 the intrinsic ability of ambiguous figures to generate perceptual bistability, it is reasonable that
91 increasing the number of ambiguous figures generates a scene in which reversals are facilitated, thus
92 inducing field effects. Conversely, a lack of increase of the number of reversals would suggest the

93 presence of top-down attentional filter mechanisms which act on the reversals and tend to reduce
94 them.

95 A further point which we face here is to investigate whether the perceived duration of a time lapse
96 can be influenced by the number of ambiguous figures presented and by their number of reversals.
97 Subjective evaluation of time duration is a fundamental point in psychology research and it has never
98 been studied in association to bistable perception, a situation in which subjective aspects of cognition
99 are of particular relevance. Thus we think that the investigation of the mechanisms underlying the
100 interactions between the two perception domains rests on solid epistemological foundations. In the
101 investigation of temporal processing, there are two main paradigms which may involve different
102 cognitive processes. In the attention-related prospective timing task, participants are informed in
103 advance that they will have to judge the duration of a period of time, for which attention plays a
104 primary role in performance. In contrast, in the memory-related retrospective timing task,
105 participants do not know until the end of the task that they will have to judge its duration, so memory
106 processes seem to be critical (see Vatakis et al., 2018). In general, evidence supports at least two
107 different systems for temporal processing: one based on senses for short durations ($\leq 1s$,
108 approximately) and one based on high-level cognitive processing for longer durations (Hellström &
109 Rammsayer, 2004; Lewis & Miall, 2003). The perception of short durations ($\leq 1s$, approximately) is
110 more influenced by the input sensory information (e.g., brightness, spatial frequencies, number of
111 elements comprising the stimulus, and neural adaptation; Xuan et al., 2007; Aaen-Stockdale et al.,
112 2011; Eagleman & Pariyadath, 2009). Instead, the judgments of duration on scales equal to or greater
113 than 3 seconds are influenced by cognitive load (Block et al., 2010). The present work uses an
114 attention-related prospective timing task and only considers durations of 3 minutes. Given that the
115 duration to be judged is long, we expect that the variation of the input sensory information (number
116 of figures presented simultaneously) will have limited effects on the judgment. On the contrary, we
117 predict that the number of perceived alternations may influence the temporal task. To our
118 knowledge, this is the first study linking this type of stimuli to temporal perception. However, it is
119 known that if during prospective temporal judgement tasks the subject performs another activity,
120 and attention is mainly focused on the non-temporal task, the estimate of elapsed time will be lower
121 than when attention is focused on the passage of time (Zakay & Block, 1996, 1997). In contrast, the
122 number of perceived reversals, when attention is paid to the bistable perception task, tends to
123 increase (Paffen et al., 2006). Consequently, we predict that the number of perceived reversals is

124 negatively correlated with duration judgments and that the direction of this correlation will remain
125 constant as the number of figures changes.

126 To join these goals, in the present study we presented 1, 2, 4 or 8 copies of the same ambiguous
127 figure (the Rubin's vase; Rubin, 1915) simultaneously, and asked subjects to press a key when they
128 experienced a perceptual reversal from the vase to the face profiles or viceversa. In Experiment 1
129 subjects had to press a specific key associated to each of the presented figures, whereas in
130 Experiment 2, to exclude an effect of the type of response on the reversal rate, participants had to
131 press a key regardless from the figure in which the reversal was perceived. In addition, to ascertain
132 that the participants looked at the fixation cross during the tasks, eye movements were recorded in
133 Experiment 2. At the end of each condition we asked subjects to estimate the duration of the session,
134 which lasted actually 3 minutes. A firm theoretical point on which this study bases is that perceptual
135 reversals elicited by ambiguous figures are pure changes in consciousness, which do not depend from
136 stimulus changes or from other "external" variables which change in synchrony with the reversals.
137 Although the literature shows that the context can influence the number of reversals (Intaitè et al.,
138 2013; Ouhnana & Kingdom, 2016), based on the above results (Brancucci et al., 2020) the main
139 hypothesis of the present piece of work is that, despite the different number of ambiguous figures
140 presented in the 4 experimental conditions, the number of perceptual reversals and the estimation
141 of time tends to remain constant.

142

143

144 **MATERIALS AND METHODS**

145 **Experiment 1**

146 *Participants*

147 Forty participants (22 females) aged between 19 and 30 years (mean age = 22.03, standard error =
148 0.38) took part in the study. The needed sample size was estimated with the G*power 3.1 software
149 (Faul et al., 2009) by performing an a priori analysis considering all results (average) of the sole
150 previously published work on the same topic (Brancucci et al., 2020). A number of 22 participants
151 was needed to achieve, with a power of .90 ($\alpha = .05$; correlation between measures = 0.68), a
152 medium/small effect size (Cohen's $f = 0.236$). Handedness scores measured by means of the
153 Edinburgh Handedness Inventory (Oldfield, 1971; according to which the handedness score ranges
154 from -100 = totally left handed, to +100 = totally right handed), showed that 37 subjects were right-
155 handed (i.e. score > 0) and 3 left-handed (mean 54.63 ± 7.22). Eyedness, measured with the same

156 test in which one item is dedicated to eye preference, showed that 15 subjects had left eye
157 preference, 6 subjects had no eye preference, and 19 subjects had right-eye preference (mean 0.17
158 \pm 0.23). Participants were enrolled if they did not complain of particular visual impairments and all of
159 them had normal or corrected-to-normal vision. The study was carried out in accordance to the
160 principles expressed in the Declaration of Helsinki of year 2013 and was approved by the local
161 research ethics committee.

162

163 *Stimuli*

164 The Rubin's face-vase figure (Rubin, 1915) was presented in 4 different conditions: single figure (1F),
165 2 figures (2F), 4 figures (4F), 8 figures (8F). In the 1F condition one Rubin's vase was presented at the
166 centre of the screen. In the 2F condition, 2 Rubin's vases were presented simultaneously, one in the
167 LVF and the other in the RVF. In the 4F condition, 4 Rubin's vases were presented simultaneously,
168 two in the RVF (one in the upper and one in the lower RVF) and two in the LVF (one in the upper and
169 one in the lower LVF). In the 8F condition, 8 Rubin's vases were presented simultaneously in a circle,
170 each at the same distance from the centre (Fig. 1).

171 The computer display was 34 cm wide and 27 cm high. The absolute dimension of the figures in the
172 1F and 2F conditions was 5 cm, in the 4F and 8F conditions it was 4 cm. In terms of visual angle, this
173 resulted in a difference of 0.8°. In the 2F, 4F, and 8F conditions the centre of each image was 6.5°
174 from the central fixation point. In each condition, stimuli were presented continuously for 3 min.

175

176 *Procedure*

177 The 4 conditions were presented in 4 blocks in a pseudorandom order counterbalanced across
178 participants, wherein the 8F condition was preceded by a training phase to familiarize the participants
179 with the answer buttons due to its greater difficulty. Subjects were tested in a quiet room and sat
180 comfortably in front of the computer monitor (approximately 70 cm from subject's head), with both
181 hands placed on the keyboard. During the experiment, participants were instructed and trained to
182 look at the centre of the screen and to not shift their gaze and their attention from the fixation cross.
183 In order to prepare participants to the experimental tasks, before starting they were familiarized with
184 a Rubin's face-vase figure. Participants were instructed to press a button when they perceived a
185 reversal in one of the presented images. In the 1F condition they were asked to press the spacebar
186 with their right hand to indicate a reversal perception in the image centrally displayed. In the 2F
187 condition participants had to press the "A" button with the left hand and the "L" button with the right

188 hand to indicate a reversal perception respectively in the LVF or RVF. In the 4F condition participants
189 had to press the “Z” or the “V” button with their left hand to indicate a reversal perception in the
190 lower LVF or lower RVF, respectively, and the “U” or the “P” button with their right hand to indicate
191 a reversal perception in the higher LVF or higher RVF, respectively. In the 8F condition, the “Z”, “X”,
192 “C”, “V” buttons were associated to the left hand, and the “P”, “O”, “I”, “U” buttons to the right hand,
193 and each button corresponded to one image of the Rubin’s vase (see Fig. 1, bottom). Before this
194 condition a specific training was performed in which particular care was devoted to the avoidance of
195 ocular movements. Participants were required to look at the centre of the screen and to press the
196 button corresponding to one of the 8 images which lighted on the monitor. The training sessions
197 were presented with increasing difficulty where in the first step the images lighted in sequence
198 following their circular arrangement, then in a spatially random way. The time limit for providing the
199 response decreased progressively (every 8 stimuli) from 5s to 3s. Participants were required to repeat
200 the training until 80% accuracy was achieved before they could start the experimental session for the
201 8F condition. No eye-tracking control was performed in Experiment 1. Participants were aware that
202 at the end of each condition they had to report the subjective evaluation of time duration and were
203 asked not to count during the conditions (Rattat & Droit-Volet, 2012). Participants wrote down their
204 answer on a sheet of paper on which the following sentence (here translated into English) was written
205 in Italian: "Indicate how much time has passed for each condition". Five out of 40 subjects did not
206 perform this task.

207 The experiment ran automatically using a software written in E-Prime (Psychology Software Tools
208 Inc., Pittsburgh, PA, United States) and participants’ responses and reaction times were registered by
209 the computer. The total duration of the experiment was approximately 20 min.

210

211 **Experiment 2**

212 *Participants*

213 Thirty-two participants (17 females) aged between 19 and 39 years (mean age = 25.33, standard error
214 = 0.87) recruited from the same population of university students as the previous sample (no subject
215 was recruited in both experiments) volunteered to take part in the study. The needed sample size
216 was estimated as in Experiment 1. Handedness and eye dominance were assessed as in Experiment
217 1 and showed that 31 subjects were right-handed (i.e. score > 0) and 1 left-handed (mean = 86.10,
218 st. dev. = 15.03) and that 12 subjects had left eye preference, 13 right eye preference, and 7 no eye
219 preference. Participants were enrolled with the same rules of Experiment 1. Seven of them were then

220 not included in the analyses due to problems related to the execution of the task (1 female) or to eye
221 movements (5 females, 1 male).

222

223 *Stimuli and procedure*

224 The Rubin's face-vase figure was presented in 4 different conditions (1F, 2F, 4F, 8F) as in Experiment
225 1. The absolute dimension of the figures was here 4 cm in all conditions and the centre of each figure
226 in the 2F, 4F, and 8F conditions was 6.5° from the central fixation point. Presentation conditions and
227 duration of stimuli were identical to those of Experiment 1. Subjects task was here to press the
228 spacebar with one hand each time they perceived a reversal in one of the presented figures
229 regardless from which figure it was. At the end of each session, subjects were requested to estimate
230 its duration.

231

232 *Eye-tracker recordings*

233 Participants were seated in front of a computer monitor (size: 1920 x 1080 pixels) approximately 70
234 cm away from their head as in Experiment 1. Eye movement data were collected with a Gazepoint
235 GP3 eye-tracker sampling at 60 Hz. OGAMA software (Voßkühler et al., 2008) was used to display the
236 stimuli and to record task responses and eye-tracker data. The eye tracking equipment was calibrated
237 using a grid of 5 points, 1 in the centre and 4 displaced in the 4 corners of the screen. During the
238 calibration, participants were required to gaze for 1 second at each dot. Throughout the experiment,
239 participants were asked to keep their gaze at the centre of the screen and to not shift it from the
240 fixation cross. For each session we obtained the spatial coordinates of each fixation and after the task
241 we analysed the percentage of total dwell time on the fixation region (a box of 4 cm around the
242 fixation cross). Then we removed 6 participants (1 male) whose gaze remained on the fixation region
243 for less than 75% in one of the conditions, and the statistical analyses were performed on subjects
244 who looked in the fixation region for an average time of 94.95% (standard error = 1.14).

245

246

247 **Results**

248 Statistical analyses were carried out with the software Statsoft Statistica 8.0. Shapiro-Wilk test for
249 normality distribution showed that the analysed variables met not normality criteria ($p < 0.01$ for all
250 variables). Since the analysis of variance (ANOVA) procedure is robust enough to tolerate violations
251 of the normality assumption (Pituch & Stevens, 2015), we preferred to avoid data transformations

252 that could bias the results. Preliminary statistical analyses indicated that the order of condition
253 administration (whether participants first received the 1F, 2F, 4F or 8F), did not influence the
254 bistability scores. This variable was therefore not included in the subsequent analyses.

255

256 **Experiment 1**

257 *Analyses on bistability scores*

258 We define as 'bistability score' the total number of reversals observed in the 180 s presentation of
259 the ambiguous figures. Across subjects (mean and standard deviation) bistability scores in the 4
260 conditions (1F, 2F, 4F, 8F) were 39.2 ± 29.8 for the 1F condition, 43.1 ± 31.1 for the 2F condition, 51.0
261 ± 42.0 for the 4F condition, and 51.1 ± 49.5 for the 8F condition (Fig. 2).

262 A repeated measures multivariate ANOVA with bistability score as a dependent variable, Handedness
263 and Eyedness as continuous predictors, and one within-subjects factor (Condition, with 4 levels 1F,
264 2F, 4F, 8F) was carried out. Results showed no significant effects for all factors: Handedness ($F=0.88$,
265 $p=0.768$, $\eta^2=0.003$), Eyedness ($F=0.69$, $p=0.412$, $\eta^2=0.022$), Condition ($F=0.78$, $p=0.507$, $\eta^2=0.025$)
266 and their interactions: Condition x Handedness ($F=0.50$, $p=0.687$, $\eta^2=0.016$), Condition x Eyedness
267 ($F=1.20$, $p=0.315$, $\eta^2=0.037$). This result indicates that the number of reversals when 1, 2, 4, or 8
268 ambiguous figures are simultaneously presented does not change statistically.

269 We performed a similar analysis excluding the condition 1F to control for a possible confound of the
270 fixation point which coincides with the figure only in that condition. Results showed no significant
271 effects for all factors: Handedness ($F=0.31$, $p=0.862$, $\eta^2<0.001$), Eyedness ($F=0.87$, $p=0.359$,
272 $\eta^2=0.027$), Condition ($F=0.05$, $p=0.951$, $\eta^2=0.002$) and their interactions: Condition x Handedness
273 ($F=0.80$, $p=0.453$, $\eta^2=0.025$), Condition x Eyedness ($F=1.75$, $p=0.183$, $\eta^2=0.053$). This results indicates
274 that the fixation point does not statistically influence the lack of reversals increment.

275

276 *Effects of figures positions on the screen*

277 Subsequently, analyses were carried out within the conditions with multiple figures presentation (2F,
278 4F, 8F) to investigate whether the position of the figure in the screen could exert some effect on the
279 reversals. In the 2F condition, one-way repeated measures ANOVA with the factor Side (LVF, RVF)
280 showed a main effect ($F=4.756$, $p=0.03$, $\eta^2=0.11$) with more perceptual reversals occurring in the
281 LVF. In the 4F condition, two-ways repeated measures ANOVA with Side (LVF, RVF) and Height (UVF,
282 WVF) as factors showed a significant interaction Side x Height ($F=8.896$, $p=0.005$, $\eta^2=0.186$) and no
283 other statistically significant effects (Side: $F=0.516$, $p=0.477$, $\eta^2=0.013$; Height: $F=2.136$, $p=0.152$,

284 $\eta^2=0.052$). Tukey post-hoc comparisons showed that concerning the LVF more reversals occurred in
285 the left WVF compared to the left UVF ($p=0.004$) and that concerning the UVF more reversals
286 occurred in the upper RVF compared to the upper LVF ($p=0.029$). Number of reversals did not differ
287 between right WVF compared to the right UVF ($p=0.946$) and between the lower RVF compared to
288 the lower LVF ($p=0.568$). In the 8F condition, two-ways repeated measures ANOVA with Side (LVF,
289 RVF) and Height (UVF, WVF) as factors showed no significant effects (Side: $F=2.546$, $p=0.119$,
290 $\eta^2=0.061$; Height: $F=0.115$, $p=0.736$, $\eta^2=0.003$; Side x Height: $F=3.756$, $p=0.060$, $\eta^2=0.088$). Factor
291 levels have been here calculated considering the sum of the 3 positions corresponding to each
292 quadrant in the visual field (e.g. the left-up quadrant was considered as the sum of the reversals in 3
293 positions corresponding to the letters Z, U, I in Fig. 1). Of note, a very similar result has been observed
294 in the analogous analysis considering only the 4 positions corresponding to the letters U, O, X, V in
295 Fig. 1 (specifically, the interaction was here identical, which is obvious from a statistical point of view).

296

297 **Experiment 2**

298 *Analyses on bistability scores*

299 Analyses were carried out after having excluded 1 participant since she did not perform the task
300 correctly and 6 subjects since they made too large eye movements. The sample of the following
301 analyses is thus $n = 25$, all subjects who gazed for at least 75% of the time in each condition to the
302 fixation cross (mean = 94.95% of the total test time, standard error = 1.14). Across subjects (mean
303 and standard deviation) bistability scores in the 4 conditions (1F, 2F, 4F, 8F) were 51.8 ± 34.8 for the
304 1F condition, 44.5 ± 33.6 for the 2F condition, 39.7 ± 22.3 for the 4F condition, and 46.3 ± 22.3 for
305 the 8F condition (Fig. 2).

306 A repeated measures multivariate ANOVA with bistability score as a dependent variable, Handedness
307 and Eyedness as continuous predictors, and one factor (Condition, with 4 levels 1F, 2F, 4F, 8F) was
308 carried out. Results showed no significant effects for all factors: Handedness ($F=2.04$, $p=0.167$,
309 $\eta^2=0.084$), Eyedness ($F=0.82$, $p=0.376$, $\eta^2=0.036$), Condition ($F=0.01$, $p=0.998$, $\eta^2=0.001$) and their
310 interactions: Condition x Handedness ($F=0.31$, $p=0.817$, $\eta^2=0.014$), Condition x Eyedness ($F=0.167$,
311 $p=0.918$, $\eta^2=0.008$) confirming the result of Experiment 1.

312 Finally, we performed a similar analysis excluding the condition 1F to control for a possible confound
313 of the fixation point which coincides with the figure only in that condition. Results showed again no
314 significant effects for all factors: Handedness ($F=1.78$, $p=0.196$, $\eta^2=0.075$), Eyedness ($F=0.71$,

315 $p=0.407$, $\eta^2=0.031$), Condition ($F=0.02$, $p=0.981$, $\eta^2=0.001$) and their interactions: Condition x
316 Handedness ($F=0.28$, $p=0.752$, $\eta^2=0.013$), Condition x Eyedness ($F=0.19$, $p=0.825$, $\eta^2=0.009$).

317

318 **Time estimation**

319 For both experiments, an analysis was carried out on the perceived duration of the 4 sessions
320 (conditions), which lasted actually 3 min. For Experiment 1, mean and standard deviation of the
321 perceived duration was 3.28 ± 1.36 min in the 1F condition, 2.73 ± 1.37 min in the 2F condition, 3.15
322 ± 1.83 min in the 4F condition, and 3.04 ± 1.45 min in the 8F condition. For Experiment 2, mean and
323 standard deviation of the perceived duration was 2.59 ± 1.35 min in the 1F condition, 2.56 ± 1.72 min
324 in the 2F condition, 2.34 ± 1.21 min in the 4F condition, and 2.25 ± 1.29 min in the 8F condition (Fig.
325 2). Repeated-measures ANOVA on the perceived duration (1F, 2F, 4F, 8F) showed no significant
326 effects (Experiment 1: $F=2.235$, $p=0.089$, $\eta^2=0.06$; Experiment 2: $F=0.846$, $p=0.473$, $\eta^2=0.03$).

327 Correlation analysis on the data of both experiments together ($n=60$; partial correlations controlling
328 for the experiment) between the bistability score and the perceived duration indicated that all
329 correlations were negative and ranged from $r = -0.14$ to $r = -0.44$. Specifically, in the single conditions
330 they were $r = -0.24$, $p = 0.070$ (1F); $r = -0.17$, $p = 0.192$ (2F); $r = -0.14$, $p = 0.278$ (4F); $r = -0.44$, $p <$
331 0.001 (8F) being this last correlation significant after Bonferroni correction for 4 comparisons. The
332 global partial correlation computed between the mean stability scores and the mean perceived
333 duration in the 4 conditions was also significant: $r = -0.30$, $p = 0.022$ (Fig. 3).

334

335 **Control analyses**

336 A further analysis was performed on all subjects of both experiments ($n=60$) to control whether the
337 number of reversals in the different subjects was consistent across conditions. To this aim, Pearson's
338 r correlation coefficients were pairwise calculated between the 4 conditions (1F, 2F, 4F, 8F). Results
339 showed correlation coefficients ranging from $r = 0.50$ to $r = 0.77$ with p always < 0.001 . This result
340 indicates that subjects who saw more reversals did this in all conditions and vice versa, and is in
341 accordance with previous similar evidence in the literature (Cao et al., 2018).

342 In the subsample of Experiment 2 in which we recorded eye movements ($n=25$), we made a further
343 control analysis across conditions to see whether the tendency of maintain the gaze correctly on the
344 fixation point was related to the number of perceived reversals. In such an analysis we correlated %
345 correct fixation time to the number of reversals and observed no relation ($r = -0.027$, $p = 0.897$).

346

347

348

349 Discussion

350 The present study was primarily aimed at finding a relation between the number of ambiguous
351 figures simultaneously presented and the perceptual reversals experienced by the subjects. Results
352 showed that changing the number of figures does not significantly alter the number of reversals
353 observed. The I/O curve (Fig. 2) shows that despite an increase of the number of simultaneously
354 presented figures, the number of reversals tends to remain constant. Specifically, since with one
355 ambiguous figure (1F condition) we observed about 40 reversals in 3 minutes, with 8 figures one
356 would have expected to observe about 320 reversals in the same time lapse, instead only about 50
357 were observed. As explained in the Introduction, a lack of increase of the number of reversals despite
358 an increase of the number of presented figures suggests a major role of top-down attention. Top-
359 down attention, as a general, non-specific, and content-limited mechanism, does not act specifically
360 on each representation of the figures remaining independent of their number, and its limited capacity
361 is possibly a cause of the decrease of the ratio reversals/number of figures. A further key role in the
362 route to consciousness is possibly played by central mechanisms related to the identification of the
363 stimulus. The lack of increase of the number of reversals when more figures are presented should
364 exclude that the underlying mechanisms operate at the level of object identification or slightly after
365 it ("object-token"; Zimmer and Ecker, 2010) as there each reversal mechanism would operate
366 independently for different objects producing instead an increase of reversals. These observations
367 speak in favor of a limited number (possibly one) of central mechanisms at the basis of bistable
368 perception and of consciousness.

369 A further aim of the present study was to investigate whether the number of ambiguous figures
370 simultaneously presented and their perceptual reversals could influence the judgement of the
371 duration of a time lapse. No differences were found between the time duration evaluations of the
372 four sessions, but an interesting and seemingly robust result emerged from the analysis of
373 correlations with the bistability score: all correlations (significant and non-significant) were negative,
374 indicating that the more reversals were perceived, the shorter the duration of the session was
375 evaluated. The most evident result was obtained in the 8F condition which showed the highest (and
376 significant) correlation value (-0.44). The overall correlations between the mean of the time
377 evaluations and the mean of the reversals in the 4 conditions showed a similar (-0.30) significant
378 correlation. In our opinion, this finding suggests the presence of a link between the perceptual

379 reversals and the internal clock which can be used to estimate time flow. If between two reversals
380 the time lapse is long (as is the case with a low number of reversals), then the elapsed time evaluation
381 tends to increase.

382 Several further results which go beyond the main scope of the present study were observed, which
383 we think are worth to be discussed. We detected more perceptual reversals in the LVF than in the
384 RVF when two ambiguous figures were presented simultaneously in the two lateral visual hemifields,
385 a condition (2F) in which the classical visual hemifield-paradigm stimulation mode was employed.
386 This result points to a major implication of the right hemisphere in the genesis of the perceptual
387 reversals, as the LVF projects mainly to the right visual cortex. A first explanation of this result is
388 possibly related to the right hemispheric parietal specialization for attention. It is known that
389 attention shows a bias towards the left in healthy subjects (pseudoneglect, Nicholls et al., 2017) and
390 the number of reversals during bistable perception has been shown to be directly proportional to
391 attentional resources (Paffen et al., 2006). A further explanation lies in the right hemispheric
392 superiority in the processing of faces (Hasson et al., 2001; Prete et al., 2015; Sergent & Bindra, 1981),
393 one of the two possible interpretations of the Rubin's vase, and on the fact that the ambiguous figure
394 presented is inherently a non-verbal stimulus. For presently unknown reasons, this superiority would
395 produce more reversals between the two interpretation of the Rubin's vase as if competition
396 between preferred stimuli would produce less perceptual stability compared to non-preferred
397 stimuli. Further research is needed to elucidate this interesting issue.

398 In the 4F condition, a more complex result was produced by an interaction between the UVF-WVF
399 and the LVF-RVF, or in other words between the concurrent horizontal and vertical divisions of the
400 visual field. Considering the left part of the visual field, more reversals occurred in the lower
401 compared to the upper quadrant, and considering the upper part of the visual field, more reversals
402 occurred in the right compared to the left quadrant. This complex result is in accordance with a
403 previous finding which shows that only the LVF has different responses between its upper and lower
404 parts (Lee et al., 2009), but it contrasts with other reports which found a right hemisphere
405 specialization for the UVF in addition to that for the LVF (Thomas et al., 2015; D'Anselmo et al., 2018).
406 More specific studies on up-low (UVF-WVF) asymmetries in visual attention suggest that these are
407 elusive or strictly dependent on the specific task requested (Thomas & Elias, 2011).

408 In the history of behavioural studies with ambiguous figures, several authors have presented two or
409 more ambiguous images more or less simultaneously (Adams & Haire, 1959; Babich & Standing, 1981;
410 Jensen & Mathewson, 2011; Long & Toppino, 1981; Mathewson, 2018). However, in most of these

411 studies the presentation techniques used were underdeveloped and the instructions given to the
412 subjects did not allow, in our opinion, sufficient control on the afferent information flow occurring
413 during the stimulation. Also the present study has potential confounds related to the lack of eye
414 movements control and the different size of the figures presented in Experiment 1 (5 cm in the 1F
415 and 2F conditions compared to 4 cm in the 4F and 8F conditions). Experiment 2 was however
416 designed to solve these issues by presenting all figures with the same size and by controlling eye
417 movements with an eye tracker instrument. In Experiment 2 we also modified the response buttons:
418 unlike Experiment 1 in which participants were asked to associate each key with a figure, in
419 Experiment 2 they had to press the same button for each perceived reversal. Hence, the results of
420 Experiment 2 allow us to exclude that the reversal rate in Experiment 1 could be influenced by the
421 difficulty due to associate each image to one specific key. Another confound concerns the fixation
422 point, which coincided with the figure only in the 1F condition raising possible different attentional
423 effects compared to the other conditions which could have biased the results. A dedicated control
424 analysis showed that the lack of reversals increase was evident also in the 3 conditions in which the
425 distance of the figures to the fixation point was identical.

426 The present results are generally in line with the studies cited at the beginning of the previous
427 paragraph and together with them show important consequences for the interpretation of the
428 experimental results obtained so far with ambiguous figures. As anticipated in the Introduction, the
429 main result of the study has some cues in the literature, in particular in the research studying the
430 relations between consciousness and attention, two aspects of cognition which are very closely
431 related, yet different (Koch & Tsuchiya, 2007). Paffen and coworkers (2006) demonstrated that
432 distracting focal attention during bistable perception slows down the number of reversals per time
433 unit. They showed that shifting attention from binocular rivalry stimuli to a simultaneously presented
434 motion-detection task reduces the rate of rivalry alternations. It appears that rivalry dynamics
435 depend on the amount of attentional resources allocated to the rival stimuli. When this amount is
436 reduced by increasing the difficulty of a concurrent task, the rivalry reversal rate slows further. This
437 drop in alternation rate is not attributable to a degraded ability to track rivalry alternations while
438 performing the detection task since under pseudo-rivalry conditions mimicking real rivalry
439 alternations, observers reliably tracked stimulus alternations. In our experiments, actually, presenting
440 more than one ambiguous figure at one time has led to a reduced amount of attention that could be
441 allocated to one single figure. This mechanism could explain why the number of reversals observed
442 does not increase with the number of simultaneous ambiguous figures presented and is in tight

443 agreement with the notion that attention allocation increases the number of reversals during bistable
444 perception.

445 Concerning the issue of time evaluation, as a limitation of the present study the number of reversals
446 was always associated to the number of the motor responses. In principle, this would not allow to
447 establish a relation between the perceptual reversals and time evaluation, as the same relation holds
448 between the number of motor responses and time evaluation. However, extant literature suggests
449 that the influence of motor activity on time perception is at best limited to the evaluation of short
450 durations in the sub-second domain (Mioni et al., 2016; Hass et al., 2012; Lewis & Miall, 2003). Here
451 the durations were one order bigger (minutes) and their evaluations are more related to cognitive
452 factors such as attention and working memory. Evidence in the literature suggests in fact the
453 existence of at least two different functional and neural mechanisms for temporal perception
454 (Hellström & Rammsayer, 2004; Lewis & Miall, 2003). The processing of smaller time intervals
455 (approximately $\leq 1s$) is sensory-based, whereas the processing of longer intervals requires the
456 support of cognitive resources. In this study, varying sensory information (understood as the number
457 of figures) did not produce a significant effect on perceived duration but it was related to the number
458 of perceived alternations. This would seem to confirm that the perception of longer durations is
459 based more on high-level processing than on sensory processing. Thus we expect a negligible
460 influence of the motor response on time evaluation in the present study. Nevertheless, further
461 studies could disentangle this issue for instance using no-report paradigms during bistable perception
462 which do not request a motor response. Although perceived duration is generally assumed to
463 correspond with objective duration, several studies suggest that time perception cannot be placed in
464 a simpleminded framework (Buetti & Walsh, 2009). Previous studies suggested several rules that
465 would govern temporal perception. It has been proposed that the experience of duration is a
466 signature of the amount of energy expended in processing the stimulus (Eagleman & Pariyadath,
467 2009) and that perceived time is positively related to perceptual vividness and to the ease of
468 extracting information from the stimulus (Matthews & Meck, 2016). Stimulus repetition has been
469 shown to reduce temporal estimation possibly due to a suppression of neural sustained responses
470 (Eagleman & Pariyadath, 2009). This outcome seems to agree with our result of a negative
471 relationship between number of spontaneous reversals and evaluation of the session duration.
472 Consistent with this interpretation, adaptation has often been considered an underlying dynamic of
473 multistable perception (Long & Toppino, 2004; Kogo et al., 2015) and would seem to explain at least
474 a small part of the variability in perceptual reversals (Pastukhov & Braun, 2011). In particular, as a

475 consequence of adaptation, reversals tend to be more frequent when an ambiguous figure is
476 continuously displayed for a few minutes. Theoretically, both low-level neural mechanisms and high-
477 level cognitive processes (as clarified in the next paragraph) can provide valid interpretations for the
478 observed effect on temporal judgment (Intaite et al., 2013; Meng & Tong, 2004; Toppino, 2003), and
479 we do not rule out that either explanation can have some bases. Resolving this issue would require
480 independent modulation of attentional and adaptation mechanisms. Future studies could resolve
481 such theoretical questions in the field of temporal perception. Intuitively, the underestimation of the
482 temporal interval might be related to a common experience, namely that in some situations time
483 seems to pass more quickly (or slowly). In truth, subjective judgments about the speed of time seem
484 to be dissociated from those of interval length estimation (e.g. Droit-Volet & Wearden, 2016; Deinzer
485 et al., 2017; Thönes & Oberfeld, 2015; Wearden, 2015); however, there are data demonstrating a
486 correlation between the two measures (Sucala et al., 2010). The paradigm used here can be easily
487 adapted to investigate these processes and whether they are influenced by the same factors.

488 Our result of a negative relationship between the number of spontaneous reversals and evaluation
489 of the session duration seems consistent with the Attentional Gate Model (Zakay & Block, 1996,
490 1997). In prospective temporal judgement tasks (when the subject is aware from the beginning that
491 he/she will have to report the duration of the event) the accuracy of the estimate is influenced by
492 the degree of attention paid to the task. If during the temporal estimation task the subject performs
493 another activity, i.e. bistable perception, and the attention is mainly placed on the non-temporal task,
494 the estimate of elapsed time will tend to have negative values. Furthermore, as already reported
495 above, focusing attention on bistable figures increases the perceived reversals. Thus, by integrating
496 the AGM theory with the effects of attention in bistable perception tasks, it follows that an increase
497 in the number of perceived reversals is negatively correlated with perceived duration. On a practical
498 level, this negative correlation could be used as an index of the subject's ability to direct attention to
499 a main task. The highest correlation was observed in the 8F condition. This probably occurred
500 because the increase in the number of figures increased the difficulty of the task and interfered more
501 with the evaluation of time. We hypothesise that this more negative correlation cannot be explained
502 solely in terms of how much attentional resources were allocated to the two tasks. In fact, a
503 higher/lower demand for attentional resources to the bistable perception task would also translate
504 into an increase/decrease in the number of perceived reversals (Paffen et al., 2006), but this was not
505 observed. This, instead, would seem to be related to the cognitive load required by the bistable
506 perception task. Indeed, increasing cognitive load in prospective paradigms leads to a decrease in

507 duration judgment. As in Block and coworkers (2010), the term cognitive load refers to the required
508 amount of information processed by attention and working memory. Prospective judgments on
509 duration can be thus used as a measure of the amount of mental load required to perform a non-
510 temporal task (Zakay et al., 1999; Block et al., 2010).

511 The relation investigated here and the associated I/O curve should not be confounded with the
512 classical relations studied by psychophysics. Psychophysics quantitatively investigates the
513 relationship between physical stimuli and the sensations they produce, studying the effect of
514 systematically varying the properties of a stimulus along one or more physical dimensions
515 (Gescheider, 1997). That is, psychophysics bases its roots on situations in which the stimulus is always
516 changed experimentally and the effects of the changes are measured from behaviour. In the present
517 study instead, the pivotal point is that the measurements start from situations in which the stimulus
518 is always constant (the ambiguous figure does not change) but it produces changes in perception,
519 which can be thus assumed as changes in consciousness (O'Regan & Noë, 2001) – an assumption that
520 cannot be made in the experiments of psychophysics.

521

522

523 **Data availability**

524 Data are available in the Open Science Framework website at URL: <https://osf.io/98scp/> .

525

526

527 **References**

- 528 Aaen-Stockdale, C., Hotchkiss, J., Heron, J., & Whitaker, D. (2011). Perceived time is spatial frequency
529 dependent. *Vision research*, 51(11), 1232–1238. <https://doi.org/10.1016/j.visres.2011.03.019>.
- 530 Adams, P., & Haire, M. (1959). The effect of orientation on the reversal of one cube inscribed in another. *The*
531 *American Journal of Psychology*, 72, 296–299. <https://doi.org/10.2307/1419384>.
- 532 Allen, M. L., & Chambers, A. (2011). Implicit and explicit understanding of ambiguous figures by adolescents
533 with autism spectrum disorder. *Autism: the international journal of research and practice*, 15(4), 457–472.
534 <https://doi.org/10.1177/1362361310393364>.
- 535 Babich, S., & Standing, L. (1981). Satiation effects with reversible figures. *Perceptual and motor skills*, 52(1),
536 203–210. <https://doi.org/10.2466/pms.1981.52.1.203>.
- 537 Balduzzi, D., & Tononi, G. (2009). Qualia: the geometry of integrated information. *PLoS computational biology*,
538 5(8), e1000462. <https://doi.org/10.1371/journal.pcbi.1000462>.
- 539 Block, R. A., Hancock, P. A., & Zakay, D. (2010). How cognitive load affects duration judgments: A meta-analytic
540 review. *Acta psychologica*, 134(3), 330–343. <https://doi.org/10.1016/j.actpsy.2010.03.006>.
- 541 Boring, E. G. (1930). A new ambiguous figure. *The American Journal of Psychology*, 42, 444–445.
542 <https://doi.org/10.2307/1415447>.

- 543 Brancucci, A., & Tommasi, L. (2011). "Binaural rivalry": dichotic listening as a tool for the investigation of the
544 neural correlate of consciousness. *Brain and cognition*, 76(2), 218–224.
545 <https://doi.org/10.1016/j.bandc.2011.02.007>.
- 546 Brancucci, A., D'Anselmo, A., Pasciucchio, M. R., & San Martini, P. (2020). Independent perceptual reversals for
547 simultaneously presented ambiguous figures. *Consciousness and cognition*, 81, 102928.
548 <https://doi.org/10.1016/j.concog.2020.102928>.
- 549 Brancucci, A., Franciotti, R., D'Anselmo, A., Della Penna, S., & Tommasi, L. (2011). The sound of consciousness:
550 neural underpinnings of auditory perception. *The Journal of neuroscience : the official journal of the Society*
551 *for Neuroscience*, 31(46), 16611–16618. <https://doi.org/10.1523/JNEUROSCI.3949-11.2011>.
- 552 Brancucci, A., Lugli, V., Perrucci, M. G., Del Gratta, C., & Tommasi, L. (2016). A frontal but not parietal neural
553 correlate of auditory consciousness. *Brain structure & function*, 221(1), 463–472.
554 <https://doi.org/10.1007/s00429-014-0918-2>.
- 555 Brancucci, A., Padulo, C., Franciotti, R., Tommasi, L., & Della Penna, S. (2018). Involvement of ordinary what
556 and where auditory cortical areas during illusory perception. *Brain structure & function*, 223(2), 965–979.
557 <https://doi.org/10.1007/s00429-017-1538-4>.
- 558 Brascamp, J., Sterzer, P., Blake, R., & Knapen, T. (2018). Multistable Perception and the Role of the
559 Frontoparietal Cortex in Perceptual Inference. *Annual review of psychology*, 69, 77–103.
560 <https://doi.org/10.1146/annurev-psych-010417-085944>.
- 561 Brascamp, J. W., Klink, P. C., & Levelt, W. J. (2015). The 'laws' of binocular rivalry: 50 years of Levelt's
562 propositions. *Vision research*, 109(Pt A), 20–37. <https://doi.org/10.1016/j.visres.2015.02.019>.
- 563 Bueti, D., & Walsh, V. (2009). The parietal cortex and the representation of time, space, number and other
564 magnitudes. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*,
565 364(1525), 1831–1840. <https://doi.org/10.1098/rstb.2009.0028>.
- 566 Cao, T., Wang, L., Sun, Z., Engel, S. A., & He, S. (2018). The Independent and Shared Mechanisms of Intrinsic
567 Brain Dynamics: Insights From Bistable Perception. *Frontiers in psychology*, 9, 589.
568 <https://doi.org/10.3389/fpsyg.2018.00589>.
- 569 D'Anselmo, A., Giuliani, F., Campopiano, F., Carta, E., & Brancucci, A. (2018). Hemispheric asymmetries in
570 setticlavio reading. *Neuropsychology*, 32(3), 337–343. <https://doi.org/10.1037/neu0000430>.
- 571 De Graaf, T. A., Hsieh, P. J., & Sack, A. T. (2012). The 'correlates' in neural correlates of consciousness.
572 *Neuroscience and biobehavioral reviews*, 36(1), 191–197.
573 <https://doi.org/10.1016/j.neubiorev.2011.05.012>.
- 574 Dehaene, S., & Naccache, L. (2001). Towards a cognitive neuroscience of consciousness: basic evidence and a
575 workspace framework. *Cognition*, 79(1-2), 1–37. [https://doi.org/10.1016/s0010-0277\(00\)00123-2](https://doi.org/10.1016/s0010-0277(00)00123-2).
- 576 Deinzer, V., Clancy, L., & Wittmann, M. (2017). The Sense of Time While Watching a Dance Performance. *SAGE*
577 *Open*. <https://doi.org/10.1177/2158244017745576>.
- 578 Droit-Volet, S., & Wearden, J. (2016). Passage of Time Judgments Are Not Duration Judgments: Evidence from
579 a Study Using Experience Sampling Methodology. *Frontiers in psychology*, 7, 176.
580 <https://doi.org/10.3389/fpsyg.2016.00176>.
- 581 Eagleman, D. M., & Pariyadath, V. (2009). Is subjective duration a signature of coding efficiency?. *Philosophical*
582 *transactions of the Royal Society of London. Series B, Biological sciences*, 364(1525), 1841–1851.
583 <https://doi.org/10.1098/rstb.2009.0026>.
- 584 Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using G*Power 3.1: tests for
585 correlation and regression analyses. *Behavior research methods*, 41(4), 1149–1160.
586 <https://doi.org/10.3758/BRM.41.4.1149>.

- 587 Gescheider, G. A. (1997). *Psychophysics: The fundamentals*. (3rd ed.) Routledge.
588 <https://doi.org/10.4324/9780203774458>.
- 589 Goolkasian, P., & Woodberry, C. (2010). Priming effects with ambiguous figures. *Attention, perception &*
590 *psychophysics*, 72(1), 168–178. <https://doi.org/10.3758/APP.72.1.168>.
- 591 Hass, J., Blaschke, S., & Herrmann, J. M. (2012). Cross-modal distortion of time perception: demerging the
592 effects of observed and performed motion. *PloS one*, 7(6), e38092.
593 <https://doi.org/10.1371/journal.pone.0038092>.
- 594 Hasson, U., Hendler, T., Ben Bashat, D., & Malach, R. (2001). Vase or face? A neural correlate of shape-selective
595 grouping processes in the human brain. *Journal of cognitive neuroscience*, 13(6), 744–753.
596 <https://doi.org/10.1162/08989290152541412>.
- 597 Hellström, A., & Rammsayer, T. H. (2004). Effects of time-order, interstimulus interval, and feedback in
598 duration discrimination of noise bursts in the 50- and 1000-ms ranges. *Acta psychologica*, 116(1), 1–20.
599 <https://doi.org/10.1016/j.actpsy.2003.11.003>.
- 600 Intaitė M., Šoliūnas A., Gurčienė O., & Rukšėnas O. (2013). Effect of Bias on the Perception of Two
601 Simultaneously Presented Ambiguous Figures. *Psichologija*, 47, 91-101.
602 <https://doi.org/10.15388/Psichol.2013.47.1403>.
- 603 Jensen, M. S., & Mathewson, K. E. (2011). Simultaneous Perception of Both Interpretations of Ambiguous
604 Figures. *Perception*, 40(8), 1009–1011. <https://doi.org/10.1068/p6880>.
- 605 Koch, C., Massimini, M., Boly, M., & Tononi, G. (2016). Neural correlates of consciousness: progress and
606 problems. *Nature reviews. Neuroscience*, 17(5), 307–321. <https://doi.org/10.1038/nrn.2016.22>.
- 607 Koch, C., & Tsuchiya, N. (2007). Attention and consciousness: two distinct brain processes. *Trends in cognitive*
608 *sciences*, 11(1), 16–22. <https://doi.org/10.1016/j.tics.2006.10.012>.
- 609 Kogo, N., Hermans, L., Stuer, D., van Ee, R., & Wagemans, J. (2015). Temporal dynamics of different cases of
610 bi-stable figure-ground perception. *Vision research*, 106, 7–19.
611 <https://doi.org/10.1016/j.visres.2014.10.029>.
- 612 Kornmeier, J., & Bach, M. (2012). Ambiguous figures - what happens in the brain when perception changes
613 but not the stimulus. *Frontiers in human neuroscience*, 6, 51. <https://doi.org/10.3389/fnhum.2012.00051>.
- 614 Lee, B., Kaneoke, Y., Kakigi, R., & Sakai, Y. (2009). Human brain response to visual stimulus between
615 lower/upper visual fields and cerebral hemispheres. *International journal of psychophysiology : official*
616 *journal of the International Organization of Psychophysiology*, 74(2), 81–87.
617 <https://doi.org/10.1016/j.ijpsycho.2009.07.005>.
- 618 Leopold, D. A., & Logothetis, N. K. (1999). Multistable phenomena: changing views in perception. *Trends in*
619 *cognitive sciences*, 3(7), 254–264. [https://doi.org/10.1016/s1364-6613\(99\)01332-7](https://doi.org/10.1016/s1364-6613(99)01332-7).
- 620 Lewis, P. A., & Miall, R. C. (2003). Distinct systems for automatic and cognitively controlled time measurement:
621 evidence from neuroimaging. *Current opinion in neurobiology*, 13(2), 250–255.
622 [https://doi.org/10.1016/s0959-4388\(03\)00036-9](https://doi.org/10.1016/s0959-4388(03)00036-9).
- 623 Long, G. M., & Toppino, T. C. (2004). Enduring interest in perceptual ambiguity: alternating views of reversible
624 figures. *Psychological bulletin*, 130(5), 748–768. <https://doi.org/10.1037/0033-2909.130.5.748>.
- 625 Long, G. M., & Toppino, T. C. (1981). Multiple representations of the same reversible figure: implications for
626 cognitive decisional interpretations. *Perception*, 10(2), 231–234. <https://doi.org/10.1068/p100231>.
- 627 Mathewson K. E. (2018). Duck Eats Rabbit: Exactly Which Type of Relational Phrase Can Disambiguate the
628 Perception of Identical Side by Side Ambiguous Figures?. *Perception*, 47(4), 466–469.
629 <https://doi.org/10.1177/0301006618756810>.
- 630 Matthews, W. J., & Meck, W. H. (2016). Temporal cognition: Connecting subjective time to perception,
631 attention, and memory. *Psychological bulletin*, 142(8), 865–907. <https://doi.org/10.1037/bul0000045>.

- 632 McManus, I. C., Freegard, M., Moore, J., & Rawles, R. (2010). Science in the Making: Right Hand, Left Hand. II:
633 The duck-rabbit figure. *Laterality*, 15(1-2), 166–185. <https://doi.org/10.1080/13576500802564266>.
- 634 Meng, M., & Tong, F. (2004). Can attention selectively bias bistable perception? Differences between binocular
635 rivalry and ambiguous figures. *Journal of vision*, 4(7), 539–551. <https://doi.org/10.1167/4.7.2>.
- 636 Mioni, G., Grassi, M., Tarantino, V., Stablum, F., Grondin, S., & Bisiacchi, P. S. (2016). The impact of a concurrent
637 motor task on auditory and visual temporal discrimination tasks. *Attention, perception & psychophysics*,
638 78(3), 742–748. <https://doi.org/10.3758/s13414-016-1082-y>.
- 639 Necker LALXI. 1832 Observations on some remarkable optical phænomena seen in Switzerland; and on an
640 optical phænomenon which occurs on viewing a figure of a crystal or geometrical solid. Lond. Edinb. Dublin
641 Philos. Mag. J. Sci. 1, 329–337. doi:10.1080/ 14786443208647909.
- 642 Nicholls, M. E., Hobson, A., Petty, J., Churches, O., & Thomas, N. A. (2017). The effect of cerebral asymmetries
643 and eye scanning on pseudoneglect for a visual search task. *Brain and cognition*, 111, 134–143.
644 <https://doi.org/10.1016/j.bandc.2016.11.006>.
- 645 O'Regan, J. K., & Noë, A. (2001). A sensorimotor account of vision and visual consciousness. *The Behavioral
646 and brain sciences*, 24(5), 939–1031. <https://doi.org/10.1017/s0140525x01000115>.
- 647 Oldfield R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia*,
648 9(1), 97–113. [https://doi.org/10.1016/0028-3932\(71\)90067-4](https://doi.org/10.1016/0028-3932(71)90067-4).
- 649 Ouhnana, M., Jennings, B. J., & Kingdom, F. A. A. (2017). Common contextual influences in ambiguous and
650 rivalrous figures. *PLoS ONE*, 12(5), Article e0176842. <https://doi.org/10.1371/journal.pone.0176842>.
- 651 Ouhnana, M., & Kingdom, F.A. (2016). Perceptual-binding in a rotating Necker cube: The effect of context
652 motion and position. *Vision Research*, 126, 59-68. <https://doi.org/10.1016/j.visres.2016.02.005>.
- 653 Overgaard M. (2017). The Status and Future of Consciousness Research. *Frontiers in psychology*, 8, 1719.
654 <https://doi.org/10.3389/fpsyg.2017.01719>.
- 655 Paffen, C. L., Alais, D., & Verstraten, F. A. (2006). Attention speeds binocular rivalry. *Psychological science*,
656 17(9), 752–756. <https://doi.org/10.1111/j.1467-9280.2006.01777.x>.
- 657 Parkkonen, L, Andersson, J, Hämäläinen, M, & Hari, R (2008). Early visual brain areas reflect the percept of an
658 ambiguous scene. *Proceedings of the National Academy of Sciences*, 105(51), 20500-20504.
- 659 Pastukhov, A., & Braun, J. (2011). Cumulative history quantifies the role of neural adaptation in multistable
660 perception. *Journal of vision*, 11(10), 12. <https://doi.org/10.1167/11.10.12>.
- 661 Pituch, K. A., & Stevens, J. P. (2016). *Applied multivariate statistics for social sciences* (6th ed.). New York:
662 Routledge. <https://doi.org/10.4324/9781315814919>.
- 663 Prete, G., Laeng, B., Fabri, M., Foschi, N., & Tommasi, L. (2015). Right hemisphere or valence hypothesis, or
664 both? The processing of hybrid faces in the intact and callosotomized brain. *Neuropsychologia*, 68, 94–106.
665 <https://doi.org/10.1016/j.neuropsychologia.2015.01.002>.
- 666 Rattat, A. C., & Droit-Volet, S. (2012). What is the best and easiest method of preventing counting in different
667 temporal tasks?. *Behavior research methods*, 44(1), 67–80. <https://doi.org/10.3758/s13428-011-0135-3>.
- 668 Rubin, E (1915). *Synsoplevede Figurer*, Doctoral thesis.
- 669 Sergent, J., & Bindra, D. (1981). Differential hemispheric processing of faces: Methodological considerations
670 and reinterpretation. *Psychological Bulletin*, 89(3), 541–554. <https://doi.org/10.1037/0033-2909.89.3.541>.
- 671 Seth, A. K., Dienes, Z., Cleeremans, A., Overgaard, M., & Pessoa, L. (2008). Measuring consciousness: relating
672 behavioural and neurophysiological approaches. *Trends in cognitive sciences*, 12(8), 314–321.
673 <https://doi.org/10.1016/j.tics.2008.04.008>.
- 674 Sterzer, P., Kleinschmidt, A., & Rees, G. (2009). The neural bases of multistable perception. *Trends in cognitive
675 sciences*, 13(7), 310–318. <https://doi.org/10.1016/j.tics.2009.04.006>.

676 Sucala, M., Scheckner, B., & David, D. (2011). Psychological Time: Interval Length Judgments and Subjective
677 Passage of Time Judgments. *Current Psychology Letters. Behaviour, Brain & Cognition*, 26(2), 1–10.
678 <http://journals.openedition.org/cpl/4998>.

679 Thomas, N. A., Castine, B. R., Loetscher, T., & Nicholls, M. E. (2015). Upper visual field distractors preferentially
680 bias attention to the left. *Cortex; a journal devoted to the study of the nervous system and behavior*, 64,
681 179–193. <https://doi.org/10.1016/j.cortex.2014.10.018>.

682 Thomas, N. A., & Elias, L. J. (2011). Upper and lower visual field differences in perceptual asymmetries. *Brain*
683 *research*, 1387, 108–115. <https://doi.org/10.1016/j.brainres.2011.02.063>.

684 Thönes, S., & Oberfeld, D. (2015). Time perception in depression: a meta-analysis. *Journal of affective*
685 *disorders*, 175, 359–372. <https://doi.org/10.1016/j.jad.2014.12.057>.

686 Tong, F., Meng, M., & Blake, R. (2006). Neural bases of binocular rivalry. *Trends in cognitive sciences*, 10(11),
687 502–511. <https://doi.org/10.1016/j.tics.2006.09.003>.

688 Toppino T. C. (2003). Reversible-figure perception: mechanisms of intentional control. *Perception &*
689 *psychophysics*, 65(8), 1285–1295. <https://doi.org/10.3758/bf03194852>.

690 Vatakis, A., Balci, F., Di Luca, M., & Correa, Á. (2018). *Timing and Time Perception: Procedures, Measures, &*
691 *Applications*. Leiden, The Netherlands: Brill. doi: <https://doi.org/10.1163/9789004280205>.

692 Voßkühler, A., Nordmeier, V., Kuchinke, L., & Jacobs, A. M. (2008). OGAMA (Open Gaze and Mouse Analyzer):
693 Open-source software designed to analyze eye and mouse movements in slideshow study designs.
694 *Behavior Research Methods*, 40(4), 1150–1162. <https://doi.org/10.3758/BRM.40.4.1150>.

695 Wang, M., Arteaga, D., & He, B. J. (2013). Brain mechanisms for simple perception and bistable perception.
696 *Proceedings of the National Academy of Sciences of the United States of America*, 110(35), E3350–E3359.
697 <https://doi.org/10.1073/pnas.1221945110>.

698 Wearden J. H. (2015). Passage of time judgements. *Consciousness and cognition*, 38, 165–171.
699 <https://doi.org/10.1016/j.concog.2015.06.005>.

700 Xuan, B., Zhang, D., He, S., & Chen, X. (2007). Larger stimuli are judged to last longer. *Journal of vision*, 7(10),
701 1–5. <https://doi.org/10.1167/7.10.2>.

702 Zakay, D., & Block, R. A. (1996). The role of attention in time estimation processes. In M. A. Pastor & J. Artieda
703 (Eds.), *Time, internal clocks and movement* (pp. 143–164). North-Holland/Elsevier Science Publishers.
704 [https://doi.org/10.1016/S0166-4115\(96\)80057-4](https://doi.org/10.1016/S0166-4115(96)80057-4).

705 Zakay, D., & Block, R. A. (1997). Temporal Cognition. *Current Directions in Psychological Science*, 6(1), 12–16.
706 <https://doi.org/10.1111/1467-8721.ep11512604>.

707 Zakay, D., Block, R. A., & Tsal, Y. (1999). Prospective duration estimation and performance. In D. Gopher & A.
708 Koriati (Eds.), *Attention and performance XVII: Cognitive regulation of performance: Interaction of theory*
709 *and application* (pp. 557–580). The MIT Press.

710 Zeman, A. (2001). Consciousness. *Brain: A Journal of Neurology*, 124(7), 1263–1289.
711 <https://doi.org/10.1093/brain/124.7.1263>.

712 Zimmer, H. D., & Ecker, U. K. (2010). Remembering perceptual features unequally bound in object and episodic
713 tokens: Neural mechanisms and their electrophysiological correlates. *Neuroscience and biobehavioral*
714 *reviews*, 34(7), 1066–1079. <https://doi.org/10.1016/j.neubiorev.2010.01.014>.

715


716

717 Figures

718

719

720



721

722

723

724


725 **Fig. 1.** The stimuli (Rubin's face-vase figures) presented in the two experiments in the 2F condition
726 (*top-left*), 4F condition (*top-right*), and 8F condition (*bottom-left*). The cross indicates the fixation
727 point. In the 1F condition only one Rubin's face-vase figure was presented instead of the cross at the
728 center of the screen. *Bottom-right*: the stimulus presented before in the training session for the 8F
729 condition. Letters above each image indicate the button that was to be pressed to indicate a reversal
730 perception in the corresponding image. See text for the relative size of the figures presented in the
731 two experiments.

732

733

734


735
736
737



738
739
740
741
742
743
744
745

Fig. 2. Means and standard errors for the bistability score (number of reversals) and for the time estimation in the 4 conditions of Experiments 1 and 2.

746
747
748



749
750
751
752

753 **Fig. 3.** Scatterplot showing the global correlation (n=60) between the bistability score (n, number of
754 perceived reversals averaged across conditions) and time estimation (min, session duration
755 estimation averaged across conditions) in both experiments together ($r = -0.30$, $p = 0.022$).