

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Factors Contributing to Volcano Lateral Collapse

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Roverato, M., Di Traglia, F., Procter, J., Paguican, E., Dufresne, A. (2021). Factors Contributing to Volcano Lateral Collapse. Cham : Springer Nature [10.1007/978-3-030-57411-6_5].

Availability:

This version is available at: <https://hdl.handle.net/11585/963195> since: 2024-02-28

Published:

DOI: http://doi.org/10.1007/978-3-030-57411-6_5

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Factors contributing to volcano lateral collapse

Matteo Roverato^{1,2}, Federico Di Traglia³, Jonathan Procter⁴, Engielle Paguican^{5,6}, Anja Dufresne⁷

- 1- School of Earth Science, Energy and Environment, Yachay Tech University, Urcuqui, Ecuador
- 2- Department of Earth Sciences, University of Geneva, Switzerland
- 3- Dipartimento di Scienze della Terra, Università degli Studi di Firenze, Firenze, Italy
- 4- School of Agriculture and Environment, Massey University, Private Bag 11 222, Palmerston North, New Zealand
- 5- School of Geography, Environment and Earth Science, University of the South Pacific, Laucala Campus, Suva, Fiji
- 6- College of Engineering and Geosciences, Caraga State University, Ampayon, Butuan City, Philippines
- 7- Engineering Geology and Hydrogeology, RWTH-Aachen University, Aachen, Germany

Abstract

Many factors can lead to volcano lateral collapse, which can produce devastating debris avalanches that travel up to several tens to over 100 km and cover hundreds to more than a thousand km² with debris. Volcanic lateral collapses are severe hazards because of their destructive power and size, and sudden onset. Although their frequency of occurrence is not as high as those of smaller volcanic mass movements, such as rock falls and lahars, globally large collapses ≥ 0.1 km³ have occurred at least five times per century over the last 500 years. A large variety of destabilizing factors such as over-steepened slopes, magma intrusions, hydrothermal activity, climate fluctuations, deformation of the basement, and faulting can create the conditions for volcano collapse. Once a volcano reaches its critical point, a mechanism is necessary to trigger the failure event. We present the state-of-the-art of the knowledge acquired in the last few decades concerning the causes of large-scale volcanic failures to better understand the triggers, preparatory factors, and timing of volcano lateral collapse.

Keywords: volcanic lateral collapse; debris avalanche; instability factors; triggering mechanisms

1 - Introduction

Large long-lived composite volcanoes are positive topographic structures characterized mainly by constructive phases. The stability of the edifice is primarily controlled by the characteristics of the outer slope, which, when it reaches a critical configuration, can become unstable. Edifice failure may also be triggered by external processes even before it reaches its critical configuration. Slope instability in volcanoes typically results in a structural failure leading to lateral collapse. Volcano flank movement can be classified as: i) persistent flank motion, typically deep-seated, steady-state movement of large sectors of a volcano edifice due to gravity; ii) transient flank motion (i.e. flank “unrest”), considered the precursor to catastrophic collapses, associated with intrusive processes, co-eruptive deformation, seismic shaking; or iii) surficial slope motion, or shallow motion such as material erosion and remobilization of volcanoclastic and lava deposits (Schaefer et al. 2019). Edifice collapses are not restricted to stratovolcanoes, but have occurred on all types of volcanoes (dome complexes, shield volcanoes, calderas and even within monogenetic fields) (Dufresne et al. 2020; this volume) and may happen in all geodynamic contexts (Ui 1983; Siebert 1984; McGuire 1996; Voight and Elsworth 1997; Voight 2000; Shea and Van Wyk de Vries 2010; Van Wyk de Vries and Davies 2015).

Such volcano sector collapses leave behind a scar that could be of a caldera size (kilometers across) and shape (Siebert, 1984; McGuire 2006). This might explain why, before the 1980 Mount St. Helens event, many wide depressions hosted in volcanic summits were considered calderas (Siebert and Roverato 2020 - this volume), which form by vertical subsidence along ring faults due to eruptive events, rather than by a lateral collapse. This erroneous interpretation is understandable because horseshoe-shaped scar can be partly obliterated by volcano regrowth covering the failure area with new eruption deposits. Belousov et al. (1999) wrote: “Evaluation of known cases of large-scale failures on volcanoes throughout the world shows that failure can occur, given the right combination of circumstances, at almost any reasonably high volcanic edifice”. It appears that this behavior is ubiquitous, with evidence for volcanic failures both within the geological record and at many currently active volcanoes. For example, Francis (1994) wrote that 75% of Andean volcanic edifices, with heights greater than 2,500 m, have experienced at least one sector collapse, while in Indonesia, most volcanoes with heights greater than 2,000 m have suffered the same fate (MacLeod 1989). Inokuchi (1988) reported that over 100 volcanic debris avalanche deposits (VDADs) have been identified around Japanese Quaternary

volcanoes. Volcanic collapses, and their associated VDAs, are severe potential hazards not only because of their destructive power and size but, as stressed by Siebert and Roverato (2020 – this volume), because volcanic edifice failures $\geq 0.1 \text{ km}^3$ have occurred at least five times per century over the last 500 years. Augustine volcano (Alaska, USA) has repeatedly collapsed, once every 150-200 years during the past 2,000 years (Begét and Kienle, 1992), although those were at relatively “small” volumes ($< 0.25 \text{ km}^3$). Mount Taranaki (New Zealand) shows a similar pattern of growth and collapse with 14 VDADs over its 120,000 years evolution with a regular 16,000 years return period (Zernack et al. 2012). Colima volcano collapsed at least nine times during the last 20,000 years (Capra et al. 2002). This frequency and cyclic behavior should not be underestimated for hazard assessment.

The study of this type of phenomenon strengthened after the collapse and eruption of Mount St. Helens volcano (USA) on 18 May 1980. This event represents the only case where lateral collapse of the volcanic edifice has been observed directly (Lipman and Mullineaux 1981). Although the eruption of Mount St. Helens represented an event of great importance for the understanding of such phenomenon linked to the collapse of a volcanic edifice, the factors that prepare and trigger these events are not simple, but can be multiple and combined. The effects of a volcanic lateral collapse are usually of great magnitude. Mass movement on volcanic slopes can occur at different scales, which can span from relatively minor rock falls consisting of a few hundred to a few thousand cubic meters of material up to the giant landslides that may involve materials more than $1,000 \text{ km}^3$ (McGuire 1996, 2003). Events involving small volumes of material may occur with time intervals of weeks or months, whereas larger events are having frequencies of 10s to 100s of thousands of years (Schaefer et al. 2019). The large volumes observed in many volcanic debris avalanches involve the collapse of part of a volcanic edifice and this can in turn influence the runout of a VDAD. These destructive flows can travel at high velocities, cover large areas, overcome topographical barriers and modify the topography in seconds. Generally, these deposits could preserve evidence of the cause of the instability, but it is often difficult to determine the triggering mechanism. Volcanic lateral collapses can be caused by a wide variety of destabilizing factors such as over-steepened slopes, increasing fluid pore pressures (consequence of several combined factors), magma intrusions, hydrothermal alteration, climate fluctuations, deformation of the basement, cataclysmic eruptions, among others (Table 1). It is important to identify the differences between those

factors that define the degree of instability of a volcanic edifice from those mechanisms that trigger the collapse. Indeed, the mechanism that triggers a sector collapse could be independent of the causes that prepare its instability. For example, the tectonic setting of a volcano may influence the direction of the collapse and prolonged hydrothermal alteration can deeply weaken the edifice, but the collapse itself could be triggered by an eruption or an earthquake.

Table 1

Sector collapses can cause tsunamis or lahars increasing the areas and population at risk. In 2018, tsunamigenic lateral collapse of Anak Krakatau, Indonesia (Williams et al. 2019) and lateral collapse of Kilauea, US (Chen et al. 2019) and Etna, Italy (Bonforte et al. 2019) occurred, with resulting loss of life and property. It is therefore important to understand the instability factors and triggers of volcano lateral collapse since that knowledge can provide very important precursory information that can directly improve hazard warnings (Capra 2006).

2 - Instability factors

A wide range of factors may contribute to the instability of a volcanic edifice, and some volcanoes are more prone to lateral failure than others, depending on their structure and the environment in which they form (McGuire 2003). Large-scale structural instability is invariably confined to major composite volcanoes, located either on continental (e.g. Shiveluch in Russia, Taranaki in New Zealand, and Colima in Mexico) or oceanic (e.g. Teide in Canary Islands - Spain, and Piton de la Fournaise on Reunion Island) crust. Destabilization of a volcanic edifice may take thousands or tens of thousands of years, or it may be achieved in only a few months (McGuire 2003). One well-known example of short-lived destabilization-to-collapse event is the Mount St. Helens collapse, where a cryptodome intruded into the edifice only few months before the collapse. That is in contrast to slow, incremental and progressive factors, such as long-term hydrothermal alteration, or long-lasting deformation that can occur individually or often act together as cumulative effects of many small events. Below, we describe the instability factors that favor the weakening of a volcanic edifice and promote sector collapses.

2.1 Basement, tectonics, and faults

Volcanism is linked to tectonically active regions, whether extensional, compressional or along transcurrent faulting, in both transcompressional and transtensional regimes (Tibaldi et al. 2005). Tectonic faults propagate from the substrate and into the volcano edifice, where their geometry and kinematics become more complex. A close relationship exists between fault movement and the stability of the overlying volcanic edifice: basement faults affect volcanoes (Francis and Self 1987; Carracedo 1994; Tibaldi 1995) and vice versa (van Wyk de Vries and Merle 1996; van Wyk de Vries 1998). Volcano edifice morphology, including collapse directions and fracture planes used as pathways for magma, respond to regional stress fields in complex ways (Delaney et al. 1986). Preferential directions of progressive flank destabilization depend not only upon the orientation, kinematics, and dip and inclination of basement faults (Tibaldi et al. 2005), but also on the position of the fault within the edifice (Norini et al. 2008; Wooller et al. 2009; Norini et al. 2020). Once volcanic failure occurs, the collapse depression and the new geometry of the volcanic edifice strongly influence the location and configuration of new craters and dykes.

Nakamura (1977) suggested that cone elongation, strike of dikes and fractures, and alignment of flank vents occur parallel to the regional maximum compressive stress direction (σ_{Hmax}) (Figurea). Moriya (1980) suggested that the elongated growth of volcanic flanks parallel to dike emplacement, steeper slopes normal to the elongation direction of volcanoes, and the dilatational effect of dike intrusion all promote failure perpendicular to σ_{Hmax} , as was observed at 30 Japanese volcanoes. The higher percentage of volcanoes with parallel dike swarms that formed amphitheater-like scars relative to volcanoes without a dominant direction of dike orientation indicates that dike swarms that develop parallel to σ_{Hmax} play an important role in volcanic lateral collapse (Siebert 1984). Francis and Wells (1988) suggested a model of failure scar opening perpendicular to the strike of normal faults and σ_{Hmax} (Figureb).

These models, however, are not consistent with the alignment of volcanoes, dikes, flank vents, and collapse directions in Indonesia, where they tend to be at a high angle to σ_{Hmax} as determined from fault-fold structures in the region (Bahar and Girod 1983). An examination of 39 Japanese volcanoes revealed failure scar openings at an acute angle relative to σ_{Hmax} (Ui et al. 1986). The same is true for the 28 examples of breached cones in the Central Andes (Figurec) (Francis and Wells 1988).

Figure 1

The inconsistencies in these studies led to the idea that the opening direction may not be influenced by the regional tectonic field, but may be controlled by temporal and local stress fields within the volcanic edifice or the different geodynamic settings where volcanoes are located (Ui et al. 1986).

In a study of 1,315 Quaternary volcanoes in different structural settings, Tibaldi (1995) established that failure scar breaching perpendicular to fault strike is prevalent in extensional regions, and collapses are usually in the direction of the downthrown block (Vidal and Merle 2000). This is consistent with the model of Moriya (1980) and Siebert (1984), where plate-shaped conduits and dikes could propagate through extensional faults and thereby promote collapse perpendicular to the regional σ_{Hmax} . Also, in extensional regimes, the regional stress field produces brittle discontinuities that crosscut the edifice by forming rift zones, used as a pathway for the magma to reach the surface. This feeds dyke swarms in a preferred orientation, which then displace the volcano flanks and may induce lateral collapse of the edifice (Elsworth and Voight 1996; Tibaldi 1996).

In contrast to extensional regions, breaching in volcanoes along transcurrent faults or transtensional tectonics is at an acute angle relative to σ_{Hmax} (Tibaldi, 1995; Norini et al., 2008). This is true for the volcanoes in Southwest and Southeast Luzon, Philippines including Iriga (Figure 1d; Lagmay et al. 2000; Lagmay and Valdivia 2006; Paguican et al. 2012), as well as Mount St. Helens, USA. Lateral collapses in transtensional tectonic regions underlain by normal faults are more diffused compared to transcompressional and reverse tectonic regions due to tilting and topographic offsets within the basement (Tibaldi 2005). Analogue experiment by Lagmay et al. (2000) and Norini and Lagmay (2005) showed that edifices on top of strike-slip faults form a pair of sigmoids composed of one reverse and one normal fault on the surface, and a flower structure in the interior of the edifice. Two destabilized regions characterized by bulging and landsliding are created on the flanks between the traces of the sigmoidal faults. Sigmoidal summit depressions, fractures, and faults developed within the edifice create space for subsequent magma intrusion, thus causing further instability (van Wyk de Vries and Merle 1996).

The variety of fault types and geometries has a general tendency to produce regions of instability parallel to fault strike, irrespective of the fault motion. Analogue experiments by Wooller et al. (2009) showed that in oblique-slip faulting, the instability is always on the downthrown side, and usually in the volcano sector facing the strike-slip sense of motion (Figure 2). The location, type, and magnitude of instability of the edifice changes with the position of the fault beneath the analogue volcano. The further the fault is from the central axis, the larger the destabilized sector. Also, with greater fault offset from the central axis, larger unstable volumes are generated. Such failures are perpendicular to the strike of the underlying fault.

These models may not work in volcanoes where an underlying sloping substrate or gravitational spreading have a stronger influence on the collapse direction (van Wyk de Vries and Merle 1996).

Figure 2

2.2 Sloping substrate and gravitational spreading

Many volcanoes grow on sloping ground, and some of them tend to slide downhill and collapse catastrophically on the downslope side (Vallance et al. 1995; Carrasco-Núñez et al. 2006; Murray et al. 2018). Many large stratovolcanoes worldwide are directly in contact with thick sedimentary or igneous sequences that have been affected by different compressional and extensional processes depending on the geotectonic evolution of the area (e.g. the Andean mountain range). These sequences often appear tilted, showing a certain grade of dipping and generating a strength anisotropy. In northern Chile, for example, widespread ignimbrite sheets and thick ductile Cenozoic sedimentary deposits dip toward the west into the Salar de Atacama basin with an angle of 5° (Wooller et al. 2004). The Tumisa volcanic complex, located northward in the same region, is underlain by several hundred meters of salt-rich lake sediments, ignimbrites and evaporite deposits, which together form a westward-dipping ductile package (Kuhn 2002). According to Wooller et al. (2004), the entire Tumisa volcanic complex has undergone a downslope sliding toward the same western direction. Analogue experiments presented by these authors demonstrate

that volcano spreading is highly dependent on substrata dip and spreading occurs even at very small substrata tilt ($< 1^\circ$).

Many other Central Andean volcanoes experienced slope collapses that left horseshoe-shaped scars oriented toward the direction of the regional slope (Francis and Wells 1988). Vallance et al. (1995) noted a similar pattern with Guatemalan volcanoes, which failed preferentially trenchward to the southwest in the direction of regional slope of the Guatemalan Highlands. At the eastern end of the Mexican Volcanic Belt, the Cofre de Perote-Citlatépetl range is constructed at the margin of the Altiplano. The basement drops dramatically to the Caribbean coastal plain, and all collapse events have been in that direction (Carrasco-Núñez et al. 2006). In Central America, Siebert et al. (2006) noted that with volcanoes at the far ends of the arc, failures tended to occur preferentially perpendicular to the trend of the arc and in a direction toward the Central-American trench. In contrast, volcanoes in El Salvador and Nicaragua in the center of the arc, where the basement topographic effect is less pronounced, have more variable failure directions. This is even more pronounced at an isolated island volcano like Augustine in Alaska, without regional topographic gradients, where failures occurred radially in all directions (Begét and Kienle 1992).

The same behavior occurs at volcanoes in Indonesia that are located on dipping ductile substrata and have undergone volcano spreading, or at the volcanoes belonging to the Lesser Antilles, in the Caribbean Sea. Deplus et al. (2001) report that the volcanoes are situated on westward-dipping substrata related to the Tertiary Caribbean arc, the same orientation in all major collapses of the region.

Numerous studies (Merle and Borgia 1996; Lundgren et al. 2004; Bonforte et al. 2011; Murray et al. 2018) reveal that the eastern and southern flanks of Mt. Etna are sliding eastward towards the Mediterranean Sea, affected by the slope and lithology of the basement, with zones of differential velocities forming several fault-bounded blocks. It is not coincidental that the “Valle del Bove” scar is oriented in the same direction as Mt. Etna creeps. Lateral collapses might also occur for those volcanoes located on steeper dip slopes, compared with low-angle dip basement mentioned above. Due to the migration of an eruptive vent, a consequence of various factors (e.g. changes in the plate subduction angle, variation in the tectonic setting, migration of the magma chamber, among others), a volcano might grow on the flank of an older volcanic edifice with slopes angles $> 20\text{-}25^\circ$. A

good example is the Fuego de Colima volcano (Fig. 3), which is located on the southern flank of the older and inactive Nevado de Colima volcano. Whilst the direction of the sector collapses of the Nevado de Colima volcano has been controlled by the Colima Rift confinement (Norini et al. 2019), the inclination of the southward dipping Nevado de Colima flank appears to be the main topographic constraint generating the instability of the Fuego de Colima edifice (Norini et al. 2010; Roverato et al. 2011; Norini et al. 2019).

Figure 3

At a lower scale, on 16 November 2006 a lateral collapse affected the unstable eastern slope of the steep South-East Crater (SEC) of Mount Etna, built on one side of the steep central craters of the volcano (Norini et al. 2009). The collapse was triggered by erosion of loose, hydrothermally altered material of the steep south-east sector of SEC from the outpour of lava and produced a landslide of 330,000 - 413,000 m³ that involved both lithic and juvenile material, resulting in a deposit emplaced on the eastern flank of the volcano (Norini et al. 2009). Thus, the substrata dip and the steepness of volcano flanks is important for volcanic edifice instability, as it can control the spreading to downslope sectors and the associated sector-collapse volumes. Differential buttressing at the volcano base is also a major factor controlling flank instability and is an important indication to expect asymmetric activity in any volcano, like flank eruptions and deformations (Norini and Acocella 2011).

Volcanic gravitational spreading is one of the most complex factors contributing to instability that can affect a volcanic edifice. This is because numerous variables are involved in the process, although the most important of those is the substratum on which the volcano sits. Gravitational spreading is the result of a set of processes that translate the vertical volcano load into horizontal movement, accommodated by high-angle normal faults on the edifice's slopes and low-angle thrust faults at its boundaries and in the substratum. In a pioneer work, van Bemmelen (1949) reported faults across volcanic edifices and anticlinal ridges and low-angle thrusts along their peripheries for several Indonesian volcanoes (e.g. Soropati, Lawu, Merapi, Bukittinggul, Tjereme). These volcanoes are built on thick sequences of marine clays and fine-grained deposits. The deformation structures of the substrata are attributed to sediment faulting beneath the volcanoes in response to

gravitational loading (Borgia et al. 2002). At Parinacota (Chile), the deformation of low-density unconsolidated fluvio-glacial and lacustrine sediments and pyroclastic basin-fill sequences in response to volcano loading is considered as the main causative factor of the ca. 8,000 years old slope collapse that produced a 6 km³ VDAD (Clavero et al. 2002). Other researchers described such typical features of deformation in the substratum sediments, and it appears that they play a fundamental role in controlling the stability of a volcanic edifice and the size of the collapsed landslide. At Mombacho volcano (Nicaragua), two Holocene collapses are well described in the literature, one toward the south (El Crater) and the other to the northeast (Las Isletas) (van Wyk de Vries and Francis 1997; Shea et al. 2008; Shea and van Wyk De Vries 2010). The base of Mombacho volcano in its northeastern side is surrounded by an anticline structure, which was likely formed as Mombacho was slowly spreading outwards (Shea and van Wyk de Vries 2010). A pumice-rich substratum has also been found at the base of the Las Isletas VDAD, much further from the original in-situ ignimbrite deposit. Sedimentary sequences in the deposit indicate extrusion of the substratum, which is eventually ripped up and transported by the VDA resulting from lateral collapse. Socompa volcano, located on the southeastern margin of the Atacama basin (Chile/Argentina), suffered a dominantly westward gravitational spreading (Wooller et al. 2004). The collapse was triggered by failure of active thrust-anticlines in sediments underlying the volcano. High amounts of the substratum sediments have been found within the Socompa VDAD (van Wyk de Vries et al. 2001).

Two distinct styles of deformation are described: 1) “volcano spreading” if the volcanic edifice spreads with the substrata, and 2) “substrata extrusion” (van Wyk de Vries and Matela 1998) when the volcano and substrata are decoupled and the substrata are extruded from under the edifice. In both cases, compressional structures are produced at the base of the volcano. At Socompa volcano, the gravitational deformation has developed mainly by substrata extrusion expressed by a series of thrust anticlines (van Wyk de Vries et al. 2001). Major structures observed on the summit of other volcanoes like Etna and Kilauea (Borgia et al. 1992; Morgan et al. 2003; Urlaub et al. 2018) suggest that, at the beginning of gravitational spreading, the volcanoes present grabens and rifts on their summits. These features are also reported by Del Camp et al. (2008) for Piton de la Fournaise volcano (Reunion Island), Concepción and Maderas volcanoes (Nicaragua), Merapi volcano (Indonesia), and other Andean Chile/Bolivian volcanoes. Recent analogue models by Shea

and van Wyk de Vries (2008), Andrade and van Wyk de Vries (2010) and Paguican et al. (2012) explored the kinematics of a volcanic failure from the early stages of a sector collapse. Andrade and van Wyk de Vries (2010) reproduced in their experiments how grabens and rift-structures develop at the summit of volcanic edifices in response to the compressional forces at the base of the edifice. Furthermore, they report that most of the viscous basal layers used in the experiments had been expelled from underneath the volcano as in natural examples (van Wyk de Vries and Francis, 1997; van Wyk de Vries et al. 2001). One important insight from these experiments is that gravitational spreading stops or is reduced after a catastrophic collapse (Andrade and van Wyk de Vries 2010). This is an obvious conclusion as the volcanic loading decreases considerably after part of the edifice is removed and spread onto the surrounding landscape. Alternatively, gravitational spreading enhances catastrophic collapse only during its initial stages. When no catastrophic collapse occurs, the volcanic edifice will continue to spread progressively, gradually reducing the slope of its flanks (Andrade and van Wyk de Vries 2010; Norini et al. 2010).

2.3 Hydrothermal alteration

Volcanoes are actively growing and evolving systems characterized by the interaction of fluids, gases and heat that can hydrothermally alter the edifice. Hydrothermal alteration causes rock dissolution and mineral-chemical modification, it can promote high pore pressures, weaken the volcanic edifice's rocks and increase the overall clay-content. Hydrothermal alteration can occur progressively over long periods (>100 yrs), allowing deformation to develop slowly before collapse (Van Wyk de Vries et al. 2000). Each hydrothermal system develops in a unique manner depending on the rate of fluids that circulate within the volcano, their temperature, the porosity and permeability of the material forming the edifice, as well as the amount of external meteoric water infiltrating into the edifice. Consequently, each hydrothermal system has its own distinctive size, volume, chemical, and physical characteristics (Cecchi et al. 2004). Many large volcanic lateral collapses have involved hydrothermally altered rocks (Siebert 1984), which can be a useful indicator of the source of the collapse, and the slope destabilization and collapse triggering mechanisms. Examples of VDADs that display hydrothermal alteration (Figure 4) are reported in the literature and some are listed in Table 2.

Figure 4

Clay minerals from hydrothermal alteration are also found in some avalanche-induced lahar deposits at Mount Rainier, including the 3.8 km³ Osceola mudflow (Vallance and Scott 1997), the 0.2 km³ Round Pass mudflow, and the 0.26 km³ Electron mudflow (Scott et al. 1995). The presence of hydrothermally altered rocks in these deposits suggests that this process partly weakened the edifice and likely promoted the slope collapse (Reid et al., 2001). Mount Rainier has large areas of hydrothermally altered rocks (Frank, 1995; Zimbelman, 1996; Reid et al., 2001; John et al., 2008). Reid et al. (2001) noted that the alteration in highly porous rocks such as scoria and pumice is less intense and present mainly as vesicle infillings. On the other hand, the alteration is more severe in dense lavas and dikes, largely obliterating original igneous minerals and textures, and contributing to an overall reduction in rock mass strength.

Table 2

The Holocene Teteltzingo deposit, a ~2 km³ clay-rich VDAD formed by the Citlaltépetl (Mexico) volcano's slope collapse, displays a high clay content in the range of 10 to 16% (Carrasco-Núñez et al., 1993). Its features suggest that it had an origin from a water-saturated hydrothermally altered edifice and that it transformed into a cohesive lahar at the very beginning of the failure event. The volume of altered rocks involved in the collapse was on the order of 10⁸ to 10⁹ m³ (Carrasco-Núñez et al. 1993). Currently, the Citlaltépetl volcano is still characterized by a widespread alteration enhanced by the glacier (melted-ice circulation), including massive zones of replacement and vein fillings along numerous fractures (Zimbelman et al. 2004). Much of the replacement alteration is of acid sulfate. During vapor condensation, the SO₂ responsible for the alteration is present in very high concentrations, and a fluid of very low pH forms (Zimbelman et al. 2004). This acid brine reacts with the host rocks, leaching it and creating an increase in permeability. The voids left by the acid dissolution are now available to be filled by fluids such as meteoric water, which may increase the pore pressure into the volcanic edifice if sufficient water exploits the new pathways. Lopez and Williams (1993) stressed the importance of hydrothermal fluids circulation into a volcanic edifice and considered the hot spring compositions, residence times and flow path through faults of fluids to model potential collapse scenarios at Nevado

del Ruiz volcano (Colombia). The continual hydrothermal activity reduces the strength of the rock (mainly due to clay mineral formation), and favors the more porous lithologies, which, in turn, can increase pore pressures when fluids fill the voids (Day 1996; Voight and Elsworth 1997; Reid 2004). Day (1996) suggests that the hydrothermal alteration is more effective in the interior of volcanic edifices where higher pressures and temperatures could dissolve the rocks more effectively (Frantz et al. 1981) creating high-permeability zones. On the contrary, in the exterior and cooler parts of the edifice, clay minerals will precipitate from the hydrothermal fluids, decreasing the permeability and promoting for example, meteoric water storage. There is growing evidence that the internal fluid pressure in a volcanic edifice is one of the main factors for potential instability. For example, Borselli et al. (2011) report that, at Colima volcano (Mexico), the fluid pressure likely produces a local reduction of shear strength along a portion or all of a potential sliding surface, promoting a global reduction of the stability factor of the volcanic edifice. This idea is supported by a recent resistivity tomography survey by Rosas-Carbajal et al. (2016) in which they investigated the structure and dynamics of the hydrothermal system at La Soufrière volcano (Guadalupe). The authors suggested that highly altered, fluid-saturated, weak zones and the low basal friction of argillized detachment planes could significantly reduce the factor of safety of the volcano slope. Finally, the recent study of Delmelle and Bernard (2015) showed how slope instabilities, associated with hydrothermal alteration, can increase by the formation of acidic crater lakes on the summits or within the flank topography of active volcanoes, such as the Kawah Ijen hyperacid lake ($\text{pH} < 0.5$) on the flank of Merapi volcano. Leakages through the flanks of the acid lake and dissolution beneath it, weakens the interior of the volcano and build a substantial body of highly altered rock within the upper part of the volcanic edifice that reduces its stability (Delmelle et al. 2015).

2.4 Dikes and magma intrusions

Historical volcanic lateral collapses at Bezymianny, Kamchatka in 1957, Mount St. Helens in 1980, and Soufrière Hills, Montserrat in 1997, all generated large eruptions with associated lateral blasts caused by large-scale deformation and steepening of flanks prior to failure (Gorshkov and Bogoyavkenskaya 1965; Gorshkov 1959; Voight et al. 1983; Siebert et al. 1987; Belousov et al. 2004). These volcanoes provide examples of lateral collapse initiated by magma intrusions and stress the importance that this represents for volcano

stability and hazard assessment. Instability can be related to the intrusion (dikes and cryptodomes), as well as extrusion (summit domes) of fresh magma, and can trigger failures through overloading and oversteepening, deformation, or mechanical- and temperature-related changes in pore pressures (Elsworth & Voight 1996; Day 1996; McGuire 2003, Norini and Acocella 2011; Samaniego et al. 2015). For example, the high eruption rates of silicic domes and lava flows at Mount St. Augustine (USA) repeatedly produced an oversteepened, unstable volcanic cone which then collapsed (Beget and Kienle 1992). At Ruapehu (New Zealand) a paleomagnetic study (McClelland and Erwin 2003) revealed that a summit dome could have played some contributory part in the destabilization of the edifice. High temperatures (350°C) of some dacite blocks into the 9.5 Ka BP Murimotu formation suggests that a dacite dome was active on Ruapehu immediately prior to the lateral collapse. In Mexico, Macias et al. (2010) report that the late Pleistocene collapse of Tacaná volcano was likely initiated by the extrusion of an andesitic lava dome that weakened the structure of the cone. A recent contribution by Rincon et al. (2018) tried to explain, with analog models, how volcanoes could be destabilized by magma intrusion. The authors noted that a large and deep listric fault crossing the volcano can form when the magma body is still located at the base of the cone. This explains why, in some cases, a volcanic edifice could collapse even if the intrusion does not reach shallower levels. If any trigger induces the slope failure at this stage, no juvenile material will be involved in the collapse. Rincon et al. (2018) also noted that, when the intrusion reaches shallower levels, an elongated subsidence forms at the summit of the cone and a bulge at the middle flank develops. The same deformation was noted at Mount St. Helens before its collapse (Lipman et al. 1981) and described also in other analogue experiments (Donnadieu and Merle, 1998). The intrusion of a cryptodome (Figure 5) into the upper part of a volcanic edifice

Figure 5

decreases its stability due to many different factors, for instance lower shear strength of the magmatic body and associated weakening structures, shallow seismicity on the flank caused by edifice fracturing in response to intrusion, outward-dipping normal faults bounding a summit graben, hydrothermal alteration of large portions of the edifice, and bulging (i.e. increase in steepness) of the destabilized flank (Donnadieu et al. 2001). Elsworth and Voight

(1996) have drawn attention to the potential role of magmatic intrusions in triggering structural failure as a consequence of their raising pore fluid pressures by means of mechanical or thermal straining (thermal expansion) of the rock-fluid medium and/or hydrothermal fluid circulation. The thermal expansion can develop large uplift forces beneath a potentially unstable block (Elsworth and Voight 1995). The magma has a gas-rich phase at depth. This, coupled with magma degassing at shallower levels, could lead to excess upward-propagating pore pressures (Thomas et al. 2004). In addition, dehydration (boiling) of active hydrothermal systems during magma emplacement can lead to considerably elevated pore pressures (Reid 2004). Once a volcanic system has reached a critical state, collapse may be initiated by any trigger or simply occur because the factor of safety is reduced sufficiently. At Shiveluch volcano, the cause of the destabilization was the ascent of a batch of viscous, water-rich magma to shallow levels (Belousov et al. 1999). The failure occurred before magma intruded into the edifice suggesting that the trigger mechanism was not magma-associated. A more likely destabilizing mechanism could have been related to the increase of pore pressure in the edifice due to pressurized gas released from the ascending magma (Belousov et al. 1999). Evidences from fumarolic gases in the thermal springs at La Soufrière of Guadeloupe volcano (Lesser Antilles) suggest that the current activity is the result of a new magma intrusion which was progressively emplaced at shallow depth in the last decades (Villemant et al. 2014). The assessment of potential hazards associated with the evolution of the current unrest must consider the implications of recurrent intrusions and further pressurization of the hydrothermal system. The role of hydrothermal pressurization, linked with magma intrusion, on the basal friction along low-strength layers within the upper part of the edifice must be evaluated with regards to slope collapses (Borselli et al. 2011; Villemant et al. 2014).

2.5 Past and present climate implications

Climate can influence volcanic activity and plays an important role in the stability of a volcanic edifice. Heavy rains or hurricanes, for example, can promote massive landsliding over large areas. During the passage of Hurricane Mitch above Central America in 1998, heavy rains promoted a lateral collapse of Casita volcano (Nicaragua), killing 2,500 people (Kerle and van Wyk de Vries 2001; Scott et al. 2005). The source volume of the landslide (~200,000 m³) was comparatively small to other giant volcanic collapse-related mass flows.

However, the landslide entrained older deposits and organic material during the flow, transforming into a devastating debris avalanche (Kerle and van Wyk de Vries 2001) and subsequently into a debris flow (Scott et al. 2005). It is common worldwide that debris avalanches of Casita-size occur every few years (McGuire 1996; Kerle and van Wyk de Vries 2001). Climatic hazards rapidly affect widespread areas contributing to volcanic instability. Heavy rains during wet seasons and glacier retreat have become more prominent over the last several decades due to changing climatic conditions as a result of global warming (Capra 2006; Roberti et al. 2020 - this volume). Recent examples in periglacial environments of ice-avalanches, rock avalanches, and a mix between both have been considered to be increasing by these changing conditions (Huggel et al. 2007; Huggel et al. 2008; Tuffen 2010; Kos et al., 2016; Coe et al. 2018; Hibert et al. 2019). The 2010 and other large landslides at Mount Meager volcano (Canada), for example, resulted, in part, from glacier retreat and erosion contributing to changes in stress of its already unstable slopes (Grämiger et al. 2017; Roberti et al. 2020 - this volume). With current accelerated rates of glacial melting, glaciated active volcanoes are at an increasing probability of generating landslides, from shallow failures to debris flows to large-volume VDAs. Well-known paleo-climate records from the last deglaciation show that melting of ice triggered a dramatic acceleration in volcanic eruptions (Capra et al. 2015). However, major glaciations (and deglaciations) have been considered not only as a triggering mechanism for large magmatic eruptions but also inducing volcano instability (Capra et al. 2013). Ice-capped volcanoes are sensitive environments to climatic changes, and large-scale glacial melting occurring at the end of the Pleistocene, caused a significant increase in the incidence of large volcanic lateral collapse and debris flows (Tormey 2010). Tormey (2010) reported that Planchon-Peteroa volcano (Chile-Argentina border) had a large slope collapse that produced a highly mobile and erosive debris avalanche 11,000 years BP, and other slope instabilities during the end of the Pleistocene to early Holocene deglaciation. Capra (2006) suggested that most volcanic lateral collapses of ice-capped volcanoes after the last main glacial peaks have been induced by changing climatic conditions. The late Pleistocene and Holocene geological record indicates that several volcanic slope collapses occurred in limited time spans at 1.4–1.3 ka BP, at 1.0–0.9 ka BP, and at ~0.8 ka BP coinciding with several abrupt deglaciations at different latitudes (Clapperton 1998; Capra 2006). Volcanic lateral collapse events in periglacial environments are conditioned by different factors such as 1) shallow

hydrothermal alteration driven by snow and ice melting (Carrasco-Nuñez et al. 1993; Waythomas et al. 1999; Roverato et al. 2011; Roverato and Capra 2013), 2) decreasing rock mass-strength by abrasion and deformation due to glacier advancing (Capra et al. 2013), 3) glacial plucking enhancing volcano weaknesses by exploiting preexisting fractures (Bennett and Glasser, 2009) and 4) increasing pore pressures by high water circulation (Roberti et al. 2017). The high circulation of water in the collapsing mass could also strongly influence the debris avalanche by transforming it into a debris flow-like rheology (Carrasco et al. 1993; Capra et al. 2002; Roverato et al. 2011). Roverato et al. (2015) reported that at the time of the Pungarehu collapse, during the Last Glacial Maximum, Taranaki volcano (New Zealand) was not glaciated, but it did experience periglacial conditions. The snow/ice water-melting coming from the glacier/snowfield or at least a high humid environment might suggest that the pre-avalanche volcanic edifice was partially water-saturated. The increase in precipitation arising from post-glacial warmer, wetter climatic conditions has been considered possible trigger of catastrophic volcanic slope collapses (Capra 2006). This ice/snow could have been an additional supply of water in the collapsed mass during the Pungarehu event permitting the long runout of the avalanche (Roverato et al. 2015). Similar conditions are reported, for example, by Alvarado et al. (2004) concerning volcanic slope collapses in Costa Rica during the last glaciation, for the Teteltzingo collapse at Citlaltepetl volcano (Carrasco-Nuñez et al. 1993) and for the Tonila event at Colima volcano (Roverato et al. 2011). In light of the potential effects of climate change on landslide hazards, it is crucial to monitor potential landslides locations and establish readily available methods for early debris avalanche prediction (Huggel et al. 2008).

3. The instability of volcanic islands

Volcanic lateral collapses have been identified around volcanic islands worldwide and their failure can trigger devastating tsunamis threatening coastal communities and infrastructure (Karstens et al. 2019; Watt et al. 2020 - this volume). The term “volcanic island” is a comprehensive term that includes the main endmembers: oceanic island and arc island. The first ones are gently sloping oceanic shield volcanoes, such the Kilauea (Hawai’i, USA) in the Pacific Ocean or the Piton de La Fournaise (Reunion Island, France) in the Indian Ocean, usually associated with persistent flank movement (Poland et al., 2018). Oceanic island volcanoes, especially in their active shield-building phase, are frequently affected by

large-scale lateral collapses (Moore and Fiske 1969; Elsworth and Day 1999). The arc volcanic islands instead are usually steep-sided composite volcanoes such as Fogo (Cabo Verde) in the Atlantic Ocean, Stromboli (Italy) in the Mediterranean Sea, or Augustine (USA) in the Aleutian arc (Reid et al. 2010), associated with minimal ground deformation at the flank-scale (González et al. 2010), sometimes with strong (volcaniclastic) material remobilization (Di Traglia et al. 2018). Failures of volcanic islands include thin-skinned, narrow collapse, and deep-seated instability events, removing much of an edifice and depositing huge amounts (several km³) of debris (Coombs et al., 2007; Montanaro et al. 2011). They represent the largest formations which can be created in a single geological moment (Blahůt et al. 2018). In addition to the instability and triggering factors that affect volcanoes in all geographical contexts (Voight and Elsworth 1997; Elsworth and Day 1999; Reid et al. 2010), observing that >10 km³-volume collapses occurred at the onset of glacial to interglacial transitions, it suggested that rapid sea level rise enhanced changing of pore pressure conditions, contributing to slope collapse (Quidelleur et al. 2008).

The classification of landslides on volcanic islands usually comprises two endmembers: volcanic debris avalanches, thought to reflect a sudden single catastrophic event, and slumps, thought to reflect progressive slope failure (Blahůt et al. 2018). Slumps affects volcanic flanks with persistent motion and consequent flank deformation, as the Hilina Slumps at Kilauea in Hawai'i (Okubo 2004) and the southeastern flank of the Pico volcanic ridge in the Azores (Hildenbrand et al. 2012). Large VDADs have been observed on the ocean floor surrounding the Hawaiian Islands (Iverson 1995; Morgan et al. 2003), the Canarian Archipelago (Carracedo 1999; Hurlimann et al. 1999), Cape Verde (Elsworth and Day 1999; Ramalho et al., 2015) and Azores (Hildebrand et al. 2018). Some of the debris avalanches may develop by partial or complete collapse of a volcano slope already oversteepened by a slump (Normark et al. 1993). Debris avalanches in arc volcanic islands are typically smaller than those at ocean islands, but they are more frequent (Day et al. 2015). The VDADs around ocean islands and island arc volcanoes have similar slope gradients in the deposition area, mobility and lithology (Karstens et al. 2019).

In historical times, tsunamigenic volcanic debris avalanches have caused many casualties, two of the most devastating cases were: i) the 1792 Unzen–Mayuyama megaslide (volume of 3.4×10^8 m³ and the maximum depth was 400 m) that killed around 15,000 people, and ii) Ritter, Papua New Guinea, in 1888 (mobilized volume of about $4.2 \times$

10⁹ m³; Day et al. 2015 or 2.2-2.6 x 10⁹ m³, Karstens et al. 2019). The slope collapse of Ritter Island is the largest historically recorded landslide at a volcanic island (Cooke 1981; Day et al. 2015; Karstens et al. 2019). The failure produced a headwall width of ~3 km and a fall in vent elevation of ~1,500 m, reducing the 780 m high, 1.5-km-wide island to a small remnant (Johnson, 1987). The landslide-induced tsunami affected islands out to ~600 km from Ritter, killing hundreds of people and producing considerable damage (Cooke 1981; Day et al. 2015; Watt et al. 2019). Recent studies (Karstens et al. 2019; Watt et al. 2019) based on high-resolution 2D and 3D seismic and bathymetric data, provided new insights into the collapse sequence and landslide-induced tsunami generation. The collapse of Ritter occurred in two phases: (1) the volcanic edifice was first affected by deep-seated gradual spreading generating compressional deformation within the volcano and the adjacent seafloor sediments; (2) the failure of about 2.4 km³ of the island and the generation of a highly energetic mass flow. This second phase is inferred to have been deposited up to 80 km from Ritter (Watt et al. 2019). The slide triggered an explosive phase that formed a crater-like depression (Karstens et al. 2019).

More recently on 22 December 2018 the lateral collapse of the Anak Krakatau occurred (Figure 6). The active volcanic island, situated within the Krakatau caldera in the Sunda Strait between Java and Sumatra, experienced a slope collapse after a period of intense volcanic activity (Williams et al. 2019). This event generated tsunami waves ranging in height from 0.27 to 1.40 m, caused the deaths of 431 people, injured a further 7,200 people, and displaced 46,646 people (Williams et al. 2019).

A good example of a tsunamigenic landslide that occurred in recent times, and is supported by many studies, is the one at Stromboli volcano during its 2002–2003 eruption (Tinti et al., 2006). The volcanic island is located in the Tyrrhenian Sea off the southern coast of Italy, and was affected by several mass-wasting events during its geological history, mainly located in the northwest part of the volcano that produced a depression called "Sciara del Fuoco" (Kokelaar and Romagnoli 1995; Tibaldi 2001). During the 2002–2003 flank eruption, a submarine–subaerial landslide sequence (total volume of 25–30 x 10⁶ m³, Bonaccorso et al. 2003; Chiocci et al. 2008) generated two tsunamis. These tsunami waves affected the coastline with a maximum run-up of 6–7 m at the Stromboli village (Tinti et al., 2006) and were caused by two initial submarine rotational slides that affected the

underwater volcanoclastic apron down to a depth of 350 m (Chiocci et al. 2008), as well as by two second subaerial translational slides (Tommasi et al. 2005).

Figure 6

Geological and geotechnical studies (Tommasi et al. 2005; Chiocci et al. 2008; Boldini et al. 2009; Verrucci et al. 2019) suggest that the submarine and sub-aerial slides were caused by large deep-seated gravitational slope deformations (DSGSD) affecting the northern part of the Sciara del Fuoco, triggered by the injection of a lateral intrusion (Bonaccorso et al. 2003). The DSGSD developed under drained conditions, preventing the collapse of the entire block (Chiocci et al. 2008). Contrariwise, during its continuous displacement, excessive pore pressure was likely generated in the submarine volcanoclastic apron (undrained load condition), reducing the effective stresses down to values compatible with the static liquefaction of the material (Tommasi et al. 2005; Boldini et al. 2009).

Another volcano showing strong flank instability is Kilauea, on the Big Island of Hawai'i (USA). Kilauea is a persistently active volcano, partially overlapping the larger neighbour Mauna Loa, located in the middle of the Pacific Plate at the leading edge of a ~6,000-km-long chain of islands and seamounts. The south flank of Kilauea volcano, bracketed by two rift zones that extend from Kilauea's summit caldera, is creeping slowly seaward along a weak, low-angle décollement (Poland et al, 2017. Large scale slope collapse (volumes of 5,000 km³) have been recognized. The Hilina Slump is the most notable of these features (Smith et al. 1999). Major rupture and offset on the southern flank have occurred along faults as recently as 1975, during the M7.2 Kalapana earthquake that struck the south coast of Hawai'i (Okubo 2004; Day et al. 2005). Kilauea's south flank experienced also transient aseismic displacements, as occurred in early November 2000, when an event of about 36 hours moved the flank at a maximum slip velocity of about 6 cm per day (Cervelli et al. 2002). The last transient flank motion occurred on early May 2018 that coincided with the lateral propagation of a dike along the East Rift Zone. The flank slip triggered a moment magnitude (M_w) 7.2 earthquake, producing ~5 meters of fault slip (Chen et al. 2019).

Other examples of failures of volcanic islands can be found in the Lesser Antilles arc, where 52 volcanic flank-collapse events are recognized, including the active volcanoes of Montserrat, Dominica, Martinique, and St. Lucia (Deplus et al. 2001; Boudon et al. 2013).

Large-scale debris avalanche deposits (10-20 km³) lying on the seafloor around the islands, implying that relatively small volcanic islands, as Montserrat, can be affected by repeated large-volume landslides that deposit very large volcanoclastic material into the surrounding basins (Lebas et al. 2011). The largest deposits are related to collapses which include large sectors of the islands, both subaerial and submarine. However, it must be considered that even smaller subaerial landslides, associated with the collapse of lava domes, such as one that occurred on 26 December 1997 (the so-called "Boxing day" collapse; Voight et al. 2002), are extremely dangerous. At Soufrière Hills Volcano on Montserrat, a phase of dome growth started in the 1995. The collapse produced a $\sim 50 \times 10^6$ m³ volcanic debris avalanche, and depressurized the interior of the lava dome, which exploded to generate a violent pyroclastic density current (Voight et al. 2002).

At Réunion Island spreading and avalanching developed in two different sectors (Le Friant et al. 2011). In particular, Réunion island is a 7,000 m-high volcanic complex with a large submarine part and two emerging volcanoes (Piton des Neiges and Piton de la Fournaise; Lénat et al. 1990; Lebas et al. 2018). Investigations by Le Friant et al. (2011) revealed that slow flank deformation is likely one of the main processes acting on the submarine flanks of Piton des Neiges, causing the continuous flank erosion by landslides, and producing turbidity flows (Bachèlery et al. 2003; Le Friant et al. 2011). Conversely, Piton de la Fournaise volcano collapsed several times in the same seaward direction, generating several voluminous debris avalanches that flowed down the submarine flanks. Le Friant et al. (2011) suggested that the difference in behaviour between the two volcanoes is likely related to the volcano basement. Piton des Neiges was built on marine sediments, promoting radial spreading, whereas Piton de la Fournaise was emplaced on the flank of a pre-existing edifice, favouring repeated collapses.

4- Discussion

Several contributing factors can affect the stability of volcanoes. These factors create the conditions that set the volcanic edifice to "ready to collapse". According to classifications proposed by previous authors (Gorshkov, 1965; Moriya 1980; Siebert et al. 1987) three main failure categories exist: Bezymianny, Bandai, and Unzen/Ontake types. The difference between them is mainly based on the presence of a magmatic or phreatic explosive activity (Bezymianny and Bandai) or an earthquake as the trigger (Unzen/Ontake). The Bezymianny-

type collapse takes its name from the event that occurred in 1957 at Bezymianny Volcano (Kamchatka), which was mainly related to a magmatic activity (Gorshkov, 1959; Bogoyavlenskaya and Kirnasov 1981). This type of activity corresponds to the destabilization of the volcanic edifice by magmatic intrusions and followed by magmatic eruptions. The edifice undergoes strong deformation due to the emplacement of cryptodomes; the same conditions that occurred at Mount St. Helens. The eruptive phase may contain highly energetic lateral blasts (Belousov et al. 2004) with associated pyroclastic density currents. The lateral blasts of Mount St. Helens and Bezymianny produced major pyroclastic density currents that devastated areas of 500-600 km² around each of the volcanoes (Crandell and Hoblitt 1986; Siebert et al. 1987). The Bezymianny activity is often followed by construction of a lava dome or a new pyroclastic cone within the collapse scarp (Siebert et al. 1987; Leyrit 2000). In the case of Bezymianny volcano, the duration and magnitude of deformation, as well as the presence of a flat lava dome in the crater prior to failure and magmatic eruptions suggest that the intrusion was well developed. Other cryptodome-induced deformations have been found at Showa-Shinzan and Usu-Shinzan (Katsui et al. 1985) of Usu volcano (Minakami et al. 1951) in Japan. They did not culminate in a debris avalanche associated with a paroxysmal eruption like at Mount St. Helens or Bezymianny (Donnadieu et al. 2001). Bandai activity (Moriya 1980) refers to the explosion that occurred in 1888 at the Bandai-San volcano (Japan). It is associated with phreatic explosions that favor the collapse of the volcanic edifice without subsequent magmatic activity. It means that no juvenile material is present neither within the debris avalanche deposit nor at the base or top of it. Massive VDADs can also occur without related explosive eruptions, as at the Unzen volcano in 1792 or at Ontake volcano in 1984, where earthquake triggered the collapse. The potential of a seismically-triggered volcanic edifice collapse to initiate a volcanic debris avalanche has commonly been recognized (Voight and Elsworth, 1997; McGuire, 2003). Edifice collapse and flank instability-derived deposits (VDADs) are now recognized as ubiquitous events associated with active and growing volcanoes, yet the identification of the triggering mechanisms is poorly correlated due to difficulties that lie in elucidating diagnostic lithological and sedimentological features that record evidence of the trigger. Kataoka et al. (2018) and Fairchild (1987) both highlight the importance of understanding the triggering mechanisms of a landslide to then understand the emplacement mechanisms and resulting hazard. Only in some cases such as Unzen-Mayuyama (1792), Bandai (1888), Bezymianny

(1956), Shiveluch (1964), Mount St. Helens (1980), and Ontake (1984) were observations of the events well-constrained enough that linkages could be made between initiation, trigger and flow through to emplacement and deposition. In the cases such as Unzen-Mayuyama (1792) and Ontake (1984) earthquakes were directly attributed to initiating the VDA. The 1792 Unzen-Mayuyama VDA is remembered as the trigger for a tsunami that impacted the Ariake Sea, killing ~14,500 people (Watanabe et al. 1993; Miyamoto 2010). Wang et al. (2019) states that from November 1791 to August 1792 earthquakes, eruptions and flows were recorded from various sites around Mt. Unzen. On the 21 May 1792 a magnitude 6.4 earthquake triggered a volcanic debris avalanche with an associated tsunami. Usami (1996) estimated that the earthquake which triggered a $3.4\text{--}4.8 \times 10^8 \text{ m}^3$ volcanic debris avalanche (Ui et al. 1986; Inoue 2000) was likely to have been of magnitude = 6.4 ± 0.2 . This was estimated based on record that more than 30% of buildings in the Shimabara area were destroyed and ground ruptures were recorded which requires seismic accelerations of 400 cm/s^2 or more (Sassa et al. 2014). Sassa et al. (2014) also reported that ground accelerations of between 370 to 507 cm/s^2 were required from an earthquake to trigger a landslide of large enough volume to generate a tsunami of the size recorded. Although the geological and volcanological relationship between other volcanoes of the Unzen volcanic complex and Mayuyama is difficult to discern, the Mayuyama volcano or lava dome is considered to be 5.1 ± 1.5 ka old (Hoshizumi et al., 1999) with the last recorded coeval activity at Unzen being a block-and-ash-flow at 4.2 ± 1.1 ka B.P. Therefore, eruptive activity or intrusion into the dome was not a contributing factor in triggering the debris avalanche sourced from older brittle and dacitic volcanic rock. Another well-known earthquake-triggered edifice collapse is Ontake type, defined from the 1984 events from Ontake-San volcano. On 14 September 1984 the southeast slope of Ontake-San collapsed producing a $32\text{--}36 \times 10^6 \text{ m}^3$ avalanche (Voight and Sousa 1994). The last previous eruptive event at this volcano was in 1979 and since then the volcano had been dormant with the 1984 debris avalanche event being triggered by a magnitude 6.8 earthquake centered at 9-10 km southeast (Endo et al. 1989). The source rock was highly saturated andesite with interbedded or intercalated pyroclasts (Endo et al. 1989). Earthquake activity in this region is strongly aligned with regional stress fields, yet Sano et al. (1998) suggest some relationship with deeper-seated magmatic processes. Whereas Voight and Sousa (1994) focused on modeling the flowing mass and constraining parameters that increase mobility, little attention was paid to modeling the

initiation of the collapsing mass or fragmentation of that mass from the dynamic loading of the earthquake.

The eruption from Mount St. Helens in 1980 is probably one of the most well-described and documented events prior to, during and post- a VDA. Lipman and Mullineaux (1982) and Christiansen et al. (1981) provide the chronology of the event identifying that the magnitude 5.1 seismic event (Glicken 1996) triggered a series of events that lead to a VDA. Endo et al. (1981) also identified that the magnitude 5+ earthquake had a hypocenter depth of 1.5 km. The series of events (with respect to the VDA) recorded from eyewitness accounts progress from the earthquake with avalanching of rocks from crater walls through to “quivering” of the flank, fracturing and then slope failure. From the analysis of photos, it is estimated that there were 7-20 seconds between the earthquake and the initiation of the collapse (Voight et al., 1981). It is important to note that seismicity increased from 20 March with seismicity being magnitude 4 or less considered to be confined to 2.5 km depth or less. Dzurisin (2018) noted that by 27 March 1980, magma had intruded the edifice with associated deformation being recorded. Voight et al. (1981) further postulated that the failure was due to several cumulative facts, including progressive strength-loss prior to slip, seismicity and gravitational movements, and increased pore pressure within rock units.

How earthquakes change the physical properties of the volcanic rock mass is not well defined yet. The complexity of a volcano structure including its lithological layering, variability of rock strengths, constructional boundaries, faulting, water storage, hydrothermal activity, and others make the task of applying simple geotechnical analysis difficult let alone introducing seismicity into the equation. The mechanisms that cause a lateral collapse are in general cumulative or inter-related (Voight and Elsworth 1997) and therefore it is difficult to identify a single initiation mechanism or trigger to edifice collapse which would in turn negate the previous oversimplification of the Bezymianny, Bandai, and Unzen/Ontake type of classifications previously used. It thus requires a more comprehensive classification of edifice collapse based on a range of pre-conditioning and triggering factors.

Voight and Elsworth (1997) identified that while the inherent properties of volcanic rocks and the structure of the volcanic edifice are factors that contribute to volcano instability, the driving factors that lead to collapse are generally changes in shear stress on the rock mass. An important associated factor that changes the shear stress is variation in

pore fluid pressure. In terms of seismicity, earthquake shaking can increase pore fluid pressure in a saturated material. Modelling or simulating these types of relationships and processes within a volcanic edifice is difficult in particular on large stratovolcanoes where rock layers are massive and varying in thickness and rock strength is juxtaposed in a multitude of positions related to the evolution of that edifice over hundred or even thousands of years. Geotechnical analyses of the volcanic edifices have been undertaken around the world with success in identifying inherent instabilities in a variety of volcano types (e.g. Zimbelman et al. 2004; Moon et al. 2005; del Porto et al. 2013; Apuani et al. 2013).

Deutschle (2013) provided a comprehensive study of examining a range of triggers that contribute to flank instabilities on Mt. Taranaki in New Zealand. This complex stratovolcano has evolved over ~150,000 years and undergone up to 14 collapse events and regrowth resulting in a structurally and lithologically complex edifice (Zernack et al. 2012). Numerous limit equilibrium and finite element simulations that involved investigating rock strength variability due to changes in basement and edifice conditions, edifice growth, and magmatic intrusion show that while weaker areas of unconsolidated pyroclastic material exist on the current cone, it is still, in general considered stable unless magmatic intrusion or significant seismic shaking (>0.3 g) occurs (Deutschle 2013). Della Pasqua et al. (2016) undertook a similar analysis to understand the hazards posed by VDAs in the Taranaki region and found that susceptibility to failure will increase when there is higher seismic loads on weak, stratified layers. Alloway et al. (2005) identified a number of faults that traverse the Mt. Taranaki edifice and reconstructed earthquakes of magnitude 6.7-7.2 with shaking intensities of magnitude >8 with a recurrence interval of 4 ka. Keefer (1984) suggests that rock and soil avalanches require strong shaking intensities of magnitude $>7-8$. It is conceivable that a number of Mt. Taranaki VDADs were initiated by ground shaking associated with movements on those faults, but no conclusive evidence is presented in the literature. Earthquakes triggering VDAs is well recorded and while the triggering effects of the ground shaking to induce collapse have been elucidated from landslide investigations through understanding the physical properties of soils and rock, the initiation of the mass that suddenly loses its stability and move downslope is not well constrained or modelled accurately for volcanic edifices.

As we can see, most of the information available in the literature regarding eruptions and earthquakes, and their role in sector collapses, arises from historical cases or is inferred by theoretical models and simulations. However, if we take a look at the statistics of real cases, there is not much pre-historic information available about the triggering mechanisms that promote a sector collapse. Dufresne et al. (2020 - this volume) reports that only 16% of VDADs in a new global database of 1001 events have a related triggering mechanism; most of them are associated with magmatic activities. It is obvious that it proves difficult to associate a volcanic collapse with a specific trigger. However, some information on factors and processes involved in past collapses is recorded in the resulting deposits. In Table 3, we summarize key features that could be useful for studying VDADs in the field by providing information on the state of the pre-collapse volcanic edifice and the processes that acted before or during a sector collapse.

Table 3

Efforts to couple lateral collapse models and flow models as proposed in Voight and Sousa (1994) need to be further developed. Being able to transfer those simulations and predictions into a hazard and risk analysis framework (e.g. Mead et al. 2018) to provide a more holistic analysis of the event is also required considering that much of the motivation for understanding volcanic debris avalanches arose from better understanding the hazardous events that occurred over the last 300 years.

5- Conclusions

Volcanic sector collapses can be prepared by a wide variety of destabilizing factors, which are followed by triggering elements that drive the collapse (Figure 7). Although we group these factors in distinctive sections, it is evident that most of them are inherently related. In fact, factors that prepare and trigger volcanic sector collapses are not simple but can be multiple and cumulative. Over-steepened slopes, magma intrusions, hydrothermal activity, climate fluctuations, deformation of the basement, eruptions, earthquakes, are just a list of elements that could promote a volcanic collapse. Although it is necessary to identify the differences between those factors that define the degree of instability from those mechanisms that trigger the collapse, it is also important to define those factors that, in

combination, could lead to different catastrophic scenarios. Indeed, the mechanism that triggers a sector collapse could be independent of the causes that prepare its instability. But it is also true that multiple factors in consort could drastically enhance the power and magnitude of the collapse event and affect its runout.

Figure 7

The intrusion of a magma body (e.g. cryptodome) into a volcanic edifice, for example, has multiple consequences for the stability of the volcano. On one hand, the magma body deforms the edifice structure and, if the magma reaches the surface, its products could overload the volcanic flanks (destabilizing factor). On the other hand, the hydrothermal fluids associated with the magma body could promote all those processes of rock leaching and dissolution that weaken the volcanic material. If the intrusion of a magma body, linked with an earthquake or volcanic tremors, could be considered a factor that induce a volcanic collapse (trigger), clay mineral formation and migration due to the presence of magmatic fluids is an important element both for the loss of rock strength (destabilizing factor) as well as for the increase of impermeability that promote fluids and water (e.g. meteoric) storage. Once the collapse occurred, the presence of fluids and water, and the clay content as well, can play a fundamental role in the transport and mobility of the avalanche body, enhancing its mobility and transforming the moving mass into a cohesive lahar. Thus, considering all the endogenous and exogenous factors, a volcanic sector collapse and debris avalanche motion depend on an intricate system of factors that promote volcano instability, trigger the collapse, and enhance the avalanche mobility. It is therefore important to understand all the antecedent factors and triggers of volcano instability since that knowledge can provide critical precursory information that can directly inform hazard warnings.

Acknowledgments

We acknowledge Marc-André Brideau and Lucia Capra reviewers for their important suggestions and corrections that helped to notably improve the manuscript.

References

- Alloway B, McComb P, Neall V, Vucetich C, Gibb J, Sherburn S, Stirling M (2005) Stratigraphy, age, and correlation of voluminous debris-avalanche events from an ancestral Egmont Volcano: implications for coastal plain construction and regional hazard assessment. *J Roy Soc New Zealand* 35(1-2): 229-267
- Alvarado GE, Veg E, Chaves J, Vasquez M (2004) Los grandes deslizamientos (volcánicos y no-volcánicos) de tipo debris avalanche en Costa Rica. *Rev Geo de Am Centr* 30:83-99
- Andrade SD, van Wyk de Vries B (2010) Structural analysis of the early stages of catastrophic stratovolcano flank-collapse using analogue models. *Bull Vulcanol* 72:771-789
- Apuani T, Corazzato C, Merri A, Tibaldi A (2013) Understanding Etna flank instability through numerical models. *J Vol Geotherm Res* 251: 112-126
- Bachèlery P, Robineau B, Courteaud M, Savin C (2003) Debris avalanches on the western flank of Piton des Neiges shield volcano (Reunion Island). *Bull Soc Géo de France* 174(2): 125-140
- Bahar I, Girod M (1983) Controle Structural Du Volcanisme Indonésie (Sumatra, Java-Bali); Application et Critique de La Methode de Nakamura." *Bull Soc Géo de France XXV* 7: 609-14
- Begét JE, Kienle J (1992) Cyclic formation of debris avalanches at Mt. St. Augustine volcano, Alaska. *Nature* 356:701-704
- Belousov A, Belousova M, Voight B (1999) Multiple edifice failures, debris avalanches and associated eruptions in the Holocene history of Shiveluch volcano, Kamchatka, Russia. *Bull Vulcanol* 61(5):324-342
- Belousov A, Voight B, Belousova M (2004) Directed blasts and blast-generated pyroclastic density currents: a comparison of the Bezymianny 1956, Mount St Helens 1980, and Soufrière Hills, Montserrat 1997 eruptions and deposits. *Bull Vulcanol* 69:701-740
- Bennett MR, Glasser NF (2009) *Glacial Geology. Ice Sheets and Landforms*. Wiley- Blackwell 385
- Blahůt J, Klimeš J, Rowberry M, Kusák M (2018) Database of giant landslides on volcanic islands—first results from the Atlantic Ocean. *Landslides* 15(4): 823-827
- Bogoyavlenskaya GE, Kirsanov IT (1981) Twenty-five years of activity of Bezymianny Volcano *Vulkanologiya i Seismologiya* 2:3-13
- Boldini D, Wang F, Sassa K, Tommasi P (2009) Application of large-scale ring shear tests to the analysis of tsunamigenic landslides at the Stromboli volcano, Italy. *Landslides* 6(3): 231-240
- Bonaccorso A, Calvari S, Garfi G, Lodato L, Patanè D (2003) Dynamics of the December 2002 flank failure and tsunami at Stromboli volcano inferred by volcanological and geophysical observations. *Geo Res Letters* 30(18): 1941.
- Bonforte A, Guglielmino F, Coltelli M, Ferretti A, Puglisi G (2011) Structural assessment of Mount Etna volcano from Permanent Scatterers analysis. *Geoch, Geophy, Geosys* 12(2): Q02002.
- Bonforte A, Guglielmino F, Puglisi G (2019) Large dyke intrusion and small eruption: The December 24, 2018 Mt. Etna eruption imaged by Sentinel-1 data. *Terra Nova* 31 (4): 405-412
- Borgia A, Delaney PT, Denlinger RP (2002) Spreading Volcanoes. *Annu Rev Earth Planet Sci* 2000 28:539-70
- Borgia A, Ferrari L, Pasquarè G (1992) Importance of gravitational spreading in the tectonic and volcanic evolution of Mount Etna. *Nature* 357:231-235
- Borselli L, Capra L, Sarocchi D, De la Cruz-Reyna S (2011) Flank collapse scenario at Volcan de Colima, Mexico: a relative instability analysis. *J Vol Geotherm Res* 208:51-65
- Boudon, G., Villemant, B., Le Friant, A., Paterne, M., & Cortijo, E. (2013). Role of large flank-collapse events on magma evolution of volcanoes. Insights from the Lesser Antilles Arc. *J Vol Geotherm Res* 263: 224-237
- Capra (2006) Abrupt climatic changes as triggering mechanism of massive volcanic collapses. *J Vol Geotherm Res* 155:329-333
- Capra L, Bernal JP, Carrasco-Nuñez G, Roverato M (2013) Climatic fluctuations as a significant contributing factor for volcanic collapses. Evidence from Mexico during the Late Pleistocene.

918 Glob and Plan Change 100:194-203
 919 Capra L, Macias JL, Scott KM, Abrams M, Garduño-Monroy VH (2002) Debris avalanche and debris
 920 flow transformed from collapses in the Trans-Mexican Volcanic Belt, Mexico - behavior, and
 921 implication for hazard assessment. J Vol Geotherm Res 113:81-110
 922 Capra L, Roverato M, Groppelli G, Caballero L, Sulpizio R, Norini G (2015) Glacier melting during lava
 923 dome growth at Nevado de Toluca volcano (Mexico): Evidences of a major threat before main
 924 eruptive phases at ice-caped volcanoes. J Vol Geotherm Res 294:1-10
 925 Carracedo JC (1994) The Canary Islands : An Example of Structural Control on the Growth of Large
 926 Oceanic-Island Volcanoes. J Vol Geotherm Res 60: 225-41
 927 Carracedo JC (1999) Growth, structure, instability and collapse of Canarian volcanoes and
 928 comparisons with Hawaiian volcanoes. J Vol Geotherm Res 94(1-4): 1-19
 929 Carrasco-Núñez G, Díaz-Castellón R, Siebert L, Hubbard R, Sheridan MF, Rodríguez SR (2006).
 930 Multiple edifice-collapse events in the Eastern Mexican Volcanic Belt: The role of sloping
 931 substrate and implications for hazard assessment. J Volcanol Geotherm Res 158:151–176
 932 Carrasco-Núñez G, Vallance JW, Rose WI (1993) A voluminous avalanche-induced lahar from
 933 Citlaltépetl volcano, Mexico: implications for hazard assessment. J Volcanol Geotherm Res 59(1-
 934 2):35-46
 935 Cecchi E, van Wyk de Vries B, Lavest JM (2004) Flank spreading and collapse of weak-cored
 936 volcanoes. Bull Volcanol 67(1):72-91
 937 Cervelli P, Segall P, Johnson K, Lisowski M, Miklius A (2002) Sudden aseismic fault slip on the south
 938 flank of Kilauea volcano. Nature 415(6875): 1014
 939 Chen K, Smith JD, Avouac JP, Liu Z, Song YT, Gualandi A (2019) Triggering of the Mw 7.2 Hawaii
 940 earthquake of May 4, 2018 by a dike intrusion. Geoph Res Lett 46(5): 2503-2510
 941 Chiocci FL, Romagnoli C, Tommasi P, Bosman A (2008) The Stromboli 2002 tsunamigenic submarine
 942 slide: characteristics and possible failure mechanisms. Journal of Geophysical Research: S Earth
 943 113(B10): B10102.
 944 Christiansen RL, Peterson DW, Lipman PW, Mullineaux DR (1981) Chronology of the 1980 eruptive
 945 activity. US Geol. Surv. Prof. Pap 1250: 17-30
 946 Clapperton CM (1998) Late Quaternary glacier fluctuations in the Andes: testing the synchrony of
 947 global change. In: Owen LA (Ed.), Mountain Glaciation. Quaternary Proceedings, 6. John Wiley
 948 & Sons, Chichester, 65-73.
 949 Clavero JE, Sparks RSJ, Huppert HE (2002) Geological constraints on the emplacement mechanism of
 950 the Paríncota avalanche, northern Chile. Bull Volcanol 64:40-54
 951 Coe JA, Bessette-Kirton EK, Geertsema M (2018) Increasing rock-avalanche size and mobility in
 952 Glacier Bay National Park and Preserve, Alaska detected from 1984 to 2016 Landsat imagery.
 953 Landslides 15: 393
 954 Cooke RJ (1981) Eruptive history of the volcano at Ritter Island. Geol Surv Papua New Guinea Mem,
 955 10: 115-123
 956 Coombs ML, White SM, Scholl DW (2007) Massive edifice failure at Aleutian arc volcanoes. Earth and
 957 Plan Scie Lett 256(3-4): 403-418
 958 Crandell DR, Hoblitt RP (2006) Lateral blasts at Mount St. Helens and hazard zonation. Bull Volcanol
 959 48:27-37 Day SJ (1996) Hydrothermal pore fluid pressure and the stability of porous, permeable
 960 volcanoes. In: McGuire, W.J., Neuberg, J., Jones, A. (Eds), Volcano Instability on the Earth and
 961 Terrestrial Planets. Geol Soc London Spec Publ 110:77-93
 962 Day SJ, Llanes P, Silver E, Hoffmann G, Ward S, Driscoll N (2015) Submarine landslide deposits of the
 963 historical lateral collapse of Ritter Island, Papua New Guinea. Marine and Petroleum Geology
 964 67: 419-438
 965 Day SJ, Watts P, Grilli ST, Kirby JT (2005) Mechanical models of the 1975 Kalapana, Hawaii
 966 earthquake and tsunamis. Marine Geology 215(1-2): 59-92
 967 Delaney PT, Pollard DD, Ziony JI, McKee EH (1986) Field Relations between dikes and joints:
 968 Emplacement processes and paleostress analysis. Journal of Geophysical Research 91(B5): 4920

969 Delcamp A, van Wyk de Vries B, James MR (2008) The influence of the edifice slope and substrata on
 970 volcano spreading. *J Vol Geotherm Res* 177:925-943
 971 Delmelle P, Bernard A (2015) The Remarkable Chemistry of Sulfur in Hyper-Acid Crater Lakes: A
 972 Scientific Tribute to Bokuichiro Takano and Minoru Kusakabe, in Rouwet D, Christenson B, Tassi
 973 F, Vandemeulebrouck J (Eds) *Volcanic Lakes, Advances in Volcanology*, Springer
 974 del Potro R, Hürlimann M, Pinkerton H (2013) Modelling flank instabilities on stratovolcanoes:
 975 parameter sensitivity and stability analyses of Teide, Tenerife. *J Vol Geotherm Res* 256: 50-60
 976 della Pasqua F, Massey CI, McSaveney M, Townsend D (2016). Preliminary assessment of some flank-
 977 failure scenarios for Mount Taranaki and recommendations for future assessment of the risk
 978 from such hazards, GNS Science Report 2015/54: 48
 979 Deplus C, Le Friant A, Boudon G, Komorowski JC, Villemant B, Harford C, Cheminée JL (2001)
 980 Submarine evidence for large-scale debris avalanches in the Lesser Antilles Arc. *Earth and Plan*
 981 *Sci Lett* 192(2): 145-157
 982 Deutschle OK (2013) Characterisation of geotechnical units on Mount Taranaki and Influence on
 983 Edifice Stability (Masters dissertation, University of Auckland).
 984 Dufresne A, Siebert L, Bernard B (2020) Distribution and geometric parameters of volcanic debris
 985 avalanche deposits, in Roverato M, Dufresne A, Procter J (Eds), *Volcanic Debris Avalanches:*
 986 *From Collapse to Hazard, Advances in Volcanology*, Springer
 987 Di Traglia F, Nolesini T, Ciampalini A, Solari L, Frodella W, Bellotti F, Fumagalli A, De Rosa G, Casagli,
 988 N (2018a) Tracking morphological changes and slope instability using spaceborne and ground-
 989 based SAR data. *Geomorphology* 300:95-112
 990 Donnadieu F, Merle O (1998) Experiments on the indentation process during cryptodome intrusions:
 991 New insights into Mount St. Helens deformation. *Geology* 26:79-82
 992 Donnadieu F, Merle O, Besson JC (2001) Volcanic edifice stability during cryptodome intrusion. *Bull*
 993 *Volcanol* 63:61-72
 994 Dzurisin D (2018) Mount St. Helens retrospective: Lessons learned since 1980 and remaining
 995 challenges. *Front in Earth Sci* 6: 142
 996 Endo ET, Malone SD, Noson LL, Weaver CS (1981) Locations, magnitudes, and statistics of the March
 997 20–May 18 earthquake sequence. In *The 1980 Eruptions of Mount St. Helens, Washington*
 998 (1250: 93-107). US Department of the Interior.
 999 Endo K, Sumita M, Machida M, Furuichi M (1989) The 1984 collapse and debris avalanche deposits of
 1000 Ontake Volcano, central Japan. In *Volcanic Hazards* (pp. 210-229). Springer, Berlin, Heidelberg.
 1001 Elsworth D, Voight B (1995) Dike intrusion as a trigger for large earthquakes and the failure of
 1002 volcano flanks. *J Geo-phys Res* 100(B4):6005-6024
 1003 Elsworth D, Voight B (1996) Evaluation of volcano flank instability triggered by dyke intrusion.
 1004 Geological Society, London, Special Publications 110 (1): 45-53
 1005 Elsworth D, Day SJ (1999) Flank collapse triggered by intrusion: the Canarian and Cape Verde
 1006 Archipelagoes. *J Vol Geotherm Res* 94(1-4): 323-340
 1007 Fairchild LH (1987) The importance of lahar initiation processes. Debris flows/avalanches: process,
 1008 recognition, and mitigation: *Geo S of Am Rev in Eng Geo* 7: 51-61
 1009 Francis PW, Self S (1987) Collapsing volcanoes. *Scientific American* 256 (6): 90-99
 1010 Francis PW, Wells GL (1988) Landsat Thematic Mapper observations of debris avalanche deposits in
 1011 the Central Andes. *Bull Volcanol* 50: 258-78
 1012 Francis PW (1994) Large volcanic debris avalanches in the central Andes. In: *Proceedings of*
 1013 *International Conference on Volcano Instability on the Earth and Other Planets*. Geol Soc
 1014 London.
 1015 Frank D (1995) Surficial extent and conceptual model of hydrothermal system at Mount Rainier,
 1016 Washington. *J Vol Geotherm Res* 65:51-80
 1017 Frantz JD, Popp RK, Boctor NZ (1981) Mineral-solution equilibria - V. Solubilities of rock forming
 1018 minerals in supercritical water. *Geoch Cosm Acta* 45:69-77
 1019 Glicken H (1996) Rockslide-debris avalanche of May 18, 1980, Mount St. Helens volcano, Washington

(No. 96-677). US Geological Survey Open File Report.

González PJ, Tiampo KF, Camacho AG, Fernández J (2010) Shallow flank deformation at Cumbre Vieja volcano (Canary Islands): Implications on the stability of steep-sided volcano flanks at oceanic islands. *Earth Plan Sci Lett* 297(3-4):545-557

Gorshkov GS (1959) Gigantic eruption of the volcano Bezymianny. *Bull Volcanol* 21:77-109

Gorshkov GS, Bogoyavlenskaya GE (1965) Bezymianny volcano and peculiarities of its last eruption (1955–1963). Moscow, Nauka (in Russian)

Grämiger LM, Moore JR, Gischig VS, Ivy-Ochs S, Loew S (2017) Beyond debuttressing: mechanics of paraglacial rock slope damage during repeat glacial cycles. *J Geophys Res Earth Surf* 122(4):1004-1036

Hibert C, Michéa D, Provost F, Malet JP, Geertsema M (2019) Exploration of continuous seismic recordings with a machine learning approach to document 20 years of landslide activity in Alaska. *Geophys J Intern* 219:1138-1147

Hildenbrand A, Marques FO, Catalão J (2018) Large-scale mass wasting on small volcanic islands revealed by the study of Flores Island (Azores). *Scientific reports* 8(1):13898

Hildenbrand A, Marques FO, Catalão J, Catita CMS, Costa ACG (2012) Large-scale active slump of the southeastern flank of Pico Island, Azores. *Geology* 40(10):939-942

Hoshizumi H, Uto K, Watanabe K (1999) Geology and eruptive history of Unzen volcano, Shimabara peninsula, Kyushu, SW Japan. *J Vol Geotherm Res* 89(1-4):81-94

Huggel C, Caplan-Auerbach J, Wessels R (2008) Recent extreme avalanches: triggered by climate change? *Eos* 89(47):469-470

Huggel C, Caplan-Auerbach J, Waythomas CF, Wessels RL (2007) Monitoring and modeling ice-rock avalanches from ice-capped volcanoes: A case study of frequent large avalanches on Iliamna Volcano, Alaska. *J Vol Geotherm Res* 168:114-136

Hürlimann M, Ledesma A, Martí J (1999) Conditions favouring catastrophic landslides on Tenerife (Canary Islands). *Terra Nova* 11(2-3): 106-111

Inoue K (2000) Shimabara-Shigatusaku Earthquake and topographic changes by Shimabara Catastrophe in 1792

Inokuchi (1988) Gigantic landslides and debris avalanches on volcanoes in Japan: case studies on Bandai, Chokai and Iwate Volcanoes. *Rep Nat Res Center Disaster Prevention* 41:163–275 (in Japanese with English abstract)

Iverson RM (1995) Can magma-injection and groundwater forces cause massive landslides on Hawaiian volcanoes? *J Vol Geotherm Res* 66(1-4):295-308

John DA, Sisson TW, Breit GN, Rye RO, Vallance JW (2008) Characteristics, extent and origin of hydrothermal alteration at Mount Rainier Volcano, Cascades Arc, USA: Implications for debris-flow hazards and mineral deposits. *J Vol Geotherm Res* 175:289-314

Johnson RW (1987) Large-scale volcanic cone collapse: the 1888 slope failure of Ritter volcano, and other examples from Papua New Guinea. *Bull Vol* 49(5):669-679

Karstens J, Berndt C, Urlaub M, Watt SF, Micallef A, Ray M, Klauke I, Muff S, Klaeschen D, Kühn M, Roth T, Böttner C, Schramm B, Elger J, Brune S (2019) From gradual spreading to catastrophic collapse—Reconstruction of the 1888 Ritter Island volcanic sector collapse from high-resolution 3D seismic data. *Earth Plan S Lett* 517:1-13

Kataoka KS, Matsumoto T, Saito T, Kawashima K, Nagahashi Y, Iyobe T, Suzuki K (2018) Lahar characteristics as a function of triggering mechanism at a seasonally snow-clad volcano: contrasting lahars following the 2014 phreatic eruption of Ontake Volcano, Japan. *Earth, Planets and Space* 70(1): 113

Katsui Y, Komuro H, Uda T (1985) Development of faults and growth of Usu-Shinzan cryptodome in 1977–1982 at Usu volcano, north Japan. *J Fac Sci Hokkaido Univ, Ser IV* 21:339-362

Keefer DK (1984) Landslides caused by earthquakes. *G S A Bull* 95(4): 406-421

Kerle N, van Wyk de Vries B (2001) The 1998 debris avalanche at Casita volcano, Nicaragua - investigation of structural deformation as the cause of slope instability using remote sensing. *J*

1071 Vol Geotherm Res 105(1-2):49-63
 1072 Kokelaar P, Romagnoli C (1995) Sector collapse, sedimentation and clast population evolution at an
 1073 active island-arc volcano: Stromboli, Italy. Bull Volcan 57(4):240-262
 1074 Kos A, Amann F, Strozzi T, Delaloye R, von Ruetten J, Springman S (2016) Contemporary glacier retreat
 1075 triggers a rapid landslide response, Great Aletsch Glacier, Switzerland. Geophys Res Lett
 1076 43(24):12466-12474
 1077 Kuhn D (2002) Fold and thrust belt structures and strike-slip faulting at the SE margin of the Salar de
 1078 Atacama basin, Chilean Andes. Tectonics 21:1026
 1079 Lagmay AMF, Valdivia W (2006) Regional stress influence on the opening direction of crater
 1080 amphitheatres in Southeast Asian volcanoes. J Vol Geotherm Res 158 (1-2): 139-50
 1081 Lagmay AMF, van Wyk de Vries B, Kerle N, Pyle DM (2000) Volcano instability induced by strike-slip
 1082 faulting. Bull Volc 62 (4-5): 331-46
 1083 Lebas E, Le Friant A, Boudon G, Watt SFL, Talling PJ, Feuillet N, Deplus C, Berndt C, Vardy ME (2011)
 1084 Multiple widespread landslides during the long-term evolution of a volcanic island: Insights
 1085 from high-resolution seismic data, Montserrat, Lesser Antilles. Geoch, Geophys, Geosyst 12(5):
 1086 Q05006.
 1087 Lebas E, Le Friant A, Deplus C, de Voogd B (2018) Understanding the Evolution of an Oceanic
 1088 Intraplate Volcano from Seismic Reflection Data: A New Model for La Réunion, Indian Ocean.
 1089 Journal of Geophysical Research: Solid Earth 123(2):1035-1059
 1090 Le Friant A, Lebas E, Clément V, Boudon G, Deplus C, De Voogd B, Bachèlery P (2011) A new model
 1091 for the evolution of La Réunion volcanic complex from complete marine geophysical surveys.
 1092 Geophys Res Lett 38(9): L09312.
 1093 Lénat JF, Bachèlery P, Bonneville A, Galdéano A, Labazuy P, Rousset D, Vincent P (1990) Structure
 1094 and morphology of the submarine flank of an active basaltic volcano: Piton de la Fournaise
 1095 (Réunion Island, Indian Ocean). Oceanol. Acta 10:211-223
 1096 Leyrit H (2000) Flank collapse and debris avalanche deposits. In: Volcaniclastic Rocks, from Magmas
 1097 to Sediments. Edited by H. Leyrit and C. Montenat 111-129
 1098 Lipman PW, Mullineaux DR, (eds) (1981) The 1980 Eruptions of Mount St Helens, Washington. U.S.
 1099 Geological Survey, Professional Paper 1250. 844 pp.
 1100 Lipman PW, Moore JG, Swanson DA (1981) Bulging of the north flank before the May 18 eruption-
 1101 Geodetic data, in Lipman PW, Mullineaux, DR (Eds), The 1980 eruptions of Mount St. Helens,
 1102 Washington: U.S. Geological Survey Professional Paper 1250-143-155
 1103 Lopez DL, Williams SN (1993) Catastrophic volcanic collapse: Relation to hydrothermal processes.
 1104 Science 260:1794-1795
 1105 Lundgren P, Casu F, Manzo M, Pepe A, Berardino P, Sansosti E, Lanari R (2004) Gravity and magma
 1106 induced spreading of Mount Etna volcano revealed by satellite radar interferometry. Geophys
 1107 Res Lett 31(4)
 1108 Macias JL, Arce JL, Garcia-Palomo A, Mora JC, Lauer PW, Espindola JM (2010) Late-Pleistocene flank
 1109 collapse triggered by dome growth at Tacaná volcano, México-Guatemala, and its relationship
 1110 to the regional stress regime. Bull Volcanol 72:33-53
 1111 MacLeod N (1989) Sector-failure eruptions in Indonesian volcanoes. Geol Indonesia 12:563-601
 1112 McClelland E, Erwin PS (2003) Was a dacite dome implicated in the 9,500 B.P. collapse of Mt
 1113 Ruapehu? A palaeomagnetic investigation. Bull Volc 65:294-305
 1114 McGuire WJ (1996) Volcano instability: a review of contemporary themes. Geo Soc, London, Special
 1115 Publications 110(1):1-23
 1116 McGuire WJ (2003) Volcano instability and lateral collapse. Revista 1: 33-45
 1117 McGuire WJ (2006) Lateral collapse and tsunamigenic potential of marine volcanoes, in Troise C, De
 1118 Natale G, Kilburn CRJ (eds), Mechanisms of Activity and Unrest at Large Calderas. Geo Soc,
 1119 London, Special Publications 269:121-140.
 1120 Mead S, Bebbington M, Procter J (2018). Emulation of the relationship between hazard intensity and
 1121 volcanic processes. In EGU General Assembly Conference Abstracts (Vol. 20, p. 11000).

1122 Merle O, Borgia A (1996) Scaled experiments of volcanic spreading. *J Geophys Res: Solid Earth*
 1123 101(B6):13805-13817
 1124 Merle O, Lenat JF (2004) Hybrid collapse mechanism at Piton de la Fournaise volcano, Reunion
 1125 Island, Indian Ocean. *J Geophys Res* 108(B3):2166
 1126 Minakami T, Ishikawa T, Yagi K (1951) The 1944 eruption of volcano Usu in Hokkaido, Japan. *Bull*
 1127 *Volcanol* 11:45-157
 1128 Miyamoto K (2010) Numerical simulation of landslide movement and Unzen-Mayuyama disaster in
 1129 1792, Japan. *J Disast Res* 5(3): 280-287
 1130 Montanaro C, Beget J, Marti J, Siebert L, Coombs M (2011) Volcano collapse along the Aleutian Ridge
 1131 (western Aleutian Arc). *Nat Haz & Earth Sys Scie* 11(3)
 1132 Moon V, Bradshaw J, Smith R, de Lange W (2005) Geotechnical characterisation of stratocone crater
 1133 wall sequences, White Island Volcano, New Zealand. *Engineering geology*, 81(2):146-178
 1134 Moore JG, Fiske RS (1969) Volcanic substructure inferred from dredge samples and ocean-bottom
 1135 photographs, Hawaii. *G S A Bulletin* 80(7):1191-1202
 1136 Morgan JK, Moore GF, Clague DA (2003) Slope failure and volcanic spreading along the submarine
 1137 south flank of Kilauea volcano, Hawaii. *J Geophys Res: Solid Earth* 108(B9)
 1138 Moriya I (1980) Bandaian Eruption and Landforms Associated with It: Collection of Articles in
 1139 Memory of Retirement of Prof. K. Hishimura. Vol. 66. Tohoku University
 1140 Murray JB, van Wyk de Vries B, Pitty A, Sargent P, Woller L (2018) Gravitational sliding of the Mt.
 1141 Etna massif along a sloping basement. *Bull Volc* 80:40
 1142 Nakamura K (1977) Volcanoes as possible indicators of tectonic stress orientation-Principle and
 1143 proposal. *J Vol Geotherm Res* 2: 1-16
 1144 Norini G, Agliardi F, Crosta G, Groppelli G, Zuluaga MC (2019) Structure of the Colima Volcanic
 1145 Complex: Origin and Behaviour of Active Fault Systems in the Edifice, in Varley N, Connor CB,
 1146 Komorowski JC (eds) *Volcanic de Colima, Active Volcanoes of the World*, Springer
 1147 Norini G, Capra L, Groppelli G, Agliardi F, Pola A, Cortes A (2010) The structural architecture of the
 1148 Colima Volcanic Complex. *J Geophys Res - Solid Earth* 115:B12209
 1149 Norini G, Acocella V (2011) Analogue modelling of flank instability at Mount Etna: understanding the
 1150 driving factors. *J Geophys Res - Solid Earth* 116:B07206.
 1151 Norini G, Bustos E, Arnosio M, Baez W, Zuluaga MC, Roverato M (2020) Unusual volcanic instability
 1152 and sector collapse configuration at Chimpa volcano, central Andes. *J Volc Geoth Res*
 1153 393:106808
 1154 Norini G, De Beni E, Andronico D, Polacci M, Burton M, Zucca F (2009) The 16 November 2006 flank
 1155 collapse of the south-east crater at Mount Etna, Italy: Study of the deposit and hazard
 1156 assessment. *J Geophys Res – Solid Earth* 114:B02204
 1157 Norini G, Capra L, Groppelli G, Lagmay AMF (2008) Quaternary sector collapses of Nevado de Toluca
 1158 volcano (Mexico) as consequence of regional tectonics and volcanic evolution. *Geosphere* 4
 1159 (5):854-871
 1160 Norini G, Lagmay AMF (2005) Deformed symmetrical volcanoes. *Geology* 33-7:605-608.
 1161 Normark WR, Moore JG, Torresan ME (1993) Giant volcano-related landslides and the development
 1162 of the Hawaiian Islands. *Submarine landslides: selected studies in the US Exclusive economic*
 1163 *zone. US Geol Surv Bull* 2002:184-196
 1164 Okubo CH (2004) Rock mass strength and slope stability of the Hilina slump, Kilauea volcano,
 1165 Hawaii'i. *J Volc Geoth Res* 138(1-2):43-76
 1166 Paguican EM, Wyk de Vries B, Lagmay AMF (2012) Volcano-tectonic controls and emplacement
 1167 kinematics of the Iriga debris avalanches (Philippines). *Bull Volc* 74 (9): 2067-81
 1168 Poland MP, Peltier A, Bonforte A, Puglisi G (2018) The spectrum of persistent volcanic flank
 1169 instability: A review and proposed framework based on Kilauea, Piton de la Fournaise, and
 1170 Etna. *J Volc Geoth Res* 339: 63-80
 1171 Quidelleur X, Hildenbrand A, Samper A (2008) Causal link between Quaternary paleoclimatic changes
 1172 and volcanic islands evolution. *Geophys Res Lett* 35(2): L02303.

1173 Ramalho RS, Winckler G, Madeira J, Helffrich GR, Hipólito A, Quartau R, Adena K, Schaefer JM (2015)
 1174 Hazard potential of volcanic flank collapses raised by new megatsunami evidence. *Sci Adv*
 1175 1:e1500456
 1176 Reid ME (2004) Massive collapse of volcano edifices triggered by hydrothermal pressurization.
 1177 *Geology* 32(5):373-376
 1178 Reid ME, Christian SB, Brien DL (2000) Gravitational stability of three-dimensional stratovolcano
 1179 edifices. *J Geophys Res - Solid Earth* 105(B3):6043-6056
 1180 Reid ME, Keith TE, Kayen RE, Iverson NR, Iverson RM, Brien DL (2010) Volcano collapse promoted by
 1181 progressive strength reduction: new data from Mount St. Helens. *Bull Volc* 72(6):761-766
 1182 Reid ME, Sisson TW, Brien DL (2001) Volcano collapse promoted by hydrothermal alteration and
 1183 edifice shape, Mount Rainier, Washington. *Geology* 29(9):779-782
 1184 Rincon M, Marquez A, Herrera R, Alonso-Torres A, Granja-Bruña JL, van Wyk de Vries B (2018)
 1185 Contrasting catastrophic eruptions predicted by different intrusion and collapse scenarios.
 1186 *Scientific Reports* 8:6178
 1187 Roberti G, Ward B, van Wyk de Vries B, Friele P, Perotti L, Clague JJ, Giardino M (2017) Precursory
 1188 slope distress prior to the 2010 Mount Meager landslide, British Columbia. *Landslides* 15:637
 1189 Roberti G, Roberts NJ, Lit C (2020) Climatic influence on volcanic landslides, in Roverato M, Dufresne
 1190 A, Procter J (Eds), *Volcanic Debris Avalanches: From Collapse to Hazard*, *Advances in*
 1191 *Volcanology*, Springer
 1192 Rosas-Carbajal M, Komorowski JC, Nicollin F, Gibert D (2016) Volcano electrical tomography unveils
 1193 edifice collapse hazard linked to hydrothermal system structure and dynamics. *Scientific*
 1194 *Reports* 6: 29899
 1195 Roverato M, Capra L (2013) Características microtexturales como indicadores del transporte y
 1196 emplazamiento de dos depósitos de avalancha de escombros del volcán de Colima. *Rev Mex*
 1197 *Ciencias Geol* 30:512-525
 1198 Roverato M, Capra L, Sulpizio R, Norini G (2011) Stratigraphic reconstruction of two debris avalanche
 1199 deposits at Colima Volcano (Mexico): insights into pre-failure conditions and climate influence.
 1200 *J Volcan Geoth Res* 207:33-46
 1201 Roverato M, Cronin S, Procter J, Capra L (2015) Textural features as indicators of debris avalanche
 1202 transport and emplacement, Taranaki volcano. *GSA Bull.* 127: 3-18
 1203 Samaniego P, Valderrama P, Mariño J, van Wyk de Vries B, Roche O, Manrique N, Chédeville C,
 1204 Liorzou, C, Fidel L, Malnati J (2015) The historical (218 ± 14 aBP) explosive eruption of Tutupaca
 1205 volcano (Southern Peru). *Bull Volcanol* 77:51
 1206 Sano Y, Nishio Y, Sasaki S, Gamo T, Nagao K (1998) Helium and carbon isotope systematics at Ontake
 1207 volcano, Japan. *Journal of Geophysical Research: Solid Earth* 103(B10): 23863-23873
 1208 Sassa K, Dang K, He B, Takara K, Inoue K, Nagai O (2014) A new high-stress undrained ring-shear
 1209 apparatus and its application to the 1792 Unzen–Mayuyama megaslide in Japan. *Landslides*
 1210 11(5): 827-842
 1211 Schaefer LN, Di Traglia F, Chaussard E, Lu Z, Nolesini T, Casagli N (2019) Monitoring volcano slope
 1212 instability with Synthetic Aperture Radar: A review and new data from Pacaya (Guatemala) and
 1213 Stromboli (Italy) volcanoes. *Earth-Scie Rev* 192:236-257
 1214 Scott KM, Vallance JW (1995) Debris flow, debris avalanche, and flood hazards at and downstream
 1215 from Mount Rainier, Washington. *US Geological Survey Hydrologic Investigation Atlas*, HA-729
 1216 Scott KM, Vallance JW, Kerle N, Macías JL, Strauch W, Devoli G (2005) Catastrophic precipitation-
 1217 triggered lahar at Casita volcano, Nicaragua: occurrence, bulking and transformation. *Earth Surf*
 1218 *Proc Landforms* 30:59-79
 1219 Scott KM, Vallance JW, Pringle PT (1995) Sedimentology, behavior and hazards of debris flows at
 1220 Mount Rainier, Washington. *U.S. Geol Surv Prof Pap* 1547:56
 1221 Shea T, van Wyk de Vries B (2010) Collapsing volcanoes: the sleeping giants' threat. *Geology Today*
 1222 26(2):72-77
 1223 Shea T, van Wyk de Vries B (2008) Structural analysis and analogue modeling of the kinematics and

1224 dynamics of rockslide avalanches. *Geosphere* 4:657-686
 1225 Shea T, van Wyk de Vries B, Pilato M, (2008) Emplacement mechanisms of contrasting debris
 1226 avalanches at Volcán Mombacho (Nicaragua), provided by structural and facies analysis. *Bull*
 1227 *Volc* 70:899-921
 1228 Siebert L (1984) Large volcanic debris avalanches: Characteristics of source areas, deposits, and
 1229 associated eruptions. *J Volcan Geoth Res* 22: 163-97
 1230 Siebert L, Alvarado GE, Vallance JW, van Wyk de Vries B (2006) Large-volume volcanic edifice failures
 1231 in Central America and associated hazards. In: Rose WI, Bluth GJS, Carr MJ, Ewert JW, Patino
 1232 LC, Vallance JW (eds) *Volcanic Hazards in Central America*. *Geol Soc Am Spec Pap* 412:1-26
 1233 Siebert L, Glicken H, Ui T (1987) Volcanic hazards from Bezymianny-and Bandai-type eruptions. *Bull*
 1234 *Volcan* 49(1):435-459
 1235 Siebert L, Roverato M (2020) A Historical Perspective on Lateral Collapse and Debris Avalanches, in
 1236 Roverato M, Dufresne A, Procter J (Eds), *Volcanic Debris Avalanches: From Collapse to Hazard*,
 1237 *Advances in Volcanology*, Springer
 1238 Smith JR, Malahoff A, Shor AN (1999) Submarine geology of the Hilina slump and morpho-structural
 1239 evolution of Kilauea volcano, Hawaii. *Journal of volcanology and geothermal research* 94(1-
 1240 4):59-88
 1241 Thomas ME, Petford N, Bromhead EN (2004) The effect of internal gas pressurization on volcanic
 1242 edifice stability: evolution towards a critical state. *Terra Nova* 16:312-317
 1243 Tibaldi A (1995) Morphology of pyroclastic cones and tectonics. *Journal of Geophysical Research* 100
 1244 (B12): 24521-24535
 1245 Tibaldi A (1996) Mutual influence of dyking and collapses at Stromboli Volcano, Italy. *Geol Soc*,
 1246 London, Special Publications 110: 55-63
 1247 Tibaldi A (2001) Multiple sector collapses at Stromboli volcano, Italy: how they work. *Bull Volcan*
 1248 63(2-3):112-125
 1249 Tibaldi A, Lagmay AMF, Ponomareva VV (2005) Articles effects of basement structural and
 1250 stratigraphic heritages on volcano behaviour and implications for human activities (the
 1251 UNESCO / IUGS / IGCP Project 455). *Episodes* 28 (3): 158-70
 1252 Tinti S, Pagnoni G, Zaniboni F (2006) The landslides and tsunamis of the 30th of December 2002 in
 1253 Stromboli analysed through numerical simulations. *Bull Volcanol* 68:462–479
 1254 Tommasi P, Baldi P, Chiocci FL, Coltelli M, Marsella M, Pompilio M, Romagnoli C (2005) The landslide
 1255 sequence induced by the 2002 eruption at Stromboli volcano. In *Landslides: Risk Analysis and*
 1256 *Sustainable Disaster Management* (251-258). Springer, Berlin, Heidelberg
 1257 Tormey D (2010) Managing the effects of accelerated glacial melting on volcanic collapse and debris
 1258 flows: Planchon-Peteroa volcano, Southern Andes. *Glob Plan Ch* 74 (2), 82–90
 1259 Tuffen H (2010) How will melting of ice affect volcanic hazards in the twenty-first century? *Phil Trans*
 1260 *R Soc A* 368:2535-2558
 1261 Ui T (1983) Volcanic dry avalanche deposits—identification and comparison with nonvolcanic debris
 1262 stream deposits. *J Volcan Geoth Res* 18(1-4):135-150
 1263 Ui T, Yamamoto H, Suzuki-Kamata K (1986) Characterization of Debris Avalanche Deposits in Japan. *J*
 1264 *Volcan Geoth Res* 29 (1-4): 231-43
 1265 Urlaub M, Petersen F, Gross F, Bonforte A, Puglisi G, Guglielmino F, Krastel S, Lange D, Kopp H (2018)
 1266 Gravitational collapse of Mount Etna's southeastern flank. *Science advances* 4(10):eaat9700.
 1267 Usami T (1996) Materials for comprehensive list of destructive earthquakes in Japan, 416-1995.
 1268 Revised and Enlarged Edition. University of Tokyo Press: Tokyo, Japan.
 1269 Valderrama P, Roche O, Samaniego P, van Wyk de Vries B, Bernard K, Mariño J (2016). Dynamic
 1270 implications of ridges on a debris avalanche deposit at Tutupaca volcano (southern Peru). *Bull*
 1271 *Volcan* 78:14
 1272 Villemant B, Komorowski JC, Dessert C, Michel A, Crispi O, Hammouya G, Beauducel F, De Chabaliér
 1273 JB (2014) Evidence for a new shallow magma intrusion at La Soufrière of Guadeloupe (Lesser
 1274 Antilles). *J Vol Geoth Res* 285:247-277

1275 Vallance JW, Scott KM (1997) The Osceola mudflow from Mount Rainier: sedimentology and hazard
1276 implication of a huge clay-rich debris flow. *GSA Bull* 109:143-163

1277 Vallance JW, Siebert L, Rose WI, Girón JR, Banks NL (1995) Edifice collapse and related hazards in
1278 Guatemala. In: Ida Y and Voight B (eds), *Models of magmatic processes and volcanic eruptions*.
1279 *J Volcanol Geotherm Res* 66:337-345

1280 Van Bemmelen RW (1949) *The geology of Indonesia: The Hague, Government Printing Office, 732*

1281 van Wyk de Vries B (1998) Extension Induced by Volcanic Loading in Regional Strike-Slip Zones.
1282 *Geology* 26(11): 983-986.

1283 van Wyk de Vries B, Davies T (2015) Landslides, debris avalanches, and volcanic gravitational
1284 deformation. In *The Encyclopedia of Volcanoes* (665-685). Academic Press.

1285 Van Wyk de Vries B, Fancis PW (1997) Catastrophic collapse at stratovolcanoes induces by gradual
1286 volcano spreading. *Nature* 387:387-390

1287 van Wyk de Vries B, Matela RJ (1998) Styles of volcano-induced deformation: numerical models of
1288 substratum flexure, spreading and extrusion. *J Vol Geoth Res* 81:1-18.

1289 van Wyk de Vries B, Merle O (1996) The effect of volcanic constructs on rift fault patterns. *Geology*
1290 24 (7): 643

1291 van Wyk de Vries B, Kerle N, Petley D (2000) A sector-collapse forming at Casita volcano. *Geology*
1292 28:167-170

1293 van Wyk de Vries B, Self S, Francis PW, Keszthelyi L (2001) A gravitational spreading origin for the
1294 Socompa debris avalanche: *J Vol Geoth Res* 105:225-247

1295 Verrucci L, Tommasi P, Boldini D, Graziani A, Rotonda T (2019) Modelling the instability phenomena
1296 on the NW flank of Stromboli Volcano (Italy) due to lateral dyke intrusion. *J Vol Geoth Res*
1297 371:245-262

1298 Vidal N, Merle O (2000) Reactivation of basement faults beneath Volcanoes: A new model of flank
1299 collapse. *J Vol Geoth Res* 99 (1-4): 9-26

1300 Villemant B, Komorowski JC, dessert c, Michel A, Crispi O, Hammouya G, Beauducel F, DE Chaballier
1301 JB (2014) Evidence for a new shallow magma intrusion at La Soufrière of Guadeloupe (Lesser
1302 Antilles) Insights form long-term geochemical monitoring of halogen-rich hydrothermal fluids. *J*
1303 *Vol Geoth Res* 285:247-277

1304 Voight B (2000) Structural stability of andesite volcanoes and lava domes. *Philosophical Transactions*
1305 *of the Royal Society of London. Series A: Math, Physic Engin Sci* 358(1770):1663-1703

1306 Voight B, Elsworth D (1997) Failure of volcano slopes. *Geotechnique*, 47(1): 1-31

1307 Voight B, Glicken H, Janda RJ, Douglass PM (1981) Catastrophic rockslide avalanche of May 18. In
1308 *The 1980 Eruptions of Mount St. Helens, Washington* (1250: 347-377). *US Geol Surv Prof Pap*

1309 Voight B, Janda RJ, Glicken H, Douglass PM (1983) Nature and mechanics of the Mount St Helens
1310 rockslide-avalanche of 18 May 1980. *Geotechnique* 33:243-273

1311 Voight B, Komorowski JC, Norton GE, Belousov AB, Belousova M, Boudon G, Young SR (2002) The 26
1312 December (Boxing Day) 1997 sector collapse and debris avalanche at Soufriere Hills volcano,
1313 Montserrat. *Geol Soc, London, Memoirs* 21(1):363-407

1314 Voight B, Sousa J (1994) Lessons from Ontake-san: a comparative analysis of debris avalanche
1315 dynamics. *Engineering Geology* 38(3-4): 261-297

1316 Wang J, Ward SN, Xiao L (2019) Tsunami Squares modeling of landslide generated impulsive waves
1317 and its application to the 1792 Unzen-Mayuyama mega-slide in Japan. *Engineering Geology*
1318 256: 121-137

1319 Watanabe K, Hoshizumi H, Itaya T (1993) K-Ar Ages of Unzen Volcano in Kyushu, Japan: With Some
1320 Aspects of Geology of Mayu-Yama.

1321 Watt SFL, Karstens J, Micallef A, Berndt C, Urlaub M, Ray M, Desai A, Sammartini M, Klauke I,
1322 Böttner C, Day S, Downes H, Kühn M, Elger J (2019) From catastrophic collapse to multi-phase
1323 deposition: Flow transformation, seafloor interaction and triggered eruption following a
1324 volcanic-island landslide. *Earth Plan Sci Lett* 517:135-147

1325 Watt SFL, Karstens J, Berndt C (2020) Volcanic-island lateral collapse and their submarine deposits, in

- Roverato M, Dufresne A, Procter J (Eds), Volcanic Debris Avalanches: From Collapse to Hazard, Advances in Volcanology, Springer
- Waythomas CF, Miller TP, Begét JE (1999) Record of late Holocene debris avalanche and lahar formation at Iliamna Volcano, Alaska. Am Geophys Union Trans 80: F1140
- Waythomas CF, Wallace KL (2002) Flank collapse at Mount Wrangell, Alaska, recorded by volcanic mass-flow deposits in the Copper River lowland. Canad J Earth Scie 39:1257-1279
- Williams R, Rowley P, Garthwaite MC, (2019) Reconstructing the Anak Krakatau flank collapse that caused the December 2018 Indonesian tsunami. Geology 47(10):973-976
- Wooller L, van Wyk de Vries B, Cecchi E, Rymer H (2009) Analogue Models of the Effect of Long-Term Basement Fault Movement on Volcanic Edifices. Bull Volc 71 (10): 1111-31
- Wooller L, van Wyk de Vries B, Murray JB, Rymer H, Meyer S (2004) Volcano spreading controlled by dipping substrata. Geology 32(7):573-576
- Zernack AV, Cronin SJ, Bebbington MS, Smith IEM, Price RC, Procter JN (2012) Forecasting catastrophic stratovolcano collapse, a model based on Mt. Taranaki, New Zealand. Geology 40:983-986
- Zimbelman DR (1996) Hydrothermal alteration and its influence on volcanic hazards Mount Rainier, Washington, a case history: PhD Thesis, Univ. Colorado, Boulder
- Zimbelman DR, Watters RJ, Firth IR, Breit GN, Carrasco-Nuñez G (2004) Stratovolcano stability assessment methods and results from Citlaltepētli, Mexico. Bull Volc 66:66-79

Figure captions:

Figure 1 (a) Model showing crater opening perpendicular to dike propagation (Nakamura 1977). (b) Model off crater opening perpendicular to the strike of normal faults and σ_{Hmax} (Francis and Self 1987). (c) Crater openings in Andean volcanoes (Francis and Wells 1988). (d) Crater opening at an acute angle to the underlying strike-slip fault (Lagmay et al. 2000).

Figure 2 Analogue experiment results by Wooller et al. 2009 showing the fault patterns created, areas of instability and the main control of the main geometrical parameters. Figures from Wooller et al., (2009).

Figure 3 Google Earth image showing the Colima Volcanic Complex formed by the youngest Fuego de Colima volcano grown into the Paleofuego volcano depression (dashed line). Both volcanoes are located on the southern flank of the older and inactive Nevado de Colima volcano.

Figure 4 Example of highly hydrothermally altered material in a VDAD at Nevado del Ruiz, Colombia. *Espeletia* plants for scale at the top of the cliff are ~1.5 m tall (photo: Matteo Roverato).

Figure 5 USGS photograph taken by Peter Lipman on April 27, 1980, showing the impressive bulge on the northeastern flank of the Mount St. Helens volcano few days before its collapse.

Figure 6 Moderate resolution, SENTINEL-2 multispectral images of the Anak Krakatau A) before (1st November 2018; true colour) and B) after (2nd February 2019; true color) the lateral collapse occurred on 22nd December 2018. After the collapse, the intense explosive volcanic activity started to re-fill the collapse scar, producing a small coastal tuff ring.

Figure 7 Sketch representation of the instability factors that can destabilize a volcano and the triggering mechanisms, such as explosive activity and earthquakes, which promote a volcanic sector

collapse. The top right rectangle represents schematically how the hydrothermal activity affects a volcanic edifice, promoting leaching and dissolution of the rock, favoring the formation of voids and the migration and accumulation of clay minerals, and consequently, fluid storage and impermeability. This promotes loss of strength and shear reduction of the rock. See the text for more details.

Tables:

Table 1 References used in the present text regarding factors that lead to the instability of volcanic edifices and trigger a volcanic sector collapse. Note that some references are underlined because they do not refer directly to volcanic lateral collapse but are still relevant for understanding aspects of the processes promoting a volcanic edifice failure.

Table 2 Examples of volcanic debris avalanche deposits that display hydrothermal alteration

Table 3 Key features in a VDAD that could suggest what happened in the volcano edifice before and/or during the lateral collapse event

Table 1

Instability factors	References
Basement tectonics and faults	Nakamura 1977; Moriya 1980; Bahar and Girod 1983; Siebert 1984; <u>Delaney et al. 1986</u> ; Ui et al. 1986; Francis and Self 1987; Francis and Wells 1988; Carracedo 1994; <u>Tibaldi 1995</u> , 1996; Elsworth and Voight 1996; van Wyk de Vries and Merle 1996, 1998; Lagmay et al. 2000; Vidal and Merle 2000; Norini and Lagmay 2005; Tibaldi et al. 2005; <u>Lagmay and Valdivia 2006</u> ; Norini et al. 2008; Wooller et al. 2009; Norini et al. 2010; Paguican et al. 2012; Norini et al. 2019; Norini et al. 2020
Sloping substrate and gravitational spreading	<u>van Bemmelen 1949</u> ; Borgia et al. 1992; Vallance et al. 1995; Merle and Borgia 1996; van Wyk de Vries and Francis 1997; Francis and Wells 1998; van Wyk de Vries and Matela 1998; Borgia et al. 2000; van Wyk de Vries et al. 2001; Deplus et al. 2001; Clavero et al. 2002; <u>Kuhn 2002</u> ; Morgan et al. 2003; Lundgren et al. 2004; Wooller et al. 2004; Carrasco-Núñez et al. 2006; Siebert et al. 2006; Del Camp et al. 2008; Shea et al. 2008; Norini et al. 2009; Andrade and van Wyk de Vries 2010; Norini et al. 2010; Shea and van Wyk De Vrie 2010; Bonforte et al. 2011; Norini and Acocella 2011; Roverato et al. 2011; Paguican et al. 2012; Murray et al. 2018; Norini et al. 2019
Hydrothermal alteration	<u>Frantz et al. 1981</u> ; Siebert 1984; Carrasco--Núñez et al. 1993; Lopez and Williams 1993; <u>Frank 1995</u> ; <u>Scott et al. 1995</u> ; Day 1996; Kerle and Zimbelman 1996; <u>Vallance and Scott 1997</u> ; Voight and Elsworth 1997; Van Wyk de Vrie et al. 2000; van Wyk de Vrie 2001; Reid et al. 2001; Waythomas and Wallace 2002; Merle et al. 2003; Reid 2004; Zimbelman et al. 2004; Alvarado et al. 2004; Cecchi et al. 2005; <u>John et al. 2008</u> ; Shea et al. 2008; Borselli et al. 2011; Roverato et al. 2011; Delmelle et al. 2015; Rosas-Carbajal et al. 2016; Valderrama et al. 2016; Norini et al. 2020

Dikes and magma intrusions	Gorshkov 1959; Gorshkov and Bogoyavenskaya 1965; Lipman et al. 1981; Voight et al. 1983; Siebert et al. 1987; Beget and Kienle 1992; Elsworth and Voight 1995; Day 1996; Elsworth & Voight 1996; Donnadieu and Merle 1998; Belousov et al. 1999; Donnadieu et al. 2001; McClelland and Erwin 2003; McGuire 2003; Belousov et al. 2004; Reid 2004; Thomas et al. 2004; Macias et al. 2010; Borselli et al. 2011; Norini and Acocella 2011; Villemant et al. 2014; Samaniego et al. 2015; Rincon et al. 2018
Past and present climate implications	Carrasco-Nuñez et al. 1993; McGuire 1996; <u>Clapperton 1998</u> ; Waythomas and Begét 2000; Kerle and van Wyk de Vries 2001; Scott et al. 2005; Capra 2006; Huggel et al. 2007; <u>Huggel et al. 2008</u> ; <u>Bennett and Glasser 2009</u> ; Tormey 2010; Tuffen 2010; Roverato et al. 2011; Capra et al. 2013; Roverato and Capra 2013; Roverato et al. 2015; <u>Kos et al. 2016</u> ; <u>Grämiger et al. 2017</u> ; Roberti et al. 2017; Roberti et al. 2020 (this volume)
Triggering mechanisms	References
Explosive eruptions (Bezymianny/Bandai types)	Gorshkov 1959; Gorshkov 1962; Moriya 1980; Bogoyavlenskaya and Kirnasov 1981; Ui 1985; Siebert et al. 1987; Voight and Elsworth 1997; Leyrit 2000; Donnadieu et al. 2001, Belousov et al. 2004; Dzurisin 2018
Earthquakes (Unzen/Ontake type)	Christiansen et al. 1981; Endo et al. 1981; Lipman and Mullineaux 1981; Voight et al. 1981; Keefer 1984; Ui et al. 1986; Fairchild 1987; Endo et al. 1989; Watanabe et al. 1993; Voight and Sousa 1994; Glicken 1996; Usami, 1996; Voight and Elsworth 1997; Hoshizumi et al. 1998; Sano et al. 1998; Inoue 2000; Donnadieu et al. 2001; McGuire 2003; Miyamoto 2010; Sassa et al. 2014; Della Pasqua et al. 2016; Kataoka et al. 2018; Wang et al. 2019

Table 2

Deposit	Volcano	Reference
Teteltzingo VDAD	Citlaltepetl volcano, Mexico	Carrasco-Núñez et al. (1993)
1998 VDAD	Casita volcano, Nicaragua	Kerle and van Wyk de Vries (2001)
Chetaslina mass-flow	Mount Wrangell, Alaska, USA	Waythomas and Wallace (2002)
Piton de la Fournaise VDAD	Reunion Island, France	Merle and Lenat (2003)
Fortuna VDAD	Miravalles volcano, Costa Rica	Alvarado et al. (2004)
El Crater VDAD	Mombacho volcano, Nicaragua	Shea et al. (2008)
Tonila VDAD	Colima volcano, Mexico	Roverato et al. (2011)

Tutupaca VDAD	Tutupaca volcano, Peru	Valderrama et al. (2016)
Chimpa Unit	Chimpa volcano, Argentina	Norini et al. (2020)


Table 3


Key features		Pre/syn-collapse setting
Hydrothermal alteration	The VDAD displays: - color stains in those reaches (e.g. proximal) where the volcanoclastic material is not mixed with other materials; - homogeneous yellowish matrix in portions (e.g. medial/distal) where the material is partially or completely mixed; - high clay content	The edifice was affected by hot fluid circulation favoring hydrothermal alteration responsible for the weakening of the strength of the rock and promoting instability of the volcano.
Water content	The VDAD is characterized by a water-saturated (or semi-saturated) aspect displaying facies with characteristics between debris-flow deposits (DFDs) and VDADs - homogenous and partially cemented matrix- preservation of millimetric porosity The VDAD is often associated to DFDs that lying directly on top of it	The edifice was affected by high water content due to different climatic conditions such as: - high precipitation in tropical settings - presence of snow and/or ice in boreal settings or at high altitudes in tropical settings These factors contribute to the instability of the volcanic edifice and could also increase the area affected by a VDA through flow transformation into more mobile debris flows
Pyroclastic deposits	Pyroclastic deposits such as pyroclastic density current (PDC) or fall-out deposits lying directly in contact on top of the VDAD displaying sharp contacts	Pyroclastic material situated on top of the VDAD suggests that an eruption likely occurred after the collapse event as consequence of the depressurization of the magmatic system. The sharp contact between PDC and VDA deposits suggests that the VDA had already ceased to move when the PDC material emplaced. The magmatic event was likely not responsible for the collapse and other triggering mechanisms are involved.

	PDC deposits such as directed blasts and/or blast-generated PDC deposits are “intimately” associated with the VDAD. Some characteristics are: - the contact between PDC and VDA deposits is very irregular - twisted clastic dikes of PDC material penetrate into the VDAD - in distal reached PDC deposits are located below and above the VDAD These features could be associates to the following section (juvenile material).	These characteristics indicate that the VDA was still moving when the PDC material started to emplace on the upper surfaces of the VDA. PDC deposits located below the VDAD in distal portions suggest that the kinetic energy of the PDC material was higher than the VDA and that it emplaced before the arrival of the collapsing mass. This section could be associates with the following one (juvenile material).
Juvenile material	Fresh fragments of volcanic material (lava fragments or other pyroclastic material) are immersed into the VDAD	It indicates that the collapse was associated to a magmatic event that triggered the collapse. Fresh lava fragments could indicate the presence of a summit dome or a shallow intrusion. Vesiculated juveniles are the product of the explosive character of the eruption.


1407

1408





1410
1411
1412
1413
1414
1415
1416
1417
1418





1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431



1432



1433



1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452

