



# Transcutaneous capnometry-guided exercise in respiratory settings

M. Polastri, M.L. Vega Pittao, J. Crotti, L. Pisani & S. Nava

To cite this article: M. Polastri, M.L. Vega Pittao, J. Crotti, L. Pisani & S. Nava (2024) Transcutaneous capnometry-guided exercise in respiratory settings, *Pulmonology*, 30:5, 498-499, DOI: [10.1016/j.pulmoe.2023.12.003](https://doi.org/10.1016/j.pulmoe.2023.12.003)

To link to this article: <https://doi.org/10.1016/j.pulmoe.2023.12.003>



© 2024 Sociedade Portuguesa de Pneumologia/SPP



Published online: 10 Mar 2025.



Submit your article to this journal [↗](#)



Article views: 12



View related articles [↗](#)



View Crossmark data [↗](#)



## LETTER TO THE EDITOR

## Transcutaneous capnometry-guided exercise in respiratory settings



Dear Editor,

Capnometry is a noninvasive method to estimate the partial pressure of carbon dioxide ( $p\text{CO}_2$ ) in respiratory gases.<sup>1,2</sup> We describe here a case of a 48-year-old woman (who provided consent for using her anonymised data) with nemaline myopathy admitted to our unit. The patient had been undergoing noninvasive ventilation (NIV) for more than twenty years. Because of discomfort following new ventilator settings prescribed at home, she was admitted to the hospital –for readaptation to NIV– presenting with compensated chronic respiratory failure (pH 7.38,  $\text{PaCO}_2$  63.5 mmHg,  $\text{PaO}_2$  76.9 mmHg,  $\text{HCO}_3$  33.6 mmol/L) while on standard oxygen therapy ( $\text{FiO}_2$  28% via nasal prongs). After a few days, the patient optimised comfort and synchronisation with the ventilator. On day 4, however, she presented with acute-on-chronic respiratory acidosis (pH 7.34,  $\text{PaCO}_2$  84.3 mmHg,  $\text{PaO}_2$  73.7 mmHg,  $\text{HCO}_3$  39.1 mmol/L). The patient, whose Hospital Anxiety and Depression Score was 21 points, reported having taken 2 mg of lorazepam to manage an anxiety crisis.

Despite the hypercapnia level during the clinical stability phase, the patient could still perform daily moderate respiratory and motor exercises (see below) at home. The recent exacerbation episode posed us the question of whether we can proceed with physiotherapy because of the elevated levels of  $\text{PaCO}_2$ . Resting hypercapnia or pulmonary function tests do not predict hypercapnia during exercise,<sup>3</sup> which in some patients may reach elevated values, theoretically putting them at risk. The possible development or worsening of alveolar gas exchange during exercise is usually assessed with a cardio-respiratory test that measures minute ventilation (VE), physiologic dead space ( $V_D/V_T$ ), and carbon dioxide output ( $\text{VCO}_2$ ). This test is somehow burdensome and has some technical biases when exercise is performed with supplemental oxygen, while capnometry may much easier depict gas derangement during exercise, but it is very seldom used. Indeed, this test is mostly used in COPD patients, while no data is available for restrictive patients.

We, therefore, monitored capnometry during the hospital physiotherapy, which consisted of respiratory (chest expansion, breathing control techniques, and inspiratory muscle training) and motor exercises (strengthening and aerobic).

The strengthening exercise consisted of active squats of the lower limbs, whereas aerobic activities focused on walking and marching on the spot with alternate arm elevation. The patient exercised at an intensity of 13–14 points (somewhat hard) in the ratings of the perceived exertion scale,<sup>4</sup> and each session lasted about 30 min as per patient's tolerance. Transcutaneous  $\text{CO}_2$  ( $\text{TcCO}_2$ ) monitoring (SenTec Digital Monitoring System, Therwil, Switzerland) was executed on two different days under two ventilatory modalities (standard oxygen therapy via nasal prongs and NIV with additional  $\text{O}_2$ ) to determine to which extent exercise contributed to a possible increase in hypercapnia. We measured physiological parameters such as oxygen saturation, heart rate, dyspnoea intensity and muscle fatigue (using the modified Borg scale).<sup>5</sup> In Table 1,  $\text{TcCO}_2$  monitoring data and physiological parameters during exercise are listed in detail. These data, taken together with  $\text{TcCO}_2$ , contributed to planning an appropriate exercise programme, suggesting that NIV cycles may contain a further increase in hypercapnia vs standard oxygen via nasal prongs and avoid taking benzodiazepine on exercise days.

This is the first report assessing exercise hypercapnia in a restrictive patient, suggesting that a simple technique such as capnometry can titrate the optimal mode of supporting exercise without applying more complex instrumentation, like during a cardiopulmonary test.

Studies have investigated the use of capnography in patients with COPD and back pain, exploring the possibility of using capnography as a biofeedback, promoting self-regulated breathing.<sup>6,7</sup> A critical feature of these studies was the psychological aspect; anxiety and depression were two of the most hidden and impactful complications in patients with COPD. In a case series of 29 patients with back or neck pain, the authors found patients had poor respiratory function as tested with capnography, and all could improve respiratory function with retraining.<sup>6</sup>

Furthermore, in the CALM project, it has been found that guiding respiratory exercise with real-time biofeedback (capnography) can promote self-regulated breathing, reducing the connections between dyspnoea and anxiety symptoms and targeting breathing efficiency and self-efficacy in COPD.<sup>7</sup>

In the case described here, we were able to define the optimal exercise modality and suggest the patient adhere to a given protocol to avoid hypercapnia. Transcutaneous

<https://doi.org/10.1016/j.pulmoe.2023.12.003>

2531-0437/© 2023 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Table 1** TcCO<sub>2</sub> monitoring during exercise under two different ventilatory modalities: SOT and NIV.

Activity (duration, minutes)	TcCO <sub>2</sub> mmHg		SpO <sub>2</sub> %		Heart rate beats x min		Dyspnoea mBorg, score		Fatigue mBorg, score	
	SOT	NIV	SOT	NIV	SOT	NIV	SOT	NIV	SOT	NIV
Pretest	76.1	59	96	96	109	95	0.5	0.5	0	0
Strengthening (10')	78.4	62.5	97	93	114	112	4	3	3	2
Rest (5')	76.6	67.9	95	86	105	95	4	4	0	2
Respiratory exercise (10')	75.1	70.1	96	100	115	95	3	4	3	3
Rest (5')	74.6	60	96	95	108	97	1	3	0	2
Walking (3')	80.6	63.8	94	97	114	113	4	3	3	4
Rest (5')	76.4	57.1	97	95	103	89	1	2	0	3

TcCO<sub>2</sub>, transcutaneous CO<sub>2</sub>; SOT, standard oxygen therapy; NIV, noninvasive ventilation; mBorg, modified Borg scale.

capnometry-guided exercise is a promising practice that could simplify and optimise physiotherapy practice in respiratory settings.

### Conflicts of interest

The authors have no conflicts of interest to declare.

### Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

### References

- Schmitz BD, Shapiro BA. Capnography. *Respir Care Clin N Am*. 1995;1(1):107–17.
- Benumof JL. Interpretation of capnography. *AANA J*. 1998;66(2):169–76.
- O'Donnell DE, D'Arsigny C, Fitzpatrick M, Webb KA. Exercise hypercapnia in advanced chronic obstructive pulmonary disease: the role of lung hyperinflation. *Am J Respir Crit Care Med*. 2002;166(5):663–8. <https://doi.org/10.1164/rccm.2201003>.
- Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*. 1982;14(5):377–81.
- Burdon JG, Juniper EF, Killian KJ, Hargreave FE, Campbell EJ. The perception of breathlessness in asthma. *Am Rev Respir Dis*. 1982;126(5):825–8. <https://doi.org/10.1164/arrd.1982.126.5.825>.
- McLaughlin L, Goldsmith CH, Coleman K. Breathing evaluation and retraining as an adjunct to manual therapy. *Man Ther*. 2011;16(1):51–2. <https://doi.org/10.1016/j.math.2010.08.006>.
- Norweg A, Hofferber B, Oh C, Spinner M, Stavrolakes K, Pavol M, et al. Capnography-assisted learned, monitored (CALM) breathing therapy for dysfunctional breathing in COPD: a bridge to pulmonary rehabilitation. *Contemp Clin Trials*. 2023;134:107340. <https://doi.org/10.1016/j.cct.2023.107340>.

M. Polastri<sup>a,\*</sup>, M.L. Vega Pittao<sup>b,c</sup>, J. Crotti<sup>b</sup>, L. Pisani<sup>b,c</sup>, S. Nava<sup>b,c</sup>

<sup>a</sup> Department of Continuity of Care and Integration, Physical Medicine and Rehabilitation, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

<sup>b</sup> Department of Medical and Surgical Sciences (DIMEC), Alma Mater Studiorum University of Bologna, Bologna, Italy

<sup>c</sup> Respiratory and Critical Care Unit, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

\* Corresponding author at: Via G. Massarenti 9, Bologna 40138, Italy.

E-mail address: [massimiliano.polastri@aosp.bo.it](mailto:massimiliano.polastri@aosp.bo.it) (M. Polastri).

Received 28 November 2023; Accepted 12 December 2023  
Available online 5 January 2024