



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Q^* -curvature flow on pseudo-Einstein CR manifolds

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Maalaoui, A., Martino, V. (2024). Q^* -curvature flow on pseudo-Einstein CR manifolds. ANNALI DI MATEMATICA PURA ED APPLICATA, 203(4), 1851-1878 [10.1007/s10231-024-01425-3].

Availability:

This version is available at: <https://hdl.handle.net/11585/969505> since: 2024-05-13

Published:

DOI: <http://doi.org/10.1007/s10231-024-01425-3>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

\overline{Q}' -curvature flow on Pseudo-Einstein CR manifolds

Ali Maalaoui⁽¹⁾ & Vittorio Martino⁽²⁾

Abstract In this paper we consider the problem of prescribing the \overline{Q}' -curvature on three dimensional Pseudo-Einstein CR manifolds. We study the gradient flow generated by the related functional and we will prove its convergence to a limit function under suitable assumptions.

Keywords: Pseudo-Einstein CR manifolds, \overline{P}' -operator

2010 MSC. Primary: 58J60, 58J05. Secondary: 58E05, 58E07 .

1 Introduction and statement of the results

Let $(M, T^{1,0}M, \theta)$ be a CR three manifold, which we will always assume smooth and closed. It is known that one can construct a pair (Q, P_θ) such that under a conformal change of the contact form $\hat{\theta} = e^u\theta$, one has

$$P_\theta u + Q_\theta = Q_{\hat{\theta}} e^{2u}$$

where the Paneitz operator $P_\theta = (\Delta_b)^2 + T^2 + l.o.t.$; in particular the operator P_θ contains the space of CR pluriharmonic functions \mathcal{P} in its kernel, moreover the total Q -curvature is always zero [16], hence it does not provide any extra geometric information.

Therefore, one considers another pair (P', Q') , see [3], where P' is a Paneitz type operator satisfying $P' = 4(\Delta_b)^2 + l.o.t.$ and is defined on the space of pluriharmonic functions and the Q' -curvature is defined implicitly so that

$$P'_\theta u + Q'_\theta - \frac{1}{2}P_\theta(u^2) = Q'_{\hat{\theta}} e^{2u},$$

which is equivalent to

$$P'_\theta u + Q'_\theta = Q'_{\hat{\theta}} e^{2u} \text{ mod } \mathcal{P}^\perp. \quad (1)$$

In the case of pseudo-Einstein three dimensional CR manifolds (we refer the reader to the next section for further details), in [10] the authors showed that the total Q' -curvature is not always zero and it is invariant under the conformal change of the contact structure; in

¹Department of Mathematics, Clark University, Worcester, MA 01610, USA.
E-mail address: amaalaoui@clarku.edu

²Dipartimento di Matematica, Università di Bologna, piazza di Porta S.Donato 5, 40126 Bologna, Italy.
E-mail address: vittorio.martino3@unibo.it

particular it is proportional to the Burns-Epstein invariant $\mu(M)$ (see [5], [13]) and if (M, J) is the boundary of a strictly pseudo-convex domain X , then

$$\int_M Q'\theta \wedge d\theta = 16\pi^2 \left(\chi(X) - \int_X (c_2 - \frac{1}{3}c_1^2) \right),$$

where c_1 and c_2 are the first and second Chern forms of the Kähler-Einstein metric on X obtained by solving Fefferman's equation.

Now, equation (1) has to be solved orthogonally to the infinite dimensional space \mathcal{P}^\perp : in order to solve this problem on a pseudo-Einstein three dimensional CR manifolds, in [9] it is introduced a new couple $(\overline{P}', \overline{Q}')$, which comes from the projection of equation (1) on to the space of L^2 CR pluriharmonic functions $\hat{\mathcal{P}}$, which is the completion of \mathcal{P} under the L^2 -norm. Since the P' -operator is only defined after projection on \mathcal{P} , we denote by $\Gamma : L^2(M) \rightarrow \hat{\mathcal{P}}$ the orthogonal projection and we let $\overline{P}' = \Gamma \circ P'$ and $\overline{Q}' = \Gamma \circ Q'$, then on a pseudo-Einstein CR manifolds, we can consider the problem of prescribing the \overline{Q}' -curvature, under a conformal change of the contact structure, in particular: for a given a function $f \in \hat{\mathcal{P}}$, we have the following equation

$$P'_\theta u + Q'_\theta = f e^{2u} \text{ mod } \mathcal{P}^\perp, \quad (2)$$

that is equivalent to

$$\overline{P}'_\theta u + \overline{Q}'_\theta = \Gamma(f e^{2u}).$$

Therefore, if u solves (2), then by setting $\tilde{\theta} = e^u \theta$, one has $\overline{Q}'_{\tilde{\theta}} = f$. Let us explicitly notice the differences between the two projections. Since the space of L^2 CR pluriharmonic functions $\hat{\mathcal{P}}$ does not depend on the contact form, thus \overline{Q}'_θ is the orthogonal projection of Q'_θ on $\hat{\mathcal{P}}$ with respect to the L^2 -inner product induced by θ , while $\overline{Q}'_{\tilde{\theta}}$ is the orthogonal projection of $Q'_{\tilde{\theta}}$ with respect to the L^2 -inner product induced by $\tilde{\theta}$; in particular $\phi \in \hat{\mathcal{P}}_\theta$ if and only if $\phi \in \hat{\mathcal{P}}_{\tilde{\theta}}$ and $\psi \in \mathcal{P}_\theta^\perp$ if and only if $e^{-2u}\psi \in \mathcal{P}_{\tilde{\theta}}^\perp$. Therefore, by denoting Γ_u the orthogonal projection induced by $\tilde{\theta}$, one has $\Gamma_u(Q'_{\tilde{\theta}}) = f$. Let us also recall that in [10], the authors show that the non-negativity of the Paneitz operator P_θ and the positivity of the CR-Yamabe invariant imply that \overline{P}' is non-negative and $\ker \overline{P}' = \mathbb{R}$. Moreover, $\int_M Q'_\theta = \int_M \overline{Q}'_\theta \leq 16\pi^2$ with equality if and only if $(M, T^{1,0}M, \theta)$ is the standard sphere; in particular the previously assumptions imply that the $(M, T^{1,0}M, \theta)$ is embeddable (see [12]). Notice that unlike the Riemannian case, it remains unclear if the non-negativity of \overline{P}' and $\ker \overline{P}' = \mathbb{R}$ is a sufficient condition for $\int_M Q'_\theta \leq 16\pi^2$. In particular, the results presented in this paper do not fully cover the case $\overline{P}' \geq 0$ and $\ker \overline{P}' = \mathbb{R}$.

Thus, from now on we will always assume that $(M, T^{1,0}M, \theta)$ is a pseudo-Einstein CR three dimensional manifold such that \overline{P}' is non-negative and $\ker \overline{P}' = \mathbb{R}$. The problem in (2) was first studied in [10] for f constant and in the subcritical case, namely $\int_M \overline{Q}'_\theta < 16\pi^2$. Then in [21] the problem was solved for $f > 0$ via a probabilistic approach again in the subcritical case; also, a solution of the problem was provided in [17] for $f > 0$ and $0 < \int_M \overline{Q}'_\theta < 16\pi^2$ via direct minimization.

In this paper we will study the equation (2) allowing f to change sign: our approach follows closely the methods in [1], where the authors study the analogous problem in the Riemannian setting. In particular, we will use a variational approach by defining a suitable functional on an appropriate space and then we will study the evolution problem along the negative gradient flow lines: the convergence at infinity will provide a solution to the initial problem.

Indeed, with respect to [1], new technical issues will appear, which are essentially due to our sub-Riemannian setting: in particular all the computations and the estimates regarding the convergence along the flow lines have to be done accordingly to the projection on the space of L^2 CR pluriharmonic functions, that we defined earlier. Moreover, some technical estimates on the sphere will be adapted to the CR setting as we will see in Section 5 and the Appendix. Therefore, let us define the following functional $E : H \rightarrow \mathbb{R}$, by

$$E(u) = \int_M u \bar{P}' u + 2 \int_M \bar{Q}' u$$

where $H = \hat{\mathcal{P}} \cap \mathcal{S}^2(M)$ and $\mathcal{S}^2(M)$ is the Folland-Stein Sobolev space equipped with the equivalent norm (see section 2), defined by

$$\|u\|^2 = \int_M u \bar{P}' u + \int_M u^2. \quad (3)$$

We consider the following space, which will serve as a constraint

$$X = \left\{ u \in H; N(u) := \int_M \Gamma(f e^{2u}) = \int_M \bar{Q}' \right\};$$

we notice that the space is well defined since $e^u \in L^2$, see [9], Theorem 3.1.

As in the classical case, we will need the following hypotheses, depending on the sign of $\int_M \bar{Q}'$, namely:

$$\left\{ \begin{array}{ll} (i) & \inf_{x \in M} f(x) < 0, \quad \text{if } \int_M \bar{Q}' < 0 \\ (ii) & \sup_{x \in M} f(x) > 0, \quad \inf_{x \in M} f(x) < 0 \quad \text{if } \int_M \bar{Q}' = 0 \\ (iii) & \sup_{x \in M} f(x) > 0, \quad \text{if } 0 < \int_M \bar{Q}' \leq 16\pi^2. \end{array} \right. \quad (4)$$

In the case when $\int_M \bar{Q}' = 0$, we let ℓ be the unique CR pluriharmonic function satisfying $\bar{P}'_\theta \ell + \bar{Q}' = 0$ and $\int_M \ell = 0$, see [9], Theorem 1.1. Notice that $\bar{Q}'_{e^{\ell\theta}} = 0$. We also recall that in the critical case $M = S^3$, there are some extra compatibility conditions of Kazdan-Warner type that f needs to satisfy in order to be the \bar{Q}' -curvature of a contact structure conformal to the standard one on the sphere (see Theorem 1.3. in [17]).

Now, in order to define the flow equation, we compute the first variation of E , N , and their (\mathcal{S}^2) gradient, respectively:

$$\begin{aligned} \langle \nabla E(u), \phi \rangle &= 2 \int_M \left(\bar{P}' u + \bar{Q}' \right) \phi, \quad \forall \phi \in H, \\ \langle \nabla N(u), \phi \rangle &= 2 \int_M \Gamma(f e^{2u}) \phi, \quad \forall \phi \in H, \\ \nabla E(u) &= 2 \left(\bar{P}' + I \right)^{-1} \left(\bar{P}' u + \bar{Q}' \right), \\ \nabla N(u) &= 2 \left(\bar{P}' + I \right)^{-1} \Gamma(f e^{2u}). \end{aligned}$$

In addition, since by hypotheses (4), $\nabla N \neq 0$ on X , then X is a regular hypersurface in H and a unit normal vector field on X is given by $\nabla N / \|\nabla N\|$. Indeed, $\nabla N(u) \neq 0$ if and only if $\Gamma(e^{2u}f) \neq 0$. This last identity is clear for the hypothesis (i) and (iii). But for (ii), recall that $f \in \hat{\mathcal{P}}$, so if $\Gamma(e^{2u}f) = 0$, then $\int_M e^{2u}f^2 = 0$, leading to a contradiction. The gradient of E restricted to X is then

$$\nabla^X E = \nabla E - \left\langle \nabla E, \frac{\nabla N}{\|\nabla N\|} \right\rangle \frac{\nabla N}{\|\nabla N\|}.$$

Finally, the (negative) gradient flow equation is given by

$$\begin{cases} \partial_t u = -\nabla^X E(u) \\ u(0) = u_0 \in X \end{cases} \quad (5)$$

Now we can state our main results.

Theorem 1.1. *Let $(M, T^{1,0}M, \theta)$ be a pseudo-Einstein CR three dimensional manifold such that \bar{P}' is non-negative and $\ker \bar{P}' = \mathbb{R}$. Let us assume that $\int_M \bar{Q}' < 0$ and let $f \in C(M) \cap \hat{\mathcal{P}}$ as in (4). Then there exists a positive constant C_0 depending on $f^- = \max\{-f, 0\}$, M and θ , such that if*

$$e^{\tau \|u_0\|^2} \sup_{x \in M} f(x) \leq C_0$$

for a constant $\tau > 1$ depending on M and θ , then as $t \rightarrow \infty$, the flow converges in H to a solution u_∞ of (1). Moreover, there exist constants $B, \beta > 0$ such that

$$\|u(t) - u_\infty\| \leq B(1+t)^{-\beta},$$

for all $t \geq 0$.

Theorem 1.2. *Let $(M, T^{1,0}M, \theta)$ be a pseudo-Einstein CR three dimensional manifold such that \bar{P}' is non-negative and $\ker \bar{P}' = \mathbb{R}$. Let us assume that $\int_M \bar{Q}' = 0$ and let $f \in C(M) \cap \hat{\mathcal{P}}$ as in (4). Then as $t \rightarrow \infty$, the flow converges in H to a function u_∞ and there exists a constant λ such that $v = u_\infty + \lambda$ satisfies*

$$\bar{P}'v + \bar{Q}' = \delta \Gamma(fe^{2v}),$$

where $\delta \in \{+1, 0, -1\}$. Moreover, there exist constants $B, \beta > 0$ such that

$$\|u(t) - u_\infty\| \leq B(1+t)^{-\beta},$$

for all $t \geq 0$. If in addition, we assume that $\int_M fe^{2\ell} \neq 0$, then $\delta \neq 0$.

Theorem 1.3. *Let $(M, T^{1,0}M, \theta)$ be a pseudo-Einstein CR three dimensional manifold such that \bar{P}' is non-negative and $\ker \bar{P}' = \mathbb{R}$. Let us assume that $0 < \int_M \bar{Q}' < 16\pi^2$ and let $f \in C(M) \cap \hat{\mathcal{P}}$ as in (4). Then as $t \rightarrow \infty$, the flow converges in H to a solution u_∞ of (1). Moreover, there exist constants $B, \beta > 0$ such that*

$$\|u(t) - u_\infty\| \leq B(1+t)^{-\beta},$$

for all $t \geq 0$.

Finally, the critical case of the sphere, which is a bit different. We will consider a group G acting on S^3 preserving the CR structure. We denote by Σ the set of points fixed by G , that is

$$\Sigma = \{x \in S^3; g \cdot x = x, \forall g \in G\}$$

and we will assume f being invariant under G , namely $f(g \cdot x) = f(x), \forall g \in G$. Then we have the following

Theorem 1.4. *Let us consider the sphere $M = S^3$ equipped with its standard contact structure and let $f \in C(M) \cap \mathcal{P}$ as in (4) and invariant under G . Let us assume that also $u_0 \in X$ is invariant under G . If $\Sigma = \emptyset$ or*

$$\sup_{x \in \Sigma} f(x) \leq e^{-\frac{E(u_0)}{16\pi^2}},$$

then as $t \rightarrow \infty$, the flow converges in H to a solution (invariant under G) u_∞ of (1). Moreover, there exist constants $B, \beta > 0$ such that

$$\|u(t) - u_\infty\| \leq B(1+t)^{-\beta},$$

for all $t \geq 0$.

Acknowledgment

The authors want to express their gratitude to the referee for his/her careful reading of the paper: the remarks and the suggestions led to a serious improvement of the paper.

2 Some definitions in pseudo-Hermitian geometry

We will follow the notations in [10]. Let M^3 be a smooth, oriented three-dimensional manifold. A CR structure on M is a one-dimensional complex sub-bundle $T^{1,0}M \subset T_{\mathbb{C}}M := TM \otimes \mathbb{C}$ such that $T^{1,0}M \cap T^{0,1}M = \{0\}$ for $T^{0,1}M := \overline{T^{1,0}M}$. Let $\mathcal{H} = \text{Re}T^{1,0}M$ and let $J: \mathcal{H} \rightarrow \mathcal{H}$ be the almost complex structure defined by $J(Z + \bar{Z}) = i(Z - \bar{Z})$, for all $Z \in T^{1,0}M$. The condition that $T^{1,0}M \cap T^{0,1}M = \{0\}$ is equivalent to the existence of a contact form θ such that $\ker \theta = \mathcal{H}$. We recall that a 1-form θ is said to be a contact form if $\theta \wedge d\theta$ is a volume form on M^3 . Since M is oriented, a contact form always exists, and is determined up to multiplication by a positive real-valued smooth function. We say that $(M^3, T^{1,0}M)$ is strictly pseudo-convex if the Levi form $d\theta(\cdot, J\cdot)$ on $\mathcal{H} \otimes \mathcal{H}$ is positive definite for some, and hence any, choice of contact form θ . We shall always assume that our CR manifolds are strictly pseudo-convex.

Notice that in a CR-manifold, there is no canonical choice of the contact form θ . A pseudo-Hermitian manifold is a triple $(M^3, T^{1,0}M, \theta)$ consisting of a CR manifold and a contact form. The Reeb vector field T is the vector field such that $\theta(T) = 1$ and $d\theta(T, \cdot) = 0$. The choice of θ induces a natural L^2 -dot product $\langle \cdot, \cdot \rangle$, defined by

$$\langle f, g \rangle = \int_M f(x)g(x) \theta \wedge d\theta.$$

A $(1,0)$ -form is a section of $T_{\mathbb{C}}^*M$ which annihilates $T^{0,1}M$. An admissible coframe is a non-vanishing $(1,0)$ -form θ^1 in an open set $U \subset M$ such that $\theta^1(T) = 0$. Let $\theta^{\bar{1}} := \overline{\theta^1}$ be

its conjugate. Then $d\theta = ih_{1\bar{1}}\theta^1 \wedge \theta^{\bar{1}}$ for some positive function $h_{1\bar{1}}$. The function $h_{1\bar{1}}$ is equivalent to the Levi form. We set $\{Z_1, Z_{\bar{1}}, T\}$ to the dual of $(\theta^1, \theta^{\bar{1}}, \theta)$. The geometric structure of a CR manifold is determined by the connection form ω_1^1 and the torsion form $\tau_1 = A_{1\bar{1}}\theta^1$ defined in an admissible coframe θ^1 and is uniquely determined by

$$\begin{cases} d\theta^1 = \theta^1 \wedge \omega_1^1 + \theta \wedge \tau^1, \\ \omega_{1\bar{1}} + \omega_{\bar{1}1} = dh_{1\bar{1}}, \end{cases}$$

where we use $h_{1\bar{1}}$ to raise and lower indices. The connection forms determine the pseudo-Hermitian connection ∇ , also called the Tanaka-Webster connection, by

$$\nabla Z_1 := \omega_1^1 \otimes Z_1.$$

The scalar curvature R of θ , also called the Webster curvature, is given by the expression

$$d\omega_1^1 = R\theta^1 \wedge \theta^{\bar{1}} \pmod{\theta}.$$

Definition 2.1. *A real-valued function $w \in C^\infty(M)$ is CR pluriharmonic if locally $w = \operatorname{Re} f$ for some complex-valued function $f \in C^\infty(M, \mathbb{C})$ satisfying $Z_{\bar{1}}f = 0$.*

Equivalently, [20], w is a CR pluriharmonic function if

$$\Phi w := \nabla_1 \nabla_1 \nabla^1 w + iA_{1\bar{1}} \nabla^1 w = 0$$

for $\nabla_1 := \nabla_{Z_1}$. We denote by \mathcal{P} the space of all CR pluriharmonic functions and $\hat{\mathcal{P}}$ the completion of \mathcal{P} in $L^2(M)$, also called the space of L^2 CR pluriharmonic functions. Let $\Gamma : L^2(M) \rightarrow \hat{\mathcal{P}}$ be the orthogonal projection on the space of L^2 pluriharmonic functions. If $S : L^2(M) \rightarrow \ker \bar{\partial}_b$ denotes the Szego kernel, then

$$\Gamma = S + \bar{S} + F, \tag{6}$$

where F is a smoothing kernel as shown in [19]. The Paneitz operator P_θ is the differential operator

$$\begin{aligned} P_\theta(w) &:= 4\operatorname{div}(\Phi w) \\ &= \Delta_b^2 w + T^2 - 4\operatorname{Im} \nabla^1 (A_{1\bar{1}} \nabla^1 f) \end{aligned}$$

for $\Delta_b := \nabla^1 \nabla_1 + \nabla^{\bar{1}} \nabla_{\bar{1}}$ the sub-Laplacian. In particular, $\mathcal{P} \subset \ker P_\theta$. Hence, $\ker P_\theta$ is infinite dimensional. For a thorough study of the analytical properties of P_θ and its kernel, we refer the reader to [19, 6, 8]. The main property of the Paneitz operator P_θ is that it is CR covariant [16]. That is, if $\hat{\theta} = e^w \theta$, then $e^{2w} P_{\hat{\theta}} = P_\theta$.

Definition 2.2. *Let $(M^3, T^{1,0}M, \theta)$ be a pseudo-Hermitian manifold. The Paneitz type operator $P'_\theta : \mathcal{P} \rightarrow C^\infty(M)$ is defined by*

$$\begin{aligned} P'_\theta f &= 4\Delta_b^2 f - 8\operatorname{Im} \left(\nabla^\alpha (A_{\alpha\beta} \nabla^\beta f) \right) - 4\operatorname{Re} (\nabla^\alpha (R \nabla_\alpha f)) \\ &\quad + \frac{8}{3} \operatorname{Re} (\nabla_\alpha R - i \nabla^\beta A_{\alpha\beta}) \nabla^\alpha f - \frac{4}{3} f \nabla^\alpha (\nabla_\alpha R - i \nabla^\beta A_{\alpha\beta}) \end{aligned} \tag{7}$$

for $f \in \mathcal{P}$.

The main property of the operator P'_θ is its “almost” conformal covariance as shown in [2, 10]. That is if $(M^3, T^{1,0}M, \theta)$ is a pseudo-Hermitian manifold, $w \in C^\infty(M)$, and we set $\hat{\theta} = e^w\theta$, then

$$e^{2w} P'_\theta(u) = P'_\theta(u) + P_\theta(uw) \quad (8)$$

for all $u \in \mathcal{P}$. In particular, since P_θ is self-adjoint and $\mathcal{P} \subset \ker P_\theta$, we have that the operator P' is conformally covariant, mod \mathcal{P}^\perp .

Definition 2.3. *A pseudo-Hermitian manifold $(M^3, T^{1,0}M, \theta)$ is pseudo-Einstein if*

$$\nabla_\alpha R - i\nabla^\beta A_{\alpha\beta} = 0.$$

Moreover, if θ induces a pseudo-Einstein structure then $e^u\theta$ is pseudo-Einstein if and only if $u \in \mathcal{P}$. The definition above was stated in [10], but it was implicitly mentioned in [16]. In particular, if $(M^3, T^{1,0}M, \theta)$ is pseudo-Einstein, then P'_θ takes a simpler form:

$$P'_\theta f = 4\Delta_b^2 f - 8\text{Im}(\nabla^1(A_{11}\nabla^1 f)) - 4\text{Re}(\nabla^1(R\nabla_1 f)).$$

In particular, one has

$$\int_M u P'_\theta u \geq 4 \int_M |\Delta_b u|^2 - C \int_M |\nabla_b u|^2.$$

Using the interpolation inequality

$$\int_M |\nabla_b u|^2 \leq C \|u\|_{L^2} \|\Delta_b u\|_{L^2},$$

and $2ab \leq \varepsilon a^2 + \frac{1}{\varepsilon} b^2$, we have the existence of $C_1 > 0$ and $C_2 > 0$, such that

$$\int_M u P'_\theta u \geq C_1 \int_M |\Delta_b u|^2 - C_2 \int_M u^2.$$

Hence, if P'_θ is non-negative, with trivial kernel, one has the equivalence of the Folland-Stein Sobolev norm and (3).

Definition 2.4. *Let $(M^3, T^{1,0}M, \theta)$ be a pseudo-Einstein manifold. The Q' -curvature is the scalar quantity defined by*

$$Q'_\theta = 2\Delta_b R - 4|A|^2 + R^2. \quad (9)$$

The main equation that we will be dealing with is the change of the Q' -curvature under conformal change. Let $(M^3, T^{1,0}M, \theta)$ be a pseudo-Einstein manifold, let $w \in \mathcal{P}$, and set $\hat{\theta} = e^w\theta$. Hence $\hat{\theta}$ is pseudo-Einstein. Then [2, 10]

$$e^{2w} Q'_\theta = Q'_\theta + P'_\theta(w) + \frac{1}{2} P_\theta(w^2). \quad (10)$$

In particular, Q'_θ behaves as the Q -curvature for P'_θ , mod \mathcal{P}^\perp . Since we are working modulo \mathcal{P}^\perp it is convenient to project the previously defined quantities on $\hat{\mathcal{P}}$. So we define the operator $\overline{P}'_\theta = \Gamma \circ P'_\theta$ and the \overline{Q}' -curvature by $\overline{Q}'_\theta = \Gamma(Q'_\theta)$. Notice that

$$\int_M Q'_\theta \theta \wedge d\theta = \int_M \overline{Q}'_\theta \theta \wedge d\theta.$$

Moreover, the operator \overline{P}'_θ has many interesting analytical properties. Indeed, $\overline{P}'_\theta : \mathcal{P} \rightarrow \hat{\mathcal{P}}$ is an elliptic pseudo-differential operator (see [9]) and if we assume that $\ker \overline{P}'_\theta = \mathbb{R}$, then its Green's function G satisfies

$$\overline{P}'_\theta G(\cdot, y) = \Gamma(\cdot, y) - \frac{1}{V},$$

where $V = \int_M \theta \wedge d\theta$ is the volume of M and $\Gamma(\cdot, \cdot)$ is the kernel of the projection operator Γ . Moreover,

$$G(x, y) = -\frac{1}{4\pi^2} \ln(|xy^{-1}|) + \mathcal{K}(x, y),$$

where \mathcal{K} is a bounded kernel as proved in [7].

3 Preliminary results on the flow

First we recall one fundamental inequality that we will be using all along this paper, namely the CR version of the Beckner-Onofri inequality. This inequality was first proved in the odd dimensional spheres in [3] and then naturally extended to pseudo-Einstein 3-manifolds in [9, Theorem 3.1].

Theorem 3.1. *Assume that \overline{P}' is non-negative and $\ker \overline{P}' = \mathbb{R}$. Then, there exists $C > 0$ such that for all $u \in \hat{\mathcal{P}} \cap \mathcal{S}^2(M)$ with $\int_M u = 0$, we have*

$$\frac{1}{16\pi^2} \int_M u \overline{P}' u + C \geq \ln \left(\int_M e^{2u} \right).$$

In the case of the sphere, C can be taken to be 0 and equality holds if and only if $u = J(h)$ with $h \in \text{Aut}(S^3)$ and $J(h) = \det(\text{Jac}(h))$ is the determinant of the Jacobian determinant of h . The dual version of the above inequality was also investigated in [22], where the existence of extremals was investigated.

Now, we prove the global existence of solutions of (5):

Lemma 3.1. *Let $(M, T^{1,0}M, \theta)$ be a pseudo-Einstein CR three dimensional manifold such that \overline{P}' is non-negative and $\ker \overline{P}' = \mathbb{R}$. Let $f \in C(M) \cap \hat{\mathcal{P}}$ as in (4). Then for any $u_0 \in X$ there exists a solution $u \in C^\infty([0, \infty), H)$ of problem (5) such that $u(t) \in X$, for all $t \geq 0$. Moreover it holds*

$$\int_0^t \|\partial_s u(s)\|^2 ds = E(u_0) - E(u(t)),$$

for all $t \geq 0$.

Proof. Since all the functionals involved are regular, the short time existence of a solution u for (5) is ensured by the Cauchy-Lipschitz Theorem. In order to extend it to all $t \geq 0$, we notice that

$$\|\partial_t u\| = \|\nabla^X E(u)\| \leq 2\|\nabla E(u)\| \leq C_1 \|u\| + C_2.$$

Thus, since

$$\partial_t \|u\|^2 = 2\langle u, \partial_t u \rangle \leq C_3 \|u\|^2 + C_4,$$

by Gronwall's lemma, the solution u exists for all $t \geq 0$. In addition

$$\partial_t N(u) = \langle \nabla N(u), \partial_t u \rangle = -\langle \nabla N(u), \nabla^X E(u) \rangle = 0,$$

therefore $u(t) \in X$, for all $t \geq 0$. Finally, we have

$$\partial_t E(u) = \langle \nabla E(u), \partial_t u \rangle = -\|\partial_t u\|^2.$$

Hence, E is decreasing along the flow and the following energy identity holds

$$\int_0^t \|\partial_s u\|^2 ds = E(u_0) - E(u(t)). \quad (11)$$

□

Next we prove the following lemma about the convergence

Lemma 3.2. *Let $(M, T^{1,0}M, \theta)$ be a pseudo-Einstein CR three dimensional manifold such that \bar{P}' is non-negative and $\ker \bar{P}' = \mathbb{R}$. Let $f \in C(M) \cap \hat{\mathcal{P}}$ as in (4) and let u be the solution of problem (5) obtained in the previous Lemma (3.1). If there exists a constant $C > 0$ such that $\|u(t)\| \leq C$, for all $t \geq 0$, then when $t \rightarrow \infty$, $u(t) \rightarrow u_\infty$ in H and u_∞ solves the equation*

$$\bar{P}'u + \bar{Q}'u = \lambda \Gamma(fe^{2u}),$$

for a certain $\lambda \in \mathbb{R}$. Moreover, there exist constants $B, \beta > 0$ such that

$$\|u(t) - u_\infty\| \leq B(1+t)^{-\beta},$$

for all $t \geq 0$.

Proof. Since $\|u\| \leq C$, we have that

$$|E(u)| \leq 2\|u\|^2 + C_2.$$

Therefore, by the previous energy estimate

$$\int_0^\infty \|\partial_t u\|^2 dt < \infty.$$

So there exists a sequence $t_k \rightarrow \infty$ such that

$$\|\partial_t u(t_k)\| = \|\nabla^X E(u(t_k))\| \rightarrow 0.$$

Now, from the boundedness of $\|u\|$, we also have the convergence $u(t_k) \rightarrow u_\infty$ strongly in $L^2(M)$ and weakly in $\mathcal{S}^2(M)$. From Theorem 3.1, we have that $e^{2u(t_k)} \in L^p(M)$, with $p \geq 1$, and $\|e^{2u(t_k)}\|_{L^p}$ is uniformly bounded. Thus by Egorov's lemma, we can deduce that

$$\left\| fe^{2u(t_k)} - fe^{2u_\infty} \right\|_{L^p} \rightarrow 0, \quad 1 \leq p < \infty.$$

Indeed, we fix $\varepsilon > 0$. Then, there exists a set A with $\text{Vol}(A) < \varepsilon$ such that $fe^{2u(t_k)}$ converges uniformly to fe^{2u_∞} on $M \setminus A$. Therefore,

$$\begin{aligned} \left\| fe^{2u(t_k)} - fe^{2u_\infty} \right\|_{L^p} &\leq C \left\| fe^{2u(t_k)} - fe^{2u_\infty} \right\|_{L^\infty(M \setminus A)} \\ &\quad + \left(\left\| fe^{2u(t_k)} \right\|_{L^{\bar{p}}} + \left\| fe^{2u_\infty} \right\|_{L^{\bar{p}}} \right) \text{Vol}(A)^{\frac{1}{p} - \frac{1}{\bar{p}}}, \end{aligned}$$

for $p < \tilde{p} < \infty$. So the conclusion follows from the uniform boundedness of $\|e^{2u(t_k)}\|_{L^p}$, for all $1 \leq p < \infty$.

Thus $u_\infty \in X$. Now we have that

$$\nabla N(u(t_k)) = 2(\bar{P}' + I)^{-1} \Gamma \left(f e^{2u(t_k)} \right),$$

and since $f e^{2u(t_k)}$ converges strongly in $L^p(M)$ and Γ maps continuously $L^p(M)$ to $L^p(M)$ (this follows from (6) and [24]), we have by the compactness of $\bar{P}' + I$ that $\nabla N(u(t_k))$ converges strongly to $\nabla N(u_\infty)$. Also,

$$\nabla E(u(t)) = 2(\bar{P}' + I)^{-1} (\bar{P}' u + \bar{Q}') = 2u(t) + 2(\bar{P}' + I)^{-1} (\bar{Q}' - u).$$

Thus, since $\partial_t u(t_k) \rightarrow 0$ in $\mathcal{S}^2(M)$, we have that $u(t_k)$ converges strongly to u_∞ and $\nabla^X E(u_\infty) = 0$. Moreover, we have

$$(\bar{P}' + I)^{-1} (\bar{P}' u_\infty + \bar{Q}') = \lambda(u_\infty) (\bar{P}' + I)^{-1} \Gamma(f e^{2u_\infty})$$

where

$$\lambda(u_\infty) = \frac{\langle \nabla E(u_\infty), \nabla N(u_\infty) \rangle}{\|\nabla N(u_\infty)\|^2}.$$

Now by integration we have that

$$\int_M \bar{Q}' = \lambda(u_\infty) \int_M \Gamma(f e^{2u_\infty}),$$

and since $u_\infty \in X$, we have if $\int_M \bar{Q}' \neq 0$, that $\lambda(u_\infty) = 1$ and hence u_∞ solves the desired equation. On the other hand, if $\int_M \bar{Q}' = 0$, either $\lambda(u_\infty) = 0$ and thus

$$\bar{P}' u_\infty + \bar{Q}' = 0,$$

or $\lambda(u_\infty) > 0$, thus setting $v = u_\infty + \frac{1}{2} \ln(\lambda(u_\infty))$ we have

$$\bar{P}' v + \bar{Q}' = \Gamma(f e^{2v}),$$

and similarly if $\lambda(u_\infty) < 0$, we have a function $v = u_\infty + \frac{1}{2} \ln(-\lambda(u_\infty))$ such that

$$\bar{P}' v + \bar{Q}' = -\Gamma(f e^{2v}).$$

In particular, if we assume that $\int_M f e^{2\ell} \neq 0$ in the case $\lambda(u_\infty) = 0$, we have $u_\infty - \ell$ is constant. Hence,

$$0 = \int_M f e^{2u_\infty} = e^{2(u_\infty - \ell)} \int_M f e^{2\ell} \neq 0,$$

which is a contradiction.

The polynomial convergence of the flow can be deduced from the Lojasiewicz-Simon inequality following Theorem 3 in [25] and Lemma 3.2 in [1]. Let $\eta : H \rightarrow T_{u_\infty} X$ be the natural

projection, where $T_{u_\infty}X$ denotes the tangent space of the manifold X at the point u_∞ . We have, for $v \in T_{u_\infty}X$

$$(\nabla^X)^2 E(u_\infty)v = \eta \left(\nabla^2 E(u_\infty)v - \frac{\langle \nabla E(u_\infty), \nabla N(u_\infty) \rangle}{\|\nabla N(u_\infty)\|^2} \nabla^2 N(u_\infty)v + R^\perp v \right)$$

where $R^\perp v$ is the component along $\nabla N(u_\infty)$. Thus, since $\eta(R^\perp v) = 0$, we have that

$$(\nabla^X)^2 E(u_\infty)v = 2 \left(I - \eta(\bar{P}' + 1) \right) v - 4 \frac{\langle \nabla E(u_\infty), \nabla N(u_\infty) \rangle}{\|\nabla N(u_\infty)\|^2} (\bar{P}' + 1)^{-1} \Gamma(fe^{2u_\infty}v).$$

It can be checked that $(\nabla^X)^2 E(u_\infty) : T_{u_\infty}X \rightarrow T_{u_\infty}X$ is a Fredholm operator, then there exists a constant $\delta > 0$ and $0 < \kappa < \frac{1}{2}$ such that if $\|u(t) - u_\infty\| < \delta$, it holds

$$\|\nabla^X E(u)\| \geq (E(u(t)) - E(u_\infty))^{1-\kappa}.$$

We note that if $E(u(t_0)) = E(u_\infty)$ for some $t_0 \geq 0$, then the flow is stationary and the estimate is trivially satisfied. So we can assume that $E(u(t)) - E(u_\infty) > 0$, for all $t \geq 0$. Since $\lim_{n \rightarrow \infty} \|u(t_n) - u_\infty\| = 0$, for a given $\varepsilon > 0$, there exists $n_0 > 0$ such that for $n \geq n_0$ we have,

$$\|u(t_n) - u_\infty\| < \frac{\varepsilon}{2}$$

and

$$\frac{1}{\kappa} (E(u(t_n)) - E(u_\infty))^\kappa < \frac{\varepsilon}{2}.$$

We set $\varepsilon = \frac{\delta}{2}$ and

$$T := \sup \{t \geq t_{n_0}; \|u(s) - u_\infty\| < \delta; s \in [t_{n_0}, t]\},$$

and we assume for the sake of contradiction that $T < \infty$. Now we have

$$-\partial_t [E(u(t)) - E(u_\infty)]^\kappa = -\kappa \partial_t E(u(t)) [E(u(t)) - E(u_\infty)]^{\kappa-1},$$

but

$$-\partial_t E(u(t)) = -\langle E(u), \partial_t u \rangle = \|\nabla^X E(u)\| \|\partial_t u\|.$$

Thus, for $t \in [t_{n_0}, T]$ we have

$$-\partial_t [E(u(t)) - E(u_\infty)]^\kappa \geq \kappa \|\partial_t u\|,$$

and since E is non-increasing along the flow, we have after integration in the interval $[t_{n_0}, T]$

$$\|u(T) - u(t_{n_0})\| \leq \int_{t_{n_0}}^T \|\partial_s u\| ds \leq \frac{1}{\kappa} [E(u(t_{n_0})) - E(u_\infty)]^\kappa < \frac{\varepsilon}{2}.$$

Hence,

$$\|u(T) - u_\infty\| \leq \|u(T) - u(t_{n_0})\| + \|u(t_{n_0}) - u_\infty\| < \varepsilon = \frac{\delta}{2}$$

which is a contradiction and so $T = +\infty$. We set now $g(t) = E(u(t)) - E(u_\infty)$, for $t \in [t_{n_0}, +\infty)$. Then we have

$$g'(t) = -\|\nabla^X E(u)\|^2 \geq g^{2\kappa-1}(t).$$

By integration we obtain

$$g^{2\kappa-1}(t) \geq g^{2\kappa-1}(t_{n_0}) + (1 - 2\kappa)(t - t_{n_0}).$$

Since $2\kappa - 1 < 0$, then

$$g(t) \leq [g^{2\kappa-1}(t_{n_0}) + (1 - 2\kappa)(t - t_{n_0})]^{\frac{1}{2\kappa-1}} \leq Ct^{\frac{1}{2\kappa-1}}.$$

Now, by taking $t' > t$, we have

$$\|u(t) - u(t')\| \leq \int_t^{t'} \|\partial_s u\| ds \leq \frac{1}{\theta} [E(u(t)) - E(u_\infty)]^\kappa \leq \frac{1}{\kappa} g^\kappa(t) \leq Ct^{\frac{\kappa}{2\kappa-1}}.$$

For $t' = t_n$, letting $n \rightarrow \infty$ and setting $\beta = \frac{\kappa}{1-2\kappa}$, we get that for $t > t_{n_0}$

$$\|u(t) - u_\infty\| \leq Ct^{-\beta}$$

Therefore, since $\|u(t) - u_\infty\|$ is bounded for $t > t_{n_0}$, we have the existence of $B > 0$ such that for all $t \geq 0$

$$\|u(t) - u_\infty\| \leq B(1+t)^{-\beta}.$$

□

Corollary 3.1. *Let $(M, T^{1,0}M, \theta)$ be a pseudo-Einstein CR three dimensional manifold such that \bar{P}' is non-negative and $\ker \bar{P}' = \mathbb{R}$. Let $f \in C(M) \cap \hat{\mathcal{P}}$ as in (4) and let u be the solution of problem (5) obtained in the Lemma (3.1). If $\bar{u} = \frac{1}{V} \int_M u$ is uniformly bounded then the flow converges. Here $V = \int_M \theta \wedge d\theta$ is the volume of M .*

Proof. From the energy identity (11) we have that

$$\int_M u \bar{P}' u + 2 \int_M \bar{Q}' u \leq E(u_0),$$

but we also know from the Poincaré-type inequality (or the non-negativity of the operator \bar{P}'), that

$$\int_M u \bar{P}' u \geq \lambda_1 \int_M (u - \bar{u})^2.$$

Here λ_1 is the first non-zero eigenvalue of the operator \bar{P}' . In particular, from Young's inequality, we obtain that

$$\int_M u \bar{P}' u \leq E(u_0) + \varepsilon \int_M (u - \bar{u})^2 + C(\varepsilon) \|\bar{Q}'\|_{L^2}^2 - 2\bar{u} \int_M \bar{Q}'.$$

Hence, for ε small enough, we get

$$\int_M u \bar{P}' u \leq C,$$

since \bar{u} is uniformly bounded, then the uniform boundedness of $\|u\|$ and the conclusion follows from Lemma 3.2. □

Therefore, in the rest of the paper, we will show the uniform boundedness of \bar{u} along the flow, in order to have convergence at infinity.

4 The sub-critical case

Along all this section we will assume that \bar{P}' is non-negative and $\ker \bar{P}' = \mathbb{R}$. Next we consider the three separate cases in which $\int_M \bar{Q}' < 16\pi^2$. Also we let $V = \int_M \theta \wedge d\theta$ be the volume of M .

4.1 Case $\int_M \bar{Q}' < 0$ and proof of Theorem 1.1

Lemma 4.1. *There exists a positive constant $C > 0$ depending on M and θ such that for any measurable subset $K \subset M$ with $\text{Vol}(K) > 0$, we have*

$$\int_M u \leq |E(u_0)| + \frac{C}{\text{Vol}(K)} + \frac{4V}{\text{Vol}(K)} \max\left(\int_K u, 0\right)$$

Proof. Without loss of generality we can assume that $\int_M u > 0$ otherwise the inequality is trivially satisfied. First

$$\int u \bar{P}' u \leq E(u_0) - 2 \int_M \bar{Q}' u$$

and

$$\|u - \bar{u}\|_{L^2}^2 \leq \frac{1}{\lambda_1} \int_M u \bar{P}' u.$$

Hence,

$$\int_M u^2 \leq \frac{1}{\lambda_1} E(u_0) - \frac{2}{\lambda_1} \int_M \bar{Q}' u + \frac{1}{V} \left(\int_M u\right)^2$$

Now if $\int_K u \leq 0$, then we have

$$\left(\int_M u\right)^2 \leq \left(\int_{K^c} u\right)^2 \leq \text{Vol}(K^c) \int_M u^2,$$

hence

$$\frac{\text{Vol}(K)}{V} \int_M u^2 \leq \frac{1}{\lambda_1} E(u_0) - \frac{2}{\lambda_1} \int_M \bar{Q}' u.$$

Again using Young's inequality we obtain

$$\int_M u^2 \leq \frac{2V}{\lambda_1 \text{Vol}(K)} E(u_0) + \frac{4\|\bar{Q}'\|_{L^2}^2 V^2}{\lambda_1^2 \text{Vol}(K)^2},$$

but

$$\begin{aligned} \left(\int_M u\right)^2 &\leq V \int_M u^2 \leq \frac{2V^2}{\lambda_1 \text{Vol}(K)} E(u_0) + \frac{4\|\bar{Q}'\|_{L^2}^2 V^3}{\lambda_1^2 \text{Vol}(K)^2} \\ &\leq |E(u_0)|^2 + \frac{V^4}{\lambda_1^2 \text{Vol}(K)^2} + \frac{4\|\bar{Q}'\|_{L^2}^2 V^3}{\lambda_1^2 \text{Vol}(K)^2}, \end{aligned}$$

which yields

$$\int_M u \leq |E(u_0)| + \frac{C}{\text{Vol}K}.$$

We assume now that $\int_K u > 0$. Then one has

$$\int_M u^2 \leq \frac{1}{\lambda_1} E(u_0) - \frac{2}{\lambda_1} \int_M \bar{Q}' u + \frac{1}{V} \left(\left(\int_K u \right)^2 + \left(\int_{K^c} u \right)^2 + 2 \int_{K^c} u \int_K u \right),$$

and

$$\frac{2}{V} \int_{K^c} u \int_K u \leq \frac{2 \text{Vol}(K^c)}{\text{Vol}(K)V} \left(\int_K u \right)^2 + \frac{\text{Vol}(K)}{2V} \int_M u^2.$$

Hence,

$$\frac{\text{Vol}(K)}{2V} \int_M u^2 \leq \frac{1}{\lambda_1} E(u_0) - \frac{2}{\lambda_1} \int_M \bar{Q}' u + \frac{3}{\text{Vol}(K)} \left(\int_K u \right)^2.$$

By using that

$$\left| \frac{2}{\lambda_1} \int_M \bar{Q}' u \right| \leq \frac{\text{Vol}(K)}{4V} \int_M u^2 + \frac{4V \|\bar{Q}'\|_{L^2}^2}{\lambda_1^2 \text{Vol}(K)},$$

we have,

$$\int_M u^2 \leq \frac{4V}{\lambda_1 \text{Vol}(K)} |E(u_0)| + \frac{16V^2 \|\bar{Q}'\|_{L^2}^2}{\lambda_1^2 \text{Vol}(K)^2} + \frac{12V}{\text{Vol}(K)^2} \left(\int_M u \right)^2.$$

Hence,

$$\left(\int_M u \right)^2 \leq \frac{4V^2}{\lambda_1 \text{Vol}(K)} |E(u_0)| + \frac{16V^3 \|\bar{Q}'\|_{L^2}^2}{\lambda_1^2 \text{Vol}(K)^2} + \frac{12V^2}{\text{Vol}(K)^2} \left(\int_M u \right)^2,$$

and therefore

$$\int_M u \leq |E(u_0)| + \frac{C}{\text{Vol}(K)} + \frac{4V}{\text{Vol}(K)} \int_K u.$$

□

Lemma 4.2. *Let K be a measurable subset of M such that $\text{Vol}(K) > 0$. Then there exists a constant $\alpha > 1$ depending on M and θ and a constant $C_K > 1$ depending on $\text{Vol}(K)$ such that*

$$\int_M e^{2u} \leq C_K e^{\alpha \|u_0\|^2} \max \left(\left(\int_K e^{2u} \right)^\alpha, 1 \right).$$

Proof. Recall that from Theorem 3.1 one has the existence of $C > 0$ such that

$$\int_M e^{2u} \leq C \exp \left(\frac{1}{16\pi^2} \int_M u \bar{P}' u + \frac{2}{V} \int_M u \right).$$

Again, by the energy identity (11) and Young's inequality, we have

$$\begin{aligned} \int_M u \bar{P}' u &\leq E(u_0) - 2 \int_M \bar{Q}'(u - \bar{u}) - 2\bar{u} \int_M \bar{Q}' \\ &\leq E(u_0) - 2\bar{u} \int_M \bar{Q}' + \frac{1}{\varepsilon} \|\bar{Q}'\|_{L^2}^2 + \frac{\varepsilon}{\lambda_1} \int_M u \bar{P}' u. \end{aligned}$$

Thus, for $\varepsilon = \frac{\lambda_1}{2}$,

$$\frac{1}{2} \int_M u \bar{P}' u \leq E(u_0) - 2\bar{u} \int_M \bar{Q}' + \frac{2}{\lambda_1} \|\bar{Q}'\|_{L^2}^2.$$

Therefore

$$\int_M e^{2u} \leq C \exp \left(\frac{1}{8\pi^2} E(u_0) + \frac{\|\bar{Q}'\|_{L^2}^2}{4\lambda_1\pi^2} + \left(2 - \frac{1}{4\pi^2} \int_M \bar{Q}' \right) \bar{u} \right).$$

Now we notice that $E(u_0) \leq \|u_0\|^2 + \|\bar{Q}'\|_{L^2}^2$, hence there exist constants C_1 and C_2 such that

$$\int_M e^{2u} \leq C_1 \exp \left(\frac{1}{8\pi^2} \|u_0\|^2 + C_2 \int_M u \right).$$

By using Lemma 4.1, we get

$$\int_M e^{2u} \leq \bar{C}_K \exp \left(A_1 \|u_0\|^2 + \frac{A_2}{\text{Vol}(K)} \max \left(\int_K u, 0 \right) \right),$$

where \bar{C}_K depends on $\text{Vol}(K)$. Now, we set $\alpha = \max(A_1, \frac{A_2}{2}, 2) > 1$, and we get

$$\int_M e^{2u} \leq \bar{C}_K \exp \left(\alpha \|u_0\|^2 + \frac{\alpha}{\text{Vol}(K)} \max \left(\int_K 2u, 0 \right) \right).$$

But Jensen's inequality yields

$$\exp \left(\frac{1}{\text{Vol}(K)} \int_K 2u \right) \leq \frac{1}{\text{Vol}(K)} \int_K e^{2u},$$

in particular

$$\exp \left(\frac{\alpha}{\text{Vol}(K)} \max \left(\int_K u, 0 \right) \right) \leq \max \left(\left(\frac{1}{\text{Vol}(K)} \int_K e^{2u} \right)^\alpha, 1 \right).$$

Therefore, by adjusting the constant eventually

$$\int_M e^{2u} \leq C_K e^{\alpha \|u_0\|^2} \max \left(\left(\int_K e^{2u} \right)^\alpha, 1 \right)$$

which completes the proof. \square

Next we move to the proof of Theorem 1.1. We set

$$K = \left\{ x \in M; f(x) \leq \frac{1}{2} \inf_{x \in M} f(x) \right\}.$$

From the compatibility condition (i) in (4), we have that $\text{Vol}(K) > 0$, and since

$$\int_M \bar{Q}' = \int_M f e^{2u_0},$$

we obtain

$$\frac{\int_M \bar{Q}'}{\inf_{x \in M} f(x)} \leq \int_M e^{2u_0}.$$

Thus, there exists $C > 0$ (we will assume $C > 1$ actually) such that

$$\int_M e^{2u_0} \leq C \exp \left[C \left(\int_M u_0 \overline{P}' u_0 + \int_M u_0^2 \right) \right] = C e^{C \|u_0\|^2}.$$

Hence,

$$\frac{\int_M \overline{Q}'}{\inf_{x \in M} f(x)} \leq C e^{C \|u_0\|^2}. \quad (12)$$

Next we will prove the following

Lemma 4.3. *Let C_K and α be the constants found in Lemma 4.2. Let*

$$r = C_K (8C)^\alpha e^{(C+1)\alpha \|u_0\|^2},$$

and let us assume that

$$e^{\tau \|u_0\|^2} \sup_{x \in M} f(x) \leq C_0,$$

where $\tau = \alpha(C+1) - C$ and

$$C_0 = -\frac{\inf_{x \in M} f(x)}{8^\alpha C_K C^{\alpha-1}}.$$

Then for all $t \geq 0$, it holds

$$\int_M e^{2u} \leq 2r.$$

Proof. Let

$$T = \sup \left\{ s \geq 0; \int_M e^{2u} \leq 2r \text{ in } [0, s] \right\}$$

and let us assume for the sake of contradiction that $T < \infty$. We notice that by continuity, we obtain that

$$\int_M e^{2u(T)} = 2r.$$

We assume first that

$$\int_M f^+ e^{2u(T)} \leq \frac{1}{2} \int_M f^- e^{2u(T)},$$

where $f^+ := \max\{f, 0\}$ and $f^- = f^+ - f$ denote the positive and negative part of f respectively. Then we get

$$\int_M f^- e^{2u(T)} \leq -2 \int_M f e^{2u(T)} = -2 \int_M \overline{Q}' \leq -4 \int_M \overline{Q}'.$$

Since in K we have $f^-(x) \geq -\frac{1}{2} \inf_{x \in M} f(x)$, we have

$$\int_K e^{2u(T)} \leq \frac{8 \int_M \overline{Q}'}{\inf_{x \in M} f(x)}$$

which combined with (12) gives

$$\int_K e^{2u(T)} \leq 8C e^{C \|u_0\|^2}.$$

But from Lemma 4.2, we have

$$\int_M e^{2u(T)} \leq C_K e^{\alpha \|u_0\|^2} \max \left(\left(\int_K e^{2u} \right)^\alpha, 1 \right).$$

Thus

$$\int_M e^{2u(T)} \leq C_K e^{\alpha \|u_0\|^2} \left(8C e^{C \|u_0\|^2} \right)^\alpha = r,$$

which is a contradiction.

So we move to the next case, where

$$\int_M f^+ e^{2u(T)} > \frac{1}{2} \int_M f^- e^{2u(T)}.$$

Then we have

$$-\frac{1}{2} \inf_{x \in M} f(x) \int_K e^{2u(T)} \leq \int_M f^- e^{2u(T)} < 2 \int_M f^+ e^{2u(T)} \leq 4r \sup_{x \in M} f(x).$$

Hence,

$$\int_K e^{2u(T)} \leq -\frac{8r \sup_{x \in M} f(x)}{\inf_{x \in M} f(x)}.$$

By using our assumption, we obtain that

$$\int_K e^{2u(T)} \leq -\frac{8r e^{-\tau \|u_0\|^2} C_0}{\inf_{x \in M} f(x)},$$

and by Lemma 4.2, we have

$$\int_M e^{2u(T)} \leq C_K e^{\alpha \|u_0\|^2} \left(\frac{8r e^{-\tau \|u_0\|^2} C_0}{-\inf_{x \in M} f(x)} \right)^\alpha \leq r,$$

leading again to a contradiction. Hence $T = +\infty$ and $\int_M e^{2u}$ is uniformly bounded. \square

Now, by Jensen's inequality we have

$$\exp \left(\frac{1}{V} \int_M 2u \right) \leq \frac{1}{V} \int_M e^{2u} \leq \frac{2r}{V},$$

thus \bar{u} is bounded from above. Now again using the energy identity (11), we get

$$\int_M u \bar{P}' u + 2 \int_M \bar{Q}' (u - \bar{u}) + 2\bar{u} \int_M \bar{Q}' \leq E(u_0),$$

and

$$\int_M u \bar{P}' u + 2 \int_M \bar{Q}' (u - \bar{u}) \geq \frac{1}{2} \int_M u \bar{P}' u - \frac{2 \|\bar{Q}'\|_{L^2}^2}{\lambda_1} \geq -C_3.$$

Therefore

$$2\bar{u} \int_M \bar{Q}' \leq E(u_0) + C_3,$$

and since $\int_M \bar{Q}' < 0$ we have that \bar{u} is uniformly bounded from below which finishes the proof of Theorem 1.1.

4.2 Case $\int_M \bar{Q}' = 0$ and proof of Theorem 1.2

Since $\int_M \bar{Q}' = 0$, we have that

$$\langle \nabla E(u), 1 \rangle = 2 \int_M \bar{P}' u = 0$$

and

$$\langle \nabla N(u), 1 \rangle = 2 \int_M \Gamma(f e^{2u}) = 2 \int_M \bar{Q}' = 0.$$

Hence,

$$0 = \int_M \partial_t u = \partial_t \int_M u,$$

which means that the average value of u is preserved. Therefore $\bar{u} = \bar{u}_0$ and by Corollary 3.1, we have the convergence of the flow. This completes the proof of Theorem 1.2.

4.3 Case $0 < \int_M \bar{Q}' < 16\pi^2$ and proof of Theorem 1.3

First, we have again from the energy identity (11)

$$\int_M u \bar{P}' u + 2 \int_M \bar{Q}' (u - \bar{u}) + 2\bar{u} \int_M \bar{Q}' \leq E(u_0). \quad (13)$$

Hence

$$2\bar{u} \int_M \bar{Q}' \leq E(u_0) - \frac{1}{2} \int_M u \bar{P}' u + \frac{2}{\lambda_1} \|\bar{Q}'\|_{L^2}^2$$

and then \bar{u} is bounded from above; we will need a bound from below. Since $u \in X$, we get

$$\int_M \bar{Q}' = \int_M f e^{2u} \leq \|f\|_\infty \int_M e^{2u},$$

and therefore

$$\ln \left(\frac{\int_M \bar{Q}'}{\|f\|_\infty} \right) \leq \ln \left(\int_M e^{2u} \right).$$

Now again from Theorem 3.1 we have

$$\ln \left(\frac{\int_M \bar{Q}'}{\|f\|_\infty} \right) \leq C + \frac{1}{16\pi^2} \int_M u \bar{P}' u + \frac{2}{V} \int_M u. \quad (14)$$

Let $\delta > 0$ to be determined later, we sum equation (13) and $-\delta$ times equation (14), obtaining

$$\ln \left(\frac{\int_M \bar{Q}'}{\|f\|_\infty} \right) - \delta E(u_0) \leq C + \left(\frac{1}{16\pi^2} - \delta \right) \int_M u \bar{P}' u + 2 \left(1 - \delta \int_M \bar{Q}' \right) \bar{u} - 2\delta \int_M \bar{Q}' (u - \bar{u}).$$

Since $\int_M \bar{Q}' < 16\pi^2$, we choose δ such that $\int_M \bar{Q}' < \frac{1}{\delta} < 16\pi^2$, and we set

$$c_1 = 2 \left(1 - \delta \int_M \bar{Q}' \right), \quad c_2 = \delta - \frac{1}{16\pi^2}.$$

We have

$$\ln \left(\frac{\int_M \bar{Q}'}{\|f\|_\infty} \right) - \delta E(u_0) - C + c_2 \int_M u \bar{P}' u + 2\delta \int_M \bar{Q}' (u - \bar{u}) \leq c_1 \bar{u}.$$

Now we notice that

$$c_2 \int_M u \bar{P}' u + 2\delta \int_M \bar{Q}' (u - \bar{u}) \geq (c_2 \lambda_1 - \delta \varepsilon) \|u - \bar{u}\|_{L^2}^2 - \frac{\delta}{\varepsilon} \|\bar{Q}'\|_{L^2}^2,$$

therefore for ε small enough we have that

$$c_2 \int_M u \bar{P}' u + 2\delta \int_M \bar{Q}' (u - \bar{u}) \geq -c_3.$$

It follows that \bar{u} is bounded from below and therefore from Corollary 3.1 this finishes the proof.

5 The critical case and proof of Theorem 1.4

Here we will study the case $\int_M \bar{Q}' = 16\pi^2$, where $M = S^3$ is the sphere equipped with its standard contact structure. We will see S^3 as a subset of \mathbb{C}^2 with coordinates (ζ_1, ζ_2) such that

$$S^3 = \{(\zeta_1, \zeta_2) \in \mathbb{C}^2; |\zeta_1|^2 + |\zeta_2|^2 = 1\}.$$

We recall, following the notations in [3, page 15], that every C^4 conformal mapping of $\mathbb{H}^1 = \mathbb{C} \times \mathbb{R}$ comes from the action of $SU(2, 1)$ and it can be written as the composition of the following four transformations:

- left translations: $(z, t) \rightarrow (z', t') * (z, t)$, where here $*$ denotes the group operation on \mathbb{H}^1 ,
- dilations: $(z, t) \rightarrow (\delta z, \delta^2 t)$ for $\delta > 0$,
- rotations: $(z, t) \rightarrow (az, t)$, where $a \in S^1 \subset \mathbb{C}$,
- inversion: $(z, t) \rightarrow \left(-\frac{z}{|z|^2 + it}, \frac{t}{t^2 + |z|^4} \right)$.

The group of conformal transformation of the Heisenberg group \mathbb{H}^1 , also called the group of CR automorphisms, will be denoted by $Aut(\mathbb{H}^1)$. Using the Cayley transform $C : \mathbb{H}^1 \rightarrow S^3 \setminus \{(0, -i)\}$ one has a clear description of the set $Aut(S^3)$:

$$Aut(S^3) = \{C \circ h \circ C^{-1}; h \in Aut(\mathbb{H}^1)\}.$$

For $p \in S^3$ and $r \geq 1$, we will write $h_{p,r}$ the element of $Aut(S^3)$ corresponding to a Cayley transform centered at p and a dilation of size r . That is, if $h \in Aut(\mathbb{H}^1)$ is a dilation with $\delta = r$ and $C_p : \mathbb{H}^1 \rightarrow S^3 \setminus \{-p\}$ is the Cayley transform sending zero to p (instead of $(0, i)$), then $h_{p,r} = C_p \circ h \circ C_p^{-1}$. Now, for $u \in X$ we set

$$v_{p,r} = u \circ h_{p,r} + \frac{1}{2} \ln(J(h_{p,r})),$$

where we denoted $J(h) = \det(Jac(h))$, the Jacobian determinant of h . We have

$$E(v_{p,r}) = E(u) \leq E(u_0),$$

and since $u \in X$

$$\int_{S^3} f \circ h_{p,r} e^{2v_{p,r}} = \int_{S^3} f e^{2u},$$

hence

$$\int_{S^3} e^{2v_{p,r}} \geq \frac{16\pi^2}{\sup_{x \in S^3} f(x)}.$$

From [3, page 38], we know that for all $t \geq 1$ there exists $r(t) \geq 1$ and $p(t) \in S^3$ such that

$$\int_{S^3} \xi_i e^{2v_{p(t),r(t)}} = 0, \quad i = 1, 2.$$

So we let $v(t) = v_{p(t),r(t)}$ and $h(t) = h_{p(t),r(t)}$. Then using Corollary A.2 in the Appendix, one has the existence of $a < \frac{1}{16\pi^2}$ and a constant C_1 such that

$$a \int_{S^3} v(t) \overline{P}' v(t) + 2 \int_{S^3} v(t) - \ln \left(\int_{S^3} e^{2v(t)} \right) + C_1 \geq 0.$$

Since $E(v(t)) \leq E(u_0)$, we find that

$$\int_{S^3} v(t) \overline{P}' v(t) \leq C,$$

and

$$\left| \int_{S^3} v(t) \right| \leq C.$$

In particular we have that for all $p \geq 1$

$$\int_{S^3} e^{2|pv(t)|} \leq C_p,$$

and hence

$$\int_{S^3} v^2(t) \leq C$$

leading to the boundedness of $v(t)$ in H . We need the following concentration-compactness lemma in order to prove uniform boundedness.

Lemma 5.1. *Either*

(i) $\|u(t)\| \leq C$, for some constant C ;
or

(ii) there exists a sequence $t_n \rightarrow \infty$ and a point $p_0 \in S^3$ such that for all $r > 0$

$$\lim_{n \rightarrow \infty} \int_{B_r(p_0)} f e^{2u(t_n)} = 16\pi^2.$$

Moreover, for any $\tilde{x} \in S^3 \setminus \{p_0\}$, and any $r < d(\tilde{x}, p_0)$, we have

$$\lim_{n \rightarrow \infty} \int_{B_r(\tilde{x})} f e^{2u(t_n)} = 0.$$

Proof. We assume first that $r(t)$ is bounded. Then we get

$$0 < C_1 \leq J(h_{p(t), r(t)}) \leq C_2.$$

Thus, from the uniform boundedness of $v(t)$ we have

$$\int_{S^3} |u(t)| \leq C.$$

Therefore, from Lemma 3.1, it follows that $\|u(t)\|$ is uniformly bounded.

So now we assume that $r(t)$ is not bounded, then there exists a sequence $t_n \rightarrow \infty$ such that $r(t_n) \rightarrow \infty$ and without loss of generality, by compactness of S^3 we can assume that $p(t_n) \rightarrow p_0$. From the uniform boundedness of $v(t)$, we can also assume that $v(t_n) \rightarrow v_\infty$ strongly in $L^2(S^3)$ and weakly in H . We let then $r > 0$ and set $K_n = h(t_n)^{-1}(B_r(p_0))$. Then we have

$$\left| \int_{S^3} f \circ h(t_n) e^{2v(t_n)} - \int_{K_n} f \circ h(t_n) e^{2v(t_n)} \right| \leq \left(\sup_{x \in S^3} f(x) \right) \left(\text{Vol}(K_n^c) \int_{S^3} e^{4|v(t_n)|} \right)^{\frac{1}{2}}.$$

Since $h(t_n)(x) \rightarrow p_0$ a.e. then $\lim_{n \rightarrow \infty} \text{Vol}(K_n) = V$, and thus

$$\int_{B_r(p_0)} f e^{2u(t_n)} = \int_{K_n} f \circ h(t_n) e^{2v(t_n)} = \int_{S^3} f \circ h(t_n) e^{2v(t_n)} + o(1).$$

We have also

$$\int_{S^3} f \circ h(t_n) e^{2v(t_n)} = 16\pi^2,$$

and then

$$\lim_{n \rightarrow \infty} \int_{B_r(p_0)} f e^{2u(t_n)} = 16\pi^2.$$

Now if we consider $\tilde{x} \in S^3 \setminus \{p_0\}$ and $r < d(p_0, \tilde{x})$ we have that $h(t_n)(x) \notin B_r(\tilde{x})$ for n big enough, since $\lim_{n \rightarrow \infty} h(t_n)(x) = p_0$ a.e.; in particular

$$\lim_{n \rightarrow \infty} \chi_{h(t_n)^{-1}(B_r(\tilde{x}))} = 0,$$

where χ is the characteristic function. Therefore

$$\lim_{n \rightarrow \infty} \int_{B_r(\tilde{x})} f e^{2u(t_n)} = \lim_{n \rightarrow \infty} \int_{h(t_n)^{-1}(B_r(\tilde{x}))} f \circ h(t_n) e^{2v(t_n)} = 0.$$

□

Let us assume now that $\Sigma = \emptyset$. By using the previous lemma, if $\|u(t)\|$ is not uniformly bounded, then there exists $p_0 \in S^3$ such that

$$\lim_{n \rightarrow \infty} \int_{B_r(p_0)} f e^{2u(t_n)} = 16\pi^2,$$

and if $p_1 \neq p_0$ and $r < d(p_0, p_1)$, then

$$\lim_{n \rightarrow \infty} \int_{B_r(p_1)} f e^{2u(t)} = 0.$$

Since $\Sigma = \emptyset$, then there exists $g \in G$ such that $p_1 = g \cdot p_0 \neq p_0$. But

$$16\pi^2 = \lim_{n \rightarrow \infty} \int_{B_r(p_0)} f e^{2u(t)} = \lim_{n \rightarrow \infty} \int_{B_r(g \cdot p_0)} f e^{2u(t)} = \lim_{n \rightarrow \infty} \int_{B_r(p_1)} f e^{2u(t)} = 0$$

which is a contradiction. Hence $\|u(t)\|$ is uniformly bounded.

Now we assume that $\Sigma \neq \emptyset$ and that $\|u(t)\|$ is not uniformly bounded. We have that the concentration point $p_0 \in \Sigma$, otherwise we reach a contradiction arguing as in the previous case. So we obtain

$$\int_{B_r(p_0)} f e^{2u(t_n)} \leq \sup_{x \in B_r(p_0)} f(x) \int_{B_r(p_0)} e^{2u(t_n)} \leq \max \left(\sup_{x \in B_r(p_0)} f(x), 0 \right) \int_{S^3} e^{2u(t_n)}.$$

By using the the sphere version of Theorem 3.1, proved in [3], we have that

$$\frac{1}{V} \int_{S^3} e^{2u(t_n)} \leq e^{\frac{E(u(t_n))}{V}}.$$

Thus

$$\int_{B_r(p_0)} f e^{2u(t_n)} < \max \left(\sup_{x \in B_r(p_0)} f(x), 0 \right) V e^{\frac{E(u_0)}{V}}.$$

Now we first let $n \rightarrow \infty$, then $r \rightarrow \infty$ and we get

$$16\pi^2 < V \max(f(p_0), 0) e^{\frac{E(u_0)}{V}}.$$

Therefore $f(p_0) > 0$ and

$$1 < f(p_0) e^{\frac{E(u_0)}{16\pi^2}},$$

hence

$$f(p_0) > e^{-\frac{E(u_0)}{16\pi^2}},$$

which leads to a contradiction of the assumption in Theorem 1.4. Therefore we get the uniform boundedness of $\|u\|$ also in this case, which yields the convergence of the flow and it ends the proof.

A Appendix: Improved Moser-Trudinger Inequality

In what follows we will consider S^3 as a subset of \mathbb{C}^2 with coordinates (ζ_1, ζ_2) such that $|\zeta_1|^2 + |\zeta_2|^2 = 1$. We recall here the Improved Moser-Trudinger inequality introduced in [3] in order to prove the existence of a minimizer:

Proposition A.1. ([3], Proposition 3.4) *Given $\frac{1}{2} < a < 1$, there exist constants $C_1(a), C_2(a)$ such that for $u \in H$ with $\int_{S^3} \zeta_i e^{2u} = 0$, $i = 1, 2$, it holds:*

$$\frac{a}{16\pi^2} \int_{S^3} u P' u + 2 \int_{S^3} u - \ln \left(\int_{S^3} e^{2u} \right) + C_1(a) \|(-\Delta_b)^{\frac{3}{4}} u\|_2^2 + C_2(a) \geq 0$$

This improved estimate will not be useful to us in our setting since it contains the term $C_1(a) \|(-\Delta_b)^{\frac{3}{4}} u\|_2^2$ that we cannot bound along the flow. Notice that in [3], the authors exploit Ekeland's principle to exhibit a good minimizing Palais-Smale sequence that allows the control of this extra term. In our setting, we will prove a result that can be seen as intermediate between Proposition A.1 and the usual Moser-Trudinger inequality in Theorem 3.1. In fact in [3] the authors gave hints on how to prove this result, knowing that this method only works in dimension 3 and 5. We will follow a technique used in [11], since it is simpler and it allows even more improved estimates.

We set

$$P_k := \left\{ \text{polynomials of } \mathbb{C}^2 \text{ with degree at most } k \right\}$$

and

$$P_{k,0} := \left\{ f \in P_k; \int_{S^3} f = 0 \right\}.$$

For a given $m \in \mathbb{N}$ we let

$$\mathcal{N}_m := \left\{ \begin{array}{l} N \in \mathbb{N}; \exists x_1, \dots, x_N \in S^3, \nu_1, \dots, \nu_N \in \mathbb{R}^+ \text{ with } \sum_{k=1}^N \nu_k = 1 \\ \text{and for any } f \in P_{m,0}; \sum_{k=1}^N \nu_k f(x_k) = 0 \end{array} \right\}.$$

We let then $N_m = \min \mathcal{N}_m$. As it was shown in [11], one has $N_1 = 2$ and $N_2 = 4$. We recall from [3] that one has the following inequality on the standard sphere:

There exists a constant $A_2 > 0$ such that

$$\int_{S^3} \exp \left[A_2 \frac{|u - \bar{u}|^2}{\|\Delta_b u\|_{L^2}^2} \right] \leq C_0.$$

In fact the sharp constant A_2 was explicitly computed in [3] and it has the value $A_2 = 32$. With this result we can easily deduce that if $u \in \mathcal{S}^2(S^3)$ then $e^{2u} \in L^p(S^3)$ for all $1 \leq p < \infty$.

Lemma A.1. *Consider a sequence of functions $u_k \in \mathcal{S}^2(S^3)$ such that*

$$\bar{u}_k = 0, \quad \|\Delta_b u_k\|_{L^2} \leq 1$$

and suppose that $u_k \rightharpoonup u$ weakly in $\mathcal{S}^2(S^3)$ and

$$|\Delta_b u_k|^2 \rightharpoonup |\Delta_b u|^2 + \sigma \text{ in measure,}$$

where σ is a measure on S^3 . Let $K \subset S^3$ be a compact set with $\sigma(K) < 1$, then for all $1 \leq p < \frac{1}{\sigma(K)}$ we have

$$\sup_k \int_K \exp \left[p A_2 u_k^2 \right] < \infty.$$

Proof. Let φ be a fixed smooth compactly supported function on S^3 . We set $v_k = u_k - u$. Then $v_k \rightarrow 0$ strongly in L^2 and weakly in $\mathcal{S}^2(S^3)$. Now we compute

$$\begin{aligned} \int_{S^3} |\Delta_b(\varphi v_k)|^2 &= \int_{S^3} (\varphi \Delta_b v_k + v_k \Delta_b \varphi + 2\nabla_H \varphi \nabla_H v_k)^2 \\ &= \int_{S^3} \varphi^2 (\Delta_b v_k)^2 + v_k^2 (\Delta_b \varphi)^2 + 4|\nabla_H v_k \nabla_H \varphi|^2 + 2\varphi v_k \Delta_b \varphi \Delta_b v_k + \\ &\quad + 4\varphi (\nabla_H \varphi \nabla_H v_k) \Delta_b v_k + 4v_k (\nabla_H v_k \nabla_H \varphi) \Delta_b \varphi. \end{aligned} \quad (15)$$

Hence,

$$\int_{S^3} |\Delta_b(\varphi v_k)|^2 \rightarrow \int_{S^3} \varphi^2 d\sigma.$$

Assume that $1 \leq p_1 < \frac{1}{\sigma(K)}$ and take φ so that $\varphi|_K = 1$, and $\int_{S^3} \varphi^2 d\sigma < \frac{1}{p_1}$. Then we have for k large,

$$\|\Delta_b(\varphi v_k)\|_{L^2}^2 < \frac{1}{p_1}.$$

Therefore,

$$\int_K \exp \left[p_1 A_2 (v_k - \overline{\varphi v_k})^2 \right] \leq \int_{S^3} \exp \left[p_1 A_2 (\varphi v_k - \overline{\varphi v_k})^2 \right] \leq \int_{S^3} \exp \left[A_2 \frac{(\varphi v_k - \overline{\varphi v_k})^2}{\|\Delta_b \varphi v_k\|_{L^2}^2} \right] \leq C_0.$$

Thus, if we fix $\varepsilon > 0$, we can write

$$\begin{aligned} u_k^2 &= (v_k - \overline{\varphi v_k} + u + \overline{\varphi v_k})^2 \\ &= (v_k - \overline{\varphi v_k})^2 + 2(v_k - \overline{\varphi v_k})(u + \overline{\varphi v_k}) + (u + \overline{\varphi v_k})^2 \\ &\leq (1 + \varepsilon)(v_k - \overline{\varphi v_k})^2 + 2(1 + \frac{1}{\varepsilon})u^2 + 2(1 + \frac{1}{\varepsilon})^2 \overline{\varphi v_k}^2. \end{aligned}$$

Hence, given $p < \frac{1}{\sigma(K)}$ we can take $p_1 \in (p, \frac{1}{\sigma(K)})$ such that

$$\int_K e^{A_2 p_1 u_k^2} < C_0,$$

which finishes the proof. \square

Corollary A.1. *We consider the same assumptions as in Lemma A.1 and we let $\ell = \max_{x \in S^3} \sigma(\{x\}) \leq 1$. Then the following hold*

- *If $\ell < 1$, then for any $1 \leq p < \frac{1}{\ell}$, $e^{A_2 u_k^2}$ is bounded in $L^p(S^3)$. In particular $e^{A_2 u_k^2} \rightarrow e^{A_2 u^2}$ in L^1 .*
- *If $\ell = 1$, then there exists $x_0 \in S^3$ such that $\sigma = \delta_{x_0}$, $u = 0$ and after passing to a subsequence if necessary, we have*

$$e^{A_2 u_k^2} \rightharpoonup 1 + c_0 \delta_{x_0},$$

for some $c_0 \geq 0$.

Proof. Assume that $\ell < 1$ and let $1 \leq p < \frac{1}{\ell}$. Then for all $x \in S^3$, $\sigma(\{x\}) < \frac{1}{p}$. By continuity, there exists $r_x > 0$ such that $\sigma(\overline{B_{r_x}(x)}) < \frac{1}{p}$. Since S^3 is compact we can find a finite collection of balls of the form $B_{r_i}(x_i)$ such that

$$S^3 = \bigcup_{i=1}^N \overline{B_{r_i}(x_i)}.$$

So using Lemma A.1, we have

$$\sup_k \int_{B_{r_i}(x_i)} \exp \left[p A_2 u_k^2 \right] < \infty.$$

Thus,

$$\sup_k \int_{S^3} \exp \left[p A_2 u_k^2 \right] < \infty.$$

We assume now that $\ell = 1$. Since $\|\Delta_b u_k\|^2 \leq 1$ we have that $\|\Delta_b u\|^2 + \sigma(S^3) \leq 1$. Therefore, we have $u = 0$ and there exists $x_0 \in S^3$ such that $\sigma = \delta_{x_0}$. Hence, for r small, we have that

$$\sup_k \int_{S^3 \setminus B_r(x_0)} \exp \left[q A_2 u_k^2 \right] < \infty,$$

for all $q \geq 1$. Therefore, $e^{A_2 u_k^2} \rightarrow 1$ in $L^1(S^3 \setminus B_r(x_0))$ for every $r > 0$ and small. Hence, after passing to a subsequence if necessary we have that $e^{A_2 u_k^2} \rightarrow 1 + c_0 \delta_{x_0}$ in measure. \square

Proposition A.2. *Let $\alpha > 0$ and consider a sequence $m_k \rightarrow \infty$ and $u_k \in \mathcal{S}^2(S^3)$ such that $\overline{u_k} = 0$ and $\|\Delta_b u_k\|_{L^2} = 1$ such that $u_k \rightarrow u$ weakly in $\mathcal{S}^2(S^3)$ and $(\Delta_b u_k)^2 \rightarrow (\Delta_b u)^2 + \sigma$ in measure. We assume moreover that*

$$\ln \left(\int_{S^3} e^{2m_k u_k} \right) \geq \alpha m_k,$$

and

$$\frac{e^{2m_k u_k}}{\int_{S^3} e^{2m_k u_k}} \rightarrow \nu \text{ in measure.}$$

We set $R = \left\{ x \in S^3; \sigma(\{x\}) \geq A_2 \alpha \right\} = \{x_1, \dots, x_N\}$. Then $\nu = \sum_{i=1}^N \nu_i \delta_{x_i}$ with $\nu_i \geq 0$ and $\sum_i \nu_i = 1$.

Proof. Let $K \subset S^3$ such that $\sigma(K) < A_2 \alpha$. By continuity, we can find a compact set K_1 such that $K \subset \text{int}(K_1)$ and $\sigma(K_1) < A_2 \alpha$. Now given $\frac{1}{A_2 \alpha} < p < \frac{1}{\sigma(K_1)}$, we have

$$\sup_k \int_{K_1} e^{p A_2 u_k^2} \leq C_0.$$

Since $2m_k u_k \leq p A_2 u_k^2 + \frac{m_k^2}{p A_2}$, we have

$$\int_{K_1} e^{2m_k u_k} \leq C e^{\frac{m_k^2}{A_2 p}}.$$

Therefore,

$$\frac{\int_{K_1} e^{2m_k u_k}}{\int_{S^3} e^{2m_k u_k}} \leq C e^{\left(\frac{1}{A_2 p} - \alpha\right) m_k^2}.$$

So $\nu(K) \leq \nu(K_1) = 0$ and $\nu(K) = 0$. Thus, if $\sigma(\{x\}) < A_2 \alpha$, then there exists $r_x > 0$ small enough so that $\sigma(\overline{B_{r_x}(x)}) < A_2 \alpha$. Hence, $\nu(\overline{B_{r_x}(x)}) = 0$. We deduce then that $\nu(S^3 \setminus R) = 0$. Therefore

$$\nu = \sum_{k=1}^N \nu_k \delta_{x_k},$$

with $\nu_k \geq 0$ and $\sum_{k=1}^N \nu_k = 1$. □

Let $f_1, \dots, f_\ell \in C(S^3)$. We define

$$\mathcal{S}_f = \left\{ u \in \mathcal{S}^2(S^3); \bar{u} = 0; \int_{S^3} f_k e^{2u} = 0; k = 1, \dots, \ell \right\}.$$

Proposition A.3. *If $f_j \in P_{m,0}$ for $j = 0, \dots, \ell$ and $\alpha > \frac{1}{A_2 N_m}$, then there exists $C \in \mathbb{R}$ such that*

$$\ln \left(\int_{S^3} e^{2u} \right) \leq \alpha \|\Delta_b u\|_2^2 + C, \forall u \in \mathcal{S}_f.$$

Proof. We assume that the inequality

$$\ln \left(\int_{S^3} e^{2u} \right) \leq \alpha \|\Delta_b u\|_2^2 + C$$

does not hold for $u \in \mathcal{S}_f$. Then there exists a sequence $u_k \in \mathcal{S}_f$ such that

$$\ln \left(\int_{S^3} e^{2u_k} \right) - \alpha \|\Delta_b u_k\|_{L^2}^2 \rightarrow \infty.$$

Therefore, it follows that $\int_{S^3} e^{2u_k} \rightarrow \infty$ and $\|\Delta_b u_k\|_{L^2} \rightarrow \infty$. So we let $m_k = \|\Delta_b u_k\|_{L^2}$ and $v_k = \frac{u_k}{m_k}$. Then $m_k \rightarrow \infty$, $\|\Delta_b v_k\|_{L^2}^2 = 1$. Hence, after passing to a subsequence, we have

$$\begin{cases} v_k \rightharpoonup v \text{ weakly in } \mathcal{S}^2(S^3), \\ |\Delta_b v_k|^2 \rightharpoonup |\Delta_b v|^2 + \sigma \text{ in measure,} \\ \frac{e^{2m_k v_k}}{\int_{S^3} e^{2m_k v_k}} \rightharpoonup \nu \text{ in measure.} \end{cases}$$

So we let $R = \{x \in S^3; \sigma(\{x\}) \geq A_2 \alpha\} = \{x_1, \dots, x_N\}$. It follows from Proposition A.2 that $\nu = \sum_{j=1}^N \nu_j \delta_{x_j}$, with $\sum_{j=1}^N \nu_j = 1$ and $\nu_j \geq 0$.

But since $u_k \in \mathcal{S}_f$, we have

$$\int_{S^3} f_j d\nu = 0.$$

Therefore,

$$\sum_{i=1}^N \nu_i f_j(x_i) = 0, \text{ for all } 1 \leq j \leq \ell.$$

On the other hand, $A_2 \alpha N \leq 1$. In particular, if $f_j \in P_{m,0}$, we have that $N \in \mathcal{N}_m$. Therefore,

$$\alpha \leq \frac{1}{A_2 N} \leq \frac{1}{A_2 N_m}.$$

Hence, if $\alpha = \frac{1}{A_2 N_m} + \varepsilon$ we get a contradiction and the result holds. \square

Therefore, if we define

$$\mathcal{S}_0 = \left\{ u \in \mathcal{S}^2(S^3); \bar{u} = 0; \int_{S^3} f e^{2u} = 0 \text{ for all } f \in P_{1,0} \right\},$$

the following corollary holds

Corollary A.2. *There exist $a < \frac{1}{16\pi^2}$ and $C > 0$ such that for all $u \in \mathcal{P} \cap \mathcal{S}_0$, we have*

$$a \int_{S^3} u \bar{P}' u + 2 \int_M u - \ln \left(\int_M e^{2u} \right) \geq -C.$$

Indeed, this corollary follows from the fact that

$$\int_{S^3} u \bar{P}' u \geq \int_{S^3} |2\Delta_b u|^2$$

for all $u \in \hat{\mathcal{P}}$ and $8A_2 > 16\pi^2$.

References

- [1] Baird, P., Fardoun, A., Regbaoui, R.; *Q-curvature flow on 4-manifolds*, Calc. Var. 27 (2006) 75-104.
- [2] Branson, T., Gover, A.; *Conformally invariant operators, differential forms, cohomology and a generalisation of Q-curvature*, Comm. Partial Differential Equations, 30(10-12): (2005), 1611-1669.
- [3] Branson, T., Fontana, L., Morpurgo, C., *Moser-Trudinger and Beckner-Onofri's inequalities on the CR sphere*, Ann. of Math. (2), 177(1): (2013), 1-52.
- [4] Brendle, S.; *Convergence of the Q-curvature flow on S^4* , Adv. Math. 205, (2006), 1–32.
- [5] Burns, D., Epstein, C. L.; *A global invariant for three-dimensional CR-manifolds*, Invent. Math., 92(2): (1988), 333-348.
- [6] Case, J.S., Chanillo, S., Yang, P.; *A remark on the kernel of the CR Paneitz operator*, Nonlinear Analysis Volume 126, (2015), 153-158.

- [7] Case, J.S., Cheng, J.H., Yang, P.; *An integral formula for the q -prime curvature in 3-dimensional CR geometry*, Proceedings of the American Mathematical Society, 147(4), (2018), 1577-1586.
- [8] Case, J.S., Hsiao, C.Y., Yang, P.C.; *Extremal metrics for the \overline{Q}' -curvature in three dimensions*. Comptes Rendus Mathematique, Vol 354 (4), (2016), 407-410.
- [9] Case, J.S., Hsiao, C.Y., Yang, P.C.; *Extremal metrics for the Q' -curvature in three dimensions*. Journal of the European Mathematical Society, Vol 21 (2), (2019), 585-626.
- [10] Case, J.S., Yang, P.; *A Paneitz-type operator for CR pluriharmonic functions*, Bull. Inst. Math. Acad. Sin. (N.S.), 8(3): (2013), 285-322.
- [11] Chang, S.-Y., Fengbo, H.; *Improved Moser-Trudinger-Onofri inequality under constraints*, Preprint, arXiv:1909.00431
- [12] Chanillo, S., Chiu, H.L., Yang, P.; *Embeddability for 3-dimensional Cauchy-Riemann manifolds and CR Yamabe invariants*, Duke Math. J., 161(15): (2012), 2909-2921.
- [13] Cheng, J.H., Lee, J.M.; *The Burns-Epstein invariant and deformation of CR structures*. Duke Math. J. 60, no. 1, (1990), 221-254.
- [14] Frank, R.L., Lieb, E.H.; *Sharp constants in several inequalities on the Heisenberg group*. Ann. of Math. (2) 176, no. 1, 349-381, (2012).
- [15] Graham, C.R., Jenne, R., Mason, L., Sparling, G.; *Embeddability for 3-dimensional Cauchy-Riemann manifolds and CR Yamabe invariants Conformally invariant powers of the Laplacian. I. Existence*, J. London Math. Soc. (2) 46, (1992), 557-565.
- [16] Hirachi, K.; *Scalar pseudo-Hermitian invariants and the Szegő kernel on three - dimensional CR manifolds*, In Complex geometry (Osaka, 1990), volume 143 of Lecture Notes in Pure and Appl. Math., pages 67-76. Dekker, New York, 1993.
- [17] Ho, P.T.; *Prescribing the \overline{Q}' -curvature in three dimension*, Discrete Contin. Dyn. Syst. 39, (2019), 2285-2294.
- [18] Ho, P.T.; *Prescribed Q -curvature flow on S^n* , J. Geom. Phys. 62, (2012), 1233-1261.
- [19] Hsiao, C.Y.; *On CR Paneitz operators and CR pluriharmonic functions*, Math. Ann. 362: (2015), 903-929.
- [20] Lee, J.M.; *Pseudo-Einstein structures on CR manifolds*, Amer. J. Math., 110, (1988), 157-178.
- [21] Maalaoui, A.; *Prescribing the \overline{Q}' -Curvature on Pseudo-Einstein CR 3-Manifolds*, Preprint.
- [22] Maalaoui, A.; *Logarithmic Hardy-Littlewood-Sobolev Inequality on Pseudo-Einstein 3-manifolds and the Logarithmic Robin Mass*, Preprint.
- [23] Malchiodi, A., Struwe, M.; *Q -curvature flow on S^4* , J. Differential Geom. 73, (2006), 1-44.

- [24] Phong, D. H., Stein, E. M.; *Estimates for the Bergman and Szego projections on strongly pseudo-convex domains*. Duke Math. J. 44 (1977), no. 3, 695-704.
- [25] Simon, L.; *Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems*. Ann. of Math. (2) 118, no. 3, (1983), 525-571.