



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Delta sustainability from the Holocene to the Anthropocene and envisioning the future

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Anthony, E., Syvitski, J., Zăinescu, F., Nicholls, R.J., Cohen, K.M., Marriner, N., et al. (2024). Delta sustainability from the Holocene to the Anthropocene and envisioning the future. *NATURE SUSTAINABILITY*, 7(10), 1235-1246 [10.1038/s41893-024-01426-3].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/996148> since: 2024-11-10

*Published:*

DOI: <http://doi.org/10.1038/s41893-024-01426-3>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

1 **Delta sustainability from the Holocene to the Anthropocene and envisioning**  
2 **the future**

3

4 **Edward Anthony<sup>1</sup>, Jaia Syvitski<sup>2</sup>, Florin Zăinescu<sup>1,3</sup>, Robert J. Nicholls<sup>4</sup>, Kim M.**  
5 **Cohen<sup>5</sup>, Nick Marriner<sup>6</sup>, Yoshiki Saito<sup>7</sup>, John Day<sup>8</sup>, Philip S.J. Minderhoud<sup>9,10,11</sup>,**  
6 **Alessandro Amorosi<sup>12</sup>, Zhongyuan Chen<sup>13</sup>, Christophe Morhange<sup>1,14</sup>, Toru**  
7 **Tamura<sup>15,16</sup>, Alfred Vespremeanu-Stroe<sup>3</sup>, Manon Besset<sup>17</sup>, François Sabatier<sup>1</sup>,**  
8 **David Kaniewski<sup>18</sup>, Vittorio Maselli<sup>19</sup>**

9

10 <sup>1</sup>*Aix Marseille University, CNRS, IRD, INRAE, Coll France, CEREGE, Aix-en-Provence, France.*

11 <sup>2</sup>*INSTAAR & Dept. Geol. Sci., University of Colorado, Boulder CO, 80309, USA.*

12 <sup>3</sup>*Faculty of Geography, University of Bucharest, Bucharest 010041, Romania.*

13 <sup>4</sup>*Tyndall Centre for Climate Change Research, University of East Anglia, Norwich, NR4 7TJ, UK.*

14 <sup>5</sup>*Utrecht University, PO Box 80125, 3508 TC Utrecht, The Netherlands.*

15 <sup>6</sup>*CNRS, ThéMA, Université de Franche-Comté, UMR 6049, MSHE Ledoux, 32 rue Mégevand, 25030 Besançon*  
16 *Cedex, France.*

17 <sup>7</sup>*Estuary Research Center, Shimane University of Matsue, 690-8504, Japan.*

18 <sup>8</sup>*Dept. of Oceanography & Coastal Sciences, College of the Coast & Environment, Louisiana State Univ., Baton*  
19 *Rouge LA, 70803, USA.*

20 <sup>9</sup>*Soil Geography and Landscape Group, Wageningen University, Droevendaalsesteeg 3, 6708 PB Wageningen,*  
21 *The Netherlands.*

22 <sup>10</sup>*University of Padova, Department of Civil, Environmental and Architectural Engineering (ICEA), Via Marzolo,*  
23 *9 - 35131 Padova, Italy.*

24 <sup>11</sup>*Department of Subsurface and Groundwater Systems, Deltares Research Institute, Daltonlaan 600, 3584 BK*  
25 *Utrecht, the Netherlands.*

26 <sup>12</sup>*Department of Biological, Geological, and Environmental Sciences, University of Bologna, Piazza di Porta*  
27 *S.Donato 1, Bologna, Italy.*

28 <sup>13</sup>*State Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200062,*  
29 *China.*

30 <sup>14</sup>*Ecole Pratique des Hautes Etudes, PSL-AOROC, Paris, France.*

31 <sup>15</sup>*Geological Survey of Japan, AIST, Tsukuba, Ibaraki 305-8567, Japan.*

32 <sup>16</sup>*Graduate School of Frontier Sciences, Graduate School of Frontier Sciences, The University of Tokyo,*  
33 *Kashiwa, Chiba 277-8561, Japan.*

34 <sup>17</sup>*i-Sea, Bordeaux Technowest, 25 rue Marcel Issartier, 33700 Mérignac, France.*

35 <sup>18</sup>*Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), Université de Toulouse, CNRS, IRD,*  
36 *Toulouse INP, Université Toulouse 3 - Paul Sabatier (UT3), Toulouse, France*

37 <sup>19</sup>Department of Chemical and Geological Sciences, University of Modena and Reggio Emilia, 41124 Modena,  
38 Italy.

39

## 40 PREFACE

41 **River deltas offer numerous ecosystem services and host an estimated global population of**  
42 **350 to >500 million in over 100 countries. To maintain their sustainability into the future,**  
43 **deltas need to withstand sea-level rise from global warming but human pressures and**  
44 **diminishing sediment supplies are exacerbating their vulnerability. We show how deltas**  
45 **served as environmental incubators for societal development over the last 7000 years, and**  
46 **how this tightly interlocked relationship now poses challenges to deltas globally. Without**  
47 **climate stabilization, the sustainability of populous low-to mid-latitude deltas will be**  
48 **difficult to maintain, probably terminating the delta-human relationship we know today.**

49

50 Coastal river deltas (Fig. 1) offer numerous ecosystem services and resources and host  
51 growing populations in more than 100 countries, underscoring the need for a better  
52 understanding of how these landforms function. This has given rise to a remarkable corpus  
53 of studies, reports, and knowledge-driven delta-resilience organizations across a spectrum  
54 of evolving geo-, climate, ecological, and social science, and from the individual delta scale to  
55 the global scale. The human footprint spans up to 7000 years of the 8000-year  
56 evolution of modern deltas across the Holocene. Coastal space, flat topography, rich  
57 ecology, and water and other resources have provided a favourable environment for  
58 human development, but human activities are leading to global-scale vulnerability of deltas  
59 and a need for anticipation and planning<sup>1-6</sup>.

60 One of the largest human migrations in history (in raw numbers) occurred during the  
61 20th century with the rapid growth of delta cities and mega cities (many now exceed 10 M  
62 inhabitants). In 1975, the 86 largest coastal river deltas were home to about 146 million  
63 people (Fig. 2), 3.5% of the total global population of 4000 M. In 2020, the global population  
64 has almost doubled to 7800 M, but the delta population has disproportionately increased to  
65 an estimated 350 to >500 M<sup>4,7,8</sup>, outpacing at ~4.5% the global population. In 2020, this  
66 population is concentrated in ~730,000 km<sup>2</sup> of deltaic lands<sup>9</sup>, yielding a density (ranging from  
67 480 to 680 inhabitants/km<sup>2</sup>) over eight times that of Earth's habitable landmass. Global delta  
68 population is concentrated in Asia (87%). Growth is driven by large cities acting as economic  
69 motors<sup>10</sup> across the largest 86 deltas (>1000 km<sup>2</sup>) that capture 84% of the global human delta

70 population, but small deltas are also often completely urbanized<sup>9</sup>. This rapid urbanization is a  
71 product of the Anthropocene<sup>11</sup> (taken here as commencing in 1950 CE)<sup>12</sup>. Although the  
72 Anthropocene is now formally rejected (perhaps only provisionally) as a unit of geological time  
73 by the International Union of Geological Sciences<sup>13</sup>, we take the timely opportunity to refer to  
74 that decision and point out that the multi-faceted Anthropocene as a concept is here to stay.  
75 It lends itself particularly well to describing delta social-ecological systems and gives us an  
76 opportunity to conceptualise delta sustainability in a time (if not an epoch) of human  
77 dominance of global environmental change. The massive urbanization of deltas that is a  
78 product of this human dominance poses challenges to climate-change adaptation<sup>4,7,8,10,14,15</sup>.  
79 The human-delta association has become locked in a quasi-irreversible situation<sup>16</sup> for many  
80 deltas, at a time when the Anthropocene planetary transition from nature-dominance to  
81 human-dominance implies a sustainability in the balance for deltas<sup>17</sup> due to aggregated  
82 human impacts, including sea-level rise (SLR). It is hard enough creating delta megacities to  
83 cope with the influx of people, let alone deal with an environment rendered ephemeral by  
84 SLR and subject to sinking, a process intrinsic to deltas but which is now exacerbated by human  
85 activities<sup>14,18</sup>. There will be no easy fixing or undoing of this urbanization. We can renourish  
86 eroding beaches but can we remove cities from sinking deltas, pour in the sediment, and move  
87 the cities back? No, we cannot. Could the future simply consist of ‘sustaining’ deltas by  
88 manipulating sediment and water? Even doing so would not necessarily make deltas  
89 sustainable.

90 We review delta sustainability from historical through present to future perspectives  
91 conceptualizing the human-environment relationship that started as global sea level stabilized  
92 after the rapid post-glacial rise, and the strengthening of which, over time, now challenges  
93 this sustainability. We show how changing delta environments in the low- to mid-latitudes  
94 served as incubators for the Earth’s earliest political entities<sup>19</sup>, sustaining transitions in human  
95 development. We chart delta resilience over the 7000-year relationship with humans, to the  
96 current stage where humans are adversely altering the trajectory of many deltas towards  
97 perilous futures. We illustrate the future challenges of global environmental change for delta  
98 sustainability. Regarding these challenges, we draw attention to the specificity of deltas as  
99 coastal landforms, but also the distinctness of each delta, how we envisage sustainability and  
100 the obstacles to this, including what revolves around who ‘owns’ deltas, and governance and  
101 management, if they exist at all, and the role of planning. Inequalities in political-social actions

102 around delta 'ownership', governance and management will influence resilience and  
103 adaptation, creating differences between the world's deltas. All deltas are intrinsically  
104 different already, even if humans had not colonized them. But human history and cultural  
105 heritage in particular create diversified delta landscapes and their capacity to cope with  
106 change. Accessing reliable data, improved modelling, and anticipating sustainability hurdles  
107 and tipping points from intensive human occupation, exploitation and alteration of deltas, and  
108 from failing sediment supplies, should help to inform delta management and adaptation  
109 regarding projected sinking/drowning due to exacerbated subsidence and climate-induced  
110 SLR. Our review briefly frames three Holocene phases (inception, expansion, upbuilding-  
111 outbuilding deltas) of the delta-human association, hinged on an historical stable sea level  
112 with changes limited to  $\sim\pm 2$  m, followed by the Anthropocene overprint (delta vulnerability).  
113 We then chart pathways of management, planning and anticipation that we confront with an  
114 outlook on the sustainability and future of deltas.

115

## 116 **Delta inception and human encroachment**

117 About 8000 years ago, as post-glacial SLR decelerated<sup>20</sup>, accommodation space in the  
118 vicinity of some large river mouths was filled more completely, stopping their landward  
119 retreat and initiating delta formation. Accommodation space is the vertical and lateral space  
120 available for clastic sediment filling, organic matter accumulation, and freshwater bodies that  
121 counterbalance rising seas<sup>21</sup>. A delta plain traversed by distributary channels gradually  
122 developed behind changing beach coastlines up to an inland apex where it graded into the  
123 lower river valley (Fig. 1). As deltas started developing, they provided space and resources for  
124 humans<sup>19,22</sup>. The oldest human settlements on these early marshy and swampy delta plains  
125 and their coasts date 5000-4000 BCE (Before the Common Era) from radiocarbon ages and  
126 archaeological artefacts in the Danube, Rhine, Rhone, Nile, Tigris-Euphrates, Yangtze<sup>22</sup>, and  
127 the Grijalva<sup>23</sup>. The early human incursions into developing deltas were motivated by the  
128 availability of favourable lands and coastal-zone resources<sup>19</sup>, notably from harvesting lagoons  
129 and salt pans, but were also conditioned by each delta's geomorphology and sediment-  
130 dispersive dynamics, involving risks, but also possibilities for resilience to river floods and  
131 marine forces. We briefly describe in the succeeding sections a number of spikes that were to  
132 mark this relationship in the course of the 7000 years following the earliest human incursions  
133 into deltas (Fig. 3).

134

## 135 **Deltas become incubators of human progress**

136 **Delta environments and Neolithic occupation.** The start of Neolithic encroachment (Fig. 3)  
137 and the shift to sedentary occupation occurred as deltas expanded, providing wetlands for  
138 agriculture, thus favouring settlements, sedentary continuity, and food security that had not  
139 been experienced before by Mesolithic fishers<sup>24</sup>. Agricultural subsistence spread from  
140 terrestrial uplands to delta wetlands, providing aquatic diets supplemented by wetland plants  
141 and fauna. As the deltas became populous and lower river valleys infilled with sediment,  
142 hunter-gatherer subsistence was replaced by grains and fibre crops<sup>25</sup> supplemented by fish,  
143 allowing power centre cities to form, with the Tigris-Euphrates having a head start<sup>19</sup>. Within a  
144 millennium of sea-level stabilization, the Nile's originally marine flood-prone initial bayhead  
145 delta had grown large and protected enough from waves to be exploited by herding  
146 communities around 5000 BCE, and agri-cultivated by predynastic Egyptians from around  
147 4700 BCE<sup>26</sup>. This time frame is similar to that from archaeological records in the Yangtze and  
148 Yellow (Huang He) deltas<sup>27,28</sup> where rice farming and exploitation of coastal resources were  
149 fostered by a wet monsoon climate<sup>29</sup>, and in the Grijalva delta where farmers domesticated  
150 maize and possibly manioc<sup>23</sup>. Neolithic expansion in the Rhine delta began about 4300 BCE<sup>30</sup> in  
151 the wake of cultivation of valley and delta-apex floodplains and loess hillslopes upstream  
152 (5500-4500 BCE: Linear Pottery Culture). A subneolithic culture practised farming (crops and  
153 cattle) along river channels (5300 and 3400 BCE: Swifterbant Culture) and beach-ridge  
154 complexes (after 3500 BCE: Valaardingen Culture).

155 A protein-rich diet of fatty acids and staple foods fostered increasing population  
156 densities within a few hundred years after sea-level stabilization, contributing to the  
157 emergence of complex societies with increased social ranking and the construction of  
158 monumental architecture<sup>19</sup>. In the Nile delta, farming and animal husbandry played a  
159 fundamental role in establishing a robust and sustainable food system that supported the  
160 construction of the pyramid chain<sup>31</sup> along a now abandoned river branch<sup>32,33</sup>. Delta avulsions (a  
161 mechanism by which new river branches and delta lobes are created progressively or  
162 suddenly, leading to abandonment of older ones) closely conditioned settlement location  
163 choices as early as 4000-3300 BCE in the Tigris-Euphrates<sup>34</sup>. Avulsions were particularly  
164 important for the perennity of settlements in the large Pacific and Indian Ocean deltas of East  
165 and Southeast Asia, allowing for occupation of abandoned lobes<sup>27</sup>. In the Indus floodplain and

166 delta, avulsions commonly left settlements and cities without water resources, leading to their  
167 abandonment<sup>35</sup>. As avulsion-exposed deltas became more populous in the Neolithic,  
168 population centres could be more easily moved to available arable lands in adjacent river  
169 valleys<sup>19</sup> with channels less subject to avulsions.

170  
171 **Deltas foster emergence of state societies.** States originated primarily in fluvial and  
172 expanding deltaic settings in currently arid areas<sup>36</sup>, where agricultural communities supported  
173 cities that served as precursors for statehood (Fig. 3): Tigris-Euphrates: 4000-3100 BCE<sup>37</sup>; Nile:  
174 3800–3100 BCE<sup>38</sup>; Indus: 3300-2800 BCE<sup>39,40</sup>. In Asia, various archaeological cultures in the  
175 middle to lower valleys and deltas of the Yellow and Yangtze developed ca. 4000-3000 BCE,  
176 but whether these late Neolithic polities are early states remains controversial<sup>41</sup>.

177 Delta expansion was favoured by high sediment influx from river basins increasingly  
178 affected by human activities, alongside climate fluctuations<sup>42</sup>. These allocyclic (external)  
179 controls are well-evidenced by climate proxies, notably the so-called “4.2 ka event” (2150  
180 BCE), essentially an Indian Ocean Monsoonal event. This has been identified as the cause of  
181 decline of societies in some Asian deltas by affecting rice cultivation<sup>29</sup>. In the Indus valley, the  
182 4.2 ka event overlapped flourishing Harappan urbanism: between 2500 and 1900 BCE  
183 aridification may have diminished the intensity of floods, thus allowing inundation agriculture  
184 to develop across the region<sup>40</sup>. The swings in the Harappan civilization (3200-1000 BCE), from  
185 urban to rural settlements, along with the abandonment of a large number of sites, occurred  
186 between 1900 and 1000 BCE as adjustments to climate variations and water availability  
187 associated with the Monsoon<sup>43</sup>.

188  
189 **Bronze Age beginnings of delta modifications.** The Bronze Age witnessed an upsurge in  
190 human occupation of deltas, notably in the Mediterranean, marked by the establishment of  
191 trading harbours in numerous deltas<sup>44</sup>, and by transformations of delta-plain hydrology to  
192 enhance agriculture and mitigate risks. In the Arno and Serchio deltas in Italy, meandering in  
193 expanding swamps strongly influenced early Etruscan (700-500 BCE) settlement patterns,  
194 culture and society, while the Roman age (from 100 BCE onwards) saw ascendancy of human  
195 influence with wetland drainage as the modern delta plains prograded<sup>45</sup>. In the Rhine delta,  
196 clusters of farms practising trade and exchanging ceremonial goods over long distances are

197 identified from middle Bronze Age (1500-800 BCE) sites<sup>46</sup>, as a mature delta plain developed.  
198 Rhine delta farm clusters persisted in the Iron Age (800-1 BCE).

199

## 200 **Humans reinforce their control over deltas**

201 **Delta vicissitudes in Europe.** The first half of the CE witnessed increasing delta instability  
202 generated by human activities. The most noteworthy aspect of the early CE was the impact of  
203 the Roman Empire, through direct engineering of deltas, but also through this empire's  
204 influence on river sediment supply through deforestation for agriculture, roads and water  
205 harnessing. The postulate of an overarching upstream anthropogenic influence on deltas via  
206 fluvial sediment loads is embodied in the Mediterranean and Black Seas in the concept of  
207 'man-made' deltas<sup>47</sup>. For small deltas, it may be postulated that hinterland deforestation by  
208 the Romans led, within a century or so, to a progradational response, whereas the fall of the  
209 Roman Empire and the Dark Ages that followed, or the massive population decline caused by  
210 the Black Death<sup>48</sup>, all resulted in agricultural regression with forests regaining area,  
211 contributing to soil stabilization in catchments and diminished delta growth<sup>49</sup>. For large  
212 European deltas (Danube, Rhine, Rhone, Po, Ebro), growth more likely reflected a longer  
213 cumulative impact of development spanning the delta expansion and upbuilding-outbuilding  
214 phases (Bronze Age, Iron Age).

215

216 **Engineering reinforces the delta-human nexus.** Historical records from courts and  
217 monastic/ecclesiastical accounts show, in the course of the Early Middle Ages in Europe, a  
218 strategy of delta conquest that was both religious and political, especially in the Rhine<sup>50</sup>,  
219 where Roman-age settlement shows relative continuity (despite population and power shifts  
220 in the Dark Ages), and new towns and churches built along newly avulsed channels. Dyke  
221 systems along all active distributaries emerged between 1050 and 1300 CE, as bishoprics and  
222 counties implemented land reclamation campaigns to secure food production for the growing  
223 town and city populations. In the central and lower delta, and especially the northern and  
224 southern distal coastal-plain sectors, embankments and drainage of areas with organic  
225 topsoils and subsoils (peat) caused land-use sustainability problems generated by human-  
226 induced subsidence<sup>51</sup>. In the Danube catchment, important sediment release from major land-  
227 use changes caused several avulsions in the delta that resulted in the development of a  
228 southern distributary, the St. George, and the incorporation of the Greek colony of Histria, a

229 former open-coast city, into the delta plain<sup>52</sup>. The northern Chilia branch, the formation of  
230 which started in Antiquity, progressively became the largest Danube distributary, attracting  
231 new settlements along its course during the Middle Ages<sup>53</sup>.

232 In Asia, human impacts on channels and dyke-building efforts have been summarized  
233 for the Yellow delta<sup>54</sup>, a spectacular example illustrating the impact of humans on delta  
234 growth. Between 1580 and 1849, human-accelerated erosion of the Loess Plateau led to a  
235 super-elevated lower Yellow River channel bed that facilitated frequent breaching (up to 280  
236 times) of the artificial river bank levees, and sediment storage, to the tune of ~312 Gt, on the  
237 river's floodplain outside these levees<sup>55</sup>. 90% of the modern delta (i.e. since 1855 CE) is due to  
238 farming and gullying of the Loess Plateau.

239 By 1670 CE, and the start of the informal pre-industrial period, global population was  
240 about 600 M, and 50 to 70% of GDP was still devoted to basic energy resources (human food,  
241 fodder for animals, and wood fuel)<sup>12</sup>. By 1850 CE and the start of the global industrial interval  
242 (100 yr earlier in Europe), population reached 1250 M (0.8% per year growth), powered by  
243 excess energy from the combustion of fossil fuels (coal, oil) and hydroelectric plants, allowing  
244 societies to mechanize<sup>12</sup>. These changes brought increasing human pressure to bear on deltas  
245 and prompted various technological developments, including hydraulic engineering in the  
246 Po<sup>56</sup>, and management of embanked fields (polders), wind mills and pumping stations in the  
247 Rhine<sup>57</sup>.

248  
249 **Globalization of the delta-human nexus.** The industrial/colonial interval (1850–1950 CE)  
250 captures the global change in human–nature interactions and widespread occupation and  
251 transformation of deltas in North America and South America, and less than 100 years ago in  
252 Africa, the Sub-Arctic and Arctic environments, although the human footprint is, in all  
253 likelihood, as ancient in African deltas as in New World deltas<sup>23,58</sup>. The millennial-scale  
254 pressures on deltas did not initiate vulnerability as deltas generally benefited from sustained  
255 fluvial sediment supplies due to catchment deforestation by growing upland populations.  
256 Under these conditions, the relatively stable Holocene sea level (Fig. 3) constituted an  
257 important background template for delta sustainability. In Europe, deforestation and soil  
258 erosion impacts on deltas are well-documented<sup>49</sup>. In the Danube, rapidly prograding lobes  
259 formed after 1800<sup>53</sup> led to ~2.5 times higher rates of area increase compared to Middle Ages  
260 rates<sup>59</sup>. Channel instability and avulsions caused by high river sediment supply during the Little

261 Ice Age in the Rhone delta were countered by engineering modifications in the late 18<sup>th</sup>  
262 century that were a prelude to massive river-damming after the 1950s<sup>60</sup>. A similar scenario  
263 played out in many river systems and their deltas worldwide in the 19<sup>th</sup> and first half of the  
264 20<sup>th</sup> century.

265

## 266 **The Anthropocene global pressure on deltas**

267 **More populous and sediment-starved deltas.** The previous sections have shown how deltas  
268 progressively served, in the course of their growth, as incubators of human development. As  
269 humans consolidated their hold on deltas, they undertook landscape and hydraulic  
270 engineering modifications that enabled better harnessing of resources and protection against  
271 floods, erosion, and avulsions, encouraging further widespread urbanization, agriculture and  
272 engineering. These developments reinforced the 'locked-in' human-delta relationship<sup>16</sup>. The  
273 already impressive human footprint of the industrial/colonial interval is dwarfed, however, by  
274 that of the Anthropocene. Pressure on low- to mid-latitude deltas has occurred through  
275 exponential population growth (Fig. 2) that brings with it dramatic changes that strain the  
276 sustainability of deltas, whatever the breadth of their Holocene relationship with humans. A  
277 now widespread and shared global pattern of delta vulnerability prevails.

278 Humans now dominate the sediment cycle, the nitrogen cycle, the terrestrial  
279 hydrological cycle, the geochemical cycles (particularly the chalcophile elements, which have  
280 an affinity for sulfide, and more recently the platinum group elements), the planet's forest  
281 covers, ocean fish stocks, atmospheric greenhouse gases (H<sub>2</sub>O, CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>), and plant and  
282 animal density and diversity. The global warming impact of burning fossil fuels results in 20  
283 times more heat being retained by our planet than from the original energy produced during  
284 combustion<sup>12</sup>. As a result, humans have overwhelmed the planetary forcings from orbital  
285 variations in insolation, warmed the planet by >1.2°C, initiated ocean acidification, reduced  
286 sea ice volume, glacial ice mass, and permafrost, and global SLR is now at ~4 mm/yr. A high-  
287 end SSP5-8.5 scenario forecasts a median global-mean SLR of nearly 1.4 m by 2150<sup>61</sup>, setting  
288 a template for increasing delta vulnerability. Beyond 2150, sea levels will keep rising for  
289 centuries even if we stabilize climate<sup>62</sup>.

290 Population growth (Fig. 2), sediment-starvation, and human exploitation of deltas are  
291 leading to broad trends of vulnerability involving shoreline erosion and land loss<sup>63,64</sup>, elevation

292 loss<sup>18,65,66</sup>, and growing dependence on engineered flood defences and ‘lock-in’ as defined  
293 earlier<sup>16</sup>. Humans currently depend so much on long-established uses and infrastructure that  
294 it becomes extremely difficult or costly to reverse the situation, weakening resilience and  
295 creating conditions of vulnerability. A synthesis of 48 deltas revealed that 46% have a ‘lock in’  
296 relationship with humans, especially in Europe and Asia, but also in the New World<sup>16</sup>. While  
297 the Earth’s sediment production (supply) from anthropogenic soil erosion, construction  
298 activities, mineral mining, aggregate mining, and sand and gravel mining increased by about  
299 467% between 1950 and 2010, sediment transport from land to the coastal ocean (the fluvial  
300 part of which underpinned 8000 years of delta growth), has decreased by 23%, largely due to  
301 sediment trapping behind dams associated with global hydropower development<sup>67</sup>, notably  
302 in Asia-Pacific, South America and Africa<sup>68</sup>. Other human activities such as subsurface resource  
303 overexploitation, notably water and hydrocarbons, but also surface extractions of aggregates  
304 and clay increasingly cause subsidence, particularly affecting delta megacities<sup>14,18,51,69</sup>. This  
305 subsidence is no longer balanced by sedimentation<sup>3,14,18</sup>, leading to transformation of  
306 permanent or seasonal delta drylands into permanent wetlands and to shoreline retreat<sup>65,66</sup>.  
307 Many deltas are no doubt overloaded with nutrients, and, increasingly, microplastics<sup>e.g.70</sup>,  
308 leading to rapid deterioration of delta ecology and eco-services<sup>71</sup>. Channel deepening caused  
309 by sediment mining and fluvial sediment starvation<sup>e.g.72</sup> exacerbates salt intrusion in many  
310 deltas<sup>73,74</sup>. Although deltas have always been subjected to fluctuations in sediment supply that  
311 guided, in part, patterns of human occupation, the current massive diminutions in catchment  
312 sediment supply, combined with increased human-driven environmental changes, are  
313 rendering many deltas being ranked as in peril<sup>18</sup> or highly vulnerable<sup>64</sup>. SLR, under these  
314 conditions, poses a sustainability issue, and ultimately an existential threat to deltas<sup>18,75</sup>.  
315 Similar sustainability issues face the world’s estuaries<sup>76</sup>.

316

### 317 **Sustainable delta futures?**

318         Humans are now masters (wittingly or unwittingly) of the flow of water (when, where,  
319 how much), nutrients, sediment supply and redistribution, land cover and land use, urban and  
320 non-urban areas, coastal structures and protection, and energy. Humans caused the SLR, the  
321 land subsidence, and the loss of wetlands in deltas. Hence, maintaining future delta  
322 sustainability will depend on how humans, as masters of the environment, can efficiently

323 manage, if at all, the complex blend of evolving geological-climate-ecological-social science  
324 relationships that has driven the delta-human relationship over the last 7000 years, and  
325 rebuild resilience, while scaling all this down locally to individual delta social-ecological  
326 systems, each of which is distinct. A relatively stable sea level formed the background for this  
327 long relationship which now unfurls in a context of global SLR at rates into the future that are  
328 uncertain, and in a time of diminishing sediment supply. Maintaining delta sustainability raises  
329 challenging questions around the river-basin-delta governance relationship, delta 'ownership'  
330 and management, long-term planning (preferably knowledge- and data-driven and -sharing),  
331 delta distinctness, and strategies or imposed approaches into the future (Fig. 4). River basin  
332 management is key to understanding the link between climate change, local precipitation  
333 sediment supply to deltas and delta governance.

334

335 **Challenges of delta ownership and management.** The issue of 'ownership' of deltas, and the  
336 embedded questions, now and into the future, of who manages/governs a delta's health, how,  
337 and with what resources, are fundamental when considering delta sustainability. Ownership  
338 is generally defined as 'the fact of owning something'. There is an explicit link between  
339 'owning' a delta and being in a position to determine how it evolves, through some form of  
340 management, including anticipation and planning, or through no management at all. Most  
341 deltas have little or no management structure. Some deltas are managed where political  
342 systems recognise them as such, but this varies extensively with engineers, elected  
343 government representatives, wildlife/nature interests, etc., having strong roles in different  
344 deltas, and sometimes exerting little management at all. When considering the river basin,  
345 delta management always involves upstream cross-border planning and management, be it  
346 national or federal (internal) boundaries. How are management decisions made? How  
347 inclusive is the decision process? How is management funded? We raise questions that merit  
348 pondering if society is ready to examine the inequalities in, and realities and challenges of,  
349 delta sustainability into the future. But we believe, unfortunately, that society is clearly not  
350 ready to do so yet.

351

352 **Towards knowledge-driven long-term planning.** Delta planning should be integrated through  
353 a systems approach<sup>3</sup>, (re)connecting river basins to deltas and rivers to floodplains, and  
354 include management of (re)sedimentation and control of human-accelerated subsidence (Fig.

355 4), something that is being attempted in only a few deltas<sup>77,78</sup>. The feasibility and implications  
356 of re-establishing delta-plain connectivity following, for instance, strategic deployment of  
357 sedimentation-enhancing strategies<sup>79,80</sup> and nature-based solutions<sup>81</sup>, involving dialogue and  
358 knowledge-sharing<sup>5</sup> from biophysics through to legislation, should be at the forefront of  
359 interdisciplinary studies<sup>82</sup> to back planning (Fig. 4). But even here, we should restrain from  
360 over-optimism. In the Mekong delta, for instance, sedimentation-enhancing strategies could  
361 be effective against SLR but are limited by the sediment-starved situation of the delta<sup>83</sup>.  
362 Current sedimentation-enhancing strategies collectively comprise only 0.1% of the global  
363 delta area<sup>79</sup>. Unlocking the full adaptation potential of nature-based, sedimentation-  
364 enhancing strategies will require a fundamental paradigm shift in delta management to  
365 surpass biophysical and societal barriers that currently impede their widespread  
366 deployability<sup>80</sup>.

367

368 **Subsisting dataset and knowledge challenges.** Insight from big data now permeates delta  
369 studies globally. Remote sensing and modelling, in particular, confronted with the  
370 global/regional issues of climate change and regional/local anthropogenic pressures, should  
371 help us to investigate the challenges and solutions to delta sustainability. Lines of progress  
372 include accurate quantifications and projections of sediment fluxes that should provide a  
373 scientific basis for basin-wide management directives and planning<sup>e.g.84</sup>, estimates of  
374 sediment connectivity and (re)distribution processes within deltas<sup>81</sup>, and natural and human-  
375 induced subsidence<sup>69</sup>. There are, however, several areas in delta research where our  
376 knowledge is still patchy and datasets too sketchy or challenging to obtain, impacting our  
377 possibilities for reliable modeling and forecasting. There is a plethora of land-cover remote-  
378 sensing datasets that are used, for instance, to identify anthropogenic delta transformations  
379 and human occupation of subaerial delta area (Fig. 2), including megacities, agriculture,  
380 aquaculture, infrastructure, land reclamation and polders (all increasingly detrimental to  
381 mangroves and marshes), engineered distributary channels, engineered coastal barriers, and  
382 the impacts of subsidence. These datasets are useful but we still need to progress on  
383 resolution, and exert caution in data analysis and interpretation<sup>85</sup>. Standardization of datasets  
384 should also be a future goal, especially relevant in the identification of the areal limits of deltas  
385 and the distinction of delta sub-environments. Accurate delta-plain elevations and reliable  
386 projections of subsidence are also crucial for quantitative assessments of future delta

387 elevation change under SLR. High-resolution data on the elevation of most of the world's  
388 deltas, including the 86 largest deltas (Fig. 2), are currently lacking. Recent attempts in tackling  
389 this problem show that high-resolution mean delta elevations are lower than estimated using  
390 lower-resolution data<sup>86-88</sup>. The Mekong delta example (with a mean elevation of ~0.8 m above  
391 sea level, dramatically lower than the earlier erroneously assumed ~2.6 m) also underscores  
392 the fact that the quality of global coastal elevation data is inadequate and the crucial need to  
393 convert to local tidal datum is often neglected<sup>86</sup>.

394 Another challenge consists in addressing delta volume change<sup>9</sup> caused by  
395 miscellaneous human actions: organic matter production through rewilding, mangrove  
396 replanting, or reforestation, oxidation through soil drainage, empoldering and engineering,  
397 groundwater mining, peat mining, sand and gravel mining, clay extraction, deforestation,  
398 anthropogenic infrastructure. Some cause surface deformation, resulting in land subsidence  
399 in growing delta megacities that can be further exacerbated by earthquake deformation,  
400 monsoon flood weight or drought-driven shrink-swell dynamics. We also need to improve our  
401 knowledge of the subaqueous domain of deltas which can store large amounts of sediments<sup>9</sup>.  
402 Deltas are major Earth sediment sinks. Beyond the need for integration of subaqueous delta  
403 erosion into sustainability evaluations, especially under the stormier conditions  
404 accompanying climate change<sup>91</sup>, fundamental questions concern the effects of delta sediment  
405 load changes on continental margin geological (e.g., volcanic activity) and sea-level feedbacks,  
406 hence providing a link between local (river basin-delta) processes and global regulation. Geo-  
407 engineering of individual deltas has been ongoing for at least 5000 years. The current global  
408 situation suggests that regulating industrial waste outputs is a necessary step in mitigating  
409 environmental damage. Managing deltas is an obvious component of such mitigation, at least  
410 as a source of data, but also as a means of managing inputs and thresholds in the Earth system.

411  
412 **Sustainability uncertainties into the future.** Innovative knowledge-driven delta management  
413 and planning strategies, but also data acquisition and modelling<sup>3,4</sup> are in their infancy but,  
414 where feasible, now and into the future, could provide sustainable options for deltas against  
415 near future (low-end SSP) projected SLR, including low rates of land subsidence (Fig. 4). There  
416 are, however, other potential obstacles here, in addition to those related to datasets and  
417 knowledge acquisition. Individual deltas are distinct entities, each with unique boundary  
418 conditions, and a unique history of human change and impacts. Large deltas may in fact

419 display physical, cultural and human-history diversity even within their individual boundary  
420 conditions. This complicates the deployment of general 'models' of sustainability. Alongside  
421 this difficult outlook, management strategies are simply not presently feasible for most deltas,  
422 reflecting lack of resources and planning and management capacity (Fig. 4). This raises the  
423 question of human capacity-building regarding better knowledge of delta functioning and  
424 management<sup>3</sup>, but also of harnessing better *indigenous knowledge*<sup>90</sup> and its perspectives.  
425 Accessing reliable data is still a problem in many deltas due to geopolitical sensitivities, and  
426 yet important not only for management, planning and anticipation, but also for gauging  
427 tipping points in the delta-human relationship. The diversity of the Earth's deltas will require  
428 high-quality field observations to inform important and often costly environmental decisions,  
429 as well as community-level information with citizens conversant with the finer-scale changes  
430 that affect their daily livelihoods<sup>6,9</sup>, especially in the populous deltas. Transferable lessons can  
431 also be identified, and, where possible, deployed to improve climate resilience<sup>5</sup>. These  
432 include, from the Ganges-Brahmputra and Mekong deltas, strategic plans to identify risk  
433 hotspots, guide decision-making and enhance grassroots resilience through community  
434 livelihood diversification in response to changing risks and land-ater conditions, and from the  
435 Yangtze and Pearl deltas forecasting and sensing technologies developed to enable effective  
436 preparedness for, and response to, hazards<sup>5</sup>.

437         Climate control to mitigate SLR<sup>62</sup>, sustained fluvial sediment supplies, control of  
438 human-induced land subsidence, and delta population, set to attain by 2050 an averaged  
439 global density<sup>4</sup> of ~700 inhabitants/km<sup>2</sup>, are, however, sources of future constraints. The  
440 sustainability of many low-to-mid-latitude deltas will be severely affected by relative  
441 SLR<sup>3,4,7,8,10,14,75</sup> in the absence of climate stabilization, and compounded by fluvial sediment  
442 starvation<sup>91</sup> and accelerated land subsidence<sup>3,14</sup>. Growing hydropower dam constructions in  
443 developing economies<sup>68</sup> will further negatively impact sediment supply to deltas in the future.  
444 Basin-wide planning of sediment releases from dams will need to be thoroughly gauged and  
445 calibrated, notably by resorting less to the widespread use of large dead storages (the portion  
446 of the reservoirs that cannot be emptied) in dam designs, and designing smaller dead storages  
447 that can ease sediment starvation in sinking deltas<sup>92</sup>.

448         In spite of a high loss of fluvial connectivity due to dams and engineering, some deltas  
449 associated with Asia-Pacific rivers still continue to gain land<sup>64</sup>, as dams are flushed of sediment  
450 to increase the calculated yield<sup>e.g.93</sup>. Rapid SLR will also outpace marsh and mangrove

451 growth<sup>21,94</sup>, important components of sedimentation in many deltas. Low-population Arctic  
452 deltas with increasing climate-change-induced sediment loads<sup>95</sup> may be a temporary  
453 exception regarding their sediment budgets but could also become increasingly exposed to  
454 anthropogenic pressures with climate warming. Sustainability will depend on our capacity to  
455 mitigate climate change and global SLR, while differences in current and future anthropogenic  
456 pressures on individual deltas and inequalities in political-social actions addressing them will  
457 also strongly influence the effectiveness of local mitigation and adaptation measures (Fig. 4).  
458 The recent United Nations Convention on Conserving River Deltas (UNCCRD) initiative  
459 proposed by engaged scientists at the COP28 in 2023 is an important global endeavour that  
460 could consolidate our efforts, but properly enacting this convention could take, at best,  
461 several years. The *International Panel on Deltas and Coastal Areas*  
462 (<https://www.deltasandcoasts.net>), launched in 2023, could also promote sustainability  
463 efforts.

464

465 **Expected outcomes without climate mitigation.** In the crucial battle against the inevitable  
466 SLR, three end-member strategies<sup>(e.g.96)</sup>, alongside a ‘laissez-faire’ (a term borrowed from  
467 economists) approach, are currently deployed and/or envisaged for coasts in general.  
468 However, we need to recognize the biophysical specificities of deltas (Box 1) which go beyond  
469 just the coastline fringe. These strategies/approaches are not mutually exclusive. ‘Protect’ and  
470 ‘accommodate’ strategies are costly, impact delta biophysics, and larger and larger areas are  
471 threatened with deeper floods if protection fails, especially for the higher-emission scenarios  
472 (SSP3-7.0 and SSP5-8.5) (Box 1). Even for wealthy economies, dedicated to containing the  
473 effects of SLR, such as the Netherlands with the Rhine delta<sup>97,98</sup>, or the United States with the  
474 Mississippi<sup>99</sup>, this outcome is undesirable (Fig. 5) and protection that works with delta  
475 processes are more desirable. Both absolute SLR and the annual rate of rise pose challenges,  
476 the latter being susceptible to reduce, for instance, the lifetime of defence constructions when  
477 the rate of SLR rise increases beyond projected values<sup>98</sup>. Assuming no protection measures,  
478 deltas globally might lose 5% (35,000 km<sup>2</sup>) of their area by 2100 and 50% by 2300 due to SLR  
479 under the high-emission scenarios<sup>75</sup>. Large-scale marine inundation, scaled against prohibitive  
480 adaptation costs<sup>10,99</sup>, will impose generalized give-up and human retreat (Fig. 5). Drastic  
481 wholesale urban migrations and landward redeployments in sinking and marine-inundated  
482 deltas may become more frequent in the future: the population of New Orleans has never

483 recovered post-Katrina. Djakarta, 40% of which is now below present sea level on the sinking  
484 Ciliwung-Citarum delta, 4<sup>th</sup> in world conurbation population ranking (30 M), and Bangkok 13<sup>th</sup>  
485 (18 M) on the sediment-starved Chao Phraya delta, are considering moving their city rather  
486 than engaging in costly engineering for their survival. In some delta areas subject to extremely  
487 high subsidence rates (>10 cm/yr) that threaten sustainability, such as the Semarang-Demak  
488 in northern Java or the Pampanga in the Philippines, reports show that entire drowning villages  
489 have simply become abandoned in 5-10 years. Significant movements of people away from  
490 deltas may be anticipated and retreat managed in delta zones likely to be most exposed to  
491 sea-level rise and/or subsidence.

492

493 The 7000-year relationship of deltas with humans fostered technological  
494 developments geared at water control and the fight against subsidence, erosion and the sea.  
495 These developments, together with new technologies, strategies, and data, will be  
496 instrumental in the battle of sustaining our deltas, and maintaining sustainable, if not entirely  
497 habitable deltas with SLR. Pathways of sustainability and survival in the populous low-to-mid-  
498 latitude deltas will need to be confronted with paradigms and tough challenges revolving  
499 around dedicated and coordinated governance, management, planning, at both river-basin  
500 and delta levels, and subsidence control, without losing sight of the distinctness of each delta,  
501 but also of the diversity within some large deltas. Without climate control, an extreme SLR  
502 scenario (rising up to and >2 m) over the next two centuries will lead to progressive delta  
503 drowning, imposing untenable conditions from both environmental and economic  
504 standpoints for human occupation, leading to global-scale human retreat from deltas. This  
505 would terminate the 7000-year mutual relationship of humans with deltas as we know and  
506 live it today, and establish a future of living with drowning and drowned deltas.

507

## 508 **References**

- 509 1. Brondizio E.S. *et al.* Catalyzing action towards the sustainability of deltas. *Current Opinion*  
510 *in Environmental Sustainability*, **19**, 182-194 (2016).  
511 <https://doi.org/10.1016/j.cosust.2016.05.001>
- 512 2. Loucks, D.P. Developed river deltas: are they sustainable? *Environ. Res. Lett.* **14**, 113004  
513 (2019). DOI 10.1088/1748-9326/ab4165
- 514 3. Schmitt, R.J.P. & Minderhoud, P. Data, knowledge, and modeling challenges for science-  
515 informed management of river deltas. *One Earth* **6**, 3 (2023).  
516 <https://doi.org/10.1016/j.oneear.2023.02.010>

517 **Review of knowledge and data needed to better promote sustainable delta management**

- 518 4. Scown, M.W. *et al.* Global change scenarios in coastal river deltas and their sustainable  
519 development implications. *Global Environmental Change*, **82**, 102736 (2023).  
520 <https://doi.org/10.1016/j.gloenvcha.2023.102736>
- 521 **Scenarios of delta change in the near future and implications for delta sustainability**
- 522 5. Chan, F.K.S. *et al.* Building resilience in Asian mega-deltas. *Nat. Rev. Earth Environ.* (2024).  
523 <https://doi.org/10.1038/s43017-024-00561-x>
- 524 **Review of challenges and transferable lessons among five Asian megadeltas to enhance**  
525 **resilience**
- 526 6. Cremin, E. *et al.* Causes and consequences of tipping points in river delta social–ecological  
527 systems. *Ambio* (2024). <https://doi.org/10.1007/s13280-023-01978-2>
- 528 7. Edmonds, D.A., Caldwell, R.L., Brondizio, E.S. & Siani, S.M.O. Coastal flooding will  
529 disproportionately impact people on river deltas. *Nat. Commun.* **11**, 4741 (2020).  
530 <https://doi.org/10.1038/s41467-020-18531-4>
- 531 **Global analysis of the potential impacts of flooding from sea-level rise on delta populations**
- 532 8. Nicholls, R.J., Adger, W.N., Hutton, C.W. & Hanson, S.E. (eds.). *Deltas in the Anthropocene*  
533 (Springer, 2020).
- 534 9. Syvitski, J. *et al.* Large deltas, small deltas: Towards a more rigorous understanding of  
535 marine deltas. *Global & Planetary Change* **103958** (2022).  
536 <https://doi.org/10.1016/j.gloplacha.2022.103958>
- 537 10. Hauer, M.E. *et al.* Sea-level rise and human migration. *Nat. Rev. Earth Environ.* **1**, 28–39  
538 (2020). <https://doi.org/10.1038/s43017-019-0002-9>
- 539 **Analysis of drivers and determinants of human migration to coasts subject to sea-level rise**
- 540 11. Wallenhurst, N. & Wulf, C. (eds.) *Handbook of the Anthropocene: Humans between*  
541 *Heritage and Future*. Springer (2020).
- 542 **Exhaustive analysis of the multi-faceted Anthropocene from all angles**
- 543 12. Syvitski, J. *et al.* Extraordinary human energy consumption and resultant geological  
544 impacts beginning around 1950 CE initiated the proposed Anthropocene Epoch. *Commun.*  
545 *Earth Environ.* **1**, 32 (2020). <https://doi.org/10.1038/s43247-020-00029-y>.
- 546 13. Witze, A. Geologists reject the Anthropocene as Earth’s new epoch — after 15 years of  
547 debate. *Nature* **627**, 249-250 (2024). doi: <https://doi.org/10.1038/d41586-024-00675-8>
- 548 14. Nicholls, R.J. *et al.* A global analysis of subsidence, relative sea-level change and coastal  
549 flood exposure. *Nat. Clim. Chang.* **11**, 338–342 (2021). [https://doi.org/10.1038/s41558-](https://doi.org/10.1038/s41558-021-00993-z)  
550 [021-00993-z](https://doi.org/10.1038/s41558-021-00993-z)
- 551 **Global analysis of coastal and delta exposure to flooding from sea-level rise and subsidence**
- 552 15. McGranahan, G., Balk, D., Colenbrander, S., Engin, H., & MacManus, K. Is rapid  
553 urbanization of low-elevation deltas undermining adaptation to climate change? A global  
554 review. *Environment and Urbanization*, **35** (2), 527-559 (2023).  
555 <https://doi.org/10.1177/09562478231192176>
- 556 **Global review of obstacles to climate change adaptation due to rapid delta urbanization**
- 557 16. Santos, M.J. & Dekker, S.C. Locked-in and living delta pathways in the Anthropocene. *Sci.*  
558 *Rep.* **10** (1), 19598 (2020). <https://doi.org/10.1038/s41598-020-76304-x>
- 559 **Analysis of the intertwined Anthropocene relationship between many deltas and humans**
- 560 17. Day, J.W. *et al.* Approaches to defining deltaic sustainability in the 21st century. *Estuarine,*  
561 *Coastal and Shelf Science*, **183**, 275-291 (2016).  
562 <https://doi.org/10.1016/j.ecss.2016.06.018>.
- 563 18. Syvitski, J. *et al.* Sinking deltas due to human activities. *Nat. Geosci.* **2**, 681–686 (2009).  
564 <https://doi.org/10.1038/ngeo629>

- 565 19. Day, J.W., Gunn, J.D., Folan, W.J., Yáñez-Arancibia, A. & Horton, B.P. The influence of  
566 enhanced post-glacial coastal margin productivity on the emergence of complex societies.  
567 *The Journal of Island and Coastal Archaeology*, **7** (1), 23-52 (2012).  
568 <http://dx.doi.org/10.1080/15564894.2011.650346>
- 569 20. Lambeck, K., Rouby, H., Purcell, A., Sun, Y. & Sambridge, M. Sea level and global ice  
570 volumes from the Last Glacial Maximum to the Holocene. *Proc. Natl. Acad. Sci. U.S.A.* **111**,  
571 15296–15303 (2014). <https://doi.org/10.1073/pnas.1411762111>
- 572 21. Osland, M.J. et al. Migration and transformation of coastal wetlands in response to rising  
573 seas. *Sci. Adv.* **8**, eabo5174 (2022). DOI: 10.1126/sciadv.abo5174
- 574 22. Stanley, D.J. & Warne, A.G. Holocene sea-level change and early human utilization of  
575 deltas. *GSA Today* **7**, 1–7 (1997).
- 576 **An early analysis of the first links between deltas and human encroachment globally**
- 577 23. Pope, K.O. et al. Origin and environmental setting of ancient agriculture in the low-lands  
578 of Mesoamerica. *Science* **292**, 1370–1373 (2001). DOI: 10.1126/science.292.5520.1370
- 579 24. Zong, Y. et al. Fire and flood management of coastal swamp enabled first rice paddy  
580 cultivation in east China. *Nature* **449**, 459–462 (2007).  
581 <https://doi.org/10.1038/nature06135>
- 582 25. Scott, J.C. *Against the Grain: A Deep History of the Earliest States*. (Yale University Press,  
583 2017).
- 584 26. Stanley, D.J. & Warne, A.G. Sea level and initiation of Predynastic culture in the Nile delta.  
585 *Nature* **363**, 435–438 (1993). <https://doi.org/10.1038/363435a0>
- 586 27. Chen, Z., Zong, Y., Wang, Z., Wang, H., & Chen, J. Migration patterns of Neolithic  
587 settlements on the abandoned Yellow and Yangtze River deltas of China. *Quaternary*  
588 *Research*, **70** (2), 301-314 (2008). doi:10.1016/j.yqres.2008.03.011
- 589 28. Wang, Z., Zhuang, C., Saito, Y., Chen, J. & Zhan, Q. Early mid-Holocene sea-level change  
590 and coastal environmental response on the southern Yangtze delta plain, China:  
591 implications for the rise of Neolithic culture. *Quaternary Science Reviews*, **35**, 51-62 (2012).  
592 <https://doi.org/10.1016/j.quascirev.2012.01.005>
- 593 29. Deng, L.J. et al. New archaeobotanical evidence reveals synchronous rice domestication  
594 7600 years ago on south Hangzhou Bay coast, eastern China. *Anthropocene*, **33**, 100280  
595 (2021). [10.1016/j.ancene.2021.100280](https://doi.org/10.1016/j.ancene.2021.100280)
- 596 30. Louwe Kooijmans, L.P., van den Broeke, W., Fokkens, H. & van Gijn, A.I. *The Prehistory of*  
597 *the Netherlands*. (Amsterdam University Press, 2005), 844 pp.
- 598 31. Sheisha, H. et al. Feeding the pyramid builders: Early agriculture at Giza in Egypt.  
599 *Quaternary Science Reviews*, **312**, 108172.  
600 <https://doi.org/10.1016/j.quascirev.2023.108172>.
- 601 32. Sheisha, H. et al. Nile waterscapes facilitated the construction of the Giza pyramids during  
602 the 3rd millennium BCE. *Proceedings of the National Academy of Sciences*, **119**,  
603 e2202530119 (2022). <https://doi.org/10.1073/pnas.2202530119>
- 604 33. Ghoneim, E. et al. The Egyptian pyramid chain was built along the now abandoned  
605 Ahramat Nile Branch. *Commun. Earth Environ.* **5**, 233 (2024).  
606 <https://doi.org/10.1038/s43247-024-01379-7>
- 607 34. Morozova, G. A review of Holocene avulsions of the Tigris and Euphrates rivers and  
608 possible effects on the evolution of civilizations in lower Mesopotamia. *Geoarchaeology*,  
609 **20** (4), 401-423 (2005). <https://doi.org/10.1002/gea.20057>

- 610 35. Syvitski, J., Overeem, I., Brakenridge, G.R. & Hamon, M. Floods, floodplains, delta plains:  
611 A satellite imaging approach. *Sedimentary Geology* **267–268**, 1–14 (2012).  
612 <https://doi.org/10.1016/j.sedgeo.2012.05.014>
- 613 36. Macklin, M.G. & Lewin, J. The rivers of civilization. *Quaternary Science Reviews* **114**, 228–  
614 244 (2015). <https://doi.org/10.1016/j.quascirev.2015.02.004>
- 615 37. Marchetti, N. *et al.* Long-Term urban and population trends in the Southern  
616 Mesopotamian floodplains. *J. Archaeol. Res.* (2024). [https://doi.org/10.1007/s10814-024-](https://doi.org/10.1007/s10814-024-09197-3)  
617 [09197-3](https://doi.org/10.1007/s10814-024-09197-3)
- 618 38. Dee, M. *et al.* An absolute chronology for early Egypt using radiocarbon dating and  
619 Bayesian statistical modelling. *Proc. R. Soc. A. Math. Phys. & Eng. Sci.* **469** 20130395,  
620 (2013). <http://doi.org/10.1098/rspa.2013.0395>
- 621 39. Meadow, R.H. & Kenoyer, J.M. Excavations at Harappa 2000-2001: new insights on  
622 chronology and city organization. *South Asian Archaeology*, CNRS, Paris 207-225 (2001).
- 623 40. Wright, R.P. *The Ancient Indus: Urbanism, Economy, and Society*. Case Studies in Early  
624 Societies, Series Number 10, 418 pp. (Cambridge University Press, 2010).
- 625 41. Liu, L. State emergence in Early China. *Annual Reviews of Anthropology* **38**, 217–32 (2009).  
626 <https://doi.org/10.1146/annurev-anthro-091908-164513>
- 627 42. Walsh, K. *et al.* Holocene demographic fluctuations, climate and erosion in the  
628 Mediterranean: A meta data-analysis. *The Holocene* **29** (5) 864–885 (2019).  
629 <https://doi.org/10.1177/0959683619826637>
- 630 43. Coningham, R., Young, R. *The Archaeology of South Asia: From the Indus to Asoka, c. 6500*  
631 *BCE – 200 CE* (Cambridge University Press, 2015).
- 632 44. Giaime, M. *et al.* Evolution of ancient harbours in deltaic contexts: A geoarchaeological  
633 typology. *Earth-Science Reviews* **191**, 141-167 (2019).  
634 <https://doi.org/10.1016/j.earscirev.2019.01.022>
- 635 45. Amorosi, A. *et al.* Middle to late Holocene environmental evolution of the Pisa coastal  
636 plain (Tuscany, Italy) and early human settlements. *Quat. Int.* **303**, 93–106 (2013).  
637 <https://doi.org/10.1016/j.quaint.2013.03.030>
- 638 46. Fontijn, D. Land at the other end of the sea? Metalwork circulation, geographical  
639 knowledge and the significance of British/Irish imports in the Bronze Age of the Low  
640 Countries. In *Bronze Age Connections: Cultural Contact in Prehistoric Europe* (ed. Clark, P.),  
641 129–148. (Oxbow Books, 2009).
- 642 47. Maselli, V., Trincardi, F. Man-made deltas. *Sci. Rep.* **3**, 1926 (2013).  
643 <https://doi.org/10.1038/srep01926>
- 644 48. Kelly, J. *The Great Mortality. An Intimate History of the Black Death, the most devastating*  
645 *Plague of all Time* (New York, Harper Collins, 2006).
- 646 49. Anthony, E.J., Marriner, N. & Morhange, C. Human influence and the changing  
647 geomorphology of Mediterranean deltas and coasts over the last 6000 years: from  
648 progradation to destruction phase? *Earth-Science Reviews* **139**, 336-361 (2014).  
649 <https://doi.org/10.1016/j.earscirev.2014.10.003>
- 650 50. van Dinter, M. The Roman Limes in the Netherlands: how a delta landscape determined  
651 the location of the military structures. *Netherlands Journal of Geosciences* **92** (1), 11-32  
652 (2013). <https://doi.org/10.1017/S0016774600000251>
- 653 51. Erkens, G. van der Meulen, M.J. & Middelkoop, H. Double trouble: subsidence and CO2  
654 respiration due to 1,000 years of Dutch coastal peatlands cultivation. *Hydrogeology*  
655 *Journal* **24**, 551 (2016). DOI: 10.1007/s10040-016-1380-4

- 656 52. Vespremeanu-Stroe, A. *et al.*, 2013. The impact of the Late-Holocene coastal changes on  
657 the rise and decay of the ancient city of Histria (Southern Danube delta). *Quaternary*  
658 *International* **293**, 245–256 (2013). <https://doi.org/10.1016/j.quaint.2012.11.039>
- 659 53. Preoteasa, L. *et al.* Late-Holocene landscape evolution and human presence in the  
660 northern Danube delta (Chilia distributary lobes). *The Holocene* **31** (9), 1459-1475 (2021).  
661 <https://doi.org/10.1177/09596836211019121>
- 662 54. Chen, Z., Syvitski, J., Gao, S. Overeem, I. & Kettner, A.J. Socio-economic impacts on  
663 flooding: a 4000-year history of the Yellow River, China. *AMBIO* **41** (7), 682-698 (2012).  
664 doi: 10.1007/s13280-012-0290-5
- 665 55. Chen, Y. *et al.* Quantifying sediment storage on the floodplains outside levees along the  
666 lower Yellow River during the years 1580–1849. *Earth Surf. Proc. & Landf.* **44** (2), 581-594.  
667 <https://doi.org/10.1002/esp.4519>
- 668 56. Simeoni, U. & Corbau, C. A review of the Delta Po evolution (Italy) related to climatic  
669 changes and human impacts. *Geomorphology* **107**, 64–71 (2009).  
670 <https://doi.org/10.1016/j.geomorph.2008.11.004>
- 671 57. van Koningsveld, M., Mulder, J.P.M., Stive, M.J.F., van der Valk, L. & van der Weck,  
672 A.W. Living with sea-level rise and climate change: a case study of the Netherlands. *Journal*  
673 *of Coastal Research*, 24 (2), 367-379 (2008). <https://doi.org/10.2112/07A-0010.1>
- 674 58. Chamberlain, E.L., Mehta, J.M., Reimann, T. & Wallinga, J. A geoarchaeological perspective  
675 on the challenges and trajectories of Mississippi Delta communities. *Geomorphology* **360**,  
676 107132 (2020). <https://doi.org/10.1016/j.geomorph.2020.107132>
- 677 59. Vespremeanu-Stroe A. *et al.* Holocene evolution of the Danube delta: An integral  
678 reconstruction and a revised chronology. *Marine Geology* **388**, 38–61 (2017).  
679 <https://doi.org/10.1016/j.margeo.2017.04.002>
- 680 60. Provansal, M. *et al.* The geomorphic evolution and sediment balance of the Lower Rhône  
681 River (southern France) over the last 130 years: hydropower dams versus other control  
682 factors. *Geomorphology* **219**, 27-41 (2015).  
683 <https://doi.org/10.1016/j.geomorph.2014.04.033>
- 684 61. Fox-Kemper, B. *et al.* Ocean, Cryosphere and Sea Level Change. In *Climate Change 2021:*  
685 *The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report*  
686 *of the Intergovernmental Panel on Climate Change* (eds. Masson-Delmotte, V. *et al.*).  
687 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1211–  
688 1362 (2021). doi:10.1017/9781009157896.011.
- 689 62. Nicholls, R.J. *et al.* Stabilization of global temperature at 1.5°C and 2.0°C: implications for  
690 coastal areas. *Phil. Trans. R. Soc. A* **376**, 20160448 (2018).  
691 <http://dx.doi.org/10.1098/rsta.2016.0448>
- 692 63. Anthony, E.J. *et al.* Linking rapid erosion of the Mekong River delta to human activities.  
693 *Scientific Reports*, **5**, 14745 (2015). <https://www.nature.com/articles/srep14745>
- 694 64. Besset, M., Anthony, E.J. & Bouchette, F. Multi-decadal variations in delta shorelines and  
695 their relationship to river sediment supply: An assessment and review. *Earth-Science*  
696 *Reviews* **193**, 199-219. <https://doi.org/10.1016/j.earscirev.2019.04.018>
- 697 65. Minderhoud, P. S. J. *et al.* The relation between land use and subsidence in the Vietnamese  
698 Mekong delta. *Science of The Total Environment*, **634**, 715–726 (2018).  
699 <https://doi.org/10.1016/j.scitotenv.2018.03.372>
- 700 66. Anthony, E.J., Besset, M., Zăinescu, F. & Sabatier, F. Multi-decadal deltaic land-surface  
701 changes: Gauging the vulnerability of a selection of Mediterranean and Black Sea river  
702 deltas. *J. Mar. Sci. Eng.* **9** (5) 512 (2021). <https://doi.org/10.3390/jmse9050512>

- 703 67. Syvitski, J. *et al.* Earth's sediment cycle during the Anthropocene. *Nat. Rev. Earth Environ.*  
704 **3**, 179–196 (2022). <https://www.nature.com/articles/s43017-021-00253-w>.
- 705 68. Gernaat, D.E.H.J. *et al.* High-resolution assessment of global technical and economic  
706 hydropower potential. *Nat. Energy* **2**, 821–828 (2017). [https://doi.org/10.1038/s41560-](https://doi.org/10.1038/s41560-017-0006-y)  
707 [017-0006-y](https://doi.org/10.1038/s41560-017-0006-y)
- 708 69. Ao, Z. *et al.* A national-scale assessment of land subsidence in China's major cities. *Science*  
709 **384**, 301–306 (2024). DOI: 10.1126/science.adl4366
- 710 70. Leslie, H.A., Brandsma, S.H., van Velzen, M.J.M., Vethaak, A.D. Microplastics en route: Field  
711 measurements in the Dutch river delta and Amsterdam canals, wastewater treatment  
712 plants, North Sea sediments and biota. *Environment International*, **101**, 133-142 (2017).  
713 <https://doi.org/10.1016/j.envint.2017.01.018>.
- 714 71. Nicholls, R.J. *et al.* (eds.). *Ecosystem services for well-being in deltas: integrated*  
715 *assessment for policy analysis* (Springer Nature, 2018).
- 716 72. Yuen, K.W. *et al.* Extent of illegal sand mining in the Mekong Delta. *Commun. Earth*  
717 *Environ.* **5**, 31 (2024). <https://doi.org/10.1038/s43247-023-01161-1>
- 718 73. Eslami, S. *et al.* Projections of salt intrusion in a mega-delta under climatic and  
719 anthropogenic stressors. *Commun. Earth Environ.* **2**, 142 (2021).  
720 <https://doi.org/10.1038/s43247-021-00208-5>
- 721 74. Rahman, M.M. *et al.* Salinization in large river deltas: Drivers, impacts and socio-  
722 hydrological feedbacks. *Water Security* **6**, 100024 (2019).  
723 <https://doi.org/10.1016/j.wasec.2019.100024>
- 724 75. Nienhuis, J.H. *et al.* River deltas and sea-level rise. *Annual Review of Earth and Planetary*  
725 *Sciences* **51**, 1, 79-104 (2023). <https://doi.org/10.1146/annurev-earth-031621-093732>
- 726 76. Jung, N.W. *et al.* Economic development drives massive global estuarine loss in the  
727 Anthropocene. *Earth's Future*, **12**, e2023EF003691.  
728 <https://doi.org/10.1029/2023EF003691>
- 729 77. Seijger, C. *et al.* 2017. An analytical framework for strategic delta planning: negotiating  
730 consent for long-term sustainable delta development. *Journal of Environmental Planning*  
731 *and Management* **60**(8), 1485-1509 (2017).  
732 <https://doi.org/10.1080/09640568.2016.1231667>
- 733 78. Zevenbergen, C., Khan, S.A., van Alphen, J., Terwisscha van Scheltinga, C. & Veerbeek, W.  
734 Adaptive delta management: a comparison between the Netherlands and Bangladesh  
735 Delta Program. *International Journal of River Basin Management* **16** (3), 299-305 (2018).  
736 <https://doi.org/10.1080/15715124.2018.1433185>
- 737 79. Cox, J.R. *et al.* A global synthesis of the effectiveness of sedimentation-enhancing  
738 strategies for river deltas and estuaries. *Global and Planetary Change* **214**, 103796 (2022).  
739 <https://doi.org/10.1016/j.gloplacha.2022.103796>
- 740 80. Dunn, F.E. *et al.* Sedimentation-enhancing strategies for sustainable deltas: An integrated  
741 socio-biophysical framework. *One Earth*, **6**(12), 1677–1691 (2023).  
742 <https://doi.org/10.1016/j.oneear.2023.11.009>
- 743 81. Moodie, A.J. & Nittrouer, J.A. Optimized river diversion scenarios promote sustainability  
744 of urbanized deltas. *Proceedings of the National Academy of Sciences*, **118** (27)  
745 e2101649118 (2021). <https://doi.org/10.1073/pnas.2101649118>
- 746 82. Du, H *et al.* Enriching the concept of solution space for climate adaptation by unfolding  
747 legal and governance dimensions. *Environmental Science and Policy* **127**, 253–262 (2022).  
748 <https://doi.org/10.1016/j.envsci.2021.10.021>

- 749 83. Dunn, F.E. & Minderhoud, P.S.J. Sedimentation strategies provide effective but limited  
750 mitigation of relative sea-level rise in the Mekong delta. *Communications Earth and*  
751 *Environment*, **3**(1) (2022). <https://doi.org/10.1038/s43247-021-00331-3>
- 752 84. Kondolf, G. M. *et al.* Save the Mekong Delta from drowning. *Science* **376** (6593), 583–585  
753 (2022). <https://doi.org/10.1126/science.abm5176155>
- 754 85. Zăinescu, F., Anthony, E., Vespremeanu-Stroe, A., Besset, M. & Tatui, F. Concerns about  
755 data linking delta land gain to human action. *Nature* **614**, E20–E25 (2023).  
756 <https://doi.org/10.1038/s41586-022-05624-x>
- 757 86. Minderhoud, P.S.J., Coumou, L., Erkens, G., Middelkoop, H., & Stouthamer, E. Mekong  
758 delta much lower than previously assumed in sea-level rise impact assessments. *Nature*  
759 *Communications* **10** (1), 3847 (2019). <https://doi.org/10.1038/s41467-019-11602-1105>.
- 760 87. Seeger, K. *et al.* Assessing land elevation in the Ayeyarwady Delta (Myanmar) and its  
761 relevance for studying sea level rise and delta flooding. *Hydrol. Earth Syst. Sci.* **27**, 2257–  
762 2281 (2023). <https://doi.org/10.5194/hess-27-2257-2023>
- 763 88. Hooijer, A. & Vernimmen, R. Global LiDAR land elevation data reveal greatest sea-level rise  
764 vulnerability in the tropics. *Nat. Commun.* **12**, 3592 (2021).  
765 <https://doi.org/10.1038/s41467-021-23810-9>
- 766 89. Zhu, Q. *et al.* Hidden delta degradation due to fluvial sediment decline and intensified  
767 marine storms. *Science Advances*, **10** (18) (2024). DOI: 10.1126/sciadv.adk1698
- 768 90. Schipper, E.L.F. *et al.* Climate resilient development pathways. In *Climate Change 2022:*  
769 *Impacts, Adaptation and Vulnerability* (eds. Pörtner H-O *et al.*). Contribution of Working  
770 Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.  
771 Cambridge, UK and New York, NY, USA: Cambridge University Press, pp. 2655–2807 (2022).
- 772 91. Dunn, F.E. *et al.* Projections of declining fluvial sediment delivery to major deltas  
773 worldwide in response to climate change and anthropogenic stress. *Environ. Res. Lett.* **14**,  
774 084034 (2019). DOI 10.1088/1748-9326/ab304e.
- 775 92. Chua, S.D.X. *et al.* Can restoring water and sediment fluxes across a mega-dam cascade  
776 alleviate a sinking river delta? *Sci. Adv.* **10** (18), eadn9731 (2024). DOI:  
777 10.1126/sciadv.adn9731
- 778 93. Wang, H. J. *et al.* Impacts of the dam-orientated water-sediment regulation scheme on the  
779 lower reaches and delta of the Yellow River (Huanghe): a review. *Glob. Planet. Change* **157**,  
780 93–113 (2017). <https://doi.org/10.1016/j.gloplacha.2017.08.005>
- 781 94. Saintilan, N. *et al.* Widespread retreat of coastal habitat is likely at warming levels above  
782 1.5 °C. *Nature* **621**, 112–119 (2023). <https://doi.org/10.1038/s41586-023-06448-z>
- 783 95. Zhang, T. *et al.* Warming-driven erosion and sediment transport in cold regions. *Nature*  
784 *Reviews Earth & Environment* **3** (12), 832–851 (2022). 10.1038/s43017-022-00362-0
- 785 96. Oppenheimer, M. *et al.* Sea Level Rise and Implications for Low-Lying Islands, Coasts and  
786 Communities. In *IPCC Special Report on the Ocean and Cryosphere in a Changing*  
787 *Climate* (eds., Pörtner, H-O *et al.*). Cambridge University Press, Cambridge, UK and New  
788 York, NY, USA, pp. 321–445. <https://doi.org/10.1017/9781009157964.006>.
- 789 97. Timmermans, J. & Storms, J. (eds.) *Panorama New Netherlands*. (2023)  
790 <https://flowsplatform.nl/#/panorama-new-netherlands-1581327249675>
- 791 98. van Alphen, J., Haasnoot, M. & Diermanse, F. Uncertain accelerated sea-level rise,  
792 potential consequences, and adaptive strategies in The Netherlands. *Water* **14**, 1527  
793 (2022). <https://doi.org/10.3390/w14101527>
- 794 99. Tessler, Z.D. *et al.* Profiling risk and sustainability in coastal deltas of the world. *Science*  
795 **349**, 638–643 (2015). doi: 10.1126/science.aab3574

- 796 100. Karra, K. *et al.* Global land use/land cover with Sentinel 2 and deep learning. In *2021 IEEE*  
797 *International Geoscience and Remote Sensing Symposium IGARSS* (pp. 4704-4707). IEEE  
798 (2021).  
799 101. Marconcini, M., Metz-Marconcini, A., Esch, T., & Gorelick, N. Understanding current  
800 trends in global urbanisation-the world settlement footprint suite. *GI\_Forum* **9** (1), 33-38  
801 (2021).  
802

### 803 **Correspondence**

804 Correspondence should be addressed to Edward Anthony (anthony@cerege.fr)  
805

### 806 **Online content**

807 Source data and Python codes for Fig. 2 are available at  
808 <https://github.com/FlorinZai/DeltaHumans>.  
809

### 810 **Acknowledgements**

811 Patrick Pentsch assisted in drawing figures 1,3,5, and Sarah Anthony drew the adaptation  
812 strategies on the delta in Box 1.  
813

### 814 **Author contributions**

815 All the authors contributed to the discussions that formed the basis for this Review, and to  
816 the writing and editing of the manuscript. F.Z. and N.M. collated the data for Figure 2.  
817

### 818 **Competing interests**

819 The authors declare no competing interests.  
820

## 821 **FIGURE CAPTIONS**

822

823 **Figure 1. Simplified sketch of a river delta.** (A) Deltas result from a river feeding sediment into  
824 a standing body of water at a rate that exceeds dispersal processes, leading, especially in large  
825 deltas, to the accumulation of a considerable sediment mass both on land and in the  
826 subaqueous zone<sup>9</sup>. The largest deltas started developing about 8000 years ago (covering much  
827 of the Holocene, i.e., the last 10,000 years of Earth's history). Delta existence has hinged on  
828 abundant sediment supply from river catchments. (B) Idealized delta. Deltas expanded and  
829 built up and out from initial bay-head settings. Their growth was favoured by a relatively stable  
830 global sea level, and most of the world's deltas have a mean elevation below 2 m above  
831 present mean sea level<sup>88</sup> although precise elevation data are lacking. Deltas undergo natural  
832 subsidence (sinking) due to sediment, including organic matter, compacting under its own  
833 weigh. Deltas are ecologically diverse with subtle elevation variations, subject to floods,  
834 channel switches (avulsions) and meandering, marine incursions during storms, and localized  
835 erosion. Deltas have provided space and resources for the development and thriving of human  
836 society alongside these hazards. Humans progressively adapted across the past 7000 years to  
837 deltas, building up a highly imbricated relationship but also generating profound delta  
838 biophysical modifications. Low- to mid-latitude deltas are increasingly subjected to sediment  
839 starvation from river-basin hydropower development involving dams and reservoirs, from

840 aggregate mining, and from aggravated subsidence caused by delta population growth and  
841 resource exploitation, all culminating in vulnerability to global sea-level rise.

842

843

844 **Figure 2. Anthropocene delta demography and land changes.** (A) Population over a total area  
845 of approximately 730,000 km<sup>2</sup> covered by the largest 86 global deltas<sup>9</sup>, with concentric circles  
846 representing 1975, 2020, and projected for 2030. (B) Anthropogenic footprint: combined  
847 fractions of built-up and cropland areas within delta plains, with juxtaposed regional averages.  
848 (C) Breakdown of delta area, population, natural area regionally, and global land cover  
849 emphasizing the disproportionate anthropogenic influences across different regions. (D)  
850 Urban development highlighted by Shanghai (Yangtze delta), the world's largest conurbation  
851 and city with respectively 80 M and 22.3 M inhabitants in 2018. (E) Land-use patterns in the  
852 Mekong delta. (F) Temporal trends illustrating population growth dynamics within deltas over  
853 the decades, underscoring the increasing anthropogenic pressures. Population data (plots  
854 A,C,F) from the GHS-POP R2023A population grid multitemporal (1975-2030) of the European  
855 Commission available at: [http://data.europa.eu/89h/2ff68a52-5b5b-4a22-8f40-](http://data.europa.eu/89h/2ff68a52-5b5b-4a22-8f40-c41da8332cfe)  
856 [c41da8332cfe](http://data.europa.eu/89h/2ff68a52-5b5b-4a22-8f40-c41da8332cfe); land cover data (plots B, E, C) from the ESRI 2020 Land Cover dataset<sup>100</sup> with  
857 the original 10 classes simplified into four classes: Cropland, Settlement, Water and Natural,  
858 and settlement area (plot D) from World Settlement Footprint: 1985-2015 and 2019<sup>101</sup>, and  
859 from the samapriya-awesome-gee community-dataset hosted at  
860 <https://doi.org/10.5281/zenodo.8223455> retrieved via Earth Engine<sup>®</sup>.

861

862

863 **Figure 3. Human intersection with delta geomorphological development over the last 7000**  
864 **years in the wake of stabilization of the postglacial SLR.** (A) Smoothed global mean sea level  
865 over the last 8000 years<sup>20</sup>. The strong association between sea level and deltas has been  
866 reviewed recently<sup>75</sup>. (B) Idealized geomorphic phases of Holocene delta development  
867 comprising inception following sea-level stabilization, expansion over much of the Neolithic  
868 and the Bronze Age, notably through active avulsions, resulting in the broad fan shape of  
869 modern deltas downstream of the apex (Fig. 1), and upbuilding-outbuilding especially in the  
870 Common Era that has, over the last three thousand years, led to burial of anthropogenic  
871 artefacts along nameless (and no doubt numerous) former delta river branches, long  
872 abandoned in classic to modern times. (C) Global population (sources are referenced in<sup>12</sup>)  
873 since 1670 CE (taken as the start of the informal pre-industrial period), showing the significant  
874 Anthropocene spike that also saw the creation of numerous delta cities and megacities. (D)  
875 Timeline showing significant phases and spikes in the 7000 year-long human-delta relationship  
876 from the earliest human occupation, through the Neolithic and formation of the world's first  
877 city states and important expansion of settlements in the Bronze Age, followed by increasing  
878 human engineering and transformation during the Common Era, accompanied by strong  
879 human influence on river catchment sediment supply. This culminated with globalization of  
880 human occupation of deltas during the industrial era that has resulted in many deltas being  
881 locked into anthropogenic transformations that have become irreversible in the  
882 Anthropocene<sup>16</sup>.



883







884



885 **Figure 4. Coordinated river and delta planning and management strategies to reduce**  
886 **vulnerability and maintain delta sustainability.** Thriving deltas in the past have done so in the  
887 framework of a complex balanced geological-climate-ecological-social science relationship.  
888 Coordinated river-basin planning and management, and delta planning and management,  
889 revolving around this blend, should be at the forefront of future delta sustainability, as they  
890 will be determinant, in assuring or not, vulnerability reduction and in maintaining delta  
891 sustainability at low-end near-future SLR scenarios (SSP1-1.9/1-2.6). River basins are  
892 fundamental to the link between climate change, local precipitation and sediment supply to  
893 deltas. Important areas of river-basin management are water and sediment fluxes to minimize  
894 fragmentation and assure connectivity, notably through rethinking of alternative solutions to  
895 hydropower and irrigation dams where feasible, and where dams are inevitable, their optimal  
896 design and operation to minimize sediment trapping by enabling sediment routing through  
897 reservoirs via sluices, sediment-drawdown gates, bypass tunnels, dredging and downstream  
898 relocation of dredged sediment. Other aspects include controlled aggregate mining, and  
899 population mobility from upland basin areas to deltas. Hence the importance of considering  
900 knowledge- and data-backed aspects revolving around what delta 'ownership' implies, and  
901 how governance, management and adaptation are deployed. Anticipation is of equal  
902 importance at a time when most deltas have no known management structure. Differences in  
903 the extent to which these actions are taken, or not, will generate inequalities among deltas  
904 and their vulnerability to global/regional SLR. Both the river-basin and delta spheres face  
905 social-ecological, political and funding challenges that will generate variability among deltas  
906 in the capacity to act. Sustainability will decline for all deltas under high SLR scenarios,  
907 underlining the overarching condition of urgent climate stabilization.  
908

909 **Figure 5. SLR and delta sustainability.** Sustainability is scaled against projected likely ranges  
910 of global SLR under different shared socio-economic pathways (SSP) from<sup>61</sup>, assuming mean  
911 delta-plain elevations below 2 m above present mean sea level<sup>86,88</sup>: (1) progressively  
912 imperiled, notably sediment-starved deltas, with no river basin-delta management or  
913 planning, even under a near-future low-end (SSP1-1.9) scenario; (2) deltas with good  
914 adaptation through basin-delta planning and management and sustainable at low-end (SPP1-  
915 1.9) and moderately-low (SPP1-2.6) scenarios; (3) increasing marine inundation and costlier  
916 and unsustainable adaptation at a moderately-high scenario (SPP2-4.5) likely to affect most  
917 world deltas; (4) large-scale inundation and drowning of world deltas at high-end scenarios  
918 (SSP3-7.0 and SPP 5-8.5) of 1-1.4 m above present mean sea-level. Action perspectives will  
919 strongly diverge between deltas, and while stronger economic means and governance may  
920 provide larger space for solution, adaptation will (rapidly) decline for all deltas under high SLR  
921 scenarios. Projection uncertainties constitute a challenge for investment planning in Protect  
922 and Accommodate strategies (Box 1). In the Netherlands, at the forefront in battling SLR<sup>57</sup>, a  
923 1 m-rise is factored into defences to 2100, following the Delta Commission Plan, and defences  
924 will continue to be raised to withstand another metre by 2200. Residual risks like storm surges  
925 and unforeseeable extremely rapid SLR cause, however, concern and raise questions about  
926 which strategy to adopt. Retreat could be selective, letting, for instance, Friesland become  
927 flooded, but protecting the Rhine delta provinces hosting most people and economic  
928 activities. Note that subsidence is as important as global SLR in any individual delta.

929







**Coordinated  
governance**



**River basin planning and management**


- *Basin water and sediment management (dams/dam storage; aggregate extraction) – controlled fragmentation*
- *Source-to-sink (basin-to-delta) sediment connectivity*
- *Controlled population migration to deltas*


**Delta planning and management**

- *Population, settlement and infrastructure management*
- *Knowledge/data collection, and anticipation (tipping points)*
- *Sediment connectivity and redistribution*
- *Sedimentation-enhancing strategies*
- *Curbed aggregate and fine-grained sediment extraction*
- *Subsidence control*
- *Nature-based solutions*


# ANTHROPOCENE

Large-scale global destabilization of populous and sediment-starved deltas  
Global climate change and global/regional sea-level rise (plus ocean warming and acidification, ocean waves, storms and surge, heat waves)





## Box 1. Adaptation strategies and approaches to SLR in deltas



**Protect.** Levees, dikes, seawalls, and storm-surge barriers offer straightforward, but costly, protection in populous deltas, sometimes with land reclamation (termed 'advance'<sup>97</sup>). Addressing SLR necessitates a strong commitment, and mass construction and ongoing raising of dikes as in the Ganges-Brahmaputra delta. As sea levels rise and land levels sink so the costs to hold the line and the consequences of failure (residual risk) increase, ultimately representing an existential disaster for the delta inhabitants. Leveed deltas buy time, but are probably not tenable in the long run<sup>98</sup>.

**Accommodate.** This strategy integrates adaptive living solutions in wetlands and sedimentation-enhancing approaches. It appears sustainable in the face of near-future SLR, aligns with historical human-delta co-existence and is favoured by some local communities<sup>90</sup>. It poses challenges in densely populated deltas, requiring alterations in planning and lifestyle.

**Retreat.** Managed or as realignment<sup>96</sup> (eventually orchestrated under delta governance) or unmanaged (spontaneous), this approach involves community relocation from high-risk zones to safer terrain, underscores classic climate adaptation, but is fraught with sociocultural and economic considerations<sup>10</sup>, particularly regarding community integrity, heritage loss, and funding. It is an alternative to costlier Protect and Accommodate strategies in urbanized deltas and suited to low-population deltas (Mississippi, Danube).

**'Laisser-faire'.** This 'give-up' approach may be cost-effective but only really workable where population is low. It implies minimal human intervention, and whether adopted in resignation or deliberately, aligns with preserving natural delta processes and ecological integrity. It is currently implemented to varying degrees in the Mississippi, Danube, and Rhone, and is pertinent to Arctic deltas.