

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

The potential impact of the Lactobacillus name change: the results of an expert meeting organised by the Lactic Acid Bacteria Industrial Platform (LABIP).

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Pot, B., Salvetti, E., Mattarelli, P., Felis, G.E. (2019). The potential impact of the Lactobacillus name change: the results of an expert meeting organised by the Lactic Acid Bacteria Industrial Platform (LABIP). *TRENDS IN FOOD SCIENCE & TECHNOLOGY*, 94(December 2019), 105-113 [10.1016/j.tifs.2019.07.006].

Availability:

This version is available at: <https://hdl.handle.net/11585/695381> since: 2019-12-04

Published:

DOI: <http://doi.org/10.1016/j.tifs.2019.07.006>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

The potential impact of the *Lactobacillus* name change: the results of an expert meeting organised by the Lactic Acid Bacteria Industrial Platform (LABIP).

Bruno Pot^{1,2*}, Elisa Salvetti³, Paola Mattarelli⁴ and Giovanna E. Felis³

Affiliations:

1. Yakult Europe BV, Science department, Schutsluisweg 1, 1332 EN Almere, The Netherlands
2. Research Group of Industrial Microbiology, Fermentation Technology and Downstream Processing (IMDO), Department of Applied Biological Sciences, Vrije Universiteit Brussel (VUB), Pleinlaan 2, 1050 Brussels, Belgium
3. University of Verona, Department of Biotechnology, Villa Lebrecht – Via della Pieve 70 – 37029 San Floriano di San Pietro in Cariano, Verona, Italy
4. Dipartimento di Scienze e Tecnologie Agro-Alimentari, Viale Fanin 42, Bologna, Italy

*Correspondence: bpot@yakult.eu ; Yakult Europe Science Department, Schutsluisweg 1, 1332 EN Almere, The Netherlands

Abstract

Background.

Taxonomy and **nomenclature** are important aspects of biological science, as they allow unambiguous communication about all living species. In contrast to higher life forms, microbes have no comparable sexual reproduction, upon which bacterial speciation can be based. Therefore, alternative criteria have to be used, which differ depending on the group of microorganisms. In the past, phenotypic criteria such as fermentation patterns, enzymatic profiles and DNA-DNA hybridisation were cornerstone techniques for speciation. But today, the wider availability of **high-throughput sequence technology** and the relatively small genome size of bacteria have allowed phenotypic testing to be replaced by genome sequencing as the main source of taxonomic information.

Scope and Approach.

Not unexpectedly, the results of phylogenetic analyses based on these new data do not always match results from phenotypic approaches. Based on a recent analysis of the **genome sequences** of 222 species of the genus

Lactobacillus and related taxa, it is expected that the genus *Lactobacillus* will be split in a considerable number of new genera.

Key Findings and Conclusions.

In October 2018 LABIP organised an expert workshop to discuss the economic, scientific and **regulatory** consequences of this taxonomic change. This report represents a summary of the considerations and outcomes of this workshop, supplemented with some later reflections and recent literature.

Keywords: *Lactobacillus*, Taxonomy, Nomenclature, Genome Sequencing, Food Regulation, QPS-FDA

Background

Members of the genus *Lactobacillus* are widely used in industrial fermentations. Fermentation has the potential to improve the taste, nutritional value and texture of foods and adds to food safety and shelf life by acidification and through the production of antibacterial compounds. Some species of the genus *Lactobacillus* have a long history of safe and legal use in foods. Other species within the genus do not benefit from this status and consequently proper nomenclature and taxonomy is a prerequisite for safety assessment.

Recently Salvetti et al. (2018) investigated the relatedness of 269 species belonging primarily to the Families Lactobacillaceae (genera *Lactobacillus* and *Pediococcus*) and Leuconostocaceae (*Convivina*, *Fructobacillus*, *Leuconostoc*, *Oenococcus* and *Weissella*) as well as reference strains of the genera *Atopobium*, *Carnobacterium*, *Kandleria* and *Olsenella*. Whole genome sequence analysis showed that many of these genera are phylogenetically interwoven and that the genus *Lactobacillus* in particular is very heterogeneous. Based on several parameters, including Average Amino acid Identity (AAI) and Percentage Of Conserved Proteins (POCP), the genus has a spread that largely exceeds the normal spread of a genus. These findings confirmed the results obtained by Sun et al. (2015), who showed that the Average Nucleotide Identity (ANI, pairwise comparison of homologous sequences) for species of the genus *Lactobacillus* was as low as values typically seen at the taxonomic level of an order. The authors therefore concluded that a formal split of the genus was unavoidable and 10 potential groups of microorganisms were proposed that could become the basis for a reclassification of the genus. In another study covering almost all the available *Lactobacillus* genome sequences, Parks et al. (2018) discriminated 16 groups within the genus. Based on 16S rRNA sequence similarity only, Salvetti et al. (2012) and Pot et al. (2014), earlier described the presence of at least 14 stable phylogenetic groups within the genus.

This aberrant taxonomic situation was discussed by the Taxonomic Subcommittee on *Bifidobacterium*, *Lactobacillus* and Related Genera at a meeting in Berlin, September 3rd, 2018. The conclusion of this discussion was that the Subcommittee would support a formal split of the genus and a working group of subcommittee members and external experts was created to collect all available genotypic and phenotypic information that would allow definition of a stable and reliable structure for the genus *Lactobacillus* and its closely related taxa. While this is an ongoing academic process, it was clear that this renaming would have considerable economic and legal consequences, which may heavily impact the industry that is using these organisms in the production of fermented foods, food supplements and feed additives. The Lactic Acid Bacteria Industrial Platform (LABIP) therefore decided to organise an expert workshop (32 participants) to (i) explain the process of reclassification, (ii) make an inventory of the possible consequences and (iii) discuss possible solutions. LABIP was established in 1995 with the purpose of securing a link between academic research and industry. Today the association accommodates 23 companies with an interest in the industrial use of lactic acid bacteria (LAB) in foods or food supplements. The topics of earlier workshops included amongst others, Food Safety (Adams and Marteau, 1995), Probiotics (Guarner and Schaafsma, 1998) and the Nagoya protocol (Johansen, 2017). As full partner in the SSA GutImpact project, LABIP has organized 3 technical expert workshops and 3 consumer related workshops, which included consumer representatives (Lahteenmaki and Ledeboer, 2006; Ledeboer and Lahteenmaki, 2007 ; Ledeboer et al., 2007).

Introduction to prokaryotic nomenclature

As mentioned above, a stable set of names (nomenclature) is needed to allow clear communication on microorganisms . While taxonomy is a matter of experience and agreement among scientists, nomenclature of prokaryotes, earlier called ‘the handmaid of taxonomy’ (Sneath, 2015), is regulated by the Nomenclatural Code (Lapage, 1992) revised after being presented and discussed at the Fourteenth International Congress of Bacteriology and Applied Microbiology (BAM), Montréal 2014 (Parker et al. 2016). The most recent version is available from Oren (2019).

The first ‘Bacteriological Code’, was developed in 1930, during the First International Congress on Taxonomy in Paris. An international Committee crafted a number of unanimously approved resolutions. The work was continued by the Nomenclature Committee on behalf of the International Society for Microbiology International

Committee. After several intermediate documents had been proposed and discussed The first formal English text was officially published in March 1948 in the Journal of Bacteriology (Buchanan et al., 1948).

This code has further evolved and is now under the auspices of the 'International Committee on the Systematics of Prokaryotes' (ICSP), which includes the Archaea in their scope. The need to improve the quality of bacterial nomenclature, with fewer duplicate names and including the description of type species and type strains, resulted in publication on January 1st, 1980 of the 'Approved List of Bacterial Names', a list of bacterial names with standing in microbiology, published in the 'International Journal of Systematic Bacteriology' (IJSB). The list, consisting of 124 names of taxa above the rank of genus, encompasses 2212 genera, species and subspecies names. Till today, new names are added to this list only after valid publication in IJSB, known since 2000 as the 'International Journal of Systematic and Evolutionary Microbiology' ([IJSEM](#); consulted last on March 29th, 2019). Valid publication means that the species description is published as a formal article in IJSEM or appeared on the so-called 'Validation Lists' in IJSEM. In order to be on this list, authors must ask ICSP for formal validation of the taxa described in other peer reviewed scientific journals.

ICSP today supervises several subcommittees, including the 'Subcommittee on the taxonomy of *Bifidobacterium*, *Lactobacillus* and related organisms'. The Subcommittees are responsible for the maintenance of the Nomenclatural Code (Parker et al., 2016) in their respective range of taxa. The 'International Code of Nomenclature of Prokaryotes' (ICNP) defines the rules for the naming of bacterial taxa according to their taxonomic ranking. While covering many possible situations and exceptions, its principles are relatively simple and comprise a limited number of rules (Tindall et al. , 2006). While it is not the intention of this report to review these rules, some aspects are important in order to better understand the reasons for name changes in the genus *Lactobacillus* (Table 1) and are presented in the supplementary material of this article

It is important to note that, in agreement with general principles in prokaryotic taxonomy, the goal is a monophyletic taxonomy, meaning that a phylogenetic analysis should demonstrate that all descendants of any given taxon (in this case a genus) are grouped together and are not mixed with members of other taxa. Also, in favour of clarity, taxa should be kept reasonably compact, comparable to analogous genera, and distinctive in terms of phenotypic and genotypic criteria. In the case of the genus *Lactobacillus*, the genetic distinctiveness has clearly been shown, but the differential phenotypic description of the different genotypic groups might be an issue, as species have been added to the genus based on a number of common phenotypic characteristics. One of the

most important tasks of the above-mentioned working group therefore is to try and find discriminative sets of parameters that might, besides genome sequence parameters, allow reliable description of the new genera. The use of polyphasic datasets, including multiple types of genome sequence analyses, should guarantee a stable and clear separation of the genus. The working group has to collect all possible information on all species of the genus *Lactobacillus* and by exploring and comparing different bioinformatic approaches, derive the best possible reclassification for the genus. For some already recognized larger subgroups, as mentioned above, different phylogenetic approaches, including genome sequence-based and 16S rRNA based trees, have already confirmed their taxonomic status, but for other, smaller clusters or individual species, the new phylogenetic position is still uncertain and needs further dedicated work.

Why the genus has to be split

For the 237 *Lactobacillus* species (208, excluding synonyms and subspecies) the Total Nucleotide Identity (TNI) was found to be between the values expected between order and family and the Average Nucleotide Identity (ANI) values between Order and Class (Sun et al, 2015, Salvetti et al, 2018). Results also revealed that representatives of other genera, such as *Pediococcus*, and members of the family Leuconostocaceae (*Convivina*, *Fructobacillus*, *Leuconostoc*, *Oenococcus* and *Weissella*) are intermixed with species of the genus *Lactobacillus*, which of course is an infringement of the monophyletic requirement. The genus continues to grow, with new isolates being added at an ever increasing speed (Figure 1a and 1b). This is unacceptable from a scientific point of view and the *status quo* was not considered an option.

The scientific rationale for the split

In the past, lactobacilli were grouped mainly based on phenotypic characteristics such as growth temperature or the fermentation of hexoses (homo/heterofermentative potential). The introduction of 16S rRNA gene sequence analysis allowed the development of a more exhaustive taxonomy for the genus, but unfortunately revealed little correlation between the traditional, phenotype-based classification and the phylogenetic relatedness. In addition, the massive description of novel *Lactobacillus* species led to the recognition of a growing number of variable phylogenetic subgroups (Vandamme et al., 1996; Felis and Dellaglio, 2007; Salvetti et al., 2012; Pot et al., 2014).

The decision to use whole genome sequences to further study the issue was based on earlier work performed on other genera (Coenye et al., 2005; Gevers et al., 2005; Konstantinidis and Tiedje, 2005a,b; Rosselló-Mora, 2005;

Wu et al., 2009; Sangal et al., 2016; Chun et al., 2018; Parks et al., 2018) with the use of a variety of new tools (Segata et al., 2013; Qin et al., 2014; Garity, 2016; Hugenholtz et al., 2016; Yoon et al., 2017).

The first comparative genomics on LAB was performed by Makarova and colleagues in 2006, which included 12 LAB genomes (encompassing genera *Lactobacillus*, *Oenococcus*, *Leuconostoc*, *Pediococcus*, *Streptococcus* and *Lactococcus*). The phylogenomic tree based on 54 ribosomal proteins showed that *Lactobacillus* species were intermixed with leuconostocs and pediococci. This topology was later observed also by Zhang and co-workers (2011), who performed a phylogenetic analysis based on 232 genes derived from 28 LAB genomes, and Salvetti and colleagues (2013), who analysed 27 LAB genomes using different datasets (mainly ribosomal proteins and carbohydrate metabolism genes). Two consistent groups were obtained: the Enterococcaceae and Streptococcaceae families on the one hand, and the families Lactobacillaceae and Leuconostocaceae on the other hand. The latter contained two clades, one encompassed the *Lactobacillus acidophilus* complex (including *Lactobacillus delbrueckii*), *Lactobacillus sakei* and *Lactobacillus casei*. The second clade contained the *Lactobacillus salivarius* subgroup, *Lactobacillus plantarum*, *Lactobacillus reuteri*, *Lactobacillus brevis*, and the genera *Pediococcus*, *Oenococcus* and *Leuconostoc*, indicating the polyphyletic nature of the genus.

Zheng et al. (2015) published a phylogenetic perspective based on the concatenated protein sequences of 172 single-copy core genes of 174 *Lactobacillus* and *Pediococcus* type strains, focusing on ecology and biochemistry properties. A clear separation of homofermenters and heterofermenters within these genera was shown and the presence of 24 subgroups (including *Pediococcus*) was revealed.

Sun et al. (2015) sequenced 213 genomes from 174 type strains belonging to 9 genera and also included available genomes from *Oenococcus* and *Leuconostoc* in their analysis (11 genera and 185 species). Results revealed not only that the *Lactobacillus* ancestor was facultatively heterofermentative but also showed that the pairwise ANI values for the genus *Lactobacillus* differed substantially from the values traditionally found for a genus or even for a family, overlapping with values generally seen at the level of an order or even class. TNI values confirmed that the genomic diversity of the genus *Lactobacillus* is at the order and family level.

More recently, Salvetti and colleagues (2018) investigated the relatedness of 269 species belonging primarily to the Families Lactobacillaceae and Leuconostocaceae through phylogenetic analyses (by the use of 16S rRNA genes, 29 ribosomal proteins and 12 housekeeping genes) and the assessment of the AAI and the POCP values. Both

distance-based and sequence-based metrics showed that the *Lactobacillus* genus was paraphyletic and revealed the presence of 10 methodologically consistent subclades, whose suitability to be nuclei of novel genera was substantiated through the detection of putative clade-specific genes and other conventional taxonomic data. Their phylogenomic tree showed the association of obligately heterofermentative lactobacilli with members of the Leuconostocaceae, well-separated from the homofermentative and facultatively heterofermentative *Lactobacillus* species. This separation is consistent with the phylogenetic trees calculated based on ribosomal proteins, housekeeping and core genes and is congruent with carbohydrate fermentation profiles.

In the same period, Parks and colleagues (2018) proposed a standardized bacterial taxonomy based on a phylogenetic analysis of 120 proteins, performed from 94,759 prokaryotic genomes, with the purpose to normalize ranks based on relative evolutionary divergence. This taxonomic analysis, publicly available at the Genome Taxonomy Database website (<http://gtdb.ecogenomic.org>), suggested 16 separate lineages within the genus *Lactobacillus*, based on 205 genomes and also confirmed that leuconostocs and pediococci intermingle with lactobacilli, indicating that the family Leuconostocaceae might need to be absorbed by the family Lactobacillaceae.

More recently, Wittouck and co-workers presented their analysis of the *Lactobacillus* Genus Complex (LGC) based on 2,459 publicly available genomes (96.1% of all genomes). They selected 'decent-quality' genomes (minimum completeness of 90% and maximum redundancy of 10%) and used a 94% Core Nucleotide Identity (CNI) threshold to show 239 discontinuous and exclusive clusters, representing *de novo* species. The analysis thereof allowed the merging of ten sets of published species and one species that can be split off. At least eight clusters could become new species. The data set also allowed classification of 98 previously unclassified genomes and re-identified 74 wrongly classified genomes, albeit the latter was done based on 16S rDNA sequences.

This whole historical discussion leads us to the following question: How many species and genera can we actually find in the genus *Lactobacillus*?

The current dilemma: how many genera?

Distance based metrics, as used by Salvetti et al. (2018) are useful because they can be applied in a pragmatic and naïve manner. The TNI and ANI operational values available in the literature that should be used as cut-offs, however, are controversial and cannot be applied to all the species of the current genus *Lactobacillus*. At the amino acid level the result is somewhat better but still not adequate. CNI values have been used most recently and

may need some further evaluation, especially when used to delineate genera rather than species. For the value of the POCP as a measure to define genera, the discussion is ongoing.

The bottom line is that depending on the methodology and cut-offs used, the resultant number of 'stable' groups in the genus *Lactobacillus* varies. The best path forward is to evaluate all these parameters, compare them and propose solutions which, based on multiple trees, represent the common clusters or groups. Other considerations, however, may need to be taken into account. In any case the methodological rationale will be mainly based on whole genome sequences in the first place, as most species of the genus show high phenotypic similarity, the reason for them to have been included in the genus. In the evaluation of genome similarity the most common criteria, such as TNI and ANI or AAI of core genome genes will be used, in accordance with the generally used standards for genus delineation.

One major concern of the LABIP workshop participants was related to the stability of the revised taxonomy. It would be unacceptable if in the next 25 years or so further taxonomic changes would be proposed for this genus. Therefore, it is important that the new scheme, in line with the rules above, is stable and able to cope with future expected descriptions of new species in this genus and family. Figures 1a and 1b show the increase of the number of species in the 'traditional' genus *Lactobacillus* and this number is expected to rise as new DNA-based research is used to investigate complex microbiota in food, feed, environment, man or animal (Wittouck et al., 2019).

There was a consensus that the most stable structure, with the least chance of future subdivision, might be a structure with a higher number of genera, allowing to more flexibly accommodate future species. Besides being more stable, it would also allow to maybe describe subgroups of lactobacilli focusing on specific characteristics common to the members of these smaller taxa. These characteristics could e.g. be applications in food or feed, geography, physiological properties, even health effects, or may refer to characteristics which are not linked to humans or animals. This stable, detailed structure resembles the benefits of the taxonomic situation in the family Enterobacteriaceae, in which a detailed taxonomy and nomenclature was created. This was the historical result of the clinical importance of these organisms and has the advantage of being able to communicate efficiently about pathogenic performance, expected health consequences, clinical outcomes or treatment options. It is generally accepted that a genus in the family Enterobacteriaceae has the span of a species in most other Families (Public

Health England, 2015). While this is considered a ‘medical’ advantage, the detailed taxonomy makes identification sometimes more difficult and more specialized (Adeolu et al., 2016). Clearly, at the level of the genus *Lactobacillus*, we currently do not have this type of detailed phenotypic support, but a finer subdivision of the genus *Lactobacillus*, based on genome sequences may represent important opportunities to improve for example, communication, assist in improved regulation and support scientific investigation of molecular mechanisms of probiotic action.

In order to assist in this decision, the expert working group may consider ecological or health related factors, as well as metabolic or mechanistic properties to define and describe the new genera. If these data are not available, they may possibly be derived from genome sequences (Wittouck et al., 2019). In this context, the use of molecular characteristics such as Conserved Signature Insertions/deletions (CSIs) found in widely distributed proteins present in related groups of organisms (Gupta, 2014, 2016; Gupta et al., 2017; Naushad et al., 2014), may provide reliable evidence of phylogenetic relationship, independent of traditional phylogenetic trees. Species with identical CSIs are related to each other since they share a common ancestor (synapomorphic characteristic) and could therefore contribute to a more reliable and more practical delineation and description of the new genera.

The naming of the future genera

In the light of the description of new genera, LABIP workshop participants suggested keeping the initial “L” for the names of the new genera in order to minimize confusion. In this way, commercially important species such as *Lactobacillus casei*, *Lactobacillus plantarum* and *Lactobacillus reuteri*, which will be no longer lactobacilli and will be included in novel genera, will still be abbreviated as *L. casei*, *L. plantarum* and *L. reuteri*. There is already such an example in the literature regarding the former human pathogen ‘*Clostridium*’ *difficile*, recently reclassified as *Clostridioides difficile* maintaining the common abbreviation *C. difficile*, preventing confusion (Lawson et al., 2016). Proposals to maintain the subpart ‘Lact-’ as a start of the new genus names are therefore currently being considered and proposals have been made to the International Committee on Systematics of Prokaryotes (ICSP).

Genera and species: the future for taxonomic descriptions

While official ‘minimal standards’ are available for the description of new taxa of LAB (Mattarelli et al., 2014), these indications are focused on the species level and cannot be considered really ‘minimal’, making the description quite cumbersome. Therefore, it would be useful to suggest a revision (simplification) of these

standards. At the same time, the Taxonomic Subcommittee should think about the requirement for culturing of the bacteria, given the fact that with increased use of sequencing technology, the discovery of new species that cannot currently be cultured, will only increase. Maybe Mol% G+C content, ANI, TNI, CNI, CSI or POCP values, digital (or wet lab measured) DNA:DNA homology values or combinations thereof, using whole genome sequence based phylogenomics should nowadays be used, while the importance of cell wall structure, chemotaxonomic or other phenotypic markers may need to be reduced in future requirements for the description of new taxa.

Examples were discussed during the workshop for genera other than *Lactobacillus*. Next-generation sequencing (NGS) combined with new bioinformatics approaches for phylogeny, based on whole genome sequences, have been used successfully for strain discrimination and for phylogenetic characterization of novel species, even of species in complex communities such as the gut (Ramasamy et al., 2014; Vernikos et al., 2015). Extensive attempts have recently been made to survey bifidobacterial populations in mammals, resulting in the identification of several putative novel bifidobacterial species. Ventura and co-workers described a methodology based on whole-genome comparisons, similar to that used for the reclassification of members of the Gram-positive genus *Bacillus* (Dunlap 2015), with the ambition to unambiguously redefine the taxonomy of members of the genus *Bifidobacterium* (Milani et al. 2014; Lugli et al., 2018 a,b).

The recent introduction of a Digital Protologue Database (DPD) by the journals 'Systematic and Applied Microbiology' and 'Antonie van Leeuwenhoek' (Rossello-Mora et al., 2017) may also be a next step in the modernisation of taxonomic publications. In the DPD, the protologues given in the taxonomic manuscripts (i.e. the formal descriptions) would be extracted from a Digital Protologue (DP) compulsorily filled by the authors. Their accumulated contributions are creating a public and interactive database (<http://imedea.uib-csic.es/dprotologue/>, now with > 700 species), which generates a unique Taxonumber. The initiative, already supported by a number of journals, also encompasses the descriptions of taxa in papers as a protologue table rather than difficult to read text, traditionally presented at the end of the manuscript. Practical details can be found on e.g. the [SAM website](#) (last consulted March 19th, 2019). The link also contains 13 other standard requirements for the submission of taxonomic manuscripts. The use of this procedure is thought to largely limit future duplicate or wrong descriptions as seen in the past (Tables 1 and 2; Wittouck, 2019). Unfortunately, however, this useful initiative has not been adopted by the IJSEM. This is unfortunate, as this journal is instrumental in valid taxonomic descriptions. In the frame of the current renaming of the genus *Lactobacillus*, the availability of a formal public website, listing all

279 official names and their taxonomic history, might avoid confusion for governmental, legal, medical and scientific
280 stakeholders.

281 **The legal and other consequences**

282 Commercially important *Lactobacillus* species, including species and strains with well-known fermentation capacity
283 or proven probiotic activity, will be found in most of the newly defined genera. The LABIP workshop participants
284 were concerned about the scale of the renaming. While frequent name changes have occurred in the past for
285 species of the genus *Lactobacillus* (Tables 1 and 2), it is likely that close to 200 species will change genus name in a
286 single operation. This is an unprecedented scale of change for the LAB, with the potential to confuse the various
287 stakeholders.

288 The renaming of the majority of the lactobacilli will impact the consumer, the scientific and medical or veterinary
289 communities, regulators, governments, lawyers and companies. Consumers may wonder why an ingredient list no
290 longer contains the name of a familiar microbe and therefore have doubts about the content of the product (feed,
291 food or food supplement). Scientific literature surveys will need to account for historical and new genus names.
292 Residual use of the historical names is likely to occur in books. Lawyers or scientists involved in the validation of
293 patent applications or trademarks may be confronted with the same issue for many years to come. Patent officers
294 will not find prior art for the new bacterial genera and the risk exists that in this way existing patents may be
295 undermined. This risk is real, as according to Espacenet there are more than 10.000 patents with the term
296 '*Lactobacillus*' in the title or the abstract. In the medical and veterinary field, specific taxonomic names are familiar,
297 referring to either innocence or danger. New names will surely require further information to be collected.
298 Regulators and governments will need to compare laboratory reports with biosafety and biosecurity lists (Laulund
299 et al., 2017), important for considerations on quarantine or certification, to decide on Nagoya protocol
300 requirements, or in the food and feed arena, to verify with QPS-, GRAS-, GMO-, Additive-, IDF inventory- or
301 national positive lists, the latter existing in Canada, China, Brazil, Malaysia, and Thailand. In the EU Register of Feed
302 Additives (Anonymous) more than 90% of the listed products with lactobacilli will be concerned by the name
303 change.

304 In transportation, International Airline Transportation Authorities and truck companies may face issues during
305 border controls, as import/export of specific products may lead to unexpected delays, because responsible officers

may not be aware of the official name changes of organisms that are normally 'approved for import/export'. For companies, taxonomic names are not only important for their research and manufacturing, but also for master regulatory dossiers, product authorizations, marketing materials, ingredient lists, safety dossiers, product websites, patent monitoring, export certificates, and other documentation.

While all these issues seem quite alarming, the general impression, based on experiences in the past, was that the impact of the name change could be manageable with time and dedicated resources but will have a significant cost for companies. As an example, the legal consequences of the name changes were explored in a presentation of a member of the European Scientific Panel on BIOHAZ and member of the EFSA Working Group on QPS (28 of 36 QPS species of the genus *Lactobacillus* might be renamed, 12 species have strains with GRAS notifications). It was clear that the normal approval procedure, as explained in Herman et al. (2019), can be used for the approval of the new names, even if the taxonomic changes concern name changes at the genus level (and not on the species level as usually dealt with).

In the USA, the FDA discriminates common or market names from scientific names. While they generally regard common names as appropriate market names, provided they are not misleading or confusing (Principle 2), they generally use the valid scientific names in accordance with the scientific literature. The new official genus names will be introduced through submission of specific dossiers to the FDA, and since companies may make individual decisions regarding the use of the new names, an abrupt disruption in the use of the old names is very unlikely.

More specific changes in labelling of probiotics may be needed as labelling with current nomenclature for genus and species is stipulated in several published guidelines (FAO /WHO guidelines, 2002; CRN and IPA guidelines, 2017).

What can be done?

A call was launched for a comprehensive and easy-to-use document or website where the official name changes are listed and updated when necessary. The site should explain in a comprehensible way why the name was changed and give a clear representation of the 'BEFORE' versus 'NEW' names, noting the date that the change was officially introduced. This document or site should be available before the publication of the new names, so that it can be activated simultaneously with the official publication of the new names in the IJSEM. The provider of the system should have a high credibility, preferably an academic or governmental institution. While similar websites

already exist (for example, <https://www.itis.gov/>; <http://www.bacterio.net/>; www.straininfo.net/), some require substantial background knowledge to be used efficiently and may also not provide the right explanation and context for the name changes in, for example, the food, veterinary or medical sector.

It was furthermore suggested that similar information should be made available at websites for and from governmental organisations including FDA, EFSA, CDC, ECDC, WHO, FAO, EMA, [public health agencies](#) (last consulted March 19th, 2019) or research organisations such as NIH - NCBI, NHS. The issue should also be taken up by ongoing education and training programs of public health agencies, as suggested by the Institute of Medicine (2003) (or homologues in other parts of the world). Also, a smartphone application was suggested as a way to collect name changes in a timely fashion. Companies will need to plan and implement name changes on their commercial products, their websites, packaging materials and ingredient lists as well as their associated marketing material, product composition and import or export certificates. They will need to communicate transparently with all their stakeholders and end-users and be ready to answer their questions in a comprehensive way. The possibility, however, to be able to refer to an independent, preferably academic source to support their communication is considered key. Finally, it should be stressed that while the genus and species name of a specific strain can change, its official strain designation (strain number as known in one or more reference culture collections) will remain the same, as the renaming procedure will only change the genus name '*Lactobacillus*' and not the species epithet. '*Lactobacillus*' may be replaced by the genus name '*Examplebacillus*', but the species epithet and the strain indication will not change: *Lactobacillus casei* strain XY123 in this example will be renamed to *Examplebacillus casei* strain XY123. Using the strain designation or strain number could therefore be an important way to also limit confusion.

What to expect in the future?

The recent finding of 25 new bifidobacterial species in non-human primates is a clear sign that the discovery of the bacterial diversity and therefore the extension and finetuning of the bacterial taxonomy and nomenclature is only at its beginning. It is to be expected that new species, new genera and even new families will be described in fields that are relevant for the food business or crucial in human and animal health or disease considerations. It is important that these new taxa are dealt with efficiently and unambiguously. Therefore, careful attention is necessary regarding the optimal criteria to use for the description of these new taxa, taking into considerations new sources of phylogenetic information and revisiting existing nomenclatural and taxonomic structures as

needed. Taxonomy is a dynamic process, able to accommodate new scientific findings and insights. The anticipated renaming in the genus *Lactobacillus*, and likely also in *Bifidobacterium*, can be considered a logical scientific reflection of these new insights. The economic absorption and the integration in society of the results, however, are known to be slow processes. In July 2004, as an example, the two species *Bifidobacterium animalis* and *Bifidobacterium lactis* were merged and became officially *Bifidobacterium animalis* subsp. *animalis* and *Bifidobacterium animalis* subsp. *lactis* respectively (Masco et al., 2004). At present, 15 years after date, there are still many “*B. lactis*” products on the market. This could be due to the commercial interest of the name *B. lactis* as well as because producers are not yet aware of the name change. It can therefore be expected that during a considerable transition period the current *Lactobacillus* nomenclature will be used in parallel to the new one. Another example, on the genus level, is the renaming of ‘*Streptococcus lactis*’ to *Lactococcus lactis* in 1985 by Schleifer et al. (1985). While more than 30 years ago, there are still products in the market with the name *Streptococcus lactis*. While this is probably unavoidable, it is definitely not desirable, as it will lead to confusion. Therefore, the above mentioned measures are important and their availability needs to be spread widely.

Conclusion

Although lactobacilli are ubiquitous and appear in diverse niches, they have common phenotypic properties that were the basis of their taxonomic grouping. The use of modern sequencing technology recently revealed their extraordinary diversity, putting a shadow on their current classification. Lactobacilli have significant scientific and economic value, but, as mentioned by a speaker, the history of the genus *Lactobacillus* has been ‘long, painful and torturous’, led by many different people, in different times, with different methodologies. Evolution is a keyword in this, both referring to the microbes themselves as well as to the methodologies used. Recent technological evolutions have enabled us to gather the complete genome sequence of any bacterium we find. In the past already the importance of the genotypic information for taxonomic work has been recognised, but the methods to study DNA were limited. Nowadays this hurdle has been largely overcome and consequences of this are major in terms of expected nomenclatural changes. The taxonomy and nomenclature of important genera such as *Bacillus* and *Clostridium* have already been revised. Next in line is *Lactobacillus*, most probably followed by the genus *Bifidobacterium*. While nomenclatural changes have been common in the past, the anticipated scale in a group of bacteria that have such an important economic impact raised some concerns. While these concerns may often be from a practical and financial point of view, there is a clear request for stability of the

new classification scheme, urging the scientific community to make the new nomenclature suitably robust to withstand future methodology changes in the field for at least the next 25 years. This is especially important to accommodate the slow pace of change evident in the regulatory compared to the scientific arenas.

The new entities should be defined by an as wide as possible bioinformatic approach, using all information available within the complete genome sequences, and possibly tracing uniform patterns of the presence/absence of specific (sets of) genes, linked to biological features relevant for their grouping and description. On the positive side, reclassification of the genus *Lactobacillus* into more uniform and stable taxa will (i) create opportunities for more accurate communication, and (ii) assist in defining more accurate molecular markers that will facilitate regulatory assessment of applications (Salveti and O'Toole, 2017). The upcoming renaming of the genus *Lactobacillus* should therefore not be regarded solely as a possible cause of confusion for stakeholders, but also as an opportunity to start a new nomenclatural situation, adapted to the new scientific reality and ready to cope with future evolutions in the field of microbiology. The translation of this scientific reality to legal, economic and societal structures may take some years but a lot can be done to smoothen that transition.

References

1. Adams M. R. & Marteau P. (1995). On the safety of lactic acid bacteria from food. *International Journal of Food Microbiology* 27, 263-264. [https://doi.org/10.1016/0168-1605\(95\)00067-T](https://doi.org/10.1016/0168-1605(95)00067-T).
2. Adeolu, M., Alnajjar S., Naushad S. & Gupta R. S. (2016). Genome-based phylogeny and taxonomy of the 'Enterobacteriales': proposal for Enterobacterales ord. nov. divided into the families Enterobacteriaceae, Erwiniaceae fam. nov., Pectobacteriaceae fam. nov., Yersiniaceae fam. nov., Hafniaceae fam. nov., Morganellaceae fam. nov., and Budviciaceae fam. nov. *International Journal of Systematic and Evolutionary Microbiology*, 66, 5575–5599. <https://doi.10.1099/ijsem.0.001485>.
3. Anonymous. European Union, Register of Feed Additives, pursuant to Regulation (EC) No 1831/2003. Annex I: List of additives. https://ec.europa.eu/food/sites/food/files/safety/docs/animal-feed-eu-reg-comm_register_feed_additives_1831-03.pdf (consulted last on March 21, 2019)
4. Back, W. (1978). Elevation of *Pediococcus cerevisiae* subsp. *dextrinicus* Coster and White to species status [*Pediococcus dextrinicus* (Coster and White) comb. nov.]. *International Journal of Systematic Bacteriology*, 28, 523–527.
5. Buchanan R. E., St. John-Brooks R. & Breed R. S. (1948). International bacteriological code of nomenclature. *Journal of Bacteriology*, 55, 287–306. (also reprinted in 1949, *Journal of General Microbiology*, 3, 444–462).
6. Code: <http://www.bacterio.net/-foreword.html#Code>
7. Chun, J., Oren, A., Ventosa, A., Christensen H., Araha, I. D. R., da Costa, M.S, Rooney, A. P., Yi, H., Xu, X. W., De Meyer, S., & Trujillo, M. E. (2018). Proposed minimal standards for the use of genome data for the taxonomy

- of prokaryotes. *International Journal of Systematic and Evolutionary Microbiology*, 68(1), 461-466.
[https://doi: 10.1099/ijsem.0.002516](https://doi.org/10.1099/ijsem.0.002516).
8. Coenye, T., Gevers, D., Van de Peer, Y., Vandamme, P. & Swings, J. (2005). Towards a prokaryotic genomic taxonomy. *FEMS Microbiology Reviews*, 29(2), 147-167. [https:// DOI:10.1016/j.femsre.2004.11.004](https://doi.org/10.1016/j.femsre.2004.11.004)
9. Collins, M. D., Farrow, J. A. E., Phillips, B. A., Feresu, S. & Jones, D. (1987). Classification of *Lactobacillus divergens*, *Lactobacillus piscicola*, and some catalase-negative, asporogenous, rod-shaped bacteria from poultry in a new genus, *Carnobacterium*. *International Journal of Systematic Bacteriology*, 37, 310-316.
10. Collins, M. D., Phillips, B. A. & Zanoni, P. (1989). Deoxyribonucleic acid homology studies of *Lactobacillus casei*, *Lactobacillus paracasei* sp. nov., subsp. *paracasei* and subsp. *tolerans*, and *Lactobacillus rhamnosus* sp. nov., comb. nov. *International Journal of Systematic Bacteriology*, 39, 105-108.
11. Collins, M. D. & Wallbanks, S. (1992). Comparative sequence analyses of the 16S rRNA genes of *Lactobacillus minutus*, *Lactobacillus rimae* and *Streptococcus parvulus*: proposal for the creation of a new genus *Atopobium*. *FEMS Microbiology Letters*, 95, 235-240. [https://DOI: 10.1111/j.1574-6968.1992.tb05372.x](https://doi.org/10.1111/j.1574-6968.1992.tb05372.x)
12. Collins, M. D., Samelis, J., Metaxopoulos, J., & Wallbanks, S. (1993). Taxonomic studies on some leuconostoc-like organisms from fermented sausages: description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *Journal of Applied Bacteriology*, 75, 595-603.
13. CRN and IPA guidelines. (2017). Best Practices Guidelines for Probiotics. Document from the Council for Responsible Nutrition (CRN) and the International Probiotics Association; IPA.
<https://www.crnusa.org/sites/default/files/pdfs/CRN-IPA-Best-Practices-Guidelines-for-Probiotics.pdf>
(consulted last on March 21, 2019).
14. Dewhirst, F.E., Paster, B. J., Tzellas, N., Coleman, B., Downes, J., Spratt, D. A., Wade, W. G. (2001). Characterization of novel human oral isolates and cloned 16S rDNA sequences that fall in the family Coriobacteriaceae: description of *Olsenella* gen. nov., reclassification of *Lactobacillus uli* as *Olsenella uli* comb. nov. and description of *Olsenella profusa* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 51, 1797-1804. [https://DOI: 10.1099/00207713-51-5-1797](https://doi.org/10.1099/00207713-51-5-1797)
15. Dellaglio, F., Torriani, S. & Felis, G. E. (2004). Reclassification of *Lactobacillus cellobiosus* Rogosa et al. 1953 as a later synonym of *Lactobacillus fermentum* Beijerinck 1901. *International Journal of Systematic and Evolutionary Microbiology*, 54, 809-812. [https://doi:10.1099/ijse.0.02947-0](https://doi.org/10.1099/ijse.0.02947-0).
16. Dellaglio, F., Vancanneyt, M., Endo, A., Vandamme, P., Felis, G. E., Castioni, A., Fujimoto, J., Watanabe, K. & Okada, S. (2006). *Lactobacillus durianis* Leisner et al. 2002 is a later heterotypic synonym of *Lactobacillus vaccinostrercus* Kozaki and Okada 1983. *International Journal of Systematic and Evolutionary Microbiology*, 56, 1721-1724. [https://DOI 10.1099/ijse.0.64316-0](https://doi.org/10.1099/ijse.0.64316-0).
17. Dunlap, C. A. (2015). Phylogenomic analysis shows that '*Bacillus vanillea*' is a later heterotypic synonym of *Bacillus siamensis*. *International Journal of Systematic and Evolutionary Microbiology* 65, 3507–3510. <https://doi.org/10.1099/ijsem.0.000444>.
18. Endo, A. & Okada, S. (2008). Reclassification of the genus *Leuconostoc* and proposals of *Fructobacillus fructosus* gen. nov., comb. nov., *Fructobacillus durionis* comb. nov., *Fructobacillus ficulneus* comb. nov. and

- Fructobacillus pseudoficulneus* comb. nov. *International Journal of Systematic and Evolutionary Microbiology*, 58, 2195–2205. <https://DOI.10.1099/ijs.0.65609-0>.
19. FAO/WHO Working Group (2002). Report on Drafting Guidelines for the Evaluation of Probiotics in Food London, Ontario, Canada, April 30 and May 1. https://www.who.int/foodsafety/fs_management/en/probiotic_guidelines.pdf (consulted last on March 21, 2019).
20. Felis, G. E., Vancanneyt, M., Snauwaert, C., Swings, J., Torriani, S., Castioni, A., & Dellaglio, F. (2006). Reclassification of *Lactobacillus thermotolerans* Niamsup et al. 2003 as a later synonym of *Lactobacillus ingluviei* Baele et al. 2003. *International Journal of Systematic and Evolutionary Microbiology*, 56, 793–795. [https://DOI: 10.1099/ijs.0.63992-0](https://DOI:10.1099/ijs.0.63992-0).
21. Felis, G. E. & Dellaglio, F. (2007). Taxonomy of lactobacilli and bifidobacteria. *Current Issues in Intestinal Microbiology*, 8, 44–61.
22. Garrity G. 2016. A new genomics-driven taxonomy of Bacteria and Archaea: Are we there yet?. *Journal of Clinical Microbiology*, 54, 1956–1963. [https:// DOI: 10.1128/JCM.00200-16](https://DOI:10.1128/JCM.00200-16).
23. Gevers, D., Cohan, F. M., Lawrence, J. G., Spratt, B. G., Coenye, T., Feil, E. J., Stackebrandt, E., Van de Peer, Y., Vandamme, P., Thompson, F. L., & Swings J. (2005). Opinion: Re-evaluating prokaryotic species. *Nature Reviews in Microbiology*, 3(9), 733–9. [https://DOI: 10.1038/nrmicro1236](https://DOI:10.1038/nrmicro1236)
24. Guarner, F. & Schaafsma, G. J. (1998) Probiotics. *International Journal of Food Microbiology*, 39, 237–238. [https://dx.doi.org/10.1016/S0168-1605\(97\)00136-0](https://dx.doi.org/10.1016/S0168-1605(97)00136-0).
25. Gupta, R. S. (2014). Chapter 8 - Identification of conserved indels that are useful for classification and evolutionary studies. *Methods in Microbiology*, 41, 153–182. Oxford, UK: Academic Press. <https://doi.org/10.1016/bs.mim.2014.05.003>.
26. Gupta, R. S. (2016). Impact of genomics on the understanding of microbial evolution and classification: the importance of Darwin's views on classification. *FEMS Microbiology Reviews*, 40, 520–553. [https://DOI: 10.1093/femsre/fuw011](https://DOI:10.1093/femsre/fuw011).
27. Gupta, R. S., Nanda, A. & Khadka, B. (2017). Novel molecular, structural and evolutionary characteristics of the phosphoketolases from bifidobacteria and Coriobacteriales. *PLoS ONE* 12(2), e0172176. <https://doi:10.1371/journal.pone.0172176>.
28. Herman, L., Chemaly M., Cocconcelli P.-S., Fernandez P., Klein G., Peixe L., Prieto M., Querol A., Suarez J. E., Sundh I., Vlak J. & Correia S. (2019). The qualified presumption of safety assessment and its role in EFSA risk evaluations: 15 years past. *FEMS Microbiology Letters*, 366, fny260; [https://doi: 10.1093/femsle/fny260](https://doi:10.1093/femsle/fny260).
29. Hugenholtz, P., Sharshewski, A., Parks, D. H. (2016). Genome-based microbial taxonomy coming of age. In Ochman, H. (Ed), *Microbial Evolution* (pp. 55–65). Cold Spring Harbor, New York, USA: Cold Spring Harbor Laboratory Press.
30. IJESM: <https://ijs.microbiologyresearch.org/content/journal/ijsem/about>
31. Institute of Medicine (US) Committee on Educating Public Health Professionals for the 21st Century. (2003). 6, Public Health Agencies: Their Roles in Educating Public Health Professionals. In Gebbie, K., Rosenstock, L., &

- 495 Hernandez, L. M. (Eds). *Who Will Keep the Public Healthy? Educating Public Health Professionals for the 21st*
496 *Century*. Washington (DC): National Academies Press (US). <https://www.ncbi.nlm.nih.gov/books/NBK221185/>.
- 497 32. Jakava-Viljanen, M., Murros, A., Palva, A., & Björkroth, K. J. (2008). *Lactobacillus sobrius* Konstantinov et al.
498 2006 is a later synonym of *Lactobacillus amylovorus* Nakamura 1981. *International Journal of Systematic and*
499 *Evolutionary Microbiology*, 58, 910-913. [https:// DOI: 10.1099/ijs.0.65432-0](https://doi.org/10.1099/ijs.0.65432-0).
- 500 33. Johansen, Eric. (2017). Future access and improvement of industrial lactic acid bacteria cultures. *Microbial*
501 *Cell Factories*, 16, 230. <https://doi.org/10.1186/s12934-017-0851-1>.
- 502 34. Kagermeier-Callaway, A. S., & Lauer, E. (1995). *Lactobacillus sake* Katagiri, Kitahara, and Fukami 1934 is the
503 senior synonym for *Lactobacillus bavaricus* Stetter and Stetter 1980. *International Journal of Systematic*
504 *Bacteriology*, 45, 398-399. [https:// DOI:10.1099/00207713-45-2-398](https://doi.org/10.1099/00207713-45-2-398).
- 505 35. Kaneuchi, C., Seki, M., & Komagata, K. (1988). Taxonomic study of *Lactobacillus mali* Carr and Davis 1970 and
506 related strains: validation of *Lactobacillus mali* Carr and Davis 1970 over *Lactobacillus yamanashiensis*
507 Nonomura. *International Journal of Systematic Bacteriology*, 38, 269-272. [https:// doi:10.1099/00207713-38-](https://doi.org/10.1099/00207713-38-3-269)
508 3-269.
- 509 36. Konstantinidis, K. T. & Tiedje, J. M. (2005a) Towards a Genome-Based Taxonomy for Prokaryotes. *Journal of*
510 *Bacteriology*, 187(18), 6258–6264. [https://doi:10.1128/JB.187.18.6258-6264.2005](https://doi.org/10.1128/JB.187.18.6258-6264.2005)
- 511 37. Konstantinidis, K. T. & Tiedje, J. M. (2005b) Genomic insights that advance the species definition for
512 prokaryotes. *Proceedings of the National Academy of Sciences USA*, 102(7), 2567–2572.
513 [https://doi:10.1073/pnas.0409727102](https://doi.org/10.1073/pnas.0409727102).
- 514 38. Koort, J., Vandamme, P., Schillinger, U., Holzapfel, W. H. & Björkroth, J. (2004). *Lactobacillus curvatus* subsp.
515 *melibiosus* is a later synonym of *Lactobacillus sakei* subsp. *carnosus*. *International Journal of Systematic and*
516 *Evolutionary Microbiology*, 54, 1621-1626. [https://DOI: 10.1099/ijs.0.63164-0](https://doi.org/10.1099/ijs.0.63164-0).
- 517 39. Kostinek, M., Pukall, R., Rooney, A. P., Schillinger, U., Hertel, C., Holzapfel, W. H. & Franz, C. M. A. P. (2005).
518 *Lactobacillus arizonensis* is a later heterotypic synonym of *Lactobacillus plantarum*. *International Journal of*
519 *Systematic and Evolutionary Microbiology*, 55, 2485-2489. [https:// DOI: 10.1099/ijs.0.63880-0](https://doi.org/10.1099/ijs.0.63880-0).
- 520 40. Lahteenmaki, L., & Ledeboer, A. M. (2006). Responsible exploitation of scientific data on pro- and prebiotics
521 for consumer health: a consensus document. *Food Science & Technology Bulletin*, 3, 47-50.
- 522 41. Lapage, S. P. (1992). International Code of Nomenclature of Bacteria. In: B. S. P. Lapage, P. H. A. Sneath, E. F.
523 Lessel, V. B. D. Skerman, H. P. R. Seeliger, & W. A. Clark (Eds). *Bacteriological Code, 1990 Revision*.
524 Washington (DC), ASM Press. ISBN-10: 1-55581-039-X. <https://www.ncbi.nlm.nih.gov/books/NBK8806/>.
- 525 42. Laulund, S., Wind, A. Derkx, P. M. F. & Zuliani V. (2017). Regulatory and Safety Requirements for Food
526 Cultures. *Microorganisms* 5, 28. [https://doi:10.3390/microorganisms5020028](https://doi.org/10.3390/microorganisms5020028).
- 527 43. Lawson, P. A., Citron, D. M., Tyrrell, K. L., & Finegold, S. M. (2016). Reclassification of *Clostridium difficile* as
528 *Clostridioides difficile* (Hall and O'Toole 1935) Prévot 1938. *Anaerobe*, 40, 95-9. [https:// DOI:](https://doi.org/10.1016/j.anaerobe.2016.06.008)
529 10.1016/j.anaerobe.2016.06.008.
- 530 44. Ledeboer, A. M. & Lahteenmaki, L. (2007) Probiotics – the consumer perspective. *Food Science and*
531 *Technology Bulletin*, 4, 43-49. [https:// DOI: 10.1616/1476-2137.14590](https://doi.org/10.1616/1476-2137.14590).

- 532 45. Ledeboer, A. M., Nauta, A., Sikkema, J., Laulund, E., Niederberger, P. & Sybesma, W. (2007). Survival of
533 probiotics. *European Dairy Magazine*, 4, 31-32.
- 534 46. Lugli, G. A., Milani, C., Duranti, S., Mancabelli, L., Mangifesta, M., Turrone, F., Viappiani, A., van Sinderen, D., &
535 Ventura, M. (2018a). Tracking the taxonomy of the genus *Bifidobacterium* based on a phylogenomic approach.
536 *Applied and Environmental Microbiology*, 84, e02249-17. <https://doi.org/10.1128/AEM.02249-17>.
- 537 47. Lugli, G. A., Mangifesta, M., Duranti, S., Anzalone, R., Milani, C., Mancabelli, L., Alessandri, G., Turrone, F.,
538 Ossiprandi, M. C., van Sinderen, D., & Ventura, M. (2018b). Phylogenetic classification of six novel species
539 belonging to the genus *Bifidobacterium* comprising *Bifidobacterium anseris* sp. nov., *Bifidobacterium criceti* sp.
540 nov., *Bifidobacterium imperatoris* sp. nov., *Bifidobacterium italicum* sp. nov., *Bifidobacterium margollesii* sp.
541 nov. and *Bifidobacterium parvae* sp. nov. *Systematic and Applied Microbiology*, 41(3), 173-183. [https://doi:](https://doi.org/10.1016/j.syapm.2018.01.002)
542 10.1016/j.syapm.2018.01.002. Epub 2018 Jan 31.
- 543 48. Makarova, K., Slesarev, A., Wolf, Y. Sorokin, A., Mirkin, B., Koonin, E., Pavlov, A., Pavlova, N., Karamychev, V.,
544 Polouchine, N., Shakhova, V., Grigoriev, I., Lou, Y., Rohksar, D., Lucas, S., Huang, K., Goodstein, D. M.,
545 Hawkins, T., Plengvidhya, V., Welker, D., Hughes, J., Goh, Y., Benson, Y., Baldwin, K., Lee, J.-H., Díaz-Muñoz, I.,
546 Dosti, B., Smeianov, V., Wechter, W., Barabote, R., Lorca, G., Altermann, E., Barrangou, R., Ganesan, R., Xie, Y.,
547 Rawsthorne, H., Tamir, D., Parker, C., Breidt, F., Broadbent, J., Hutkins, R., O'Sullivan, D., Steele, J., Unlu, G.,
548 Saier, M., Klaenhammer, T., Richardson, P., Kozyavkin, S., Weimer, B., & Mills, D. (2006). Comparative
549 genomics of the lactic acid bacteria. *Proceedings of the National Academy of Sciences U S A.*, 103(42), 15611–
550 15616. doi: 10.1073/pnas.0607117103.
- 551 49. Masco, L., Ventura, M., Zink, R., Huys, G., & Swings, J. (2004). Polyphasic taxonomic analysis of
552 *Bifidobacterium animalis* and *Bifidobacterium lactis* reveals relatedness at the subspecies level:
553 reclassification of *Bifidobacterium animalis* as *Bifidobacterium animalis* subsp. *animalis* subsp. nov. and
554 *Bifidobacterium lactis* as *Bifidobacterium animalis* subsp. *lactis* subsp. nov. *International Journal of Systematic
555 and Evolutionary Microbiology*, 54, 1137–1143. <https://DOI: 10.1099/ij.s.0.03011-0>.
- 556 50. Mattarelli, P., Holzapfel, W., Franz, C. M., Endo, A., Felis, G. E., Hammes, W., Pot, B., Dicks, L., & Dellaglio, F.
557 (2014). Recommended minimal standards for description of new taxa of the genera *Bifidobacterium*,
558 *Lactobacillus* and related genera. *International Journal of Systematic and Evolutionary Microbiology*, 64(Pt 4),
559 1434-1451. <https://doi: 10.1099/ij.s.0.060046-0>.
- 560 51. Milani, C., Lugli, G. A., Duranti, S., Turrone, F., Bottacini, F., Mangifesta, M., Sanchez, B., Viappiani, A.,
561 Mancabelli, L., Taminiau, B., Delcenserie, V., Barrangou, R., Margolles, A., van Sinderen, D., & Ventura, M.
562 (2014). Genomic encyclopedia of type strains of the genus *Bifidobacterium*. *Applied and Environmental
563 Microbiology*, 80, 6290 – 6302. <https://doi.org/10.1128/AEM.02308>.
- 564 52. Mora, D., Scarpellini, M., Franzetti, L., Colombo, S. & Galli, A. (2003). Reclassification of *Lactobacillus*
565 *maltaromicus* (Miller et al. 1974) DSM 20342T and DSM 20344 and *Carnobacterium piscicola* (Collins et al.
566 1987) DSM 20730T and DSM 20722 as *Carnobacterium maltaromaticum* comb. nov. *International Journal of
567 Systematic and Evolutionary Microbiology*, 53, 675-678. <https://DOI: 10.1099/ij.s.0.02405-0>.

53. Naser, S. M., Vancanneyt, M., Hoste, B., Snauwaert, C. & Swings, J. (2006). *Lactobacillus cypricasei* Lawson et al. 2001 is a later heterotypic synonym of *Lactobacillus acidipiscis* Tanasupawat et al. 2000. *International Journal of Systematic and Evolutionary Microbiology*, 56, 1681-1683. <https://DOI: 10.1099/ijs.0.64229-0>.
54. Naushad, H. S., Lee, B. & Gupta, R. S. (2014). Conserved signature indels and signature proteins as novel tools for understanding microbial phylogeny and systematics: identification of molecular signatures that are specific for the phytopathogenic genera *Dickeya*, *Pectobacterium* and *Brenneria*. *International Journal of Systematic and Evolutionary Microbiology*, 64, 366–383. <https://DOI: 10.1099/ijs.0.054213-0>.
55. Oren, A. (2019). Prokaryotic Nomenclature. Theory and Practice of Classification Practice. In Bergey's Manual of Systematics of Archaea and Bacteria. Published by John Wiley & Sons, Inc., in association with Bergey's Manual Trust. <https://doi.org/10.1002/9781118960608.bm00004.pub2>. (consulted last on March 21, 2019).
56. Pang, H., Kitahara, M., Tan, Z., Wang, Y., Qin, G., Ohkuma, M. & Cai, Y. (2012). Reclassification of *Lactobacillus kimchii* and *Lactobacillus bobalius* as later subjective synonyms of *Lactobacillus paralimentarius*. *International Journal of Systematic and Evolutionary Microbiology*, 62, 2383-2387. <https://DOI: 10.1099/ijs.0.035329-0>.
57. Parker, T. C., Tindall, B. J., & Garrity, G. M. (2016). International Code of Nomenclature of Prokaryotes - Prokaryotic Code (2008 Revision). *International Journal of Systematic and Evolutionary Microbiology*, 69, S1-S111. <https://DOI 10.1099/ijsem.0.000778>.
58. Parks, D. H. , Chuvochina, M., Waite, D. W., Rinke, C., Skarszewski, A., Chaumeil, P.-A. & Hugenholtz, P. (2018). A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nature Biotechnology*, 36, 996–1004. doi: <http://dx.doi.org/10.1101/256800>
59. Pot, B., Felis G. E., De Bruyne, K., Tsakalidou, E., Papadimitriou, K., Leisner, J., & Vandamme P. (2014). Chapter 19. The genus *Lactobacillus*. In W. H. Holzapfel B. J. B. Wood (Eds). *Lactic Acid Bacteria: Biodiversity and Taxonomy*; Wiley Online Library. <https://doi.org/10.1002/9781118655252.ch19>.
60. Public health agencies: https://en.wikipedia.org/wiki/List_of_national_public_health_agencies.
61. Public Health England, 2015. UK Standards for Microbiology Investigations, Identification of Enterobacteriaceae. Issued by the Standards Unit, Microbiology Services, PHE, Bacteriology – Identification | ID 16 | Issue no: 4 | Issue date: 13.04.15 | Page: 1 - 34. (https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/423601/ID_16i4.pdf). (consulted last on March 21, 2019).
62. Qin, Q. L., Xie, B. B., Zhang, X. Y., Chen, X. L., Zhou, B. C., Zhou, J., Oren, A., & Zhang, Y. Z. (2014). A proposed genus boundary for the prokaryotes based on genomic insights. *Journal of Bacteriology*, 196(12), 2210-2215. doi: 10.1128/JB.01688-14.
63. Ramasamy, D., Mishra, A. K., Lagier, J. C., Padhmanabhan, R., Rossi, M., Sentausa, E., Raoult, D., Fournier, P. E. (2014). A polyphasic strategy incorporating genomic data for the taxonomic description of novel bacterial species *International Journal of Systematic and Evolutionary Microbiology*, 64, 384–391. <https://doi.org/10.1099/ijs.0.057091-0>.
64. Rosselló-Mora, R. (2005). Updating Prokaryotic Taxonomy. *Journal of Bacteriology*, 187(18), 6255–6257. <https://doi: 10.1128/JB.187.18.6255-6257.2005>

65. Rosselló-Mora, R., Trujillo, M., & Sutcliffe, I. (2017). Introducing a digital protologue: a timely move towards a database-driven systematics of Archaea and Bacteria. *Systematic and Applied Microbiology*, 40, 121–122. <https://doi.org/10.1016/j.syapm.2017.02.001>
66. Salvetti, E., Felis, G. E., Dellaglio, F., Castioni, A., Torriani, S. & Lawson, P. A. (2011). Reclassification of *Lactobacillus cateniformis* (Eggerth 1935) Moore and Holdeman 1970 and *Lactobacillus vitulinus* Sharpe et al. 1973 as *Eggerthia cateniformis* gen. nov., comb. nov. and *Kandleria vitulina* gen. nov., comb. nov., respectively. *International Journal of Systematic and Evolutionary Microbiology*, 61, 2520–2524. <https://doi.org/10.1099/ijs.0.029231-0>.
67. Salvetti, E., Torriani, S. & Felis, G. E. (2012). The Genus *Lactobacillus*: A Taxonomic Update. *Probiotics & Antimicrobial Proteins*, 4(4), 217–226. <https://doi.org/10.1007/s12602-012-9117-8>.
68. Salvetti, E., Fondi, M., Fani, R., Torriani, S., & Felis, G. E. (2013). Evolution of lactic acid bacteria in the order Lactobacillales as depicted by analysis of glycolysis and pentose phosphate pathways. *Systematic and Applied Microbiology*, 36(5), 291–305. doi: 10.1016/j.syapm.2013.03.009.
69. Salvetti, E., Harris, H. M. B., Felis, G. E., & O'Toole, P. W. (2018). Comparative genomics of the genus *Lactobacillus* reveals robust phylogroups that provide the basis for reclassification. *Applied and Environmental Microbiology*, 84, e00993–18. <https://doi.org/10.1128/AEM.00993-18>.
70. Salvetti, E., & O'Toole, P. W. (2017). When regulation challenges innovation: The case of the genus *Lactobacillus*. *Trends in Food Science & Technology* 66, 187–194. <https://doi.org/10.1016/j.tifs.2017.05.009>.
71. SAM website: <https://www.elsevier.com/journals/systematic-and-applied-microbiology/07232020/guide-for-authors>.
72. Sangal, V., Goodfellow, M., Jones, A. L., Schwalbe, E. C., Blom, J., Hoskisson, P. A., & Sutcliffe, I. C. (2016). Next-generation systematics: An innovative approach to resolve the structure of complex prokaryotic taxa. *Scientific Reports*, 6, 38392. <https://doi.org/10.1038/srep38392>.
73. Schleifer, K. H., Kraus, J., Dvorak, C., Kilpper-Bälz, R., Collins, M. D. & Fischer, W. (1985). Transfer of *Streptococcus lactis* and related streptococci to the genus *Lactococcus* gen. nov. *Systematic and Applied Microbiology*, 6, 183–195. [https://doi.org/10.1016/S0723-2020\(85\)80052-7](https://doi.org/10.1016/S0723-2020(85)80052-7).
74. Segata, N., Börnigen, D., Morgan, X. C., Huttenhower, C. (2013). PhyloPhlAn is a new method for improved phylogenetic and taxonomic placement of microbes. *Nature Communication*, 4, 2304. <https://doi.org/10.1038/ncomms3304>.
75. Sneath, P. H. A. (2015) *Bergey's Manual of Systematics of Archaea and Bacteria*, Online © 2015 Bergey's Manual Trust, pp.83–88. This article is © 2001 Bergey's Manual Trust. Published by John Wiley & Sons, Inc., in association with Bergey's Manual Trust. <https://onlinelibrary.wiley.com/doi/pdf/10.1002/9781118960608.bm00004>. (consulted last on March 21, 2019).
76. Sun, Z., Harris, H. M. B., McCann, A., Guo, C., Argimón, S., Zhang, W., Yang, X., Jeffery, I. B., Cooney, J. C., Kagawa, T. F., Liu, W., Song, Y., Salvetti, E., Wroble, A., Rasinkanga, P., Parkhil, J., Rea, M. C., O'Sullivan, O., Ritari, J., Douillard, F. P., Ross, R. P., Yang, R., Briner, A. E., Felis, G. E., de Vos, W. M., Barrangou, R., Klaenhammer, T. R., Caufield, P. W., Cui, Y., Zhang, H. & O'Toole, P. W. (2015). Expanding the biotechnology potential of lactobacilli

- through comparative genomics of 213 strains and associated genera. *Nature Communications*, 6, 8322.
https://DOI: 10.1038.
77. Tindall, B. J., Kämpfer, P., Euzéby, J.-P. & Oren, A. (2006). Valid publication of names of prokaryotes according to the rules of nomenclature: past history and current practice. *International Journal of Systematic and Evolutionary Microbiology*, 56, 2715–2720. https://DOI 10.1099/ijs.0.64780-0
 78. Vancanneyt, M., Mengaud, J., Cleenwerck, I., Vanhonacker, K., Hoste, B., Dawyndt, P., Degivry, M. C., Ringuet, D., Janssens, D. & Swings, J. (2004). Reclassification of *Lactobacillus kefirgranum* Takizawa et al. 1994 as *Lactobacillus kefirnofaciens* subsp. *kefirgranum* subsp. nov. and emended description of *L. kefirnofaciens* Fujisawa et al. 1988. *International Journal of Systematic and Evolutionary Microbiology*, 54, 551-556.
https://DOI: 10.1099/ijs.0.02912-0.
 79. Vancanneyt, M., Engelbeen, K., De Wachter, M., Vandemeulebroecke, K., Cleenwerck, I. & Swings, J. (2005). Reclassification of *Lactobacillus ferintoshensis* as a later heterotypic synonym of *Lactobacillus parabuchneri*. *International Journal of Systematic and Evolutionary Microbiology*, 55, 2195-2198. https://DOI: 10.1099/ijs.0.63779-0.
 80. Vancanneyt, M., Naser, S. M., Engelbeen, K., De Wachter, M., Van Der Meulen, R., Cleenwerck, I., Hoste, B., De Vuyst, L. & Swings, J. (2006). Reclassification of *Lactobacillus brevis* strains LMG 11494 and LMG 11984 as *Lactobacillus parabrevis* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 56, 1553-1557. https://DOI: 10.1099/ijs.0.64215-0.
 81. Vandamme, P., Pot, B., Gillis, M., de Vos, P., Kersters, K. & Swings J. (1996). Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiology Reviews*, 60(2), 407-38
 82. Vernikos, G., Medini, D., Riley, D. R., Tettelin, H. (2015). Ten years of pangenome analyses. *Current Opinion in Microbiology*, 23, 148 –154. <https://doi.org/10.1016/j.mib.2014.11.016>.
 83. Weiss, N., Schillinger, U. & Kandler, O. (1983a). *Lactobacillus lactis*, *Lactobacillus leichmannii* and *Lactobacillus bulgaricus* subjective synonyms of *Lactobacillus delbrueckii* subsp. *lactis* comb. nov. and *Lactobacillus delbrueckii* subsp. *bulgaricus* comb. nov. *Systematic and Applied Microbiology*, 4, 552-557. https://DOI: 10.1016/S0723-2020(83)80012-5.
 84. Weiss, N., Schillinger, U. & Kandler, O. (1983b). *Lactobacillus trichodes*, and *Lactobacillus heterohiochii*, subjective synonyms of *Lactobacillus fructivorans*. *Systematic and Applied Microbiology* 4, 507-511.
https://DOI: 10.1016/S0723-2020(83)80008-3.
 85. Wittouck, S., Wuyts, S., Meehan, C. J, van Noort, V., & Lebeer, S. (2019). A genome-based species taxonomy of the *Lactobacillus* Genus Complex. bioRxiv preprint first posted online Jan. 31, 2019;
https://dx.doi.org/10.1101/537084.
 86. Wu, D., Hugenholtz, P., Mavromati, K., Pukall, R., Dalin, E., Ivanova, N. N., Kunin, V., Goodwin, L., Wu? M, Tindall, B. J., Hooper, S. D., Pati, A., Lykidi, A., Spring, S., Anderson, I. J., D’haeseleer, P., Zemla, A., Singer, M., Lapidus, A., Nolan, M., Copeland, A., Han, C., Chen, F., Cheng, J.-F., Lucas, S., Kerfeld, C., Lang, E., Gronow, S., Chain, P., Bruce, D., Rubin, E. M., Kyrpides, N. C., Klenk, H.-P. & Eisen, J. A. (2009). A phylogeny-driven genomic encyclopaedia of Bacteria and Archaea. *Nature*, 462, 24-31. https://doi:10.1038/nature08656

87. Yang, S. J., Kim, B. Y. & Chun, J. (2017). Rejection of reclassification of *Lactobacillus kimchii* and *Lactobacillus bobalius* as later subjective synonyms of *Lactobacillus paralimentarius* using comparative genomics. *International Journal of Systematic and Evolutionary Microbiology*, 67, 4515-4517. [https://DOI: 10.1099/ijsem.0.002326](https://doi.org/10.1099/ijsem.0.002326).
88. Yoon, S. H., Ha, S. M., Kwon, S., Lim, J., Kim, Y., Seo, H., & Chun, J. (2017). Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *International Journal of Systematic and Evolutionary Microbiology*, 67, 1613-1617. [https://DOI: 10.1099/ijsem.0.001755](https://doi.org/10.1099/ijsem.0.001755).
89. Zhang, Z.-G., Ye, Z.-Q., Yu, L., Shi, P. (2011). Phylogenomic reconstruction of lactic acid bacteria: an update. *BMC Evolutionary Biology* 11, 1. [https://DOI: 10.1186/1471-2148-11-1](https://doi.org/10.1186/1471-2148-11-1).
90. Zheng, J., Ruan, L., Sun, M., & Gänzle, M. (2015). A genomic view of lactobacilli and pediococci demonstrates that phylogeny matches ecology and physiology. *Applied and Environmental Microbiology*, 81, 7233–7243. [https://doi:10.1128/AEM.02116-15](https://doi.org/10.1128/AEM.02116-15).

Acknowledgements

This paper describes the outcome of an expert workshop entitled “Major changes in the taxonomy of

Bifidobacterium and *Lactobacillus*; consequences for industry” (<http://www.labip.com/news/>), held in Verona, October 4-5, 2018.

Presentations were given by invited speakers Giovanna Felis, Charles Franz, Marco Ventura, Paola Mattarelli, Paul W. O’Toole, Pier-Sandro Cocconcelli, and by E. Laulund and S. Duboux, representing LABIP members. Other participants were Aat Ledeboer, Alain Sourabie, Arjen Nauta, Bruno Pot, Christoph Bolten, Elisa Salvetti, Ingeborg Haagsma-Boels, Junji Fujimoto, Kaihei Oki, Koichi Watanabe, Luc De Vuyst, Marie-Eve Boyte, Marina Elli, Markus Brandt, Mary-Ellen Sanders, Monica Modesto, Patrice Antoine, Patrice Laforce, Patrick Boyaval, Peter Bron, Sander Sieuwerts, Sandra Torriani, Sophie Legrain-Raspaud, Svend Laulund, and Tadhg O’Sullivan.

The opinions expressed herein are those of LABIP and are not necessarily endorsed by all participants of the workshop, although all above approved the publication of this report.

This paper was written by BP on behalf of the Lactic Acid Bacteria Industrial Platform (LABIP) and complemented with research elements by ES, PM and GF, local organisers of the workshop. The authors wish to sincerely thank all the Workshop participants for their active and constructive contribution to the discussion and the reviewing of the manuscript.

Competing interests


BP is an employee of Yakult Europe BV.

Availability of data and materials

- 
- 713 Not applicable.
- 714 **Consent for publication**
- 715 Publication of this Communication was approved by LABIP and agreed to by the workshop participants.
- 716 **Ethics approval and consent to participate**
- 717 Not applicable.
- 718 **Funding**
- 719 The workshop was organised and sponsored by LABIP.


720 Figure 1: Historical evolution of the species number in the genus *Lactobacillus*.

721 Figure 1a: number of newly described *Lactobacillus* species per year.



722

723 Figure 1b: number of newly described *Lactobacillus* species per decade.



724

Invalid species names	Synonymous with existing species	Reference
<i>'Lactobacillus' arizonensis</i>	<i>Lactobacillus plantarum</i>	Kostinek et al. 2005
<i>'Lactobacillus' bavaricus</i>	<i>Lactobacillus sakei</i> subsp. <i>sakei</i>	Kagermeier-Callaway and Lauer 1995
<i>'Lactobacillus' bobalius</i>	<i>Lactobacillus paralimentarius</i>	Pang et al. 2012; but questioned by Yang et al. 2017
<i>'Lactobacillus' brevis</i>	<i>Lactobacillus parabrevis</i>	Vancanneyt et al. 2006
<i>'Lactobacillus' bulgaricus</i>	<i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i>	Weiss et al. 1983a
<i>'Lactobacillus' carnis</i>	<i>'Lactobacillus piscicola'</i> (see Table 2)	Mora et al. 2003
<i>'Lactobacillus' cypricasei</i>	<i>Lactobacillus acidipiscis</i>	Naser et al. 2006
<i>'Lactobacillus' casei</i> subsp. <i>alactosus</i>	<i>Lactobacillus paracasei</i> subsp. <i>paracasei</i>	Collins et al. 1989
<i>'Lactobacillus' casei</i> subsp. <i>pseudoplantarum</i>	<i>Lactobacillus paracasei</i> subsp. <i>pseudoplantarum</i>	Collins et al. 1989
<i>'Lactobacillus' casei</i> subsp. <i>rhamnosus</i>	<i>Lactobacillus rhamnosus</i>	Collins et al. 1989
<i>'Lactobacillus' casei</i> subsp. <i>tolerans</i>	<i>Lactobacillus paracasei</i> subsp. <i>tolerans</i>	Collins et al. 1989
<i>'Lactobacillus' cellobiosus</i>	<i>Lactobacillus fermentum</i>	Dellaglio et al. 2004
<i>'Lactobacillus' curvatus</i> subsp. <i>melibiosus</i>	<i>Lactobacillus sakei</i> subsp. <i>carnosus</i>	Koort et al. 2004
<i>'Lactobacillus' cypricasei</i>	<i>Lactobacillus acidipiscis</i>	Naser et al. 2006
<i>'Lactobacillus' durianis</i>	<i>Lactobacillus vaccिनosterus</i>	Dellaglio et al. 2006
<i>'Lactobacillus' ferintoshensis</i>	<i>Lactobacillus parabuchneri</i>	Vancanneyt et al. 2005
<i>'Lactobacillus' heterohiochii</i>	<i>Lactobacillus fructivorans</i>	Weiss et al. 1983b
<i>'Lactobacillus' kefirgranum</i>	<i>Lactobacillus kefiranofaciens</i> subsp. <i>kefirgranum</i>	Vancanneyt et al. 2004
<i>'Lactobacillus' kimchii</i>	<i>Lactobacillus paralimentarius</i>	Pang et al. 2012; but questioned by Yang et al. 2017
<i>'Lactobacillus' lactis</i>	<i>Lactobacillus delbrueckii</i> subsp. <i>lactis</i>	Weiss et al. 1983a
<i>'Lactobacillus' leichmannii</i>	<i>Lactobacillus delbrueckii</i> subsp. <i>lactis</i>	Weiss et al. 1983a
<i>'Lactobacillus' sobrius</i>	<i>Lactobacillus amylovorus</i>	Jakava-Viljanen et al. 2008
<i>'Lactobacillus' suntoryeus</i>	<i>Lactobacillus helveticus</i>	Naser et al. 2006
<i>'Lactobacillus' thermotolerans</i>	<i>Lactobacillus ingluviei</i>	Felis et al. 2006
<i>'Lactobacillus' trichodes</i>	<i>Lactobacillus fructivorans</i>	Weiss et al. 1983b
<i>'Lactobacillus' yamanashiensis</i>	<i>Lactobacillus mali</i>	Kaneuchi et al. 1988

Table 1: List of ‘invalid’ species names, as these species were later shown to be homologous to species that had already been described and validated earlier.

732
733
734
735

Old name	Synonymous with	Renamed to	Synonymous with (Final name)	Reference
' <i>Lactobacillus</i> ' <i>carnis</i>	' <i>Lactobacillus piscicola</i> '	' <i>Carnobacterium piscicola</i> '	<i>Carnobacterium maltaromaticum</i>	Mora et al. 2003
' <i>Lactobacillus</i> ' <i>catenaformis</i>			<i>Eggerthia catenaformis</i>	Salvetti et al. 2011
' <i>Lactobacillus</i> ' <i>confusus</i>			<i>Weissella confusa</i>	Collins et al. 1993
' <i>Lactobacillus</i> ' <i>dextrinicus</i>			<i>Pediococcus dextrinicus</i>	Back 1978
' <i>Lactobacillus</i> ' <i>divergens</i>			<i>Carnobacterium divergens</i>	Collins et al. 1987
' <i>Lactobacillus</i> ' <i>fructosus</i>		' <i>Leuconostoc</i> ' <i>fructosum</i>	<i>Fructobacillus fructosus</i>	Endo and Okada 2008
' <i>Lactobacillus</i> ' <i>halotolerans</i>			<i>Weissella halotolerans</i>	Collins et al. 1993
' <i>Lactobacillus</i> ' <i>kandleri</i>			<i>Weissella kandleri</i>	Collins et al. 1993
' <i>Lactobacillus</i> ' <i>maltaromicus</i>			<i>Carnobacterium maltaromaticum</i>	Mora et al. 2003
' <i>Lactobacillus</i> ' <i>minor</i>			<i>Weissella minor</i>	Collins et al. 1993
' <i>Lactobacillus</i> ' <i>minutus</i>			<i>Atopobium minutum</i>	Collins and Wallbanks 1992
' <i>Lactobacillus</i> ' <i>piscicola</i>		' <i>Carnobacterium piscicola</i> '	<i>Carnobacterium maltaromaticum</i>	Mora et al. 2003
' <i>Lactobacillus</i> ' <i>rimae</i>			<i>Atopobium rimae</i>	Collins and Wallbanks 1992
' <i>Lactobacillus</i> ' <i>uli</i>			<i>Olsenella uli</i>	Dewhirst et al. 2001
' <i>Lactobacillus</i> ' <i>viridescens</i>			<i>Weissella viridescens</i>	Collins et al. 1993
' <i>Lactobacillus</i> ' <i>vitulinus</i>			<i>Kandleria vitulina</i>	Salvetti et al. 2011
' <i>Lactobacillus</i> ' <i>xylosus</i>			<i>Lactococcus lactis</i> subsp. <i>lactis</i>	Schleifer et al. 1985

736
737
738
739
740
741
742
743
744
745
746
747
748

Table 2: List of *Lactobacillus* species that have already been transferred to other genera.

749
750
751 Supplementary information for the paper

752

753 The potential impact of the *Lactobacillus* name change: the results of an
754 expert meeting organised by the Lactic Acid Bacteria Industrial Platform
755 (LABIP).

756
757 Bruno Pot^{1,2*}, Elisa Salvetti³, Paola Mattarelli⁴ and Giovanna E. Felis³
758

759

760

761 Bacterial names are composed of a genus and a species name (rule 12a), always to be mentioned together. An additional subgenus or subspecies name can
762 be provided if this improves the clarity of the taxonomic situation or improves communication. Once validly published, as mentioned above, names cannot be
763 changed and an existing name can never be used again to describe another organism. Even if the organism is transferred to another species, e.g. because of
764 duplicate names (Table 1), the name that was described later in time, and therefore is redundant based on the rule that states that older names have priority,
765 cannot disappear (Rule 23) but can also not be used again later to name another organism, as a name is associated to the type strain of the species (Rule 15).
766 While these rules aim to improve the stability of taxonomy, it is one of the reasons why the number of species in a genus can become quite high.

767 Furthermore, a taxonomic name should be unambiguous and therefore a genus should always be described with a type species. The type species will remain
768 the core of the genus, regardless of future proposed taxonomic changes (Rules 39a and 39b of the [Code](#); last consulted 29th March 2019). For the genus
769 *Lactobacillus*, the type species is *Lactobacillus delbrueckii*. This is important as the revised genus *Lactobacillus* must be anchored around this species.

770 Furthermore, for each species, a type strain is chosen and deposited in a public culture collection (if the species was described before 31 December 2000) or
771 at least two culture collections (if the species was described after 1st of January 2001) (Rule 30). In absence of such type strain, a taxon cannot be validated.
772 Also, the choice of the type strain has practical importance, as illustrated by the taxonomic issues around the description of *Lactobacillus paracasei* for
773 aberrant strains of the species *L. casei* (Hill et al., 2018). The type strain of *L. delbrueckii*, and therefore of the current and future genus *Lactobacillus*, is ATCC®
774 9649D-5™.

775

776 Reference

777 Hill, D., Sugrue, I., Tobin, C., Hill, C., Stanton, C. & Ross, R. P. (2018) The *Lactobacillus casei* Group: History and Health Related Applications. *Frontiers in*
778 *Microbiology* 9, 2107. [https://doi: 10.3389/fmicb.2018.02107](https://doi.org/10.3389/fmicb.2018.02107).
779
780

Highlights Pot et al.

- Genome sequences reveal a high heterogeneity within the genus *Lactobacillus*
- What is the effect for the industry of renaming 200 lactobacilli?
- Industry, regulators, doctors and scientists prefer a soft but profound nomenclatural correction