

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Functional and sensory properties of phenolic compounds from unripe grapes in vegetable food prototypes

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Bucalossi G., Fia G., Dinnella C., De Toffoli A., Canuti V., Zanoni B., et al. (2020). Functional and sensory properties of phenolic compounds from unripe grapes in vegetable food prototypes. FOOD CHEMISTRY, 315, 1-9 [10.1016/j.foodchem.2020.126291].

Availability:

This version is available at: <https://hdl.handle.net/11585/805410> since: 2024-05-14

Published:

DOI: <http://doi.org/10.1016/j.foodchem.2020.126291>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **Functional and sensory properties of phenolic compounds from unripe grapes in vegetable** 2 **food prototypes**

3

4 Ginevra Bucalossi^a, Giovanna Fia^a, Caterina Dinnella^a, Alessandra De Toffoli^a, Valentina Canuti^a,
5 Bruno Zanoni^a, Maurizio Servili^b, Ella Pagliarini^c, Tullia Gallina Toschi^d, Erminio Monteleone^{a*}

6

7 ^a Department of Agricultural, Food, Environmental and Forestry Sciences and Technologies
8 (DAGRI) ,University of Florence, Piazzale delle Cascine, 18 - 50144 - Firenze, Italy

9 ^b Department of Agricultural, Food and Environmental Sciences, University of Perugia, Borgo XX
10 Giugno, 74 - 06121 - Perugia, Italy

11 ^c Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milan, Via
12 Giovanni Celoria, 2, - 20133 – Milano, Italy

13 ^d Department of Agricultural and Food Sciences (DiSTAL), Alma Mater Studiorum - University of
14 Bologna, Italy

15 ginevra.bucalossi@unifi.it, giovanna.fia@unifi.it, caterina.dinnella@unifi.it,
16 alessandra.detoffoli@unifi.it, valentina.canuti@unifi.it, bruno.zanoni@unifi.it,
17 maurizio.servili@unipg.it, ella.pagliarini@unimi.it, tullia.gallinatoschi@unibo.it

18

19 * Corresponding author: erminio.monteleone@unifi.it

20

21 **Abstract**

22 Unripe grapes (UGs) from thinning are an unexploited source of phenols useful as functional
23 ingredient. However, phenols may negative affect sensory quality of food. Chemical and sensory
24 properties of UG phenols in plant-based foods were not investigated before.

25 With this aim, an extract from UGs, obtained by a green extraction technique, was used to fortify
26 three plant-based food models: carbohydrates/acidic pH/sweet - beetroot purée, proteins/neutral
27 pH/sweet - pea purée and starch/neutral pH - potato purée.
28 Functional and sensory properties of phenol-enriched foods varied as a function of their
29 composition and original taste. The amount of UG phenols recovered from potato purée was higher
30 than that recovered from beetroot and pea purée, while the antioxidant activity detected in beetroot
31 purée was higher than that in potato and pea purée. Significant variations of sourness, saltiness,
32 bitterness and astringency were induced by UG phenols added to food models. Beetroot purée
33 resulted more appropriate to counteract the negative sensations induced by UG phenols.

34

35 **Keywords:** functional food; unripe grapes; polyphenols; antioxidant activity; sourness; sweetness.

36

37 1. Introduction

38 By-products of the wine industry are rich in phenols and other valuable elements for the human diet
39 such as mineral salts, fibres and vitamins. There are emerging evidences of the potential preventive
40 effects of grape polyphenols towards cardiovascular diseases, diabetes, and degenerative diseases
41 such as cancer (Guilford & Pezzuto, 2011; Mihaylova, Popova, Alexieva, Krastanov & Lante,
42 2018). The role of phenols from grapes in the prevention of various diseases associated with
43 oxidative stress is primarily related to their antioxidant properties (Guilford & Pezzuto, 2011;
44 Villaño, Fernández-Pachón, Moyá, Troncoso, & García-Parrilla, 2007; Rasines-Perea & Teissedre,
45 2017).

46

47 The sustainability of the winemaking process could be improved by the recovery of high-value
48 bioactive compounds from by-products. Indeed, extensive studies have been made of the biological
49 properties, extraction techniques and applications in the food system of phenols from grape pomace,
50 the main by-product of the wine industry (Beres et al., 2017; Yu & Ahmedna, 2013).

51 Unripe grapes (UGs) discarded during thinning are an undervalued by-product of vineyard
52 management for the production of high-quality wine (Gatti, Bernizzoni, Civardi, & Poni, 2012;
53 Keller, Mills, Wample, & Spayd, 2005; Ough Cs, 1984). In unripe berries, the most important
54 classes of grape antioxidants (phenolic acids, flavan-3-ols, flavonols, anthocyanins, stilbenes and
55 glutathione) are present to variable extents in function of some factors such as variety, maturity
56 level and season (Adams, 2006) but their anti
57
58
59 oxidant activity and potential application have received scarce scientific attention (Fia, Gori,
60 Bucalossi, Borghini, & Zanoni, 2018; Tinello & Lante, 2017). Low-quality unripe grapes are
61 processed into various traditional juices and sauces with a low pH and variable levels of antioxidant
62 activity ((Dupas de Matos, Magli, Marangon, Curioni, Pasini & Vincenzi, 2018; Öncül &
63 Karabiyikli, 2015). The added value of thinned grapes is higher than the one of other by-products of
64 wine industry that were largely studied and proposed as source of antioxidants. That is because, the
65 thinned grapes have not been exploited to make wine and therefore contain an intact complex of
66 bio-active compounds. Recently, a green extraction technique (i.e. performed without solvents and
67 preservatives) was patented (Fia & Gori, 2016) and applied at an industrial level with the aid of a
68 patented oenological machine (Gori, C., Menichetti, S., & Fia, G. 2014) to obtain an extract from
69 unripe grapes.

70
71 Functional food is essentially a marketing term with different definitions and regulations depending
72 on the country (Henry, 2010). Recently in Europe, there has been a growing interest in functional
73 foods. A scientific consensus document was drafted to develop a science-based approach for the
74 emerging concepts in functional food (Europe, 1999). Foods that have been modified by enrichment
75 with bioactive substances are included in the functional food categories and the health benefits of

76 phenols, beyond basic nutritional values of plant-based food and beverages containing phenols, are
77 reported in a recent review (Shahidi & Ambigaipalan, 2015).

78 Phenols from plant by-products have been proposed as ingredients for functional foods and
79 beverages preparation to improve their nutritional characteristics (De Toffoli et al., 2019; Torri et
80 al., 2015; Nirmala, Bisht, Bajwa, & Santosh, 2018; Świeca, Gawlik-Dziki, Sęczyk, Dziki, & Sikora,
81 2018). Some examples of functional food enriched with phenols from tea and Guava are already
82 included in the “food for specified health uses” (FOSHU) and regulated as functional food in Japan
83 (Iwatani & Yamamoto, 2019).

84 In developing a phenol-enriched functional food, two main aspects need to be investigated: the first
85 concerns the phenols’ stability after their addition to the food system, affecting the preservation of
86 their biological activities; the second concerns oral sensations, such as astringency, bitterness and
87 sourness, which can arise after the addition of phenols to food and impair the acceptability of the
88 product to consumers.

89

90 From a sensory point of view, it is well documented that phenolic compounds contribute to the
91 bitter and astringent oral sensation of food and beverages (Hufnagel & Hofmann, 2008) and this
92 significantly affects the preference and choice of phenol-rich vegetable foods (Dinnella, Recchia,
93 Tuorila, & Monteleone, 2011). Monomeric and polymeric phenols have been widely studied
94 because of their contribution to wine sensory perception. Monomeric flavan-3-ols, procyanidin
95 dimers and trimers seem to be involved in the perception of astringency and bitterness in red wine
96 (Peleg, Gacon, Schlich, & Noble, 1999). Several authors have studied the bitterness of polyphenols
97 in red wine, demonstrating that larger molecules tend to be less bitter and more astringent (Peleg et
98 al., 1999). More recently, in reconstruction studies it was observed that the puckering astringent
99 offset was caused by a polymeric fraction exhibiting molecular masses above >5 kDa and it was
100 found to be amplified by organic acids (Hufnagel & Hofmann, 2008). Some factors such as pH,
101 acidity, carbohydrate content and saliva characteristics could affect oral sensations (Dinnella,

102 Recchia, Fia, Bertuccioli, & Monteleone, 2009; Fia, Dinnella, Bertuccioli, & Monteleone, 2009; de
103 Freitas & Mateus, 2012).

104

105 To mitigate functional phenol's bitter and astringent potential, the naturally occurring interactions
106 phenols/biopolymers in vegetable foods (Zhang et al., 2014) are an effective strategy (De Toffoli et
107 al., 2019). Plant biopolymers can act as a physical barrier for the phenol stimuli utilized, thus
108 hindering their interactions with sensory receptors and saliva. Many factors affect
109 phenol/biopolymer binding, including pH and reagent features such as chemical compositions,
110 structure, and hydrophobic/hydrophilic characteristics (Kroll, Rawel, & Rohn, 2003). Furthermore,
111 several studies have investigated the chemical features of phenol/biopolymer interactions and their
112 consequences on sensory attributes (Jakobek, 2015).

113

114 The health effects of phenols depend on the consumed amount and on their bioavailability. The
115 bioavailability of phenols may vary depending on their bioaccessibility, referred as the release from
116 the food matrix, their stability against several biochemical factors, and their later intestinal
117 absorption (Sengul, Surek & Nilufer-Erdil, 2014). The bioavailability of phenols from many
118 different vegetable sources, including grapes, was systematically studied by Manach, Scalbert,
119 Morand, Rémésy, & Jiménez (2004). In humans, among the most well absorbed phenols there are
120 gallic acid, catechins and quercetin glucosides (Manach et al., 2004). Recently, a phenol extract
121 from grape pomace was included in the diet of Wistar rats by Olivero-David et al., (2018). The
122 same authors observed a partial bioavailability of the phenol extract and an improvement in lipid
123 metabolism of rats.

124 During food processing, bioactive compounds may undergo chemical degradation and lose their
125 activities. Thermal processing and long-term storage can lead to a decrease in both polyphenol
126 content and antioxidant activity (Yu & Ahmedna, 2013). Other factors such as pH and interactions
127 with other macromolecular food constituents can affect the chemical stability and antioxidant

128 activity of phenolic compounds (Jakobek, 2015). It is emerging that the bioaccessibility and
129 bioavailability of phenolic compounds are affected by interaction with other macromolecules such
130 as proteins, carbohydrates and lipids. These interactions could give phenolic compounds protection
131 from oxidation during their passage through the gastrointestinal tract (Saura-Calixto, 2011). On the
132 other hand, phenol/protein interactions can lead to a loss of nutritional values due to protein
133 precipitation and enzyme inactivation (Rohn, Petzke, Rawel & Kroll, 2006).

134

135 Variations in chemical composition, antioxidant activity and sensory profiles in food-base
136 vegetables with added phenols from unripe grapes have never been investigated before.

137

138 This paper explores the chemical and sensory properties of phenols extracted from UGs and the
139 consequences of phenol/biopolymer interactions on the chemical and sensory properties of plant-
140 base foods. With this aim, three food models with variable macro-compositions in which different
141 phenol/biopolymer interactions might occur were functionalised with an extract from unripe grapes
142 (UGs).

143

144 **2. Material & Methods**

145 *2.1. UG extract and UG-water solutions preparation*

146 The unripe grapes (UGs), cv Merlot, were hand-picked in August 2017 in a commercial vineyard
147 located in Velletri, Rome, Italy. To obtain the UG extract, maceration was performed as previously
148 described by Fia et al. (2018), with some modifications (**Fig. S1**). After decantation and filtration of
149 the liquid extract, sugar was eliminated by ultrafiltration, using a spiral wound configuration
150 membrane, with a molecular weight cut-off of 2500 Dalton (General Electrix, Boston,
151 Massachusetts, United States). The liquid extract was dehydrated by lyophilization with the addition
152 of arabic gum (2% w/v) (Nexira Food, Rouen Cedex, France) as a support and stored in
153 polyethylene pouches under vacuum, in a desiccator, at room temperature, protected from the light.

154 The UG extract (334 g) was diluted in distilled water to a total volume of 1L. This suspension was
155 centrifuged at 1646 g, for 10 min, to eliminate the excess arabic gum. The phenol concentration in
156 the supernatant UG stock solution (SS) was 6.81 g/L. The SS was daily prepared and used to
157 prepare UG-water solutions at different phenol concentrations to be added to the plant-based food
158 models (**Fig. S1**).

159 The UG-water solutions were filtered through a membrane (\varnothing 0.45 μ m) and the phenolic
160 compounds were purified using a C18 Sep-pak cartridge (1 g) (Waters, Milan, Italy) before the
161 evaluation of the total polyphenol content.

162

163 *2.2 Food models*

164 Three food models were selected on the basis of their composition (**Table S1**) and taste: beetroot
165 purée (BP) characterized by high carbohydrate content, acidic pH and sweet taste; pea purée (PeP)
166 characterized by high proteins content, neutral pH and sweet taste; potato purée (PoP) characterized
167 by high carbohydrates content and neutral pH. Canned or powdered ingredients produced by large
168 food companies were used to prepare the food models, since they are not subject to seasonal
169 restriction and their composition is constant. Purées of beetroot, pea and potato were prepared as
170 following: a) 500 g of peeled and steamed beetroots were blended at maximum speed, for about 1
171 min, using a Kenwood FDM 780 mixer (Kenwood, Treviso, Italy), until it was obtained a
172 homogeneous product; b) 310 g of steamed peas were rinsed under cold water for 30 sec and
173 drained for 30 sec to eliminate the water, then 7 g of water were added and the mix was blended at
174 maximum speed for 2 min in a mixer Kenwood; c) 75 g of dehydrated potatoes were added to 340 g
175 of water brought to 80°C and the product was mixed until it became homogeneous, then it was
176 cooled for 30 min before using. Each food model was prepared at five levels of phenol
177 concentration (0.00, 0.21, 0.44, 1.11 and 1.93 g/kg) (**Fig. S1**).

178

179 *2.3. Chemicals*

180 All solvent and reagents were supplied from Sigma-Aldrich (Milan, Italy), except for methanol and
181 ethanol which were supplied by Carlo Erba (Milan, Italy). Ultrapure water was obtained using a
182 Milli-Q Gradient water purification system (Thermo Scientific, Waltham, Massachusetts, USA).

183

184 2.4. *Physical-chemical analysis*

185 2.4.1 *General analysis*

186 Total acidity and pH were evaluated according to the methods recommended by the International
187 Organization of Vine and Wine (OIV) (International Organization of Vine and Wine Website,
188 2014).

189 2.4.2. *Moisture content and water activity*

190 The powder moisture content was determined gravimetrically by drying in a vacuum oven, at 70°C,
191 until a constant weight was reached (A.O.A.C. , 1990) . Powder water activity (A_w) was measured
192 using a Rotronic Hygroskop *DT* hygrometer (Michell Italia Srl, Milan, Italy).

193 2.4.3. *Solubility*

194 Water solubility was determined according to (Cano-Chauca, Stringheta, Ramos, & Cal-Vidal,
195 2005). A volume of 100 mL of distilled water was transferred into a blender jar. The sample (1g,
196 dry basis) was carefully added to the blender while operating at high speed for 5 min. The solution
197 was centrifuged at 3000 g for 5 min. An aliquot of 25 mL of the supernatant was transferred to pre-
198 weighed Petri dishes and immediately oven-dried at 105°C for 5 h. The solubility (%) was
199 calculated by weight difference.

200 2.4.4. *Hygroscopicity*

201 Hygroscopicity was evaluated following the method described by Callahan et al. (1982), with some
202 modifications. The equilibrium moisture content (EMC) of the samples (1 g, dry basis) was
203 evaluated following storage in desiccators containing saturated salt solutions with a relative
204 humidity ranging from 8% to 84% at 25°C until a constant weight was reached (approx. 21 days).
205 The hygroscopicity was expressed as g of adsorbed water per 100 g of dry matter (g/100g dm).

206 2.4.5. Phenol extraction

207 Extracts were obtained from the food models (FMs) following the method described by Turkmen,
208 Sari, & Velioglu (2005). For each food matrix, 1 g was homogenized and extracted twice with 4.5
209 mL of 80% aqueous methanol solution in a mechanical shaker, for 2 h. The mixture was centrifuged
210 at 13440 g, for 15 min, at room temperature, and the supernatant decanted into polypropylene tubes.
211 The supernatant was filtered through Whatman No.1 filter paper. The extraction procedure was
212 performed in triplicate.

213 2.4.6. Total polyphenol

214 The total polyphenols (TP) were quantified according to the Folin-Ciocalteu method (Singleton,
215 Rossi Jr., & Rossi J A Jr., 1965). A Perkin Elmer Lambda 10 spectrophotometer (Waltham, MA,
216 USA) was used to measure the absorbance of the reaction mixture at 700 nm. A standard curve was
217 obtained with (+)-catechin solutions at concentrations ranging from 5 to 500 mg/L. The TP was
218 expressed as mg of (+)-catechin equivalents/L of the UG-water solution or kg of the food model
219 extracts.

220 2.4.7. Antioxidant activity

221 Antioxidant activity was evaluated by 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay (Brand-
222 Williams, Cuvelier, & Berset, 1995). Trolox standard solutions were prepared daily in absolute
223 ethanol at concentrations ranging from 10 to 600 µmol/L. Antioxidant activity was expressed as
224 µmol of Trolox equivalent antioxidant capacity (TEAC)/L of the solution or kg of the food model
225 extract.

226 2.4.8. LC-HRMS analysis

227 Analysis of the phenolic compounds and glutathione was performed via liquid chromatography –
228 high-resolution mass spectrometry (LC-HRMS), according to Fia et al. (2018) using an Accela
229 1250 (Thermo Fisher Scientific) coupled with an LTQ OrbitrapExactive mass spectrometer
230 (Thermo Fisher Scientific) equipped with an electrospray ionization (ESI) source in negative mode.
231 The standards were purchased from Sigma-Aldrich (Milan, Italy), except for the quercetin 3-O-

glucoside which was supplied by Analytik GmbH (Rülzheim, Germany). Coumaric and ferulic acids were used as standards for coumaric and ferulic acids due to the lack of reference materials. Data were expressed as mg of phenols/kg of the UGs or food models.

2.5. Sensory evaluations

The present data were collected as part of a larger study aimed at investigating factors affecting the acceptability of health foods (PRIN 2015: Individual differences in the acceptability of health foods: focus on phenol and fat content). This multisession study consisted of a home questionnaire session and one-on-one testing in a sensory laboratory across two days. This paper will only present a selection of these data. The sensory tests are further detailed in De Toffoli et al. (2019). Two respondent groups were recruited to evaluate the UG extract (Group 1: n=29; 59% females; mean age 27.5 ± 7.1) or functionalized food prototypes (Group 2: n=27; 70% females; mean age 31.5 ± 9.4). The participants received a gift to compensate for their time. The respondents gave their written informed consent at the beginning of the test according to the principles of the Declaration of Helsinki. In brief, training was performed as described by Monteleone et al., (2017) using the general Labelled Magnitude Scale - gLMS (0: no sensation-100: the strongest imaginable sensation of any kind) (Green et al., 2007). Eight water solutions of UG extract were prepared as sensory stimuli with increasing phenol concentration: 0.14, 0.21, 0.30, 0.41, 0.59, 1.11, 1.27 and 1.93 g/L of phenol (**Fig. S1**). The data were collected using Fizz software (ver.2.51. A86, Biosystèmes, Couternon, France).

2.6. Data analysis

A one-way ANOVA model was used to assess the storage effect on the variation of phenol content and antioxidant activity of the UG extract. Two-way ANOVA models were used to assess the effect of both phenol concentration and replicates on the antioxidant activity in the UG solutions and to

257 assess the effect of both the amount of phenol added and replicates on the recovery of UG phenols
258 from food models.

259 The UG phenols recovered (recovery %) from the functionalized food samples were calculated as
260 the difference between the total phenol content of the functionalized food and that of the non-
261 functionalized food, then it was expressed as percentage of the phenols added. Two-way ANOVA
262 models were used to assess the effect of phenol concentration on the intensity of the target
263 sensations in UG solutions and food prototype samples (phenol concentration were used as fixed
264 factor; subjects were considered as random factor). Three-way ANOVA were used to assess the
265 effect of the food matrix on the perceived intensity of the target sensations models (fixed factors:
266 food matrix and phenol concentration; random factor: subjects and interactions). A p -value of 0.05
267 was considered as the threshold for statistical significance.

268 Data analysis was performed using XLSTAT statistical software package (Addinsoft - version
269 19.02).

270

271 **3. Results**

272

273 *3.1. Physical-chemical characterization*

274 *3.1.1. UG extract*

275 The solubility of the UG extract was $88.1 \pm 1.2\%$. The moisture content of the UG extract, at 25°C ,
276 was $8.1 \pm 0.3\%$ and the water activity was $38.7 \pm 0.1\%$. The adsorption isotherm of the UG extract
277 at 25°C was determined (**Fig. S2**). The experimental data for water activity (A_w) as a function of the
278 moisture content fitted well with the Halsey model (Okos et al., 1992), as follows:

279

$$280 \quad A_w = \exp\left(-\frac{B}{n_s^A}\right) \quad (r^2 = 0.98)$$

281

282 where n_s (g water/g dry matter), $A = 0.039$ and $B = 1.461$.

283 The powder displayed little hygroscopic behaviour up to A_w values < 0.80 , while for A_w values
284 greater than 0.85 the hygroscopicity increased exponentially.

285

286 The total phenol content of the UG extract was 20403 ± 943 mg/kg. The total phenol content of the
287 UG extract was evaluated monthly until to nine months of storage. After this period, the UG extract
288 displayed the same phenolic concentration as the outset. No significant differences ($p = 0.05$) were
289 assessed among phenolic content values during storage.

290

291 The phenolic composition of the UG extract was analysed by LC-HRMS. Nineteen phenolic
292 compounds were identified in the UG extract (**Table 1**). Phenolic acids were the most abundant
293 class of phenolic compounds and they accounted for 89% of the amount of phenols identified in the
294 UG extract. Caftaric acid accounted for 85% of the phenolic acid content. Flavonols, flavan-3-ols,
295 procyanidins, trans-resveratrol and 2-S-glutathionyl fertaric acid accounted for the remaining 11%
296 of the amount of phenols detected in the UG extract.

297

298 The antioxidant activity of the UG extract was 33829 ± 949 TEAC $\mu\text{mol/kg}$, and the specific
299 activity of the phenols was 1.66 ± 0.04 TEAC $\mu\text{mol/mg}$. The antioxidant activity of the UG extract
300 was evaluated monthly, up to nine months of storage. After this period, the antioxidant activity of
301 the UG extract remained at 99.4%. No significant differences ($p = 0.05$) were assessed in the
302 antioxidant activity values at different times of storage.

303 3.1.2. UG water solutions

304 The total phenol content of the stock solution was 6.81 ± 0.04 g/L. The stock solution was
305 characterized for total acidity (7.6 ± 0.26 g/L as tartaric acid) and pH (3.21 ± 0.02). The solutions
306 from the UG extract were tested for antioxidant activity at increasing phenol concentration levels
307 (0.14, 0.21, 0.30, 0.41, 0.59, 1.11, 1.27 and 1.93 g/L) (**Fig. S3**). The UG phenol concentration
308 significantly affected the level of antioxidant activity of the water solutions ($p \leq 0.001$) while the

309 replicates were not significant ($p < 0.05$). A significant positive relationship ($r = 0.978$) was found
310 between the total phenol content and the antioxidant activity of the UG water solutions.

311

312 3.1.3. Functionalized food models

313 After the addition of an increasing amount (0.00, 0.21, 0.44, 1.11 and 1.93 g/kg) of UG phenols to
314 the food models, the phenol concentration in the FM extracts was determined (**Fig. 1A**). The non-
315 functionalized food models showed different phenolic content, with the highest level detected in the
316 beetroot purée and the lowest in the potato purée. The amount of phenols added to the food models
317 significantly affected the concentration of phenols found in the FM extracts ($p \leq 0.05$).

318

319 The phenols recovered from food models significantly varied as a function of both the food model
320 and the amount of phenols added. The recovered amount ranged from 27.7% to 81.3% in the
321 beetroot purée, from 34.0% to 53.6% in the pea purée and from 52.7% to 86.4% in the potato purée.
322 The mean phenol value recovered with the highest added amount of phenols was highest in the
323 potato purée (68.7%), followed by the beetroot purée (57.8%), and the pea purée (43.3%). (**Fig.**
324 **1B**).

325

326 The food samples functionalized with the highest amount of phenols (1.93 g/kg) were extracted and
327 the extracts analysed via LC-HRMS to evaluate their phenol composition. The FM extracts
328 contained almost all of the phenolic compounds identified in the original UG extract, except for
329 kaempferol-3-*O*-glucoside, quercetin-3-*O*-hexoside and 2-S-glutathionyl caftaric acid (**Table 1**).
330 Caftaric acid was the most abundant phenolic compound assayed in the FM extracts of the three
331 food models. Ferulic acid was not detected in the potato purée. The phenol profiles of the food
332 model functionalized with 1.93 g/kg of UG phenols were compared to the profile of the UG extract
333 (**Fig. 1C**). The relative amounts of each phenolic class in functionalized beetroot purée was similar
334 to that observed in the UG extract, while slight differences were observed in the functionalized pea

335 and potato purées. Phenolic acids represented the most abundant class of phenols in the UG extract
336 (90.3%) and the beetroot purée almost retained this same high percentage (88.9%), while in the pea
337 and potato purées a slight loss was observed (80.6 and 83.9%, respectively). The proportion of other
338 phenolic classes (flavonols, flavan-3-ols, procyanidins and stilbenes) was slightly higher in the pea
339 and potato purées compared to the figure observed in the UG extract and the beetroot purée.

340

341 The antioxidant activity of the food models with an increasing added amount (0.00, 0.21, 0.44, 1.11
342 and 1.93 g/kg) of UG phenols was determined after extraction (**Fig. 2A**). The non-functionalized
343 beetroot and pea purées had similar values of antioxidant activity while it was much lower in the
344 potato purée. A significant increase in antioxidant activity was observed in the beetroot purée as
345 function of the UG phenol concentration. No significant difference was observed between the
346 antioxidant activity of the pea purée functionalized with 0.44 or 1.11 g/kg of UG phenols.

347 The difference between the antioxidant activity of functionalized food and that of food without
348 added phenol was calculated to assess the contribution of UG phenols to the food models' final
349 antioxidant activity. The relationship between the antioxidant activity of UG phenols in the water
350 solution and in the FM extracts is shown in **Figure 2B**. The antioxidant activity was always
351 significantly higher in the extracts of beetroot purée compared to that detected in the potato and pea
352 purée extracts. The mean antioxidant activity was 3794 $\mu\text{mol/kg}$ in the BP, 1722 $\mu\text{mol/kg}$ in the
353 PoP and 1127 $\mu\text{mol/kg}$ in the PeP extracts.

354

355 3.2. Sensory evaluation

356 3.2.1. UG extract solutions

357 The phenol concentration of the UG solutions significantly affected the intensity of the target
358 sensations (**Fig. 3A and Table S2**). According to the F values, the increase in phenol concentration
359 had the strongest effect on sourness while it influenced the other target sensations much less.
360 Significant intensity increases were observed in the samples with phenols from the UG extract

361 compared to the sample without added phenol (0.00 g/L). Sourness increased from weak to strong
362 across the phenol concentration range. Bitterness, astringency and saltiness showed limited intensity
363 increases, from barely detectable to weak.

364 Four concentration levels, which cover the whole range of significant variations of intensity of
365 target sensations, were selected to fortify the vegetable matrices: 0.00, 0.21, 0.41, 1.11 and 1.93
366 g/L.

367 3.2.2. *Functionalized foods*

368 The intensity of target sensations significantly changed in all of the three vegetable prototypes as a
369 function of the increasing phenol concentrations, the only exception being sweetness in the PoP
370 (**Table 2**). Phenol concentration induced the strongest effect on sourness in all of the three food
371 models as showed by F-values. The intensity of the other sensations was influenced by both the
372 increase in phenol concentration and, to a lesser extent, by the macro-composition of the matrix. All
373 of the sensations were barely detectable in the beetroot purée sample without added phenol, while in
374 the rest of the samples, sourness increased from weak to strong, sweetness showed a significant
375 decrease from moderate to weak, while saltiness, astringency and bitterness increased slightly from
376 barely detectable to weak (**Fig. 3 B-Beetroot purée**). The variation in intensity of the target
377 sensation in the pea purée as a function of the phenol concentration was similar to that observed in
378 the beetroot purée (**Fig. 3 C-Pea purée**). The increase in sourness from barely detectable to
379 moderate was associated with a significant decrease in sweetness, from moderate to weak, while the
380 rest of the sensations were perceived at a weak intensity or even lower. In the potato purée sample
381 without added phenols, all the sensations were rated at a barely detectable/weak intensity, while
382 only sourness showed a remarkable increase from barely detectable to strong as the phenol
383 concentration increased (**Fig. 3 D-Potato purée**).

384 Bitterness, astringency and saltiness were not further investigated since these sensations were
385 marginally affected by addition of phenols and perceived at a weak intensity across the whole range
386 of concentrations.

387

388 Sourness and sweetness perceived in the food functionalized at different UG concentration were
389 compared to further explore the effect of food macro-composition on UG phenol sensory properties.
390 While the vegetable matrix and phenol concentration significantly affected the intensity of sourness
391 and sweetness, the vegetable matrix*concentration interaction was never significant (**Table S3**).
392 Significant differences were found upon comparing sourness from the three matrices at phenol
393 concentrations of 0.41, 1.11 and 1.93 g/L. The highest sourness intensity was rated in the PoP,
394 whereas no significant differences were found between the BP and PeP (**Fig. 4-A**). Sweetness was
395 rated as more intense in the BP and PeP than in the PoP across the 0.0 to 0.41 g/kg concentration
396 range of spiked phenols. At the highest concentration levels, sweetness was perceived at the highest
397 intensity in the BP (**Fig. 4-B**).

398

399 **4. Discussion**

400

401 Physical-chemical characterization was carried out to evaluate the attitude of UG extract towards
402 rehydration and stability during storage, in terms of phenolic content and antioxidant activity. The
403 solubility value of the UG extract was similar to those (86% - 88%) obtained by Kuck & Noreña
404 (2016) on grape skin extracts lyophilized with arabic gum and partially hydrolysed guar gum as
405 supports.

406

407 The moisture content and water activity value of the UG extract were in agreement with the results
408 obtained on grape skin extracts by Kuck & Noreña (2016). The UG extract showed similar
409 hygroscopic behaviour to the absorption isotherm of an aqueous solution of salts and simple sugars.
410 Therefore, the powder has to be protected from humidity during storage to avoid water absorption,
411 thus preserving the extract's stability.

412

413 The total phenol content of the UG extract was similar to that obtained by Kuck & Noreña (2016)
414 on aqueous extracts of grape skin microencapsulated with different agents while the antioxidant
415 activity was slightly lower. In general, the phenol content and antioxidant activity of extracts vary
416 mainly depending on the origin of grape by-products and extraction conditions (Trigo, Alexandre,
417 Saraiva, & Pintado, 2019). Indeed, when ethanol or methanol were used for the extraction, the
418 phenolic content and antioxidant activity of the extracts were higher than those detected in aqueous
419 extracts (Trigo et al., 2019; Tournour, Segundo, Magalhães, Costa & Cunha, 2017). After nine
420 months, the high percentage of both residual phenols and antioxidant activity in the UG extract
421 indicated that the adopted storage conditions were suitable to protect the UG phenols from
422 degradation.

423

424 When a different amount of the UG phenols was used to enrich the food models, the increase of
425 phenol concentration in the FM extracts was expected. Similar results were obtained by other
426 authors who studied the addition of phenolic extracts from different by-products to some food and
427 beverages (Trigo et al., 2019). Chemical-physical characteristics of food models explored in these
428 study significantly affect phenol recovery thus indicating clear reactivity differences between UG
429 phenols and food components. The lowest amount of phenols was recovered from the protein-rich
430 model (pea purée). A similar effect of the interaction phenol/biopolymers on the bioactivity of
431 phenols from olive mill waste waters in plant-based food has already been observed by other
432 authors (De Toffoli et al., 2019).

433 The formation of phenol/protein aggregates significantly lowers the phenol bio-activity both in
434 terms of extractability from raw material and antioxidant activity (Ozdal et al., 2013). Proteins bind
435 plant polyphenols through hydrophobic and hydrogen interactions; the preferred sites of interaction
436 plant phenol/food protein in *in vitro* conditions are the proline-rich regions of leguminous proteins
437 characterized by high basic-residue contents as well as open and flexible structures (Kroll et al.,
438 2003; Zhang et al., 2014).

439 Phenol chemical structure, size and composition, including number of OH groups, play an
440 important role in phenol/protein interactions, and phenolic compounds with a low molecular weight
441 are inefficient to bond proteins (de Freitas & Mateus, 2012). It is known that upon extraction, the
442 acidic condition of grape juice promotes the depolymerization of proanthocyanidins (Vidal,
443 Cartalade, Souquet, Fulcrand, & Cheynier, 2002). However, these reactions begin during
444 maceration and proceed slowly in wine, but they have never been highlighted in grape juice.
445

446 The quite high percentages of UG phenols recovered, mainly in the carbohydrate-rich potato and
447 beetroot purée food models, indicated that moderate/weak chemical interactions take place among
448 UG phenols and food components. These findings, associated with the significant increase in
449 antioxidant activity detected in the functionalized food models after the addition of UG phenols,
450 indicate that most of the potential biological activity and the extractability of UG phenols were
451 maintained after blending.

452 Phenolic compounds can bridge or cross-link with polysaccharides, and a large fraction of the not
453 extractable polyphenols consist phenol associated with polysaccharides (Pérez-Jiménez, Díaz-
454 Rubio, & Saura-Calixto, 2013). The consequences of phenol/carbohydrate interactions on phenol
455 biological activity depends on the chemical characteristics of both phenols and carbohydrates
456 (Zhang et al., 2014).

457 Other authors have described a competition between the arabic gum and other carbohydrates and
458 the proteins to bind to the tannin (Gonçalves, Mateus, & de Freitas, 2011). The mechanism was
459 previously investigated by tasting the influence of several carbohydrates on the formation of
460 polyphenols/protein complexes. Polygalacturonic acid, arabic gum and pectin prevented the
461 association of procyanidin B3 with trypsin, and that of salivary proteins with grape seed
462 procyanidins. The interruption of polyphenol-protein association by carbohydrates can prevent
463 some of the negative effects of these complexes, such as enzyme activity inhibition, and it can
464 influence the perceived astringency of some food products.

465

466 The antioxidant activity of UG phenols was influenced by the food composition. The highest level
467 of antioxidant activity was found in the carbohydrate-rich/acidic pH beetroot purée. The antiradical
468 capacity of phenols depends on several factors such as their concentration and structures, and the
469 physical-chemical characteristics of the solvent. The role of acidity in the kinetics of phenol/radical
470 reactions was previously investigated by (Musialik, Kuzmich, Pawcowski, & Litwinienko, 2009). In
471 general, it is known that deprotonated flavonoids are more potent electron donors and are better
472 radical scavengers than neutral molecules. However, the ability of phenols to scavenge reactive
473 oxygen species such as peroxy and hydroxyl radicals is still far from being fully understood.
474 Valgimigli et al. (2009) described an unexpected dramatic acceleration of phenol-peroxy radical
475 reaction with the addition of acid. The best performance, in terms of antioxidant activity, of UG
476 phenols when added to beetroot purée could be due to the acidic pH of the beetroot food model.

477

478 Sensory profiles of the three matrices were significantly affected by the addition of UG extracts.
479 Sourness intensity increased as a function of the UG phenol concentration. The natural sweetness
480 of the beetroot and pea purées was reduced by the spiked phenols due to the intermodal interaction
481 between sour and bitter tastes, which induced the suppression of perceived sweetness as the
482 sourness intensity increased (Keast & Breslin, 2002). The bitterness, saltiness and astringency
483 intensities were significantly modified by the UG extract, but the extent of these effects appears
484 marginal since these sensations are perceived at a weak intensity across the whole range of
485 concentrations.

486 The different compositions of the vegetable matrices affect the UG phenols' contribution to
487 sourness. Furthermore, the observed increasing intensity range differed across the series of samples
488 indicating that their macro-component plays an active role in modulating the sensory impact of UG
489 phenols.

490

491 5. Conclusions

492

493 An extract from unripe grapes showed suitable physical-chemical characteristics for its inclusion in
494 plant-based foods. Food composition influenced the functional and sensory properties of phenols
495 from unripe grapes. The strongest effect in terms of recovered phenol and antioxidant activity was
496 observed in protein-based food. The use of matrices high in carbohydrates, with acidic pH and
497 characterized by sweet taste appears a suitable strategy to counteract the impact of the negative
498 sensory properties of added phenol on plant-based food. The use of phenolic extracts from unripe
499 grapes can be useful to improve potential health benefits when formulating plant-based functional
500 food.

501

502 Acknowledgements

503 This research was support by the Ministero dell'Istruzione, dell' Università e della Ricerca (MIUR),
504 ITALY - Research Project : 20158YJW3W Programmi di Ricerca Scientifica di Rilevante Interesse
505 Nazionale – PRIN 2015: “Individual differences in the acceptability of health foods: focus on
506 phenol and fat content”.

507507

508 References

509

- 510 Adams, D. O. (2006). Phenolics and ripening in grape berries. *American Journal of Enology and*
511 *Viticulture*, 57(3), 249–256.
- 512 A.O.A.C., 1990. Official Methods of Analysis, 14th ed. Association of Official Analytical
513 Chemists, Washington, D.C. Bhandari.
- 514 Beres, C., Costa, G. N. S., Cabezudo, I., da Silva-James, N. K., Teles, A. S. C., Cruz, A. P. G.,
515 Freitas, S. P. (2017). Towards integral utilization of grape pomace from winemaking process:
516 A review. *Waste Management*, 68, 581–594. <https://doi.org/10.1016/J.WASMAN.2017.07.017>
- 517 Brand-Williams, W., Cuvelier, M. E., & Berset, C. L. W. T. (1995). Use of free radical method to
518 evaluate antioxidant activity. *LWT - Food Science and Technology*, 28, 25–30.
519 [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)

520 Callahan, J. C., Cleary, G. W., Elefant, M., Kaplan, G., Kensler, T., & Nash, R. A. (1982).
 521 Equilibrium Moisture Content of Pharmaceutical Excipients. *Drug Development and*
 522 *Industrial Pharmacy*, 8(3), 355–369. <https://doi.org/10.3109/03639048209022105>

523 Cano-Chauca, M., Stringheta, P. C., Ramos, A. M., & Cal-Vidal, J. (2005). Effect of the carriers on
 524 the microstructure of mango powder obtained by spray drying and its functional
 525 characterization. *Innovative Food Science and Emerging Technologies*, 6(4), 420–428.
 526 <https://doi.org/10.1016/j.ifset.2005.05.003>

527 de Freitas, V., & Mateus, N. (2012). Protein/Polyphenol Interactions: Past and Present
 528 Contributions. Mechanisms of Astringency Perception. *Current Organic Chemistry*, 16(6),
 529 724-746. <https://doi.org/http://dx.doi.org/10.2174/138527212799958002>

530 De Toffoli, A., Monteleone, E., Bucalossi, G., Veneziani, G., Fia, G., Servili, M., Zanoni, B.,
 531 Pagliarini, E., Gallina Toschi, T., & Dinnella, C. (2019). Sensory and chemical profile of a
 532 phenolic extract from olive mill waste waters in plant-based food with varied macro-
 533 composition. *Food Research International*, 119(October 2018), 236–243.
 534 <https://doi.org/10.1016/j.foodres.2019.02.005>

535 Dinnella, C., Recchia, A., Fia, G., Bertuccioli, M., & Monteleone, E. (2009). Saliva Characteristics
 536 and Individual Sensitivity to Phenolic Astringent Stimuli. *Chemical Senses*, 34, 295–304.
 537 <https://doi.org/10.1093/chemse/bjp003>

538 Dinnella, C., Recchia, A., Tuorila, H., & Monteleone, E. (2011). Individual astringency
 539 responsiveness affects the acceptance of phenol-rich foods. *Appetite*, 56(3), 633–642.
 540 <https://doi.org/10.1016/j.appet.2011.02.017>

541 Dupas de Matos, A., Magli, M., Marangon, M., Curioni, A., Pasini, G., & Vincenzi, S. (2018). Use
 542 of verjuice as an acidic salad seasoning ingredient: evaluation by consumers' liking and Check-
 543 All-That-Apply. *European Food Research and Technology*, 244, 2117-2125.

544 Europe, I. (1999). Scientific Concepts of Functional Foods in Europe Consensus Document. *British*
 545 *Journal of Nutrition*, 81, S1-S27.

546 Fia, G., Dinnella, C., Bertuccioli, M., & Monteleone, E. (2009). Prediction of grape polyphenol
 547 astringency by means of a fluorimetric micro-plate assay. *Food Chemistry*, 113(1), 325–330.
 548 <https://doi.org/10.1016/j.foodchem.2008.07.058>

549 Fia, G., Gori, C., Bucalossi, G., Borghini, F., & Zanoni, B. (2018). A naturally occurring
 550 antioxidant complex from unripe grapes: The case of sangiovese (v. *Vitis vinifera*).
 551 *Antioxidants*, 7(2). <https://doi.org/10.3390/antiox7020027>

552 Fia, G., & Gori, C. (2016) Process for the extraction of antioxidants from plant matrices (Italian
 553 Patent number 102016000022015).

554 Gatti, M., Bernizzoni, F., Civardi, S., & Poni, S. (2012). Effects of cluster thinning and

555 preflowering leaf removal on growth and grape composition in cv. Sangiovese. *American*
556 *Journal of Enology and Viticulture*, 63(3), 325–332. <https://doi.org/10.5344/ajev.2012.11118>

557 Gonçalves, R., Mateus, N., & de Freitas, V. (2011). Inhibition of α -amylase activity by condensed
558 tannins. *Food Chemistry*, 125(2), 665–672. <https://doi.org/10.1016/j.foodchem.2010.09.061>

559 Green, B. G., Higgins, J., Cowart, B., Rankin, K., Dalton, P., & Shaffer, G. (2007). Evaluating the
560 ‘Labeled Magnitude Scale’ for Measuring Sensations of Taste and Smell. *Chemical Senses*,
561 21(3), 323–334. <https://doi.org/10.1093/chemse/21.3.323>

562 Gori, C., Menichetti, S., & Fia, G. (2014). Multi-functional oenological machine and use in the
563 oenological production chain (European Patent number 2957627).

564 Guilford, J. M., & Pezzuto, J. M. (2011). Wine and health: A review. *American Journal of Enology*
565 *and Viticulture*, 62(4), 471–486. <https://doi.org/10.5344/ajev.2011.11013>

566 Henry, C. J. (2010). Functional foods. *European Journal of Clinical Nutrition*, 64, 657–659.
567 <https://doi.org/10.1038/ejcn.2010.101>

568 Hufnagel, J. C., & Hofmann, T. (2008). Quantitative Reconstruction of the Nonvolatile
569 Sensometabolome of a Red Wine. *Journal of Agricultural and Food Chemistry*, 56, 9190–
570 9199.

571 Iwatani, S., & Yamamoto, N. (2019). Functional food products in Japan : A review. *Food Science*
572 *and Human Wellness*. 8, 96–101.

573 Jakobek, L. (2015). Interactions of polyphenols with carbohydrates, lipids and proteins. *Food*
574 *Chemistry*, 175, 556–567. <https://doi.org/10.1016/j.foodchem.2014.12.013>

575 Keller, M., Mills, L. J., Wample, R. L., & Spayd, S. E. (2005). Cluster thinning effects on three
576 deficit-irrigated *Vitis vinifera* cultivars. *American Journal of Enology and Viticulture*, 56(2),
577 91–103. <https://doi.org/10.1071/CH9540055>

578 Kroll, J., Rawel, H. M., & Rohn, S. (2007). Reactions of Plant Phenolics with Food Proteins and
579 Enzymes under Special Consideration of Covalent Bonds. *Food Science and Technology*
580 *Research*, 9(3), 205–218. <https://doi.org/10.3136/fstr.9.205>

581 Kuck, L. S., & Noreña, C. P. Z. (2016). Microencapsulation of grape (*Vitis labrusca* var. Bordo)
582 skin phenolic extract using gum arabic, polydextrose, and partially hydrolyzed guar gum as
583 encapsulating agents. *Food Chemistry*, 194, 569–576.
584 <https://doi.org/10.1016/j.foodchem.2015.08.066>

585 Manach, C., Scalbert, A., Morand, C., Rémésy, C., & Jiménez, L. (2004). Polyphenols: food
586 sources and bioavailability. *American Journal of Clinical Nutrition*, 79, 727-747.

587 Mihaylova, D., Popova, A., Alexieva, I., Krastanov, A., & Lante, A. (2018). Polyphenols as suitable
588 control for obesity and diabetes. *The Open Biotechnology Journal*, 12, 2019-228.

589 Monteleone, E., Spinelli, S., Dinnella, C., Endrizzi, I., Laureati, M., Pagliarini, E., ... Tesini, F.

590 (2017). Exploring influences on food choice in a large population sample: The Italian Taste
 591 project. *Food Quality and Preference*, 59, 123–140.
 592 <https://doi.org/10.1016/j.foodqual.2017.02.013>

593 Musialik, M., Kuzmich, R., Pawcowski, T. S., & Litwinienko, G. (2009). Acidity of Hydroxyl Grps:
 594 Overlooked Influence on Antioxidant Properties of Flavonoid. *Journal of Organic Chemistry*,
 595 74(7), 2699–2709. <https://doi.org/10.1021/jo802716v>

596 Nirmala, C., Bisht, M. S., Bajwa, H. K., & Santosh, O. (2018). Bamboo: A rich source of natural
 597 antioxidants and its applications in the food and pharmaceutical industry. *Trends in Food*
 598 *Science and Technology*, 77, 91–99. <https://doi.org/10.1016/j.tifs.2018.05.003>

599 Olivero-David, R., Ruiz-Roso, M., Caporaso, N., Perez-Olleros, L., De las Heras, N., Lahera, V., &
 600 Ruiz-Roso, B. (2018). In vivo bioavailability of polyphenols from grape by-product extracts,
 601 and effect on lipemia of normocholesterolemic Wistar rats. *Journal of the Science of Food and*
 602 *Agriculture*, 98, 5581–5590.

603 Öncül, N., & Karabiyikli, Ş. (2015). Factors Affecting the Quality Attributes of Unripe Grape
 604 Functional Food Products. *Journal of Food Biochemistry*, 39(6), 689–695.
 605 <https://doi.org/10.1111/jfbc.12175>

606 Ough C. S., Nagaoka, R. (1984). Effect of Cluster Thinning and Vineyard Yields on Grape and
 607 Wine Composition and Wine Quality of Cabernet Sauvignon. *American Journal of Enology*
 608 *and Viticulture*, 35(1), 30–34.

609 Ozdal, T., Capanoglu, E., & Altay, F. (2013). A review on protein-phenolic interactions and
 610 associated changes. *Food Research International*, 51(2), 954–970.
 611 <https://doi.org/10.1016/j.foodres.2013.02.009>

612 Peleg, H., Gacon, K., Schlich, P., & Noble, A. C. (1999). *Bitterness and astringency of flavan-3-ol*
 613 *monomers, dimers and trimers*. 1128, 1123–1128.

614 Pérez-Jiménez, J., Díaz-Rubio, M. E., & Saura-Calixto, F. (2013). Non-extractable polyphenols, a
 615 major dietary antioxidant: Occurrence, metabolic fate and health effects. *Nutrition Research*
 616 *Reviews*, 26(2), 118–129. <https://doi.org/10.1017/S0954422413000097>

617 Rasines-Perea, Z., & Teissedre, P. L. (2017). Grape Polyphenols' effects in human cardiovascular
 618 diseases and diabetes. *Molecules*, 22(1), 1–19. <https://doi.org/10.3390/molecules22010068>

619 Sengul, H., Surek, E., & Nilufer-Erdil, D. (2014). Investigating the effect of food matrix and food
 620 component on bioaccessibility of pomegranate (*Punica granatum*) phenolics and anthocyanins
 621 using an *in-vitro* gastrointestinal digestion model. *Food Research International*, 62, 1069–
 622 1079.

623 Shahidi, F., & Ambigaipalan, P. (2015). Phenolics and polyphenolics in foods, beverages and
 624 spices: Antioxidant activity and health effects. *Journal of Functional Foods*, 18, 820–897.

<https://doi.org/10.1016/j.jff.2015.06.018>
 Singleton, V. L., Rossi Jr., J. A., & Rossi J A Jr. (1965). Colorimetry of Total Phenolics with
 Phosphomolybdic-Phosphotungstic Acid Reagents. *American Journal of Enology and
 Viticulture*, 16(3), 144–158. <https://doi.org/10.12691/ijebb-2-1-5>
 Sengul, H., Surek, E., & Nilufer-erdil, D. (2014). Investigating the effects of food matrix and food
 components on bioaccessibility of pomegranate (*Punica granatum*) phenolics and
 anthocyanins using an in-vitro gastrointestinal digestion model. *Food Research International*,
 62, 1069–1079. <https://doi.org/10.1016/j.foodres.2014.05.055>
 Świeca, M., Gawlik-Dziki, U., Sęczyk, Ł., Dziki, D., & Sikora, M. (2018). Interactions of green
 coffee bean phenolics with wheat bread matrix in a model of simulated in vitro digestion. *Food
 Chemistry*, 258(March), 301–307. <https://doi.org/10.1016/j.foodchem.2018.03.081>
 Tinello, F., & Lante, A. (2017). Evaluation of antibrowning and antioxidant activities in unripe
 grapes recovered during bunch thinning. *Australian Journal of Grape and Wine Research*,
 23(1), 33–41. <https://doi.org/10.1111/ajgw.12256>
 Trigo, J. P., Alexandre, E. M. C., Saraiva, J. A., & Pintado M. E. (2019). High value-added
 compounds from fruit and vegetable by-products – Characterization , bioactivities , and
 application in the development of novel food products food products. *Critical Reviews in Food
 Science and Nutrition*, 0(0), 1–29. <https://doi.org/10.1080/10408398.2019.1572588>
 Torri, L., Piochi, M., Marchiani, R., Zeppa, G., Dinnella, C., & Monteleone, E. (2015). A sensory-
 and consumer-based approach to optimize cheese enrichment with grape skin powders. *Journal
 of Dairy Science*, 99(1), 194–204. <https://doi.org/10.3168/jds.2015-9922>
 Tournour, H., Segundo, M. A., Magalhães, L. M. Costa, A. S. G., & Cunha, L. M. (2017). Effect of
 Touriga nacional grape extract on characteristics of mechanically deboned chicken meat kept
 under frozen storage. *Journal of Food Process Engineering*, 40, 1-10.
<https://doi.org/10.1111/jfpe.12434>
 Turkmen, N., Sari, F., & Velioglu, Y. S. (2005). The effect of cooking methods on total phenolics
 and antioxidant activity of selected green vegetables. *Food Chemistry*, 93(4), 713-718.
<https://doi.org/10.1016/j.foodchem.2004.12.038>
 Valgimigli, L., Amorati, R., Petrucci, S., Pedulli, G. F., Hu, D., Hanthorn, J. J., & Pratt, D. A.
 (2009). Unexpected Acid Catalysis in Reactions of Peroxyl Radicals with Phenols.
Angewandte Chemie International Edition, 48(44), 8348–8351.
<https://doi.org/10.1002/anie.200903360>
 Vidal, S., Cartalade, D., Souquet, J.-M., Fulcrand, H., & Cheynier, V. (2002). Changes in
 Proanthocyanidin Chain Length in Winelike Model Solutions. *Journal of Agricultural and
 Food Chemistry*, 50(8), 2261–2266. <https://doi.org/10.1021/jf011180e>

Villaño, D., Fernández-Pachón, M. S., Moyá, M. L., Troncoso, A. M., & García-Parrilla, M. C. (2007). Radical scavenging ability of polyphenolic compounds towards DPPH free radical. *Talanta*, 71(1), 230–235. <https://doi.org/10.1016/j.talanta.2006.03.050>

Yu, J., & Ahmedna, M. (2013). Functional components of grape pomace: Their composition, biological properties and potential applications. *International Journal of Food Science and Technology*, 48(2), 221–237. <https://doi.org/10.1111/j.1365-2621.2012.03197.x>

Zhang, H., Yu, D., Sun, J., Liu, X., Jiang, L., Guo, H., & Ren, F. (2014). Interaction of plant phenols with food macronutrients: characterisation and nutritional–physiological consequences. *Nutrition Research Reviews*, 27(01), 1–15. <https://doi.org/10.1017/s095442241300019x>

Figure legend

Figure 1. Total phenols (A) of food models, mean values of UG phenols recovered (B) from beetroot purée (BP), pea purée (PeP) and potato purée (PoP) functionalized with increasing amounts (0.00, 0.21, 0.44, 1.11 and 1.93 g/kg of food) of phenols and percentage of each phenolic class (C) detected in the UG extract (UG ext) and food models functionalized with 1.93 g/kg phenols from UG extract. The bars represent standard deviation. Different letters represent significant different values ($p \leq 0.001$).

Figure 2. Antioxidant activity (A) of beetroot purée (BP), pea purée (PeP) and potato purée (PoP) functionalized with increasing amounts of phenols (0, 0.21, 0.44, 1.11 and 1.93 g/kg of food) from UG extract and antioxidant activity (B) of UG phenols in water solution vs antioxidant activity in the FM extracts. The bars represent standard deviation. Different letters represent significant different values ($p \leq 0.001$).

Figure 3. Mean intensity of target sensations (A) in the UG solutions with increasing phenol concentration and food models (B, C and D) functionalized with increasing concentrations of phenols from UG extract. The bars represent standard error.

689 **Figure 4.** Effect of the vegetable matrix on the perceived intensity of sourness (A) and sweetness
690 (B) in foods spiked with different concentrations of phenols from UG extract. Different letters
691 represent significant different values ($p \leq 0.038$).
692

Table 1. Phenol profile of the UG extract and phenols detected in the FM extracts. Beetroot purée (BP), pea purée (PeP) and potato purée (PoP) functionalized with 1.93 g/kg of phenols from the UG extract.

Compound	mg/kg			
	UG extract	BP*	PeP*	PoP*
<i>Phenolic acid</i>				
Caffeic acid	11.0 ± 0.4	1.04 ± 0.07 ^c	1.55 ± 0.14 ^a	1.28 ± 0.14 ^b
Caftaric acid	704 ± 33	48.7 ± 1.2 ^a	35.7 ± 6.5 ^b	36.5 ± 4.0 ^b
Coumaric acid	19.6 ± 0.6	1.80 ± 0.13 ^b	2.30 ± 0.12 ^a	1.79 ± 0.14 ^b
Coutaric acid	34.3 ± 1.1	2.31 ± 0.17 ^a	2.03 ± 0.18 ^{ab}	1.81 ± 0.15 ^b
Ferulic acid	4.63 ± 0.59	2.51 ± 0.04 ^a	0.44 ± 0.03 ^b	nd
Fertaric acid	52.0 ± 2.0	3.44 ± 0.10 ^a	3.54 ± 0.28 ^a	3.71 ± 0.19 ^a
Gallic acid	1.63 ± 0.03	0.03 ± 0.01 ^b	0.24 ± 0.02 ^a	0.05 ± 0.01 ^b
<i>Flavonols</i>				
Isorhamnetin	1.41 ± 0.03	0.05 ± 0.01 ^b	0.09 ± 0.01 ^a	0.06 ± 0.02 ^b
Kaempferol	0.78 ± 0.04	0.06 ± 0.01 ^a	0.06 ± 0.01 ^a	0.0 ± 0.01 ^a
Kaempferol-3- <i>O</i> -glucoside	0.54 ± 0.03	nd	nd	nd
Myricetin	3.79 ± 0.11	0.39 ± 0.03 ^b	0.47 ± 0.04 ^a	0.45 ± 0.03 ^{ab}
Quercetin	14.0 ± 0.4	1.26 ± 0.11 ^b	1.48 ± 0.13 ^{ab}	1.57 ± 0.14 ^a
Quercetin-3- <i>O</i> -hexoside	1.32 ± 0.08	nd	nd	nd
<i>Flavan-3-ols</i>				
(+)-Catechin	13.6 ± 0.8	1.23 ± 0.07 ^c	2.28 ± 0.12 ^a	1.51 ± 0.11 ^b
(-)-Epicatechin	8.23 ± 0.29	0.70 ± 0.03 ^c	1.09 ± 0.08 ^a	0.83 ± 0.05 ^b
<i>Procyanidins</i>				
Procyanidin B1	4.55 ± 0.19	0.44 ± 0.04 ^b	0.56 ± 0.04 ^a	0.47 ± 0.06 ^{ab}
Procyanidin B2	9.74 ± 0.37	1.13 ± 0.05 ^c	1.66 ± 0.05 ^a	1.33 ± 0.07 ^b
<i>Stilbenes</i>				
Trans-resveratrol	31.3 ± 1.6	2.18 ± 0.13 ^b	3.33 ± 0.48 ^a	2.36 ± 0.36 ^b
2- <i>S</i> -Glutathionyl caftaric acid	16.8 ± 0.6	nd	nd	nd

Data are expressed as mean ± standard deviation (n=3); nd, not detected. Different letters represent significant different values ($p \leq 0.001$) among the columns.

Table 2. Two-way ANOVA mixed model (random effect: assessors): phenol concentration effect on intensity of target sensations in food models. Mean, F and p values.

			Concentration of phenols from UG (g/kg)				
			0.00	0.21	0.41	1.11	1.93
	F	p					
Bitterness							
Beetroot Purée	4.92	0.0011	0.97 b	1.34 b	0.62 b	1.34 b	3.31 a
Pea Purée	6.78	< 0.0001	1.28 b	1.31 b	1.41 b	3.72 a	5.28 a
Potato Purée	2.53	0.0445	2.61 b	3.00 b	3.25 b	4.11 ab	5.46 a
Sourness							
Beetroot Purée	26.22	< 0.0001	2.38 c	3.07 c	4.41 c	13.86 b	21.86 a
Pea Purée	39.02	< 0.0001	3.48 b	3.34 b	5.62 b	16.31 a	19.72 a
Potato Purée	48.39	< 0.0001	3.07 e	8.54 d	13.46 c	20.43 b	27.68 a
Saltiness							
Beetroot Purée	4.85	0.0012	1.17 b	1.38 b	2.38 b	2.86 ab	4.55 a
Pea Purée	3.63	0.0081	4.52 c	4.31 c	5.79 bc	7.24 ab	8.55 a
Potato Purée	5.78	0.0003	2.29 bc	1.96 c	3.89 bc	4.00 b	6.14 a
Sweetness							
Beetroot Purée	3.07	0.0194	16.31 a	17.79 a	15.21 ab	13.83 ab	11.28 b
Pea Purée	10.01	< 0.0001	12.72 a	13.69 a	11.41 a	7.31 b	5.52 b
Potato Purée	1.56	0.1865	4.18	3.21	3.43	2.36	2.54
Astringency							
Beetroot Purée	4.64	0.0017	4.31 bc	4.07 c	31 c	7.38 a	6.34 ab
Pea Purée	4.16	0.0035	5.48 bc	3.72 c	3.97 bc	6.76 ab	8.72 a
Potato Purée	6.01	0.0001	2.86 c	4.93 bc	6.86 ab	7.64 a	8.43 a

Different letters indicate significantly different values ($p \leq 0.05$).

Highlights

- A strategy was outlined for the exploitation of high-quality unripe grapes
- The food composition affected both the phenol recovered and antioxidant activity
- The highest recovery of phenols was from the starch/neutral pH food model
- The highest antioxidant activity was from the carbohydrates/acidic pH food model
- The models' sensory properties are modulated by phenol content and food composition

715
716
717
718
719
720
721
722

723
724
725
726
727

728
729
730
731
732
733
734
735
736
737
738
739

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Ginevra Bucalossi: Investigation, Visualization
Giovanna Fia: Writing- Original draft preparation
Caterina Dinnella: Conceptualization, Methodology, Writing - Review & Editing
Erminio Monteleone : Conceptualization, Methodology, Funding acquisition
Alessandra De Toffoli: Investigation, Visualization
Valentina Canutia: Investigation
Bruno Zanoni: Writing - Review & Editing
Maurizio Servili: Investigation
Ella Pagliarini: Investigation
Tullia Gallina Toschi: Investigation