

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Molecular and biological characterization of phytoplasmas from coconut palms affected by the lethal yellowing disease in Africa

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Contaldo, N., D'Amico, G., Paltrinieri, S., Diallo, H.A., Bertaccini, A., Arocha Rosete, Y. (2019). Molecular and biological characterization of phytoplasmas from coconut palms affected by the lethal yellowing disease in Africa. MICROBIOLOGICAL RESEARCH, 223-225(June-August 2019), 51-57 [10.1016/j.micres.2019.03.011].

Availability:

This version is available at: <https://hdl.handle.net/11585/686048> since: 2019-05-06

Published:

DOI: <http://doi.org/10.1016/j.micres.2019.03.011>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Accepted Manuscript

Title: Molecular and biological characterization of
phytoplasmas from coconut palms affected by the lethal
yellowing disease in Africa

Authors: Nicoletta Contaldo, Gianfranco D'Amico, Samanta
Paltrinieri, Hortense A. Diallo, Assunta Bertaccini, Yaima
Arocha Rosete

PII: S0944-5013(19)30220-4
DOI: <https://doi.org/10.1016/j.micres.2019.03.011>
Reference: MICRES 26275

To appear in:

Received date: 21 February 2019
Revised date: 26 March 2019
Accepted date: 29 March 2019

Please cite this article as: Contaldo N, D'Amico G, Paltrinieri S, Diallo HA, Bertaccini A, Arocha Rosete Y, Molecular and biological characterization of phytoplasmas from coconut palms affected by the lethal yellowing disease in Africa, *Microbiological Research* (2019), <https://doi.org/10.1016/j.micres.2019.03.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Molecular and biological characterization of phytoplasmas from coconut palms affected by the lethal yellowing disease in Africa

Nicoletta Contaldo^{a*}, Gianfranco D'Amico^a, Samanta Paltrinieri^a, Hortense A. Diallo^b, Assunta Bertaccini^a, Yaima Arocha Rosete^c

^aDepartment of Agricultural and Food Sciences, Plant Pathology, *Alma Mater Studiorum*, University of Bologna, viale G. Fanin, 40, 40127 Bologna, Italy

^bUniversité Nangui Abrogoua, 02 BP 801 Abidjan 02, Côte d'Ivoire

^cSporometrics, 219 Dufferin Street, Suite 20C, Toronto, ON M6K 3J1, Canada

Corresponding author

Nicoletta Contaldo

DISTAL, viale G. Fanin, 40, 40127 Bologna, Italy

Phone +39-0512096723

Email address: nicoletta.contaldo2@unibo.it

Research Highlights

- Phytoplasma identification in coconut
- Phytoplasma isolation in media
- Phytoplasma identification in solid/liquid media
- Biochemical characterization of two phytoplasma isolates
- Antibiotic susceptibility of two phytoplasma isolates

Abstract

Côte d'Ivoire lethal yellowing (CILY) is a devastating disease associated with phytoplasmas and has recently rapidly spread to several coconut-growing areas in the Country. Phytoplasmas are phloem-restricted bacteria that affect plant species worldwide. These bacteria are transmitted by plant sap-feeding insects, and their cultivation was recently achieved in complex artificial media. In this study, phytoplasmas were isolated for the first time from coconut palm trunk borings in ~~both~~ solid and liquid media from both CILY symptom-bearing and symptomless coconut palms. The colony morphology, PCR and sequencing analyses indicated the presence of phytoplasmas ~~from~~ different ribosomal groups. This study reports the first biochemical characterization of two of these phytoplasma isolates. Moreover, a disc-diffusion antibiotic susceptibility assay revealed that these bacteria exhibit tobramycin susceptibility and cephalexin hydrate and rifampicin resistance. Urea and arginine hydrolysis, and glucose fermentation tests that were performed on colonies of phytoplasmas and *Acholeplasma laidlawii* indicated that both phytoplasmas tested were negative for urea and positive for glucose and arginine, whereas *A. laidlawii* was positive for glucose and negative for urea and arginine. The growth of coconut phytoplasma in both solid and liquid artificial media and the biological characterization of these isolates are novel and important advancements in the field of disease management and containment measures for the CILY disease. The characterization of isolated phytoplasmas will allow for more efficient management strategies in both the prevention of a coconut phytoplasma epidemic and the reduction of the economic impact of the disease in the affected areas.

Abbreviations:

‘Ca. P.’: ‘*Candidatus* Phytoplasma’

CFU: colony forming unit

CILY: Côte d'Ivoire lethal yellowing

CSPWD: Cape St. Paul Wilt Disease

CTAB: hexadecyltrimethylammonium bromide

DDSW: distilled, deionize, sterile water

DNA: deoxyribonucleic acid

dNTP: deoxynucleotide triphosphate

ICSB: International Committee for Systematic Bacteriology

LY: Lethal yellowing

MIC: minimal inhibitory concentration

PCR: polymerase chain reaction

rDNA: ribosomal DNA

RFLP: restriction fragment length polymorphism

RNA: ribonucleic acid

rRNA: ribosomal RNA

UK: United Kingdom

UV: ultra violet

Keywords

Coconut lethal yellowing disease; antibiotics susceptibility; molecular detection; metabolism; microbial phylogenetics.

Introduction

Phytoplasmas are phloem-restricted bacteria from the class *Mollicutes* that affect more than one thousand crop and plant species worldwide; these bacteria are transmitted by plant sap-feeding insect species (Weintraub and Beanland, 2006). Since their discovery (Doi et al., 1967) and until recently, phytoplasmas were long reported to be unculturable in cell-free media; thus, ~~phytoplasmas have been well~~ characterized only from the genetic perspective. However, the cultivation of phytoplasma was recently achieved using micro-propagated periwinkle plants as a source of phytoplasma-infected shoots (Contaldo et al., 2012). Later, phytoplasma colonies from naturally phytoplasma-infected grapevine plants were isolated in artificial media ~~using a complex medium~~ (Contaldo et al., 2016).

Côte d'Ivoire lethal yellowing phytoplasma disease (CILY) was shown to be associated with a phytoplasma in 2013 in Grand-Lahou, Côte d'Ivoire (Konan Konan et al., 2013); since then, this disease has rapidly spread to several coconut-growing villages where circa 400 ha were destroyed, and an additional 7,000 ha are under threat (Arocha-Rosete et al., 2014). Lethal yellowing (LY)-like diseases of palms have been associated with several phytoplasmas worldwide and have killed millions of palms in the last 40 years (Sullivan and Harrison, 2013).

The CILY phytoplasma was recently classified as a member of group 16SrXXII, subgroup – B, '*Candidatus* Phytoplasma palmicola'–related strain (Harrison et al., 2014); this group also includes the Cape St. Paul Wilt Disease phytoplasma (CSPWD) strain from Ghana, which collapsed the Ghanaian coconut industry in the last 20 years (Danyo, 2011).

Bila et al. (2015) identified three phytoplasma strains that were associated with the LY in Mozambique, which included '*Ca. P. palmicola*' (16SrXXII-A), the Tanzanian LD strain (16SrIV-C), and a '*Ca. P. pini*' – related strain (16SrXXI-A) found in co-infection with a '*Ca. P. palmicola*' strain. Phytoplasmas of the 16SrI group, '*Ca. P. asteris*'-related, were recently identified from CILY phytoplasma-infected coconut palms from ~~coconut-growing~~ villages in Grand-Lahou (Kwadjo et al., 2018), indicating that mixed phytoplasma infection occurs in coconut palms affected by CILY along the south coastal littoral of ~~Grand-Lahou~~, Côte d'Ivoire.

The present work was conducted to elucidate some of the biological properties of phytoplasma isolated from coconut palms affected by CILY disease. Samples from fifteen coconut palms were collected ~~from coconut-growing farms in Grand-Lahou in Côte d'Ivoire~~ during 2015 and 2016 and used for the isolation and cultivation of phytoplasmas in artificial media. Data were collected on the colony morphology and the biochemical performance of the phytoplasma strains isolated from these coconut palms, which were previously confirmed to be CILY phytoplasma-infected. These results represent the first steps of phytoplasma species description, following the ICSB minimal standard criteria (1995).

Material and methods

Plant material

During 2015 and 2016, surveys were carried out in the following seven coconut-growing villages of Grand-Lahou: Baffredon, Badadon, Palmindustrie V1, Yaokro, Doudougbaouzou, Adjadon and Amanikro (Fig. 1).

Thirty-nine samples of leaves, inflorescences, and trunk borings were collected from 15 coconut palms of West African Tall (WAT) or PB121 (Malayan Yellow Dwarf × WAT) ecotypes that were approximately 30 years old and exhibited CILY symptoms corresponding to the disease stages 1, 2 and 3 (Table 1) (Arocha-Rosete et al., 2017); these ecotypes were representative of the ecotypes cultivated in the area. A sample from one symptomless palm was also collected from each village to serve as a negative control. Trunk borings were collected by boring the trunk using an 8-cm drill bit sterilised in 70% ethanol prior to each sample collection. The trunk was also surface-sterilized at approximately 1.20 m from the soil, where it was drilled, and the sawdust was collected in a sterile plastic bag that was placed in a cool container until it reached the laboratory. The sawdust was then transferred in a sterile manner to sterile polycarbonate tubes (50 mL) and maintained under cool conditions (6 to 8°C) until further processing.

PCR and RFLP analyses

Total DNA was extracted from 1 g of each sample type using the CTAB (Cetyl trimethylammonium bromide) method (Angelini et al., 2001). Phytoplasma DNA was amplified by PCR using the 16S rDNA universal primer pairs R16F2n/R2 (Gundersen and Lee, 1996) nested with U5/U3 primers (Lorenz et al., 1995). Deionized distilled sterile water (DDSW) (Sigma-Aldrich, St Louis, USA) was used as a non-template control for each PCR reaction. Each PCR reaction was performed in a total volume of 25 µL, containing 2.5 µL of the 10× buffer, 200 µM dNTPs, 0.625 U of *Taq* polymerase (Sigma-Aldrich, St Louis, USA), and 0.4 µM of each primer. The PCR cycling conditions were set as previously described (Schaff et al., 1995). Six microliters of the PCR products were separated by electrophoresis on 1% agarose gels, stained with ethidium bromide, and UV photographed. The preliminary

classification of the phytoplasmas at the group/subgroup level was performed with RFLP analyses, as previously reported (Gundersen and Lee, 1996).

Isolation in artificial media

A previously described phytoplasma isolation method (Contaldo et al., 2016) was employed for the symptom-bearing and symptomless palms, using leaves, inflorescences and trunk boring tissues. The samples tested were from four out of the seven villages surveyed; samples that were more representative of the various stages of the disease were selected (Table 1).

The leaf and inflorescence pieces were soaked in 5 mL of CBI (liquid medium CB) ~~orange liquid medium~~ (Contaldo et al., 2016), which was examined daily for an acidification-induced colour change from orange to yellow. ~~Non-inoculated tubes were subjected to the same conditions.~~

For the trunk borings, approximately 100 mg from each plant was placed in two separate sterile mortars and mixed with 1 mL of CBI ~~orange liquid medium~~ (Contaldo et al., 2016) using a sterile pestle. Immediately afterward, the mixtures were transferred into 1.5-mL sterile tubes. ~~In a sterile laminar hood,~~ 100 µL of the mixtures were transferred into 8-mL Monovette Urine tubes (Sigma-Aldrich, St Louis, USA) and incubated at 25±1°C. The medium was examined daily for an acidification-induced colour change from orange to yellow. Non-inoculated tubes were prepared following the same procedure.

After ~~the medium exhibited a~~ colour change, 100 µL of medium per sample was pipetted from each tube and used to inoculate plates containing 8 mL of CBs solid medium (Contaldo et al., 2016).

Strain purification and molecular identification

Single phytoplasma colonies (Contaldo et al, 2012, 2016) grown for 24 to 36 hours were picked with sterile toothpicks and transferred into 8-mL ~~Monovette tubes (Sigma-Aldrich, St Louis, USA)~~ containing 2 mL of ~~fresh~~ CBI ~~liquid medium~~ for the subsequent purification

steps. The colony purification was performed by gentle filtration of the entire liquid culture through 0.8-µm membrane filters (Whatman, Maidstone, UK) to avoid plugging, followed by $10^{-3}/10^{-4}$ serial dilutions in CBI ~~liquid medium~~. Both the undiluted filtrate and the filtrate dilutions were cultured on solid CBs plates and incubated at 25 ± 1 °C. This process, including the filtration, was repeated three times from single descending colonies. The final plates, which contained colonies that had grown for 72 h, were assessed and photographed under an optical bifocal microscope at 40~~x~~ magnification (Nikon, Tokyo, Japan).

Single colonies from the exponential growth phase were marked, picked and dissolved in 100 µL of DDSW, and then subjected to nucleic acid extraction (Contaldo et al., 2016). Aliquots of CBI ~~liquid media~~ containing the isolation material were processed for DNA extraction using the alkaline lysis/phenol/chloroform method (Pourbakhsh et al., 2013). The detection and identification of phytoplasmas were performed with specific PCR/RFLP assays ~~based upon~~ the 16S rRNA gene sequence. One µL of the total extracted DNA was used as template for PCR with the primers 758f (=M1)/B6, followed by nested amplification using the primers M1/1232r (=M2) (Gibb et al., 1995; Padovan et al., 1995). Further PCR analyses were then performed using the primer pairs R16F2n/R2 (Gundersen and Lee, 1996) nested with U5/U3 primers (Lorenz et al., 1995), as for the DNA from original plant materials. For each PCR reaction, DDSW and non-inoculated media were used as negative controls, and the samples were then processed as previously reported (Contaldo et al., 2016).

Sequencing and virtual RFLP analyses

The direct sequencing of ~~the~~ selected M1/M2 and U5/U3 PCR products was performed as previously described (Contaldo et al., 2012, 2016). The sequences ~~were subjected to~~ virtual RFLP analyses ~~based upon~~ portions of the 16S ribosomal gene ~~from the obtained sequences of the coconut and colony samples and ‘Candidatus Phytoplasma’ sequences for the corresponding subgroups available in GenBank,~~ using pDRAW32 (<http://www.acaclone.com/>). In particular, elm yellows (EY) ‘*Ca. P. ulmi*’ (GenBank

accession number, AC, AY197655), “stolbur” (STOL11), ‘*Ca. P. solani*’ (AC, AF248959), Cape St. Paul wilt phytoplasma (CSPWB), ‘*Ca. P. palmicola*’ (AC, JQ868442), Coconut lethal yellowing (LYDM, Mozambique), ‘*Ca. P. palmicola*’ (AC EU549768), aster yellows (MAY) ‘*Ca. P. asteris*’ (AC M30790) were included.

Antimicrobial susceptibility

To evaluate the *in vitro* antimicrobial activity, seven antibiotics were tested against two phytoplasma isolates from trunk borings, CILY-XII-A (9T-16) and CILY-XXII-B (17T-16). The standard disc diffusion method was employed to determine the level of antibiotic susceptibility following both, the guidelines of the Clinical and Laboratory Standard Institute and Hannan’s protocols (Hannan, 2000; CLSI, 2002). The assays were performed using rifampicin (2.5 mg/mL in DDSW), 5-fluorouracil [0.5 mg/mL in C₂H₆O, ethanol (Sigma-Aldrich, St Louis, USA)], tetracycline (0.5 mg/mL in C₂H₆O), tobramycin (0.5 mg/mL in DDSW), polymyxin (0.5 mg/mL in DDSW) and cephalixin hydrate (0.5 mg/mL in C₂H₆O). All antibiotics were obtained from Sigma-Aldrich (St Louis, USA). Standard sterile 6-mm paper discs (Whatman, Maidstone, UK) were gently placed on the surface of the agar plates, which were previously inoculated with 10⁸ CFU/mL of phytoplasma isolates and 25 µL of each antibiotic at the indicated concentrations. The plates were then incubated at 25±1°C for 24 h in triplicate. Diffusion distances were calculated by subtracting half of the disc diameter from the inhibition zone diameter (measured with a centimetre ruler). For comparison, the negative control disc containing DDSW without any added antibiotic was included in the experiment.

Biochemical properties

The biochemical tests that were previously described for *Mollicutes* were employed (Aluotto et al., 1970) to assess the specific properties of the abovementioned two isolates. These tests

included analyses of the breakdown of glucose, arginine and urea. Three types of media were prepared, each containing 10 mL of the 10% w/v test substrate (either arginine, urea or glucose) (Merck, Darmstadt, Germany). All media included 10 mL of horse serum (Oxoid, Basingstoke, Hampshire, UK), 5 mL of 10% w/v yeast extract stock solution (Oxoid, Basingstoke, Hampshire, UK), and 1 mL of 0.5% phenol red (Sigma-Aldrich, St Louis, USA) in 74 mL of CBI medium. The pH values were adjusted to 7.6 (for the glucose medium), 7.0 (for the arginine medium), and 7.0 (for the urea medium). The test tubes were inoculated with 500 μ L of overnight CBI culture and incubated at $25 \pm 1^\circ\text{C}$. The following two sets of controls were incubated at the same time: one set of three media, ~~which~~ each contained either glucose, arginine or urea was inoculated with DDSW; and a second set of CBI media adjusted to the indicated pH values (~~as mentioned above~~) with 5 N NaOH (Sigma-Aldrich, St Louis, USA) or 5 N HCl (Sigma-Aldrich, St Louis, USA) was inoculated with the CBI ~~liquid~~ cultures. The pH values for each tube were measured daily with pH strips (Macherey-Nagel, Düren, Germany). The growth was measured by a decrease of 0.5 pH units or more in the glucose-containing tubes compared to the negative control, or an increase of 0.5 pH units or more in the arginine- and urea-containing tubes. The same biochemical tests were performed on a pure culture of *Acholeplasma laidlawii* strain PG8 (ATCC# 1039) that was kindly provided by Dr. M. Kube (Thünen-Institute of Forest Genetics, Großhansdorf, Germany) and used as reference control.

Growth curves

Two flasks containing 20 mL of CBI medium were inoculated with 2 mL of 9T-16 and 17T-16 culture suspensions (ca. 10^8 CFU/mL) that were measured with a previously calibrated Beckman Coulter DU730 spectrophotometer (Brea, USA) ($0.1 \text{ OD}_{600 \text{ nm}}$). The flasks were then incubated at 24°C on a KS10 rotational shaker (Edmund Buhler, Tübingen, Germany) set at 80 oscillations per min for 2 to 3 weeks. Individual suspensions corresponding to the two isolates were six-fold serially diluted in CBI medium. Two 100- μ L aliquots of each dilution were plated onto three ~~individual~~ CB plates and incubated at 24°C for 2 to 3 days before the

colonies were counted. Two independent replicates for each phytoplasma strain were tested 30 days apart. For each growth curve, during the exponential phase, the number of phytoplasma generations (n) and the time elapsed between generations (duplication time) were calculated with the Pedersen formula (Panikov, 1995).

$$n = (\log N_t - \log N_0) / \log 2 = (\log N_t - \log N_0) / 0,301$$

- $N(t)$ = the number of cells at time t
- $N(0)$ = the number of cells at time 0

qPCR analysis

The phytoplasma presence and titre were verified in 9T-16 and 17T-16 culture suspensions at the inoculation time (T_0) and after 18 hours (T_{18}) following DNA extraction (Pourbakhsh et al., 2013) by qPCR analysis in an ABI PRISM®7000 cycler (Applied Biosystems, Waltham, USA). The primer pair 16S(RT)F1/16S(RT)R1 was used, and a previously reported protocol was followed (Saccardo et al., 2012). The assays were carried out in duplicate, each experiment was repeated three times and the mean comparisons were evaluated. An additional sample was added to each plate as a DNA-free negative control.

For the quantification of phytoplasma genes, a standard curve was produced using the amplicons obtained from 10-fold serial dilutions ranging from 10 to 1×10^{-7} ng / μ L of the phytoplasma strain STOL (16SrXII-A) (<http://www.ipwgnnet.org/collection>). The standard curve was constructed by the interpolation of the $\log_{10}$ of the copy number of each standard dilution with the respective threshold cycle (C_t) values. Relative quantification of gene expression in the phytoplasma samples was estimated from the extrapolation of the standard curve values.

Results

Phytoplasma identification

Prior to the isolation and cultivation trials, PCR testing yielded amplicons for all of the CILY symptom-bearing coconut palms corresponding to the 16SrXXII-B phytoplasma. Two of the palms showed mixed infection with 16SrI phytoplasmas. PCR testing yielded no amplicons for any of the symptomless palms or the negative controls (Table 1).

A total of 39 samples corresponding to trunk borings, leaves and inflorescences collected from 15 coconut palms were used for the isolation trials. Inflorescences were not collected from palms exhibiting CILY symptoms of stage 3, as they were rotten before reaching the laboratory. The CBI liquid media changed colour from orange to yellow at varying times after inoculation, depending on the plant tissue. Particularly for trunk borings, the colour change was visible from 24 to 36 hours post-inoculation. For leaves, medium acidification was not visible until 1 week post-inoculation. For inflorescences, the colour change times post-inoculation were extremely variable due to the contamination of the medium by several other microorganisms, primarily bacteria. This contamination rendered the inflorescences unsuitable for phytoplasma isolation under the current procedures. The non-inoculated tubes did not exhibit colour changes or colony growth throughout the experiments. The plating from tubes, immediately after the medium acidification (~~colour change~~), produced phytoplasma colonies 1 to 2 days post-inoculation (Fig. 2). The single colony purification method was used to rule out contamination with larger bacteria and to prepare the isolates for further analyses.

Phytoplasma DNA amplification was performed on a total of 18 samples from liquid and/or solid media (Table 1). The results revealed that 7 of the tested tubes with isolation

materials from symptom-bearing palms were positive; of these, 6 were from trunk borings, and 1 was from a leaf midrib. An inoculated CBI tube corresponding to a trunk boring sample from one symptomless coconut palm was positive for ~~the presence of~~ 16SrI phytoplasmas ('*Ca. P. asteris*'), whereas the trunk borings from the three remaining symptomless plants did not exhibit colour changes or colony growth in the medium. Phytoplasma DNA was also amplified from the colonies obtained from the 14 inoculated tubes at the exponential phase. The nested PCR was necessary since the yield of the recovered DNA with the kit used was about 10 ng/μl.

The results of blast analyses of U5/U3 and/or M1/M2 ~~amplicons~~, deposited in GenBank under accession numbers MF419688 to MF419695, confirmed the presence of ~~various strains of 'Candidatus Phytoplasma' including~~ '*Ca. P. asteris*', '*Ca. P. ulmi*', '*Ca. P. solani*,' which had 99% identity with the corresponding '*Candidatus Phytoplasma*' strains; the identity to the corresponding '*Candidatus Phytoplasma*' was 100% only for '*Ca. P. palmicola*'. The results of the virtual RFLP analyses further confirmed the phytoplasma classification in groups 16SrI, 16SrV, 16SrXII and 16SrXXII (Figure 3). The isolation of one '*Ca. P. asteris*' strain with SNPs to strains enclosed in subgroups 16SrI-B was revealed by *Mse*I restriction analyses (Figure 3) and sequence comparison (data not shown).

Antimicrobial susceptibility of two phytoplasma isolates

The results of the antibiotic susceptibility tests revealed that tobramycin exhibited the maximum activity against the two phytoplasma isolates, followed by polymyxin and tetracycline. The two isolates displayed intermediate susceptibility to 5-fluorouracil but were completely resistant to cephalexin hydrate and rifampicin (Table 2).

Biochemical properties of two phytoplasma isolates

The results of the biochemical tests for urea and arginine hydrolysis and glucose fermentation confirmed that both phytoplasma strains were negative for urea and positive for glucose and arginine (Fig. 4). For both phytoplasma strains, a decrease in the pH from 7.6 to 6.0 was observed in the glucose test, and a decrease in the pH from 7.0 to 5.5 was observed in the urea tests, whereas an increase in the pH of 1.5 (from 7.0 to 8.5) was estimated for the arginine test. On the other hand, *A. laidlawii* was positive for glucose and negative for urea and arginine, as previously reported [26] (Fig. 4).

Growth curves of two phytoplasma isolates

Growth curves were determined for the 9T-16 and 17T-16 phytoplasma isolates (Fig. 5a). The growth rate of the 17T-16 (16SrXXII-B) isolate was higher than that of the isolate 9T-16 (16SrXII-A) by a factor of 10 at 12 hrs/growth. Both phytoplasma isolates reached a final growth plateau between 48 and 72 hrs with a concentration of approximately 10^{10} CFU/mL, but showed also intermediate growth plateau at about 24 hrs. Furthermore, the number of generations and the time elapsed between each generation were lower for the phytoplasma isolate 17T-16 compared with isolate 9T-16 (Fig. 5b). The duplication time (time elapsed between two generations) of the two phytoplasma isolates, as estimated by colony count, was 250 and 500 minutes for 9T-16 (16SrXII-A) and 17T-16 (16SrXXII-B), respectively. In conclusion, the isolate 17T-16 was shown to multiply less efficiently in CBI or CBs media after 24 hours of cultivation compared with the isolated 9T-16.

qPCR analysis of two phytoplasma isolates

From the qPCR assay, an increase in the phytoplasma titre was detected for both isolates. The T_m of the standard curves exhibited a single peak at $83 \pm 1^\circ\text{C}$, and all of the dissociation curves from phytoplasma DNA exhibited a single peak at the same temperature. The C_t values ranged from 6.5 to 35.0 for the 10-fold STOL dilutions and from 31.4 to 35 for the phytoplasma dilutions (Table 3). An increase of the phytoplasma titre was therefore detected between time 0 and 18 hours; the C_t values were not determined in the following 24 – 48 considering the growth plateau registered at about 24 hours.

Discussion

This is the first report of the isolation and growth of phytoplasmas from coconut palms in artificial media. The trunk borings, compared to leaves and inflorescences, proved to be the most suitable plant tissue for phytoplasma isolation, as the majority of phytoplasma colonies were obtained from these materials. This finding is in agreement with the fact that trunk borings have previously been shown to be the most optimal plant tissue for phytoplasma sampling and detection in coconut and other palm species (Harrison et al., 2013). The 16SrXXII-B isolated phytoplasmas correspond to the prevalent subgroup associated with CILY in Côte d'Ivoire (Arocha-Rosete et al., 2017). Furthermore, the detection of phytoplasmas of diverse ribosomal subgroups, such as 16SrI-B, 16SrV-A, 16SrXII-A and 16SrXXII-B from typical phytoplasma colonies, confirmed for the first time the natural occurrence of a mixed phytoplasma infection in coconut palms affected by CILY. The comparison between the phytoplasma subgroups found in the plant material prior to culture and those found in the liquid and solid media indicates the presence of different phytoplasmas

in the tissues used for isolation. These somewhat contradictory results on phytoplasma identification obtained from plant material, CBI and CB₁ medium, confirm the presence of a mixed infection. While it is very likely that the prevalent strain in the plant material is the strain that determines the PCR/RFLP pattern in plants, the medium appears to favour other phytoplasma strains. Moreover, the differences in the identity of the phytoplasma that were detected from testing the DNA extracts from the CB liquid cultures and plate colonies further indicate the presence of mixtures of phytoplasmas. This finding was further confirmed by the detection of a mixed infection in both liquid and solid media. Moreover, the symptomless and initially PCR-negative sample, 2T-16, exhibited no phytoplasma in the starting plant material when analysed by PCR. However, this sample later exhibited a single phytoplasma subgroup in the liquid medium and a mixture of two subgroups on the solid medium. In particular, the presence of 16SrXII-A and 16SrI in the latter case confirms that the media used in this study are especially suitable for the growth of these phytoplasma, as previously reported (Contaldo et al., 2016). Recently, the CILY phytoplasma was detected in a mixed infection with 16SrI phytoplasmas in both CILY-affected coconut palms and the potential leafhopper vector, *Nedotepa curta* Dimitriev, in coconut-growing villages of Grand-Lahou (Kwadjjo et al., 2018); this finding partly confirms the results obtained from cultivation.

The presence of multiple phytoplasmas in a single plant host may modulate the expression of disease symptoms, as was suggested regarding the presence of multiple apple proliferation phytoplasma strains in apple trees or for other phytoplasma-associated diseases (Seemüller and Schneider, 2007; Rid et al., 2016).

The biochemical characterization of 16SrXII-A, ‘*Ca. P. solani*’ and 16SrXXII-B ‘*Ca. P. palmicola*’ isolates revealed that both 9T-16 (16SrXII-A) and 17T-16 (16SrXXII-B) shared metabolic features with *Mycoplasma* species group III (glucose positive/arginine positive), which are represented by several mycoplasma species (i.e., *M. fermentans*) (Brown et al., 2007; Freundt et al., 1997). The ability to metabolize both glucose and arginine differentiates these phytoplasma isolates from *Acholeplasma* and *Mycoplasma* species, which were isolated

from the surfaces of several tropical ~~floral~~ plants (Rose et al., 1990). From this study, both phytoplasma isolates, 9T-16 and 17T-16 were shown to be positive for arginine, which suggests that these isolates may possess either the arginine dehydrolase pathway or, at least, certain components of ~~this~~ pathway for the ~~hydrolysis of~~ arginine. This finding may be an indicator that supports the differentiation of phytoplasmas from other *Mollicutes* species that inhabit the phloem of coconut palms, such as *A. palmarum* (Tully et al., 1994). Previous biochemical tests performed on various palm cultivars grown in Florida revealed that arginine was readily detected in lethal yellowing susceptible palm ecotypes, but it was absent (or present at minimal detectable levels) in LY-resistant palms, which led to the hypothesis that LY phytoplasmas may hydrolyse arginine (Mc Coy, 1984).

On the other hand, the medium-dependent growth kinetics for each isolate (9T-16 and 17T-16) indicated their different performances in ~~each media~~. This phenomenon was clearly evidenced by the longer growth timeframe for the 16SrXII-A (9T-16) isolate and partly verified by the qPCR assays at the initial and 18 hours growing times. The 16SrXXII-B phytoplasma correspond to the prevalent subgroup associated with CILY in Côte d'Ivoire [13]; however, the isolate 17T-16 was shown to multiply less efficiently in CBI or CBs media after 24 hours of cultivation. The shorter survival time in the artificial medium for isolate 17T-16 may be associated with a limiting factor of the CILY phytoplasma multiplication rate in cell-free media. Further biochemical comparative studies on CILY phytoplasma-free and CILY phytoplasma-infected trunk borings may lead to the identification of new chemicals or bio-compounds which may improve the media composition to support ~~a~~ better growth performance and survival rate of 16SrXXII-B phytoplasmas.

The assessment of the phytoplasma susceptibility to antibiotics performed in this study was the first ~~such assessment~~ performed on cultured phytoplasmas and provided ~~a clear~~ indication of the resistance/susceptibility of the two studied isolates. This work is also the first report of phytoplasma growth kinetics, clearly demonstrating that the various isolates exhibited different performances in the same growth medium, confirming that different phytoplasmas

exhibit different biological behaviour and very likely possess different plant colonization abilities. This phenomenon was clearly evidenced and confirmed by the longer growth timeframe registered for the 16SrXII-A (9T-16) isolate, which was not detected in the original coconut tissues by nested-PCR.

Conclusions

The first assessment of the antibiotic susceptibility of 16SrXII-A and 16SrXXII-B phytoplasmas isolated from coconut palm and their biochemical characterization provided experimental evidence about certain metabolic features of these strains. The availability of phytoplasma colonies associated with CILY disease represents a novel and important tool for the further study of their biology, pathogenicity and epidemiology and to support their species classification. This newly achieved coconut phytoplasma growth in both solid and liquid artificial media and the biological characterization of two isolates are important advancements in the field of disease management and containment measures.

Acknowledgements

This work was supported by the International Development Research Centre (IDRC), Ottawa, Canada, www.idrc.ca, [Grant No. 107789], with financial support from the Government of Canada, provided through Foreign Affairs, Trade and Development Canada (DFATD), www.international.gc.ca

References

- Aluotto, B.B., Wittler, R.G., Williams, C.O., Faber, J.E., 1970. Standardized bacteriologic techniques for the characterization of mycoplasma species. *Int. J. Syst. Bacteriol.* 20(1), 35-58.
- Angelini, E., Clair, D., Borgo, M., Bertaccini, A., Boudon-Padieu, E., 2001. “Flavescence dorée” in France and Italy - Occurrence of closely related phytoplasma isolates and their near relationships to Palatinate grapevine yellows and an alder yellows phytoplasma. *Vitis* 40(2), 79-86.
- Arocha-Rosete, Y., Konan Konan, J.L., Diallo, A.H., Allou, K., Scott, J.A., 2014. Identification and molecular characterization of the phytoplasma associated with a lethal yellowing-type disease of coconut in Côte d’Ivoire. *Can. J. Plant Pathol.* 36(2), 141-150.
- Arocha Rosete, Y., Atta Diallo, H., Konan Konan, J.L., Yankey, N., Saleh, M., Pilet, F., Contaldo, N., Paltrinieri, S., Bertaccini, A., Scott, J.A., 2017. Improving sampling and detection for the coconut lethal yellowing phytoplasma in coconut-growing villages of Grand-Lahou, Côte d’Ivoire. *Ann. Appl. Biol.* 170, 333–347.
- Bila, J., Mondjana, A., Samils, B., Högborg, N., 2015. High diversity, expanding populations and purifying selection in phytoplasmas causing coconut lethal yellowing in Mozambique. *Plant Pathol.* 64(3), 597-604.
- Brown, D.R., Whitcomb, R.F., Bradbury, J.M., 2007. Revised minimal standards for description of new species of the class Mollicutes (division Tenericutes). *Int. J. Syst. Evol. Microbiol.* 57, 2703–2719.
- CLSI, 2002. Clinical and Laboratory Standard Institute, development of in vitro susceptibility testing criteria and quality control parameters for veterinary antimicrobial agents. CLSI document, approved guideline, 22 Wayne, Pennsylvania, USA.

- Contaldo, N., Bertaccini, A., Paltrinieri, S., Windsor, H.M., Windsor, D.G., 2012. Axenic culture of plant pathogenic phytoplasmas. *Phytopath. Medit.* 51(3), 607–617.
- Contaldo, N., Satta, E., Zambon, Y., Paltrinieri, S., Bertaccini, A., 2016. Development and evaluation of different complex media for phytoplasma isolation and growth. *J. Microbiol. Meth.* 127, 105-110.
- Danyo, G., 2011. Review of scientific research into the Cape Saint Paul wilt disease (CSPWD) of coconut in Ghana. *Afr. J. Agr. Res.* 6, 4567–4578.
- Doi, Y., Teranaka, M., Yora, K., Asuyama, H., 1967. Mycoplasma or PLT group-like microorganisms found in the phloem elements of plants infected with mulberry dwarf, potato witches' broom, aster yellows, or *Paulownia* witches' broom. *Ann. Phytopath. Soc. Jpn.* 33, 259–266.
- Freundt, E.A., Erno, H., Lemcke, R.M., 1979. Identification of *Mycoplasma*. In Bergan & Norris (Eds.), *Methods in microbiology*. Academic Press London. pp 396.
- Gibb, K.S., Padovan, A.C., Mogen, B.A., 1995. Studies on sweet potato little-leaf phytoplasmas detected in sweet potato and other plant species growing in Northern Australia. *Phytopathology* 85, 169-174.
- Gundersen, D.E., Lee, I.-M., 1996. Ultrasensitive detection of phytoplasmas by nested-PCR assays using two universal primer pairs. *Phytopath. Medit.* 35, 144-151.
- Hannan, P.C., 2000. Guidelines and recommendations for antimicrobial minimum inhibitory concentration (MIC) testing against veterinary *Mycoplasma* species. *Vet. Res.* 31, 373–395.
- Harrison, N.A., Davis, R.E., Helmick, E., 2013. DNA extraction from arborescent monocots and how to deal with other challenging hosts. In M. Dickinson and J. Hodgetts (Eds.), *Phytoplasmas: Methods and Protocols*. Humana Press, Springer: New York, pp. 147–158.
- Harrison, N., Davis, R.E., Oropeza, C., Helmick, E., Narvaez, M., Eden-Green, S., Dollet, M., Dickinson, M., Konan Konan, J.L. 2014. '*Candidatus Phytoplasma palmicola*', a novel

- taxon associated with a lethal yellowing-type disease (LYD) of coconut (*Cocosnucifera* L.) in Mozambique. Int. J. Syst. Evol. Microbiol. 64, 1890-1899.
- ICSB, International Committee of Systematic Bacteriology, Subcommittee on the Taxonomy of Mollicutes, 1995. Revised minimum standards for descriptions of new species of the class Mollicutes (Division Tenericutes). Int. J. Syst. Bacteriol. 45, 605–612.
- Konan Konan, J.L., Allou, K., Atta Diallo, H., Saraka Yao, D., Koua, B., Kouassi, N., Benabid, R., Michelutti, R., Scott, J.A., Arocha-Rosete, Y., 2014. First report on the molecular identification of the phytoplasma associated with a lethal yellowing-type disease of coconut palms in Cote d'Ivoire. New. Dis. Rep. 28, 3.
- Kwadjo, K.E., N'Djiha Beugré, I., Dietrich, C., Thierry Kodjo, A.T., Atta Diallo, H., Konan Konan, J.L., Contaldo, N., Paltrinieri, S., Bertaccini, A., Arocha-Rosete, Y., 2017. Identification of *Nedotepacurta* Dmitriev as a potential vector of the Côte d'Ivoire lethal yellowing phytoplasma in coconut palms sole or in mixed infection with a 'Candidatus Phytoplasma asteris'-related strain. Crop Protect. 110, 48-56.
- Lorenz, K.H., Schneider, B., Ahrens, U., Seemüller, E., 1995. Detection of the apple proliferation and pear decline phytoplasmas by PCR amplification of ribosomal and nonribosomal DNA. Phytopathology 85, 771-776.
- Mc Coy, R.E., 1984. Presence of free arginine associated with susceptibility to lethal yellowing of palms. Isr. J. Med. Sci. 20(10), 1016-1018.
- Padovan, A.C., Gibb, K.S., Bertaccini, A., Vibio, M., Bonfiglioli, R.E., Magarey, P.A., Sears, B.B., 1995. Molecular detection of the Australian grapevine yellows phytoplasma and comparison with a grapevine yellows phytoplasma from Emilia-Romagna in Italy. Austral. J. Grape Wine Res. 1, 25-31.
- Panikov, N.S., 1995. Microbial Growth Kinetics. Chapman & Hall eds. pp 378, ISBN 0-412-56630-3.

- Pourbakhs, S.A., Maghami, M., Ashtari, A., Bayatzadeh, M.A., Ahangaran, S., 2013. The *vlhA* gene sequencing of Iranian *Mycoplasma synoviae* isolates. Arch. Razi. Inst. 68(2), 117–124.
- Rid, M., Mesca, C., Ayasse, M., Gross, J., 2016. Apple proliferation phytoplasma influences the pattern of plant volatiles emitted depending on pathogen virulence. Front. Ecol. and Evol. 3, 152.
- Rose, D.L., Kocka, J.P., Somerson, N.L., Tully, J.G., Whitcomb, R.F., Carle, P., Bové, J.M., Colflesh, D.E., Williamson, D.L., 1990. *Mycoplasma lactucae* sp. nov. a sterol-requiring Mollicute from a plant surface. Int. J. Syst. Bact. 40, 138-142.
- Saccardo, F., Martini, M., Palmano, S., Ermacora, P., Scortichini, M., Loi, N., Firrao, G., 2012. Genome drafts of four phytoplasma strains of the ribosomal group 16SrIII. Microbiology 158, 2805–2814.
- Schaff, D.A., Lee, I.-M., Davis, R.E., 1995. Sensitive detection and identification of mycoplasma-like organisms by polymerase chain reactions. Biochem. Biophys. Res. Comm. 186, 503-1509.
- Seemüller, E., Schneider, B., 2007. Differences in virulence and genomic features of strains of ‘*Candidatus Phytoplasma mali*’, the apple proliferation agent. Phytopathology 97, 964–970.
- Sullivan, M., Harrison, N., 2013. CPHST Pest Datasheet for ‘*Candidatus Phytoplasma palmae*’ and related strains. USDA-APHIS-PPQ-CPHST. <http://caps.ceris.purdue.edu>
- Tully, J.G., Whitcomb, R.F., Rose, D.L., Bové, J.M., Carle, P., Somerson, N.L., Williamson, D.L., Eden-Green, S., 1994. *Acholeplasma brassicae* sp. nov. and *Acholeplasma palmae* sp. nov., two non-sterol-requiring mollicutes from plant surfaces. Int. J. Syst. Bacteriol. 44(4), 680-684.
- Weintraub, P.G., Beanland, L., 2006. Insect vectors of phytoplasmas. Ann. Rev. Entomol. 51, 91-111.



Fig. 1. A map showing the villages of Grand-Lahou that were surveyed for the presence of CILY phytoplasma in Côte d'Ivoire; red circles indicate the villages from which phytoplasma isolation was performed.




Fig. 2. Phytoplasma colonies. A: colonies corresponding to 16SrV-A phytoplasma, B: colonies corresponding to 16SrXXII-B phytoplasma, C: colonies corresponding to 16SrI (left) and 16SrXII-A (right) phytoplasmas, D: magnification of a single phytoplasma colony.






Fig. 3 Virtual RFLP 3% gels showing the polymorphisms of the diverse ‘*Ca. Phytoplasma*’ detected in coconut and culture samples obtained in this work and retrieved from GenBank. The enzyme used was *MseI*. In A and B the sequences used are from U5/U3 amplicons, while in C and D the sequences are from M1/M2 amplicons. The acronyms are reported in materials and methods and Table 1. P, marker phiX174 *HaeIII* digested with fragment sizes in base pairs from top to bottom of 1,353; 1,078; 872; 603; 310; 281; 271; 234; 194; 118 and 72.



A



B

Fig. 4. Biochemical tests. **A.** Numbers 1-2, liquid medium inoculated with DDSW and adjusted to a pH of 7.0. and 7.6, respectively. Numbers 3-4-5, glucose fermentation: 3 (*A. laidlawii*), 4 (isolate 9T-16), 5 (isolate 17T-16), pH variation at 48 hrs from 7.6 to 6.0. Numbers 6-7-8, urea hydrolysis: 6 (*A. laidlawii*), 7 (isolate 9T-16), 8 (isolate 17T-16). Numbers 9-10-11, pH variation at 48 hrs from 7.6 to 6.0; arginine hydrolysis: 9 (*A. laidlawii*),

10 (isolate 9T-16), 11 (isolate 17T-16), pH variation only in tubes 10 and 11 ~~at 48 hrs~~ from 7.0 to 8.5. **B.** Tubes U1, G1 and A1, two replicas of the isolate 9T-16 and U2, G2 and A2 two replicas of the isolate 17T-16 both inoculated in media with the three different substrates (urea, glucose and arginine, respectively). Tubes U3, G3 and A3, liquid medium containing either urea (U) glucose (G) or arginine (A) inoculated with DDSW. Tubes U4, G4 and A4, media adjusted to a pH of 7.0 and inoculated with the isolate 17T-16. Tubes U5, G5 and A5, media adjusted to a pH of 7.6 and inoculated with the isolate 17T-16.




Fig. 5. Growth kinetics of phytoplasma isolates from coconut palms 9T-16 (16SrXII-A) and 17T-16 (16SrXXII-B). A, growth curves. B, number of generations and time elapsed between each generation (calculated with the Pedersen formula).

Table 1. Phytoplasma identification in coconut palm samples.

Sample code	Sample type	Disease stage	Village	Sampling month	Phytoplasma PCR/RFLP identification		
					From plant material	From CBI	From CBs*
2T-15	Trunk boring	1	Badadon	03/ 2015	16SrXXII-B	16SrXII-A	16SrXII-A
20T-15	Trunk boring	3	Badadon	03/ 2015	16SrXXII-B	16SrXXII-B	16SrI
40T-15	Trunk boring	1	Badadon	06/ 2015	16SrXXII-B	-	16SrI
41T-15	Trunk boring	2	Badadon	06/ 2015	16SrXXII-B	-	16SrI
11F-15	Inflorescence	2	Badadon	06/ 2015	16SrXXII-B	-	16SrI
15F-15	Inflorescence	2	Badadon	06/ 2015	16SrXXII-B	-	16SrXII-A
43T-15	Trunk boring	2-3	Badadon	06/ 2015	16SrXXII-B	-	16SrV-A
70L-15	Leaf midribs	2	Badadon	06/ 2015	16SrXXII-B	-	16SrXII-A
2T-16	Trunk boring	Asympt.	Baffredon	04/ 2016	-	16SrI	16SrI+ 16SrXII-A
24T-16	Trunk boring	3	Yaokro	04/ 2016	16SrXXII-B	-	16SrI
20T-16	Trunk boring	3	Plamindustrie V1	04/ 2016	16SrXXII-B	16SrI	16SrI
L9-16	Leaf midribs	2	Baffredon	04/ 2016	16SrXXII-B	16SrI	16SrI
30T-16	Trunk boring	3	Palmindustrie V2	06/ 2016	16SrXXII-B + 16SrI	-	16SrI
9T-16	Trunk boring	3	Badadon	06/ 2016	16SrXXII-B + 16SrI	16SrXII-A	16SrXII-A
17T-16	Trunk boring	3	Palmindustrie V1	06/ 2016	16SrXXII-B	16SrI	16SrXXII-B

*Isolates in bold were used for biochemical and growth curve testing.

Table 2. Phytoplasma antibiotic susceptibility measured by the disc diffusion method.

Antibiotic	Diameter (mm)	$\pm$ standard error (mm)
5-fluorouracil	13	0.43
Cephalexin hydrate	2.5	0.45
Polymyxin	21	0.45
Rifampicine	2	0.2
Tetracycline	20	0.35
Tobramycin	23	0.71

Table 3. Phytoplasma titre quantification by qPCR analysis.

Sample	Ct Mean	Quantity	Quantity
		ng/ μ l	copy number
STOL- 10	6.5	10	1.54×10^{13}
STOL- 10^{-1}	12.2	10^{-1}	1.54×10^{11}
STOL- 10^{-2}	17.0	10^{-2}	1.54×10^{10}
STOL- 10^{-3}	21.5	10^{-3}	1.54×10^9
STOL- 10^{-4}	25.8	10^{-4}	1.54×10^8
STOL- 10^{-5}	29.4	10^{-5}	1.54×10^7
STOL- 10^{-6}	32.3	10^{-6}	1.54×10^6
STOL- 10^{-7}	35.0	10^{-7}	1.54×10^5
9T-16 T0*	35.0	2.1×10^{-7}	3.24×10^5
9T-16 T18**	33.3	6.0×10^{-7}	9.26×10^5
17T-16 T0*	34.2	3.5×10^{-7}	5.44×10^5
17T-16 T18**	31.4	1.7×10^{-6}	2.62×10^6

*, strain at the inoculation time; **, strain after 18 hours