



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Availability:

This version is available at: <https://hdl.handle.net/11585/792593> since: 2021-01-28

Published:

DOI: <http://doi.org/10.1016/j.marpolbul.2021.112005>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Nicolas Greggio; Marco Capolupo; Filippo Donnini; Manfred Birke; Elena Fabbri; Enrico Dinelli:
*Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon
biomonitoring*

MARINE POLLUTION BULLETIN. Vol. 164 ISSN 0025-326X

DOI: 10.1016/j.marpolbul.2021.112005

The final published version is available online at:

<https://dx.doi.org/10.1016/j.marpolbul.2021.112005>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Marine Pollution Bulletin

Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring --Manuscript Draft--

Manuscript Number:	MPB-D-20-01666R2
Article Type:	Research Paper
Keywords:	Coastal lagoon; Sediment geochemistry; Metal bioaccumulation; biomarkers; Biomonitoring.
Corresponding Author:	Nicolas Greggio, Ph.D University of Bologna Ravenna, ITALY
First Author:	Nicolas Greggio, Ph.D
Order of Authors:	Nicolas Greggio, Ph.D Marco Capolupo, Ph.D. Filippo Donnini, Ph.D. Manfred Birke, Ph.D. Elena Fabbri, Prof. Enrico Dinelli, Prof.
Abstract:	Coastal lagoons are complex environments threatened by natural and anthropogenic stressors. Here, we tested the effectiveness of combining physical, geochemical and chemical measurements with biomarker data obtained in field-exposed marine mussels (<i>Mytilus galloprovincialis</i>) as a biomonitoring strategy for a highly pressured lagoon (Pialassa Baiona, Ravenna, Italy). Data showed a spatial trend of sediment contamination by Hg, Pt, Au, Ag, Mo, Re, Cd, Pd and Zn. Local conditions of high water temperature/low conductivity were detected among selected sites. After a 30-day in situ exposure, Ag and Hg were the most bioaccumulated elements (10 and 5 folds, respectively) in mussels followed by Sb, Al, Ti and Fe. Decreased survival, lysosomal dysfunctions, increased metallothionein content and peroxisome proliferation were observed in mussels in relation to metal spatial distribution and physicochemical fluctuations. Overall, this study provides a further confirmation of the role of biomonitoring to reliably assess the environmental quality of highly pressured lagoons.

Greggio et al. Manuscript Number: **MPB-D-20-01666R1**

Dear Editor

Marine Pollution Bulletin

Dec 28th, 2020

Please find enclosed the revised manuscript authored by Greggio, Capolupo, Donnini, Birke, Fabbri and Dinelli entitled: *Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring*

We are grateful for the confirmed overall appreciation given to our work, and for the pertinent comments we received to improve the manuscript quality.

Almost all suggestions have been accepted and very clear explanations have been provided where necessary. Justification to the use of only inorganic pollutants has been included. Detailed argumentation on the organic pollutants has been provided in order to clarify their role in the work. We do hope more clarity and better focus have been reached.

Specific responses to the Reviewers' comments are provided in a separate file and changes have been highlighted on a second copy of the manuscript for easy of reading.

We do hope the revised manuscript meets the Reviewers' requests and may be considered suitable for publication on Marine Pollution Bulletin

Best Regards,

Your Sincerely

Corresponding Author
Nicolas Greggio



Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring


Authors:

Greggio Nicolas, Capolupo Marco, Filippo Donnini, Manfred Birke, Fabbri Elena, Dinelli Enrico

REVISION 2 - RESPONSE TO REVIEWERS

COMMENT	RESPONSE	POSITION
Reviewer #1		
<p>Reviewer #1: The manuscript reports an interesting study on the application of an integrated approach (geochemical and chemical measurements with biomarker data obtained in field-exposed marine mussels) for the evaluation of the environmental quality of a lagoon and despite the necessary amount of information to be reported, the authors have made an great effort to condense all the information.</p> <p>I noted that some recommendations of the other referees were accepted, improving the manuscript. The strong points of this article are: i) the strategy based on the integration of the water physico-chemical parameters and the sediment geochemical data with the measurements of the metal bioaccumulation and biomarkers in mussels. ii) the description and discussion of the results of the various lines of evidence carried out iii) the choice of to determine elements seldom analysed in sediments for environmental aims, and well described in the discussion also linking them with the possible sources of contamination.</p> <p>However, there are some aspects that are missing or need to be more detailed that I would like to highlight</p>	<p>Thanks for the appreciation</p>	
<p>The study is focused on the determination of only metals in sediments and their bioaccumulation in mussels, but in my opinion the reason for this choice should be well explained in the manuscript, having described the lagoon as an environment contaminated by both metals and organic pollutants , and therefore also the latter can contribute to the biological effects on mussels.</p>	<p>A part has been added in the introduction chapter in order to justify the choice of working only with inorganic pollutants. In fact, most studies in the area involved organic pollutants and no complete information exist on inorganic pollutants in sediments and in mussels. Moreover, few information exists on the integration of a complete inorganic pollutants suite and biomarkers .</p>	<p>Introduction chapter Lines 86-101</p>
<p>With regard to organic contaminants, in the final part of the discussion when the overall health status of the mussels of each station is evaluated, the presence of organic xenobiotics (page 23) plays a decisive role in some stations, even if haven't been reported the type of contaminants nor their concentrations. The role of organic contaminants seems a bit weak in this discussion because it relates to trends in measured AOX (Acyl-Co A oxidase), but positive correlated also with metals, and to data on PCB and PAH concentrations measured in previous studies, not reporting however when they were conducted and citing referenes of some years ago. It is difficult to correlate biomarker outcomes with contaminant concentrations in sediments of</p>	<p>The presence of organic contaminants was not evaluated in this study; we apologize if we did not correctly explain this; perhaps, we did not clarify the principle of the data integration system used to assess the overall mussel health status. The aim of this study was to evaluate the validity of combining geochemical data (i.e. metal concentrations in sediments), metal bioaccumulation and biomarker responses as a suitable tool for coastal lagoon biomonitoring; consequently, the assessment of other organics was not included in the analytical design. However, since we analyzed a wide battery of biomarkers validated in mussels, and since some of these parameters (i.e. AOX) is activated prevalently by POPs exposure, the data integration through the Mussel Expert System (MES) identified the onset of stressful conditions associated to the possible presence of POPs, notably in the sites where mussels showed increase AOX. For these reasons, the plot reporting the MES output (Fig.6) presents symbols corresponding to the "presence of POPs" as suggested by the</p>	<p>Page 10, section 2.6.5., Lines 276-285 and pages 24-25, Lines 604 – 610</p>

<p>which it is not known when and where they were sampled and analyzed.</p>	<p>Expert System output. Moreover, we did not perform any statistical correlation analyses with data from previous studies, but rather pointed out that the stress level identified by the MES in relation to the presence of POPs is coherent with previous reports of PAHs across the lagoon. In line with the Reviewer's comments, we added clarifications about these points in the Materials and Method section (Page 10, section 2.6.5., L276-285) and in Results & Discussion (MES paragraph, page 24-25, L 604 – 610).</p>	
<p>For these reasons, if it is considered important to introduce the role of organic contaminants in the overall assessment then it should review all discussion on a stronger basis and be reported i.e when the sampling was performed, information not present in the materials and methods. This information is not important in the discussion of the data relating to metals since all the measurements were performed on the same samples and in the same period.</p>	<p>Based on the considerations reported in the previous comment, we could not add any method or data on organics, since they were not analysed nor this was included as an objective of this study. However, we acknowledge that POPs, such as PAHs, have previously been reported in the lagoon and that, apart from AOX, some biomarkers of general stress (i.e. LMS, NL and/or LYS/CYT) are known to be modulated in mussels by pollutants of diverse nature, including organic and inorganic compounds, as well as their co-exposure (e.g. Capolupo et al., 2017, in bibliography). Hence, to meet the Reviewer's arguments, we added notions in the Results & Discussion section to address i) the potential relationships between chemical data from this study and those previously reported for other (organic) pollutants y (page 14, L382 – 390) and ii) the possible cumulative effects of metals and organics on the magnitude of biological effects observed in transplanted organisms (page 21, L 534 – 538)</p>	<p>page 14, Lines 382 – 390 and page 21, Lines 534 – 538</p>
<p>Another gap, in my opinion, is not having considered the grain size analysis of the sediments, even if most likely the lagoon sediments collected in the different stations are homogeneous, this information could be important considering how the concentrations of metals are related to the silt/clay fraction.</p>	<p>Authors agree with reviewer and a section has been added at the beginning of the sediment data section. Sediment grain size has been estimated through the major elements geochemical composition as established by Dinelli et al. (2007)</p>	<p>Section 3.2 Page 11 Lines 319-326</p>
<p>line 30 page 11 replace S3 and S5 at sites 3 and 5</p>	<p>Modified</p>	



Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring

Authors:

Greggio Nicolas*^{a, b}, Capolupo Marco^{a, b}, Filippo Donnini^a, Manfred Birke^c, Fabbri Elena^{a, b}, Dinelli Enrico^{a, b}

^a University of Bologna, Inter-Departmental Research Centre for Environmental Science (CIRSA), Via S. Alberto 163, 48123 Ravenna, Italy

^b University of Bologna, Department of Biological, Geological, and Environmental Sciences, Piazza S. Donato 1, 40100 Bologna, Italy

^c Federal Institute for Geosciences and Natural Resources, Stilleweg 2, 30655, Hannover, Germany

* Corresponding Author: Nicolas Greggio e-mail: nicolas.greggio2@unibo.it. University of Bologna, Inter-Departmental Research Centre for Environmental Science (CIRSA), Via S. Alberto 163, 48123 Ravenna, Italy.

Highlights

- Total Zn, Cu and Pb in sediments exceed the natural background concentrations;
- Sediments are enriched in mobile Hg, Pt, Au and Ag, with a strong S-N gradient;
- Exposed mussels bioaccumulated Ag (x10) and Hg (x5) compare to reference;
- Sediment concentration of Ag, Au, Ba, Bi, Sb affect mussel physiological parameters;
- Geochemical and physiological endpoints proved suitable for lagoon biomonitoring;

Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring

Authors:

Greggio Nicolas*^{a, b}, Capolupo Marco^{a, b}, Filippo Donnini^a, Manfred Birke^c, Fabbri Elena^{a, b}, Dinelli Enrico^{a, b}

^a University of Bologna, Inter-Departmental Research Centre for Environmental Science (CIRSA), Via S. Alberto 163, 48123 Ravenna, Italy

^b University of Bologna, Department of Biological, Geological, and Environmental Sciences, Piazza S. Donato 1, 40100 Bologna, Italy

^c Federal Institute for Geosciences and Natural Resources, Stilleweg 2, 30655, Hannover, Germany

* Corresponding Author: Nicolas Greggio e-mail: nicolas.greggio2@unibo.it.

University of Bologna, Inter-Departmental Research Centre for Environmental Science (CIRSA), Via S. Alberto 163, 48123 Ravenna, Italy.

Abstract:

Coastal lagoons are complex environments threatened by natural and anthropogenic stressors. Here, we tested the effectiveness of combining physical, geochemical and chemical measurements with biomarker data obtained in field-exposed marine mussels (*Mytilus galloprovincialis*) as a biomonitoring strategy for a highly pressured lagoon (Pialassa Baiona, Ravenna, Italy). Data showed a spatial trend of sediment contamination by Hg, Pt, Au, Ag, Mo, Re, Cd, Pd and Zn. Local conditions of high water temperature/low conductivity were detected among selected sites. After a 30-day in situ exposure, Ag and Hg were the most bioaccumulated elements (10 and 5 folds, respectively) in mussels followed by Sb, Al, Ti and Fe. Decreased survival, lysosomal dysfunctions, increased metallothionein content and peroxisome proliferation were observed in mussels in relation to metal spatial distribution and physico-chemical fluctuations. Overall, this study provides a further confirmation of the role of biomonitoring to reliably assess the environmental quality of highly pressured lagoons.

Keywords: Coastal lagoon; Sediment geochemistry; Metal bioaccumulation; Biomarkers; Biomonitoring;

Funding sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

1 Introduction

Being at the boundary between terrestrial and marine ecosystems coastal lagoons are, by definition, unstable and complex systems morphologically dependent and shaped by natural continental evolution (rivers sediment transport and discharge), sea level variations, coastal dynamics, storminess and local geologic processes (e.g. subsidence) (Bird, 1994; Newton et al., 2014). In natural conditions, coastal lagoons receive fresh and nutrient-rich inputs from the land-side, that combined with shallow, warm and relatively stable water, sustain a complex food chain (from algae, to macro benthos, fish and swamps birds), providing a multitude of life-supporting (Aliaume et al., 2007; Dominik et al., 2014; Petry et al., 2016).

The naturally complicated equilibrium of these environments has often been impacted by strong anthropic development in the last century, especially in those lagoons close to major cities and harbours. With the beginning of demographic boom and of the correlated industrial development, lagoons have been converted into commercial ports or industrial areas and in the worst case as processing water disposal sites (Guerra, 2012; Guerra et al., 2014).

Water and sediment geochemistry of lagoon environments have been widely explored in order to define water status (Mollema et al., 2013; Greggio et al., 2020), sediment provenance and local background concentrations (Migani et al., 2015; Borghesi et al., 2016; Greggio et al., 2018a) or levels of occurring pollutants, notably heavy metals (Nazneen et al., 2019; Saidi et al., 2019), organic compounds (León et al., 2017; Mattes et al., 2018) and emerging contaminants (Pignotti et al., 2017; Pignotti and Dinelli, 2018).

The geochemical characterization of sediments and water is important, but alone often does not allow for an accurate and reliable evaluation of the environmental status of lagoon ecosystems (Carvalho et al., 2014; Dahms, 2014; Moreno-González et al., 2015). Physico-chemical properties and geological background of lagoons can influence the response of exposed organisms to pollutants, inducing potential interactive effects due to changes in bioavailability and toxicity (Migani et al., 2015; Piggott et al., 2015; Borghesi et al., 2016). Chemical assessments fail in detecting biotransformation products, eventually more toxic than the parental compounds (Buryškova et al., 2006; Braga et al., 2018), and in estimating synergistic/antagonistic effects induced on the fitness of the exposed biota (Lari et al., 2017). For these considerations, multidisciplinary applications integrating geological/chemical parameters, pollutants bioaccumulation and biological effects may represent a promising approach to assess the environmental status of highly pressured coastal systems.

71 In the framework of biomonitoring programs of marine areas, particular attention has recently
1
22 been addressed to the assessment of biomarkers of environmental stress (Turja et al., 2014;
3
43 Capolupo et al., 2017; Mansour et al., 2020). Biomarkers are defined as indices of either exposure
5
674 or effect of pollutants induced at biochemical, cellular and physiological level on exposed organisms
7
75 (Viarengo et al., 2007a). They have successfully been used in association to bioaccumulation
9
1076 analyses to characterize “sub-organism” toxicological dynamics occurring in highly contaminated
11
1177 environments and provide crucial and early-warning information for planning urgent intervention
12
1378 or following the effects of remediation plans (Donnini et al., 2007; Capolupo et al., 2017). The
14
1579 simultaneous analysis of a wide set of biomarkers offers accurate information on the sub-lethal
16
1780 alterations induced by multiple stressors and may allow to correlate observed effects with specific
18
1981 classes of pollutants (Shaw et al., 2011; Turja et al., 2014). Moreover, biomarkers provide a unique
20
2182 contribution to determine the synergistic effect of pollutant mixtures, even when single
22
2383 contaminants occur at low concentrations. Biomarker-based assessments are thus currently
24
2584 promoted under the current EU legislation for monitoring surveys of marine and transitional
26
2785 environments (2000/60/EC; 2008/56/EC; Piló et al., 2017).

28
2986 The aim of this work was to evaluate the effectiveness of integrating water physico-chemical
30
3187 data, sediment geochemistry, metal bioaccumulation and biomarker measurements as a
32
3388 biomonitoring strategy for highly contaminated coastal lagoons. The study was conducted in a
34
3589 shallow and brackish basin located along the North-Western coast of the Adriatic Sea (Pialassa
36
3790 Baiona, Ravenna, Italy). In this area, previous works highlighted the presence of organic pollutants
38
3991 in sediments (Trombini et al., 2003; Vassura et al., 2005; Fabbri et al., 2006; Guerra, 2012; Guerra
40
4192 et al., 2014;) and their potential for being bioaccumulated by mussels (Fabbri et al., 2006) and clams
42
4393 (Vassura et al., 2005). Inorganic pollutants were less investigated and the current information is
44
4594 limited to few elements, as Cd, Cu, Hg, Ni, Pb and Zn (Trombini et al., 2003; Matteucci et al., 2005;
46
4795 Guerra et al., 2014). A recent hydrogeochemical study reported water physico-chemical alterations
48
4996 and the abundance of seldom analysed metals (Ag and Sb among others) in surface drainage water
50
5197 flowing into the investigated lagoon (Greggio et al., 2020). Therefore, a wide spectrum of elements
52
5398 was measured in sediments collected from different sites within the lagoon and in adult specimens
54
5599 of the Mediterranean mussel *Mytilus galloprovincialis* transplanted *in situ* at the same locations.
56
5790 Finally, a battery of biological endpoints was measured in mussels, including lysosomal responses,
58
5991 oxidative stress parameters and biomarkers of exposure to specific classes of pollutants.

102 **2 Materials and methods**


103 **2.1 Study area and sampling sites**

104 The Pialassa Baiona (Fig. 1) is a brackish lagoon located northward the town of Ravenna (Italy),
105 between 44°27'55.5"N and 44°31'31.3"N and 12°14'9.7"E and 12°16'10"E that cover an area of
106 about 10 km². The lagoon is part of a barrier-lagoon system that characterized the historical
107 evolution of the Adriatic Sea coastline in the area (Amorosi et al., 1999). It is formed by small,
108 shallow ponds (0.5–1 m depth) and deeper artificial channels (2–3 m depth) that permit the
109 incessant water exchange with the Adriatic Sea driven by tides. It is connected to the Adriatic Sea
110 only through the Candiano Channel that also is the Ravenna operative industrial harbour. The
111 Pialassa Baiona is mainly adopted by local citizens as recreational fishing and hunting site and few
112 clam farming activities exist in the northernmost portion.

113 The lagoon receives freshwaters from the west through three main channels (Fossatone,
114 Cerba and Cupa-Magni channels) which drain agricultural lands and collect effluents from industrial
115 and municipal treatment plants, located in the southernmost part (Fig. 1). The drainage water
116 entering the lagoon are often rich in nutrients and dissolved metals (Mollema et al., 2013; Greggio
117 et al., 2018b). The highest concentration of nutrients, as well as the highest bacteriological content,
118 are observed in the southern part of the lagoon, in proximity of the via Cupa channel (Soprani et al.,
119 1994). In addition, during the 60s and 70s, high quantities of chemical compounds were discharged
120 into the lagoon through the via Cupa and Magni channels, carrying wastewater from the industrial
121 area close to the southern border (Fabbri et al., 1998).

122 Previous investigations showed that high levels of metals, polycyclic aromatic hydrocarbons
123 (PAHs), polychlorinated biphenyls (PCBs) and other pollutants occurred in surface, subsurface and
124 suspended sediments with N-S increasing gradient (Trombini et al., 2003; Matteucci et al., 2005;
125 Guerra, 2012; Guerra et al., 2014). On these bases, seven sites were selected within the Pialassa
126 Baiona lagoon in order to cover its overall geochemical complexity and spatial trend of
127 anthropogenic pressures (Fig. 1). All sites (S1 to S7) were investigated for sediment geochemistry,
128 while metal bioaccumulation and biomarker responses were measured on mussels transplanted
129 into six sites (S1, S2, S3, S5, S6 and S7), since *in situ* transplantation into site S4 was not possible for
130 technical reasons. Mussels were exposed into the lagoon for 30 days, which is acknowledged as a
131 suitable time for sessile organisms to bioaccumulate contaminants at environmentally
132 representative levels and to develop physiological responses to “in site” pollution (Viarengo et al.,
133 2007b). Since previous investigations highlighted a lesser incidence of metals and organic

134 compounds in the north-western part of the lagoon (Trombini et al., 2003; Fabbri et al., 2006;
135 Donnini et al., 2007; Franzellitti et al., 2010), Site 7 (S7), located at about 0.5 km from the north-
136 western border, was defined as internal control site for both geochemical and biological
137 assessments.



538
539 **Figure 1. Location map and sampling points (S1 to S7) within the Pialassa Baiona lagoon (Ravenna, Italy).** The
540 abbreviations "R." and "Ch." refer to the main rivers and/or channels which supply freshwater to the lagoon. The
541 yellow shape represents the industrial area settled at the south-eastern boundary of the Pialassa Baiona lagoon.

143 2.2 *Field activities and measurements*

1
1244 In order to have a better characterization of the environmental conditions influencing the
3
1445 organisms' response, measurements for physico-chemical parameters were conducted at each site.
5
146 During mussel field exposure (from June 1st to June 30th) six vertical profiles (10 cm resolution) of
7
147 electrical conductivity, temperature and pH, using a CTD probe were conducted.

9
148
11
1249 2.3 *Sediment geochemistry*

13
1450 The bottom sediments were sampled at each site with a manual corer using a Plexiglas tube.
15
1461 After the extrusion, the sections 0 - 5 cm was retained for analysis, placed in polyethylene bottles,
17
1462 were stored at -25 °C. For geochemical analysis, sediment aliquots were dried at 60 °C and
19
1463 homogenized by grinding in agate mortar. On the bottom sediments total elements concentration
21
1464 by X-ray fluorescence spectrometry (XRF) and pseudo-total (mobile/available) concentrations by
23
1465 aqua regia digestion (AR) have been performed.

25
1466 The total metal content of 29 chemical elements (SiO₂, TiO₂, Al₂O₃, Fe₂O₃, MnO, MgO, CaO,
27
1467 Na₂O, K₂O, P₂O₅, S, As, Ba, Ce, Co, Cr, Cu, La, Nb, Ni, Pb, Rb, Sc, Sr, Th, V, Y, Zn, Zr) has been measured
29
1468 by XRF, using a Philips PW 1480 spectrometer equipped with a Rh tube on pressed powder pellets
31
1469 following matrix correction methods suggested by Franzini et al. (1972; 1975) Leoni and Saitta
33
1470 (1976) and Leoni et al. (1982). Thirty international reference materials were used for the instrument
35
1471 calibration, while accuracy was evaluated using 4 international reference samples, namely BR, BCR-
37
1472 1, TB and AGV-1 (Govindaraju, 1984). The average accuracy was higher than 5 % for trace-element
39
1473 determinations. In order to quantify the pseudo total concentration, 15 g of sediment were digested
41
1474 with a modified aqua regia solution of equal parts concentrated HCl, HNO₃ and deionized H₂O for
43
1475 one hour in a heating block of hot water bath at the Bureau Veritas Laboratories in Vancouver
45
1476 (Canada). Fifty-three chemical elements (Ag, Al, As, Au, B, Ba, Be, Bi, Ca, Cd, Ce, Co, Cr, Cs, Cu, Fe,
47
1477 Ga, Ge, Hf, Hg, In, K, La, Li, Mg, Mn, Mo, Na, Nb, Ni, P, Pb, Pd, Pt, Rb, Re, S, Sb, Sc, Se, Sn, Sr, Ta, Te,
49
1478 Th, Ti, Tl, U, V, W, Y, Zn, Zr) were analysed by Inductively Coupled Plasma Mass Spectrometry (ICP-
51
1479 MS).

53
1480 The volatile content (including humidity, organic matter, water incorporated in the lattice of
55
1481 clay minerals, carbon dioxide in the carbonate minerals) was evaluated by thermal analysis using a
57
1482 Setaram TAG24 double furnace apparatus, with simultaneous registration of thermogravimetric
59
1483 (TG), derivative thermogravimetric (DTG), differential thermal analysis (DTA). A CO₂ atmosphere

174 was used to increase the temperature of carbonate decomposition and the carbonate content was
175 estimated by quantifying the weight loss at temperature > 700 °C.

176

177 2.4 Mussel collection and handling

178 Adult specimens of the Mediterranean mussel (*M. galloprovincialis*) with a shell-length of 5 ±
179 1 cm were collected by professional fishermen of the “Cooperativa CONISUB” (Marina di Ravenna,
180 Italy) along the north-western Adriatic Sea coast in high quality marine areas (Zone A) and
181 immediately transferred to the laboratory in seawater tanks with continuous aeration. Prior to the
182 *in situ* deployment, all animals were acclimatized for 6 days in aquaria containing 60 L of aerated
183 natural seawater (at a ratio of 1 mussel/L) at 16 °C and natural photoperiod, and fed daily on an
184 algal slurry (Coral Diet Filtrator, Xaquia, Italy) after water renewal. Seventy organisms were dissected
185 prior to the *in situ* transplantations (time 0, T0) for bioaccumulation and biological endpoint
186 analyses. Digestive glands and gills were immediately collected, frozen in liquid nitrogen and stored
187 at –80 °C until analysed. At the same time six groups of seventy organisms each were put in fishnet
188 bags, fixed to piles and driven into the sediment at sites S1, S2, S3, S5, S6 and S7 (Fig. 1) at about 50
189 cm from the bottom to ensure continuous submersion during the whole exposure period. Deployed
190 mussels were regularly controlled at three days intervals and dead mussels were eventually
191 recorded for survival assessment. Since previous investigations identified the Magni channel as a
192 heavily polluted site (Fabbri et al., 2000; 2003; Matteucci et al., 2005), three additional fishnet bags,
193 with seventy organisms each, were transplanted into S1 to execute biomarkers with shorter
194 timestep.

195

196 2.5 Metal bioaccumulation analysis

197 Metal bioaccumulation was evaluated on mussels prior to the *in situ* deployment (Time 0, T0)
198 and after 30 days of field exposure into the lagoon. Metal bioaccumulation from each site was used
199 to calculate the bioaccumulation factor as enrichment/depletion compared to T0. Samples (0.2 g
200 each) were digested in a microwave system with ultrapure HNO₃. The resulting fluid was diluted by
201 adding deionized water to a final volume of 30 mL, at the Department of Environmental Science of
202 the University of Siena. Analytical blanks and certified reference material were included in the
203 analysis, specifically the mussel tissue ERM-CE278 (*Mytilus edulis*) from the Institute for Reference
204 Materials and Measurements, Geel, Belgium.

205

206

207

208

205 The samples were analysed by inductively coupled plasma quadrupole mass spectrometry
206 (ICP-QMS) using an Agilent 7500ce instrument at the laboratories of the Federal Institute for
207 Geosciences and Natural Resources in Berlin. Atomic fluorescence spectrometry was used for the
208 determination of mercury (Instrument PSA 10.035 Millennium Merlin 1631). The Table S1 reports
209 the results for the investigated elements in the certified reference material. Additional details can
210 be found in Capolupo et al. (2017).

211
212

213 2.6 *Biological endpoints*

214 2.6.1 Survival percentage and Stress on Stress

215 The survival (SURV) was calculated as the percentage of mussels found alive over the exposure
216 period with respect to the initial number of individuals (70 at each site).

217 For Stress on Stress (SOS) assessment, immediately after collection, 15 mussels per site were
218 subjected to anoxia by air exposure at 15 °C in a humidified chamber. Death symptoms were open
219 valves and absence of muscular activity. The survival was assessed daily, and results expressed as
220 the time at which 100% mortality was achieved (Viarengo et al., 2007a).

221
222

223 2.6.2 Cytochemical parameters

224 Lysosomal membrane stability (LMS), lysosome to cytoplasm volume ratio (LYS/CYT) and
225 lipofuscin content (LIF) were assessed in the digestive gland from 20 mussels per each site (N = 20).
226 After dissection, glands were mounted on specific aluminium supports for cryostat microtomy
227 (chucks), frozen in N-hexane at -70 °C, and stored at -80 °C. All assessments were performed
228 according to the UNEP/RAMOGGE manual (1999) on 10 µm-thick gland sections obtained using a
229 cryostat (MICROM HM 505 N) at -30 °C. The LMS was measured as the time necessary to produce
230 the maximum staining reaction between the lysosomal hydrolase N-acetyl-β-hexosaminidase and a
231 specific substrate (Capolupo et al., 2016). Briefly, sections were incubated at 37 °C in a shaking water
232 bath with a destabilization buffer (0.1 M Na-citrate, pH 4.5) for different times (0, 3, 5, 10, 15, 20,
233 30 and 40 min), exposed to the substrate naphtol AS-BI N-acetyl-β-D-glucosaminide (Sigma, N4006)
234 for 20 min and finally stained with the diazonium dye Fast Violet B (Sigma, F1631) (1 mg/ml in 0.1
235 M phosphate buffer, pH 7.4). For each site, slides not exposed to the destabilization buffer (0 min)
236 were screened for the LYS/CYT assessment, which was determined by evaluating the cytoplasmic
237 and lysosomal areas according to Capolupo et al. (2017). The LIF determination was performed

238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265

236 according to Martin-Diaz et al. (2009). Gland sections were fixed in calcium formol and immersed
1
237 for 5 min in an aqueous solution of 1 % ferric chloride and 1 % potassium ferricyanide in a 3:1 ratio.

238 For all lysosomal parameters, tissue sections were assessed under a light microscope
3
239 (Axioskop 40, Carl Zeiss, Milan, Italy) equipped with a 40X objective and a digital camera (AxioCam
5
240 MRc, Carl Zeiss, Milan, Italy) as reported by Capolupo et al. (2016). Results were expressed as
7
241 labilization period for LMS, percentage of lysosomal volume inside digestive cells for LYS/CYT and
9
242 arbitrary units for LIF.

13 14 15 2.6.3 Enzymatic biomarkers 16

17 The acetylcholinesterase activity (AChE) activity was determined in mussel gills as previously
18
19 reported (Valbonesi et al., 2003). Gills from 5 mussels per site (N = 5) were homogenized in a 0.1 M
20
21 phosphate buffer, pH 7.4, and centrifuged at 9000 ×g at 4 °C for 30 min. Supernatant aliquots were
22
23 incubated at 25 °C with 0.5 mM acetylthiocholine iodide (ASCh) and 0.33 mM DTNB (Ellman et al.,
24
25 1961), and read at 405 nm for 10 min. Results are reported as nmoles/min/mg protein. The sample
26
27 protein concentration was measured according to Lowry et al. (1951).

28
29 For catalase (CAT) assessment, about 50 mg of digestive gland was dissected from 5 mussels
30
31 per site (N = 5), homogenized in 50 mM potassium-phosphate buffer (pH 7.0, 0.5 mM Na₂EDTA) and
32
33 centrifuged at 15,000 g at 4 °C. The CAT activity was determined by measuring the time-dependent
34
35 decrease of absorbance at 240 nm in the presence of 55 mM H₂O₂. Data were expressed as
36
37 μmol/min/mg of total protein.

38 The Peroxisomal acyl-CoA oxidase (AOX) activity was determined on five pools of 2 digestive
39
40 glands per site (N = 5) according to Orbea et al. (2006). Tissues were homogenized in four volumes
41
42 of TVBE buffer (1 mM sodium bicarbonate, 1 mM EDTA, 0.1 % ethanol and 0.01 % Triton X-100), pH
43
44 7.6 and centrifuged at 500 × g for 15 min at 4 °C. Supernatants were diluted 1:10 in TVBE buffer and
45
46 spectrophotometrically assayed for AOX activity at a wavelength of 502 nm. Data were expressed
47
48 as milliunits/mg of AOX protein (equivalent to nmol H₂O₂/min x mg protein).

50 51 52 2.6.4 Metallothionein content (MT) 53

54 The metallothionein (MT) content was analysed on five pools of 5 mussel digestive glands
55
56 (about 1.5 g of tissue) per site according to Viarengo et al. (1997). Final absorbance was measured
57
58 at 410 nm, using reduced glutathione as reference standard. Data were expressed as μg of MT/mg
59
60 tissue.

268
1
269
3
270
5
271
7
272
9
273
11
274
13
275
14
276
15
277
18
278
20
279
22
280
24
281
26
282
28
283
30
284
32
285
34
286
36
287
38
288
40
289
42
290
44
291
46
292
48
293
50
294
52
295
54
296
56
57
58
59
60
61
62
63
64
65

2.6.5 Mussel Expert System

Data from biological responses were integrated using the Mussel Expert System (MES) developed by Dagnino et al. (2007) in a unique and synoptic five-level Health Status Index (HSI) ranging from A (healthy) to E (pathological status). The system classification is based on (i) the number of biomarkers altered in the exposed organisms (significant changes greater than 20 % were utilized to evaluate alterations in the health status of the organisms); (ii) the magnitude of the alteration; (iii) the role of the autophagic process and (iv) the level of biological organization affected (i.e., cell, tissue and organism level). For each biomarker/site, the system calculates an Alteration Factor (AF), which is the ratio between biomarker levels from treatments and control, and an Alteration Level (AL), which defines the magnitude of observed alterations based on the AFs magnitude and the biomarker toxicological profile (i.e. increasing, decreasing, bell-shaped). ALs and AFs are then processed by the system applying a set of rules in the “if...then” form in order to calculate the overall HSI associated to the exposure to each site. The simultaneous alterations of general stress parameters (i.e. lysosomal responses) and biomarkers of exposure to specific classes of compounds (e.g. MT for metals; AOX for POPs) is further computed and eventually recognized by the system as an early warning signal regarding the presence or excess of those pollutants in the monitored sites or tested condition.

2.7 Data analysis

All data were tested for normal distribution using the Shapiro-Wilk test, while the Levene's test was employed to test the equal variance. Significant differences between S1 to S6 compared to the control site S7 were estimated by a One-Way ANOVA (SigmaStat, SPSS) followed by the Duncan's post-hoc test. The evaluation of potential linear relationships between biological endpoints and physico-chemical parameters (including pH, temperature, electrical conductivity, metal concentrations in sediments and mussel tissues) were conducted through the Pearson's correlation analysis. For all performed tests, significance was established at $p < 0.05$. All analyses were performed using the software SigmaPlot 13 (Systat Software Inc. San Jose, CA, USA).

297 **3 Results and discussion**

1
298 **3.1 Physico-chemical data**

299 The vertical profiles of temperature, pH and electrical conductivity (EC) measured during
300 mussels field exposure as well as a graphical representation of the temperature, pH and EC vertical
301 profiles for June 14th are presented respectively in Table S2 and in Figure S1 of supplementary
302 material. The site S1, located in the Magni Channel at the southern side of the lagoon (Fig. 1), is
303 different from the other sites for all monitored parameters. It shows higher water column
304 temperature, reduced EC and a lower pH value, for the entire monitoring period (Table S2).
305 Specifically, S1 shows the lowest EC among the sites with a marked vertical gradient (Fig. S1A) and
306 it is also the site with the highest temperature, up to 32.3 °C. Moreover, it records the highest
307 average profile temperature during the surveying period, at least 2 °C constantly higher compared
308 to all the other sites (Fig. S1B). Site S1 receives freshwater from artificial drainage channel network
309 surrounding the town of Ravenna (Greggio et al., 2018b). Previous investigations also reported
310 water temperature fluctuation wider (10-24 °C) in the southernmost channel with respect to the
311 lagoon average (16-18 °C), which was subjected to variations of about 10 °C within the 24 hours
312 owing to discharges of cooling water from the nearby industrial site (Franzellitti et al., 2010). A
313 thermal vertical gradient is also present in S5 probably associated to local limited water circulation
314 favouring water warming up (Fig. S1A). The pH is the lowest in S1, ranging from 7.2 to 7.8, and with
315 a marked vertical profile. In all the other sites the pH is instead similar and fairly constant around 8.
316 Only at S5 the observed pH values are higher (8.0-8.3) for the entire surveying period (Table S2).

317
318 **3.2 Sediment data**

319 The results of the geochemical analyses are presented in Table 1. According to the
320 geochemical composition of the major elements (expressed as %), and referencing to the work of
321 Dinelli et al. (2007) which compared geochemical proxies with grain size analysing borehole samples
322 in the same geographic area, the Pialassa Baiona bed sediment samples are classified as fine
323 grained-sediments (with a recalculated SiO₂/Al₂O₃ ratio < 3) enriched in carbonates. The sediments
324 are also rich in organic matter (OM in Table 1, ranging from 1.1 to 3.4) as already reported for
325 adjacent coastal lagoons (Migani et al., 2015). The bulk geochemical composition is quite
326 homogeneous among the different sites, with S1 being slightly enriched in carbonates (Table 1).

327 The geochemical results, in terms of total sediment composition, are comparable with those
 328 reported by Donnini et al. (2007) although the number of sites was different. As far as total
 329 concentrations are concerned, these are within the range of regional background, as proposed by
 330 Migani et al. (2015), taking into consideration Holocene data from boreholes in the area not affected
 331 by anthropogenic inputs (Amorosi et al., 2002; Curzi et al., 2006; Dinelli et al., 2012, Greggio et al.,
 332 2018a). Except for sites S6 and S7, Zn and Cu concentrations exceed the maximum value observed
 333 in the background sediments, respectively of 124 and 41 mg/kg (Table 1). In particular, for S1 site
 334 Zn and Cu total concentrations are double respect to background, respectively 316 and 87 mg/kg,
 335 in line with concentration detected in the neighbouring Pialassa Piomboni by Pignotti et al. (2018).
 336 Pb concentrations in S3 and S5 are 30 mg/kg being slightly higher respect to background level
 337 established at 29 mg/kg by Migani et al. (2015), but certainly lower than Pb concentration in the
 338 Pialassa Piomboni by Pignotti et al. (2018) where a maximum concentration of 257 mg/kg was
 339 measured.

340
 341 **Table 1. Results of the chemical analyses on sediments. The table includes either pseudo total Aqua Regia (AR) and**
 342 **total XRF data. Table includes also the ranges observed in boreholes in the area, considered representative of the**
 343 **natural background (Amorosi et al., 2002; Curzi et al., 2006). Bold elements exceed natural background. AR DL**
 344 **indicates Detection Limit for Aqua Regia extraction.**

		S1		S2		S3		S4		S5		S6		S7		Natural background	
		AR DL	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	
34	Ag µg/kg	2	991	-	269	-	268	-	173	-	275	-	179	-	62	-	-
35	Al %	0.01	1.39	6.31	1.41	7.02	1.23	6.98	1.10	6.97	1.35	7.09	1.39	7.07	1.41	6.55	4.4-9.7
36	As mg/kg	0.1	7.8	-	5.9	-	6.1	-	5.9	-	7.8	-	6.8	-	6.7	-	-
37	Au µg/kg	0.2	16.2	-	5.4	-	11.1	-	2.6	-	5.1	-	3.4	-	1	-	-
38	B mg/kg	1	25	-	28	-	26	-	24	-	38	-	33	-	22	-	-
39	Ba mg/kg	0.5	61.9	263	43.7	218	41.8	260	42.6	322	40.1	258	38.7	294	36.5	258	220-530
40	Be mg/kg	0.1	0.4	-	0.7	-	0.7	-	0.8	-	0.5	-	0.6	-	0.6	-	-
41	Bi mg/kg	0.02	0.6	-	0.38	-	0.35	-	0.26	-	0.38	-	0.31	-	0.28	-	-
42	Cd %	0.01	11.27	11.37	7.45	7.51	7.87	7.96	7.34	7.87	6.10	6.65	7.24	7.43	8.61	9.52	0.3-17.2
43	Ce mg/kg	0.01	0.68	-	0.29	-	0.33	-	0.21	-	0.5	-	0.35	-	0.11	-	-
44	Ce mg/kg	0.1	19.4	42	17	55	17.6	71	16.8	52	17.8	64	18.3	42	20.1	58	14-83
45	Co mg/kg	0.1	10.2	12	10.9	11	10.7	12	9.9	11	11.7	14	11.5	12	10.6	11	5-29
46	Cr mg/kg	0.5	70.6	131	57.5	103	51.4	132	47.7	131	59.9	138	59.3	130	47	100	96-270
47	Cs mg/kg	0.02	1.36	-	1.15	-	1.08	-	1.16	-	1.11	-	1.39	-	1.57	-	-
48	Cu mg/kg	0.01	84.1	87	49.1	52	49.5	53	36.7	44	51.1	52	37.5	38	22.4	24	4-41
49	Fe %	0.01	2.17	3.42	2.46	3.92	2.23	3.86	2.00	3.69	2.54	3.83	2.30	3.64	2.36	3.72	2.1-5.3
50	Ga mg/kg	0.1	4.4	-	4.2	-	4	-	3.9	-	4	-	4.2	-	4.4	-	-
51	Ge mg/kg	0.1	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	-
52	Hf mg/kg	0.02	0.03	-	0.03	-	0.04	-	0.03	-	0.03	-	0.03	-	0.05	-	-
53	Hg mg/kg	0.005	25.7	-	1.42	-	1.48	-	1.89	-	2.38	-	2.26	-	0.1	-	-
54	In mg/kg	0.02	0.03	-	0.03	-	0.02	-	0.02	-	0.03	-	0.02	-	0.03	-	-
55	K %	0.01	0.33	1.77	0.36	1.98	0.31	2.06	0.28	2.04	0.39	1.97	0.36	2.04	0.35	2.03	0.9-3.3
56	La mg/kg	0.5	8.3	19	7.7	26	7.3	22	6.9	18	8	20	8.5	34	9.5	20	17-47
57	Li mg/kg	0.1	26.1	-	30.6	-	27.8	-	24.9	-	28.7	-	28	-	28.1	-	-
58	Mg %	0.01	1.37	2.41	1.34	2.38	1.30	2.38	1.22	2.44	1.42	2.54	1.55	2.55	1.58	2.35	2.2-4.2
59	Mn mg/kg	1	531	893	527	865	549	896	513	949	511	801	550	897	644	1396	230-1470
60	Mo mg/kg	0.01	2.67	-	3.33	-	2.61	-	2.28	-	3.34	-	1.57	-	0.54	-	-
61	Na %	0.001	0.85	0.68	0.98	0.69	0.90	0.75	0.65	0.99	1.18	0.67	0.85	0.78	0.42	0.85	0.5-2.1
62	Nb mg/kg	0.02	0.37	15	0.28	13	0.25	13	0.29	14	0.34	16	0.37	10	0.29	13	8-20
63	Ni mg/kg	0.1	50.9	73	56.5	73	50.7	88	46.3	72	55.3	82	51.5	74	46.4	57	39-148
64	P %	0.001	0.11	0.10	0.06	0.06	0.06	0.07	0.06	0.06	0.08	0.07	0.07	0.06	0.06	0.06	0.03-0.08
65	Pb mg/kg	0.01	27.92	28	22.71	26	22.14	30	18.16	19	26.22	30	20.71	22	11.99	18	5-29

1	Pd	µg/kg	10	46	-	10	-	12	-	11	-	10	-	<10	-	<10	-	-
2	Pt	µg/kg	2	37	-	6	-	4	-	4	-	4	-	<2	-	<2	-	-
3	Rb	mg/kg	0.1	24.9	113	22.2	128	20.8	128	22.4	113	22.3	130	25.1	114	29.8	89	56-200
4	Re	µg/kg	1	7	-	1	-	2	-	<1	-	2	-	2	-	<1	-	-
5	S	%	0.02	0.6	-	0.55	-	0.6	-	0.54	-	0.72	-	0.71	-	0.34	-	-
6	Sb	mg/kg	0.02	0.51	-	0.29	-	0.25	-	0.19	-	0.37	-	0.29	-	0.19	-	-
7	Sc	mg/kg	0.1	3.4	11	3.6	11	3.4	6	3.2	8	3.7	10	3.7	8	4.1	13	12-21
8	Se	mg/kg	0.1	0.9	-	0.7	-	0.7	-	0.5	-	1.1	-	0.7	-	0.5	-	-
9	Si	%	-	-	17.72	-	19.43	-	19.90	-	20.61	-	19.31	-	20.95	-	20.03	14.7-29.3
10	Sn	mg/kg	0.1	3.2	-	1.8	-	1.5	-	1.3	-	1.9	-	1.5	-	0.8	-	-
11	Sr	mg/kg	0.5	353	368	274.4	300	224.8	252	218	265	259.1	281	281.6	291	290.5	310	113-442
12	Th	mg/kg	0.1	3.5	4	4.1	5	3.6	17	3.6	25	4	11	4.2	10	4.7	7	<3-19
13	Ti	%	0.001	0.009	0.308	0.005	0.328	0.006	0.347	0.008	0.349	0.006	0.324	0.010	0.334	0.015	0.336	0.02-0.58
14	Tl	mg/kg	0.02	0.26	-	0.17	-	0.21	-	0.19	-	0.27	-	0.2	-	0.14	-	-
15	U	mg/kg	0.1	1.3	-	1.5	-	1.2	-	1.1	-	1.3	-	1.2	-	1.1	-	-
16	V	mg/kg	2	32	85	33	85	30	99	26	94	35	113	33	94	31	76	54-169
17	Y	mg/kg	0.01	10.02	18	9.39	23	9.28	24	8.86	20	9.7	24	9.63	19	10.58	19	18-43
18	Zn	mg/kg	0.1	306.6	316	129.8	152	133.1	154	101.2	126	146.9	174	109.3	124	58.2	67	43-124
19	Zr	mg/kg	0.1	1.1	108	1.2	91	1	105	1	124	1.1	86	1.2	92	1.9	106	76-260
20	H₂O⁻	%	-	-	2.7	-	4.0	-	2.8	-	2.4	-	3.7	-	2.7	-	2.1	-
21	OM	%	-	-	2.8	-	1.8	-	2.0	-	1.7	-	3.4	-	2.2	-	1.1	-
22	H₂O⁺	%	-	-	6.0	-	6.5	-	5.4	-	4.4	-	6.5	-	5.8	-	2.3	-
23	CO₂	%	-	-	11.6	-	8.9	-	9.1	-	9.3	-	8.6	-	7.4	-	12.3	-
24	LOI	%	-	-	23.1	-	21.2	-	19.3	-	17.8	-	22.3	-	18.1	-	17.8	4.8-29.3

As regards the Aqua Regia extraction, data showed the presence of elements seldom analysed in sediments of the area. The most striking features are the differences observed between S1 and all the other sites for a number of elements, in particular Ag, As, Au, Bi, Cd, Cr, Cu, Hg, Pb, Pd, Pt, Re, Zn (Table 1). According to Donnini et al. (2007), S7 should be considered as unpolluted site and if its results are taken as reference, a strong enrichment is clear for several elements (Fig. 2). Mercury, which is a well-known pollutant in the area (Miserocchi et al., 1993; Fabbri et al., 1998; Fabbri et al., 2001a, b; Trombini et al., 2003; Matteucci et al., 2005; Guerra et al., 2007; Covelli et al., 2011; Guerra, 2012; Dominik et al., 2014; Borghesi et al., 2016), is the element with the largest enrichment (up to 250 times compared to S7) and reaches relatively high concentrations in the Magni Channel site (S1, up to 25 mg/kg). Mercury dynamics in the lagoon is complex and although the maxima in concentrations are not at the surface, the element is heavily mobilized at the water/sediment interface in the southern part of the lagoon (Covelli et al., 2011). Borghesi et al. (2016) highlighted similar behaviour among Hg and Ag, Au, Cd, Cu and Zn for six different coastal lagoons in the same Region due to metal affinity for the fine fraction of the sediment.

Notable enrichments are then evident in Figure 2 for valuable metals like Pt, Au, Ag, Re and Pd, which showed substantial increase (up to 10-fold) compared to S7 and to median values reported by Cicchella et al. (2015), that represent a useful reference for metal concentrations in mineral matrix obtained with the same analytical technique and at the same laboratory. In particular, Platinum and Palladium are of recent technogenic use and their anthropogenic fluxes greatly exceed the natural ones (Ruchter and Surer, 2015; McGillicuddy et al., 2017). Platinum and Palladium are largely used in automobile catalytic converters (Bossi and Gediga, 2017) and their

367 dispersion increases particularly around urban areas in roadside soils (Ravindra et al., 2004; Ruchter
368 and Surer, 2015; Leopold et al., 2018; Zuzolo et al., 2018), but also in marine areas close to urban
369 settings (Cobelo-García et al., 2011; Abdou et al., 2016).

370 The Silver concentration reaches a peak at S1 with 0.99 mg/kg, at least four times more than
371 the other sites in the Pialassa Baiona and ten times more than other coastal lagoons in the same
372 Region as resulted by Borghesi et al. (2016). The Ag enrichment is controlled by many factors such
373 as pH, OM, as well as the source material. At typical environmental pH values, Ag is adsorbed onto
374 Fe hydroxides, in preference to Cu and Zn (Lottermoser et al., 1999). In the near Venice lagoon,
375 Giusti and Zang (2002) measured Ag sediment concentrations 5-6 fold higher than this study,
376 justifying the abundance with both famous glassmaking firms in Murano island and the large volume
377 of untreated sewage sludges discharged into the lagoon from the nearby Marghera industrial area.

378 Moderate enrichments (four times compared to S7) are shown by Mo, Cd, Zn, Sn and Cu that
379 can be more directly linked to industrial effluents as reported by Giusti and Zang (2002) in Venice
380 lagoon and Borghesi et al. (2016) for the same Region. Enrichments in all these elements also
381 characterize the other sites, although at an order of magnitude lower than S1 (around 2 times).
382 Many of the metals detected in sediments show a decreasing concentration with increasing distance
383 from the industrial area located in the southern part of the lagoon. This suggests that industrial
384 discharge effluents play a major role in the lagoon contamination compared to effluents from urban
385 and agricultural activities. In facts, the distribution of some elements, including Zn, Cr, Cu, Mo, Ag,
386 and Hg, follow trends similar to those reported for other organic pollutants of industrial origin in
387 the same area (Vassura et al., 2005; Fabbri et al., 2006). Vassura et al. (2005), for instance, detected
388 a clear South-North increasing pattern of prioritized PAHs, with sediments from the Magni channel
389 area (the present S1) showing concentrations of pyrene, fluoranthene, anthracene and
390 phenanthrene up to 10,000 times higher compared to the northern sites.

391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465




Figure 2. Metal enrichment in sediments from selected sites within the Pialassa Baiona lagoon. Data are normalized on Site 7 (S7), which was selected as internal reference based on previous literature (Trombini et al., 2003; Fabbri et al., 2006; Donnini et al., 2007; Franzellitti et al., 2010). Elements in italics (*Pt, Au, Re, Pd*) were not investigated in the mussel tissues.

3.3 Metal bioaccumulation

Results of the chemical analyses on mussel tissues are presented in Table 2. Data were obtained for all sites except S1, which was characterized by a high mortality rate (more details in section 3.4). T0 data, originating from mussels growing in the Adriatic Sea, are in line with basal level for As, Ba, Cd, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, V, Zn identified by Fattorini et al. (2008). The trend of concentrations of trace elements in soft tissue of *Mytilus galloprovincialis* belonging to T0 is (Zn, Fe) > (As, Mn, Cu, Ni, Pb) > (Cd, Cr, Ag) > (Hg, Sb). The trend is confirmed also for average trace elements concentration among all sites: (Fe, Zn) > (As, Mn, Cu) > (Pb, Ag, Ni, Cd, Cr, Hg) > Sb. The only notable change is the Ag enrichment from 0.12 mg/kg in T0 tissues to a mean concentration of 0.80 mg/kg in tissues from experimental sites. Although Cd, Hg, Sb were not included, the same abundance trend was found by Giusti and Zang (2002) in the Venice Lagoon: Fe > Zn > (Cu, Mn, As) > Ag > Pb > (Ni, Cr). Absolute concentrations of Zn, Cu, Pb and Cd found in this study (Table 2) fall in the lowest part of the ranges summarized by Benali et al. (2017) for 10 locations in Mediterranean Sea.

Table 2. Results of the chemical analyses on mussel tissues (bioaccumulation). Data refers to T0 and to the 5 sites where mussels survived or remained after the exposure period.

	unit	T0	S2	S3	S5	S6	S7
Ag	mg/kg	0.12	0.99	0.98	1.21	0.62	0.82
Al	mg/kg	36	39	56	98	82	47
As	mg/kg	9.83	12.8	9.98	11.9	12.4	11.4
B	mg/kg	25.7	22.9	19.7	22.8	25.9	19.4

	Ba	mg/kg	1.95	0.35	0.49	0.54	0.52	0.36
1	Be	mg/kg	0.003	0.003	0.003	0.005	0.006	0.004
2	Bi	mg/kg	0.057	0.009	0.009	0.007	0.011	0.008
3	Ca	mg/kg	2004	1322	1299	1726	1679	1157
4	Cd	mg/kg	0.70	0.71	0.78	0.76	0.77	0.50
5	Ce	mg/kg	0.118	0.051	0.092	0.124	0.166	0.075
6	Co	mg/kg	0.54	0.323	0.3	0.303	0.511	0.315
7	Cr	mg/kg	0.57	0.6	0.53	0.67	0.53	0.64
8	Cs	mg/kg	0.022	0.014	0.016	0.022	0.019	0.015
9	Cu	mg/kg	4.65	2.71	2.63	2.81	2.66	1.46
10	Fe	mg/kg	67.5	68.1	77.4	116.0	103.4	91.7
11	Ga	mg/kg	0.018	0.007	0.011	0.019	0.013	0.009
12	Hg	mg/kg	0.082	0.35	0.32	0.41	0.27	0.16
13	K	mg/kg	11084	10988	9729	11129	11244	9535
14	La	mg/kg	0.138	0.035	0.057	0.074	0.109	0.06
15	Li	mg/kg	0.88	0.46	0.41	0.51	0.58	0.34
16	Mg	mg/kg	4198	3851	3467	3676	4422	2506
17	Mn	mg/kg	4.84	5.19	7.50	8.16	8.03	5.03
18	Mo	mg/kg	0.645	0.353	0.368	0.317	0.394	0.274
19	Na	mg/kg	22885	20201	15986	19714	24751	13438
20	Nb	mg/kg	0.002	0.001	0.002	0.003	0.003	0.002
21	Ni	mg/kg	3.61	0.72	0.69	0.73	0.92	0.64
22	Pb	mg/kg	1.41	0.87	0.52	0.57	0.94	0.64
23	Rb	mg/kg	5.29	4.08	4.32	4.27	4.3	3.78
24	Sb	mg/kg	0.018	0.038	0.031	0.03	0.048	0.039
25	Sc	mg/kg	0.04	0.04	0.03	0.05	0.07	0.04
26	Se	mg/kg	2.27	2.10	2.10	2.01	1.83	1.52
27	Sn	mg/kg	0.12	0.12	0.12	0.10	0.09	0.08
28	Sr	mg/kg	31.2	17.7	16.5	20.6	22.8	14.3
29	Th	mg/kg	0.019	0.011	0.014	0.025	0.031	0.016
30	Ti	mg/kg	0.49	0.34	0.62	1.07	0.88	0.50
31	Tl	mg/kg	0.003	0.001	0.002	0.002	0.001	0.001
32	U	mg/kg	0.127	0.049	0.040	0.054	0.065	0.032
33	V	mg/kg	1.15	0.44	0.49	0.55	0.80	0.47
34	Y	mg/kg	0.080	0.046	0.071	0.085	0.094	0.04
35	Zn	mg/kg	107.1	116	62.4	95.6	134	86.9
36	Zr	mg/kg	0.064	0.013	0.025	0.025	0.021	0.018

36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The bioaccumulation factor is graphically showed in Figure 3. Enrichments are notable for Ag, Hg and Sb in all the sites, with higher bioaccumulation at S5 located in the centre of the lagoon in a pond close to the outlet of a canal from the southern part of the lagoon, the most contaminated one. Although not high as those measured in mussels from other sites such as Northern England (Giusti et al., 1999), these three elements could be the most critical since other elements known for toxicity to mussels (e.g. Cd, Pb, Ni) (Benali et al. 2017; Yigit et al. 2017) are similar or depleted compared to T0 (Fig. 3).


The enrichment in mercury was relatively high, as expected and already verified both in living clams (Trombini et al., 2003) and mussels (Fabbri et al., 2006; Donnini et al., 2007). Moreover, the Hg cycling at the sediment-water interface (Covelli et al., 2011), could be a reason for the mercury enrichment observed in all sediment samples and may also reflect the high Hg concentrations observed in mussels exposed at S5 (Fig. 3).

428 Silver has recently attracted much interest since its increasing application as bactericidal and
429 usage in numerous consumer products lead to contamination of aquatic ecosystems, often in form
430 of Ag nanoparticles (Yuan et al., 2018; Outa et al., 2020). Silver speciation in environment is
431 dependent by local environmental conditions and usually forms stable complexes with Lewis bases
432 such as amines, halides, and thiolates and with dissolved organic matter altering its toxicity
433 (Mousavi et al., 2015). In Pialassa Baiona, Greggio et al. (2020) showed dissolved concentrations
434 ranging from 0.02 to 10 µg/L, with higher concentrations in basins with drainage water input.
435 Concerning bioaccumulation, recent works reported the effect of Ag nanoparticles on soft tissues
436 of mussels. *Mytilus galloprovincialis* accumulated Ag in digestive gland tubule cells and gills during
437 the entire year (Duroudier et al., 2019) and intralysosomal metal accumulation and lysosomal
438 membrane destabilisation were observed (Jimeno-Romero et al., 2017). Compare to this study (max
439 1.21 mg/kg dry wt), Giusti and Zang (2002) in Venice lagoon found silver concentrations up to 6.2
440 mg/kg dry wt that are aligned with outcomes in other species of the genus *Mytilus* as reported by
441 Martin et al. (1988).

442 Antimony is pervasive in synthetic polymers (including textiles, plastics and rubbers) (Filella et
443 al., 2019) and it is widely applied in alloy industries as flame retardant, as pigments for colour or
444 colour protection (Filella et al., 2002; James and Turner 2020). Although rock weathering and soil
445 runoff supply Sb to the environment, anthropogenic source is crucial. James and Turner (2020)
446 demonstrated that Sb could also be mobilised via digestion and bioturbation by deposit-feeders in
447 sediment contaminated by plastic or directly adsorbed from microplastics taken up by mussels (Van
448 Cauwenberghe et al., 2015). Antimony concentrations in unpolluted waters are less than 1 µg/L
449 (Filella et al., 2002). Greggio et al. (2020) reported Sb water concentrations in western ponds of the
450 Pialassa Baiona ranging from 0.05 to 0.5 µg/L. Limited is the knowledge of Antimony concentration
451 in mussels. In this study Sb in mussel tissues ranged from 0.01 to 0.05 mg/kg (Table 2) with a 2-3 as
452 enrichment factor (Fig. 3). De Gregori et al. (2007) found tissue concentration ranging from 0.007 to
453 0.06 mg/kg in unpolluted area, while 0.23 to 1.030 mg/kg in harbour dock protected by antimony-
454 lead alloy cover. Garcia (2015) measured Sb concentration ranging from 0.003 to 0.023 mg/kg in the
455 tissues of *Corbicula fluminea* collected in a river affected by a decommissioned Antimony smelter.

456 It is worth noting the depletion relative to T0 of elements like Cu, Pb and Ni, (Fig. 3 and Table
457 2). This might suggest that these elements are not strongly mobilized in the lagoon environment,
458 although slight differences can be recognized between S7 and the other sites. This is in contrast with
459 results by Pignotti et al. (2018) that using sequential extraction, defined Cu and Pb as the most

460 mobile elements in sediments (~70 % and ~80 %, of the total, respectively) in the neighbouring
 461 Pialassa Piomboni. In agreement with Pignotti et al. (2018) is the Ni behaviour here characterised
 462 by low mobility and predominantly bound to the residual fraction of the sediment, suggesting a
 463 lithogenic origin.



365 **Figure 3. Bioaccumulation factor of metals in mussel tissues.** The analysis has been performed on the tissue of mussels
 366 deployed for 30 days into different sites of the Pialassa Baiona lagoon compared to mussels analysed before the *in situ*
 367 deployment (T0). Elements are ranked according to average enrichment.

471 3.4 Biological effects on mussels

472 The mussel 30-day exposure within the Pialassa Baiona lagoon resulted in significant
 473 alterations of biological endpoints measured at different levels of the hierarchical ladder (from cell
 474 to organism) (Fig. 4). Among the investigated sites, adverse conditions for the mussel survival were
 475 observed in S1, with about 80 % mortality *versus* 10 % recorded elsewhere (Fig. 4A). Different from
 476 S2 – S7, survival (and biomarker) data were measured in S1 after just three days of exposure, in line
 477 with previous observations on mussels transplanted in the same area (Franzellitti et al., 2010). The
 478 short time spent *in situ* within S1 was not such to allow mussels to bioaccumulate metals at levels
 479 considered environmentally realistic, for which ≥ 28 days are usually required (Viarengo et al.,
 480 2007a); thus, data of metal bioaccumulation are missing for this site. Sediment concentrations of
 481 Hg, Cr, Zn, Cu and Ba, known for their toxicity on marine mussels (De Los Ríos et al., 2013) were

482 negatively and significantly correlated with mussel survival (Fig. 5). Relationships were also detected
1
483 between survival and average salinity and temperature, which showed minima and maxima in S1,
2
3
484 respectively (Table S3). Both parameters are known to be responsible for the mussel physiological
4
485 fitness and to influence their response to toxicants synergistically (Fabbri and Dinelli, 2014;
5
6
486 DeCourten and Brander, 2017). It is thus plausible that physico-chemical conditions of both natural
7
8
487 and anthropogenic nature simultaneously influenced the survival of mussels across the lagoon.
9
10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Data of *in situ* survival follow trends similar to those from the SOS test, which showed significant reduction of survival functions in mussels from S1 (Fig. 4B). SOS is measured as the survival of mussels in dry conditions, which is an event regularly experienced by intertidal organisms due to tidal fluctuations. The SOS data display significant and negative correlation with a suite of metals measured in both sediments and mussel tissues, including Ba, Bi, Cu, Sr and Zn, among others (Fig. 5). These findings are in line with previous evidence obtained in *M. galloprovincialis*, which showed significant SOS decrease after 3 to 7 days of exposure to trace levels of Cu and Zn-based antifouling agents (Viarengo et al., 1995; Marcheselli et al., 2011). Further elements showing marked enrichment in S1 sediments, as Cd (0.68 mg/kg) and Hg (25.7 µg/kg), were significantly correlated with SOS (Fig. 4). SOS and survival data were found to be significantly and positively correlated to each other and showed positive and negative correlation with electrical conductivity and temperature, respectively (Table S3). This indicates that higher temperatures and salinity measured at S1 throughout the exposure period may have cumulatively increased the impact of metals on the mussel vital functions. Although it is acknowledged that SOS sensitivity is lower than other sub-lethal and general stress biomarkers (Viarengo et al., 2007a), data from this study highlight its suitability to detect causal relationships between metal contamination/bioavailability and adverse effects on mussels at the “organism” level.




Figure 4. Biomarker modulation in mussels transplanted into the Pialassa Baiona lagoon. Data from all sites (mean \pm SEM) have been measured in mussels collected after 30 days of exposure, except for S1, referred to mussels collected after 3 days of exposure due to high mortality. A, survival (SURV); B, Stress on stress (SOS); C, lysosomal membrane stability (LMS); D, lysosome to cytoplasm volume ratio (LYS/CYT); E, lipofuscin (LIF), F, catalase activity (CAT); G, metallothionein content (MT); H, acetylcholinesterase activity (ACHE); I, peroxisomal acyl-CoA oxidase (AOX). Asterisks indicate statistically significant differences compared to the control site S7 (One-Way ANOVA, Duncan's post-hoc test).

LMS is the most sensitive biomarker of general stress in bivalves and previous studies highlighted strict relationships between the xenobiotic-driven decrease of this parameter in *Mytilus*

515 spp. and alterations to higher functions, such as growth and/or reproduction (Moore et al., 2006;
516 Martinez-Gomez et al., 2008). Data reported in Figure 4C indicate a significant LMS reduction in
517 digestive glands from mussels collected at S1 (3-day exposure), S3 and S6 (30-day exposure)
518 compared to control mussels from S7. Interestingly, none of the measured metals, physico-chemical
519 factors and/or biological endpoints was correlated with trends of LMS in exposed mussels (Fig. 5,
520 Table S3). Although positive relationships between (dissolved) metal exposure and LMS have been
521 reported in mussels under laboratory conditions (Viarengo et al., 1997), no correlation has been
522 highlighted following 28 days of *in situ* exposure into a nearby lagoon (i.e. Pialassa Piomboni,
523 Ravenna, Italy; Capolupo et al., 2017).

524 The mussel exposure to contaminated environment is frequently associated to an augmented
525 permeability of the lysosomal membrane and an overall lysosomal swelling. In the present study,
526 mussels deployed into S3, S5 and S6 showed a significantly higher LYS/CYT compared to the control
527 site S7 (Fig. 4D), following trends significantly correlated to tissue levels of Ba, Mn and Zr (Fig. 5).
528 The S3 and S5 were located in proximity of streams ensuring the flowing of industrial wastewater
529 from southern and eastern settlements to the Adriatic Sea, while S6 was located along a channel
530 that receive water from Lamone river after having crossed the natural areas located to the west (Fig.
531 1); thus, these data outline the need for a better understanding of the spatial/temporal relationships
532 between different source of lagoon water (industrial and continental), metal release into the
533 environment and lysosomal dysfunction in the exposed biota.

534 Lysosomal biomarker of general stress, as LMS and LYS/CYT, describe the overall health status
535 of tested organisms and may thus be modulated by a wide range of natural/anthropogenic stressors;
536 therefore, considering the diverse array of human activities directly and indirectly linked to the
537 Pialassa Baiona lagoon, it cannot be ruled out that further organic pollutants may have played a role
538 in observed lysosomal responses.

539
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 5. Correlation between biological data and metal concentrations. Pearson's coefficients calculated between biological data and levels of metals measured in mussel tissues (bioaccumulation) and sediment. Only coefficients showing a significant correlation ($p < 0.05$) are reported in the figure.

546 Mussels from all sites, except S2, showed a significant accumulation of LIF compared to those
1
547 deployed into the reference site (S7) (Fig. 4E). LIF represent the ultimate products of the lipid
2
548 peroxidation of biological membranes; they are composed by oxidatively modified proteins, lipid
3
549 degradation by-products, carbohydrates and metals, and appear as granules within lysosomes of
4
550 cells composing the digestive tubules epithelium (Moore et al., 2008). Previous biomonitoring
5
551 studies highlighted clear relationships between LIF accumulation in mussels, the metal uptake and
6
552 the metal spatial/temporal distribution into enclosed marine systems as bays and lagoons
7
553 (Moschino et al., 2011; Capolupo et al., 2017). Accordingly, we identified significant and positive
8
554 correlations between LIF, Al and Ti tissue content in mussels, as well as with As, Cd, Se and TI
9
555 measured in sediments (Fig. 5). In addition, a significant and negative correlation has been observed
10
556 between LIF and the activity of the antioxidant enzyme CAT (Table S3). The latter showed a
11
557 significant decrease *versus* control (S7) in mussels exposed at S1 (Fig. 4F), suggesting that beside
12
558 ROS formation, the exposure to relatively high metal concentrations might also down-regulate the
13
559 overall mussel antioxidant defences. CAT also showed negative correlation with elements previously
14
560 found to elicit pro-oxidant effects in mussels, as As, Cd, Cr, Sb (Franzellitti et al., 2012; Benali et al.,
15
561 2017; Coppola et al., 2018), and with temperature trends within the lagoon. This indicates that both
16
562 chemical and physical factors might have cumulatively influenced the mussel ROS scavenging
17
563 activity over the exposure within the lagoon.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 MT have been used as biomarker of exposure in this study owing to their high heavy-metal-
36
37 binding capacity (Lavradas et al., 2016). Compared to control site S7, a significant MT synthesis
38
39 induction has been recorded in the digestive gland of mussels from S3 and S1 (Fig. 4H), both located
40
41 in the southern part of the lagoon in proximity of a large industrial complex (Fig.1). MT trends
42
43 showed a positive and significant correlation with the sediment concentration of metals known as
44
45 MT up-regulator, including Cd, Cu, Zn, Pd, Pt, Sn and Hg (Viarengo et al., 2001; Rocha et al., 2016;
46
47 Zimmermann and Sures, 2018), and other less studied such as Ag, Au, Ba, Bi, Re, and Sb (Fig. 5).
48
49 Positive relationships were detected MT trends and fluctuations of temperature and conductivity
50
51 (Table S3), indicating that physico-chemical factors may interactively modulate MT in mussels from
52
53 contaminated areas.
54


55 The activity of the enzyme AChE, which mediates the termination of the nervous impulses
56
57 driven by the neurotransmitter acetylcholine, was significantly inhibited in mussels from the
58
59 northern site S6 (Fig. 4H). AChE is typically inhibited by pesticides, although in mussels it may also
60
61 be inhibited by high levels of metals (Lionetto et al., 2003). Apart from a negative relationship with
62
63
64
65

578 Be, AChE trends did not show significant relationships with metals measured in sediments and
579 mussel tissues (Fig. 5), nor with other biological endpoints measured in deployed mussels (Table
580 S3). Compared to the S7, sediments collected in S6 showed a 20 times higher enrichment of Hg (Fig.
581 2), which is known as a potent AChE inhibitor in mussels, notably when co-exposed to fluctuations
582 of other physico-chemical parameters, such as temperature (Coppola et al., 2017).

583 The possible impact of organic xenobiotics on the physiology of transplanted mussels has been
584 screened through the analysis of the Acyl-Co A oxidase (AOX) activity. *In situ* deployed mussels
585 showed a significant AOX increase following exposure at all sites except S2 (Fig. 4I). These data
586 corroborate previous evidence of a diffuse PAH contamination (notably pyrene, fluoranthene and
587 anthracene) in sediment and mussels collected from the same area (Vassura et al., 2005; Fabbri et
588 al., 2006). Vassura et al. (2005) also found clear relationships between patterns of PAHs and Hg in
589 the lagoon sediments. Accordingly, many metals herein measured in sediments, including among
590 others Ag, Ba; Hg, Cu, Cr, Cd and Zn, as well as mussel tissues concentrations of B, Cu, Li, Mo and Sr
591 showed positive correlation with AOX (Fig. 5), suggesting a possible metal-induced peroxisome
592 proliferation and a similar spatial distribution of metals and POPs within the lagoon.

593 The output produced by biomarkers data integration performed through the MES is reported
594 in Figure 6. The LMS was chosen as guide parameter for the data integration. A good health status
595 (HSI = A) was assigned by the Expert System to mussels *in situ* exposed at the reference site S7 and
596 at S2. This output confirms the good environmental quality of the lagoon north-eastern area, as
597 previously identified based on chemical and biological evidence (Franzellitti et al., 2010), and is in
598 line with the lack of biomarker alterations observed in mussels from S2. Although this site is located
599 in the southern part of the lagoon, i.e. in proximity of industrial settlements, it is directly exposed
600 to the marine water entering the lagoon indicating the relevance of high water circulation and
601 oxygenation to the fitness of the resilient biota. A moderate stress level (HSI = C) was assigned to
602 mussels exposed at S5, while mussels from S1, S3 and S6 were classified as heavily stressed (HSI =
603 D) (Fig. 6). In line with the metal contamination/bioaccumulation and coherently with MT levels
604 measured in mussels, an excess of metal contamination was identified in S1 and S3. Moreover, the
605 MES associated the health status alterations of mussels exposed at S1, S3, S5 and S6 to the presence
606 of persistent organic compounds (POPs), in line with measured AOX trends. Although organics were
607 not included in the analytical design of this study, this output is consistent with previous evidence
608 of PCB and PAH contamination across the lagoon (Vassura et al., 2005; Fabbri et al., 2006; Guerra,

609 2012) and may confirm the co-occurrence of multiple stressors in different areas of the Pialassa
 610 Baiona lagoon



611 **Figure 6. Biomarker data integration.** A unique Health status index (HSI) has been generated using the Mussel Expert
 612 System (MES, developed by Dagnino et al., 2007) based on biomarker alterations observed at each site. The LMS was
 613 chosen as guide parameter for the HSI assessment.
 614

617 4 Conclusions

618 The integrated approach implemented in this study provided accurate and reliable clues on
 619 the distribution of metals in the Pialassa Baiona lagoon, their bioavailability for exposed mussels
 620 and their effects on key biological functions, posing also attention to seldom considered elements
 621 such as Ag and Sb.

622 The physico-chemical data revealed clear differences among S1 and other investigated sites.
 623 The water in Magni Channel (S1) showed higher column temperature, reduced EC and a lower pH
 624 value, for the entire monitoring period generating local conditions aligned with the worst forecasted
 625 climate change scenarios.

626 The geochemical analysis of total elements indicated that sediments in Pialassa Baiona are a
627 mixture of carbonate and silicates with variable amount of organic matter. The total concentrations
628 of the investigated elements are within the range of regional background. Exception is made for Zn
629 and Cu, and Pb respectively double (in S1) and slightly above (S3 and S5) the local natural
630 background values.

631 The aqua regia digestion identified a south-north decreasing contamination trend, with
632 highest enrichments of Hg, Pt, Au, Ag, Mo, Re, Cd, Pd, Zn; Sn, and Cu (in enrichment order, Fig. 2)
633 being at least two folds enriched than S7 (internal control). Moreover, the most affected sites are
634 in the south-western areas, which consistently receives wastewater effluents from a near
635 petrochemical settlement (south) and drainage water from the inland (west).

636 Bioaccumulation analysis performed in transplanted mussels proved suitable to evaluate the
637 bioavailability of metals in relation to their distribution across the lagoon. Integrating physical,
638 chemical and biological measures allowed for defining causal relationships between observed
639 mortality rates, lysosomal dysfunctions, oxidative stress induction, biomarker of exposure
640 modulations and the metal distribution within the lagoon. Besides this, the different channels
641 examined within the lagoon proved to be an excellent study system for evaluating climate change
642 effects (i.e. modified temperature, pH, and salinity) combined with pollution. In addition, the use of
643 the MES proved suitable to identify the impact induced by specific substances on the overall mussel
644 fitness and to estimate the toxicological pathways occurring in highly pressure lagoons.

645 Overall, data from this study assign a general scarce environmental quality to the Pialassa
646 Baiona lagoon; considering the heterogeneity of this area in terms of geological, physico-chemical
647 and anthropogenic factors, regular biomonitoring surveys should be performed to control the onset
648 of conditions which might exacerbate the impact of metals and other pollutants, notably on a
649 seasonal timeframe. In this respect, the integrated strategy undertaken in this investigation
650 represents a promising approach for the biomonitoring of this area and, more generally, for reliably
651 assessing the environmental quality of highly pressured lagoons.

653 **Acknowledgement**

654 The Authors are grateful to Municipality of Ravenna for supporting the work and allowing field
655 activity within the Piailassa Baiona lagoon. Furthermore, the Authors thank the anonymous
656 Reviewers who helped in improving the quality of the article.

657
1
2
3
658
5
659
7
660
8
661
9
662
11
663
12
664
14
665
15
666
17
667
18
668
20
669
21
670
22
671
23
672
24
673
26
674
27
675
29
676
30
677
32
678
33
679
35
680
36
681
38
682
39
683
41
684
42
685
43
686
45
687
46
688
48
689
49
690
51
691
52
692
53
693
54
694
55
695
57
696
58
59
60
61
62
63
64
65

5 References

- Abdou, M., Schäfer, J., Cobelo-García, A., Neira, P., Petit, J.C.J., Auger, D., Chiffoleau, J.-F., Blanc, G., 2016. Past and present platinum contamination of a major European fluvial–estuarine system: Insights from river sediments and estuarine oysters. *Mar. Chem.* 185, 104-110. <https://doi.org/10.1016/j.marchem.2016.01.006>
- Aliaume, C., Do Chi, T., Viaroli, P., Zaldivar, J.M., 2007. Coastal lagoons of southern Europe: recent changes and future scenarios. *Transitional Waters Monogr.* 1, 1-12, 10.1285/i18252273v1n1p1
- Amorosi, A., Colalongo, M.L., Pasini, G., Preti, D., 1999. Sedimentary response to Late Quaternary sea-level changes in the Romagna coastal plain (northern Italy). *Sedimentology* 46, 99-121.
- Amorosi, A., Centineo, M.C., Dinelli, E., Lucchini, F., Tateo, F., 2002. Geochemical and mineralogical variations as indicators of provenance changes in Late Quaternary deposits of SE Po Plain. *Sediment. Geol.* 151, 273-292. [https://doi.org/10.1016/S0037-0738\(01\)00261-5](https://doi.org/10.1016/S0037-0738(01)00261-5).
- Benali, I., Boutiba, Z., Grandjean, D., De Alencastro, L.F., Rouane-Hacene, O., Chèvre, N., 2017. Spatial distribution and biological effects of trace metals (Cu, Zn, Pb, Cd) and organic micropollutants (PCBs, PAHs) in mussels *Mytilus galloprovincialis* along the Algerian west coast. *Mar. Pollut. Bull.* 115(1-2), 539-550. <https://doi.org/10.1016/j.marpolbul.2016.12.028>
- Bird, E.C.F., 1994. Physical setting and geomorphology of coastal lagoons. In: Kjerfve, B. (Ed.), *Coastal Lagoon Processes*, Elsevier Oceanography Series, vol. 60. Elsevier, Amsterdam, pp. 9-30. [https://doi.org/10.1016/S0422-9894\(08\)70007-2](https://doi.org/10.1016/S0422-9894(08)70007-2)
- Borghesi, F., Migani, F., Dinelli, E., 2016. Geochemical characterization of surface sediments from the northern Adriatic wetlands around the Po River delta. Part II: aqua regia results. *J. Geochem. Explor.* 169, 13-29. <https://doi.org/10.1016/j.gexplo.2016.06.012>
- Bossi, T., Gediga, J., 2017. The environmental profile of Platinum Group Metals. *Johnson Matthey Technol. Rev.* 61, 111-121, <https://doi.org/10.1595/205651317x694713>
- Braga, A.C., Camacho, C., Marques, A., Gago-Martínez, A., Pacheco, M., Costa, P.R., 2018. Combined effects of warming and acidification on accumulation and elimination dynamics of paralytic shellfish toxins in mussels *Mytilus galloprovincialis*. *Environ. Res.* 164, 647-654. <https://doi.org/10.1016/j.envres.2018.03.045>.
- Buryskova, B., Hilscherova, K., Babica, P., Vrskova, D., Marsalek, B., Blaha, L., 2006. Toxicity of complex cyanobacterial samples and their fractions in *Xenopus laevis* embryos and the role of microcystins. *Aquat. Toxicol.* 80, 346-354. <https://doi.org/10.1016/j.aquatox.2006.10.001>
- Capolupo, M., Valbonesi, P., Kiwan, A., Buratti, S., Franzellitti, S., Fabbri, E., 2016. Use of an integrated biomarker-based strategy to evaluate physiological stress responses induced by environmental concentrations of caffeine in the Mediterranean mussel *Mytilus galloprovincialis*. *Sci. Total Environ.* 563-564, 538-548. doi:10.1016/j.scitotenv.2016.04.125
- Capolupo, M., Franzellitti, S., Kiwan, A., Valbonesi, P., Dinelli, E., Pignotti, E., Birke, M., Fabbri, E., 2017. A comprehensive evaluation of the environmental quality of a coastal lagoon (Ravenna, Italy): Integrating chemical and physiological analyses in mussels as a biomonitoring strategy. *Sci. Total Environ.* 598, 146-159. <https://doi.org/10.1016/j.scitotenv.2017.04.119>

- 697 Carvalho, R.N., Arukwe, A., Ait-Aissa, S., Bado-Nilles, A., Balzamo, S., Baun, A., Belkin, S., Blaha, L.,
698 Brion, F., Conti, D., Creusot, N., Essig, Y., Ferrero, V.E.V., Flander-Putrlle, V.V., Fürhacker, M.,
699 Grillari-Voglauer, R., Hogstrand, C., Jonáš, A., Kharlyngdoh, J.B., Loos, R., Lundebye, A.K., Modig,
700 C., Olsson, P.E., Pillai, S., Polak, N., Potalivo, M., Sanchez, W., Schifferli, A., Schirmer, K., Sforzini,
701 S., Stürzenbaum, S.R., Søfteland, L., Turk, V., Viarengo, A., Werner, I., Yagur-Kroll, S.S.,
702 Zounkova, R., Lettieri, T., 2014. Mixtures of chemical pollutants at European legislation safety
703 concentrations: how safe are they? *Toxicol. Sci.* 141, 218-233.
704 <http://dx.doi.org/10.1093/toxsci/kfu118>.
- 705 Cicchella, D., Giaccio, L., Dinelli, E., Albanese, S., Lima, A., Zuzolo, D., Valera, P., De Vivo, B., 2015.
706 GEMAS: Spatial distribution of chemical elements in agricultural and grazing land soil of Italy. *J.*
707 *Geochem. Explor.* 154, 129-142. <https://doi.org/10.1016/j.gexplo.2014.11.009>
- 708 Cobelo-García, A., Neira, P., Mil-Homens, M., Caetano, M., 2011. Evaluation of the contamination
709 of platinum in estuarine and coastal sediments (Tagus Estuary and Prodelta, Portugal). *Mar.*
710 *Pollut. Bull.* 62 (3), 646-650. <https://doi.org/10.1016/j.marpolbul.2010.12.018>
- 711 Coppola, F., Almeida, Â., Henriques, B., Soares, A.M., Figueira, E., Pereira, E., Freitas, R., 2017.
712 Biochemical impacts of Hg in *Mytilus galloprovincialis* under present and predicted warming
713 scenarios. *Sci. Total Environ.* 601, 1129-1138. <https://doi.org/10.1016/j.scitotenv.2017.05.201>
- 714 Coppola, F., Almeida, Â., Henriques, B., Soares, A.M.V.M., 2018. Ecotoxicology and Environmental
715 Safety Biochemical responses and accumulation patterns of *Mytilus galloprovincialis* exposed
716 to thermal stress and Arsenic contamination. *Ecotox. Environ. Saf.* 147, 954-962. <https://doi.org/10.1016/j.ecoenv.2017.09.051>
- 717
- 718 Covelli, S., Emili, A., Acquavita, A., Koron, N., Faganeli, J., 2011. Benthic biogeochemical cycling of
719 mercury in two contaminated northern Adriatic coastal lagoons. *Cont. Shelf Res.* 31(16), 1777-
720 1789. <https://doi.org/10.1016/j.csr.2011.08.005>.
- 721 Curzi, P.V., Dinelli, E., Ricci Lucchi, M., Vaiani, S.C., 2006. Palaeoenvironmental control on sediment
722 composition and provenance in the late Quaternary deltaic successions: A case study from the
723 Po delta area (Northern Italy). *Geol. J.* 41 (5), 591-612. <https://doi.org/10.1002/gj.1060>
- 724 Dagnino, A., Allen, J.I., Moore, M.N., Broeg, K., Canesi, L., Viarengo, A., 2007. Development of an
725 expert system for the integration of biomarker responses in mussels into an animal health
726 index. *Biomarkers* 12, 155-172. <http://dx.doi.org/10.1080/13547500601037171>.
- 727 Dahms, H.U., 2014. The grand challenges in marine pollution research. *Front. Mar. Sci.* 1, 1-5.
728 <http://dx.doi.org/10.3389/fmars.2014.00009>
- 729 De Gregori, I., Quiroz, W., Pinochet, H., Pannier, F., Potin-Gautier, M., 2007. Speciation analysis of
730 antimony in marine biota by HPLC-(UV)-HG-AFS: Extraction procedures and stability of
731 antimony species. *Talanta* 73(3), 458-465. <https://doi.org/10.1016/j.talanta.2007.04.015>
- 732 De Los Ríos, A., Pérez, L., Ortiz-zarragoitia, M., Serrano, T., Barbero, M.C., Echavarri-erasun, B., 2013.
733 Assessing the effects of treated and untreated urban discharges to estuarine and coastal waters
734 applying selected biomarkers on caged mussels. *Mar. Pollut. Bull.* 77, 251-265.
735 <https://doi.org/10.1016/j.marpolbul.2013.09.050>
- 736 Decourten, B.M., Brander, S.M., 2017. Combined effects of increased temperature and endocrine
737 disrupting pollutants on sex determination, survival, and development across generations. *Sci.*
738 *Rep.* 7, 1-9. <https://doi.org/10.1038/s41598-017-09631-1>

60
61
62
63
64
65

- 739 Dinelli, E., Tateo F., Summa V. 2007. Geochemical and mineralogical proxies for grain size in
740 mudstones and siltstones from the Pleistocene and Holocene of the Po River alluvial plain, Italy
741 J. Arribas, S. Critelli, M.J. Johnsson (Eds.), Sedimentary Provenance and Petrogenesis:
742 Perspectives from Petrography and Geochemistry, Geological Society of America Special Paper,
743 vol. 420, pp. 25-36
- 744 Dinelli, E., Ghosh, A., Rossi, V., Vaiani, S.C., 2012. Multiproxy reconstruction of Late Pleistocene-
745 Holocene environmental changes in coastal successions: microfossil and geochemical evidences
746 from the Po Plain (Northern Italy). Stratigraphy 9(2), 153-167.
- 747 Dominik, J., Tagliapietra, D., Bravo, A.G., Sigovini, M., Spangenberg, J.E., Amouroux, D., Zonta, R.
748 2014. Mercury in the food chain of the Lagoon of Venice, Italy. Mar. Pollut. Bull. 88(1), 194-206.
749 <https://doi.org/10.1016/j.marpolbul.2014.09.005>.
- 750 Donnini, F., Dinelli, E., Sangiorgi, F., Fabbri, E., 2007. A biological and geochemical integrated
751 approach to assess the environmental quality of a coastal lagoon (Ravenna, Italy). Environ. Int.
752 33 (7), 919-928. <https://doi.org/10.1016/j.envint.2007.05.002>.
- 753 Duroudier, N., Cardoso, C., Mehennaoui, K., Mikolaczyk, M., Schäfer, J., Gutleb, A.C., Giamberini, L.,
754 Bebianno, M.J., Bilbao, E., Cajaraville, M.P., 2019. Changes in protein expression in mussels
755 *Mytilus galloprovincialis* dietarily exposed to PVP/PEI coated silver nanoparticles at different
756 seasons. Aquat. Toxicol. 210, 56-68. <https://doi.org/10.1016/j.aquatox.2019.02.010>.
- 757 European Commission Directive 2000/60/EC of the European Parliament and of the Council of 23
758 October 2000 establishing a framework for community action in the field of water policy (Water
759 Framework Directive). Off. J. Eur. Community, 2000.
- 760 European Commission Directive 2008/56/EC of the European Parliament and of the Council of 17
761 June 2008 Establishing a Framework for Community Action in the Field of Marine Environmental
762 Policy (Marine Strategy Framework Directive) (OJ L 164/19. 22 pp. 25.06).
- 763 Ellman, G.L., Courtney, K.D., Andres Jr., V., Feather-Stone, R.M., 1961. A new and rapid colorimetric
764 determination of acetylcholinesterase activity. Biochem. Pharmacol. 7, 88-95.
765 [https://doi.org/10.1016/0006-2952\(61\)90145-9](https://doi.org/10.1016/0006-2952(61)90145-9)
- 766 Fabbri, D., Felisatti, O., Lombardo, M., Trombini, C., Vassura, I., 1998. The Lagoon of Ravenna Italy:
767 characterisation of mercury-contaminated sediments. Sci. Total Environ. 213, 121-128.
768 [https://doi.org/10.1016/S0048-9697\(98\)00083-7](https://doi.org/10.1016/S0048-9697(98)00083-7)
- 769 Fabbri, D., Tartari, D., Trombini, C., 2000. Analysis of poly (vinyl chloride) and other polymers in
770 sediments and suspended matter of a coastal lagoon by pyrolysis-gas chromatography-mass
771 spectrometry. Anal. Chim. Acta 413(1-2), 3-11. [https://doi.org/10.1016/S0003-2670\(00\)00766-2](https://doi.org/10.1016/S0003-2670(00)00766-2)
- 772 2
- 773 Fabbri, D., Gemelli, S., Langone, L., Misericocchi, S., Trombini, C., Vassura, I., 2001a. Anthropogenically
774 derived mercury in sediments of Piasa Baiona, Ravenna, Italy. Annali di Chimica 91(9-10),
775 563-575.
- 776 Fabbri, D., Locatelli, C., Snape, C.E., Tarabusi, S., 2001b. Sulfur speciation in mercury-contaminated
777 sediments of a coastal lagoon: The role of elemental sulfur. J. Environ. Monitor. 5, 483-486.
778 <https://doi.org/10.1039/b104477j>
- 779 Fabbri, D., Baravelli, V., Giannotti, K., Donnini, F., Fabbri, E., 2006. Bioaccumulation of
780 cyclopenta[cd]pyrene and benzo[ghi]fluoranthene by mussels transplanted in a coastal lagoon.
781 Chemosphere 64, 1083-1092. <https://doi.org/10.1016/j.chemosphere.2005.11.071>

- 782 Fabbri, E., Dinelli, E., 2014. Physiological responses of marine animals towards adaptation to climate
783 changes. The Mediterranean Sea: Its History and Present Challenges, pp. 401-417.
784 https://doi.org/10.1007/978-94-007-6704-1_23
- 785 Fattorini, D., Notti, A., Di Mento, R., Cicero, A.M., Gabellini, M., Russo, A., Regoli, F., 2008. Seasonal,
786 spatial and inter-annual variations of trace metals in mussels from the Adriatic Sea: a regional
787 gradient for arsenic and implications for monitoring the impact of off-shore activities.
788 *Chemosphere* 72(10), 1524-1533. <https://doi.org/10.1016/j.chemosphere.2008.04.071>.
- 789 Filella, M., Belzile, N., Chen, Y.W., 2002. Antimony in the environment: a review focused on natural
790 waters: I. Occurrence. *Earth-Sci. Rev.* 57(1-2), 125-176. [https://doi.org/10.1016/S0012-
791 8252\(01\)00070-8](https://doi.org/10.1016/S0012-8252(01)00070-8)
- 792 Filella, M., Hennebert, P., Okkenhaug, G., Turner, A., 2019. Occurrence and fate of antimony in
793 plastics. *J. Hazard. Mater.* 121764. <https://doi.org/10.1016/j.jhazmat.2019.121764>
- 794 Franzellitti, S., Buratti, S., Donnini, F., Fabbri, E., 2010. Exposure of mussels to a polluted
795 environment: Insights into the stress syndrome development. *Comp. Biochem. Physiol. - C
796 Toxicol. Pharmacol.* 152, 24-33. <https://doi.org/10.1016/j.cbpc.2010.02.010>
- 797 Franzellitti, S., Viarengo, A., Dinelli, E., Fabbri, E., 2012. Molecular and cellular effects induced by
798 hexavalent chromium in Mediterranean mussels. *Aquat. Toxicol.* 124-125, 125-132.
799 <https://doi.org/10.1016/j.aquatox.2012.07.011>
- 800 Franzini, M., Leoni, L., Saitta, M., 1972. A simple method to evaluate the matrix effects in X-ray
801 fluorescence analysis. *X-Ray Spectrom.* 1, 151-154. <https://doi.org/10.1002/xrs.1300010406>
- 802 Franzini, M., Leoni, L., Saitta, M., 1975. Revisione di una metodologia analitica per fluorescenza-X
803 basata sulla correzione completa degli effetti di matrice. *Rend. Soc. Ital. Mineral. Petrol.* 31,
804 365-378. (in italian)
- 805 Garcia, N., 2015. The Asiatic Clam (*Corbicula fluminea*) as a bio-monitor for determining the
806 distribution of antimony, arsenic and thallium in the water column and sediments of Manadas
807 Creek, Laredo, Texas. M.Sc thesis, Texas A&M International University, pp. 59. [https://tamiu-
808 ir.tdl.org/handle/2152.4/58](https://tamiu-ir.tdl.org/handle/2152.4/58). Accessed July 2020
- 809 Giusti, L., Williamson, A.C., Mistry, A., 1999. Biologically available trace metals in *Mytilus edulis* from
810 the coast of northeast England. *Environ. Int.* 25(8), 969-981. [https://doi.org/10.1016/S0160-
811 4120\(99\)00066-5](https://doi.org/10.1016/S0160-4120(99)00066-5)
- 812 Giusti, L., Zhang, H., 2002. Heavy metals and arsenic in sediments, mussels and marine water from
813 Murano (Venice, Italy). *Environ. Geochem. Health* 24(1), 47-65.
814 <https://doi.org/10.1023/A:1013945117549>
- 815 Govindaraju, K., 1984. 1984 compilation of working values and sample description for 170
816 international reference samples of mainly silicate rocks and minerals. *Geostandard. Newslett.*
817 8, 3-16. <https://doi.org/10.1111/j.1751-908X.1984.tb01069.x>
- 818 Greggio, N., Giambastiani, B.M.S., Campo, B., Dinelli, E., Amorosi, A., 2018a. Sediment composition,
819 provenance, and Holocene paleoenvironmental evolution of the Southern Po River coastal plain
820 (Italy). *Geol. J.* 53(3), 914-928. <https://doi.org/10.1002/gj.2934>
- 821 Greggio, N., Giambastiani, B.M.S., Antonellini, M., 2018b. Infiltration/irrigation trench for
822 sustainable coastal drainage management: Emilia-Romagna (Italy). *Environ. Eng. Manag. J.*
823 17(10). <https://doi.org/10.30638/eemj.2018.236>

- 824 Greggio, N., Giambastiani, B., Mollema, P., Laghi, M., Capo, D., Gabbianelli, G., Antonellini, M.,
825 Dinelli, E., 2020. Assessment of the Main Geochemical Processes Affecting Surface Water and
826 Groundwater in a Low-Lying Coastal Area: Implications for Water Management. *Water* 12(6),
827 1720. <https://doi.org/10.3390/w12061720>
- 828 Guerra, R., Pasteris, A., Ponti, M., Fabbri, D., Bruzzi, L., 2007. Impact of dredging in a shallow coastal
829 lagoon: Microtox® Basic Solid-Phase Test, trace metals and Corophium bioassay. *Environ. Int.*
830 33 (4), 469-473. <https://doi.org/10.1016/j.envint.2006.10.006>
- 831 Guerra R., 2012. Polycyclic Aromatic Hydrocarbons, Polychlorinated Biphenyls and Trace Metals in
832 Sediments from a Coastal Lagoon (Northern Adriatic, Italy). *Water Air Soil Poll.* 223, 85-98.
833 <https://doi.org/10.1007/s11270-011-0841-6>
- 834 Guerra R., Pasteris A., Lee Seok-hyung, Park No-jin, Ok Gon, 2014. Spatial patterns of metals,
835 PCDDs/Fs, PCBs, PBDEs and chemical status of sediments from a coastal lagoon (Pialassa Baiona,
836 NW Adriatic, Italy). *Mar. Pollut. Bull.* 89, 407-416.
837 <https://doi.org/10.1016/j.marpolbul.2014.10.024>
- 838 James, E., Turner, A., 2020. Mobilisation of antimony from microplastics added to coastal sediment.
839 *Environ. Pollut.* 114696. <https://doi.org/10.1016/j.envpol.2020.114696>
- 840 Jimeno-Romero, A., Bilbao, E., Izagirre, U., Cajaraville, M.P., Marigómez, I., Soto, M., 2017. Digestive
841 cell lysosomes as main targets for Ag accumulation and toxicity in marine mussels, *Mytilus*
842 *galloprovincialis*, exposed to maltose-stabilised Ag nanoparticles of different sizes.
843 *Nanotoxicology* 11(2), 168-183. <https://doi.org/10.1080/17435390.2017.1279358>
- 844 Lari, E., Gauthier, P., Mohaddes, E., Pyle, G.G., 2017. Interactive toxicity of Ni, Zn, Cu, and Cd on
845 *Daphnia magna* at lethal and sub-lethal concentrations. *J. Hazard. Mater.* 334, 21-28.
846 <http://dx.doi.org/10.1016/j.jhazmat.2017.03.060>
- 847 Lavradas, R.T., Rocha, R.C.C., Bordon, I.C.A.C., Saint’Pierre, T.D., Godoy, J.M., Hauser-Davis, R.A.,
848 2016. Differential metallothionein, reduced glutathione and metal levels in *Perna perna* mussels
849 in two environmentally impacted tropical bays in southeastern Brazil. *Ecotoxicol. Environ. Saf.*
850 129, 75-84. <http://dx.doi.org/10.1016/j.ecoenv.2016.03.011>
- 851 León, V.M., Moreno-González, R., García, V., Campillo, J.A., 2017. Impact of flash flood events on
852 the distribution of organic pollutants in surface sediments from a Mediterranean coastal lagoon
853 (Mar Menor, SE Spain). *Environ. Sci. Pollut. Res.* 24(5), 4284-4300.
854 <https://doi.org/10.1007/s11356-015-4628-y>
- 855 Leoni, L., Saitta, M., 1976. X-ray fluorescence analysis of 29 trace elements in rock and mineral
856 standard. *Rend. Soc. Ital. Mineral.* 72, 175-192.
- 857 Leoni, L., Menichini, M., Saitta, M., 1986. Determination of S, Cl, and F in silicate rocks by X-ray
858 fluorescence analyses. *X-Ray Spectrom.* 11, 156-158. <https://doi.org/10.1002/xrs.1300110404>
- 859 Leopold, K., Denzel, A., Gruber, A., Malle, L., 2018. Mobility of traffic-related Pd and Pt species in
860 soils evaluated by sequential extraction. *Environ. Pollut.* 242, 1119-1127.
861 <https://doi.org/10.1016/j.envpol.2018.07.130>
- 862 Lionetto, M.G., Caricato, R., Giordano, M.E., Pascariello, M.F., Marinosci, L., Schettino, T., 2003.
863 Integrated use of biomarkers (acetylcholinesterase and antioxidant enzymes activities) in
864 *Mytilus galloprovincialis* and *Mullus barbatus* in an Italian coastal marine area. *Mar. Pollut. Bull.*
865 46(3), 324-330. [https://doi.org/10.1016/S0025-326X\(02\)00403-4](https://doi.org/10.1016/S0025-326X(02)00403-4)

- 866 Lottermoser, B.G., Ashley, P.M., Lawie, D.C., 1999. Environmental geochemistry of the Gulf Creek
867 copper mine area, north-eastern New South Wales, Australia. *Environ. Geol.* 39(1), 61-74.
868 <https://doi.org/10.1007/s002540050437>
- 869 Lowry, O.H., Rosenbrough, N.J., Farr, A.L., Randall, R.J., 1951. Protein measurement with the Folin
870 phenol reagent. *J. Biol. Chem.* 193, 265-275.
- 871 Mansour, C., Guibbolini, M., Rouane, O., Dalila, H., Mosbahi, S., De Faverney, C.R., 2020. Oxidative
872 stress and damage biomarkers in clam *ruditapes decussatus* exposed to a polluted site: the
873 reliable biomonitoring tools in hot and cold seasons. *Arch. Environ. Contam. Toxicol.* 78, 478-
874 494. <https://doi.org/10.1007/s00244-020-00713-2>
- 875 Marcheselli, M., Azzoni, P., Mauri, M., 2011. Novel antifouling agent-zinc pyrithione : Stress
876 induction and genotoxicity to the marine mussel *Mytilus galloprovincialis*. *Aquat. Toxicol.* 102,
877 39-47. <https://doi.org/10.1016/j.aquatox.2010.12.015>
- 878 Martin, M., Stephenson, M.D., Smith, D.R., Gutierrez-Galindo, E.A., Munoz, G.F., 1988. Use of silver
879 in mussels as a tracer of domestic wastewater discharge, *Mar. Pollut. Bull.* 19, 512-520
880 [https://doi.org/10.1016/0025-326X\(88\)90541-3](https://doi.org/10.1016/0025-326X(88)90541-3)
- 881 Martin-Diaz, L., Franzellitti, S., Buratti, S., Valbonesi, P., Capuzzo, A., Fabbri, E., 2009. Effects of
882 environmental concentrations of the antiepileptic drug carbamazepine on biomarkers and
883 cAMP-mediated cell signaling in the mussel *Mytilus galloprovincialis*. *Aquat. Toxicol.* 94, 177-
884 185. <https://doi.org/10.1016/j.aquatox.2009.06.015>
- 885 Martinez-Gomez, C., Benedicto, J., Campillo, J.A., Moore, M., 2008. Application and evaluation of
886 the neutral red retention (NRR) assay for lysosomal stability in mussel populations along the
887 Iberian Mediterranean coast. *J. Environ. Monit.* 10, 490-499.
888 <https://doi.org/10.1039/C2EM90056D>
- 889 Mattes, T.E., Ewald, J.M., Liang, Y., Martinez, A., Awad, A., Richards, P., Hornbuckle, K.C., Schnoor,
890 J.L., 2018. PCB dechlorination hotspots and reductive dehalogenase genes in sediments from a
891 contaminated wastewater lagoon. *Environ. Sci. Pollut. Res.*, 25(17), 16376-16388.
892 <https://doi.org/10.1007/s11356-017-9872-x>
- 893 Matteucci, G., Rossini, P., Guerzoni, S., Arcangeli, A., Fonti, P., Langone, L., Miseroocchi, S., 2005.
894 Recent evolution of sedimentary heavy metals in a coastal lagoon contaminated by industrial
895 wastewaters (Pialassa Baiona, Ravenna, Italy). *Hydrobiologia* 550, 167-173.
896 <https://doi.org/10.1007/s10750-005-4374-0>
- 897 McGillicuddy, E., Murray, I., Kavanagh, S., Morrison, L., Fogarty, A., Cormican, M., Dockery, P.,
898 Prendergast, M., Rowan, N., Morris, D., 2017. Silver nanoparticles in the environment: sources,
899 detection and ecotoxicology. *Sci. Total. Environ.* 575, 231-246.
900 <https://doi.org/10.1016/j.scitotenv.2016.10.041>
- 901 Migani, F., Borghesi, F., Dinelli, E., 2015 Geochemical characterization of surface sediments from the
902 northern Adriatic wetlands around the Po river delta. Part I: Bulk composition and relation to
903 local background. *J. Geochem. Explor.* 156, 72-88. DOI: 10.1016/j.gexplo.2015.05.003
- 904 Miseroocchi, S., Langone, L., Guerzoni, S., 1993. The fate of Hg contaminated sediments of the
905 Ravenna Lagoon (Italy): Final burial or potential remobilization. *Water Sci. Technol.* 28 (8-9),
906 349-358. <https://doi.org/10.2166/wst.1993.0633>
- 907 Mollema, P.N., Antonellini, M., Dinelli, E., Gabbianelli, G., Greggio, N., Stuyfzand, P.J., 2013.
908 Hydrochemical and physical processes influencing salinization and freshening in Mediterranean

- low-lying coastal environments. *Appl. Geochem.* 34, 207-221.
<https://doi.org/10.1016/j.apgeochem.2013.03.017>
- Moore, M.N., Allen, J.I., McVeigh, A., 2006. Environmental prognostics: an integrated model supporting lysosomal stress responses as predictive biomarkers of animal health status. *Marine Environ. Res.* 61(3), 278-304. <https://doi.org/10.1016/j.marenvres.2005.10.005>
- Moore, M.N., 2008. Autophagy as a second level protective process in conferring resistance to environmentally-induced oxidative stress. *Autophagy* 4, 254-256.
<https://doi.org/10.4161/auto.5528>
- Moreno-González, R., Rodríguez-Mozaz, S., Gros, M., Barceló, D., León, V.M., 2015. Seasonal distribution of pharmaceuticals in marine water and sediment from a mediterranean coastal lagoon (SE Spain). *Environ. Res.* 138, 326-344. <https://doi.org/10.1016/j.envres.2015.02.016>
- Moschino, V., Delaney, E., Meneghetti, F., Da, L., 2011. Biomonitoring approach with mussel *Mytilus galloprovincialis* (Lmk) and clam *Ruditapes philippinarum* (Adams and Reeve, 1850) in the Lagoon of Venice. *Environ. Monit. Assess.* 177, 649-663. <https://doi.org/10.1007/s10661-010-1663-5>
- Mousavi, M.P., Gunsolus, I.L., De Jesús, C.E.P., Lancaster, M., Hussein, K., Haynes, C.L., Bühlmann, P., 2015. Dynamic silver speciation as studied with fluoros-phase ion-selective electrodes: Effect of natural organic matter on the toxicity and speciation of silver. *Sci. Total Environ.* 537, 453-461. <https://doi.org/10.1016/j.scitotenv.2015.07.151>
- Nazneen, S., Singh, S., Raju, N.J., 2019. Heavy metal fractionation in core sediments and potential biological risk assessment from Chilika lagoon, Odisha state, India. *Quatern. Int.* 507, 370-388. <https://doi.org/10.1016/j.quaint.2018.05.011>
- Newton, A., Icely, J., Cristina, S., Brito, A., Cardoso, A.C., Colijn, F., Riva, S.D., Gertz, F., Hansen, J.W., Holmer, M., Ivanova, K., Leppäkoski, E., Canu, D.M., Mocenni, C., Mudge, S., Murray, N., Pejrup, M., Razinkovas, A., Reizopoulou, S., Pérez-Ruzafa, A., Schernewski, G., Schubert, H., Carr, L., Solidoro, C., Viaroli, P., Zaldívar, J.-M., K., 2014. An overview of ecological status, vulnerability and future perspectives of European large shallow, semi-enclosed coastal systems, lagoons and transitional waters. *Estuar. Coast. Shelf Sci.* 140, 95-122.
<https://doi.org/10.1016/j.ecss.2013.05.023>
- Orbea, A., Garmendia, L., Marigómez, I., Cajaraville, M.P., 2006. Effects of the “Prestige” oil spill on cellular biomarkers in intertidal mussels: Results of the first year of studies. *Mar. Ecol. Prog. Ser.* 306, 177-189. <https://doi.org/10.3354/meps306177>
- Outa, J.O., Kowenje, C.O., Plessl, C., Jirsa, F., 2020. Distribution of arsenic, silver, cadmium, lead and other trace elements in water, sediment and macrophytes in the Kenyan part of Lake Victoria: spatial, temporal and bioindicative aspects. *Environ. Sci. Pollut. Res.* 27(2), 1485-1498. <https://doi.org/10.1007/s11356-019-06525-9>
- Ozkan, D., Dagdeviren, M., Katalay, S., Guner, A., Yavaşoğlu, N.Ü.K., 2017. Multi-biomarker responses after exposure to pollution in the mediterranean mussels (*Mytilus galloprovincialis* L.) in the Aegean Coast of Turkey. *B. Environ. Contam. Tox.* 98(1), 46-52. <https://doi.org/10.1007/s00128-016-1988-z>
- Petry, A.C., Guimarães, T.F., Vasconcellos, F.M., Hartz, S.M., Becker, F.G., Rosa, R.S., Goyenola, G., Caramaschi, E.P., Díaz de Astarloa, J.M., Sarmiento-Soares, L.M., Vieira, J.P., Garcia, A.M., Teixeira de Mello, F., de Melo, F.A., Meerhoff, M., Attayde, J.L., Menezes, R.F., Mazzeo, N., Di

- 952 Dario, F., 2016. Fish composition and species richness in eastern South American coastal
953 lagoons: additional support for the freshwater ecoregions of the world. *J. Fish Biol.* 89(1), 280-
954 314. <https://doi.org/10.1111/jfb.13011>
- 955 Piggott, J.J., Townsend, C.R., Matthaei, C.D., 2015. Reconceptualizing synergism and antagonism
956 among multiple stressors. *Ecol. Evol.* 5, 1538-1547. <http://dx.doi.org/10.1002/ece3.1465>.
- 957 Pignotti, E., Casas, G., Llorca, M., Tellbüscher, A., Almeida, D., Dinelli, E., Farré, M., Barceló, D., 2017.
958 Seasonal variations in the occurrence of perfluoroalkyl substances in water, sediment and fish
959 samples from Ebro Delta (Catalonia, Spain). *Sci. Total Environ.* 607-608, 933-943.
960 <https://doi.org/10.1016/j.scitotenv.2017.07.025>
- 961 Pignotti, E., Dinelli, E., 2018. Distribution and partition of endocrine disrupting compounds in water
962 and sediment: Case study of the Romagna area (North Italy). *J. Geochem. Explor.* 195, 66-77.
963 <https://doi.org/10.1016/j.gexplo.2018.02.008>
- 964 Pignotti, E., Guerra, R., Covelli, S., Fabbri, E., Dinelli, E., 2018. Sediment quality assessment in a
965 coastal lagoon (Ravenna, NE Italy) based on SEM-AVS and sequential extraction procedure. *Sci.*
966 *Total Environ.* 635, 216-227. <https://doi.org/10.1016/j.scitotenv.2018.04.093>
- 967 Piló, D., Carvalho, S., Pereira, P., Gaspar, M.B., Leitão, A., 2017. Is metal contamination responsible
968 for increasing aneuploidy levels in the Manila clam *Ruditapes philippinarum*?. *Sci. Total Environ.*
969 577, 340-348. <http://dx.doi.org/10.1016/j.scitotenv.2016.10.193>.
- 970 Ravindra, K., Bencs, L., Van Grieken, R., 2004. Platinum group elements in the environment and their
971 health risk. *Sci. Total Environ.* 318 (1-3), 1-43. [https://doi.org/10.1016/S0048-9697\(03\)00372-3](https://doi.org/10.1016/S0048-9697(03)00372-3)
- 972 Rocha, T.L., Gomes, T., Durigon, E.G., Bebianno, M.J., 2016. Subcellular partitioning kinetics,
973 metallothionein response and oxidative damage in the marine mussel *Mytilus galloprovincialis*
974 exposed to cadmium-based quantum dots. *Sci. Total Environ.* 554-555, 130-141.
975 doi:10.1016/j.scitotenv.2016.02.168
- 976 Ruchter, N., Sures, B., 2015. Distribution of platinum and other traffic related metals in sediments
977 and clams (*Corbicula* sp.). *Water Res.* 70, 313-324.
978 <https://doi.org/10.1016/j.watres.2014.12.011>
- 979 Saidi, I., Said, O.B., Abdelmalek, J.B., Chicharo, L., Beyrem, H., 2019. Assessment of Heavy Metal
980 Contamination in the Sediment of the Bizerte Lagoon in Northern Tunisia. In *Recent Advances*
981 *in Geo-Environmental Engineering, Geomechanics and Geotechnics, and Geohazards*, pp. 41-
982 44. Springer, Cham. https://doi.org/10.1007/978-3-030-01665-4_10
- 983 Shaw, J.P., Dondero, F., Moore, M.N., Negri, A., Dagnino, A., Readman, J.W., Lowe, D.R., Frickers,
984 P.E., Beesley, A., Thain, J.E., Viarengo, A., 2011. Integration of biochemical, histochemical and
985 toxicogenomic indices for the assessment of health status of mussels from the Tamar Estuary,
986 U.K. *Mar. Environ. Res.* 72, 13-24. <https://doi.org/10.1016/j.marenvres.2011.05.003>
- 987 Soprani, S., Ricci, C., Farina, D., Tarroni, C., Zanetti, S., 1994. Indagine sulle caratteristiche delle
988 acque. In: Soprani S, Ricci C, editors. *Analisi dello stato ambientale e sanitario nelle valli*
989 *ravennati. La Pialassa Baiona. Ravenna: Azienda U.S.L., Dipartimento dei Servizi di Prevenzione;*
990 1994. pp. 201-254.
- 991 Trombini, C., Fabbri, D., Lombardo, M., Vassura, I., Zavoli, E., Horvat, M., 2003. Mercury and
992 methylmercury contamination in surficial sediments and clams of a coastal lagoon (Pialassa
993 Baiona Ravenna Italy). *Cont. Shelf Res.* 23, 1821-1831.
994 <https://doi.org/10.1016/j.csr.2003.06.010>

- 995 Turja, R., Höher, N., Snoeijls, P., Baršiene, J., Butrimavičiene, L., Kuznetsova, T., Kholodkevich, S. V.,
996 Devier, M.H., Budzinski, H., Lehtonen, K.K., 2014. A multibiomarker approach to the assessment
997 of pollution impacts in two Baltic Sea coastal areas in Sweden using caged mussels (*Mytilus*
998 *trossulus*). Sci. Total Environ. 473-474, 398-409.
999 <http://doi.org/10.1016/j.scitotenv.2013.12.038>
- 1000 UNEP/RAMOGGE: Manual on the Biomarkers Recommended for the MED POL Biomonitoring
1001 Programme. UNEP, Athens, 1999.
1002 http://wedocs.unep.org/bitstream/handle/20.500.11822/1771/ramoge_english.pdf?sequence=1&isAllowed=y (Accessed October 2019).
- 1003
1004 Valbonesi, P., Sartor, G., Fabbri, E., 2003. Characterization of cholinesterase activity in three bivalves
1005 inhabiting the North Adriatic Sea and their possible use as sentinel organisms for biosurveillance
1006 programmes. Sci. Total Environ. 312, 79-88. [http://doi.org/10.1016/S0048-9697\(03\)00227-4](http://doi.org/10.1016/S0048-9697(03)00227-4).
- 1007 Van Cauwenberghe, L., Claessens, M., Vandegheuchte, M.B., Janssen, C.R., 2015. Microplastics are
1008 taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural habitats.
1009 Environ. Pollut., 199, 10-17. <https://doi.org/10.1016/j.envpol.2015.01.008>
- 1010 Vassura, I., Foschini, F., Baravelli, V., Fabbri, D., 2005. Distribution of alternant and non-alternant
1011 polycyclic aromatic hydrocarbons in sediments and clams of the Pialassa Baiona Lagoon
1012 (Ravenna, Italy). Chem. Ecol., 21, 415-424. <https://doi.org/10.1080/02757540500438490>.
- 1013 Viarengo, A., Moore, M.N., Mancinelli, G., Mazzucotelli, A., Pipe, R.K., Farrar, S.V., 1987.
1014 Metallothioneins and lysosomes in metal toxicity and accumulation in marine mussels: the
1015 effect of cadmium in the presence and absence of phenanthrene. Mar. Biol. 94(2), 251-257.
1016 <https://doi.org/10.1007/BF00392937>
- 1017 Viarengo, A., Canesi, L., Pertica, M., Mancinelli, G., Accomando, R., Smaalb, A.C., Orunesua, M.,
1018 1995. Stress on Stress Response: A Simple Monitoring Tool in the Assessment of a General
1019 Stress Syndrome in Mussels. Mar. Environ. Res. 39, 245-248. [https://doi.org/10.1016/0141-1136\(94\)00075-Z](https://doi.org/10.1016/0141-1136(94)00075-Z)
- 1020
1021 Viarengo, A., Ponzano, E., Dondero, F., Fabbri, R., 1997. A simple spectrophotometric method for MT
1022 evaluation in marine organisms: an application to Mediterranean and Antarctic molluscs. Mar.
1023 Environ. Res. 44, 69-84. [https://doi.org/10.1016/S0141-1136\(96\)00103-1](https://doi.org/10.1016/S0141-1136(96)00103-1)
- 1024 Viarengo, A., Burlando, B., Evangelisti, V., Mozzone, S., Dondero, F., 2001. Sensitivity and specificity
1025 of metallothionein as a biomarker for aquatic environment biomonitoring. Biomarkers in
1026 Marine Organisms, A Practical Approach. Ed. Elsevier; Chapt. 2, pp 29-43.
1027 <https://doi.org/10.1016/B978-044482913-9/50004-3>
- 1028 Viarengo, A., Lowe, D., Bolognesi, C., Fabbri, E., Koehler, A., 2007a. The use of biomarkers in
1029 biomonitoring: a 2-tier approach assessing the level of pollutant-induced stress syndrome in
1030 sentinel organisms. Comparative Biochemistry and Physiology Part C: Toxicology &
1031 Pharmacology 146(3), 281-300. <https://doi.org/10.1016/j.cbpc.2007.04.011>.
- 1032 Viarengo, A., Dondero, F., Pampanin, D.M., Fabbri, R., Poggi, E., Malizia, M., Bolognesi, C., Perrone,
1033 E., Gollo, E., Cossa, G.P., 2007b. A biomonitoring study assessing the residual biological effects
1034 of pollution caused by the HAVEN wreck on marine organisms in the Ligurian sea (Italy). Arch.
1035 Environ. Contam. Toxicol., 53, 607-616. <https://doi.org/10.1007/s00244-005-0209-2>.
- 1036 Yigit, M., Celikkol, B., Yilmaz, S., Bulut, M., Ozalp, B., Dwyer, R.L., Maita, M., Kizilkaya, B., Yigit, Ü.,
1037 Ergün, S., Gürses, K., Buyukates, Y., 2018. Bioaccumulation of trace metals in Mediterranean

1038 mussels (*Mytilus galloprovincialis*) from a fish farm with copper-alloy mesh pens and potential
1039 risk assessment. Hum. Ecol. Risk Assess. 24(2), 465-481.
1040 <https://doi.org/10.1080/10807039.2017.1387476>
1041 Yuan, L., Richardson, C.J., Ho, M., Willis, C.W., Colman, B.P., Wiesner, M.R., 2018. Stress responses
1042 of aquatic plants to silver nanoparticles. Environ. Sci. Technol. 52, 2558-2565.
1043 <https://doi.org/10.1021/acs.est.7b05837>
1044 Zimmermann, S., Sures, B., 2018. Lessons learned from studies with the freshwater mussel
1045 *Dreissena polymorpha* exposed to platinum, palladium and rhodium. Sci. Total Environ. 615,
1046 1396-1405. <https://doi.org/10.1016/j.scitotenv.2017.09.204>
1047 Zuzolo, D., Cicchella, D., Doherty, A. L., Albanese, S., Lima, A., De Vivo, B., 2018. The distribution of
1048 precious metals (Au, Ag, Pt, and Pd) in the soils of the Campania Region (Italy). J. Geochem.
1049 Explor, 192, 33-44. <https://doi.org/10.1016/j.gexplo.2018.03.009>

1050
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring

Authors:

Greggio Nicolas* ^{a, b}, Capolupo Marco^{a, b}, Filippo Donnini^a, Manfred Birke^c, Fabbri Elena^{a, b}, Dinelli Enrico^{a, b}

^a University of Bologna, Inter-Departmental Research Centre for Environmental Science (CIRSA), Via S. Alberto 163, 48123 Ravenna, Italy

^b University of Bologna, Department of Biological, Geological, and Environmental Sciences, Piazza S. Donato 1, 40100 Bologna, Italy

^c Federal Institute for Geosciences and Natural Resources, Stilleweg 2, 30655, Hannover, Germany

* Corresponding Author: Nicolas Greggio e-mail: nicolas.greggio2@unibo.it.

University of Bologna, Inter-Departmental Research Centre for Environmental Science (CIRSA), Via S. Alberto 163, 48123 Ravenna, Italy.

Abstract:

Coastal lagoons are complex environments threatened by natural and anthropogenic stressors. Here, we tested the effectiveness of combining physical, geochemical and chemical measurements with biomarker data obtained in field-exposed marine mussels (*Mytilus galloprovincialis*) as a biomonitoring strategy for a highly pressured lagoon (Pialassa Baiona, Ravenna, Italy). Data showed a spatial trend of sediment contamination by Hg, Pt, Au, Ag, Mo, Re, Cd, Pd and Zn. Local conditions of high water temperature/low conductivity were detected among selected sites. After a 30-day in situ exposure, Ag and Hg were the most bioaccumulated elements (10 and 5 folds, respectively) in mussels followed by Sb, Al, Ti and Fe. Decreased survival, lysosomal dysfunctions, increased metallothionein content and peroxisome proliferation were observed in mussels in relation to metal spatial distribution and physico-chemical fluctuations. Overall, this study provides a further confirmation of the role of biomonitoring to reliably assess the environmental quality of highly pressured lagoons.

Keywords: Coastal lagoon; Sediment geochemistry; Metal bioaccumulation; Biomarkers; Biomonitoring;

Funding sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Formatted: Numbering: Continuous

Field Code Changed

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Introduction

Being at the boundary between terrestrial and marine ecosystems coastal lagoons are, by definition, unstable and complex systems morphologically dependent and shaped by natural continental evolution (rivers sediment transport and discharge), sea level variations, coastal dynamics, storminess and local geologic processes (e.g. subsidence) (Bird, 1994; Newton et al., 2014). In natural conditions, coastal lagoons receive fresh and nutrient-rich inputs from the land-side, that combined with shallow, warm and relatively stable water, sustain a complex food chain (from algae, to macro benthos, fish and swamps birds), providing a multitude of life-supporting (Aliaume et al., 2007; Dominik et al., 2014; Petry et al., 2016).

The naturally complicated equilibrium of these environments has often been impacted by strong anthropic development in the last century, especially in those lagoons close to major cities and harbours. With the beginning of demographic boom and of the correlated industrial development, lagoons have been converted into commercial ports or industrial areas and in the worst case as processing water disposal sites (Guerra, 2012; Guerra et al., 2014).

Water and sediment geochemistry of lagoon environments have been widely explored in order to define water status (Mollema et al., 2013; Greggio et al., 2020), sediment provenance and local background concentrations (Migani et al., 2015; Borghesi et al., 2016; Greggio et al., 2018a) or levels of occurring pollutants, notably heavy metals (Nazneen et al., 2019; Saidi et al., 2019), organic compounds (León et al., 2017; Mattes et al., 2018) and emerging contaminants (Pignotti et al., 2017; Pignotti and Dinelli, 2018).

The geochemical characterization of sediments and water is important, but alone often does not allow for an accurate and reliable evaluation of the environmental status of lagoon ecosystems (Carvalho et al., 2014; Dahms, 2014; Moreno-González et al., 2015). Physico-chemical properties and geological background of lagoons can influence the response of exposed organisms to pollutants, inducing potential interactive effects due to changes in bioavailability and toxicity (Migani et al., 2015; Piggott et al., 2015; Borghesi et al., 2016). Chemical assessments fail in detecting biotransformation products, eventually more toxic than the parental compounds (Buryškova et al., 2006; Braga et al., 2018), and in estimating synergistic/antagonistic effects induced on the fitness of the exposed biota (Lari et al., 2017).

For these considerations, multidisciplinary applications integrating geological/chemical parameters, pollutants bioaccumulation and biological effects may represent a promising approach to assess the environmental status of highly pressured coastal systems.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In the framework of biomonitoring programs of marine areas, particular attention has recently been addressed to the assessment of biomarkers of environmental stress (Turja et al., 2014; Capolupo et al., 2017; Mansour et al., 2020). Biomarkers are defined as indices of either exposure or effect of pollutants induced at biochemical, cellular and physiological level on exposed organisms (Viarengo et al., 2007a). They have successfully been used in association to bioaccumulation analyses to characterize “sub-organism” toxicological dynamics occurring in highly contaminated environments and provide crucial and early-warning information for planning urgent intervention or following the effects of remediation plans (Donnini et al., 2007; Capolupo et al., 2017). The simultaneous analysis of a wide set of biomarkers offers accurate information on the sub-lethal alterations induced by multiple stressors and may allow to correlate observed effects with specific classes of pollutants (Shaw et al., 2011; Turja et al., 2014). Moreover, biomarkers provide a unique contribution to determine the synergistic effect of pollutant mixtures, even when single contaminants occur at low concentrations. Biomarker-based assessments are thus currently promoted under the current EU legislation for monitoring surveys of marine and transitional environments (2000/60/EC; 2008/56/EC; Piló et al., 2017).

In the investigated area previous works mainly focused on organic pollutants and mercury in sediments (Trombini et al 2003; Guerra, 2012; Guerra et al., 2014). The bioaccumulation of PHAs in mussels (Fabbri et al., 2006) as well as mercury and methylmercury in clams (Trombini et al., 2003) are the only studies involving living organisms in the area. Inorganic pollutants are poor investigated in the area and when considered they are limited to the ordinary Cd, Cu, Hg, Ni, Pb and Zn in sediments (Matteucci et al., 2005; Guerra et al., 2014) without integration with biomarkers. Only a recent study by Capolupo et al. (2017) included, among others, metals bioaccumulation on mussels in an adjacent coastal lagoon. Moreover, a recent hydrogeological study reported water physico-chemical alterations and the abundance of seldom analysed metals (Ag and Sb among others) in surface drainage water entering in the investigated lagoon (Greggio et al., 2020).

For these reasons, the aim of this work was to evaluate the effectiveness of integrating water physico-chemical data, sediment geochemistry, metal bioaccumulation and biomarker measurements as a biomonitoring strategy for highly contaminated coastal lagoons. The study was conducted in a shallow and brackish basin located along the North-Western coast of the Adriatic Sea (Pialassa Baiona, Ravenna, Italy). In this area, previous works highlight the presence of organic pollutants in sediments (Fabbri et al., 2006; Guerra, 2012; Guerra et al., 2014; Trombini et al 2003; Vassura et al. 2005) and their potential for being bioaccumulated by mussels (Fabbri et al., 2006)

Formatted: English (United Kingdom)

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

1
2
3
4
5
6
7
8 and clams (Vassura et al., 2005). Inorganic pollutants were less investigated and the current
9 information is limited to few elements, as Cd, Cu, Hg, Ni, Pb and Zn (Matteucci et al., 2005; Trombini
10 et al., 2003; Guerra et al., 2014). A recent hydrogeological study reported water physico-chemical
11 alterations and the abundance of seldom analysed metals (Ag and Sb among others) in surface
12 drainage water flowing into the investigated lagoon (Greggio et al., 2020). Therefore, a wide
13 spectrum of elements was measured in sediments collected from different sites within the lagoon
14 and biological parameters were assessed in adult specimens of the Mediterranean mussel *Mytilus*
15 *galloprovincialis* transplanted into different sites in situ within at the same location the lagoon.
16 Finally, a battery of biological endpoints was measured in mussels, including, and encompassed
17 lysosomal endpoints responses, oxidative stress parameters and biomarkers of exposure to specific
18 classes of pollutants.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Font: Italic, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

2 Materials and methods

2.1 Study area and sampling sites

The Pialassa Baiona (Fig. 1) is a brackish lagoon located northward the town of Ravenna (Italy), between 44°27'55.5"N and 44°31'31.3"N and 12°14'9.7"E and 12°16'10"E that cover an area of about 10 km². The lagoon is part of a barrier-lagoon system that characterized the historical evolution of the Adriatic Sea coastline in the area (Amorosi et al., 1999). It is formed by small, shallow ponds (0.5–1 m depth) and deeper artificial channels (2–3 m depth) that permit the incessant water exchange with the Adriatic Sea driven by tides. It is connected to the Adriatic Sea only through the Candiano Channel that also is the Ravenna operative industrial harbour. The Pialassa Baiona is mainly adopted by local citizens as recreational fishing and hunting site and few clam farming activities exist in the northernmost portion.

The lagoon receives freshwaters from the west through three main channels (Fossatone, Cerba and Cupa-Magni channels) which drain agricultural lands and collect effluents from industrial and municipal treatment plants, located in the southernmost part (Fig. 1). The drainage water entering the lagoon are often rich in nutrients and dissolved metals (Mollema et al., 2013; Greggio et al., 2018b). The highest concentration of nutrients, as well as the highest bacteriological content, are observed in the southern part of the lagoon, in proximity of the via Cupa channel (Soprani et al., 1994). In addition, during the 60s and 70s, high quantities of chemical compounds were discharged

1
2
3
4
5
6
7
133 into the lagoon through the via Cupa and Magni channels, carrying wastewater from the industrial
134 area close to the southern border (Fabbri et al., 1998).

135
136 Previous investigations showed that high levels of metals, polycyclic aromatic hydrocarbons
137 (PAHs), polychlorinated biphenyls (PCBs) and other pollutants occurred in surface, subsurface and
138 suspended sediments with N-S increasing gradient (Trombini et al., 2003; Matteucci et al., 2005;
139 Guerra, 2012; Guerra et al., 2014). On these bases, seven sites were selected within the Pialassa
140 Baiona lagoon in order to cover its overall geochemical complexity and spatial trend of
141 anthropogenic pressures (Fig. 1). All sites (S1 to S7) were investigated for sediment geochemistry,
142 while metal bioaccumulation and biomarker responses were measured on mussels transplanted
143 into six sites (S1, S2, S3, S5, S6 and S7), since *in situ* transplantation into site S4 was not possible for
144 technical reasons. Mussels were exposed into the lagoon for 30 days, which is acknowledged as a
145 suitable time for sessile organisms to bioaccumulate contaminants at environmentally
146 representative levels and to develop physiological responses to "in site" pollution (Viarengo et al.,
147 2007b). Since previous investigations highlighted a lesser incidence of metals and organic
148 compounds in the north-western part of the lagoon (Trombini et al., 2003; Fabbri et al., 2006;
149 Donnini et al., 2007; Franzellitti et al., 2010), Site 7 (S7), located at about 0.5 km from the north-
150 western border, was defined as internal control site for both geochemical and biological
151 assessments.

152
153
154
155
156
157
158
159
160
161
162
163
164
165




Figure 1. Location map and sampling points (S1 to S7) within the Pialassa Baiona lagoon (Ravenna, Italy). The abbreviations "R." and "Ch." refer to the main rivers and/or channels which supply freshwater to the lagoon. The yellow shape represents the industrial area settled at the south-eastern boundary of the Pialassa Baiona lagoon.

2.2 Field activities and measurements

In order to have a better characterization of the environmental conditions influencing the organisms' response, measurements for physico-chemical parameters were conducted at each site.

During mussel field exposure (from June 1st to June 30th) six vertical profiles (10 cm resolution) of electrical conductivity, temperature and pH, using a CTD probe were conducted.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.3 Sediment geochemistry

The bottom sediments were sampled at each site with a manual corer using a Plexiglas tube. After the extrusion, the sections 0 - 5 cm was retained for analysis, placed in polyethylene bottles, were stored at -25 °C. For geochemical analysis, sediment aliquots were dried at 60 °C and homogenized by grinding in agate mortar. On the bottom sediments total elements concentration by X-ray fluorescence spectrometry (XRF) and pseudo-total (mobile/available) concentrations by aqua regia digestion (AR) have been performed.

The total metal content of 29 chemical elements (SiO₂, TiO₂, Al₂O₃, Fe₂O₃, MnO, MgO, CaO, Na₂O, K₂O, P₂O₅, S, As, Ba, Ce, Co, Cr, Cu, La, Nb, Ni, Pb, Rb, Sc, Sr, Th, V, Y, Zn, Zr) has been measured by XRF, using a Philips PW 1480 spectrometer equipped with a Rh tube on pressed powder pellets following matrix correction methods suggested by Franzini et al. (1972; 1975) Leoni and Saitta (1976) and Leoni et al. (1982). Thirty international reference materials were used for the instrument calibration, while accuracy was evaluated using 4 international reference samples, namely BR, BCR-1, TB and AGV-1 (Govindaraju, 1984). The average accuracy was higher than 5 % for trace-element determinations. In order to quantify the pseudo total concentration, 15 g of sediment were digested with a modified aqua regia solution of equal parts concentrated HCl, HNO₃ and deionized H₂O for one hour in a heating block of hot water bath at the Bureau Veritas Laboratories in Vancouver (Canada). Fifty-three chemical elements (Ag, Al, As, Au, B, Ba, Be, Bi, Ca, Cd, Ce, Co, Cr, Cs, Cu, Fe, Ga, Ge, Hf, Hg, In, K, La, Li, Mg, Mn, Mo, Na, Nb, Ni, P, Pb, Pd, Pt, Rb, Re, S, Sb, Sc, Se, Sn, Sr, Ta, Te, Th, Ti, Tl, U, V, W, Y, Zn, Zr) were analysed by Inductively Coupled Plasma Mass Spectrometry (ICP-MS).

Field Code Changed

The volatile content (including humidity, organic matter, water incorporated in the lattice of clay minerals, carbon dioxide in the carbonate minerals) was evaluated by thermal analysis using a Setaram TAG24 double furnace apparatus, with simultaneous registration of thermogravimetric (TG), derivative thermogravimetric (DTG), differential thermal analysis (DTA). A CO₂ atmosphere was used to increase the temperature of carbonate decomposition and the carbonate content was estimated by quantifying the weight loss at temperature > 700 °C.

2.4 Mussel collection and handling

Adult specimens of the Mediterranean mussel (*M. galloprovincialis*) with a shell-length of 5 ± 1 cm were collected by professional fishermen of the "Cooperativa CONISUB" (Marina di Ravenna,

1
2
3
4
5
6
7
193 Italy) along the north-western Adriatic Sea coast in high quality marine areas (Zone A) and
194 immediately transferred to the laboratory in seawater tanks with continuous aeration. Prior to the
195 *in situ* deployment, all animals were acclimatized for 6 days in aquaria containing 60 L of aerated
196 natural seawater (at a ratio of 1 mussel/L) at 16 °C and natural photoperiod, and fed daily on an
197 algal slurry (Coral Diet Filtrator, Xaquia, Italy) after water renewal. Seventy organisms were dissected
198 prior to the *in situ* transplantations (time 0, T0) for bioaccumulation and biological endpoint
199 analyses. Digestive glands and gills were immediately collected, frozen in liquid nitrogen and stored
200 at -80 °C until analysed. At the same time six groups of seventy organisms each were put in fishnet
201 bags, fixed to piles and driven into the sediment at sites S1, S2, S3, S5, S6 and S7 (Fig. 1) at about 50
202 cm from the bottom to ensure continuous submersion during the whole exposure period. Deployed
203 mussels were regularly controlled at three days intervals and dead mussels were eventually
204 recorded for survival assessment. Since previous investigations identified the Magni channel as a
205 heavily polluted site (Fabbri et al., 2000; 2003; Matteucci et al., 2005), three additional fishnet bags,
206 with seventy organisms each, were transplanted into S1 to execute biomarkers with shorter
207 timestep.

209 2.5 Metal bioaccumulation analysis

210 Metal bioaccumulation was evaluated on mussels prior to the *in situ* deployment (Time 0, T0)
211 and after 30 days of field exposure into the lagoon. Metal bioaccumulation from each site was used
212 to calculate the bioaccumulation factor as enrichment/depletion compared to T0. Samples (0.2 g
213 each) were digested in a microwave system with ultrapure HNO₃. The resulting fluid was diluted by
214 adding deionized water to a final volume of 30 mL, at the Department of Environmental Science of
215 the University of Siena. Analytical blanks and certified reference material were included in the
216 analysis, specifically the mussel tissue ERM-CE278 (*Mytilus edulis*) from the Institute for Reference
217 Materials and Measurements, Geel, Belgium.

218 The samples were analysed by inductively coupled plasma quadrupole mass spectrometry
219 (ICP-QMS) using an Agilent 7500ce instrument at the laboratories of the Federal Institute for
220 Geosciences and Natural Resources in Berlin. Atomic fluorescence spectrometry was used for the
221 determination of mercury (Instrument PSA 10.035 Millennium Merlin 1631). The Table S1 reports
222 the results for the investigated elements in the certified reference material. Additional details can
223 be found in Capolupo et al. (2017).

1
2
3
4
5
6
7
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265

2.6 *Biological endpoints*

2.6.1 Survival percentage and Stress on Stress

The survival (SURV) was calculated as the percentage of mussels found alive over the exposure period with respect to the initial number of individuals (70 at each site).

For Stress on Stress (SOS) assessment, immediately after collection, 15 mussels per site were subjected to anoxia by air exposure at 15 °C in a humidified chamber. Death symptoms were open valves and absence of muscular activity. The survival was assessed daily, and results expressed as the time at which 100% mortality was achieved (Viarengo et al., 2007a).

2.6.2 Cytochemical parameters

Lysosomal membrane stability (LMS), lysosome to cytoplasm volume ratio (LYS/CYT) and lipofuscin content (LIF) were assessed in the digestive gland from 20 mussels per each site (N = 20). After dissection, glands were mounted on specific aluminium supports for cryostat microtomy (chucks), frozen in N-hexane at -70 °C, and stored at -80 °C. All assessments were performed according to the UNEP/RAMOGGE manual (1999) on 10 µm-thick gland sections obtained using a cryostat (MICROM HM 505 N) at -30 °C. The LMS was measured as the time necessary to produce the maximum staining reaction between the lysosomal hydrolase N-acetyl-β-hexosaminidase and a specific substrate (Capolupo et al., 2016). Briefly, sections were incubated at 37 °C in a shaking water bath with a destabilization buffer (0.1 M Na-citrate, pH 4.5) for different times (0, 3, 5, 10, 15, 20, 30 and 40 min), exposed to the substrate naphtol AS-BI N-acetyl-β-D-glucosaminide (Sigma, N4006) for 20 min and finally stained with the diazonium dye Fast Violet B (Sigma, F1631) (1 mg/ml in 0.1 M phosphate buffer, pH 7.4). For each site, slides not exposed to the destabilization buffer (0 min) were screened for the LYS/CYT assessment, which was determined by evaluating the cytoplasmic and lysosomal areas according to Capolupo et al. (2017). The LIF determination was performed according to Martin-Diaz et al. (2009). Gland sections were fixed in calcium formol and immersed for 5 min in an aqueous solution of 1 % ferric chloride and 1 % potassium ferricyanide in a 3:1 ratio.

For all lysosomal parameters, tissue sections were assessed under a light microscope (Axioskop 40, Carl Zeiss, Milan, Italy) equipped with a 40X objective and a digital camera (AxioCam MRC, Carl Zeiss, Milan, Italy) as reported by Capolupo et al. (2016). Results were expressed as labilization period for LMS, percentage of lysosomal volume inside digestive cells for LYS/CYT and arbitrary units for LIF.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.6.3 Enzymatic biomarkers

The acetylcholinesterase activity (AChE) activity was determined in mussel gills as previously reported (Valbonesi et al., 2003). Gills from 5 mussels per site (N = 5) were homogenized in a 0.1 M phosphate buffer, pH 7.4, and centrifuged at 9000 ×g at 4 °C for 30 min. Supernatant aliquots were incubated at 25 °C with 0.5 mM acetylthiocholine iodide (ASCh) and 0.33 mM DTNB (Ellman et al., 1961), and read at 405 nm for 10 min. Results are reported as nmoles/min/mg protein. The sample protein concentration was measured according to Lowry et al. (1951).

For catalase (CAT) assessment, about 50 mg of digestive gland was dissected from 5 mussels per site (N = 5), homogenized in 50 mM potassium-phosphate buffer (pH 7.0, 0.5 mM Na₂EDTA) and centrifuged at 15,000 g at 4 °C. The CAT activity was determined by measuring the time-dependent decrease of absorbance at 240 nm in the presence of 55 mM H₂O₂. Data were expressed as μmol/min/mg of total protein.

The Peroxisomal acyl-CoA oxidase (AOX) activity was determined on five pools of 2 digestive glands per site (N = 5) according to Orbea et al. (2006). Tissues were homogenized in four volumes of TVBE buffer (1 mM sodium bicarbonate, 1 mM EDTA, 0.1 % ethanol and 0.01 % Triton X-100), pH 7.6 and centrifuged at 500 × g for 15 min at 4 °C. Supernatants were diluted 1:10 in TVBE buffer and spectrophotometrically assayed for AOX activity at a wavelength of 502 nm. Data were expressed as milliunits/mg of AOX protein (equivalent to nmol H₂O₂/min x mg protein).

2.6.4 Metallothionein content (MT)

The metallothionein (MT) content was analysed on five pools of 5 mussel digestive glands (about 1.5 g of tissue) per site according to Viarengo et al. (1997). Final absorbance was measured at 410 nm, using reduced glutathione as reference standard. Data were expressed as μg of MT/mg tissue.

2.6.5 Mussel Expert System

Data from biological responses were integrated using the Mussel Expert System (MES) developed by Dagnino et al. (2007) in a unique and synoptic five-level Health Status Index (HSI) ranging from A (healthy) to E (pathological status). The system classification is based on (i) the number of biomarkers altered in the exposed organisms (significant changes greater than 20 % were utilized to evaluate alterations in the health status of the organisms); (ii) the magnitude of the alteration; (iii) the role of the autophagic process and (iv) the level of biological organization affected

(i.e., cell, tissue and organism level). For each biomarker/site, the system calculates an Alteration Factor (AF), which is the ratio between biomarker levels from treatments and control, and an Alteration Level (AL), which defines the magnitude of observed alterations based on the AFs magnitude and the biomarker toxicological profile (i.e. increasing, decreasing, bell-shaped). ALs and AFs are then processed by the system applying a set of rules in the "if...then" form in order to calculate the overall HSI associated to the exposure to each site. The simultaneous alterations of general stress parameters (i.e. lysosomal responses) and biomarkers of exposure to specific classes of compounds (e.g. MT for metals; AOX for POPs) is further computed and eventually recognized by the system as an early warning signal regarding the presence or excess of those pollutants in the monitored sites or tested condition.

Formatted: Highlight

Formatted: Highlight

2.7 Data analysis

All data were tested for normal distribution using the Shapiro-Wilk test, while the Levene's test was employed to test the equal variance. Significant differences between S1 to S6 compared to the control site S7 were estimated by a One-Way ANOVA (SigmaStat, SPSS) followed by the Duncan's post-hoc test. The evaluation of potential linear relationships between biological endpoints and physico-chemical parameters (including pH, temperature, electrical conductivity, metal concentrations in sediments and mussel tissues) were conducted through the Pearson's correlation analysis. For all performed tests, significance was established at $p < 0.05$. All analyses were performed using the software SigmaPlot 13 (Systat Software Inc. San Jose, CA, USA).

3 Results and discussion

3.1 Physico-chemical data

The vertical profiles of temperature, pH and electrical conductivity (EC) measured during mussels field exposure as well as a graphical representation of the temperature, pH and EC vertical profiles for June 14th are presented respectively in Table S2 and in Figure S1 of supplementary material. The site S1, located in the Magni Channel at the southern side of the lagoon (Fig. 1), is different from the other sites for all monitored parameters. It shows higher water column temperature, reduced EC and a lower pH value, for the entire monitoring period (Table S2). Specifically, S1 shows the lowest EC among the sites with a marked vertical gradient (Fig. S1A) and

1
2
3
4
5
6
7
8 it is also the site with the highest temperature, up to 32.3 °C. Moreover, it records the highest
9 average profile temperature during the surveying period, at least 2 °C constantly higher compared
10 to all the other sites (Fig. S1B). Site S1 receives freshwater from artificial drainage channel network
11 surrounding the town of Ravenna (Greggio et al., 2018b). Previous investigations also reported
12 water temperature fluctuation wider (10-24 °C) in the southernmost channel with respect to the
13 lagoon average (16-18 °C), which was subjected to variations of about 10 °C within the 24 hours
14 owing to discharges of cooling water from the nearby industrial site (Franzellitti et al., 2010). A
15 thermal vertical gradient is also present in S5 probably associated to local limited water circulation
16 favouring water warming up (Fig. S1A). The pH is the lowest in S1, ranging from 7.2 to 7.8, and with
17 a marked vertical profile. In all the other sites the pH is instead similar and fairly constant around 8.
18 Only at S5 the observed pH values are higher (8.0-8.3) for the entire surveying period (Table S2).

25 3.2 Sediment data

27 The results of the geochemical analyses are presented in Table 1. According to the
28 geochemical composition of the major elements (expressed as %), and referencing to the work of
29 Dinelli et al. (2007) which compared geochemical proxies with grain size analysing borehole samples
30 in the same geographic area, the Pialassa Baiona bed sediment samples are classified as fine
31 grained-sediments (with a recalculated $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio < 3) enriched in carbonates. The sediments
32 are also rich in organic matter (OM in Table 1, ranging from 1.1 to 3.4) as already reported for
33 adjacent coastal lagoons (Migani et al., 2015). The bulk geochemical composition is quite
34 homogeneous among the different sites, with S1 being slightly enriched in carbonates (Table 1).

35 The Pialassa Baiona bed sediments composition is a mixture of carbonate and silicates with
36 variable amount of organic matter. The geochemical results, in terms of total sediment composition,
37 are comparable with those reported by Donnini et al. (2007) although the number of sites was
38 different. As far as total concentrations are concerned, these are within the range of regional
39 background, as proposed by Migani et al. (2015), taking into consideration Holocene data from
40 boreholes in the area not affected by anthropogenic inputs (Amorosi et al., 2002; Curzi et al., 2006;
41 Dinelli et al., 2012, Greggio et al., 2018a). Except for sites S6 and S7, Zn and Cu concentrations
42 exceed the maximum value observed in the background sediments, respectively of 124 and 41
43 mg/kg (Table 1). In particular, for S1 site Zn and Cu total concentrations are double respect to
44 background, respectively 316 and 87 mg/kg, in line with concentration detected in the neighbouring
45 Pialassa Piomboni by Pignotti et al. (2018). Pb concentrations in sites S3 and S5 are 30 mg/kg being

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom), Subscript

Formatted: English (United Kingdom)

Formatted: English (United Kingdom), Subscript

Formatted: English (United Kingdom)

Formatted: English (United Kingdom), Subscript

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Highlight

1
2
3
4
5
6
7
8 slightly higher respect to background level established at 29 mg/kg by Migani et al. (2015), but
9 certainly lower than Pb concentration in the Pialassa Piomboni by Pignotti et al. (2018) where a
10 maximum concentration of 257 mg/kg was measured.
11

12
13 **Table 1. Results of the chemical analyses on sediments. The table includes either pseudo total Aqua Regia (AR) and**
14 **total XRF data. Table includes also the ranges observed in boreholes in the area, considered representative of the**
15 **natural background (Amorosi et al., 2002; Curzi et al., 2006). Bold elements exceed natural background. AR DL**
16 **indicates Detection Limit for Aqua Regia extraction.**

		S1		S2		S3		S4		S5		S6		S7		Natural background	
		AR DL	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR		XRF
Ag	µg/kg	2	991	-	269	-	268	-	173	-	275	-	179	-	62	-	
Al	%	0.01	1.39	6.31	1.41	7.02	1.23	6.98	1.10	6.97	1.35	7.09	1.39	7.07	1.41	6.55	4.4-9.7
As	mg/kg	0.1	7.8	-	5.9	-	6.1	-	5.9	-	7.8	-	6.8	-	6.7	-	-
Au	µg/kg	0.2	16.2	-	5.4	-	11.1	-	2.6	-	5.1	-	3.4	-	1	-	-
B	mg/kg	1	25	-	28	-	26	-	24	-	38	-	33	-	22	-	-
Ba	mg/kg	0.5	61.9	263	43.7	218	41.8	260	42.6	322	40.1	258	38.7	294	36.5	258	220-530
Be	mg/kg	0.1	0.4	-	0.7	-	0.7	-	0.8	-	0.5	-	0.6	-	0.6	-	-
Bi	mg/kg	0.02	0.6	-	0.38	-	0.35	-	0.26	-	0.38	-	0.31	-	0.28	-	-
Ca	%	0.01	11.27	11.37	7.45	7.51	7.87	7.96	7.34	7.87	6.10	6.65	7.24	7.43	8.61	9.52	0.3-17.2
Cd	mg/kg	0.01	0.68	-	0.29	-	0.33	-	0.21	-	0.5	-	0.35	-	0.11	-	-
Ce	mg/kg	0.1	19.4	42	17	55	17.6	71	16.8	52	17.8	64	18.3	42	20.1	58	14-83
Co	mg/kg	0.1	10.2	12	10.9	11	10.7	12	9.9	11	11.7	14	11.5	12	10.6	11	5-29
Cr	mg/kg	0.5	70.6	131	57.5	103	51.4	132	47.7	131	59.9	138	59.3	130	47	100	96-270
Cs	mg/kg	0.02	1.36	-	1.15	-	1.08	-	1.16	-	1.11	-	1.39	-	1.57	-	-
Cu	mg/kg	0.01	84.1	87	49.1	52	49.5	53	36.7	44	51.1	52	37.5	38	22.4	24	4-41
Fe	%	0.01	2.17	3.42	2.46	3.92	2.23	3.86	2.00	3.69	2.54	3.83	2.30	3.64	2.36	3.72	2.1-5.3
Ga	mg/kg	0.1	4.4	-	4.2	-	4	-	3.9	-	4	-	4.2	-	4.4	-	-
Ge	mg/kg	0.1	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	-
Hf	mg/kg	0.02	0.03	-	0.03	-	0.04	-	0.03	-	0.03	-	0.03	-	0.05	-	-
Hg	mg/kg	0.005	25.7	-	1.42	-	1.48	-	1.89	-	2.38	-	2.26	-	0.1	-	-
In	mg/kg	0.02	0.03	-	0.03	-	0.02	-	0.02	-	0.03	-	0.02	-	0.03	-	-
K	%	0.01	0.33	1.77	0.36	1.98	0.31	2.06	0.28	2.04	0.39	1.97	0.36	2.04	0.35	2.03	0.9-3.3
La	mg/kg	0.5	8.3	19	7.7	26	7.3	22	6.9	18	8	20	8.5	34	9.5	20	17-47
Li	mg/kg	0.1	26.1	-	30.6	-	27.8	-	24.9	-	28.7	-	28	-	28.1	-	-
Mg	%	0.01	1.37	2.41	1.34	2.38	1.30	2.38	1.22	2.44	1.42	2.54	1.55	2.55	1.58	2.35	2.2-4.2
Mn	mg/kg	1	531	893	527	865	549	896	513	949	511	801	550	897	644	1396	230-1470
Mo	mg/kg	0.01	2.67	-	3.33	-	2.61	-	2.28	-	3.34	-	1.57	-	0.54	-	-
Na	%	0.001	0.85	0.68	0.98	0.69	0.90	0.75	0.65	0.99	1.18	0.67	0.85	0.78	0.42	0.85	0.5-2.1
Nb	mg/kg	0.02	0.37	15	0.28	13	0.25	13	0.29	14	0.34	16	0.37	10	0.29	13	8-20
Ni	mg/kg	0.1	50.9	73	56.5	73	50.7	88	46.3	72	55.3	82	51.5	74	46.4	57	39-148
P	%	0.001	0.11	0.10	0.06	0.06	0.06	0.07	0.06	0.06	0.08	0.07	0.06	0.06	0.06	0.06	0.03-0.08
Pb	mg/kg	0.01	27.92	28	22.71	26	22.14	30	18.16	19	26.22	30	20.71	22	11.99	18	5-29
Pd	µg/kg	10	46	-	10	-	12	-	11	-	10	-	<10	-	<10	-	-
Pt	µg/kg	2	37	-	6	-	4	-	4	-	4	-	<2	-	<2	-	-
Rb	mg/kg	0.1	24.9	113	22.2	128	20.8	128	22.4	113	22.3	130	25.1	114	29.8	89	56-200
Re	µg/kg	1	7	-	1	-	2	-	<1	-	2	-	2	-	<1	-	-
S	%	0.02	0.6	-	0.55	-	0.6	-	0.54	-	0.72	-	0.71	-	0.34	-	-
Sb	mg/kg	0.02	0.51	-	0.29	-	0.25	-	0.19	-	0.37	-	0.29	-	0.19	-	-
Sc	mg/kg	0.1	3.4	11	3.6	11	3.4	6	3.2	8	3.7	10	3.7	8	4.1	13	12-21
Se	mg/kg	0.1	0.9	-	0.7	-	0.7	-	0.5	-	1.1	-	0.7	-	0.5	-	-
Si	%	-	-	17.72	-	19.43	-	19.90	-	20.61	-	19.31	-	20.95	-	20.03	14.7-29.3
Sn	mg/kg	0.1	3.2	-	1.8	-	1.5	-	1.3	-	1.9	-	1.5	-	0.8	-	-
Sr	mg/kg	0.5	353	368	274.4	300	224.8	252	218	265	259.1	281	281.6	291	290.5	310	113-442
Th	mg/kg	0.1	3.5	4	4.1	5	3.6	17	3.6	25	4	11	4.2	10	4.7	7	<3-19
Ti	%	0.001	0.009	0.308	0.005	0.328	0.006	0.347	0.008	0.349	0.006	0.324	0.010	0.334	0.015	0.336	0.02-0.58
Tl	mg/kg	0.02	0.26	-	0.17	-	0.21	-	0.19	-	0.27	-	0.2	-	0.14	-	-
U	mg/kg	0.1	1.3	-	1.5	-	1.2	-	1.1	-	1.3	-	1.2	-	1.1	-	-
V	mg/kg	2	32	85	33	85	30	99	26	94	35	113	33	94	31	76	54-169
Y	mg/kg	0.01	10.02	18	9.39	23	9.28	24	8.86	20	9.7	24	9.63	19	10.58	19	18-43
Zn	mg/kg	0.1	306.6	316	129.8	152	133.1	154	101.2	126	146.9	174	109.3	124	58.2	67	43-124
Zr	mg/kg	0.1	1.1	108	1.2	91	1	105	1	124	1.1	86	1.2	92	1.9	106	76-260
H ₂ O ⁻	%	-	-	2.7	-	4.0	-	2.8	-	2.4	-	3.7	-	2.7	-	2.1	-
OM	%	-	-	2.8	-	1.8	-	2.0	-	1.7	-	3.4	-	2.2	-	1.1	-
H ₂ O ⁺	%	-	-	6.0	-	6.5	-	5.4	-	4.4	-	6.5	-	5.8	-	2.3	-
CO ₂	%	-	-	11.6	-	8.9	-	9.1	-	9.3	-	8.6	-	7.4	-	12.3	-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

LOI	%	-	-	23.1	-	21.2	-	19.3	-	17.8	-	22.3	-	18.1	-	17.8	4.8-29.3
-----	---	---	---	------	---	------	---	------	---	------	---	------	---	------	---	------	----------

As regards the Aqua Regia extraction, data showed the presence of elements seldom analysed in sediments of the area. The most striking features are the differences observed between S1 and all the other sites for a number of elements, in particular Ag, As, Au, Bi, Cd, Cr, Cu, Hg, Pb, Pd, Pt, Re, Zn (Table 1). According to Donnini et al. (2007), S7 should be considered as unpolluted site and if its results are taken as reference, a strong enrichment is clear for several elements (Fig. 2). Mercury, which is a well-known pollutant in the area (Misericocchi et al., 1993; Fabbri et al., 1998; Fabbri et al., 2001a, b; Trombini et al., 2003; Matteucci et al., 2005; Guerra et al., 2007; Covelli et al., 2011; Guerra, 2012; Dominik et al., 2014; Borghesi et al., 2016), is the element with the largest enrichment (up to 250 times compared to S7) and reaches relatively high concentrations in the Magni Channel site (S1, up to 25 mg/kg). Mercury dynamics in the lagoon is complex and although the maxima in concentrations are not at the surface, the element is heavily mobilized at the water/sediment interface in the southern part of the lagoon (Covelli et al., 2011). Borghesi et al. (2016) highlighted similar behaviour among Hg and Ag, Au, Cd, Cu and Zn for six different coastal lagoons in the same Region due to metal affinity for the fine fraction of the sediment.

Notable enrichments are then evident in Figure 2 for valuable metals like Pt, Au, Ag, Re and Pd, which showed substantial increase (up to 10-fold) compared to S7 and to median values reported by Cicchella et al. (2015), that represent a useful reference for metal concentrations in mineral matrix obtained with the same analytical technique and at the same laboratory. In particular, Platinum and Palladium are of recent technogenic use and their anthropogenic fluxes greatly exceed the natural ones (Ruchter and Surer, 2015; McGillicuddy et al., 2017). Platinum and Palladium are largely used in automobile catalytic converters (Bossi and Gediga, 2017) and their dispersion increases particularly around urban areas in roadside soils (Ravindra et al., 2004; Ruchter and Surer, 2015; Leopold et al., 2018; Zuzolo et al., 2018), but also in marine areas close to urban settings (Cobelo-García et al., 2011; Abdou et al., 2016).

The Silver concentration reaches a peak at S1 with 0.99 mg/kg, at least four times more than the other sites in the Pialassa Baiona and ten times more than other coastal lagoons in the same Region as resulted by Borghesi et al. (2016). The Ag enrichment is controlled by many factors such as pH, OM, as well as the source material. At typical environmental pH values, Ag is adsorbed onto Fe hydroxides, in preference to Cu and Zn (Lottermoser et al., 1999). In the near Venice lagoon, Giusti and Zang (2002) measured Ag sediment concentrations 5-6 fold higher than this study,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

justifying the abundance with both famous glassmaking firms in Murano island and the large volume of untreated sewage sludges discharged into the lagoon from the nearby Marghera industrial area.

Moderate enrichments (four times compared to S7) are shown by Mo, Cd, Zn, Sn and Cu that can be more directly linked to industrial effluents as reported by Giusti and Zang (2002) in Venice lagoon and Borghesi et al. (2016) for the same Region. Enrichments in all these elements also characterize the other sites, although at an order of magnitude lower than S1 (around 2 times).

Many of the metals detected in sediments show a decreasing concentration with increasing distance from the industrial area located in the southern part of the lagoon. This suggests that industrial discharge effluents play a major role in the lagoon contamination compared to effluents from urban and agricultural activities. In fact, the distribution of some elements, including Zn, Cr, Cu, Mo, Ag, and Hg, follow trends similar to those reported for other organic pollutants of industrial origin in the same area (Fabbri et al. 2006; Vassura et al. 2005). Vassura et al.(2005), for instance, detected a clear South-North increasing pattern of prioritized PAHs, with sediments from the Magni channel area (the present S1) showing concentrations of pyrene, fluoranthene, anthracene and phenanthrene up to 10,000 times higher compared to the northern sites.




Figure 2. Metal enrichment in sediments from selected sites within the Pialassa Baiona lagoon. Data are normalized on Site 7 (S7), which was selected as internal reference based on previous literature (Trombini et al., 2003; Fabbri et al., 2006; Donnini et al., 2007; Franzellitti et al., 2010). Elements in italics (*Pt, Au, Re, Pd*) were not investigated in the mussel tissues.

1
2
3
4
5
6
7
8 3.3 Metal bioaccumulation

9 Results of the chemical analyses on mussel tissues are presented in Table 2. Data were
10 obtained for all sites except S1, which was characterized by a high mortality rate (more details in
11 section 3.4). T0 data, originating from mussels growing in the Adriatic Sea, are in line with basal level
12 for As, Ba, Cd, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, V, Zn identified by Fattorini et al. (2008). The trend of
13 concentrations of trace elements in soft tissue of *Mytilus galloprovincialis* belonging to T0 is (Zn, Fe)
14 > (As, Mn, Cu, Ni, Pb) > (Cd, Cr, Ag) > (Hg, Sb). The trend is confirmed also for average trace elements
15 concentration among all sites: (Fe, Zn) > (As, Mn, Cu) > (Pb, Ag, Ni, Cd, Cr, Hg) > Sb. The only notable
16 change is the Ag enrichment from 0.12 mg/kg in T0 tissues to a mean concentration of 0.80 mg/kg
17 in tissues from experimental sites. Although Cd, Hg, Sb were not included, the same abundance
18 trend was found by Giusti and Zang (2002) in the Venice Lagoon: Fe > Zn > (Cu, Mn, As) > Ag > Pb >
19 (Ni, Cr). Absolute concentrations of Zn, Cu, Pb and Cd found in this study (Table 2) fall in the lowest
20 part of the ranges summarized by Benali et al. (2017) for 10 locations in Mediterranean Sea.
21
22
23
24
25

26
27
28 **Table 2. Results of the chemical analyses on mussel tissues (bioaccumulation). Data refers to T0 and to the 5 sites
29 were mussels survived or remained after the exposure period.**

	unit	T0	S2	S3	S5	S6	S7	
30	Ag	mg/kg	0.12	0.99	0.98	1.21	0.62	0.82
31	Al	mg/kg	36	39	56	98	82	47
32	As	mg/kg	9.83	12.8	9.98	11.9	12.4	11.4
33	B	mg/kg	25.7	22.9	19.7	22.8	25.9	19.4
34	Ba	mg/kg	1.95	0.35	0.49	0.54	0.52	0.36
35	Be	mg/kg	0.003	0.003	0.003	0.005	0.006	0.004
36	Bi	mg/kg	0.057	0.009	0.009	0.007	0.011	0.008
37	Ca	mg/kg	2004	1322	1299	1726	1679	1157
38	Cd	mg/kg	0.70	0.71	0.78	0.76	0.77	0.50
39	Ce	mg/kg	0.118	0.051	0.092	0.124	0.166	0.075
40	Co	mg/kg	0.54	0.323	0.3	0.303	0.511	0.315
41	Cr	mg/kg	0.57	0.6	0.53	0.67	0.53	0.64
42	Cs	mg/kg	0.022	0.014	0.016	0.022	0.019	0.015
43	Cu	mg/kg	4.65	2.71	2.63	2.81	2.66	1.46
44	Fe	mg/kg	67.5	68.1	77.4	116.0	103.4	91.7
45	Ga	mg/kg	0.018	0.007	0.011	0.019	0.013	0.009
46	Hg	mg/kg	0.082	0.35	0.32	0.41	0.27	0.16
47	K	mg/kg	11084	10988	9729	11129	11244	9535
48	La	mg/kg	0.138	0.035	0.057	0.074	0.109	0.06
49	Li	mg/kg	0.88	0.46	0.41	0.51	0.58	0.34
50	Mg	mg/kg	4198	3851	3467	3676	4422	2506
51	Mn	mg/kg	4.84	5.19	7.50	8.16	8.03	5.03
52	Mo	mg/kg	0.645	0.353	0.368	0.317	0.394	0.274
53	Na	mg/kg	22885	20201	15986	19714	24751	13438
54	Nb	mg/kg	0.002	0.001	0.002	0.003	0.003	0.002
55	Ni	mg/kg	3.61	0.72	0.69	0.73	0.92	0.64
56	Pb	mg/kg	1.41	0.87	0.52	0.57	0.94	0.64
57	Rb	mg/kg	5.29	4.08	4.32	4.27	4.3	3.78
58	Sb	mg/kg	0.018	0.038	0.031	0.03	0.048	0.039
59	Sc	mg/kg	0.04	0.04	0.03	0.05	0.07	0.04
60	Se	mg/kg	2.27	2.10	2.10	2.01	1.83	1.52
61	Sn	mg/kg	0.12	0.12	0.12	0.10	0.09	0.08

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Sr	mg/kg	31.2	17.7	16.5	20.6	22.8	14.3
Th	mg/kg	0.019	0.011	0.014	0.025	0.031	0.016
Ti	mg/kg	0.49	0.34	0.62	1.07	0.88	0.50
Tl	mg/kg	0.003	0.001	0.002	0.002	0.001	0.001
U	mg/kg	0.127	0.049	0.040	0.054	0.065	0.032
V	mg/kg	1.15	0.44	0.49	0.55	0.80	0.47
Y	mg/kg	0.080	0.046	0.071	0.085	0.094	0.04
Zn	mg/kg	107.1	116	62.4	95.6	134	86.9
Zr	mg/kg	0.064	0.013	0.025	0.025	0.021	0.018

The bioaccumulation factor is graphically showed in Figure 3. Enrichments are notable for Ag, Hg and Sb in all the sites, with higher bioaccumulation at S5 located in the centre of the lagoon in a pond close to the outlet of a canal from the southern part of the lagoon, the most contaminated one. Although not high as those measured in mussels from other sites such as Northern England (Giusti et al., 1999), these three elements could be the most critical since other elements known for toxicity to mussels (e.g. Cd, Pb, Ni) (Benali et al. 2017; Yigit et al. 2017) are similar or depleted compared to T0 (Fig. 3).

The enrichment in mercury was relatively high, as expected and already verified both in living clams (Trombini et al., 2003) and mussels (Fabbri et al., 2006; Donnini et al., 2007). Moreover, the Hg cycling at the sediment-water interface (Covelli et al., 2011), could be a reason for the mercury enrichment observed in all sediment samples and may also reflect the high Hg concentrations observed in mussels exposed at S5 (Fig. 3).

Silver has recently attracted much interest since its increasing application as bactericidal and usage in numerous consumer products lead to contamination of aquatic ecosystems, often in form of Ag nanoparticles (Yuan et al., 2018; Outa et al., 2020). Silver speciation in environment is dependent by local environmental conditions and usually forms stable complexes with Lewis bases such as amines, halides, and thiolates and with dissolved organic matter altering its toxicity (Mousavi et al., 2015). In Pialassa Baiona, Greggio et al. (2020) showed dissolved concentrations ranging from 0.02 to 10 µg/L, with higher concentrations in basins with drainage water input. Concerning bioaccumulation, recent works reported the effect of Ag nanoparticles on soft tissues of mussels. *Mytilus galloprovincialis* accumulated Ag in digestive gland tubule cells and gills during the entire year (Duroudier et al., 2019) and intralysosomal metal accumulation and lysosomal membrane destabilisation were observed (Jimeno-Romero et al., 2017). Compare to this study (max 1.21 mg/kg dry wt), Giusti and Zang (2002) in Venice lagoon found silver concentrations up to 6.2 mg/kg dry wt that are aligned with outcomes in other species of the genus *Mytilus* as reported by Martin et al. (1988).

Field Code Changed

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Antimony is pervasive in synthetic polymers (including textiles, plastics and rubbers) (Filella et al., 2019) and it is widely applied in alloy industries as flame retardant, as pigments for colour or colour protection (Filella et al., 2002; James and Turner 2020). Although rock weathering and soil runoff supply Sb to the environment, anthropogenic source is crucial. James and Turner (2020) demonstrated that Sb could also be mobilised via digestion and bioturbation by deposit-feeders in sediment contaminated by plastic or directly adsorbed from microplastics taken up by mussels (Van Cauwenberghe et al., 2015). Antimony concentrations in unpolluted waters are less than 1 µg/L (Filella et al., 2002). Greggio et al. (2020) reported Sb water concentrations in western ponds of the Pialassa Baiona ranging from 0.05 to 0.5 µg/L. Limited is the knowledge of Antimony concentration in mussels. In this study Sb in mussel tissues ranged from 0.01 to 0.05 mg/kg (Table 2) with a 2-3 as enrichment factor (Fig. 3). De Gregori et al. (2007) found tissue concentration ranging from 0.007 to 0.06 mg/kg in unpolluted area, while 0.23 to 1.030 mg/kg in harbour dock protected by antimony–lead alloy cover. Garcia (2015) measured Sb concentration ranging from 0.003 to 0.023 mg/kg in the tissues of *Corbicula fluminea* collected in a river affected by a decommissioned Antimony smelter.

It is worth noting the depletion relative to T0 of elements like Cu, Pb and Ni, (Fig. 3 and Table 2). This might suggest that these elements are not strongly mobilized in the lagoon environment, although slight differences can be recognized between S7 and the other sites. This is in contrast with results by Pignotti et al. (2018) that using sequential extraction, defined Cu and Pb as the most mobile elements in sediments (~70 % and ~80 %, of the total, respectively) in the neighbouring Pialassa Piomboni. In agreement with Pignotti et al. (2018) is the Ni behaviour here characterised by low mobility and predominantly bound to the residual fraction of the sediment, suggesting a lithogenic origin.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 3. Bioaccumulation factor of metals in mussel tissues. The analysis has been performed on the tissue of mussels deployed for 30 days into different sites of the Pialassa Baiona lagoon compared to mussels analysed before the *in situ* deployment (T0). Elements are ranked according to average enrichment.

3.4 Biological effects on mussels

The mussel 30-day exposure within the Pialassa Baiona lagoon resulted in significant alterations of biological endpoints measured at different levels of the hierarchical ladder (from cell to organism) (Fig. 4). Among the investigated sites, adverse conditions for the mussel survival were observed in S1, with about 80 % mortality *versus* 10 % recorded elsewhere (Fig. 4A). Different from S2 – S7, survival (and biomarker) data were measured in S1 after just three days of exposure, in line with previous observations on mussels transplanted in the same area (Franzellitti et al., 2010). The short time spent *in situ* within S1 was not such to allow mussels to bioaccumulate metals at levels considered environmentally realistic, for which ≥ 28 days are usually required (Viarengo et al., 2007a); thus, data of metal bioaccumulation are missing for this site. Sediment concentrations of Hg, Cr, Zn, Cu and Ba, known for their toxicity on marine mussels (De Los Ríos et al., 2013) were negatively and significantly correlated with mussel survival (Fig. 5). Relationships were also detected between survival and average salinity and temperature, which showed minima and maxima in S1, respectively (Table S3). Both parameters are known to be responsible for the mussel physiological fitness and to influence their response to toxicants synergistically (Fabbri and Dinelli, 2014;

Formatted: English (United Kingdom)

1
2
3
4
5
6
7
8 DeCourten and Brander, 2017). It is thus plausible that physico-chemical conditions of both natural
9 and anthropogenic nature simultaneously influenced the survival of mussels across the lagoon.
10

11 Data of *in situ* survival follow trends similar to those from the SOS test, which showed
12 significant reduction of survival functions in mussels from S1 (Fig. 4B). SOS is measured as the
13 survival of mussels in dry conditions, which is an event regularly experienced by intertidal organisms
14 due to tidal fluctuations. The SOS data display significant and negative correlation with a suite of
15 metals measured in both sediments and mussel tissues, including Ba, Bi, Cu, Sr and Zn, among others
16 (Fig. 5). These findings are in line with previous evidence obtained in *M. galloprovincialis*, which
17 showed significant SOS decrease after 3 to 7 days of exposure to trace levels of Cu and Zn-based
18 antifouling agents (Viarengo et al., 1995; Marcheselli et al., 2011). Further elements showing
19 marked enrichment in S1 sediments, as Cd (0.68 mg/kg) and Hg (25.7 µg/kg), were significantly
20 correlated with SOS (Fig. 4). SOS and survival data were found to be significantly and positively
21 correlated to each other and showed positive and negative correlation with electrical conductivity
22 and temperature, respectively (Table S3). This indicates that higher temperatures and salinity
23 measured at S1 throughout the exposure period may have cumulatively increased the impact of
24 metals on the mussel vital functions. Although it is acknowledged that SOS sensitivity is lower than
25 other sub-lethal and general stress biomarkers (Viarengo et al., 2007a), data from this study
26 highlight its suitability to detect causal relationships between metal contamination/bioavailability
27 and adverse effects on mussels at the “organism” level.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 4. Biomarker modulation in mussels transplanted into the Pialassa Baiona lagoon. Data from all sites (mean ± SEM) have been measured in mussels collected after 30 days of exposure, except for S1, referred to mussels collected after 3 days of exposure due to high mortality. A, survival (SURV); B, Stress on stress (SOS); C, lysosomal membrane stability (LMS); D, lysosome to cytoplasm volume ratio (LYS/CYT); E, lipofuscin (LIF), F, catalase activity (CAT); G, metallothionein content (MT); H, acetylcholinesterase activity (ACHE); I, peroxisomal acyl-CoA oxidase (AOX). Asterisks indicate statistically significant differences compared to the control site S7 (One-Way ANOVA, Duncan's post-hoc test).

LMS is the most sensitive biomarker of general stress in bivalves and previous studies highlighted strict relationships between the xenobiotic-driven decrease of this parameter in *Mytilus*

1
2
3
4
5
6
7
528 spp. and alterations to higher functions, such as growth and/or reproduction (Moore et al., 2006;
530 Martinez-Gomez et al., 2008). Data reported in Figure 4C indicate a significant LMS reduction in
531 digestive glands from mussels collected at S1 (3-day exposure), S3 and S6 (30-day exposure)
532 compared to control mussels from S7. Interestingly, none of the measured metals, physico-chemical
533 factors and/or biological endpoints was correlated with trends of LMS in exposed mussels (Fig. 5,
534 Table S3). Although positive relationships between (dissolved) metal exposure and LMS have been
535 reported in mussels under laboratory conditions (Viarengo et al., 1997), no correlation has been
536 highlighted following 28 days of *in situ* exposure into a nearby lagoon (i.e. Pialassa Piomboni,
537 Ravenna, Italy; Capolupo et al., 2017).
538

539 The mussel exposure to contaminated environment is frequently associated to an augmented
540 permeability of the lysosomal membrane and an overall lysosomal swelling. In the present study,
541 mussels deployed into S3, S5 and S6 showed a significantly higher LYS/CYT compared to the control
542 site S7 (Fig. 4D), following trends significantly correlated to tissue levels of Ba, Mn and Zr (Fig. 5).
543 The S3 and S5 were located in proximity of streams ensuring the flowing of industrial wastewater
544 from southern and eastern settlements to the Adriatic Sea, while S6 was located along a channel
545 that receive water from Lamone river after having crossed the natural areas located to the west (Fig.
546 1); thus, these data outline the need for a better understanding of the spatial/temporal relationships
547 between different source of lagoon water (industrial and continental), metal release into the
548 environment and lysosomal dysfunction in the exposed biota.
549

550 Lysosomal biomarker of general stress, as LMS and LYS/CYT, describe the overall health status
551 of tested organisms and may thus be modulated by a wide range of natural/anthropogenic stressors;
552 therefore, considering the diverse array of human activities directly and indirectly linked to the
553 Pialassa Baiona lagoon, it cannot be ruled out that further organic pollutants may have played a role
554 in observed lysosomal responses.
555
556
557
558
559
560
561
562
563
564
565

Formatted: Highlight

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
54
55
56
57
58
59
60
61
62
63
64
65




Figure 5. Correlation between biological data and metal concentrations. Pearson's coefficients calculated between biological data and levels of metals measured in mussel tissues (bioaccumulation) and sediment. Only coefficients showing a significant correlation ($p < 0.05$) are reported in the figure.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Mussels from all sites, except S2, showed a significant accumulation of LIF compared to those deployed into the reference site (S7) (Fig. 4E). LIF represent the ultimate products of the lipid peroxidation of biological membranes; they are composed by oxidatively modified proteins, lipid degradation by-products, carbohydrates and metals, and appear as granules within lysosomes of cells composing the digestive tubules epithelium (Moore et al., 2008). Previous biomonitoring studies highlighted clear relationships between LIF accumulation in mussels, the metal uptake and the metal spatial/temporal distribution into enclosed marine systems as bays and lagoons (Moschino et al., 2011; Capolupo et al., 2017). Accordingly, we identified significant and positive correlations between LIF, Al and Ti tissue content in mussels, as well as with As, Cd, Se and TI measured in sediments (Fig. 5). In addition, a significant and negative correlation has been observed between LIF and the activity of the antioxidant enzyme CAT (Table S3). The latter showed a significant decrease *versus* control (S7) in mussels exposed at S1 (Fig. 4F), suggesting that beside ROS formation, the exposure to relatively high metal concentrations might also down-regulate the overall mussel antioxidant defences. CAT also showed negative correlation with elements previously found to elicit pro-oxidant effects in mussels, as As, Cd, Cr, Sb (Franzellitti et al., 2012; Benali et al., 2017; Coppola et al., 2018), and with temperature trends within the lagoon. This indicates that both chemical and physical factors might have cumulatively influenced the mussel ROS scavenging activity over the exposure within the lagoon.

MT have been used as biomarker of exposure in this study owing to their high heavy-metal-binding capacity (Lavradas et al., 2016). Compared to control site S7, a significant MT synthesis induction has been recorded in the digestive gland of mussels from S3 and S1 (Fig. 4H), both located in the southern part of the lagoon in proximity of a large industrial complex (Fig.1). MT trends showed a positive and significant correlation with the sediment concentration of metals known as MT up-regulator, including Cd, Cu, Zn, Pd, Pt, Sn and Hg (Viarengo et al., 2001; Rocha et al., 2016; Zimmermann and Sures, 2018), and other less studied such as Ag, Au, Ba, Bi, Re, and Sb (Fig. 5). Positive relationships were detected MT trends and fluctuations of temperature and conductivity (Table S3), indicating that physico-chemical factors may interactively modulate MT in mussels from contaminated areas.

The activity of the enzyme AChE, which mediates the termination of the nervous impulses driven by the neurotransmitter acetylcholine, was significantly inhibited in mussels from the northern site S6 (Fig. 4H). AChE is typically inhibited by pesticides, although in mussels it may also be inhibited by high levels of metals (Lionetto et al., 2003). Apart from a negative relationship with

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Be, AChE trends did not show significant relationships with metals measured in sediments and mussel tissues (Fig. 5), nor with other biological endpoints measured in deployed mussels (Table S3). Compared to the S7, sediments collected in S6 showed a 20 times higher enrichment of Hg (Fig. 2), which is known as a potent AChE inhibitor in mussels, notably when co-exposed to fluctuations of other physico-chemical parameters, such as temperature (Coppola et al., 2017).

The possible impact of organic xenobiotics on the physiology of transplanted mussels has been screened through the analysis of the Acyl-Co A oxidase (AOX) activity. *In situ* deployed mussels showed a significant AOX increase following exposure at all sites except S2 (Fig. 4I). These data corroborate previous evidence of a diffuse PAH contamination (notably pyrene, fluoranthene and anthracene) in sediment and mussels collected from the same area (Vassura et al., 2005; Fabbri et al., 2006). Vassura et al. (2005) also found clear relationships between patterns of PAHs and Hg in the lagoon sediments. Accordingly, many metals herein measured in sediments, including among others Ag, Ba; Hg, Cu, Cr, Cd and Zn, as well as mussel tissues concentrations of B, Cu, Li, Mo and Sr showed positive correlation with AOX (Fig. 5), suggesting a possible metal-induced peroxisome proliferation and a similar spatial distribution of metals and POPs within the lagoon.

The output produced by biomarkers data integration performed through the MES is reported in Figure 6. The LMS was chosen as guide parameter for the data integration. A good health status (HSI = A) was assigned by the Expert System to mussels *in situ* exposed at the reference site S7 and at S2. This output confirms the good environmental quality of the lagoon north-eastern area, as previously identified based on chemical and biological evidence (Franzellitti et al., 2010), and is in line with the lack of biomarker alterations observed in mussels from S2. Although this site is located in the southern part of the lagoon, i.e. in proximity of industrial settlements, it is directly exposed to the marine water entering the lagoon indicating the relevance of high water circulation and oxygenation to the fitness of the resilient biota. A moderate stress level (HSI = C) was assigned to mussels exposed at S5, while mussels from S1, S3 and S6 were classified as heavily stressed (HSI = D) (Fig. 6). In line with the metal contamination/bioaccumulation and coherently with MT levels measured in mussels, ~~the MES associated the health status alteration to a significant excess of metal contamination~~ was identified in S1 and S3. Moreover, the MES associated the health status alteration of mussels exposed at S1, S3, S5 and S6 to the presence of organic xenobiotics/persistent organic compounds (POPs) has been identified in S1, S3, S5 and S6, in line with measured AOX trends. Although organics were not included in the analytical design of this study, this output is consistent with previous and evidence of PCB and PAH contamination across the

Formatted: Highlight

Formatted: Highlight

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

lagoon reported in previous studies (Guerra, 2012; Vassura et al., 2005; Fabbri et al., 2006; Vassura et al., 2005) and may confirm the co-occurrence of multiple stressors in different areas of the Pialassa Baiona lagoon:-




Figure 6. Biomarker data integration. A unique Health status index (HSI) has been generated using the Mussel Expert System (MES, developed by Dagnino et al., 2007) based on biomarker alterations observed at each site. The LMS was chosen as guide parameter for the HSI assessment.

4 Conclusions

The integrated approach implemented in this study provided accurate and reliable clues on the distribution of metals in the Pialassa Baiona lagoon, their bioavailability for exposed mussels and their effects on key biological functions, posing also attention to seldom considered elements such as Ag and Sb.

The physico-chemical data revealed clear differences among S1 and other investigated sites. The water in Magni Channel (S1) showed higher column temperature, reduced EC and a lower pH

1
2
3
4
5
6
7
640 value, for the entire monitoring period generating local conditions aligned with the worst forecasted
641 climate change scenarios.

642 The geochemical analysis of total elements indicated that sediments in Pialassa Baiona are a
643 mixture of carbonate and silicates with variable amount of organic matter. The total concentrations
644 of the investigated elements are within the range of regional background. Exception is made for Zn
645 and Cu, and Pb respectively double (in S1) and slightly above (S3 and S5) the local natural
646 background values.

647 The aqua regia digestion identified a south-north decreasing contamination trend, with
648 highest enrichments of Hg, Pt, Au, Ag, Mo, Re, Cd, Pd, Zn; Sn, and Cu (in enrichment order, Fig. 2)
649 being at least two folds enriched than S7 (internal control). Moreover, the most affected sites are
650 in the south-western areas, which consistently receives wastewater effluents from a near
651 petrochemical settlement (south) and drainage water from the inland (west).

652 Bioaccumulation analysis performed in transplanted mussels proved suitable to evaluate the
653 bioavailability of metals in relation to their distribution across the lagoon. Integrating physical,
654 chemical and biological measures allowed for defining causal relationships between observed
655 mortality rates, lysosomal dysfunctions, oxidative stress induction, biomarker of exposure
656 modulations and the metal distribution within the lagoon. Besides this, the different channels
657 examined within the lagoon proved to be an excellent study system for evaluating climate change
658 effects (i.e. modified temperature, pH, and salinity) combined with pollution. In addition, the use of
659 the MES proved suitable to identify the impact induced by specific substances on the overall mussel
660 fitness and to estimate the toxicological pathways occurring in highly pressure lagoons.

661 Overall, data from this study assign a general scarce environmental quality to the Pialassa
662 Baiona lagoon; considering the heterogeneity of this area in terms of geological, physico-chemical
663 and anthropogenic factors, regular biomonitoring surveys should be performed to control the onset
664 of conditions which might exacerbate the impact of metals and other pollutants, notably on a
665 seasonal timeframe. In this respect, the integrated strategy undertaken in this investigation
666 represents a promising approach for the biomonitoring of this area and, more generally, for reliably
667 assessing the environmental quality of highly pressured lagoons.

668
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
668
670
671
672
673
674
15
16
674
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgement

The Authors are grateful to Municipality of Ravenna for supporting the work and allowing field activity within the Piallassa Baiona lagoon. Furthermore, the Authors thank the anonymous Reviewers who helped in improving the quality of the article.

5 References

- Abdou, M., Schäfer, J., Cobelo-García, A., Neira, P., Petit, J.C.J., Auger, D., Chiffolleau, J.-F., Blanc, G., 2016. Past and present platinum contamination of a major European fluvial-estuarine system: Insights from river sediments and estuarine oysters. *Mar. Chem.* 185, 104-110. <https://doi.org/10.1016/j.marchem.2016.01.006>
- Aliaume, C., Do Chi, T., Viaroli, P., Zaldivar, J.M., 2007. Coastal lagoons of southern Europe: recent changes and future scenarios. *Transitional Waters Monogr.* 1, 1-12, 10.1285/i18252273v1n1p1
- Amorosi, A., Colalongo, M.L., Pasini, G., Preti, D., 1999. Sedimentary response to Late Quaternary sea-level changes in the Romagna coastal plain (northern Italy). *Sedimentology* 46, 99-121.
- Amorosi, A., Centineo, M.C., Dinelli, E., Lucchini, F., Tateo, F., 2002. Geochemical and mineralogical variations as indicators of provenance changes in Late Quaternary deposits of SE Po Plain. *Sediment. Geol.* 151, 273-292. [https://doi.org/10.1016/S0037-0738\(01\)00261-5](https://doi.org/10.1016/S0037-0738(01)00261-5).
- Benali, I., Boutiba, Z., Grandjean, D., De Alencastro, L.F., Rouane-Hacene, O., Chèvre, N., 2017. Spatial distribution and biological effects of trace metals (Cu, Zn, Pb, Cd) and organic micropollutants (PCBs, PAHs) in mussels *Mytilus galloprovincialis* along the Algerian west coast. *Mar. Pollut. Bull.* 115(1-2), 539-550. <https://doi.org/10.1016/j.marpolbul.2016.12.028>
- Bird, E.C.F., 1994. Physical setting and geomorphology of coastal lagoons. In: Kjerfve, B. (Ed.), *Coastal Lagoon Processes*, Elsevier Oceanography Series, vol. 60. Elsevier, Amsterdam, pp. 9-30. [https://doi.org/10.1016/S0422-9894\(08\)70007-2](https://doi.org/10.1016/S0422-9894(08)70007-2)
- Borghesi, F., Migani, F., Dinelli, E., 2016. Geochemical characterization of surface sediments from the northern Adriatic wetlands around the Po River delta. Part II: aqua regia results. *J. Geochem. Explor.* 169, 13-29. <https://doi.org/10.1016/j.gexplo.2016.06.012>
- Bossi, T., Gediga, J., 2017. The environmental profile of Platinum Group Metals. *Johnson Matthey Technol. Rev.* 61, 111-121, <https://doi.org/10.1595/205651317x694713>
- Braga, A.C., Camacho, C., Marques, A., Gago-Martínez, A., Pacheco, M., Costa, P.R., 2018. Combined effects of warming and acidification on accumulation and elimination dynamics of paralytic shellfish toxins in mussels *Mytilus galloprovincialis*. *Environ. Res.* 164, 647-654. <https://doi.org/10.1016/j.envres.2018.03.045>.
- Buryskova, B., Hilscherova, K., Babica, P., Vrskova, D., Marsalek, B., Blaha, L., 2006. Toxicity of complex cyanobacterial samples and their fractions in *Xenopus laevis* embryos and the role of microcystins. *Aquat. Toxicol.* 80, 346-354. <https://doi.org/10.1016/j.aquatox.2006.10.001>
- Capolupo, M., Valbonesi, P., Kiwan, A., Buratti, S., Franzellitti, S., Fabbri, E., 2016. Use of an integrated biomarker-based strategy to evaluate physiological stress responses induced by

Formatted: English (United Kingdom)

Formatted: Spanish (Spain)

1
2
3
4
5
6
7
8 environmental concentrations of caffeine in the Mediterranean mussel *Mytilus*
9 *galloprovincialis*. *Sci. Total Environ.* 563-564, 538-548. doi:10.1016/j.scitotenv.2016.04.125

10 Capolupo, M., Franzellitti, S., Kiwan, A., Valbonesi, P., Dinelli, E., Pignotti, E., Birke, M., Fabbri, E.,
11 2017. A comprehensive evaluation of the environmental quality of a coastal lagoon (Ravenna,
12 Italy): Integrating chemical and physiological analyses in mussels as a biomonitoring strategy.
13 *Sci. Total Environ.* 598, 146-159. <https://doi.org/10.1016/j.scitotenv.2017.04.119>

14 Carvalho, R.N., Arukwe, A., Ait-Aissa, S., Bado-Nilles, A., Balzamo, S., Baun, A., Belkin, S., Blaha, L.,
15 Brion, F., Conti, D., Creusot, N., Essig, Y., Ferrero, V.E.V., Flander-Putrlé, V.V., Fürhacker, M.,
16 Grillari-Voglauer, R., Hogstrand, C., Jonáš, A., Kharlyngdoh, J.B., Loos, R., Lundebye, A.K., Modig,
17 C., Olsson, P.E., Pillai, S., Polak, N., Potalivo, M., Sanchez, W., Schifferli, A., Schirmer, K., Sforzini,
18 S., Stürzenbaum, S.R., Sjøfteland, L., Turk, V., Viarengo, A., Werner, I., Yagur-Kroll, S.S.,
19 Zounkova, R., Lettieri, T., 2014. Mixtures of chemical pollutants at European legislation safety
20 concentrations: how safe are they? *Toxicol. Sci.* 141, 218-233.
21 <http://dx.doi.org/10.1093/toxsci/kfu118>.

22 Cicchella, D., Giaccio, L., Dinelli, E., Albanese, S., Lima, A., Zuzolo, D., Valera, P., De Vivo, B., 2015.
23 GEMAS: Spatial distribution of chemical elements in agricultural and grazing land soil of Italy. *J.*
24 *Geochem. Explor.* 154, 129-142. <https://doi.org/10.1016/j.gexplo.2014.11.009>

25 Cobelo-García, A., Neira, P., Mil-Homens, M., Caetano, M., 2011. Evaluation of the contamination
26 of platinum in estuarine and coastal sediments (Tagus Estuary and Prodelta, Portugal). *Mar.*
27 *Pollut. Bull.* 62 (3), 646-650. <https://doi.org/10.1016/j.marpolbul.2010.12.018>

28 Coppola, F., Almeida, Â., Henriques, B., Soares, A.M., Figueira, E., Pereira, E., Freitas, R., 2017.
29 Biochemical impacts of Hg in *Mytilus galloprovincialis* under present and predicted warming
30 scenarios. *Sci. Total Environ.* 601, 1129-1138. <https://doi.org/10.1016/j.scitotenv.2017.05.201>

31 Coppola, F., Almeida, Â., Henriques, B., Soares, A.M.V.M., 2018. Ecotoxicology and Environmental
32 Safety Biochemical responses and accumulation patterns of *Mytilus galloprovincialis* exposed
33 to thermal stress and Arsenic contamination. *Ecotox. Environ. Saf.* 147, 954-962. <https://doi.org/10.1016/j.ecoenv.2017.09.051>

34 Covelli, S., Emili, A., Acquavita, A., Koron, N., Faganeli, J., 2011. Benthic biogeochemical cycling of
35 mercury in two contaminated northern Adriatic coastal lagoons. *Cont. Shelf Res.* 31(16), 1777-
36 1789. <https://doi.org/10.1016/j.csr.2011.08.005>.

37 Curzi, P.V., Dinelli, E., Ricci Lucchi, M., Vaiani, S.C., 2006. Palaeoenvironmental control on sediment
38 composition and provenance in the late Quaternary deltaic successions: A case study from the
39 Po delta area (Northern Italy). *Geol. J.* 41 (5), 591-612. <https://doi.org/10.1002/gj.1060>

40 Dagnino, A., Allen, J.I., Moore, M.N., Broeg, K., Canesi, L., Viarengo, A., 2007. Development of an
41 expert system for the integration of biomarker responses in mussels into an animal health
42 index. *Biomarkers* 12, 155-172. <http://dx.doi.org/10.1080/13547500601037171>.

43 Dahms, H.U., 2014. The grand challenges in marine pollution research. *Front. Mar. Sci.* 1, 1-5.
44 <http://dx.doi.org/10.3389/fmars.2014.00009>

45 De Gregori, I., Quiroz, W., Pinochet, H., Pannier, F., Potin-Gautier, M., 2007. Speciation analysis of
46 antimony in marine biota by HPLC-(UV)-HG-AFS: Extraction procedures and stability of
47 antimony species. *Talanta* 73(3), 458-465. <https://doi.org/10.1016/j.talanta.2007.04.015>

48 De Los Ríos, A., Pérez, L., Ortiz-zarragoitia, M., Serrano, T., Barbero, M.C., Echavarri-erasun, B., 2013.
49 Assessing the effects of treated and untreated urban discharges to estuarine and coastal waters
50

1
2
3
4
5
6
7
8 applying selected biomarkers on caged mussels. Mar. Pollut. Bull. 77, 251-265.
9 <https://doi.org/10.1016/j.marpolbul.2013.09.050>

10 Decourten, B.M., Brander, S.M., 2017. Combined effects of increased temperature and endocrine
11 disrupting pollutants on sex determination, survival, and development across generations. Sci.
12 Rep. 7, 1-9. <https://doi.org/10.1038/s41598-017-09631-1>

13 Dinelli, E., Tateo F., Summa V. 2007. Geochemical and mineralogical proxies for grain size in
14 mudstones and siltstones from the Pleistocene and Holocene of the Po River alluvial plain, Italy
15 J. Arribas, S. Critelli, M.J. Johnsson (Eds.), Sedimentary Provenance and Petrogenesis:
16 Perspectives from Petrography and Geochemistry, Geological Society of America Special Paper,
17 vol. 420, pp. 25-36

18 Dinelli, E., Ghosh, A., Rossi, V., Vaiani, S.C., 2012. Multiproxy reconstruction of Late Pleistocene-
19 Holocene environmental changes in coastal successions: microfossil and geochemical evidences
20 from the Po Plain (Northern Italy). Stratigraphy 9(2), 153-167.

21 Dominik, J., Tagliapietra, D., Bravo, A.G., Sigovini, M., Spangenberg, J.E., Amouroux, D., Zonta, R.
22 2014. Mercury in the food chain of the Lagoon of Venice, Italy. Mar. Pollut. Bull. 88(1), 194-206.
23 <https://doi.org/10.1016/j.marpolbul.2014.09.005>

24 Donnini, F., Dinelli, E., Sangiorgi, F., Fabbri, E., 2007. A biological and geochemical integrated
25 approach to assess the environmental quality of a coastal lagoon (Ravenna, Italy). Environ. Int.
26 33 (7), 919-928. <https://doi.org/10.1016/j.envint.2007.05.002>

27 Duroudier, N., Cardoso, C., Mehennaoui, K., Mikolaczyk, M., Schäfer, J., Gutleb, A.C., Giamberini, L.,
28 Bebianno, M.J., Bilbao, E., Cajaraville, M.P., 2019. Changes in protein expression in mussels
29 *Mytilus galloprovincialis* dietarily exposed to PVP/PEI coated silver nanoparticles at different
30 seasons. Aquat. Toxicol. 210, 56-68. <https://doi.org/10.1016/j.aquatox.2019.02.010>

31 European Commission Directive 2000/60/EC of the European Parliament and of the Council of 23
32 October 2000 establishing a framework for community action in the field of water policy (Water
33 Framework Directive). Off. J. Eur. Community, 2000.

34 European Commission Directive 2008/56/EC of the European Parliament and of the Council of 17
35 June 2008 Establishing a Framework for Community Action in the Field of Marine Environmental
36 Policy (Marine Strategy Framework Directive) (OJ L 164/19. 22 pp. 25.06).

37 Ellman, G.L., Courtney, K.D., Andres Jr., V., Feather-Stone, R.M., 1961. A new and rapid colorimetric
38 determination of acetylcholinesterase activity. Biochem. Pharmacol. 7, 88-95.
39 [https://doi.org/10.1016/0006-2952\(61\)90145-9](https://doi.org/10.1016/0006-2952(61)90145-9)

40 Fabbri, D., Felisatti, O., Lombardo, M., Trombini, C., Vassura, I., 1998. The Lagoon of Ravenna Italy:
41 characterisation of mercury-contaminated sediments. Sci. Total Environ. 213, 121-128.
42 [https://doi.org/10.1016/S0048-9697\(98\)00083-7](https://doi.org/10.1016/S0048-9697(98)00083-7)

43 Fabbri, D., Tartari, D., Trombini, C., 2000. Analysis of poly (vinyl chloride) and other polymers in
44 sediments and suspended matter of a coastal lagoon by pyrolysis-gas chromatography-mass
45 spectrometry. Anal. Chim. Acta 413(1-2), 3-11. [https://doi.org/10.1016/S0003-2670\(00\)00766-](https://doi.org/10.1016/S0003-2670(00)00766-2)
46 2

47 Fabbri, D., Gemelli, S., Langone, L., Miserocchi, S., Trombini, C., Vassura, I., 2001a. Anthropogenically
48 derived mercury in sediments of Pialassa Baiona, Ravenna, Italy. Annali di Chimica 91(9-10),
49 563-575.

Formatted: English (United Kingdom)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

1
2
3
4
5
6
7
792 Fabbri, D., Locatelli, C., Snape, C.E., Tarabusi, S., 2001b. Sulfur speciation in mercury-contaminated
793 sediments of a coastal lagoon: The role of elemental sulfur. *J. Environ. Monitor.* 5, 483-486.
794 <https://doi.org/10.1039/b104477j>

795 Fabbri, D., Baravelli, V., Giannotti, K., Donnini, F., Fabbri, E., 2006. Bioaccumulation of
796 cyclopenta[cd]pyrene and benzo[ghi]fluoranthene by mussels transplanted in a coastal lagoon.
797 *Chemosphere* 64, 1083-1092. <https://doi.org/10.1016/j.chemosphere.2005.11.071>

798 Fabbri, E., Dinelli, E., 2014. Physiological responses of marine animals towards adaptation to climate
799 changes. *The Mediterranean Sea: Its History and Present Challenges*, pp. 401-417.
800 https://doi.org/10.1007/978-94-007-6704-1_23

801 Fattorini, D., Notti, A., Di Mento, R., Cicero, A.M., Gabellini, M., Russo, A., Regoli, F., 2008. Seasonal,
802 spatial and inter-annual variations of trace metals in mussels from the Adriatic Sea: a regional
803 gradient for arsenic and implications for monitoring the impact of off-shore activities.
804 *Chemosphere* 72(10), 1524-1533. <https://doi.org/10.1016/j.chemosphere.2008.04.071>.

805 Filella, M., Belzile, N., Chen, Y.W., 2002. Antimony in the environment: a review focused on natural
806 waters: I. Occurrence. *Earth-Sci. Rev.* 57(1-2), 125-176. [https://doi.org/10.1016/S0012-8252\(01\)00070-8](https://doi.org/10.1016/S0012-8252(01)00070-8)

807 Filella, M., Hennebert, P., Okkenhaug, G., Turner, A., 2019. Occurrence and fate of antimony in
808 plastics. *J. Hazard. Mater.* 121764. <https://doi.org/10.1016/j.jhazmat.2019.121764>

809 Franzellitti, S., Buratti, S., Donnini, F., Fabbri, E., 2010. Exposure of mussels to a polluted
810 environment: Insights into the stress syndrome development. *Comp. Biochem. Physiol. - C
811 Toxicol. Pharmacol.* 152, 24-33. <https://doi.org/10.1016/j.cbpc.2010.02.010>

812 Franzellitti, S., Viarengo, A., Dinelli, E., Fabbri, E., 2012. Molecular and cellular effects induced by
813 hexavalent chromium in Mediterranean mussels. *Aquat. Toxicol.* 124-125, 125-132.
814 <https://doi.org/10.1016/j.aquatox.2012.07.011>

815 Franzini, M., Leoni, L., Saitta, M., 1972. A simple method to evaluate the matrix effects in X-ray
816 fluorescence analysis. *X-Ray Spectrom.* 1, 151-154. <https://doi.org/10.1002/xrs.1300010406>

817 Franzini, M., Leoni, L., Saitta, M., 1975. Revisione di una metodologia analitica per fluorescenza-X
818 basata sulla correzione completa degli effetti di matrice. *Rend. Soc. Ital. Mineral. Petrol.* 31,
819 365-378. (in italian)

820 Garcia, N., 2015. The Asiatic Clam (*Corbicula fluminea*) as a bio-monitor for determining the
821 distribution of antimony, arsenic and thallium in the water column and sediments of Manadas
822 Creek, Laredo, Texas. M.Sc thesis, Texas A&M International University, pp. 59. <https://tamiu-ir.tdl.org/handle/2152.4/58>. Accessed July 2020

823 Giusti, L., Williamson, A.C., Mistry, A., 1999. Biologically available trace metals in *Mytilus edulis* from
824 the coast of northeast England. *Environ. Int.* 25(8), 969-981. [https://doi.org/10.1016/S0160-4120\(99\)00066-5](https://doi.org/10.1016/S0160-4120(99)00066-5)

825 Giusti, L., Zhang, H., 2002. Heavy metals and arsenic in sediments, mussels and marine water from
826 Murano (Venice, Italy). *Environ. Geochem. Health* 24(1), 47-65.
827 <https://doi.org/10.1023/A:1013945117549>

828 Govindaraju, K., 1984. 1984 compilation of working values and sample description for 170
829 international reference samples of mainly silicate rocks and minerals. *Geostandard. Newslett.*
830 8, 3-16. <https://doi.org/10.1111/j.1751-908X.1984.tb01069.x>

1
2
3
4
5
6
7
834 Greggio, N., Giambastiani, B.M.S., Campo, B., Dinelli, E., Amorosi, A., 2018a. Sediment composition,
835 provenance, and Holocene paleoenvironmental evolution of the Southern Po River coastal plain
836 (Italy). *Geol. J.* 53(3), 914-928. <https://doi.org/10.1002/gj.2934>
837
838 Greggio, N., Giambastiani, B.M.S., Antonellini, M., 2018b. Infiltration/irrigation trench for
839 sustainable coastal drainage management: Emilia-Romagna (Italy). *Environ. Eng. Manag. J.*
840 17(10). <https://doi.org/10.30638/eemj.2018.236>
841
842 Greggio, N., Giambastiani, B., Mollema, P., Laghi, M., Capo, D., Gabbianelli, G., Antonellini, M.,
843 Dinelli, E., 2020. Assessment of the Main Geochemical Processes Affecting Surface Water and
844 Groundwater in a Low-Lying Coastal Area: Implications for Water Management. *Water* 12(6),
845 1720. <https://doi.org/10.3390/w12061720>
846
847 Guerra, R., Pasteris, A., Ponti, M., Fabbri, D., Bruzzi, L., 2007. Impact of dredging in a shallow coastal
848 lagoon: Microtox® Basic Solid-Phase Test, trace metals and Corophium bioassay. *Environ. Int.*
849 33 (4), 469-473. <https://doi.org/10.1016/j.envint.2006.10.006>
850
851 Guerra R., 2012. Polycyclic Aromatic Hydrocarbons, Polychlorinated Biphenyls and Trace Metals in
852 Sediments from a Coastal Lagoon (Northern Adriatic, Italy). *Water Air Soil Poll.* 223, 85-98.
853 <https://doi.org/10.1007/s11270-011-0841-6>
854
855 Guerra R., Pasteris A., Lee Seok-hyung, Park No-jin, Ok Gon, 2014. Spatial patterns of metals,
856 PCDDs/Fs, PCBs, PBDEs and chemical status of sediments from a coastal lagoon (Pialassa Baiona,
857 NW Adriatic, Italy). *Mar. Pollut. Bull.* 89, 407-416.
858 <https://doi.org/10.1016/j.marpolbul.2014.10.024>
859
860 James, E., Turner, A., 2020. Mobilisation of antimony from microplastics added to coastal sediment.
861 *Environ. Pollut.* 114696. <https://doi.org/10.1016/j.envpol.2020.114696>
862
863 Jimeno-Romero, A., Bilbao, E., Izagirre, U., Cajaraville, M.P., Marigómez, I., Soto, M., 2017. Digestive
864 cell lysosomes as main targets for Ag accumulation and toxicity in marine mussels, *Mytilus*
865 *galloprovincialis*, exposed to maltose-stabilised Ag nanoparticles of different sizes.
866 *Nanotoxicology* 11(2), 168-183. <https://doi.org/10.1080/17435390.2017.1279358>
867
868 Lari, E., Gauthier, P., Mohaddes, E., Pyle, G.G., 2017. Interactive toxicity of Ni, Zn, Cu, and Cd on
869 *Daphnia magna* at lethal and sub-lethal concentrations. *J. Hazard. Mater.* 334, 21-28.
870 <http://dx.doi.org/10.1016/j.jhazmat.2017.03.060>
871
872 Lavradas, R.T., Rocha, R.C.C., Bordon, I.C.A.C., Saint’Pierre, T.D., Godoy, J.M., Hauser-Davis, R.A.,
873 2016. Differential metallothionein, reduced glutathione and metal levels in *Perna perna* mussels
874 in two environmentally impacted tropical bays in southeastern Brazil. *Ecotoxicol. Environ. Saf.*
875 129, 75-84. <http://dx.doi.org/10.1016/j.ecoenv.2016.03.011>
876
877 León, V.M., Moreno-González, R., García, V., Campillo, J.A., 2017. Impact of flash flood events on
878 the distribution of organic pollutants in surface sediments from a Mediterranean coastal lagoon
879 (Mar Menor, SE Spain). *Environ. Sci. Pollut. Res.* 24(5), 4284-4300.
880 <https://doi.org/10.1007/s11356-015-4628-y>
881
882 Leoni, L., Saitta, M., 1976. X-ray fluorescence analysis of 29 trace elements in rock and mineral
883 standard. *Rend. Soc. Ital. Mineral.* 72, 175-192.
884
885 Leoni, L., Menichini, M., Saitta, M., 1986. Determination of S, Cl, and F in silicate rocks by X-ray
886 fluorescence analyses. *X-Ray Spectrom.* 11, 156-158. <https://doi.org/10.1002/xrs.1300110404>

Formatted: English (United Kingdom)

1
2
3
4
5
6
7
875 Leopold, K., Denzel, A., Gruber, A., Malle, L., 2018. Mobility of traffic-related Pd and Pt species in
876 soils evaluated by sequential extraction. Environ. Pollut. 242, 1119-1127.
877 <https://doi.org/10.1016/j.envpol.2018.07.130>

878
879 Lionetto, M.G., Caricato, R., Giordano, M.E., Pascariello, M.F., Marinosci, L., Schettino, T., 2003.
880 Integrated use of biomarkers (acetylcholinesterase and antioxidant enzymes activities) in
881 *Mytilus galloprovincialis* and *Mullus barbatus* in an Italian coastal marine area. Mar. Pollut. Bull.
882 46(3), 324-330. [https://doi.org/10.1016/S0025-326X\(02\)00403-4](https://doi.org/10.1016/S0025-326X(02)00403-4)

883
884
885 Lottermoser, B.G., Ashley, P.M., Lawie, D.C., 1999. Environmental geochemistry of the Gulf Creek
886 copper mine area, north-eastern New South Wales, Australia. Environ. Geol. 39(1), 61-74.
887 <https://doi.org/10.1007/s002540050437>

888
889 Lowry, O.H., Rosenbrough, N.J., Farr, A.L., Randall, R.J., 1951. Protein measurement with the Folin
890 phenol reagent. J. Biol. Chem. 193, 265-275.

891
892
893 Mansour, C., Guibbolini, M., Rouane, O., Dalila, H., Mosbahi, S., De Faverney, C.R., 2020. Oxidative
894 stress and damage biomarkers in clam ruditapes decussatus exposed to a polluted site: the
895 reliable biomonitoring tools in hot and cold seasons. Arch. Environ. Contam. Toxicol. 78, 478-
896 494. <https://doi.org/10.1007/s00244-020-00713-2>

897
898
899 Marcheselli, M., Azzoni, P., Mauri, M., 2011. Novel antifouling agent-zinc pyrithione : Stress
900 induction and genotoxicity to the marine mussel *Mytilus galloprovincialis*. Aquat. Toxicol. 102,
901 39-47. <https://doi.org/10.1016/j.aquatox.2010.12.015>

902
903
904 Martin, M., Stephenson, M.D., Smith, D.R., Gutierrez-Galindo, E.A., Munoz, G.F., 1988. Use of silver
905 in mussels as a tracer of domestic wastewater discharge, Mar. Pollut. Bull. 19, 512-520
906 [https://doi.org/10.1016/0025-326X\(88\)90541-3](https://doi.org/10.1016/0025-326X(88)90541-3)

907
908
909 Martin-Diaz, L., Franzellitti, S., Buratti, S., Valbonesi, P., Capuzzo, A., Fabbri, E., 2009. Effects of
910 environmental concentrations of the antiepileptic drug carbamazepine on biomarkers and
911 cAMP-mediated cell signaling in the mussel *Mytilus galloprovincialis*. Aquat. Toxicol. 94, 177-
912 185. <https://doi.org/10.1016/j.aquatox.2009.06.015>

913
914
915 Martinez-Gomez, C., Benedicto, J., Campillo, J.A., Moore, M., 2008. Application and evaluation of
916 the neutral red retention (NRR) assay for lysosomal stability in mussel populations along the
917 Iberian Mediterranean coast. J. Environ. Monit. 10, 490-499.
918 <https://doi.org/10.1039/C2EM90056D>

919
920
921 Mattes, T.E., Ewald, J.M., Liang, Y., Martinez, A., Awad, A., Richards, P., Hornbuckle, K.C., Schnoor,
922 J.L., 2018. PCB dechlorination hotspots and reductive dehalogenase genes in sediments from a
923 contaminated wastewater lagoon. Environ. Sci. Pollut. Res., 25(17), 16376-16388.
924 <https://doi.org/10.1007/s11356-017-9872-x>

925
926
927 Matteucci, G., Rossini, P., Guerzoni, S., Arcangeli, A., Fonti, P., Langone, L., Misericchi, S., 2005.
928 Recent evolution of sedimentary heavy metals in a coastal lagoon contaminated by industrial
929 wastewaters (Pialassa Baiona, Ravenna, Italy). Hydrobiologia 550, 167-173.
930 <https://doi.org/10.1007/s10750-005-4374-0>

931
932
933 McGillicuddy, E., Murray, I., Kavanagh, S., Morrison, L., Fogarty, A., Cormican, M., Dockery, P.,
934 Prendergast, M., Rowan, N., Morris, D., 2017. Silver nanoparticles in the environment: sources,
935 detection and ecotoxicology. Sci. Total. Environ. 575, 231-246.
936 <https://doi.org/10.1016/j.scitotenv.2016.10.041>

1
2
3
4
5
6
7
917 Migani, F., Borghesi, F., Dinelli, E., 2015 Geochemical characterization of surface sediments from the
918 northern Adriatic wetlands around the Po river delta. Part I: Bulk composition and relation to
919 local background. *J. Geochem. Explor.* 156, 72-88. DOI: 10.1016/j.gexplo.2015.05.003
920
921 Miserocchi, S., Langone, L., Guerzoni, S., 1993. The fate of Hg contaminated sediments of the
922 Ravenna Lagoon (Italy): Final burial or potential remobilization. *Water Sci. Technol.* 28 (8-9),
923 349-358. <https://doi.org/10.2166/wst.1993.0633>
924
925 Mollema, P.N., Antonellini, M., Dinelli, E., Gabbianelli, G., Greggio, N., Stuyfzand, P.J., 2013.
926 Hydrochemical and physical processes influencing salinization and freshening in Mediterranean
927 low-lying coastal environments. *Appl. Geochem.* 34, 207-221.
928 <https://doi.org/10.1016/j.apgeochem.2013.03.017>
929
930 Moore, M.N., Allen, J.I., McVeigh, A., 2006. Environmental prognostics: an integrated model
931 supporting lysosomal stress responses as predictive biomarkers of animal health status. *Marine*
932 *Environ. Res.* 61(3), 278-304. <https://doi.org/10.1016/j.marenvres.2005.10.005>
933
934 Moore, M.N., 2008. Autophagy as a second level protective process in conferring resistance to
935 environmentally-induced oxidative stress. *Autophagy* 4, 254-256.
936 <https://doi.org/10.4161/auto.5528>
937
938 Moreno-González, R., Rodriguez-Mozaz, S., Gros, M., Barceló, D., León, V.M., 2015. Seasonal
939 distribution of pharmaceuticals in marine water and sediment from a mediterranean coastal
940 lagoon (SE Spain). *Environ. Res.* 138, 326-344. <https://doi.org/10.1016/j.envres.2015.02.016>
941
942 Moschino, V., Delaney, E., Meneghetti, F., Da, L., 2011. Biomonitoring approach with mussel *Mytilus*
943 *galloprovincialis* (Lmk) and clam *Ruditapes philippinarum* (Adams and Reeve, 1850) in the
944 Lagoon of Venice. *Environ. Monit. Assess.* 177, 649-663. <https://doi.org/10.1007/s10661-010-1663-5>
945
946 Mousavi, M.P., Gunsolus, I.L., De Jesús, C.E.P., Lancaster, M., Hussein, K., Haynes, C.L., Bühlmann,
947 P., 2015. Dynamic silver speciation as studied with fluoruous-phase ion-selective electrodes:
948 Effect of natural organic matter on the toxicity and speciation of silver. *Sci. Total Environ.* 537,
949 453-461. <https://doi.org/10.1016/j.scitotenv.2015.07.151>
950
951 Nazneen, S., Singh, S., Raju, N.J., 2019. Heavy metal fractionation in core sediments and potential
952 biological risk assessment from Chilika lagoon, Odisha state, India. *Quatern. Int.* 507, 370-388.
953 <https://doi.org/10.1016/j.quaint.2018.05.011>
954
955 Newton, A., Icely, J., Cristina, S., Brito, A., Cardoso, A.C., Colijn, F., Riva, S.D., Gertz, F., Hansen, J.W.,
956 Holmer, M., Ivanova, K., Leppäkoski, E., Canu, D.M., Mocenni, C., Mudge, S., Murray, N., Pejrup,
957 M., Razinkovas, A., Reizopoulou, S., Pérez-Ruzafa, A., Schernewski, G., Schubert, H., Carr, L.,
958 Solidoro, C., Viaroli, P., Zaldívar, J.-M., K., 2014. An overview of ecological status, vulnerability
959 and future perspectives of European large shallow, semi-enclosed coastal systems, lagoons and
960 transitional waters. *Estuar. Coast. Shelf Sci.* 140, 95-122.
961 <https://doi.org/10.1016/j.ecss.2013.05.023>
962
963 Orbea, A., Garmendia, L., Marigómez, I., Cajaraville, M.P., 2006. Effects of the "Prestige" oil spill on
964 cellular biomarkers in intertidal mussels: Results of the first year of studies. *Mar. Ecol. Prog. Ser.*
965 306, 177-189. <https://doi.org/10.3354/meps306177>
966
967 Outa, J.O., Kowenje, C.O., Plessl, C., Jirsa, F., 2020. Distribution of arsenic, silver, cadmium, lead and
968 other trace elements in water, sediment and macrophytes in the Kenyan part of Lake Victoria:
969

Formatted: Spanish (Spain)

Formatted: Spanish (Spain)

1
2
3
4
5
6
7
958 spatial, temporal and bioindicative aspects. Environ. Sci. Pollut. Res. 27(2), 1485-1498.
960 <https://doi.org/10.1007/s11356-019-06525-9>

961 Ozkan, D., Dagdeviren, M., Katalay, S., Guner, A., Yavaşoğlu, N.Ü.K., 2017. Multi-biomarker
962 responses after exposure to pollution in the mediterranean mussels (*Mytilus galloprovincialis*
963 L.) in the Aegean Coast of Turkey. B. Environ. Contam. Tox. 98(1), 46-52.
964 <https://doi.org/10.1007/s00128-016-1988-z>

965 Petry, A.C., Guimarães, T.F., Vasconcellos, F.M., Hartz, S.M., Becker, F.G., Rosa, R.S., Goyenola, G.,
966 Caramaschi, E.P., Díaz de Astarloa, J.M., Sarmiento-Soares, L.M., Vieira, J.P., Garcia, A.M.,
967 Teixeira de Mello, F., de Melo, F.A., Meerhoff, M., Attayde, J.L., Menezes, R.F., Mazzeo, N., Di
968 Dario, F., 2016. Fish composition and species richness in eastern South American coastal
969 lagoons: additional support for the freshwater ecoregions of the world. J. Fish Biol. 89(1), 280-
970 314. <https://doi.org/10.1111/jfb.13011>

971 Piggott, J.J., Townsend, C.R., Matthaei, C.D., 2015. Reconceptualizing synergism and antagonism
972 among multiple stressors. Ecol. Evol. 5, 1538-1547. <http://dx.doi.org/10.1002/ece3.1465>.

973 Pignotti, E., Casas, G., Llorca, M., Tellbüscher, A., Almeida, D., Dinelli, E., Farré, M., Barceló, D., 2017.
974 Seasonal variations in the occurrence of perfluoroalkyl substances in water, sediment and fish
975 samples from Ebro Delta (Catalonia, Spain). Sci. Total Environ. 607-608, 933-943.
976 <https://doi.org/10.1016/j.scitotenv.2017.07.025>

977 Pignotti, E., Dinelli, E., 2018. Distribution and partition of endocrine disrupting compounds in water
978 and sediment: Case study of the Romagna area (North Italy). J. Geochem. Explor. 195, 66-77.
979 <https://doi.org/10.1016/j.gexplo.2018.02.008>

980 Pignotti, E., Guerra, R., Covelli, S., Fabbri, E., Dinelli, E., 2018. Sediment quality assessment in a
981 coastal lagoon (Ravenna, NE Italy) based on SEM-AVS and sequential extraction procedure. Sci.
982 Total Environ. 635, 216-227. <https://doi.org/10.1016/j.scitotenv.2018.04.093>

983 Piló, D., Carvalho, S., Pereira, P., Gaspar, M.B., Leitão, A., 2017. Is metal contamination responsible
984 for increasing aneuploidy levels in the Manila clam *Ruditapes philippinarum*? Sci. Total Environ.
985 577, 340-348. <http://dx.doi.org/10.1016/j.scitotenv.2016.10.193>.

986 Ravindra, K., Bencs, L., Van Grieken, R., 2004. Platinum group elements in the environment and their
987 health risk. Sci. Total Environ. 318 (1-3), 1-43. [https://doi.org/10.1016/S0048-9697\(03\)00372-3](https://doi.org/10.1016/S0048-9697(03)00372-3)

988 Rocha, T.L., Gomes, T., Durigon, E.G., Bebianno, M.J., 2016. Subcellular partitioning kinetics,
989 metallothionein response and oxidative damage in the marine mussel *Mytilus galloprovincialis*
990 exposed to cadmium-based quantum dots. Sci. Total Environ. 554-555, 130-141.
991 doi:10.1016/j.scitotenv.2016.02.168

992 Ruchter, N., Sures, B., 2015. Distribution of platinum and other traffic related metals in sediments
993 and clams (*Corbicula* sp.). Water Res. 70, 313-324.
994 <https://doi.org/10.1016/j.watres.2014.12.011>

995 Saidi, I., Said, O.B., Abdelmalek, J.B., Chicharo, L., Beyrem, H., 2019. Assessment of Heavy Metal
996 Contamination in the Sediment of the Bizerte Lagoon in Northern Tunisia. In Recent Advances
997 in Geo-Environmental Engineering, Geomechanics and Geotechnics, and Geohazards, pp. 41-
998 44. Springer, Cham. https://doi.org/10.1007/978-3-030-01665-4_10

999 Shaw, J.P., Dondero, F., Moore, M.N., Negri, A., Dagnino, A., Readman, J.W., Lowe, D.R., Frickers,
1000 P.E., Beesley, A., Thain, J.E., Viarengo, A., 2011. Integration of biochemical, histochemical and

Formatted: Italian (Italy)

Formatted: Spanish (Spain)

1
2
3
4
5
6
7
1008 toxicogenomic indices for the assessment of health status of mussels from the Tamar Estuary,
1009 U.K. Mar. Environ. Res. 72, 13-24. <https://doi.org/10.1016/j.marenvres.2011.05.003>

1000 Soprani, S., Ricci, C., Farina, D., Tarroni, C., Zanetti, S., 1994. Indagine sulle caratteristiche delle
1001 acque. In: Soprani S, Ricci C, editors. Analisi dello stato ambientale e sanitario nelle valli
1002 ravennati. La Pialassa Baiona. Ravenna: Azienda U.S.L., Dipartimento dei Servizi di Prevenzione;
1003 1994. pp. 201-254.

1004 Trombini, C., Fabbri, D., Lombardo, M., Vassura, I., Zavoli, E., Horvat, M., 2003. Mercury and
1005 methylmercury contamination in surficial sediments and clams of a coastal lagoon (Pialassa
1006 Baiona Ravenna Italy). Cont. Shelf Res. 23, 1821-1831.
1007 <https://doi.org/10.1016/j.csr.2003.06.010>

1008 Turja, R., Höher, N., Snoeijs, P., Baršiene, J., Butrimavičiene, L., Kuznetsova, T., Kholodkevich, S. V.,
1009 Devier, M.H., Budzinski, H., Lehtonen, K.K., 2014. A multibiomarker approach to the assessment
1010 of pollution impacts in two Baltic Sea coastal areas in Sweden using caged mussels (*Mytilus*
1011 *trossulus*). Sci. Total Environ. 473-474, 398-409.
1012 <http://doi.org/10.1016/j.scitotenv.2013.12.038>

1013 UNEP/RAMOG: Manual on the Biomarkers Recommended for the MED POL Biomonitoring
1014 Programme. UNEP, Athens, 1999.
1015 http://wedocs.unep.org/bitstream/handle/20.500.11822/1771/ramoge_english.pdf?sequence=1&isAllowed=y (Accessed October 2019).

1016 Valbonesi, P., Sartor, G., Fabbri, E., 2003. Characterization of cholinesterase activity in three bivalves
1017 inhabiting the North Adriatic Sea and their possible use as sentinel organisms for biosurveillance
1018 programmes. Sci. Total Environ. 312, 79-88. [http://doi.org/10.1016/S0048-9697\(03\)00227-4](http://doi.org/10.1016/S0048-9697(03)00227-4).

1019 Van Cauwenberghe, L., Claessens, M., Vandegehuchte, M.B., Janssen, C.R., 2015. Microplastics are
1020 taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural habitats.
1021 Environ. Pollut., 199, 10-17. <https://doi.org/10.1016/j.envpol.2015.01.008>

1022 Vassura, I., Foschini, F., Baravelli, V., Fabbri, D., 2005. Distribution of alternant and non-alternant
1023 polycyclic aromatic hydrocarbons in sediments and clams of the Pialassa Baiona Lagoon
1024 (Ravenna, Italy). Chem. Ecol., 21, 415-424. <https://doi.org/10.1080/02757540500438490>.

1025 Viarengo, A., Moore, M.N., Mancinelli, G., Mazzucotelli, A., Pipe, R.K., Farrar, S.V., 1987.
1026 Metallothioneins and lysosomes in metal toxicity and accumulation in marine mussels: the
1027 effect of cadmium in the presence and absence of phenanthrene. Mar. Biol. 94(2), 251-257.
1028 <https://doi.org/10.1007/BF00392937>

1029 Viarengo, A., Canesi, L., Pertica, M., Mancinelli, G., Accomando, R., Smaal, A.C., Orunesua, M.,
1030 1995. Stress on Stress Response: A Simple Monitoring Tool in the Assessment of a General
1031 Stress Syndrome in Mussels. Mar. Environ. Res. 39, 245-248. [https://doi.org/10.1016/0141-1136\(94\)00075-Z](https://doi.org/10.1016/0141-1136(94)00075-Z)

1032 Viarengo, A., Ponzano, E., Dondero, F., Fabbri, R., 1997. A simple spectrophotometric method for MT
1033 evaluation in marine organisms: an application to Mediterranean and Antarctic molluscs. Mar.
1034 Environ. Res. 44, 69-84. [https://doi.org/10.1016/S0141-1136\(96\)00103-1](https://doi.org/10.1016/S0141-1136(96)00103-1)

1035 Viarengo, A., Burlando, B., Evangelisti, V., Mozzone, S., Dondero, F., 2001. Sensitivity and specificity
1036 of metallothionein as a biomarker for aquatic environment biomonitoring. Biomarkers in
1037 Marine Organisms, A Practical Approach. Ed. Elsevier; Chapt. 2, pp 29-43.
1038 <https://doi.org/10.1016/B978-044482913-9/50004-3>

Field Code Changed

Formatted: Spanish (Spain)

1
2
3
4
5
6
7
1044 Viarengo, A., Lowe, D., Bolognesi, C., Fabbri, E., Koehler, A., 2007a. The use of biomarkers in
1045 biomonitoring: a 2-tier approach assessing the level of pollutant-induced stress syndrome in
1046 sentinel organisms. *Comparative Biochemistry and Physiology Part C: Toxicology &*
1047 *Pharmacology* 146(3), 281-300. <https://doi.org/10.1016/j.cbpc.2007.04.011>.
1048
1049 Viarengo, A., Dondero, F., Pampanin, D.M., Fabbri, R., Poggi, E., Malizia, M., Bolognesi, C., Perrone,
1050 E., Gollo, E., Cossa, G.P., 2007b. A biomonitoring study assessing the residual biological effects
1051 of pollution caused by the HAVEN wreck on marine organisms in the Ligurian sea (Italy). *Arch.*
1052 *Environ. Contam. Toxicol.*, 53, 607-616. <https://doi.org/10.1007/s00244-005-0209-2>.
1053
1054 Yigit, M., Celikkol, B., Yilmaz, S., Bulut, M., Ozalp, B., Dwyer, R.L., Maita, M., Kizilkaya, B., Yigit, Ü.,
1055 Ergün, S., Gürses, K., Buyukates, Y., 2018. Bioaccumulation of trace metals in Mediterranean
1056 mussels (*Mytilus galloprovincialis*) from a fish farm with copper-alloy mesh pens and potential
1057 risk assessment. *Hum. Ecol. Risk Assess.* 24(2), 465-481.
1058 <https://doi.org/10.1080/10807039.2017.1387476>
1059
1060 Yuan, L., Richardson, C.J., Ho, M., Willis, C.W., Colman, B.P., Wiesner, M.R., 2018. Stress responses
1061 of aquatic plants to silver nanoparticles. *Environ. Sci. Technol.* 52, 2558-2565.
1062 <https://doi.org/10.1021/acs.est.7b05837>
1063
1064 Zimmermann, S., Sures, B., 2018. Lessons learned from studies with the freshwater mussel
1065 *Dreissena polymorpha* exposed to platinum, palladium and rhodium. *Sci. Total Environ.* 615,
1066 1396-1405. <https://doi.org/10.1016/j.scitotenv.2017.09.204>
1067
1068 Zuzolo, D., Cicchella, D., Doherty, A. L., Albanese, S., Lima, A., De Vivo, B., 2018. The distribution of
1069 precious metals (Au, Ag, Pt, and Pd) in the soils of the Campania Region (Italy). *J. Geochem.*
1070 *Explor.* 192, 33-44. <https://doi.org/10.1016/j.gexplo.2018.03.009>

Formatted: English (United Kingdom)

Tables of the paper:

Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring

Authors:

Greggio Nicolas*, Capolupo Marco, Filippo Donnini, Manfred Birke, Fabbri Elena, Dinelli Enrico

Table 1. Results of the chemical analyses on sediments. The table includes either pseudo total Aqua Regia (AR) and total XRF data. Table includes also the ranges observed in boreholes in the area, considered representative of the natural background (Amorosi et al., 2002; Curzi et al., 2006). Bold elements exceed natural background. AR DL indicates Detection Limit for Aqua Regia extraction.

		S1		S2		S3		S4		S5		S6		S7		Natural background	
		AR DL	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	AR	XRF	
Ag	µg/kg	2	991	-	269	-	268	-	173	-	275	-	179	-	62	-	-
Al	%	0.01	1.39	6.31	1.41	7.02	1.23	6.98	1.10	6.97	1.35	7.09	1.39	7.07	1.41	6.55	4.4-9.7
As	mg/kg	0.1	7.8	-	5.9	-	6.1	-	5.9	-	7.8	-	6.8	-	6.7	-	-
Au	µg/kg	0.2	16.2	-	5.4	-	11.1	-	2.6	-	5.1	-	3.4	-	1	-	-
B	mg/kg	1	25	-	28	-	26	-	24	-	38	-	33	-	22	-	-
Ba	mg/kg	0.5	61.9	263	43.7	218	41.8	260	42.6	322	40.1	258	38.7	294	36.5	258	220-530
Be	mg/kg	0.1	0.4	-	0.7	-	0.7	-	0.8	-	0.5	-	0.6	-	0.6	-	-
Bi	mg/kg	0.02	0.6	-	0.38	-	0.35	-	0.26	-	0.38	-	0.31	-	0.28	-	-
Ca	%	0.01	11.27	11.37	7.45	7.51	7.87	7.96	7.34	7.87	6.10	6.65	7.24	7.43	8.61	9.52	0.3-17.2
Cd	mg/kg	0.01	0.68	-	0.29	-	0.33	-	0.21	-	0.5	-	0.35	-	0.11	-	-
Ce	mg/kg	0.1	19.4	42	17	55	17.6	71	16.8	52	17.8	64	18.3	42	20.1	58	14-83
Co	mg/kg	0.1	10.2	12	10.9	11	10.7	12	9.9	11	11.7	14	11.5	12	10.6	11	5-29
Cr	mg/kg	0.5	70.6	131	57.5	103	51.4	132	47.7	131	59.9	138	59.3	130	47	100	96-270
Cs	mg/kg	0.02	1.36	-	1.15	-	1.08	-	1.16	-	1.11	-	1.39	-	1.57	-	-
Cu	mg/kg	0.01	84.1	87	49.1	52	49.5	53	36.7	44	51.1	52	37.5	38	22.4	24	4-41
Fe	%	0.01	2.17	3.42	2.46	3.92	2.23	3.86	2.00	3.69	2.54	3.83	2.30	3.64	2.36	3.72	2.1-5.3
Ga	mg/kg	0.1	4.4	-	4.2	-	4	-	3.9	-	4	-	4.2	-	4.4	-	-
Ge	mg/kg	0.1	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	<0.1	-	-
Hf	mg/kg	0.02	0.03	-	0.03	-	0.04	-	0.03	-	0.03	-	0.03	-	0.05	-	-
Hg	mg/kg	0.005	25.7	-	1.42	-	1.48	-	1.89	-	2.38	-	2.26	-	0.1	-	-
In	mg/kg	0.02	0.03	-	0.03	-	0.02	-	0.02	-	0.03	-	0.02	-	0.03	-	-
K	%	0.01	0.33	1.77	0.36	1.98	0.31	2.06	0.28	2.04	0.39	1.97	0.36	2.04	0.35	2.03	0.9-3.3
La	mg/kg	0.5	8.3	19	7.7	26	7.3	22	6.9	18	8	20	8.5	34	9.5	20	17-47
Li	mg/kg	0.1	26.1	-	30.6	-	27.8	-	24.9	-	28.7	-	28	-	28.1	-	-
Mg	%	0.01	1.37	2.41	1.34	2.38	1.30	2.38	1.22	2.44	1.42	2.54	1.55	2.55	1.58	2.35	2.2-4.2
Mn	mg/kg	1	531	893	527	865	549	896	513	949	511	801	550	897	644	1396	230-1470
Mo	mg/kg	0.01	2.67	-	3.33	-	2.61	-	2.28	-	3.34	-	1.57	-	0.54	-	-
Na	%	0.001	0.85	0.68	0.98	0.69	0.90	0.75	0.65	0.99	1.18	0.67	0.85	0.78	0.42	0.85	0.5-2.1
Nb	mg/kg	0.02	0.37	15	0.28	13	0.25	13	0.29	14	0.34	16	0.37	10	0.29	13	8-20
Ni	mg/kg	0.1	50.9	73	56.5	73	50.7	88	46.3	72	55.3	82	51.5	74	46.4	57	39-148
P	%	0.001	0.11	0.10	0.06	0.06	0.06	0.07	0.06	0.06	0.08	0.07	0.07	0.06	0.06	0.06	0.03-0.08
Pb	mg/kg	0.01	27.92	28	22.71	26	22.14	30	18.16	19	26.22	30	20.71	22	11.99	18	5-29
Pd	µg/kg	10	46	-	10	-	12	-	11	-	10	-	<10	-	<10	-	-
Pt	µg/kg	2	37	-	6	-	4	-	4	-	4	-	<2	-	<2	-	-
Rb	mg/kg	0.1	24.9	113	22.2	128	20.8	128	22.4	113	22.3	130	25.1	114	29.8	89	56-200
Re	µg/kg	1	7	-	1	-	2	-	<1	-	2	-	2	-	<1	-	-
S	%	0.02	0.6	-	0.55	-	0.6	-	0.54	-	0.72	-	0.71	-	0.34	-	-
Sb	mg/kg	0.02	0.51	-	0.29	-	0.25	-	0.19	-	0.37	-	0.29	-	0.19	-	-
Sc	mg/kg	0.1	3.4	11	3.6	11	3.4	6	3.2	8	3.7	10	3.7	8	4.1	13	12-21
Se	mg/kg	0.1	0.9	-	0.7	-	0.7	-	0.5	-	1.1	-	0.7	-	0.5	-	-
Si	%	-	-	17.72	-	19.43	-	19.90	-	20.61	-	19.31	-	20.95	-	20.03	14.7-29.3
Sn	mg/kg	0.1	3.2	-	1.8	-	1.5	-	1.3	-	1.9	-	1.5	-	0.8	-	-
Sr	mg/kg	0.5	353	368	274.4	300	224.8	252	218	265	259.1	281	281.6	291	290.5	310	113-442
Th	mg/kg	0.1	3.5	4	4.1	5	3.6	17	3.6	25	4	11	4.2	10	4.7	7	<3-19
Ti	%	0.001	0.009	0.308	0.005	0.328	0.006	0.347	0.008	0.349	0.006	0.324	0.010	0.334	0.015	0.336	0.02-0.58
Tl	mg/kg	0.02	0.26	-	0.17	-	0.21	-	0.19	-	0.27	-	0.2	-	0.14	-	-
U	mg/kg	0.1	1.3	-	1.5	-	1.2	-	1.1	-	1.3	-	1.2	-	1.1	-	-
V	mg/kg	2	32	85	33	85	30	99	26	94	35	113	33	94	31	76	54-169
Y	mg/kg	0.01	10.02	18	9.39	23	9.28	24	8.86	20	9.7	24	9.63	19	10.58	19	18-43
Zn	mg/kg	0.1	306.6	316	129.8	152	133.1	154	101.2	126	146.9	174	109.3	124	58.2	67	43-124
Zr	mg/kg	0.1	1.1	108	1.2	91	1	105	1	124	1.1	86	1.2	92	1.9	106	76-260
H₂O⁻	%	-	-	2.7	-	4.0	-	2.8	-	2.4	-	3.7	-	2.7	-	2.1	-
OM	%	-	-	2.8	-	1.8	-	2.0	-	1.7	-	3.4	-	2.2	-	1.1	-
H₂O⁺	%	-	-	6.0	-	6.5	-	5.4	-	4.4	-	6.5	-	5.8	-	2.3	-
CO₂	%	-	-	11.6	-	8.9	-	9.1	-	9.3	-	8.6	-	7.4	-	12.3	-
LOI	%	-	-	23.1	-	21.2	-	19.3	-	17.8	-	22.3	-	18.1	-	17.8	4.8-29.3

Table 2. Results of the chemical analyses on mussel tissues (bioaccumulation). Data refers to T0 and to the 5 sites were mussels survived or remained after the exposure period.

	unit	T0	S2	S3	S5	S6	S7
Ag	mg/kg	0.12	0.99	0.98	1.21	0.62	0.82
Al	mg/kg	36	39	56	98	82	47
As	mg/kg	9.83	12.8	9.98	11.9	12.4	11.4
B	mg/kg	25.7	22.9	19.7	22.8	25.9	19.4
Ba	mg/kg	1.95	0.35	0.49	0.54	0.52	0.36
Be	mg/kg	0.003	0.003	0.003	0.005	0.006	0.004
Bi	mg/kg	0.057	0.009	0.009	0.007	0.011	0.008
Ca	mg/kg	2004	1322	1299	1726	1679	1157
Cd	mg/kg	0.70	0.71	0.78	0.76	0.77	0.50
Ce	mg/kg	0.118	0.051	0.092	0.124	0.166	0.075
Co	mg/kg	0.54	0.323	0.3	0.303	0.511	0.315
Cr	mg/kg	0.57	0.6	0.53	0.67	0.53	0.64
Cs	mg/kg	0.022	0.014	0.016	0.022	0.019	0.015
Cu	mg/kg	4.65	2.71	2.63	2.81	2.66	1.46
Fe	mg/kg	67.5	68.1	77.4	116.0	103.4	91.7
Ga	mg/kg	0.018	0.007	0.011	0.019	0.013	0.009
Hg	mg/kg	0.082	0.35	0.32	0.41	0.27	0.16
K	mg/kg	11084	10988	9729	11129	11244	9535
La	mg/kg	0.138	0.035	0.057	0.074	0.109	0.06
Li	mg/kg	0.88	0.46	0.41	0.51	0.58	0.34
Mg	mg/kg	4198	3851	3467	3676	4422	2506
Mn	mg/kg	4.84	5.19	7.50	8.16	8.03	5.03
Mo	mg/kg	0.645	0.353	0.368	0.317	0.394	0.274
Na	mg/kg	22885	20201	15986	19714	24751	13438
Nb	mg/kg	0.002	0.001	0.002	0.003	0.003	0.002
Ni	mg/kg	3.61	0.72	0.69	0.73	0.92	0.64
Pb	mg/kg	1.41	0.87	0.52	0.57	0.94	0.64
Rb	mg/kg	5.29	4.08	4.32	4.27	4.3	3.78
Sb	mg/kg	0.018	0.038	0.031	0.03	0.048	0.039
Sc	mg/kg	0.04	0.04	0.03	0.05	0.07	0.04
Se	mg/kg	2.27	2.10	2.10	2.01	1.83	1.52
Sn	mg/kg	0.12	0.12	0.12	0.10	0.09	0.08
Sr	mg/kg	31.2	17.7	16.5	20.6	22.8	14.3
Th	mg/kg	0.019	0.011	0.014	0.025	0.031	0.016
Ti	mg/kg	0.49	0.34	0.62	1.07	0.88	0.50
Tl	mg/kg	0.003	0.001	0.002	0.002	0.001	0.001
U	mg/kg	0.127	0.049	0.040	0.054	0.065	0.032
V	mg/kg	1.15	0.44	0.49	0.55	0.80	0.47
Y	mg/kg	0.080	0.046	0.071	0.085	0.094	0.04
Zn	mg/kg	107.1	116	62.4	95.6	134	86.9
Zr	mg/kg	0.064	0.013	0.025	0.025	0.021	0.018

Figures of the paper:

Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring

Authors:

Greggio Nicolas*, Capolupo Marco, Filippo Donnini, Manfred Birke, Fabbri Elena, Dinelli Enrico




Figure 1. Location map and sampling points (S1 to S7) within the Pialassa Baiona lagoon (Ravenna, Italy). The abbreviations “R.” and “Ch.” refer to the main rivers and/or channels which supply freshwater to the lagoon. The yellow shape represents the industrial area settled at the south-eastern boundary of the Pialassa Baiona lagoon.




Figure 2. Metal enrichment in sediments from selected sites within the Pialassa Baiona lagoon. Data are normalized on Site 7 (S7), which was selected as internal reference based on previous literature (Trombini et al., 2003; Fabbri et al., 2006; Donnini et al., 2007; Franzellitti et al., 2010). Elements in italics (*Pt*, *Au*, *Re*, *Pd*) were not investigated in the mussel tissues.




Figure 3. Bioaccumulation factor of metals in mussel tissues. The analysis has been performed on the tissue of mussels deployed for 30 days into different sites of the Pialassa Baiona lagoon compared to mussels analysed before the *in situ* deployment (T0). Elements are ranked according to average enrichment.




Figure 4. Biomarker modulation in mussels transplanted into the Pialassa Baiona lagoon. Data from all sites (mean \pm SEM) have been measured in mussels collected after 30 days of exposure, except for S1, referred to mussels collected after 3 days of exposure due to high mortality. A, survival (SURV); B, Stress on stress (SOS); C, lysosomal membrane stability (LMS); D, lysosome to cytoplasm volume ratio; E, lipofuscin (LIF), F, catalase activity (CAT); G, metallothionein content (MT); H, acetylcholinesterase activity (ACHE); I, peroxisomal acyl-CoA oxidase (AOX). Asterisks indicate statistically significant differences compared to the control site S7 (One-Way ANOVA, Duncan's post-hoc test).




Figure 5. Correlation between biological data and metal concentrations. Pearson's coefficients calculated between biological data and levels of metals measured in mussel tissues (bioaccumulation) and sediment. Only coefficients showing a significant correlation ($p < 0.05$) are reported in the figure.





Figure 6. Biomarker data integration. A unique Health status index (HSI) has been generated using the Mussel Expert System (MES, developed by Dagnino et al., 2007) based on biomarker alterations observed at each site. The LMS was chosen as guide parameter for the HSI assessment.



[Click here to access/download](#)

Supplementary Data

Revised Supplementary Material Greggio et al.
2020.docx



Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

TITLE: Integration of physical, geochemical and biological analyses as a strategy for coastal lagoon biomonitoring

Authors:

Greggio Nicolas*^{a, b}, Capolupo Marco^{a, b}, Filippo Donnini^a, Manfred Birke^c, Fabbri Elena^{a, b}, Dinelli Enrico^{a, b}

^a University of Bologna, Inter-Departmental Research Centre for Environmental Science (CIRSA), Via S. Alberto 163, 48123 Ravenna, Italy

^b University of Bologna, Department of Biological, Geological, and Environmental Sciences, Piazza S. Donato 1, 40100 Bologna, Italy

^c Federal Institute for Geosciences and Natural Resources, Stilleweg 2, 30655, Hannover, Germany

* Corresponding Author: Nicolas Greggio e-mail: nicolas.greggio2@unibo.it

Credit Author Statement

Nicolas Greggio: Conceptualization, Data Curation, Formal analysis, Investigation, Visualization, Writing - Original Draft, Writing - Review & Editing. **Marco Capolupo:** Data Curation, Formal analysis, Investigation, Visualization, Writing - Original Draft, Writing - Review & Editing. **Filippo Donnini:** Conceptualization, Formal analysis, Investigation. **Manfred Birke:** Formal analysis Writing - Review & Editing. **Elena Fabbri:** Funding acquisition, Project administration, Resources, Supervision, Validation, Writing - Review & Editing. **Enrico Dinelli:** Funding acquisition, Project administration, Resources, Supervision, Validation, Writing - Review & Editing.