



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Plant foods in the Late Palaeolithic of Southern Italy and Sicily: Integrating carpological and dental calculus evidence

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Marialetizia Carra, A.Z. (2023). Plant foods in the Late Palaeolithic of Southern Italy and Sicily: Integrating carpological and dental calculus evidence. QUATERNARY INTERNATIONAL, special issue, 1-16 [10.1016/j.quaint.2022.06.007].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/919724> since: 2024-06-18

*Published:*

DOI: <http://doi.org/10.1016/j.quaint.2022.06.007>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

# Plant foods in the Late Palaeolithic of southern Italy and Sicily: Integrating carpological and dental calculus evidence

Marialetizia Carra<sup>a</sup>, Andrea Zupancich<sup>a</sup>, Elena Fiorin<sup>a</sup>, Lucia Sarti<sup>b</sup>, Domenico Lo Vetro<sup>c,d</sup>, Fabio Martini<sup>c,d</sup>, Emanuela Cristiani<sup>a,\*</sup>

<sup>a</sup> DANTE, Diet and ANcient TEchnology Laboratory, Department of Oral and Maxillo-Facial Sciences, SAPIENZA University of Rome | Via Caserta 6, 00161, Rome, Italy

<sup>b</sup> Dipartimento di Scienze Storiche e dei Beni Culturali, Università degli Studi di Siena, Via Roma 47, 53100, Siena, Italy

<sup>c</sup> Unità di Archeologia Preistorica, Dipartimento SAGAS, Università di Firenze, Via Sant'Egidio 21, Firenze, Italy

<sup>d</sup> Museo e Istituto Fiorentino di Preistoria, Via Sant'Egidio 21, Firenze, Italy

\* *Corresponding author.*

*E-mail address: [emanuela.cristiani@uniroma1.it](mailto:emanuela.cristiani@uniroma1.it) (E. Cristiani).*

## Abstract

Several caves from southern Italy and Sicily provided invaluable evidence, including several human burials, for reconstructing human adaptations and subsistence in the area during the Upper Palaeolithic. A wealth of information is available concerning the exploitation of animal resources as food. However, little is still known about the role of plants in the diet of the ancient hunter-gatherers of the region. By combining the carpological data with vegetal micro-debris entrapped in human dental calculus, we provide new clues about the dietary role of plant foods in the analysed area during the Late Glacial. Our study focused on five key sites from southern Italy and Sicily: Grotta della Serratura in Campania, Grotta del Romito in Calabria, Grotta del Cavallo in Apulia, Grotta di San Teodoro and Grotta d'Oriente in Sicily. Our results demonstrate that several plant resources were exploited at these sites, including seeds, grasses, and forest fruits. Amongst the carpological remains recovered, several plant taxa are known for their medical properties. Overall, our study explores how critical climatic and environmental changes characterising the timeframe between the end of the Last Glacial Maximum and the beginning of the Holocene affected Late Upper Palaeolithic dietary habits. Moreover, it shows the methodological relevance of combining different strands of archaeological evidence for reconstructing the Palaeolithic diet.

*Keywords:* Upper Palaeolithic; Plant foods; Dental Calculus; Carpological analysis; Late Glacial; southern Italy and Sicily

## 1. Introduction

In southern Italy and Sicily, numerous caves and rock-shelters yielded a wealth of prehistoric evidence, including several human burials, and offer a unique opportunity to explore how the critical climatic and environmental changes of the Late Glacial affected human/environment interaction, eventually impacting human dietary strategies (Martini et al., 2009b; 2009a; Mannino et al., 2011a; 2016; Colonese et al., 2018).

A series of climate events mark this period in Mediterranean Europe, leading to the oscillations of temperate/humid and cold/arid climatic phases (Watts et al., 1996; Seguinot et al., 2018). In particular, at the beginning of the Bølling-Allerød interstadial (around 14700 cal BP), we assist in a rapid rise in temperatures and rainfall. Conversely, during the Older and the Younger Dryas cold stadials, a decrease in the forest cover is documented along with an increase in open areas. At the end of Pleistocene, rainfall intensification together with a strong temperature growth (ca. 7° C) marked the Younger Dryas-Preboreal transition, dated approximately 11650 cal BP, (Orbelli and Ravazzi, 1996). On a palaeoenvironmental ground, phases dominated by a rich vegetation cover alternate with conditions of a more open environment. The latter corresponds to a lower variety in Mediterranean arboreal and shrub species, which show a decrease in the presence of forest species in the arid climatic phases (Martini et al., 2009b).

52

53 The diet of Palaeolithic hunter-gatherer communities in southern Italy and Sicily during the Late Glacial has  
54 been the subject of intense research in the last decade through the application of two main methodologies:  
55 archaeozoological analysis and bone chemistry (namely carbon and nitrogen isotopic composition of bulk  
56 collagen) (Drucker and Bocherens, 2004; Naito et al., 2013). These methods primarily rely on the study of  
57 bones and teeth, which generally preserve better than plant remains in ancient sediments due to their high  
58 mineral/inorganic content. Based on the standard dietary reconstructions carried out for cave sites such as  
59 Grotta del Romito and Grotta Paglicci in southern Italy, as well as Grotta di San Teodoro and Grotta d'Oriente  
60 in Sicily, we know that late Upper Palaeolithic hunter-gatherers primarily exploited the hinterland for the  
61 specialized hunting of large herbivores, such as red deer, aurochs and hydruntine ass (Craig et al., 2010; Martini  
62 et al., 2012b; Vacca, 2012; Lugli et al., 2019). Based on the isotopic data, marine or coastal lagoon resources,  
63 such as fish, limpets, and cockle shells, seem to have contributed seldomly to the diet of Palaeolithic human  
64 groups of the Tyrrhenian regions of southern Italy and Sicily (Francalacci, 1988; Vigne, 2004; Martini et al.,  
65 2007; Mannino and Thomas, 2009; Paine et al., 2009; Craig et al., 2010; Lightfoot et al., 2011; Mannino et al.,  
66 2011b; 2011a; 2012; Goude et al., 2017; Colonese et al., 2018). Supposedly low productivity of the central  
67 Mediterranean Sea in respect to other contexts and the absence of specific technological equipment (e.g., fish  
68 hooks) were suggested for explaining such a behaviour (Mannino et al., 2011b; Colonese et al., 2014).  
69 However, this scenario changed with the beginning of the Early Holocene, when marine resources were made  
70 more available to Mesolithic groups due to the rise of the sea level (Colonese et al., 2018). Whether human  
71 dietary habits also involved plant foods during the final stages of the Late-Glacial is still unknown, primarily  
72 due to the limits in assessing plant isotopic composition (Drucker and Bocherens, 2004; Craig et al., 2010;  
73 Mannino et al., 2011a; 2016; Gazzoni et al., 2013). Nevertheless, micro-wear patterns observed on the teeth  
74 and trace element chemistry on bones have provided indirect evidence of the possible consumption of plants  
75 foods in the region during the Late Upper Palaeolithic (Carnieri and Mallegni, 2006).

76

77 Recently, new methods of analysis applied to human remains and material culture revealed that grasses and  
78 plant underground storage organs might have supplemented Upper Palaeolithic diet in Italy from the Late  
79 Glacial Maximum onwards. At the site of Bilacino II in Tuscany (central Italy), use-wear and residues have  
80 been identified on both non-flaked and flaked tools coming from the Gravettian levels dated to ca. 33,000 cal  
81 BP and suggested the processing of starchy rhizomes by local hunter-gatherer communities (Revedin et al.,  
82 2010). Plant microremains were preserved on grinding tools from Grotta Paglicci, revealing the consumption  
83 of oat and acorns in the same period in northern Apulia (southern Italy) (Revedin et al., 2010; 2015; Lippi et  
84 al., 2015). Starch granules were also identified in the dental calculus of Late Upper Palaeolithic individuals  
85 from Grotta Continenza, Riparo Tagliente, and Riparo Villabruna, together with microwear patterns supporting  
86 the consumption of plant resources (Oxilia et al., 2020; Nava et al., 2021). These works have provided the first  
87 direct evidence of a diffuse broadening in the subsistence base of Upper Palaeolithic groups from the Last  
88 Glacial and throughout the phases that preceded the onset of the Holocene (Revedin et al., 2010; 2015; Lippi  
89 et al., 2015; Oxilia et al., 2020; Nava et al., 2021).

90 Remarkably, one of the most accredited methods for recovering plant food remains in ancient deposits (i.e.,  
91 flotation) has only sporadically been applied to Italian Palaeolithic excavations (Carra and Marínval, 2011;  
92 Carra and Cristiani, 2020), being time-consuming and characterized by a low recovery rate. However,  
93 systematic flotation applied to Palaeolithic deposits has demonstrated that carpological macroremains can  
94 preserve quite well even without particular depositional conditions (Piperno et al., 2004; Weiss et al., 2004;  
95 2008; Carra and Marínval, 2011; Pryor et al., 2013; Carra and Cristiani, 2020).

96

97 In this paper, we contribute to the debate on the role of plant foods in the diet of Upper Palaeolithic hunter-  
98 gatherers from southern Italy and Sicily by exploring complementary evidence from five renowned  
99 archaeological contexts, which yielded long-lasting evidence of Late Glacial occupation: Grotta della Serratura  
100 in Campania, Grotta del Romito in Calabria, Grotta del Cavallo in Apulia, Grotta di San Teodoro and Grotta

101 d'Oriente in Sicily. Over the years, the evidence from these sites has been the object of interdisciplinary  
102 investigations, including palaeoenvironmental and archaeozoological studies, human anthropology and  
103 genomics, lithic technology and ornamental traditions, and meat-based dietary strategies (Martini et al.,  
104 2009b). However, direct evidence regarding dietary plant use still lacks for this region. We aim to fill this gap  
105 by integrating two different strands of evidence for plant consumption: (a) carpological data in the form of  
106 charred plant remains and (b) plant micro-debris entrapped in human dental calculus.  
107 Based on our complimentary data for southern Italy and Sicily, we suggest that a broader range of dietary  
108 resources characterized the Late Upper Palaeolithic diet and included a wide selection of plant taxa.  
109

## 110 **2. Archaeological contexts and materials**

111

112 In this paper, we consider five key Palaeolithic caves located along the Tyrrhenian coast of southern Italy  
113 (Grotta della Serratura, Grotta del Romito), in Sicily (Grotta di San Teodoro and Grotta d'Oriente), and by the  
114 Ionian coast (Grotta del Cavallo) (fig. 1). These sites were all discovered during the middle of the last century,  
115 except for Grotta di San Teodoro which was found in 1859. These contexts have a long history of excavations  
116 and numerous multidisciplinary analyses have been conducted on their Palaeolithic record, including a good  
117 assessment of their stratigraphy and environmental reconstructions detailing the various moments of their  
118 occupation. During the Late Pleistocene, variations of the sea levels caused by changes in the climate  
119 conditions affected their distance from the shore. The stratigraphy of the sites spans the Middle Palaeolithic  
120 up to the Holocene. Their Late Upper Palaeolithic occupations represent the focus of this work (Table 1).  
121

122 **FIG.1 HERE**

123

### 124 *2.1 Grotta della Serratura*

125 Grotta della Seratura is part of a complex of caves that opens along the coast, from Capo Palinuro to Torre  
126 Mozza, and is located ca. 2m a.s.l at the foot of Monte Bulgheria, the village of Marina di Camerota. The site  
127 was discovered during the of the last century, however systematic excavation began in 1984 and yielded a  
128 wealth of evidence related to the occupation of the cave during the Upper Palaeolithic, the Mesolithic and the  
129 Neolithic (Martini, 1993).

130 Excavations took place at two areas of the site: the atrium and the of the cave. The atrium is located 12 m from  
131 the current entrance of the cave. Here occupational layers to the Middle Palaeolithic and the Upper Palaeolithic  
132 have been identified, while the effects of the marine erosion led to the disappearance of deposit associated with  
133 the latest phases of the Pleistocene and the early Holocene. Layers C and D in the atrium trench are attributed  
134 to the Epigravettian while layer E relates to the Gravettian and layer F the Early Upper Palaeolithic. Some  
135 radiocarbon dates of some of these levels are unpublished, the chronological attribution of other ones is based  
136 on the comparison with the stratigraphy of the innermost area. (Martini et al., 2005).

137 In this internal area, located 40 m from the current entrance, a sequence of 10 layers related to Late Pleistocene  
138 and Early Holocene was discovered (Martini, 1993). Specifically, layers 8 to 10 refer to the Late Epigravettian,  
139 layers 7 to 4 correspond to the Early Mesolithic, while layers 3 to 1 are associated with the Neolithic occupation  
140 of the site (Martini, 1993; Colonese et al., 2010; Vadillo Conesa et al., 2021). Several 14C dates are available  
141 for this sequence (Table 1).  
142

143 A total of 56 samples were collected from Grotta della Serratura in both the Gravettian and the Epigravettian  
144 levels (Table 2). The botanical samples come from both the atrium, where layers are identified with the letters  
145 of the alphabet and the back of the cave where levels are distinguished with progressive numbers.  
146  
147  
148

### 149 *2.2 Grotta del Romito*

150 Grotta del Romito is located at 275 m a.s.l, in the Lao Valley, a mountainous region with peaks reaching 2000  
151 m, situated ca. 25 km away from the Tyrrhenian coast. The cave is sited in proximity of a narrow tributary of  
152 the Lao river. Excavations at the site exposed a thick deposit of more than 8 meters consisting in a long  
153 sequence of archaeological layers ranging from the Late Pleistocene to the Early Holocene; more than 40  
154 radiocarbon dates are available for the whole sequence (Blockley et al., 2018; Martini and Lo Vetro, 2018).  
155 Due to intense episodes of water runoff, an intermittent occupation of the site is documented during the Last  
156 Glacial, from Evolved Gravettian to Evolved Epigravettian (ca. 23000 to 16000 uncal BP) (Ghinassi et al.,  
157 2009), while an intense human activity characterises the Late Glacial (Late Epigravettian) and the Early  
158 Holocene (Early Mesolithic - Sauveterrian) (Martini, 2002; Martini et al., 2004; 2006; Martini and Lo Vetro,  
159 2005; 2018; Craig et al., 2010) (Table 1). Nine well preserved individuals have been found at the site and  
160 chronologically associated with the Epigravettian occupation of the cave (Graziosi, 1962; 1966; Fabbri et al.,  
161 1989; Mallegni and Fabbri, 1995; Martini, 2006; Craig et al., 2010; Martini and Lo Vetro, 2011). Eight of  
162 these inhumations refer to Late Epigravettian: Romito 1-2 and Romito 5-6, double burials, and Romito 3,4,7,8  
163 single burials; the ninth individual, Romito 9, dating back to the Evolved Epigravettian (De Silva et al., 2015;  
164 Martini and Lo Vetro, 2018) (Table 3).

165 At Grotta del Romito a total of 59 samples were collected inside the cave from Late Epigravettian levels dated  
166 between  $12170 \pm 60$  and  $11660 \pm 70$  BP (Martini and Lo Vetro, 2018) and underwent flotation. Within the  
167 analysed samples, only one comes from the visual screening of the sediments from the 2016 field season (Table  
168 2).

169  
170 A total of 76 samples originally destined for the study of pollen were used to recover macroremains. They  
171 come from 3 different locations: the excavation trench (40), the Romito 8 burial pit (22) and the southern  
172 section within the excavation trench (14). The chronology of the samples is mainly referable to the Late  
173 Epigravettian (D), the deeper layers (F-L) instead refer to the Ancient Epigravettian (17500-18500 BP) and  
174 the evolved Gravettian (20000-24000 BP) (Table 2) (Martini and Lo Vetro, 2018).

175  
176 Dental calculus samples were collected from 5 individuals of Grotta del Romito (Romito 4,5,6,7,8) (Table 4).

### 177 178 *2.3 Grotta del Cavallo*

179 Located at 15 m a.s.l., Grotta del Cavallo is part of a karstic systems of the Uluzzo Bay in southern Apulia.  
180 The site was first discovered in 1961 and excavations exposed a long stratigraphic sequence including  
181 Mousterian (layers M-F), and Uluzzian (layers D and E) occupations separated by a thin ash layer dated to  
182  $45500 \pm 1,0$  ky (Romagnoli, 2015; Martini and Sarti, 2017; 2021). The Uluzzian is separated from the Upper  
183 Palaeolithic by a sterile tephra lens (layer C) while the uppermost layer of the site (layer B) is associated with  
184 the final phases of the Epigravettian (Romanellian facies) and lies below a disturbed Neolithic level (layer A).

185 A total of 53 samples were analyzed from Grotta del Cavallo. The samples come from layers that are attributed  
186 the Neolithic (layer A1), the Mesolithic (layer A2) and the Late Upper Palaeolithic/Epiromanellian (layer B1)  
187 and Romanellian (layer B2) (Table 2), including a small pit with material of uncertain dating.

### 188 *2.4 Grotta di San Teodoro*

189 Located in NE Sicily between the cities of Palermo and Messina, Grotta di San Teodoro stands at an altitude  
190 of 135 m a.s.l on the northern slope of the Nebrodi mountains, about 2km from the sea. Excavations at the site  
191 began at the end of the XIX century and lasted until the first decade the current century. Two main  
192 archaeological units were identified: the upper unit (layers A-D) attributed to late Upper Palaeolithic (Late  
193 Epigravettian) and characterized by intense occupation, abundant lithic artefacts, mammal bones and mollusc  
194 remains (Vigliardi, 1968; Bonfiglio et al., 2008; Catalano et al., 2020; Garilli et al., 2020); and a lower unit  
195 (layers E-G), which yielded remains from Upper Pleistocene endemic mammals. During the excavations

196 carried out between 1937 and 1942, seven human burials (ST1-7) were unearthed, all belonging to adult  
197 individuals, two males (ST 3 and, probably, 7), four females (ST 1, 4, 5, 6) and one indeterminable (Fabbri  
198 and Lo Vetro, 2021). Based on the association with faunal and lithic remains six individuals (ST1-4,6,7) were  
199 chronologically attributed to the Late Epigravettian (ca. 15000- 11000 years BP) (Lo Vetro 2012, D'Amore et  
200 al 2009). Recently this attribution has been confirmed by tree almost contemporary AMS dates, all falling  
201 around 12600-12500 uncal BP, obtained on bones from individuals ST1 (Lo Vetro and Martini, 2006; Craig  
202 et al., 2010; Colonese et al., 2011; Martini et al., 2012a; Colonese et al., 2014; 2018) and ST4 and ST5 (Fabbri  
203 and Lo Vetro, 2021) (Table 3). Dental calculus samples were collected from 3 individuals buried at Grotta di  
204 San Teodoro (ST 3,5,6) (Table 4).

205

## 206 *2.5 Grotta d'Oriente*

207 Grotta d'Oriente is located 40 m a.s.l. On the island of Favignana, part of the Egadi archipelago 5km from the  
208 NW coast of Sicily. The cave is made of two main areas, a small chamber at the entrance and a large gallery.  
209 The first excavation at the site dates back to 1972, while more recent and systematic excavations were carried  
210 out in 2005 (Lo Vetro and Martini, 2006; Craig et al., 2010; Colonese et al., 2011; 2014; 2018; Martini et al.,  
211 2012a). The latest excavations exposed a 2-m sequence which is divided in 8 archaeological layers. Layers 7  
212 to 3 are associated with the prehistoric human occupation of the cave and relate to the Late Upper Palaeolithic  
213 (layer 7), Early Mesolithic (layer 6), Late Mesolithic or Early Neolithic (layer 5) and the Bronze Age (layers  
214 4-3) (Martini et al., 2012a; Catalano et al., 2020). Three human burials have been found at the site, namely  
215 Oriente A, Oriente B and Oriente C. Oriente A and B were discovered in 1972 by G. Mannino (2002). Oriente  
216 A is a Mesolithic or Late Palaeolithic adult male burial, while Oriente B is a Mesolithic adult female burial  
217 recently AMS dated to the 10<sup>th</sup> millennium uncal BP (Mannino et al., 2012). Oriente C, found in layer 7, has  
218 not been directly dated, however two AMS dates on charcoal fragments from the layers in which the individual  
219 stands (sublayers 7D, 12149 ± 65 BP, and 7E, 12132 ± 80) are consistent with the cultural attribution to the  
220 Late Epigravettian based on stone tools assemblages (Table 1). Dental calculus from the individual Oriente C  
221 were collected for this study (Table 4).

222

## 223 **3. Materials and Methods**

224

### 225 *3.1 Carpological analysis*

226 A total of 245 archaeobotanical samples were analysed: 136 from the Grotta del Romito, 56 from Grotta della  
227 Serratura, and 53 from Grotta del Cavallo. Samples for carpological study were selected from: (a) sieving  
228 carried out during the excavations (47); (b) systematic flotation of the archaeological sediments performed  
229 during the excavations between 2018 and 2019 (112); and (c) samples previously collected for pollen analysis,  
230 yet no longer suitable for such investigation (86) (see Table 2).

231

232 Samples collected during the 2018 and 2019 field seasons at Grotta del Romito and Grotta del Cavallo are the  
233 most abundant (Table 2). Each of these samples has a volume between 4 and 10 litres of sediment and their  
234 sampling followed the rules of capillarity and homogeneity. The samples were subjected to both manual and  
235 mechanic flotation at Grotta del Romito while only manual flotation was applied to Grotta del Cavallo (500µm  
236 and 300µm mesh sieves). Subsequently the residue was sieved and screened to recover archaeological  
237 materials. Only the light fraction was sifted under the stereomicroscope. The concentration of the finds and the  
238 number of carpological remains was greater at Grotta del Cavallo; at Grotta del Romito, the carpological  
239 remains were mostly diluted in the sediments. No specific structures linked to the accumulation of plants were  
240 found in the two sites; vegetal remains represent usual waste scattered in sediments.

241

242 Old pollen samples, each about 500 ml in volume, come from Grotta del Romito and Grotta della Serratura. It  
243 is known that pollen samples tend to deteriorate the micro-remains over time, especially if not properly stored.

244 Conversely, plant remains can be recovered even from samples that have been stored for a long time in  
245 warehouses. For this reason, we analysed samples collected between 1984 and 1990 at Grotta della Serratura  
246 and at Grotta del Romito during the 2000-2004 field seasons (Table 2).

247 The samples were subjected to manual flotation, using 500µm and 300µm mesh sieves, and residue sieving.  
248 Residues were sifted under a stereomicroscope and both light and heavy fractions were collected.

249  
250 Last, 47 samples are represented by carpological findings recovered during the visual screening of residues  
251 coming from sieving activity carried out using a 1 mm mesh at Grotta della Serratura and Grotta del Romito  
252 (Table 2). These samples were selected during the excavation and only the work of determination was carried  
253 out on them. It was not possible to calculate the concentration of remains per litre in these samples.

254  
255 For the determination of the finds, we used a collection of seeds and fruits of the flora of Italy and the Balkans  
256 is present at Sapienza University (DANTE – Diet and ANcient TEchnology laboratory) (Fig.3) as well as  
257 botanical atlases (Cappers and Neef, 2012; Cappers et al., 2012). The botanical nomenclature refers to Pignatti  
258 2017 (Pignatti, 2017). A multifocal stereo microscope (Zeiss Axio Zoom V16) was used for determination of  
259 remains with magnifications ranging from 10X to 100X. Photographs of the specimens were taken using a  
260 high-definition Zeiss AxioCam 506 colour.

261

**FIG.2 HERE**

262

263

264

### 265 *3.2 Dental calculus – sampling, extraction, decontamination and examination procedures*

266 Dental calculus samples were removed from a total of 10 individuals dated to the Late Upper Palaeolithic (Late  
267 Epigravettian) and buried at Grotta del Romito (Romito 4,5,6,7,8), Grotta di San Teodoro (ST 3,5,6) and Grotta  
268 d’Oriente (Oriente C) (Table 4).

269 The sampling was conducted following the protocol systematized by (Sabin and Fellow Yates, 2020) with  
270 some variation (e.g., disposable blades were changed after each sample extraction).

271 Extraction and decontamination procedures followed standard published protocols (Cristiani et al., 2016;  
272 Radini et al., 2017; Sabin and Fellow Yates, 2020) and were conducted in dedicated clean spaces not connected  
273 to modern botanical work and under strict environmental monitoring of the DANTE laboratory of Sapienza  
274 University of Rome.

275 Cleaning was carried out daily in order to prevent any type of modern contamination. Bench space surfaces  
276 were cleaned prior to the analysis of each sample, using soap and ethanol, followed by covering of the surfaces  
277 by aluminium foil, and using of clean starch-free nitrile gloves at all times. Calculus cleaning was carried out  
278 under the stereomicroscope, on a Petri dish previously washed, with magnifications up to 100X. The removal  
279 of the mineralized soil adhering on the surface of the calculus was meticulously carried out using sterile  
280 tweezers to hold the sample and a fine acupuncture needle to gently scrape off the soil attached to the external  
281 layer of the mineralized plaque. The procedure was performed using drops of 0.5 M HCl acid to dissolve the  
282 mineralized flecks of soil and ultrapure water to block the demineralization, as well as to wash and remove the  
283 contaminants. The mineralized soil removed was stored in plastic tubes for monitoring analysis. Then, the  
284 clean calculus was washed in ultrapure water up to three times to remove any trace of loose sediment, dissolved  
285 in a solution of 0.5 M HCl and subsequently mounted on slides using a solution of 50:50 glycerol and ultrapure  
286 water.

287 Furthermore, control samples from the clean working tables and dust-traps were collected and analysed for  
288 comparative purposes in order to prevent any type of modern contamination– this is a practice routinely done  
289 in our laboratories, even at times where no archaeological analysis occur, to allow a better understanding of  
290 the flow of contaminations through seasons. We did not retrieve any debris morphologically similar to any of  
291 the remains in the environmental control samples. Starch granules amounted to a very minor fraction of the  
292 laboratory ‘dust’.

293 The examination of micro-debris embedded in the calculus matrix was performed using a Zeiss Imager2 cross  
294 with magnifications ranging from 100X to 630X. For the identification of archaeological starch granules, a  
295 modern reference collection of 300 plants native to the Mediterranean region, and Europe was used along with  
296 published literature.

297 **FIG.3 HERE**

## 299 4. Results

### 301 4.1 Carpological remains

302 A total of 587 plant macroremains were retrieved from our study. Specifically, 106 carpological remains were  
303 recovered at Grotta della Serratura; 55 remains at Grotta del Romito; and 426 at Grotta del Cavallo (Table 2,  
304 Figs.4,5).

305 **FIG.4 HERE**

#### 307 4.1.1 Grotta della Serratura

308 The analysis and determination of the samples was carried out separately, depending on the origin of the  
309 sample origins: 46 from the visual screening operations (Supplementary Information 1), and 10 from the  
310 flotation and microscope screening (Supplementary Information 2). The first group contains the largest and  
311 most intact finds, while in the other samples the carpological remains are more fragmentary. Observations after  
312 the determination were made jointly.

313 Overall, 106 carpological remains were found, all belonging to herbaceous plants. Finds related to fruits of  
314 woody plants are instead completely missing. The majority of the findings come from the Final Epigravettian  
315 (layers 8F-G), while only indeterminable fragments were retrieved from the Gravettian deposit (layer 9 and  
316 E3, Fig. 4). Overall, our results reveal an exploitation of vegetable resources deriving from herbaceous plants  
317 most likely available in the palaeoenvironment neighbouring the site and, in some cases, still present and used  
318 today (Guarino et al., 2008; Salerno and Guarrera, 2008).

319 Vetch seeds (*Vicia* spp., Fig. 4) are the most common remains, their statistically significant presence indicates  
320 a possible food use and not just a chance. In fact, vetch is a pulse found in other contemporary Mediterranean  
321 sites and linked to food (Asouti et al., 2018; Martínez Varea and Badal García, 2018; Martínez-Varea, 2020).  
322 The state of preservation of the seeds affected the visibility of the hilum, the key feature for species  
323 determination. Other less preserved finds have morphologies that can be confused with grass pea (*Lathyrus*  
324 sp.) or the pea (*Pisum* sp.) ([www.actaplantarm.org](http://www.actaplantarm.org)). Pulses are well represented in the Epigravettian trench at  
325 the bottom of the cave while they are missing in the samples coming from the atrium).

326 Undeterminable fragments of *Poaceae* grains have also been identified and ascribed to food species.  
327 Unfortunately, the lack of a specific determination does not allow other considerations and comparisons.  
328 Several genera are included among the *Poaceae* food species; however, caution is needed, due to the great  
329 diffusion in the environment of this botanical family (Pignatti 2017).

330 Amongst the findings, of note is the finding of *Galium* sp., *Arctium* sp., *Echium* sp. and *Lithospermum*  
331 *officinale* (Guarino et al., 2008). Given the low occurrence of these remains, their use is only suggested here.  
332 Besides their medical properties, these plants are very common in nature and provide data on the vegetation in  
333 the area adjacent the site (Fig. 4).

334  
335  
336  
337 **FIG.5 HERE**  
338  
339  
340  
341  
342



343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391

#### 4.1.2 Grotta del Romito

Grotta del Romito has provided the lowest number of finds considering the sample size, probably because of the incoherent sediment, which limited the preservation of plant remains (Fig.6). Analogously, many of the samples examined for pollen turned out to be sterile (Ricciardi 2004).

The sample referable to layer D3 (Square F8 II) returned 6 fragments of bloodtwig dogwood (*Cornus sanguinea* L.), almost certainly parts of a single stone. The interest of this find concerns the possible food and/or medical use of the fruits, as well as the extraction of oil and colouring pigments (Lanzara 1997).

Carpological fragments that could not be determined due to the poor state of conservation come from the 59 samples that undergone flotation during the 2018 field season (supplementary information3). Most of the finds identified are related to fruits of tree and shrub species: hazel (*Corylus avellana* L.), blackthorn (*Prunus* sp. L.), bramble (cf. *Rubus* gr. *ulmifolius*), elder (*Sambucus nigra* L., *Sambucus* sp. L.) and probably a pine seed (cf. *Pinus* sp. L.) (Fig.4). Many of these fruits are edible, even if they have not been found inside specific structures. Carpological findings related to herbaceous species such as dock (*Rumex* sp. L.) and a fragment of the same botanical family (*Polygonaceae*) were also identified. All the plants identified are present in the environment, and may have a dietary and/or medicinal use (Lanzara 1997).

The study of the 76 pollen samples is much more complex (Supplementary Information 4). The samples come from 2 different locations of the cave: the interior of the cave and a southern section (always in the internal trench) in which the samples are identified by their metric location (from -58 to -350).

The poorest samples of plant remain are those of the southern section. Only 2 out of 14 samples tested returned seeds or fruits. The chronology of the layers with plant remains (-80 and -137) refers to the Late Epigravettian; only three carpological remains were found: a seed and a fragment of cardoon (*Galium* sp. L.) and an indeterminable fragment.

Cardoon is a weed that grows in meadows and undergrowth (Pignatti 2017) and some species have medical properties ([www.actaplantarum.org](http://www.actaplantarum.org)). The presence of the cardoon may be due to intentionality but also to chance; in fact, the rough hairs that cover the seed make it adhere to humans and animals, favouring its unintentional transport to the anthropic areas.

62 samples come from the trench inside the cave and their study has allowed the discovery of 30 carpological fragments. Originally, these samples were taken for pollen analysis but, given the scarcity of results, after a preliminary search for pollen (Ricciardi, 2005), the samples were used for the recovery of macroremains.

A fair amount of plant finds (16 carpo-remains in 22 sediment samples) comes from the filling of the burial pit of the individual called "Romito 8". Discovered in 2002, this grave consists of a narrow and deep pit, filled with numerous stones placed over the body of the deceased. The study of the burial did not reveal any grave goods. The anthropological analysis revealed an impairment of the individual, or a paralysis that must have affected the left upper limb. The great dental wear proves that the individual must have compensated for the lack of use of the arm with the mouth, used as a real tool (Martini and Lo Vetro, 2018).

The plant macroremains found include 3 botanical families: *Polygonaceae*, *Poaceae* and *Fabaceae* (among which two clover seeds - *Trifolium* sp. are identifiable), to which we must add a fragment of the indeterminable fruit epidermis and 9 indeterminate carpological fragments (Fig. 4). The botanical families identified enclose numerous edible and medical genera, including clover (Pieroni et al., 2002). We cannot exclude that the presence of stones filling the pit burial might have favoured the preservation of plant materials. The Romito 7 and Romito 8 burials were subjected to pollen analysis (Ricciardi 2004), but no sporopollinic content was identified.

**FIG.6 HERE**

In addition to the finds found in the burial, other carpological remains were identified in unit D; it consists in over 35 levels (layers) dated to the Late Epigravettian (ca. 11000-13000 uncal BP, Martini 2012). A vetch

392 fragment (*Vicia* sp.) was found in layer D6, along with some indeterminable carpological fragments from  
393 layers D3, D4, D5 and D13.

394 A pit (pit 1) found in Layer D19 (dated around 12400 uncal BP), deserves a particular mention. It is an ovoid-  
395 shaped pit at the top of which there are some limestone blocks intentionally placed and emerging from the pit  
396 filling. A fragment of a webbed deer stag (*Cervus elaphus palmilodactyloceros*), other red deer remains, a  
397 skullcap, a cervical vertebra, a perforated atrophic canine, were found inside the pit, along with some  
398 perforated shells and a few other faunal fragments. All of this seems to outline an intentional deposition for  
399 ritual purposes (Martini et al., 2012b), but it does not contain important evidence of plant remains.

400 At Romito, the stratigraphy includes more ancient levels, in which other carpological remains have been found,  
401 such as the layer F1 (AMS date: 17376±90 uncal BP) related to the Early Epigravettian, and the layer H4  
402 (AMS date: 20210±245 uncal BP) which refers to the Evolved Gravettian. The presence of a mineralized fig  
403 stone (*Ficus carica* L.) in the layer F needs further discussion. The presence of the fig tree in this chronological  
404 horizon contrasts with the palaeoclimatic reconstructions of southern Italy starting from the Last Glacial  
405 Maximum (Incarbona et al., 2010) marked by a much lower temperature than today, which would not favour  
406 the presence of a thermophilic plant such as the fig tree. The pollen analysis of layer F identified specimens  
407 altered by erosive phenomena or collapses (Ricciardi 2004). The latter ones might have caused vertical  
408 movements within the stratigraphy, which would explain the presence of a single mineralized find (the fig) in  
409 a context of charred fragments. Alternatively, anthropogenic activity preceding the glacial peak, i.e. pits  
410 excavated in more ancient layers (e.g. H), could also be considered to explain the presence of the fig in layer  
411 F.

#### 412 413 4.1.3 Grotta del Cavallo

414 At Grotta del Cavallo, the archaeological materials are well placed within the relative stratigraphic structure,  
415 in some cases the lithic remains are well arranged horizontally, to testify the original position of the deposit  
416 (Martini, Pers. Comm. 2021). However, remains of cultivated plants (mainly cereals) have been identified  
417 throughout the deposit, even in the layers attributed to the Late Upper Palaeolithic (Fig. 4). The presence of  
418 homogeneous plant material throughout the stratigraphic column has been explained considering some  
419 peculiarities of the depositional sequence, which is characterised by small burrows of ground wasps. The lairs  
420 of these insects can reach a depth of 20 cm. Since plant material is very light, the activity of the insects led to  
421 the displacement of remains of cultivated species into the deeper layers.

422 The data obtained from the carpological study of Grotta del Cavallo provides a picture of the Apulian Neolithic,  
423 with fragments of grains and glume bases of barley (*Hordeum vulgare*) and wheat, emmer in particular  
424 (*Triticum turgidum* ssp. *dicoccon*) (Castelletti et al., 1987; Fiorentino et al., 2013) (Fig. 5,7).

#### 425 426 **FIG.7 HERE** 427

428 The spontaneous vegetation identified in various layers deserves further comments. While the fig (*Ficus*  
429 *carica*) is present only in the most superficial layers and the grapevine (mainly fragments of grape stones of  
430 *Vitis vinifera*) predominates in the intermediate ones (B1B-B1B/4), different herbaceous plants, including fat-  
431 hen (*Chenopodium* spp.), sedge (*Carex* spp.), common porcelain (*Portulaca oleracea*) and many fragments of  
432 *Poaceae* grains have been identified in the oldest layers yielded (Fig.4). These findings attest to a series of  
433 possible plant foods that could have been used during the late Upper Palaeolithic phases.

#### 434 435 436 4.2 Dental calculus analysis

437 Archaeological dental calculus preserved various vegetal and animal micro-debris (Table 5). Few and very  
438 small charcoal fragments and vegetal tissues were retrieved in the matrix. However, their dimensions,  
439 frequency and typology did not allow to formulate any identification or hypothesis about their presence in the  
440 matrix (Fig.2). Starch granules were recovered in the dental calculus of 5 individuals only (Romito 5, 7; San

441 Teodoro 3, 6; Oriente C). Overall, 4 morpho-types have been retrieved in this study (Fig. 8). In order to avoid  
442 misinterpretation granules smaller than 5 µm (transitory starches) were not considered (Haslam, 2004).

443 **Type I.** Type I granules were retrieved in 2 individuals (Romito 5 and Oriente C) (Table 4) in small quantities  
444 and some of them were found still partly embedded within the calculus remains (Fig. 8a,b,g,m,o,q,r). Size,  
445 shape, morphology and bimodal distribution that characterize granules of this type are encountered in the  
446 members of the plant tribe *Triticeae* (*Poaceae* family) and considered diagnostic features for taxonomic  
447 identification (Stoddard, 1999; Henry and Piperno, 2008; Yang and Perry, 2013). Starch distribution involves  
448 the presence of large granules (A-Type), round to sub-oval in 2D shape, ranging between 21.1 and 45.1 µm in  
449 maximum dimensions (mean size of 33.1 µm), lenticular 3D shape, a central hilum, a visible equatorial groove  
450 and deep lamellae concentrated in the mesial part; and small (< 10 µm) granules (B-Type) with round/sub-oval  
451 shapes and a central hilum (Stoddard, 1999; Geera et al., 2006; Yang and Perry, 2013). A-Type granules  
452 possess diagnostic features while smaller B-Type granules are rarely diagnostic to taxa (Yang and Perry, 2013).  
453 However, in our archaeological population, variability in the proportion and dimension of small B-type  
454 granules has been noticed. Based on literature (Henry et al., 2011; Yang and Perry, 2013) and our modern  
455 reference collection, this characteristic is common in the species of the genus *Aegilops* of *Triticeae* tribe (Fig.  
456 8s) and can be attributed to both environment (Blumenthal et al., 1994; 1995) and genetics (Stoddard and  
457 Sarker, 2000; Howard et al., 2011).

458 **Type II.** Starch granules attributed to this type are characterized by a polyhedral to sub-polyhedral 3D  
459 morphology, a central hilum, and fine cracks. They were recovered in 2 individuals (Romito 5 and San Teodoro  
460 6) (Fig. 8e,f,h). These features are consistent with starch granules of the tribe *Paniceae* of the grass family  
461 *Poaceae*, which is well known in ancient starch research (Madella et al., 2016). Dimensions of starch granules  
462 assigned to type II in our samples reach 21µm of maximum width, a size range found in several species of  
463 *Setaria* spp., *Panicum* spp., and *Echinochloa* spp. (Lucarini et al., 2016; Nava et al., 2020).

464 **Type III.** Starch granules attributed to this type were recovered in one individual (Romito 5) (Fig. 8c,d). They  
465 are characterized by round to sub-polyhedral 3D morphology, a central open hilum and a maximum width of  
466 20,8 µm. These features may be consistent with *Andropogoneae* tribe (Madella et al., 2016).

467 **Type IV.** Granules belonging to this morpho-type have been identified in the individual 3 of Grotta di San  
468 Teodoro. They are characterized by a sub-polyhedral 2D morphology, a central depression and marked cracks  
469 (Fig. 8i,j). The maximum recorded width is 16,8 µm. These features are consistent with starch granules of the  
470 family *Adoxaceae* and has tentatively been attributed to the genus *Sambucus* through comparison with our  
471 modern reference (Fig. 8k,l).

472  
473 Together with vegetal material also two barbules were recovered in the dental calculus of individual 5 from  
474 Grotta del Romito and individual 6 of San Teodoro. Unfortunately, microscopic characters which generally  
475 aid in the identification of groups of birds from their barbs were not visible in our remains.

476  
477 **Figure 2. Micro-debris in dental calculus:** a,b) Starch granules attributed to type I from individual 5 of Grotta  
478 del Romito; c, d) Starch granule attributed to type III from individual 5 of Grotta del Romito ; e) Starch granule  
479 attributed to type II from individual 5 of Grotta del Romito; f) Starch granule attributed to type II from  
480 individual 5 of Grotta del Romito; g) Starch granule attributed to type I; h) Starch granule attributed to type  
481 II from individual 6 of Grotta di San Teodoro; i, j) Starch granule attributed to type IV from individual 3 of  
482 Grotta di San Teodoro; k, l) modern experimental starch granule of *Sambucus nigra*; m) Starch granule  
483 attributed to type I; n) Starch granule attributed to type I from individual 7 of Grotta di San Teodoro; o) Starch  
484 granule attributed to type I from individual 3 of Grotta di San Teodoro; p) Starch granule attributed to type I  
485 trapped in dental calculus from individual 3 of Grotta di San Teodoro; q) Starch granule attributed to type I

486 from Oriente C; r) Starch granule attributed to type I from Oriente C; s) modern experimental starch granules  
487 of *Aegilops crassa*.

488

489

490

491

## 5. Discussion and conclusions

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

By integrating macro-botanical remains from the sites of Grotta della Serratura, Grotta del Romito, and, to a less extent, Grotta del Cavallo, and micro-botanical evidence preserved in the dental calculus of the individuals buried at Grotta di San Teodoro, Grotta d'Oriente, and Grotta del Romito, we show that hunter-gatherer communities that occupied these sites during the Late Glacial collected different plant resources, including seeds, grasses and forest fruits, and used them for nutritional or medical purposes.

Grotta della Serratura yielded most of the plant remains analysed in this study. The abundance of vetch seeds (*Vicia* spp.) at this site, like in other coastal sites of Greece (Franchti cave) and Spain (Cueva de les Cendres) occupied in the same period, reveals a specific interest in this protein-rich plant foods by hunter-gatherer communities of the Mediterranean area (Asouti et al., 2018; Martínez Varea and Badal García, 2018; Martínez-Varea, 2020). Fragments of *Poaceae* plants are also documented at this site although an attribution to a species level was not possible. Along with the food species, some plants recovered at Grotta della Serratura are also renowned for their healing properties, such as *Galium* sp., *Arctium* sp., *Echium* sp. and *Lithospermum officinale*. All these plants are common in the Mediterranean environment and their presence in the stratigraphy of the cave can testify to their possible use as medicinal remedies at the end of the Palaeolithic.

Our results are congruent with the palaeoenvironmental data available for Grotta della Serratura. In particular, the anthracological study emphasises the presence of oak forest surrounding the cave (Abbate Edlmann et al., 1997) characterised by the presence of oaks, pines and various broad-leaved trees (*Acer*, *Carpinus*, *Fraxinus*, *Sorbus*), and more specifically Mediterranean species (*Ilex*, *Laurus*). However, no remains of fruits from tree plants were recovered at this site.

In the Late Epigravettian layer 8C and 8G, pollen sequence supports the carpological examination and indicate a predominantly herbaceous environment with few forest *taxa*, possibly influenced by the proximity of the cave to a series of coastal dunes. The botanical family *Asteraceae*, also represented in the carpological remains, stands out amongst the herbaceous plants (Cattani, 1993).

Our data from Grotta del Romito testify to the deep interaction between Upper Palaeolithic communities and the surrounding environment, which reflects in the exploitation of various available plant *taxa*.

The greater distance of the site from the coast is reflected in the larger exploitation of fruits that are typical of the forest environment surrounding the cave: bloodtwig dogwood (*Cornus sanguinea* L.), hazel (*Corylus avellana* L.), blackthorn (*Prunus* sp. L.), bramble (cf. *Rubus* gr. *ulmifolius*), elder (*Sambucus nigra* L., *Sambucus* sp. L.) and probably a pine seed (cf. *Pinus* sp. L.). Analogous macro-remains were recovered in the Late Upper Palaeolithic sediments of Grotta di Pozzo in the Fucino region of Abruzzo (central Italy), where forest fruits were the only plants identified (Lubell et al., 1999). Amongst the macro-botanical remains recovered at Grotta del Romito fragments of herbaceous edible plants, such as dock (*Rumex* sp. L.) and fragments belonging to the same botanical family (*Polygonaceae*) are documented, together with the vetch (*Vicia* sp. L.) and fragments of *Poaceae* grains. Plants with medicinal properties, such as the cardoon (*Galium* sp. L.) were also recovered.

The palaeoclimatic record documented at Grotta del Romito for the Late Glacial highlights a predominantly pine tree-like environment, with a lower presence of herbaceous plants of the *Rosaceae*, *Boraginaceae*, *Plantaginaceae*, *Ranunculaceae*, *Fabaceae*, and *Poaceae* botanical families (Ricciardi 2004). Our analysis of the corresponding levels shows a similar distribution of these botanical families, while the presence of edible fruits in the upper stratigraphy indicates the spread of broad-leaved trees at the end of the Late Glacial.

535 Similarly, woodland mammals appear in the microfauna while the malacofauna suggests a more open  
536 environment (Martini et al., 2009b).

537

538 *Poaceae* remains were also identified in the burial filling of Romito 8, together with *Polygonaceae* and  
539 *Fabaceae* remains (including clover seeds) and a fragment of the epidermis of an indeterminable fruit. The  
540 recovery of *Poaceae* grasses at both Grotta del Romito and Grotta della Serratura suggest that Upper  
541 Palaeolithic communities of southern Italy knew and possibly consumed plants of this botanical family.  
542 Unfortunately, the numerous genera of *Poaceae* family widespread in Southern Italy and Sicily (Huntley et  
543 al., 1999; Sadori and Narcisi, 2001; Ermolli and di Pasquale, 2002; Pignatti, 2017) and the absence of specific  
544 diagnostic parts in the macro-remains recovered at the investigated sites prevented from a precise  
545 determination of the remains at the species level. Nonetheless, starch grains belonging to *Poaceae* family were  
546 recovered in dental calculus of two individuals buried at Grotta del Romito (Romito 5 and Romito 7) and  
547 attributed to *Triticeae* tribe, based on their morpho-dimensional features. Complementary evidence for a diet  
548 rich in starchy foods comes also from the anthropological analysis of dental remains, which identifies the  
549 presence of one interproximal caries at the neck of the first molar of the individual Romito 1 (Fabbri and  
550 Mallegni, 1988).

551 In the most recent phases (D3-D5), the carpological remains are more closely related to fruit of tree species,  
552 while in the earlier layers (D6-D7) herbaceous plants seem to predominate. Starch granules retrieved from  
553 dental calculus complement this picture by suggesting the consumption of grasses of *Triticeae* tribe during the  
554 Late Glacial period. This highlights an expansion in the plants used, under the pressure of an environment that  
555 is changing in a thermophilic way, with the increase in deciduous forest cover and the exploitation of edible  
556 fruits.

557

558 Also in Sicily, starch granules recovered in the dental calculus of the individuals 3 and 6 of Grotta di San  
559 Teodoro and Oriente C suggest grasses of *Poaceae* family, and more likely grasses of *Paniceae* and *Triticeae*  
560 tribes, were consumed since the end of the Oldest Dryas. *Poaceae* botanical family is very widespread and  
561 also includes many genera suitable for living in colder climatic conditions (e.g. *Avena*) and therefore their use  
562 can be suggested even for cold phases (Lippi et al., 2015). *Setaria* is one of the possible genera to which starch  
563 granules retrieved from San Teodoro individuals were assigned, and some species (e.g. *Setaria verticillata*)  
564 grow in Sicily at altitudes up to 1000 m above sea level (Pignatti, 2017). Based on regional  
565 palaeoenvironmental record (Sadori and Narcisi, 2001; Ermolli and di Pasquale, 2002), the presence of the  
566 herbaceous taxa is confirmed in southern Italy during the Late Glacial and at the onset of the warm Bølling-  
567 Allerød interstadial phase and linked to an increase of the moisture and temperatures. Grasses of *Paniceae*  
568 tribes prefer mild climates and became more widespread in proximity of water courses and in coastal zones  
569 during the Late Glacial. At San Teodoro, the high rate of precipitations would have guaranteed water to  
570 numerous streams and rivers, hence providing ideal environment for such species. The faunal assemblage at  
571 Grotta d'Oriente suggests the presence of both forest and open environments where species of the *Triticeae*  
572 tribe could have grown.

573 A Late Upper Palaeolithic familiarity with *Poaceae* taxa seems to emerge as a part of a well-documented  
574 dietary pattern also in other regional case studies that document the use of many genera of this botanical family  
575 as edible plants from the first part of the Late Glacial until the early phases of the Holocene.

576 In Italy, the consumption of plants of this botanical family was suggested based on the starch remains identified  
577 in the dental calculus of one individual buried at Riparo Tagliente (Oxilia et al 2020), the oldest known  
578 archaeological site on the southern slope of the Alps to be re-occupied by human groups after the Late Glacial  
579 Maximum. Similar evidence was also retrieved in Late Upper Palaeolithic individuals of Grotta Continenza,  
580 in central Italy (Nava et al., 2021). A diet, including meat as well as plant foods was also hypothesised on the  
581 basis of the dental microwear and trace element analysis for the individuals of San Teodoro (Carnieri and  
582 Mallegni, 2006). A caries identified during the dental calculus sampling on the neck of the upper left second

583 molar of the individual 6 from the same site would suggest the consumption of carbohydrates (Fig. 3d). In the  
584 same period, evidence from Grotte des Pigeons in Morocco show the high occurrence of caries in individuals  
585 dated between 15,000 and 13,700 cal BP and yielded evidence of systematic harvesting and processing of  
586 edible wild plants, including acorns and pine nuts (Humphrey et al., 2014).

587 The dental calculus of the individual 3 of San Teodoro also preserved evidence for the consumption of forest  
588 fruits. Based on the comparison with fruits, drupes and berries from our modern reference collection, ancient  
589 starch granules attributed in ST3 were possibly attributed to elderflower (*Sambucus* sp.). This forest fruit is  
590 rich in carbohydrate and proteins and renowned for peculiar bioactive properties (Sidor and Gramza-  
591 Michałowska, 2015). Interestingly, a broadening in the local resources exploited at San Teodoro was already  
592 suggested based on the isotopic study for the individual 7, which purportedly consumed a wide range of foods,  
593 including terrestrial and marine resources. The specific topography and hydrology of the area and the  
594 proximity of the site to the Nebrodi mountain chain, characterised by high precipitation rate, might have made  
595 plenty of plant resources available. Forest diffused in this area would have certainly provided a perfect habitat  
596 for elderflowers and herbaceous species, as documented also today (Martini et al., 2005; Colonese et al., 2018).

597 Complimentary evidence for assessing the role of plant foods in Late Glacial human diet comes from Grotta  
598 della Serratura and Grotta d'Oriente. Here, the presence of non-food shell debris suggests that aquatic plants  
599 growing in the coastal lagoons and estuaries close to the sites were collected and likely used as dietary  
600 supplement (Martini et al., 2005; Colonese et al., 2018). One of the most common species of seagrass meadows  
601 growing in the Tyrrhenian littoral and in Sicily, the *Zostera* spp., is rhizomatous herb which produces seeds  
602 known for being a good source of carbohydrate and proteins.

603 More complex to interpret are the remains from Grotta del Cavallo, due to their stratigraphic position.  
604 However, the oldest remains might indicate the exploitation of a series of herbaceous edible plants such as fat-  
605 hen (*Chenopodium*), common porcelain (*Portulaca oleracea*) and many fragments of *Poaceae* grains, which  
606 are typical in the Mediterranean environment.

607  
608 Based on the rich data discussed in this paper, we infer a broadening in the dietary base since the earliest phases  
609 of the deglaciation in the Southern region of Italy and Sicily. Herbaceous species were used since the earliest  
610 phases of the Late Glacial together with drupes and fruits from tree plants. While it is possible that the spread  
611 of the deciduous forest cover prompted humans to explore new plant resources in the deglaciation period, our  
612 results underline the interpretative limits of dietary reconstructions based on the application of single  
613 methodologies, e.g. stable isotope study or archaeozoological analysis, when attempting the reconstruction of  
614 dietary habits in Palaeolithic contexts.

615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629

630 **Author contributions**

631

632 EC., M.C. conceived the study and designed the experiments. M.C., performed the carpological analysis, A.Z.,  
633 E.F. and E.C. performed the laboratory work on dental calculus and modern reference; F.M., L.S. and D.L.V.  
634 provided the samples; E.C., F.M., L.S. and D.L.V. supervised the work; E.C., provided funding for the  
635 analyses. All authors contributed to writing the manuscript.

636

637 **Data Availability**

638

639 Not applicable

640

641 **Declaration of competing interest**

642

643 The authors declare that they have no known competing financial interests or personal relationships that could  
644 have appeared to influence the work reported in this paper

645

646 **Acknowledgments**

647

648 The study was funded by the European Research Council (ERC) as part of the Research and Innovation  
649 program of the European Community Horizon 2020 (HIDDEN FOODS no.639286).

650 We thank Dr Giuseppina Mutri for her help in sampling residues for carpological analysis during the 2016  
651 field season. We thank Dr. Federico Poggiali for his help in the flotation activity during the 2018 field season.

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740

## References

- Abbate Edlmann, M.L., Giachi, G., Mariotti, M., 1997. Indagine antracologica. Querciola. Insediamento campaniforme a Sesto Fiorentino. *Garlatti e Razzai, Montespertoli (Firenze)* 33–39.
- Asouti, E., Ntinou, M., Kabukcu, C., 2018. The impact of environmental change on Palaeolithic and Mesolithic plant use and the transition to agriculture at Franchthi Cave, Greece. *PLoS one* 13, e0207805.
- Blockley, S., Pellegrini, M., Colonese, A.C., Lo Vetro, D., Albert, P.G., Brauer, A., Di Giuseppe, Z., Evans, A., Harding, P., Lee-Thorp, J., Lincoln, P., Martini, F., Pollard, M., Smith, V., Donahue, R., 2018. Dating human occupation and adaptation in the southern European last glacial refuge: The chronostratigraphy of Grotta del Romito (Italy). *Quaternary science reviews* 184, 5–25.
- Blumenthal, C., Wrigley, C.W., Batey, I.L., Barlow, E.W.R., 1994. The Heat-Shock Response Relevant to Molecular and Structural Changes in Wheat Yield and Quality. *Functional plant biology: FPB* 21, 901–909.
- Blumenthal, C., Bekes, F., Gras, P.W., Barlow, E.W.R., Wrigley, C.W., 1995. Identification of wheat genotypes tolerant to the effects of heat stress on grain quality. *Cereal chemistry*.
- Bonfiglio, L., Esu, D., Mangano, G., Masini, F., Petruso, D., Soligo, M., Tuccimei, P., 2008. Late Pleistocene vertebrate-bearing deposits at San Teodoro Cave (North-Eastern Sicily): Preliminary data on faunal diversification and chronology. *Quaternary international: the journal of the International Union for Quaternary Research* 190, 26–37.
- Cappers, R.T.J., Neef, R., 2012. *Handbook of Plant Palaeoecology*. Barkhuis.
- Cappers, R.T.J., Bekker, R.M., Jans, J.E.A., 2012. *Digital seed atlas of the Netherlands*. Barkhuis.
- Carnieri, E., Mallegni, F., 2006. Le microusure dentarie di resti umani appartenenti a *Homo sapiens* del Paleolitico Superiore italiano. *La cultura del morire nelle società preistoriche e protostoriche italiane. Origines, Firenze* 293–303.
- Carra, M., Cristiani, E., 2020. I reperti carpologici. In: Sarti, L., Martini, F. (Eds.), *Il Musteriano Di Grotta Del Cavallo Nel Salento (Scavi 1986-2005). Culture e Ambienti, Millenni. Museo e Istituto Fiorentino di Preistoria “Paolo Graziosi,”* pp. 111–122.
- Carra, M., Marival, P., 2011. I bulbi di avena altissima (*Arrhenatherum elatius* var. *bulbosum*) da Riparo Dalmeri (TN): offerta votiva o cibo quotidiano? *Preistoria Alpina* 45, 147–157.
- Castelletti, L., Constantini, L., Tozzi, C., 1987. Considerazioni sull’economia e l’ambiente durante il Neolitico in Italia. *Istituto Italiano di Preistoria e Protostoria*.
- Catalano, G., Modi, A., Mangano, G., Sineo, L., Lari, M., Bonfiglio, L., 2020. A mitogenome sequence of an *Equus hydruntinus* specimen from Late Quaternary site of San Teodoro Cave (Sicily, Italy). *Quaternary science reviews* 236, 106280.
- Colonese, A.C., Zanchetta, G., Fallick, A.E., Martini, F., Manganelli, G., Drysdale, R.N., 2010. Stable isotope composition of *Helix ligata* (Müller, 1774) from Late Pleistocene–Holocene archaeological record from Grotta della Serratura (Southern Italy): Palaeoclimatic implications. *Global and planetary change* 71, 249–257.
- Colonese, A.C., Zanchetta, G., Drysdale, R.N., Fallick, A.E., Manganelli, G., Vetro, D.L., Martini, F., Di Giuseppe, Z., 2011. Stable isotope composition of Late Pleistocene–Holocene *Eobania vermiculata* (Müller, 1774)(Pulmonata, Stylommatophora) shells from the Central Mediterranean basin: Data from Grotta d’Oriente (Favignana, Sicily). *Quaternary international: the journal of the International Union for Quaternary Research* 244, 76–87.
- Colonese, A.C., Lo Vetro, D., Martini, F., 2014. Holocene coastal change and intertidal mollusc exploitation in the central Mediterranean: variations in shell size and morphology at Grotta d’Oriente (Sicily). *Archaeofauna* 23, 181–192.
- Colonese, A.C., Lo Vetro, D., Landini, W., Di Giuseppe, Z., Hausmann, N., Demarchi, B., Angelo, C. d’, Leng, M.J., Incarbona, A., Whitwood, A.C., Martini, F., 2018. Late Pleistocene–Holocene coastal adaptation in central Mediterranean: Snapshots from Grotta d’Oriente (NW Sicily). *Quaternary international: the journal of the International Union for Quaternary Research* 493, 114–126.
- Craig, O.E., Biazzo, M., Colonese, A.C., Di Giuseppe, Z., Martinez-Labarga, C., Lo Vetro, D., Lelli, R., Martini, F., Rickards, O., 2010. Stable isotope analysis of Late Upper Palaeolithic human and faunal remains from Grotta del Romito (Cosenza), Italy. *Journal of archaeological science* 37, 2504–2512.
- Cristiani, E., Radini, A., Edinborough, M., Borić, D., 2016. Dental calculus reveals Mesolithic foragers in the Balkans consumed domesticated plant foods. *Proceedings of the National Academy of Sciences* 113, 10298–10303.



- 741 De Silva, M., Pizziolo, G., Vetro, D.L., De Troia, V., Machetti, P., Ortisi, E.F., Martini, F., 2015. Ritual use  
742 of Romito Cave during the Late Upper Palaeolithic: an integrated approach for spatial reconstruction.  
743 CAA2015 685.
- 744 Drucker, D., Bocherens, H., 2004. Carbon and nitrogen stable isotopes as tracers of change in diet breadth  
745 during Middle and Upper Palaeolithic in Europe. *International journal of osteoarchaeology* 14, 162–  
746 177.
- 747 Ermolli, E.R., Pasquale, G. di, 2002. Vegetation dynamics of south-western Italy in the last 28 kyr inferred  
748 from pollen analysis of a Tyrrhenian Sea core. *Vegetation history and archaeobotany* 11, 211–220.
- 749 Fabbri, P.F., Lo Vetro, D., 2021. Le sepolture e i resti umani del Paleolitico superiore e del Mesolitico ritrovati  
750 in Sicilia. In: Martini, F., Sarti, L. (Eds.), *Una Preziosa Eredità. Scritti in Ricordo Di Arturo Palma Di*  
751 *Cesnola, Millenni. Museo e Istituto Fiorentino di Preistoria*, pp. 221–241.
- 752 Fabbri, P.F., Mallegni, F., 1988. Dental anthropology of the Upper Palaeolithic remains from Romito cave at  
753 Papisidero (Cosenza, Italy)/Anthropologie dentaire du Paléolithique supérieur, Grotte del Romito à  
754 Papisidero (Cosenza, Italie). *Bulletins et memoires de la Societe d'anthropologie de Paris* 5, 163–177.
- 755 Fabbri, P.F., Graziosi, P., Guerri, M., Mallegni, F., 1989. Les hommes des sépultures de la grotte du Romito à  
756 Papisidero (Cosenza, Italie). In: *Hominidae: Proceedings of the 2nd International Congress of Human*  
757 *Paleontology*, Jaca, Milan. pp. 487–494.
- 758 Fiorentino, G., Caldara, M., De Santis, V., D'Oronzo, C., Muntoni, I.M., Simone, O., Primavera, M., Radina,  
759 F., 2013. Climate changes and human–environment interactions in the Apulia region of southeastern  
760 Italy during the Neolithic period. *Holocene* 23, 1297–1316.
- 761 Francalacci, P., 1988. Comparison of archaeological, trace element, and stable isotope data from two Italian  
762 coastal sites. *Rivista di antropologia* 66, 239–250.
- 763 Garilli, V., Vita, G., La Parola, V., Pinto Vraca, M., Giarrusso, R., Rosina, P., Bonfiglio, L., Sineo, L., 2020.  
764 First evidence of Pleistocene ochre production from bacteriogenic iron oxides. A case study of the  
765 Upper Palaeolithic site at the San Teodoro Cave (Sicily, Italy). *Journal of archaeological science* 123,  
766 105221.
- 767 Gazzoni, V., Goude, G., Herrscher, E., Guerreschi, A., Antonioli, F., Fontana, F., 2013. Late Upper  
768 Palaeolithic human diet: first stable isotope evidence from Riparo Tagliente (Verona, Italy). *Bulletins*  
769 *et memoires de la Societe d'anthropologie de Paris* 25, 103–117.
- 770 Geera, B.P., Nelson, J.E., Souza, E., Huber, K.C., 2006. Composition and properties of A- and B-type starch  
771 granules of wild-type, partial waxy, and waxy soft wheat. *Cereal chemistry* 83, 551–557.
- 772 Ghinassi, M., Colonese, A.C., Giuseppe, Z.D., Govoni, L., Vetro, D.L., Malavasi, G., Martini, F., Ricciardi,  
773 S., Sala, B., 2009. The Late Pleistocene clastic deposits in the Romito Cave, southern Italy: a proxy  
774 record of environmental changes and human presence. *Journal of Quaternary Science* 24, 383–398.
- 775 Goude, G., Willmes, M., Wood, R., Courtaud, P., Leandri, F., Cesari, J., Grün, R., 2017. New insights into  
776 Mesolithic human diet in the Mediterranean from stable isotope analysis: The sites of campu stefanu  
777 and Torre d'Aquila, Corsica. *International journal of osteoarchaeology* 27, 707–714.
- 778 Graziosi, P., 1962. Découverte de gravures rupestres de type paléolithique dans l'abri del Romito (Italie).  
779 *L'Anthropologie* 66, 262–268.
- 780 Graziosi, P., 1966. Riparo del Romito (Papisidero, prov. di Cosenza). *Rivista di Scienze Preistoriche,*  
781 *Notiziario* 21, 422.
- 782 Guarino, C., De Simone, L., Santoro, S., 2008. Ethnobotanical study of the Sannio area, Campania, southern  
783 Italy.
- 784 Haslam, M., 2004. The decomposition of starch grains in soils: implications for archaeological residue  
785 analyses. *Journal of archaeological science* 31, 1715–1734.
- 786 Henry, A.G., Piperno, D.R., 2008. Using plant microfossils from dental calculus to recover human diet: a case  
787 study from Tell al-Raqā'i, Syria. *Journal of archaeological science* 35, 1943–1950.
- 788 Henry, A.G., Brooks, A.S., Piperno, D.R., 2011. Microfossils in calculus demonstrate consumption of plants  
789 and cooked foods in Neanderthal diets (Shanidar III, Iraq; Spy I and II, Belgium). *Proceedings of the*  
790 *National Academy of Sciences of the United States of America* 108, 486–491.
- 791 Howard, T., Rejab, N.A., Griffiths, S., Leigh, F., Leverington-Waite, M., Simmonds, J., Uauy, C., Trafford,  
792 K., 2011. Identification of a major QTL controlling the content of B-type starch granules in *Aegilops*.  
793 *Journal of experimental botany* 62, 2217–2228.
- 794 Humphrey, L.T., De Groote, I., Morales, J., Barton, N., Collcutt, S., Bronk Ramsey, C., Bouzouggar, A., 2014.  
795 Earliest evidence for caries and exploitation of starchy plant foods in Pleistocene hunter-gatherers

- 796 from Morocco. *Proceedings of the National Academy of Sciences of the United States of America*  
797 111, 954–959.
- 798 Huntley, B., Watts, W.A., Allen, J.R.M., Zolitschka, B., 1999. Palaeoclimate, chronology and vegetation  
799 history of the Weichselian Lateglacial: comparative analysis of data from three cores at Lago Grande  
800 di Monticchio, southern Italy. *Quaternary science reviews* 18, 945–960.
- 801 Incarbona, A., Agate, M., Arisco, G., Bonomo, S., Buccheri, G., Di Patti, C., Di Stefano, E., Greco, A.,  
802 Madonia, G., Masini, F., Others, 2010. Ambiente e clima della Sicilia durante gli ultimi 20 mila anni.  
803 *Il Quaternario, Italian Journal of Quaternary Sciences* 23, 21–36.
- 804 Lightfoot, E., Boneva, B., Miracle, P.T., Šlaus, M., O’Connell, T.C., 2011. Exploring the Mesolithic and  
805 Neolithic transition in Croatia through isotopic investigations. *Antiquity* 85, 73–86.
- 806 Lippi, M.M., Foggi, B., Aranguren, B., Ronchitelli, A., Revedin, A., 2015. Multistep food plant processing at  
807 Grotta Paglicci (Southern Italy) around 32,600 cal BP. *Proceedings of the national Academy of*  
808 *Sciences* 112, 12075–12080.
- 809 Lo Vetro, D., Martini, F., 2006. La nuova sepoltura epigravettiana di Grotta d’Oriente (Favignana, Trapani).  
810 *La cultura del morire nelle società preistoriche italiane: studio interdisciplinare dei dati e loro*  
811 *trattamento informatico*. Florence: Origines 58–66.
- 812 Lubell, D., Mussi, M., Musacchio, A., Agostini, S., Coubray, S., Giraudi, C., 1999. Exploitation of seasonal  
813 resources in the mountains of Abruzzo (central Italy): Epigravettian to Neolithic. *L’Europe des*  
814 *derniers Chass. Epipaléolithique Mésolithique* 5, 465–472.
- 815 Lucarini, G., Radini, A., Barton, H., Barker, G., 2016. The exploitation of wild plants in Neolithic North Africa.  
816 Use-wear and residue analysis on non-knapped stone tools from the Haua Fteah cave, Cyrenaica,  
817 Libya. *Quaternary International* 410, 77–92.
- 818 Lugli, F., Cipriani, A., Capecchi, G., Ricci, S., Boschin, F., Boscato, P., Iacumin, P., Badino, F., Mannino,  
819 M.A., Talamo, S., Richards, M.P., Benazzi, S., Ronchitelli, A., 2019. Strontium and stable isotope  
820 evidence of human mobility strategies across the Last Glacial Maximum in southern Italy. *Nature*  
821 *ecology & evolution* 3, 905–911.
- 822 Madella, M., Lancelotti, C., García-Granero, J.J., 2016. Millet microremains—an alternative approach to  
823 understand cultivation and use of critical crops in Prehistory. *Archaeological and anthropological*  
824 *sciences* 8, 17–28.
- 825 Mallegni, F., Fabbri, P.F., 1995. The human skeletal remains from the Upper Palaeolithic burials found in  
826 Romito cave (Papasidero, Cosenza, Italy). *Bulletins et memoires de la Societe d’anthropologie de Paris*  
827 7, 99–137.
- 828 Mannino, M.A., Thomas, K.D., 2009. 22. Current research on prehistoric human coastal ecology: Late  
829 Pleistocene and Early Holocene hunter-gatherer transitions in north-west Sicily.
- 830 Mannino, M.A., Thomas, K.D., Leng, M.J., Di Salvo, R., Richards, M.P., 2011a. Stuck to the shore?  
831 Investigating prehistoric hunter-gatherer subsistence, mobility and territoriality in a Mediterranean  
832 coastal landscape through isotope analyses on marine mollusc shell carbonates and human bone  
833 collagen. *Quaternary international: the journal of the International Union for Quaternary Research*  
834 244, 88–104.
- 835 Mannino, M.A., Di Salvo, R., Schimmenti, V., Di Patti, C., Incarbona, A., Sineo, L., Richards, M.P., 2011b.  
836 Upper Palaeolithic hunter-gatherer subsistence in Mediterranean coastal environments: an isotopic  
837 study of the diets of the earliest directly-dated humans from Sicily. *Journal of archaeological science*  
838 38, 3094–3100.
- 839 Mannino, M.A., Catalano, G., Talamo, S., Mannino, G., Di Salvo, R., Schimmenti, V., Lalueza-Fox, C.,  
840 Messina, A., Petruso, D., Caramelli, D., Richards, M.P., Sineo, L., 2012. Origin and diet of the  
841 prehistoric hunter-gatherers on the mediterranean island of Favignana (Ègadi Islands, Sicily). *PloS*  
842 *one* 7, e49802.
- 843 Mannino, M.A., Lightfoot, E., Stevens, R., 2016. Novel isotopic approaches to investigating human  
844 palaeoecology: An introduction. *Environmental Archaeology* 21, 193–198.
- 845 Martínez Varea, C.M., Badal García, E., 2018. Plant use at the end of the Upper Palaeolithic: archaeobotanical  
846 remains from Cova de les Cendres (Teulada-Moraira, Alicante, Spain). *Vegetation history and*  
847 *archaeobotany* 27, 3–14.
- 848 Martínez-Varea, C.M., 2020. Gathering in the dunes. Seeds and fruits from the Gravettian levels of Cova de  
849 les Cendres (Teulada-Moraira, Alicante, Spain). *Journal of Archaeological Science: Reports* 33,  
850 102540.

- 851 Martini, F., 1993. Grotta della Serratura a Marina di Camerota. Culture e ambienti dei complessi olocenici.  
852 Firenze, Italy.
- 853 Martini, F., 2002. Grotta del Romito. Guide del Museo e Istituto Fiorentino di Preistoria. Museo e Istituto  
854 Fiorentino di Preistoria, Firenze.
- 855 Martini, F., 2006. Le evidenze funerarie nella grotta e nel Riparo del Romito (Papasidero, Cosenza). La cultura  
856 del morire nelle società preistoriche e protostoriche italiane. Studio interdisciplinare dei dati e loro  
857 trattamento informatico, Origines 1.
- 858 Martini, F., Lo Vetro, D., 2005. Il passaggio Gravettiano-Epigravettiano a Grotta del Romito (scavi 2003-  
859 2004): prime osservazioni. Il passaggio Gravettiano-Epigravettiano a Grotta del Romito (scavi 2003-  
860 2004): prime osservazioni 151–176.
- 861 Martini, F., Lo Vetro, D., 2011. Grotta del Romito a Papasidero: uomo, ambiente e cultura nel Paleolitico della  
862 Calabria: ricerche 1961-2011, Guide del Museo e Istituto fiorentino di Preistoria, Firenze. Editoriale  
863 Progetto 2000, Cosenza.
- 864 Martini, F., Lo Vetro, D., 2018. Grotta del Romito a Papasidero: una storia calabrese da 24.000 anni fa / Romito  
865 Cave in Papasidero: A Calabrian history from 24.000 years ago. Ente Parco Nazionale del Pollino.
- 866 Martini, F., Sarti, L., 2017. Nuove ricerche nei livelli romanelliani di Grotta del Cavallo (Lecce): le produzioni  
867 litiche e le figurazioni mobiliari. Nuove ricerche nei livelli romanelliani di Grotta del Cavallo (Lecce):  
868 le produzioni litiche e le figurazioni mobiliari 87–93.
- 869 Martini, F., Sarti, L., 2021. Grotta del Cavallo con e dopo Palma di Cesnola. In: Martini, F., Sarti, L. (Eds.),  
870 Una Preziosa Eredità. Scritti in Ricordo Di A. Palma Di Cesnola, Studi Di Archeologia Preistorica.  
871 Millenni, pp. 97–118.
- 872 Martini, F., Beccaro, P., Ghinassi, M., Martino, G., 2003. Le materie prime delle industrie litiche del Paleolitico  
873 superiore e del Mesolitico di Grotta della Serratura (Marina di Camerota): Prima caratterizzazione  
874 degli areali e delle modalità di raccolta. Rivista di scienze preistoriche 163–191.
- 875 Martini, F., Cattani, L., Colamussi, V., Colonese, A., Martino, G., Mallegni, F., Noto, F., Ricciardi, S.,  
876 Rickards, O., Rolle, R., 2004. Primi risultati delle nuove ricerche nei livelli epigravettiani di Grotta  
877 del Romito a Papasidero (scavi 2000-2002). Atti della 37, 85–100.
- 878 Martini, F., Colonese, A.C., Wilkens, B., 2005. Grotta della Serratura (Marina di Camerota-Salerno) La  
879 malacofauna dei livelli gravettiani ed epigravettiani. Considerazioni paleoecologiche. In: Atti Della  
880 IV Convegno Nazionale Di Archeozoologia. portalrecherche.uab.cat, pp. 87–96.
- 881 Martini, F., Beccaro, P., Ghinassi, M., Martino, G., 2006. Caratterizzazione degli areali e modalità di raccolta  
882 della materia prima litica nel Paleolitico superiore e nel Mesolitico in area calabro-campana: i casi  
883 studio di Grotta del Romito e di Grotta della Serratura. Caratterizzazione degli areali e modalità di  
884 raccolta della materia prima litica nel Paleolitico superiore e nel Mesolitico in area calabro-campana:  
885 i casi studio di Grotta del Romito e di Grotta della Serratura 241–252.
- 886 Martini, F., Vetro, D.L., Colonese, A.C., De Curtis, O., Di Giuseppe, Z., Locatelli, E., Sala, B., 2007.  
887 L'Epigravettiano finale in Sicilia. In: L'Italia Tra 15.000 e 10.000 Anni Fa. Cosmopolitismo e  
888 Regionalita Nel Tardoglaciale. portalrecherche.uab.cat, pp. 209–254.
- 889 Martini, F., Colonese, A.C., Di Giuseppe, Z., Lo Vetro D., G.M., Ricciardi, S., 2009a. Human-environment  
890 relationships during Late Glacial-Early Holocene transition some examples from Campania, Calabria  
891 and Sicily. Mediterr. Rev. Geogr. Pays Mediterr./J. Mediterr. Geogr. 112, 89–94.
- 892 Martini, F., Colonese, A.C., Giuseppe, Z. di, Ghinassi, M., Vetro, D.L., Ricciardi, S., 2009b. Human-  
893 environment relationships during the Late Glacial-Early Holocene transition: some examples from  
894 Campania, Calabria and Sicily. Mediterranee. Revue géographique des pays méditerranéens/Journal  
895 of Mediterranean geography 89–94.
- 896 Martini, F., Lo Vetro, D., Colonese, A.C., Cilli, C., De Curtis, O., 2012a. Primi risultati sulle nuove ricerche  
897 stratigrafiche a Grotta d'Oriente (Favignana, Trapani), Scavi 2005. Primi risultati sulle nuove ricerche  
898 stratigrafiche a Grotta d'Oriente (Favignana, Trapani), Scavi 2005 319–331.
- 899 Martini, F., Lo Vetro, D., Di Giuseppe, Z., Baglioni, L., Colonese, A.C., 2012b. Strutture e sottostrutture del  
900 Paleolitico superiore di Grotta del Romito tra funzionalità e simbolismo. Strutture e sottostrutture del  
901 Paleolitico superiore di Grotta del Romito tra funzionalità e simbolismo 33–66.
- 902 Naito, Y.I., Chikaraishi, Y., Ohkouchi, N., Drucker, D.G., Bocherens, H., 2013. Nitrogen isotopic composition  
903 of collagen amino acids as an indicator of aquatic resource consumption: insights from Mesolithic and  
904 Epipalaeolithic archaeological sites in France. World archaeology 45, 338–359.
- 905 Nava, A., Fiorin, E., Zupancich, A., Carra, M., Ottoni, C., Di Carlo, G., Voza, I., Brugnoletti, O., Alhague,  
906 F., Cremonesi, R.G., Coppa, A., Bondioli, L., Borić, D., Cristiani, E., 2021. Multipronged dental

- 907 analyses reveal dietary differences in last foragers and first farmers at Grotta Continenza, central Italy  
908 (15,500–7000 BP). *Scientific reports* 11, 4261.
- 909 Orombelli, G., Ravazzi, C., 1996. The Late Glacial and early Holocene: chronology and paleoclimate. II  
910 *Quaternario* 9, 439–444.
- 911 Oxilia, G., Bortolini, E., Badino, F., Bernardini, F., Gazzoni, V., Lugli, F., Romandini, M., Radini, A., Terlato,  
912 G., Marciani, G., Silvestrini, S., Menghi Sartorio, J.C., Thun Hohenstein, U., Fiorenza, L., Kullmer,  
913 O., Tuniz, C., Moggi Cecchi, J., Talamo, S., Fontana, F., Peresani, M., Benazzi, S., Cristiani, E., 2020.  
914 Exploring late Paleolithic and Mesolithic diet in the Eastern Alpine region of Italy through multiple  
915 proxies. *American journal of physical anthropology* n/a. doi:10.1002/ajpa.24128
- 916 Paine, C., O’Connell, T., Miracle, P.T., McCartan, S., Schulting, R., Warren, G., Woodman, P., 2009. Stable  
917 isotopic reconstruction of Early Mesolithic diet at Pupicina Cave. *Mesolithic Horizons*. Oxford:  
918 Oxbow Books 210–216.
- 919 Pieroni, A., Nebel, S., Quave, C., Münz, H., Heinrich, M., 2002. Ethnopharmacology of liakra: traditional  
920 weedy vegetables of the Arbëreshë of the Vulture area in southern Italy. *Journal of ethnopharmacology*  
921 81, 165–185.
- 922 Pignatti, S., 2017. *Flora d’Italia*, 2nd ed. Edagricole, Bologna, Italy.
- 923 Piperno, D.R., Weiss, E., Holst, I., Nadel, D., 2004. Processing of wild cereal grains in the Upper Palaeolithic  
924 revealed by starch grain analysis. *Nature* 430, 670–673.
- 925 Pryor, A.J.E., Steele, M., Jones, M.K., Svoboda, J., Beresford-Jones, D.G., 2013. Plant foods in the Upper  
926 Palaeolithic at Dolní Věstonice? *Parenchyma redux*. *Antiquity* 87, 971–984.
- 927 Radini, A., Nikita, E., Buckley, S., Copeland, L., Hardy, K., 2017. Beyond food: The multiple pathways for  
928 inclusion of materials into ancient dental calculus. *American journal of physical anthropology* 162,  
929 71–83.
- 930 Revedin, A., Aranguren, B., Becattini, R., Longo, L., Marconi, E., Lippi, M.M., Skakun, N., Sinitsyn, A.,  
931 Spiridonova, E., Svoboda, J., 2010. Thirty thousand-year-old evidence of plant food processing.  
932 *Proceedings of the National Academy of Sciences of the United States of America* 107, 18815–18819.
- 933 Revedin, A., Longo, L., Mariotti Lippi, M., Marconi, E., Ronchitelli, A., Svoboda, J., Anichini, E., Gennai,  
934 M., Aranguren, B., 2015. New technologies for plant food processing in the Gravettian. *Quaternary*  
935 *international: the journal of the International Union for Quaternary Research* 359–360, 77–88.
- 936 Ricciardi, S., 2005. Le fluttuazioni climatico-ambientali dell’ultimo ciclo glaciale nell’Italia meridionale:  
937 archeopalinologia delle sequenze paleolitiche di Grotta del Cavallo, Grotta del Romito e Grotta della  
938 Serratura (Preistoria: Ambiente e Culture). Università degli Studi di Siena.
- 939 Romagnoli, F., 2015. A second life: Recycling production waste during the Middle Palaeolithic in layer L at  
940 Grotta del Cavallo (Lecce, Southeast Italy). *Quaternary international: the journal of the International*  
941 *Union for Quaternary Research* 361, 200–211.
- 942 Sabin, S., Fellow Yates, J., 2020. Dental Calculus Field-Sampling Protocol (Sabin version) v2  
943 (protocols.io.bqecmtaw). protocols.io. doi:10.17504/protocols.io.bqecmtaw
- 944 Sadori, L., Narcisi, B., 2001. The Postglacial record of environmental history from Lago di Pergusa, Sicily.  
945 *Holocene* 11, 655–671.
- 946 Salerno, G., Guarrera, P.M., 2008. Ricerche etnobotaniche nel Parco Nazionale del Cilento e Vallo di Diano:  
947 il territorio di Castel San Lorenzo (Campania, Salerno). *Informatore botanico italiano* 40, 165–181.
- 948 Seguinot, J., Ivy-Ochs, S., Juvet, G., Huss, M., Funk, M., Preusser, F., 2018. Modelling last glacial cycle ice  
949 dynamics in the Alps. *The Cryosphere* 12, 3265–3285.
- 950 Sidor, A., Gramza-Michałowska, A., 2015. Advanced research on the antioxidant and health benefit of  
951 elderberry (*Sambucus nigra*) in food – a review. *Journal of functional foods* 18, 941–958.
- 952 Stoddard, F.L., 1999. Survey of starch particle-size distribution in wheat and related species. *Cereal chemistry*  
953 76, 145–149.
- 954 Stoddard, F.L., Sarker, R., 2000. Characterization of Starch in *Aegilops* Species. *Cereal Chemistry Journal* 77,  
955 445–447.
- 956 Vacca, B.B., 2012. The hunting of large mammals in the upper palaeolithic of southern Italy: A diachronic  
957 case study from Grotta del Romito. *Quaternary international: the journal of the International Union*  
958 *for Quaternary Research* 252, 155–164.
- 959 Vadillo Conesa, M., Ricci, G., Bencomo Viala, M., Aura Tortosa, J.E., Martini, F., 2021. A review of  
960 splintered pieces from two lithic assemblages in Mediterranean Europe: Grotta Della serratura (Marina  
961 di camerota, Italy) and Cueva de nerja (Málaga, Spain). *Lithic technology* 1–16.

962 Vigliardi, A., 1968. L'industria litica della grotta di S. Teodoro, in provincia di Messina (scavi Graziosi -  
963 Maviglia). *Rivista di scienze preistoriche* 33–144.

964 Vigne, J.-D., 2004. Accumulation de lagomorphes et de rongeurs dans le sites mésolithiques corso-sardes:  
965 origines taphonomiques implications anthropologiques. In: Brugal, J.P., Desse, J. (Eds.), *Petits*  
966 *Animaux et Sociétés Humaines: Du Complément Alimentaire Aux Ressources Utilitaires*. Presented  
967 at the XXIVe Rencontres Internationales D'archéologie et D'histoire d'Antibes, pp. 261–281.

968 Watts, W.A., Allen, J.R.M., Huntley, B., Fritz, S.C., 1996. Vegetation history and climate of the last 15,000  
969 years at Laghi di Monticchio, southern Italy. *Quaternary science reviews* 15, 113–132.

970 Weiss, E., Wetterstrom, W., Nadel, D., Bar-Yosef, O., 2004. The broad spectrum revisited: evidence from  
971 plant remains. *Proceedings of the National Academy of Sciences of the United States of America* 101,  
972 9551–9555.

973 Weiss, E., Kislev, M.E., Simchoni, O., Nadel, D., Tschauner, H., 2008. Plant-food preparation area on an  
974 Upper Paleolithic brush hut floor at Ohalo II, Israel. *Journal of archaeological science* 35, 2400–2414.

975 Yang, X., Perry, L., 2013. Identification of ancient starch grains from the tribe Triticeae in the North China  
976 Plain. *Journal of archaeological science* 40, 3170–3177.

977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017

1018 **Captions**

1019  
1020 **Figure 1.** The location of the five analysed contexts from southern Italy and Sicily

1021  
1022 **Figure 2.** Sample of experimental carpological remains from modern reference collection. a)  
1023 *Hordeum vulgare*, glume base; b) *Triticum turgidum* ssp. *dicoccon*, glume base; c) *Vicia sylvatica*, seed; d)  
1024 *Trifolium campestre*, seed.; e) *Prunus spinosa*, stone; f) *Sambucus nigra*, stone; g) *Quercus pubescens*,  
1025 *cicatrix*; h) *Corylus avellana*, stone fragment; i) *Rumex crispus*, fruit; j) *Chenopodium album*, seed; k) *Echium*  
1026 *vulgare*, seed; l) *Ficus carica*, stones

1027  
1028 **Figure 3.** Some of the teeth sampled in this study. a) San Teodoro 5; b, c) Romito 3; d) Sant  
1029 Teodoro 6

1030  
1031 **Figure 4.** Carpological remains found at Grotta del Cavallo (a-h), Grotta del Romito (i-p) and Grotta della  
1032 Serratura (q-x). a) *Ficus carica*; b) *Triticum* sp., glume bases; c) *Vitis vinifera* fragments; d) *Triticum turgidum*  
1033 ssp. *dicoccon*; e) *Carex* sp.; f) *Quercus* sp.; g) *Chenopodium gr. album*; h) *Hordeum vulgare*, glume bases; i)  
1034 *Sambucus nigra* sp.; j) *Galium* sp. ; k) *Corylus avellana* fragment; l) *Rumex* sp.; m) *Rubus gr. ulmifolius*; n)  
1035 *Prunus* sp; o) *Sambucus* sp.; p) *Trifolium* sp. ; q) *Lithospermum officinale* (side a); r) *Lithospermum officinale*  
1036 (side b); s) *Echium* sp. (ventral view); t) *Arctium* sp.; u) *Vicia/Lathyrus*; v) *Vicia* sp.; w) *Vicia* sp., seed  
1037 fragments; x) *Vicia* sp., seeds

1038  
1039 **Figure 5.** Occurrence of carpological findings in each of the analysed sites.

1040  
1041 **Figure 6.** Distribution of tree/shrub, weeds, and indeterminable findings within the analysed layers at Grotta  
1042 del Romito.

1043  
1044 **Figure 7.** Distribution of domestic species, wild species, and indeterminable findings within the analysed  
1045 layers of Grotta del Cavallo.

1046  
1047 **Figure 8. Micro-debris in dental calculus:** a,b) Starch granules attributed to type I from individual 5 of Grotta  
1048 del Romito; c, d) Starch granule attributed to type III from individual 5 of Grotta del Romito ; e) Starch granule  
1049 attributed to type II from individual 5 of Grotta del Romito; f) Starch granule attributed to type II from  
1050 individual 5 of Grotta del Romito; g) Starch granule attributed to type I; h) Starch granule attributed to type  
1051 II from individual 6 of Grotta di San Teodoro; i, j) Starch granule attributed to type IV from individual 3 of  
1052 Grotta di San Teodoro; k, l) modern experimental starch granule of *Sambucus nigra*; m) Starch granule  
1053 attributed to type I; n) Starch granule attributed to type I from individual 7 of Grotta di San Teodoro; o) Starch  
1054 granule attributed to type I from individual 3 of Grotta di San Teodoro; p) Starch granule attributed to type I  
1055 trapped in dental calculus from individual 3 of Grotta di San Teodoro; q) Starch granule attributed to type I  
1056 from Oriente C; r) Starch granule attributed to type I from Oriente C; s) modern experimental starch granules  
1057 of *Aegilops crassa*.

1058  
1059 **Table 1.** Radiocarbon dates available from the Late Palaeolithic sites analyzed in this paper.

1060  
1061 **Table 2.** Finding numbers, stratigraphic provenance, chronology, and relative frequencies.

1062  
1063 **Table 3.** Radiocarbon dates available from the Late Palaeolithic burials analysed in this paper. The dates were  
1064 calibrated with Oxcal 4.4 using the IntCal20 calibration curve in the burial (Catalano et al 2020; Lo Vetro and  
1065 Martini 2012).

1066  
1067 **Table 4.** Details of dental calculus sampled for the study (n=9).

1068

Figure 1

























Site	Lab label	Cultural attribution	AMS dating (IntCal 20)			Material	
			Lab Code	<sup>14</sup> C age (BP)	95.4% confidence, cal BP		
Oriente C	7E	Late Palaeolithic	LTL873A	12132 ± 80	14235-13796	Charcoal	Lo Vetro and Martini, 2006
	Sublayer 7D		LTL14260A	12149 ± 65	14217-13803	Charcoal	Lo Vetro and Martini, 2006
Serratura	Unit 4	Undif. Epipaleolithic	UtC - 750	10000±200	12469-10875	Charcoal	Martini et al. 2005
	Unit 5		UtC - 751	9790±170	11814-10693	Charcoal	Martini et al. 2005
	Unit 5		Bln - 3568	9720±60	11248-10800	Charcoal	Martini et al. 2005
	Unit 6	Sauveterrian	Bln - 3569	9620±60	11185-10760	Charcoal	Martini et al. 2005
	Unit 6		UtC - 752	9770±140	11690-10715	Charcoal	Martini et al., 2005
	Unit 7		Bln - 3570	9870±70	11612-11177	Charcoal	Martini et al., 1993
	Unit 7		UtC - 753	10230±130	12578 -11398	Charcoal	Martini et al., 2005
	Unit 8A		UtC - 754	10000±130	11941 -11198	Charcoal	Martini et al., 2005
	Unit 8B		UtC - 755	10270±140	12610-11403	Charcoal	Martini et al., 2005
			Bln - 3571	10220±60	12441-11642	Charcoal	Martini et al., 2005
	Unit 8C	UtC - 1418	11290±90	13405-13067	Charcoal	Martini et al., 2005	
	Unit 8E	UtC - 1420	11490±160	13743 -13095	Charcoal	Martini et al., 2005	
	Unit 8F	UtC - 1463	11460±80	13476-13175	Charcoal	Martini et al., 2005	
	Unit 8FI	Beta - 63289	11240±90	13314-12929	Charcoal	Martini et al., 2005	
	Unit 8FII	Beta - 63290	12100±100	14311-13762	Charcoal	Martini et al., 2005	
	Unit 8FIII	Beta - 63291	11880±120	14037-13503	Charcoal	Martini et al., 2005	
	Unit 8G	Beta - 63292	12060±90	14165-13756	Charcoal	Martini et al., 2005	
	Unit 9	Beta - 63293	13100±120	16055-15315	Charcoal	Martini et al., 2005	
	Unit 10A	Beta - 63294	15350±200	18997-18240	Charcoal	Martini et al., 2005	
	Unit 10C	UtC - 1421	15700±110	19228-18776	Charcoal	Martini et al., 2005	
Unit 11	Beta - 88907	24380±1530	32889-25927	Charcoal	Martini et al., 2005		
Unit 12	Beta - 88908	29020±2650	44010-28708	Charcoal	Martini et al., 2005		
Romito	D1	Late Epigravettian	Beta-160,300	11660 ± 70	13742-13345	Charcoal	Blockley et al., 2018

D1	Late Epigravettian	ROM 5-1	11765±50	13760-13502	Charcoal	Blockley et al., 2018
D1	Late Epigravettian	ROM 5-2b	12415±50	14907-14246	Charcoal	Blockley et al., 2018
D5a	Late Epigravettian	Beta-160,302	12060±90	14165-13756	Charcoal	Blockley et al., 2018
D5b	Late Epigravettian	Beta-160,303	12160±50	14297-13813	Charcoal	Blockley et al., 2018
D8	Late Epigravettian	LTL234A	12170 ± 60	14308-13813	Charcoal	Blockley et al., 2018
D11	Late Epigravettian	LTL238A	12334±75	14844-14080	Charcoal	Blockley et al., 2018
D13	Late Epigravettian	LTL607A	12258±75	14821-14026	Charcoal	Blockley et al., 2018
D14	Late Epigravettian	LTL603A	12377±95	14933-14103	Charcoal	Blockley et al., 2018
D15	Late Epigravettian	LTL608A	12331±55	14829-14091	Charcoal	Blockley et al., 2018
D16	Late Epigravettian	LTL601A	12369±100	14935-14091	Charcoal	Blockley et al., 2018
D20	Late Epigravettian	LTL602A	12438±85	14996-14188	Charcoal	Blockley et al., 2018
F1	Early Epigravettian	LTL1593A	17376±90	21345-20760	Charcoal	Blockley et al., 2018
F2	Early Epigravettian	LTL239A	18978±130	23152-22511	Charcoal	Blockley et al., 2018
F3I	Early Epigravettian	LTL606A	18483±95	22618-22185	Charcoal	Blockley et al., 2018
G1	Late Gravettian	LTL236A	19351±180	23761-22987	Charcoal	Blockley et al., 2018
G2	Late Gravettian	LTL237A	19373±90	23742-23043	Charcoal	Blockley et al., 2018
H4	Late Gravettian	LTL604A	20210±245	24974-23800	Charcoal	Blockley et al., 2018
I	Evolved Gravettian	LTL1048A	23475±190	27925-27295	Charcoal	Blockley et al., 2018



	Sample Origin	Site and sampling area	Layer	Chronology	N. sample	N. remains	Frequencies (remains per litre)	
<i>Grotta della Serratura</i>	Old Pollen Samples	Back of the Cave trench	4	Undifferentiated. Epipaleolithic	1	0		
			8	Final Epigravettian	3	24		
			9	Late Epigravettian	1	17		
			11	Gravettian	2	3		
			13	Gravettian	1	0		
		Atrium trench	E	Gravettian	1	1		
			F	Gravettian	1	0		
						<b>Total 10</b>	<b>Total 45</b>	<b>9</b>
	Sifting operation	Back of the Cave trench	8	Final Epigravettian	33	51		
			9	Late Epigravettian	1	1		
10			Late Epigravettian	3	7			
Atrium trench		C	Final Epigravettian	1	1			
		D	Gravettian	2	1			
		E	Gravettian	6	0			
					<b>Total 46</b>	<b>Total 61</b>	<b>0</b>	
<i>Grotta del Romito</i>	Manual flotation	2018 excavations	D (D4-D7)	Late Epigravettian	59	21		
						<b>Total 59</b>	<b>Total 21</b>	<b>0,07</b>
	Old Pollen Samples	2000-2004 excavation	D (2-22)	Late Epigravettian	24	10		
			F	Early Gravettian	3	1		
			G	Late Gravettian	2	0		
			H	Late Gravettian	8	3		
			I	Evolved Gravettian	2	0		
			L	Gravettian	1	0		
		Southern section	-58/60	Late Epigravettian	1	0		
			-75		1	0		
-80				1	2			
-95				1	0			
-110				1	0			
-133				1	1			
-137				1	0			
-190				1	0			
-255				1	0			
-290				1	0			
-330			1	0				
-345		1	0					
-350		1	0					
Burial – Romito 8	D7 pit	Late Epigravettian	22	16				
					<b>Total 76</b>	<b>Total 33</b>	<b>0,87</b>	
<i>Grotta del Cavallo</i>	Sifting operation	2016 excavation	D (D3)	Epigravettian	1	6		
						<b>Total 1</b>	<b>Total 6</b>	<b>0</b>

Manual flotation	Excavation 2018/2019	A1	Neolithic	2	35	
		A2	Mesolithic	2	13	
		B1	Epiromanellian	25	196	
		B2	Romanellian	15	115	
		B2 Pit	Upper Paleolithic (?)	5	67	
				<b>Total 53</b>	<b>Total 426</b>	<b>1,08</b>
<b>TOTAL</b>				<b>245</b>	<b>592</b>	

Site	Lab label	Cultural attribution	AMS dating (IntCal 20)			Material	References
			Lab Code	$^{14}\text{C}$ age (BP)	95.4% confidence, cal BP		
Oriente C	7E	Late Epigravettian	LTL873A	12132 ± 80	14235-13796	Charcoal	Lo Vetro and Martini, 2006
Oriente C	Sublayer 7D	Late Epigravettian	LTL14260 A	12149 ± 65	14217-13803	Charcoal	Mannino et al., 2011
San Teodoro	1	Late Epigravettian	ETH-34451	12580 ± 130	15315-14255	Human bone	Mannino et al., 2011
San Teodoro	4	Late Epigravettian	LTL14278 A	12531 ± 80	15130-14229	Human bone	Mannino et al., 2011
San Teodoro	5	Late Epigravettian	LTL14277 A	12637 ± 100	15322-14432	Human bone	Martini et al., 2013
Romito	4	Late Epigravettian	LTL3032A	11340±90	13366-13095	Human bone	Martini and Lo Vetro, 2018
Romito	5	Late Epigravettian	LTL3033A	10862±70	12930-12716	Human bone	Martini and Lo Vetro, 2018

	Site	Individual	Cultural attribution	Calculus location		
				<i>Tooth</i>	<i>Surface</i>	<i>Weight (mg)</i>
1	Romito	Romito 4	Late Epigravettian	43	buccal	3.4
2		Romito 5	Late Epigravettian	43	buccal	10.6
3		Romito 6	Late Epigravettian	46, 14, 42	lingual	3.1, 2.9, 1.5
4		Romito 7	Late Epigravettian		buccal	0.5
5		Romito 8	Late Epigravettian		lingual	2.2
6	San Teodoro	ST 3	Late Epigravettian	38, 42, 47	lingual	4.1, 2.4, 2.2
7		ST 5	Late Epigravettian	32, 44	lingual	0.5, 0.3
8		ST 6	Late Epigravettian	12; 14	lingual	4, 3.1
10	Oriente C	Oriente	Late Epigravettian	34	buccal	2.1

	Site	Burial label	Chrono-cultural attribution	Type I <i>Triticeae</i>	Type II <i>Paniceae</i>	Type IV <i>Andropogoneae</i>	Type IV <i>Adoxaceae</i>
2	Grotta del Romito	<b>ROMITO 5</b>	Late Upper Palaeolithic	3	3	4	
4	Grotta del Romito	<b>ROMITO 7</b>	Late Upper Palaeolithic	>50			
6	Grotta di San Teodoro	<b>ST3</b>	Late Upper Palaeolithic	>15	11		4
7	Grotta di San Teodoro	<b>ST5</b>	Late Upper Palaeolithic	2			
8	Grotta di San Teodoro	<b>ST6</b>	Late Upper Palaeolithic				
9	Grotta di Oriente	<b>ORIENTE C</b>	Late Upper Palaeolithic	10			
	<b>Total</b>			>80	14	4	4