

APPENDIX

Supplemental table A1 Data collection during study period

Data	Reported by	Tool	T0	T1-n						TD	
			Baseline	Every month or 2 months, as per clinical practice (for max 12 months), at the time of evidence of instability and at the change of the care setting						Patient's death	
Socio-demographic	Patient's Electronic Medical Record	Ad hoc form (eCRF)	X								
Medical History	Patient's Electronic Medical Record	Ad hoc form (eCRF)	X								
PaP Score	MD	Pap Score (eCRF)	X	X	X	X	X	X	X	X	
Necpal	MD	NECPAL questionnaire (eCRF)	X	X	X	X	X	X	X	X	
ECOG-PS	MD	Ad hoc form (eCRF)	X	X	X	X	X	X	X	X	
ESAS scale	Patient	ESAS Scale (eCRF)	X	X	X	X	X	X	X	X	
Dyspnea and Cough (only for pulmonary pop.)	MD	M-Borg and NRS Scale (eCRF)	X	X	X	X	X	X	X	X	
Use of health services and end-of-life care	Patient's Electronic Medical Record	Ad hoc form (eCRF)									X
Date and location of death	Patient's Electronic Medical Record	Ad hoc form (eCRF)									X

MD, medical doctor; eCRF, electronic case report forms.