

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Steroid reference intervals in women: influence of menopause, age and metabolism

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Mezzullo, M., Gambineri, A., Di Dalmazi, G., Fazzini, A., Magagnoli, M., Baccini, M., et al. (2021). Steroid reference intervals in women: influence of menopause, age and metabolism. EUROPEAN JOURNAL OF ENDOCRINOLOGY, 184(3), 395-407 [10.1530/EJE-20-1147].

Availability:

This version is available at: https://hdl.handle.net/11585/858738 since: 2022-02-15

Published:

DOI: http://doi.org/10.1530/EJE-20-1147

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Marco Mezzullo, Alessandra Gambineri, Guido Di Dalmazi, Alessia Fazzini, Matteo Magagnoli, Margherita Baccini, Valentina Vicennati, Carla Pelusi, Uberto Pagotto, Flaminia Fanelli, Steroid reference intervals in women: influence of menopause, age and metabolism, European Journal of Endocrinology, Volume 184, Issue 3, Mar 2021, Pages 395–407

The final published version is available online at: <u>https://doi.org/10.1530/EJE-20-</u> <u>1147</u>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

Steroid reference intervals in women: influence of menopause, age and metabolism

Marco Mezzullo, Alessandra Gambineri, Guido Di Dalmazi, Alessia Fazzini, Matteo Magagnoli, Margherita Baccini, Valentina Vicennati, Carla Pelusi, Uberto Pagotto and Flaminia Fanelli

Affiliations: Unit of Endocrinology and of Prevention and Care of Diabetes, Center for Applied Biomedical Research, Department of Medical and Surgical Sciences, University of Bologna, S.Orsola Policlinic, via Massarenti 9, 40138 Bologna, Italy

Correspondence and reprint requests to: Flaminia Fanelli, Junior Assistant Professor, Department of Medical and Surgical Sciences S.Orsola Policlinic, Via Massarenti 9, 40138 Bologna, Italy. Tel and Fax: +39-051-2143902; E-mail: flaminia.fanelli2@unibo.it

Short title: LC-MS/MS steroid reference interval in women

Keywords: liquid chromatography tandem mass spectrometry (LC-MS/MS), steroid profile, reference intervals, obesity, dysmetabolism, menopause, menstrual phase

1

Word Count: 3501 Number of Tables and Figures: 6 Number of Supplemental files: 2

Abstract

Objective. To investigate the impact of age, obesity and metabolic parameters on thirteen circulating steroids in reproductive and menopausal age. To define reference intervals (RI).

Design. Cross-sectional.

Methods. 325 drug-free, healthy and eumenorrheic women were selected from the general population. Independent relationships of LC-MS/MS-determined steroid levels with age, body mass index (BMI) and metabolic parameters were estimated. Reference sub-cohorts were defined for calculating upper and lower limits in reproductive age, menstrual phases and menopause, and these were compared with limits in dysmetabolic sub-cohorts.

Results. Lower androgens, pro-androgens and estrogens, but higher cortisol and metabolites were found in menopausal compared to reproductive age women. Androgens and precursors decreased during reproductive age (P<0.001-P=0.002) but not after menopause. 17OH-progesterone decreased with BMI (P=0.006) and glucocorticoids with waist circumference (P<0.001-P=0.002) in reproductive age, but increased with triglycerides (P=0.011-P=0.038) after menopause. Inverse associations of dihydrotestosterone with BMI (P=0.004) and HDL-cholesterol (P=0.010), estrone with total cholesterol (P=0.033) and estradiol with triglycerides (P=0.011) were found in reproductive age. After menopause, estrone increased with waist circumference (P<0.001) and decreased with insulin resistance (P=0.012). Ovarian steroid RI were estimated in menstrual phases and menopause. Age- and reproductive age cortisol (P=0.020) and menopausal 11-deoxycortisol (P=0.003) in dysmetabolic sub-cohorts were reduced and increased, respectively, compared to reference limits.

Conclusions: Obesity and dysmetabolism differently influence circulating steroids in reproductive and menopausal status. Age, menstrual and menopausal status-specific RI were provided by LC-MS/MS for a broad steroid panel.

Introduction

LC-MS/MS is used by an increasing number of clinical laboratories, providing reliable determinations of steroid hormones for the diagnosis and management of endocrine diseases [1; 2]. By enlarging the panel of measurable precursors and metabolites, LC-MS/MS is also boosting renewed interest in understanding the steroid system derangement in several conditions. However, translating research findings into clinical advancements requires the definition of pathophysiologic states influencing the circulating steroid levels, as well as of appropriate reference intervals (RI) allowing the effective interpretation of laboratory results [3]. Recently, LC-MS/MS has been applied to study the steroid profile of adrenal tumors [4], type I diabetes [5], or female hyperandrogenism [1; 6; 7; 8; 9]. In contrast, only a few studies were purposely designed to define steroid RI. Moreover, some studies relied on healthcare-seeking subjects. This represents an intrinsic bias for RI estimation, given the adaptive nature of steroid hormones to stressful or debilitating conditions. Other studies did not report an exhaustive characterization of subjects, for example aimed at detecting subtle hyperandrogenic states manifesting as menstrual irregularities or hirsutism, while others tolerated excess weight [10].

Overweight and obesity affect half of the female population [11; 12]. Steroids are involved in systemic and intracrine mechanisms that, once dysregulated, contribute to excess fat, mostly of abdominal type, hypertension, impaired glucose control and dyslipidaemia. However, the interplay among glucocorticoids, androgens and estrogens in such mechanisms varies with the menopausal transition [12; 13]. Androgen excess, ovarian dysfunction and metabolic impairment represent a vicious cycle in young women, however, a variegated spectrum of manifestations is observed, for which effective hormonal markers are yet undefined [14]. The androgen decline and the loss of ovarian hormones contribute to the central redistribution of body fat, the increased cardiometabolic risk, sarcopenia and bone frailty experienced after menopause. Besides, the modulation of adrenal function throughout women lifespan, and the interplay between the hypothalamus-pituitary-adrenal (HPA) axis dysregulation and obesity before and after menopause were not elucidated [15; 16].

In a recent study, we demonstrated that the circulating steroid profile in men is not only influenced by ageing, but also by obesity and metabolic derangement, and we provided age-specific RI estimated in appropriate reference cohorts [17]. In the present study, by using a similarly standardized procedural

approach, we aimed at describing the relationships of age, obesity and metabolic status with thirteen circulating steroids in women in reproductive and menopausal age. Based on the depicted associations, we generated age, menopause and menstrual phase specific RI in selected reference sub-cohorts, and investigated whether unrestricted inclusion criteria could bias steroid RI estimation.

Materials and methods

Subjects

Women aged 18–86 years were recruited from the general population [18]. The study was approved by the S.Orsola Policlinic ethical committee (85/2008/O/Tess). All women signed the informed consent before they were interviewed and examined by a trained endocrinologist, between 08:00 and 10:00 am. Waist circumference and body mass index (BMI) were recorded. Systolic (SBP) and diastolic (DBP) blood pressures were measured in supine position after 3 min rest. Inclusion criteria were: BMI \geq 18.5 kg/m², weight stability in previous 3 months and complete sexual development. Exclusion criteria included signs of clinical hyperandrogenism, history of menstrual irregularities, steroidal (including estro-progestogen) and non-steroidal drug assumption in previous 3 months (except antipyretic or non-steroidal anti-inflammatory compounds tolerated before the previous month, and thyroxine replacement in compensated hypothyroidism), present or past endocrine, hepatic, renal, oncologic, autoimmune, cardiovascular, hematologic, neurologic or psychiatric diseases, sleep disorders, shift working, frequent flying or allergies requiring treatment. Among 653 women examined, 328 were excluded because of present or previous diseases (n=32), irregular sleeping (n=80), glucose-lowering (n=3), cholesterol-lowering (n=111), anti-hypertensive (n=171) and anti-depression (n=40) drug assumption. Therefore, 325 women were included in the present study.

Biochemical and hormonal evaluation

Blood was withdrawn in overnight fasting condition in Vacuette Z serum beads clot activator tubes (Greiner Bio-One, Kremsmunster, Austria) after 10 min saline infusion for minimizing venepuncture stress. After 20 min settling, tubes were centrifuged (2000 g, 10 min, room temperature) and serum for LC-MS/MS was

stored at -80° C. Routine hormones and biochemicals were measured in fresh blood as previously reported [17]. The homeostatic model assessment-insulin resistance (HOMA-IR) was computed [19].

Steroid measurement by LC-MS/MS

We applied two in-house LC-MS/MS assays (**Supplemental Table 1**) including 17-hydroxypregnenolone (170HP5), dehydroepiadrosterone (DHEA), progesterone (P4), 17-hydroxyprogesterone (170HP4), 11-deoxycorticosterone (DOC), corticosterone (B), 11-deoxycortisol (11S), cortisol (F), androstenedione (A4), testosterone (T), dihydrotestosterone (DHT), estrone (E1) and estradiol (E2) [18; 20; 21].

Study design

Women were classified in reproductive (regular menses, age 18-54 years; n=186) or menopausal (amenorrhea in previous 12 months or more, age 48-86 years; n=127) age, and further subdivided in normal weight (NW, 18.5<BMI ≤ 25.0 kg/m²; n=123 and 63, respectively), overweight (OW, 25.0<BMI ≤ 30.0 kg/m²; n=35 and 49, respectively) and obese (OB, BMI>30.0 kg/m²; n=28 and 15, respectively). Women in reproductive age were further stratified in early follicular (day 1-6; n=31), pre-ovulatory (day 9-13; n=30) and mid-luteal (day 18-24; n=33) menstrual phases. Perimenopausal women were also identified (<6 menstrual bleedings in previous 6 months or more, age 48-54 years; n=12). Anthropometric, metabolic and steroid values were compared between reproductive and menopausal cohorts, among BMI classes and among menstrual phases. Afterward, we assessed the independent effect of age, BMI, waist circumference, SBP, DBP, HOMA-IR, total cholesterol, HDL-cholesterol and triglycerides on each steroid in reproductive and menopausal ages. According to the results, we defined steroid-specific reference sub-cohorts by excluding women displaying alterations in metabolic parameters influencing the steroid levels. Therefore, we identified subjects with normal (≤88 cm, norWC; n=216) or elevated (>88 cm, dysWC; n=106) waist circumference, normal (<2.5, norHOMA; n=194) or elevated (≥2.5, dysHOMA; n=36) HOMA-IR, normal (<5.17 mmol/L, norTC; n=194) or elevated (\geq 5.17 mmol/L, dysTC; n=123) total cholesterol, normal (\geq 1.29 mmol/L, norHDL; n=203) or reduced (<1.29 mmol/L, dysHDL; n=55) HDL-cholesterol, and normal (<1.69 mmol/L, norTG; n=302) or elevated (\geq 1.69 mmol/L, norTG; n=15) triglycerides. Lower (LRL) and upper (URL) reference limits defining the central 95% of steroid distribution were estimated in reference sub-cohorts. Age-specific LRL and URL were estimated when required. Finally, to evaluate whether altered metabolic parameters could influence RI estimation, LRL and URL were compared with lower and upper limits calculated in dysmetabolic sub-cohorts including subjects with alterations in the parameters influencing that particular steroid, respectively.

Statistical analysis

Box-Cox transformation was used for variables showing a significant skewness at the Kolmogorov–Smirnov test [22]. Far outliers at the Tukey's test were removed [23]. Variables were compared between reproductive age and menopausal cohorts by T-test. The ANOVA trend test was used for comparing BMI classes (SPSS package v.20, IBM Co). Comparisons among menstrual phases were performed by one-way ANOVA. The stepwise multiple regression included age, BMI, waist circumference, SBP, DBP, HOMA-IR, total-cholesterol, HDL-cholesterol and triglyceride levels as covariates, and each steroid as dependent variable. The menstrual phase was added as cofactor for steroids varying with the menstrual cycle. The effect size (f^2) was estimated as $f^2 = \frac{sr^2}{(1-R^2 full)}$, where sr^2 is the semipartial correlation coefficient for the predictor of

interest and R is the full correlation coefficient obtained by the multiple regression model [24].

LRL and URL were estimated as the mean – and + (1.96×s.d.) of the transformed variables, respectively, then, values were back-transformed to the original unit [25]. Age-specific RI were estimated by modelling the transformed steroid variable on age distribution, according to the fractional polynomial regression by Royston and Wright [26]. Age (*X*) was transformed in order to stabilize the steroid variable (*Y*) for large values of *X* according to the formula: $eX = \exp \frac{(\log (0.01) \times (X - \min (X))}{(\max (X) - \min (X))}$. Then, we selected the optimal model providing the lowest polynomial degree (parsimony) with maximum decrease in deviance (goodness of fit). Best-fit polynomial coefficients were selected by fp syntax, and RI were visually inspected by xrigls syntax in STATA (v.13.0, StataCorp LLC).

A large number of cases showed values below the sensitivity limit for DOC in the whole cohort (n=276 of 325), and P4, DHT and E2 in menopausal cohort (n=90, 75 and 115 of 127, respectively) (**Supplemental Table 1**). Therefore, DOC comparisons were performed by Kruskal-Wallis and Mann-Whitney tests, while no multiple regression was performed for DOC and for P4, DHT and E2 in menopausal cohort. RI for DOC and P4, and menopausal RI for DHT and E2 were estimated as the 2.5–97.5 centiles of distribution. Lower and upper limits were compared between reference and dysmetabolic sub-cohorts by z distribution. Two-

tailed P values <0.05 were considered significant. Data were analysed by MedCalc Software (v.18.2.1, Mariakerke, Belgium) except where specified.

Results

Anthropometric, metabolic and hormonal features of the cohort

Table 1 reports the anthropometric, metabolic and hormonal features of our cohort. Compared to reproductive age, menopausal women showed worse BMI, waist circumference, SBP, DBP, glucose, total cholesterol, triglycerides (all *P*<0.001) and HOMA-IR (*P*=0.013), lower 17OHP5 (-31.2%), DHEA (-42.6%), P4 (-84.4%), 17OHP4 (-60.1%), A4 (-51.3%), T (-22.5%), DHT (-23.1%), E1 (-63.6%) and E2 (-87.4%) (all *P*<0.001), but higher DOC (n.d., *P*=0.044), 11S (18.2%, *P*=0.029) and F (11.3%, *P*=0.007) (**Table 1**).

Worsening metabolic parameters at increasing BMI were observed both in reproductive age (min-max BMI: 18.5-42.9 kg/m²) and menopausal women (min-max BMI: 18.9-41.2 kg/m²). In reproductive age, levels of B (P<0.001), F (P=0.008) and DHT (P=0.006) decreased with increasing BMI classes, with OB showing lower B, F and DHT compared to NW (P=0.002, P=0.023 and P=0.017) and lower B and F compared to OW (both P<0.001) women. In menopausal women, E1 levels increased with BMI classes (P=0.002), with both OW (P=0.046) and OB (P=0.007) displaying higher values than NW women (**Table 1**). Women in different menstrual phases displayed similar anthropometric and metabolic parameters, but different levels of P4, 17OHP4, E1, E2 (all P<0.001), DOC (P=0.043) and 11S (P=0.048). In particular, women in mid-luteal phase had higher P4 and 17OHP4 as compared to early follicular and pre-ovulatory (all P<0.001), and higher DOC compared to pre-ovulatory (P=0.047) women. Moreover, lower E1 and E2 levels were found in early follicular compared to pre-ovulatory and mid-luteal phases (all P<0.001) (**Supplemental Table 2**).

Independent impact of age, anthropometric and metabolic parameters on steroid levels

Stepwise multiple regression results are detailed in **Table 2**. In women in reproductive age, 17OHP5, DHEA, A4, T (all P<0.001) and DHT (P=0.002) decreased with age, with a large effect size for A4 (f^2 =0.28), DHEA and T (both f^2 =0.22), and moderate for 17OHP5 (f^2 =0.12) and DHT (f^2 =0.11). DHT was also negatively associated with BMI (P=0.004, f^2 =0.09) and HDL-cholesterol (P=0.010, f^2 =0.08). 17OHP4 inversely

associated with BMI (P=0.006, $f^2=0.22$), while B (P<0.001, $f^2=0.20$), F (P<0.001, $f^2=0.17$) and 11S (P=0.002, $f^2=0.09$) inversely associated with waist circumference. Finally, E1 and E2 negatively associated with increasing total cholesterol (P=0.033, $f^2=0.10$) and triglycerides (P=0.011, $f^2=0.15$), respectively. In menopausal women, no age and BMI influence were detected on steroid levels. 17OHP4 (P=0.026, $f^2=0.06$), 11S (P=0.011, $f^2=0.07$) and F (P=0.038, $f^2=0.05$) directly associated with triglycerides. E1 directly associated with waist circumference (P<0.001, $f^2=0.16$), and inversely associated with HOMA-IR (P=0.012, $f^2=0.08$).

Steroid reference intervals

Age- and menopause-specific RI were estimated for 17OHP5, DHEA, A4, T and DHT (**Table 3**). All women were included in the reference sub-cohort for 17OHP5, DHEA, A4 and T (n=325). For DHT, the reference sub-cohort included NW and norHDL women in reproductive age and all menopausal women (n=65 and n=127, respectively). Androgens and precursors peak around age 25 years and progressively decline. A reduction in LRL and URL was found for A4 (-83.2 and -57.5%, respectively), DHEA (-67.6 and -56.0%, respectively), 17OHP5 (-49.9 and -43.1%, respectively) and T (-24.9 and -17.2%, respectively), as well as in DHT URL (-56.3%) from age 25 to 65 years (**Figure 1**).

Reproductive age RI were estimated in the whole cohort for DOC (n=186), and in norWC sub-cohort for B, 11S and F (n=132). Menopausal RI were estimated in the whole cohort for DOC, B, P4 and E2 (n=127), in norTG sub-cohort for 17OHP4, 11S and F (n=112), and in women who were both norWC and norHOMA for E1 (n=50) (**Table 4**).

Finally, menstrual phase specific RI were estimated for P4, 17OHP4, E1 and E2. Of 32 women in mid-luteal phase, 10 (age 39.2±10.3 years) exhibited P4 levels <10 nmol/L suggestive of incorrect classification or of anovulatory cycle [27], and were therefore excluded. The reference sub-cohorts for early follicular, pre-ovulatory and mid-luteal phase included 31, 30 and 22 women for P4; 22, 26 and 17 NW women for 17OHP4; 20, 25 and 17 norTC women for E1; and 25, 28 and 21 norTG women for E2, respectively (**Table 5**).

Impact of metabolic risk factors on the estimation of reference limits

LRL and URL of steroids influenced by metabolic parameters were compared with lower and upper limits calculated in dysmetabolic subjects. Higher 11S values were found in the lower limit calculated in dysTG (n=10) compared to LRL in norTG (n=112) menopausal women (+0.343 nmol/L, +48.4%, P=0.003). At variance, reduced F values were found in lower limit in dysWC (n=52) compared with LRL in norWC (n=132) women in reproductive age (-35.5 nmol/L, -24.0%; P=0.020, respectively) (**Table 4**).

Discussion

In the present study, to obtain RI as effective as possible when applied to the study of women health, we selected from the general population women who were drug- and disease-free and having no signs or symptoms of androgen excess. Nonetheless, our cohort included women with unmedicated excess weight, dyslipidaemia, impaired insulin sensitivity and hypertension, overall affecting a relevant portion of the general population.

Our results confirmed the dramatic decline in circulating estrogens, progestins, androgens and precursors with menopause [28]. No age dependency was detected for estrogen and progestin levels in reproductive age, therefore, the 60-90% hormone reduction observed after menopause could totally be attributed to ovarian senescence. At variance, the circulating androgens and precursors peaked around age 25 years and started a declining trend long before the menopausal transition. A steep decrease from age 25 to 65 years was observed for A4, DHEA 170HP5, and DHT, ranging 40 to 85%, while a moderate 20% decrease was found for T, overall in good agreement with previous estimates [29; 30; 31; 32; 33; 34; 35; 36]. Given the relevance of peripheral androgen generation from adrenal precursors in females, this finding may be due to the combined effect of ovarian and adrenal senescence [30]. Interestingly, recent studies showed that *zona reticularis*, but not *zona fasciculata*, undergoes a large involution with ageing, thus explaining the impairment of pro-androgen secretion [33; 36].

In our cohort, corticosteroid levels were not influenced by age, however, in keeping with a recent study [33], DOC, 11S and F slightly increased after menopause. This is in contrast with studies reporting an agedependent reduction of glucocorticoids [31; 32]. The modulation of steroid metabolites we observed suggests an increased adrenal secretion with menopause rather than a peripheral reactivation of F [37], as supported by an elegant study demonstrating a slight increase in menopausal F secretion in response to ACTH [30]. When we tried to disentangle which specific metabolic component related to a particular steroid level, we found a different network of relationships before and after menopause. Notably, in our non-hyperandrogenic women, no associations of androgens and precursors with metabolic parameters were found, except for DHT, inversely correlating with BMI and HDL-cholesterol in reproductive age. Unfortunately, the limited sensitivity of our assay prevented a similar DHT evaluation in menopausal women. The link between androgen levels and excess weight in women still has not been clarified. Indeed, a variegated combination of unchanged or reduced levels of DHEA, A4, T and DHT was associated with increasing adiposity in non-PCOS women by studies using high-specificity MS-based assays [31; 33; 34; 36; 38, 39; 40; 41]. Interestingly, low plasma DHT was associated with increasing dimension and lipogenesis function of omental adipocytes [38]. Our findings, therefore, support the concept that, in obese women, low levels of the most active androgen associate with the lipid storage capacity of visceral depots.

Interestingly, increasing estrogen levels contributed to the healthy lipid profile of our young women, underlying the importance of a balanced orchestration among active sex steroids. Besides, after menopause, E1 inversely associated with insulin resistance and directly associated with waist circumference. While, on the one hand, this is consistent with adipose tissue being the predominant estrogen source after ovarian senescence [13; 28], on the other hand, it contrasts with the detrimental link between abdominal obesity and insulin resistance. The beneficial involvement of estrogens in energy metabolism and inflammatory response was widely described [28; 42], and is evidenced by the vicious circle among abdominal adiposity, systemic low-grade inflammation, insulin resistance and increasing cardiovascular risk occurring with menopause [12]. It is recognized that estrogen levels in menopause are proportional to body fat [13; 28], however, whether visceral or subcutaneous fat is the predominant source, and whether estrogens are associated with a favourable or unfavourable metabolic status in this life epoch, is still unclear. A study using LC-MS/MS found that visceral fat secreted E1 more than the subcutaneous fat in menopausal women, however, no correlations with the metabolic profile were performed [43]. Another study found stronger associations of plasma estrogens with total rather than abdominal adiposity; moreover, direct associations between estrogens and insulin resistance were found, but these depended on the amount of fat [39]. Given this scenario, our findings could be explained by the fact that, as we excluded medicated and diabetic subjects, our menopausal women are overall only mildly dysmetabolic. Unfortunately, in our population study we could not use

techniques such as computed tomography to distinguish between abdominal visceral and subcutaneous depots. Nonetheless, it is possible that fat amount reflected by waist circumference, be it subcutaneous or visceral, is prevalently metabolically healthy, so that the quantitative relationship between circulating E1, spilling-over from fat depots, and the beneficial result of its function in terms of insulin sensitivity, is still detectable. Unfortunately, due to the limited analytical sensitivity, we could not investigate the metabolic associations of E2 in menopausal age.

Waist circumference was the only independent factor negatively affecting glucocorticoid levels in reproductive age. Moreover, menstrual phase-adjusted 17OHP4 levels, possibly reflecting its role as adrenal glucocorticoid precursor, diminished with increasing BMI, as reported in a recent study [31]. This may appear in contrast with the direct link expected between glucocorticoid tone and visceral obesity. Nonetheless, the dynamic of HPA axis in obesity is complex, and early morning F levels were previously found unchanged or inversely related with central adiposity, as in consequence of diminished ACTH-sensitivity of the adrenal, of flattening of HPA circadian rhythmicity, or of increased F clearance [44; 45]. At variance, the positive association of glucocorticoids with triglycerides we found in menopausal women is in line with their role in the derangement of energy substrate utilization [46]. Notably, we demonstrated that even mild metabolic dysfunction, such as high waist circumference or triglycerides, can significantly alter the estimation of 11S and F limits, underlying the importance of a proper metabolic characterization when generating glucocorticoid RI.

Menstrual fluctuation challenges the definition of robust RI for ovarian steroids [47]. A limit of our study is that gonadotropins were not measured, and menstrual classification was performed according to the menstrual date, which may represent a source of bias. We could partially cope with this unpredictable bias in the mid-luteal phase, by excluding women showing P4 levels <10 nmol/L [27]. However, we could not assess whether these women were actually misclassified or they had an occasional anovulatory cycle. Similarly, we cannot exclude that misclassification or anovulatory cycles were affecting women in early follicular and pre-ovulatory phases. Therefore, our menstrual phase RI are to be interpreted with cautions, and need to be refined in future studies including gonadotropin evaluation. Nonetheless, values we observed in early follicular phase are in reasonable agreement with previous reports [27; 31; 48]. Androgen fluctuation throughout the menstrual cycle is still debated. Skiba et al. found slightly higher T and A4 in mid-cycle and

luteal compared to follicular phase [34]. Bui et al. observed minimal T fluctuation, with increasing values at mid-cycle in some women [49]. We did not detect any androgen fluctuation, however, we found small but significant variations in DOC and 11S levels among menstrual phases. A proper exploration of this phenomenon would require repeated intra-subject evaluation across the menstrual cycle. Even though data are still inconclusive, we recommend to standardize the menstrual phase when studying the steroid dynamics in young women for clinical or research purposes.

Our steroid values are generally lower compared to other LC-MS/MS studies, except 17OHP5 and DHEA values which are higher, [27; 31; 32; 34; 35; 48; 50; 51; 52]. Differences in analytical methods, sampling procedures, cohort selection and study design may account for variabilities in steroid levels and relationships with women's metabolic health. Time of sampling [52] as well as needle stress [53; 54] represent non-negligible sources of variability. In addition, though harmonization of LC-MS/MS measures seems an affordable goal, collective strategies are still awaited in this direction.

In conclusion, our study focusing on a carefully selected female cohort highlighted a different network of relationships between circulating steroid profile, obesity and metabolic status of women in reproductive and menopausal age. RI specific for age, reproductive and menstrual status were generated that will be useful for the effective interpretation of the steroid involvement in women's health and disease.

Declaration of Interest: the Authors report no conflict of interest in this work. Alessandra Gambineri is on the editorial board of EJE. Alessandra Gambineri was not involved in the review or editorial process for this paper, on which he/she is listed as an author.

Funding: the study was supported by the European Union (REPROBESITY, FPVII-223713), by the Emilia-Romagna Region – University Program 2007-2009 (g.a. PRUa1a-2007-006), by the Emilia-Romagna Region, Alessandro Liberati Young Researcher Grants (g.a. PRUA 1-2012-004).

Author contribution: MM1 measured study samples, performed the statistical analysis and wrote the manuscript; AG, GDD, MB, VV and CP performed cohort recruitment and examination; AF and MM2

measured study samples; UP designed the population study and wrote the manuscript; FF designed the study and wrote the manuscript.

Acknowledgements: we thank Shimadzu for kindly providing the LC-MS/MS platform used to generate part of the data used in this study.

References

1. Ketha SS, Singh RJ & Ketha H. Role of mass spectrometry in clinical endocrinology. *Endocrinology and Metabolism Clinics of North America* 2017 **46** 593–613.

2. Wudy SA, Schuler G, Sánchez-Guijo A & Hartmann MF. The art of measuring steroids: principles and practice of current hormonal steroid analysis. *The Journal of Steroid Biochemistry and Molecular Biology* 2018 **179** 88-103.

3. Ceriotti F, Hinzmann R & Panteghini M. Reference intervals: the way forward. Annals of Clinical Biochemistry 2009 **46** 8–17.

4. Fanelli F & Di Dalmazi G. Serum steroid profiling by mass spectrometry in adrenocortical tumors: diagnostic implications. *Current Opinion in Endocrinology, Diabetes, and Obesity* 2019 **26** 160–165.

5. Gunness A, Pazderska A, Ahmed M, McGowan A, Phelan N, Boran G, Taylor AE, O'Reilly MW, Arlt W, Moore K et al. Measurement of selected androgens using liquid chromatography-tandem mass spectrometry in reproductive-age women with Type 1 diabetes. *Human Reproduction* 2018 **33** 1727-1734.

6. Pasquali R, Zanotti L, Fanelli F, Mezzullo M, Fazzini A, Morselli-Labate AM, Repaci A, Ribichini D & Gambineri A. Defining Hyperandrogenism in Women With Polycystic Ovary Syndrome: A Challenging Perspective. *The Journal of Clinical Endocrinology & Metabolism* 2016 **101** 2013-2022.

7. O'Reilly MW, Kempegowda P, Jenkinson C, Taylor AE, Quanson JL, Storbeck KH & Arlt W. 11-Oxygenated C19 Steroids Are the Predominant Androgens in Polycystic Ovary Syndrome. *The Journal of Clinical Endocrinology & Metabolism* 2017 **102** 840-848. 8. Sharma A, Kapoor E, Singh RJ, Chang AY & Erickson D. Diagnostic Thresholds for Androgen-Producing Tumors or Pathologic Hyperandrogenism in Women by Use of Total Testosterone Concentrations Measured by Liquid Chromatography-Tandem Mass Spectrometry. *Clinical Chemistry* 2018 **64** 1636-1645.

9. Turcu AF, El-Maouche D, Zhao L, Nanba AT, Gaynor A, Veeraraghavan P, Auchus RJ & Merke DP. Androgen excess and diagnostic steroid biomarkers for nonclassic 21-hydroxylase deficiency without cosyntropin stimulation. *The European Journal of Endocrinology* 2020 **183** 63-71.

10. Fanelli F, Baronio F, Ortolano R, Mezzullo M, Cassio A, Pagotto U & Balsamo A. Normative basal values of hormones and protein of gonadal and adrenal functions from birth to adulthood. *Sexual Development* 2018 **12** 50–94.

11. Piché ME, Poirier P, Lemieux I & Després JP. Overview of epidemiology and contribution of obesity and body fat distribution to cardiovascular disease: an update. *Progress in Cardiovascular Diseases* 2018 **61** 103–113.

12. Gerdts E & Regitz-Zagrosek V. Sex differences in cardiometabolic disorders. *Nature Medicine* 2019 **25** 1657-1666.

13. Guarner-Lans V, Rubio-Ruiz ME, Pérez-Torres I & Baños de MacCarthy G. Relation of aging and sex hormones to metabolic syndrome and cardiovascular disease. *Experimental Gerontology* 2011 **46** 517-23.

14. Pasquali R & Gambineri A. Polycystic ovary syndrome: a multifaceted disease from adolescence to adult age. *Annals of the New York Academy of Sciences* 2006 **1092** 158-174.

15. Pasquali R, Vicennati V, Gambineri A & Pagotto U. Sex-dependent role of glucocorticoids and androgens in the pathophysiology of human obesity. *International Journal of Obesity* 2008 **32** 1764–1779.

16. Nieuwenhuizen AG & Rutters F. The hypothalamic-pituitary-adrenal axis in the regulation of energy balance. *Physiology and Behavior* 2009 **4** 169–177.

17. Mezzullo M, Di Dalmazi G, Fazzini A, Baccini M, Repaci A, Gambineri A, Vicennati V, Pelusi C, Pagotto U & Fanelli F. Impact of age, body weight and metabolic risk factors on steroid reference intervals in men. *European Journal of Endocrinology* 2020 **182** 459-471.

18. Fanelli F, Belluomo I, Di Lallo VD, Cuomo G, De Iasio R, Baccini M, Casadio E, Casetta B, Vicennati V, Gambineri A et al. Serum steroid profiling by isotopic dilution-liquid chromatography-mass spectrometry: comparison with current immunoassays and reference intervals in healthy adults. *Steroids* 2011 **76** 244–253.

19. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF & Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985 **28** 412–419.

20. Büttler RM, Martens F, Fanelli F, Pham HT, Kushnir MM, Janssen MJ, Owen L, Taylor AE, Soeborg T, Blankenstein MA et al. Comparison of published LC-MS/MS methods for the simultaneous measurement of testosterone, androstenedione, and dehydroepiandrosterone in serum. *Clinical Chemistry* 2015 **61** 1475–1483.

21. Mezzullo M, Pelusi C, Fazzini A, Repaci A, Di Dalmazi G, Gambineri A, Pagotto U & Fanelli F. Serum reference intervals for challenging sex and precursor steroids by liquid chromatography tandem mass spectrometry. *Journal of Steroid Biochemistry and Molecular Biology* 2020 **197** 105538.

22. Box GEP & Cox DR. An analysis of transformations. *Journal of the Royal Statistical Society: Series B* 1964 **26** 211–243

23. Tukey JW. Exploratory Data Analysis. Addison-Wesley. 1977 ISBN 978-0-201-07616-5. OCLC 3058187.

24. Cohen J. Statistical Power Analysis for the Behavioural Sciences. Routledge, 1988.

25. Clinical and Laboratory Standard Institute. Defining, Establishing, and Verifying Reference Intervals in the Clinical Laboratory; Approved Guideline, 3rd ed. CLSI document EP28-A3c: Wayne, PA: Clinical and Laboratory Standards Institute, 2008.

26. Royston P & Wright EM. Method for estimating age-specific reference intervals (normal ranges) based on fractional polynomials and exponential transformation. *Journal of the Royal Statistical Society: Series A* 1998 **161** 79–101.

27. Frederiksen H, Johannsen TH, Andersen SE, Albrethsen J, Landersoe SK, Petersen JH, Andersen AN, Vestergaard ET, Schorring ME, Linneberg A et al. Sex-specific Estrogen Levels and Reference Intervals from Infancy to Late Adulthood Determined by LC-MS/MS. *The Journal of Clinical Endocrinology & Metabolism* 2020 **105** 754–768.

28. Freeman EW, Sammel MD, Lin H & Gracia CR. Obesity and reproductive hormone levels in the transition to menopause. *Menopause* 2010 **17** 718-726.

29. Kroboth PD, Salek FS, Pittenger AL, Fabian TJ & Frye RF. DHEA and DHEA-S: a review. *The Journal* of *Clinical Pharmacology* 1999 **39** 327-348.

30. Parker CR Jr, Slayden SM, Azziz R, Crabbe SL, Hines GA, Boots LR & Bae S. Effects of aging on adrenal function in the human: responsiveness and sensitivity of adrenal androgens and cortisol to adrenocorticotropin in premenopausal and postmenopausal women. *The Journal of Clinical Endocrinology* & *Metabolism* 2000 **85** 48-54.

31. Eisenhofer G, Peitzsch M, Kaden D, Langton K, Pamporaki C, Masjkur J, Tsatsaronis G, Mangelis A, Williams TA, Reincke M et al. Reference intervals for plasma concentrations of adrenal steroids measured by LC-MS/MS: impact of gender, age, oral contraceptives, body mass index and blood pressure status. *Clinica Chimica Acta: International Journal of Clinical Chemistry* 2017 **470** 115–124.

32. Bae YJ, Zeidler R, Baber R, Vogel M, Wirkner K, Loeffler M, Ceglarek U, Kiess W, Körner A, Thiery J *et al.* Reference intervals of nine steroid hormones over the life-span analyzed by LC-MS/MS: Effect of age,

gender, puberty, and oral contraceptives. *Journal of Steroid Biochemistry and Molecular Biology* 2019 **193** 105409.

33. Nanba AT, Rege J, Ren J, Auchus RJ, Rainey WE & Turcu AF. 11-Oxygenated C19 Steroids Do Not Decline With Age in Women. *The Journal of Clinical Endocrinology & Metabolism* 2019 **104** 2615-2622.

34. Skiba MA, Bell RJ, Islam RM, Handelsman DJ, Desai R & Davis SR. Androgens During the Reproductive Years: What Is Normal for Women? *The Journal of Clinical Endocrinology & Metabolism* 2019 **104** 5382-5392.

35. van der Veen A, van Faassen M, de Jong WHA, van Beek AP, Dijck-Brouwer DAJ & Kema IP. Development and validation of a LC-MS/MS method for the establishment of reference intervals and biological variation for five plasma steroid hormones. *Clinical Biochemistry* 2019 **68** 15-23.

36. Davio A, Woolcock H, Nanba AT, Rege J, O'Day P, Ren J, Zhao L, Ebina H, Auchus R, Rainey WE, Turcu AF, Sex Differences in 11-Oxygenated Androgen Patterns Across Adulthood. *The Journal of Clinical Endocrinology & Metabolism* 2020 **105** 1-9.

37. Yamatani H, Takahashi K, Yoshida T, Takata K & Kurachi H. Association of estrogen with glucocorticoid levels in visceral fat in postmenopausal women. *Menopause* 2013 **20** 437-442.

38. Côté JA, Lessard, J, Mailloux J, Laberge P, Rhéaume C, & Tchernof, A. Circulating 5αdihydrotestosterone, abdominal obesity and adipocyte characteristics in women. *Hormone Molecular Biology and Clinical Investigation* 2012 **12** 391-400.

39. Marchand GB, Carreau AM, Weisnagel SJ, Bergeron J, Labrie F, Lemieux S & Tchernof A. Increased body fat mass explains the positive association between circulating estradiol and insulin resistance in postmenopausal women. *American Journal of Physiology-Endocrinology and Metabolism*. 2018 **314** E448-E456.

40. Marchand GB, Carreau A, Laforest S, Côté J, Daris M, Cianflone K, Prehn C, Adamski J & Tchernof A. Circulating steroid levels as correlates of adipose tissue phenotype in premenopausal women. *Hormone Molecular Biology and Clinical Investigation* 2018 **34** 20170082.

41. Vihma V, Heinonen S, Naukkarinen J, Kaprio J, Rissanen A, Turpeinen U, Hämäläinen E, Hakkarainen A, Lundbom J, Lundbom N et al. Increased body fat mass and androen metabolism - A twin study in healthy young women. *Steroids* 2018 **140** 24-31.

42. Faulds MH, Zhao C, Dahlman-Wright K & Gustafsson JA. The diversity of sex steroid action: regulation of metabolism by estrogen signaling. *Journal of Endocrinology* 2012 **212** 3–12.

43. Hetemäki N, Savolainen-Peltonen H, Tikkanen MJ, Wang F, Paatela H, Hämäläinen E, Turpeinen U, Haanpää M, Vihma V & Mikkola TS. Estrogen Metabolism in Abdominal Subcutaneous and Visceral Adipose Tissue in Postmenopausal Women. *The Journal of Clinical Endocrinology & Metabolism* 2017 **102** 4588-4595.

44. Walker BR, Soderberg S, Lindahl B & Olsson T. Independent effects of obesity and cortisol in predicting cardiovascular risk factors in men and women. *Journal of Internal Medicine* 2000 **247** 198–204.

45. Roelfsema F, Pereira AM & Veldhuis JD. Impact of Adiposity and Fat Distribution on the Dynamics of Adrenocorticotropin and Cortisol Rhythms. *Current Obesity Report* 2014 **3** 387-395.

46. Björntorp P & Rosmond R. Obesity and cortisol. Nutrition 2000 16 924-936.

48. Verdonk SJE, Vesper HW, Martens F, Sluss PM, Hillebrand JJ & Heijboer AC. Estradiol reference intervals in women during the menstrual cycle, postmenopausal women and men using an LC-MS/MS method. *Clinica Chimica Acta* 2019 **495** 198-204.

49. Bui HN, Sluss PM, Blincko S, Knol DL, Blankenstein MA & Heijboer AC. Dynamics of serum testosterone during the menstrual cycle evaluated by daily measurements with an ID-LC-MS/MS method and a 2nd generation automated immunoassay. *Steroids* 2013 **78** 96-101.

50. Kushnir MM, Rockwood AL, Roberts WL, Pattison EG, Owen WE, Bunker AM & Meikle AW. Development and performance evaluation of a tandem mass spectrometry assay for 4 adrenal steroids. *Clinical Chemistry* 2006 **52** 1559–1567.

52. Parikh TP, Stolze B, Ozarda Y, Jonklaas J, Welsh K, Masika L, Hill M, DeCherney A & Soldin SJ. Diurnal variation of steroid hormones and their reference intervals using mass spectrometric analysis. *Endocrine Connection* 2018 **7** 1354-1361.

53. Weckesser LJ, Plessow F, Pilhatsch M, Muehlhan M, Kirschbaum C & Miller R. Do venepuncture procedures induce cortisol responses? A review, study, and synthesis for stress research. *Psychoneuroendocrinology* 2014 **46** 88–99.

54. Mezzullo M, Fanelli F, Di Dalmazi G, Fazzini A, Ibarra-Gasparini D, Mastroroberto M, Guidi J, Morselli-Labate AM, Pasquali R, Pagotto U et al. Salivary cortisol and cortisone responses to short-term psychological stress challenge in late adolescent and young women with different hyperandrogenic states. *Psychoneuroendocrinology* 2018 **91** 31-40.

Figure Legend

Figure 1. Distribution of steroid hormone serum levels by age.

Continuous lines: lower and upper reference limits; dashed lines: 90% confidence intervals.

Table 1. Anthropometric, metabolic and hormonal features of the cohort. Data are expressed as mean±SD, except 11-deoxycorticosterone where median (75th centile) are shown. Mean values are back transformed means of the transformed variables. Blood was withdrawn between 8:00–10:00 am after overnight fasting.

		Women in reproductive age						Women in menopausal age					
	All	Normal weight	Overweight	Obese	ANOVA trend, P	All	Normal weight	Overweight	Obese	ANOVA trend, P			
n	186	123	35	28		127	63	49	15				
Age, years	38.2±8.4	37.1±8.6	40.4±8.0	41.0±6.8	0.186	60.0±7.8	58.2±7.4	62.4±8.2	59.4±6.9	0.515	< 0.001		
Body mass index, kg/m ²	21.8±5.3	20.4±1.7	25.8±1.4ª	33.8±3.7 ^{a,b}	< 0.001	23.4±3.9	21.6±1.4	25.8±1.4ª	$32.2{\pm}3.4^{a,b}$	< 0.001	< 0.001		
Waist circumference, cm	81.1±12.8	75.8±6.7	88.2 ± 8.8^{a}	$104.7 \pm 9.2^{a,b}$	< 0.001	86.8±9.8	81.5±7.1	90.0±5.6ª	$104.7{\pm}8.2^{a,b}$	< 0.001	< 0.001		
Systolic blood pressure, mmHg	117.5±14.8	112.5±10.6	125.6±17.2ª	124.6±15.3ª	< 0.001	130.7±17.6	129.1±15.1	129.7±17.7	140.7±22.2	0.027	< 0.001		
Diastolic blood pressure, mmHg	77.5±8.6	74.9±8.1	81.7±7.9°	80.7±8.5ª	< 0.001	82.8±8.3	81.7±7.0	81.7±7.8	$88.8{\pm}11.2^{c,d}$	0.003	< 0.001		
Glucose, mmol/L	4.53±0.73	4.42±0.67	4.45±0.75	5.11±0.74 ^{,d}	< 0.001	4.87±0.80	4.72±0.79	4.93±0.77	5.08 ± 0.88	0.114	< 0.001		
Insulin, µU/mL	6.2±4.0	5.1±2.4	7.1±3.2ª	10.0±5.3 ^{a,e}	< 0.001	7.1±4.1	5.9±2.5	7.7±4.3°	9.5±5.3ª	< 0.001	0.052		
HOMA-IR	1.29±0.95	$1.04{\pm}0.55$	1.34±0.68ª	$2.24{\pm}1.24^{a,b}$	< 0.001	1.55±1.12	1.27±0.65	1.67 ± 1.19^{f}	2.14±1.50°	< 0.001	0.013		
Total cholesterol, mmol/L	4.52±0.82	4.41 ± 0.80	4.79±0.92	4.67±0.73	0.133	5.39±0.84	5.30±0.95	5.50±0.76	5.37±0.58	0.677	< 0.001		
HDL-cholesterol, mmol/L	1.50±0.37	1.57±0.35	1.51±0.31	$1.31{\pm}0.44^{a,d}$	< 0.001	1.57±0.39	1.65 ± 0.40	1.53±0.39	1.46±0.31	0.089	0.117		
Triglycerides, mmol/L	$0.70{\pm}0.34$	0.64±0.22	0.80±0.38ª	0.86±0.53ª	< 0.001	0.94±0.47	0.81 ± 0.40	1.10±0.53	0.97±0.30	0.253	< 0.001		
17-Hydroxypregnenolone (17OHP5), nmol/L	5.51±7.22	5.64±7.31	6.97±8.13	3.84±4.97	0.087	3.79±3.77	3.51±2.77	4.38±4.72	3.35±3.48	0.815	< 0.001		
Dehydroepiandrosterone (DHEA), nmol/L	15.50±14.60	15.64±14.20	17.11±14.52	13.16±16.65	0.324	8.90±6.35	8.94±5.30	9.09±7.42	8.19±6.83	0.644	< 0.001		
Progesterone (P4), nmol/L	1.93±0.91	1.91±16.81	2.02±13.23	1.87±11.52	0.364	0.17±0.07	$0.17{\pm}0.04$	$0.22{\pm}0.03$	0.21±0.01	0.067	<0.001§		
17-Hydroxyprogesterone (17OHP4), nmol/L	1.776±1.887	1.873±2.039	1.834±1.566	1.362±1.393	0.142	0.710±0.466	0.631±0.391	0.856 ± 0.541	0.635±0.339	0.968	< 0.001		
11-Deoxycorticosterone (DOC), nmol/L	<0.236 (<0.236)	<0.236 (<0.236)	<0.236 (<0.236)	<0.236 (<0.236)	0.133#	<0.236 (<0.236)	<0.236 (<0.236)	<0.236 (<0.236)	<0.236 (<0.236)	0.550#	0.044§		
Corticosterone (B), nmol/L	8.15±9.56	8.94±9.74	8.61±10.50	$5.07 \pm 5.83^{b,c}$	< 0.001	8.49±7.14	7.79±7.29	10.31±7.29	6.35±4.16	0.313	0.662		
11-Deoxycortisol (11S), nmol/L	0.755±0.710	0.743 ± 0.743	0.956±0.689	0.608±0.517	0.338	0.892 ± 0.605	0.817±0.559	1.002 ± 0.655	0.892 ± 0.592	0.613	0.029		
Cortisol (F), nmol/L	286.4±105.8	296.3±102.2	292.3±118.7	$241.1 \pm 94.7^{b,f}$	0.008	318.9±101.9	302.3±93.7	351.5±104.2	283.5±104.8	0.469	0.007		
Androstenedione (A4), nmol/L	2.34±1.09	2.40±1.09	2.48±1.21	1.96±0.85	0.369	1.14±0.54	1.06 ± 0.50	1.21±0.53	1.27±0.67	0.165	< 0.001		
Testosterone (T), nmol/L	0.768 ± 0.328	0.790±0.326	0.730±0.325	0.717±0.342	0.154	0.596±0.309	0.552±0.256	0.684 ± 0.368	0.524±0.219	0.700	< 0.001		
Dihydrotestosterone (DHT), nmol/L	0.183 ± 0.128	0.193±0.124	0.191±0.155	$0.151{\pm}0.079^{\rm f}$	0.006	0.140 ± 0.080	$0.142{\pm}0.063$	0.139 ± 0.076	0.139±0.154	0.597	< 0.001		
Estrone (E1), nmol/L	0.212±0.116	0.216±0.118	0.201±0.122	0.209±0.099	0.319	0.077±0.037	0.069±0.032	$0.083{\pm}0.038^{\rm f}$	0.098±0.044°	0.002	< 0.001		
Estradiol (E2), nmol/L	0.292±0.277	0.323±0.274	0.238±0.300	0.251±0.246	0.334	0.037±0.022	0.037±0.028	0.036±0.008	0.038±0.023	0.640	< 0.001		

HOMA-IR: homeostatic model assessment insulin resistance; HDL: high density lipoprotein. ^a vs NW, P<0.001; ^b vs OW, P<0.001; ^c vs NW, P<0.010; ^d vs OW, P<0.050; ^e vs OW P<0.010; ^f vs NW, P<0.050; [#] Kruskal-Wallis test. [§] Mann-Whitney test; †Reproductive vs Menopausal by T=test

Table 2. Impact of age, anthropometric and metabolic parameters on steroid circulating levels in women in
reproductive and menopausal status. Data are shown as Cohen's effect size for multiple linear regression (f^2) and
P value resulting from the stepwise multiple regression. The negative (-) or positive (+) nature of the relationship
is reported.

Status	Age		BMI		WC		HOM	1A-IR	TC		HDL	-C	TG	
	f^2	Р	f^2	Р	f^2	Р	f^2	Р	f^2	Р	f^2	Р	f^2	Р
17OHP5														
Reproductive	0.12	<0.001 (-)		NS		NS		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS		NS
DHEA														
Reproductive	0.22	<0.001 (-)		NS		NS		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS		NS
P4														
Reproductive*		NS		NS		NS		NS		NS		NS		NS
17OHP4														
Reproductive*		NS	0.22	0.006 (-)		NS		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS	0.06	0.026 (+)
В														
Reproductive		NS		NS	0.20	<0.001 (-)		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS		NS
11S														
Reproductive		NS		NS	0.09	0.002 (-)		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS	0.07	0.011 (+)
F														
Reproductive		NS		NS	0.17	<0.001 (-)		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS	0.05	0.038 (+)
A4														
Reproductive	0.28	<0.001 (-)		NS		NS		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS		NS
TS														
Reproductive	0.22	<0.001 (-)		NS		NS		NS		NS		NS		NS
Menopausal		NS		NS		NS		NS		NS		NS		NS
DHT														
Reproductive	0.11	0.002 (-)	0.09	0.004 (-)		NS		NS		NS	0.08	0.010 (-)		NS
E1														
Reproductive*		NS		NS		NS		NS	0.10	0.033 (-)		NS		NS
Menopausal		NS		NS	0.16	< 0.001 (+)	0.08	0.012 (-)		NS		NS		NS
E2														
Reproductive*		NS		NS		NS		NS		NS		NS	0.15	0.011 (-)

BMI: body mass index; HOMA-IR: homeostatic model assessment insulin resistance; HDL: high density lipoprotein; *NS*, not

significant. *data adjusted by the menstrual phase.

17OHP5, 17-Hydroxypregnenolone; DHEA, Dehydroepiandrosterone; P4, Progesterone; 17OHP4, 17-Hydroxyprogesterone; B, Corticosterone; 11S, 11-Deoxycortisol; F, Cortisol; A4, Androstenedione; TS, testoseterone; DHT, Dihydrotestosterone; E1, Estrone; E2, Estradiol; WC, waist circumference; TC, total cholesterol; HDL-C, HDL cholesterol; TG, triglycerides

Table 3 Reference intervals of	serum androgens and	pro-androgens accordi	ng to age- and menopausal status
	ser ann an a sens ana	pro ununogeno uccorun	

Age (y)/Status	17-Hydroxypregnenolone		Dehydroep	iandrosterone	Androst	enedione	Testos	terone	Dihydrotestosterone		
	(17	OHP5)	(D	HEA)	(A	4)	()	[]		(DHT)	
	LRL (90CI)	URL (90CI)	LRL (90CI)	URL (90CI)	LRL (90CI)	URL (90CI)	LRL (90CI)	URL (90CI)	LRL	URL (90CI)	
20	2.33 (1.65-3.21)	27.65 (17.63-51.44)	8.5 (6.2-11.0)	59.8 (44.8-92.8)	1.29 (1.06-1.50)	4.91 (4.38-5.69)	0.324 (0.234-0.392)	1.646 (1.418-2.086)	ND	ND	
25	1.92 (1.57-2.30)	30.09 (22.51-44.45)	6.6 (5.5-7.7)	60.3 (49.2-81.5)	1.54 (1.36-1.71)	5.66 (5.09-6.48)	0.315 (0.257-0.357)	1.664 (1.514-1.894)	≤0.135	0.535 (0.463-0.604)	
30	1.68 (1.46-1.90)	29.96 (24.07-39.28)	5.5 (4.8-6.1)	57.1 (48.4-71.5)	1.37 (1.23-1.51)	5.28 (4.78-5.92)	0.308 (0.268-0.339)	1.664 (1.549-1.816)	≤0.135	0.508 (0.453-0.575)	
35	1.51 (1.35-1.67)	28.59 (23.88-35.50)	4.6 (4.1-5.1)	52.5 (45.5-62.8)	1.12 (1.01-1.22)	4.64 (4.27-5.10)	0.301 (0.271-0.327)	1.652 (1.546-1.790)	≤0.135	0.472 (0.423-0.539)	
40	1.38 (1.24-1.52)	26.70 (22.74-32.58)	4.0 (3.6-4.4)	47.5 (41.9-55.4)	0.88 (0.80-0.95)	4.03 (3.76-4.37)	0.294 (0.267-0.316)	1.630 (1.523-1.773)	≤0.135	0.434 (0.391-0.496)	
45	1.27 (1.13-1.41)	24.65 (21.11-30.00)	3.5 (3.1-3.9)	42.6 (37.9-48.9)	0.69 (0.62-0.75)	3.52 (3.31-3.79)	0.285 (0.259-0.307)	1.597 (1.485-1.747)	≤0.135	0.396 (0.358-0.449)	
50	1.18 (1.03-1.32)	22.59 (19.30-27.76)	3.1 (2.7-3.5)	38.0 (33.9-43.3)	0.54 (0.46-0.60)	3.12 (2.92-3.36)	0.275 (0.247-0.298)	1.556 (1.430-1.712)	≤0.135	0.357 (0.325-0.406)	
55	1.10 (0.94-1.25)	20.64 (17.41-25.59)	2.7 (2.3-3.1)	33.8 (30.1-38.7)	0.42 (0.33-0.50)	2.81 (2.59-3.05)	0.264 (0.232-0.290)	1.505 (1.357-1.679)	≤0.135	0.317 (0.288-0.371)	
60	1.03 (0.87-1.18)	18.81 (15.54-23.64)	2.4 (2.0-2.8)	30.0 (26.5-34.5)	0.33 (0.22-0.41)	2.58 (2.31-2.85)	0.251 (0.214-0.284)	1.446 (1.269-1.649)	≤0.135	0.277 (0.245-0.342)	
65	0.96 (0.80-1.12)	17.14 (13.87-21.86)	2.1 (1.8-2.6)	26.5 (23.2-31.1)	0.26 (0.15-0.34)	2.41 (2.08-2.74)	0.236 (0.191-0.277)	1.377 (1.168-1.614)	≤0.135	0.234 (0.194-0.312)	
70	0.90 (0.74-1.07)	15.61 (12.38-20.18)	1.9 (1.5-2.3)	23.5 (20.1-27.9)	0.20 (0.09-0.29)	2.29 (1.90-2.69)	0.221 (0.168-0.272)	1.301 (1.055-1.576)	≤0.135	0.189 (0.135-0.281)	
75	0.85 (0.68-1.02)	14.22 (11.05-18.71)	1.7 (1.3-2.1)	20.8 (17.5-25.2)	0.15 (0.05-0.25)	2.22 (1.74-2.71)	0.203 (0.144-0.266)	1.217 (0.938-1.544)	≤0.135	0.141 (0.070-0.247)	
Menopausal	1.15 (0.99-1.33)	16.06 (13.15-19.69)	2.7 (2.3-3.2)	27.8 (24.0-32.1)	0.39 (0.32-0.46)	2.45 (2.25-2.66)	0.248 (0.222-0.276)	1.444 (1.280-1.629)	≤0.135*	0.462*	

LRL: lower reference limit; 90CI: 90% confidence interval; URL: upper reference limit; ND: not determined because of insufficient data points. Data are reported in nmol/L. LRL and URL were calculated as the mean - 1.96xSD and mean + 1.96xSD of hormone distribution, respectively, according to the fractional polynomial regression by Royston and Wright. * LRL and URL calculated as the 2.5th and 97.5th of hormone distribution, respectively.

the calculated as the 2.5° and 77.5° of normone distribution, respectively.

Steroid hormone	Status	Reference sub-cohort			Dysmetabolic sub-cohort						
		Features	n	LRL (90CI) (nmol/L)	URL (90CI) (nmol/L)	Features	n	LL (90CI) (nmol/L)	P value [§]	UL (90CI) (nmol/L)	P value [#]
11-Deoxycorticosterone (DOC)*	Reproductive	all	186	<0.236 (<0.236)	0.404 (0.352-0.747)						
	Menopausal	all	127	<0.236 (<0.236)	0.308						
Corticosterone (B)	Reproductive	norWC	132	1.99 (1.65-2.40)	41.87 (34.31-51.14)	dysWC	52	1.39 (1.03-1.89)	0.102	31.16 (22.52-43.21)	0.204
	Menopausal	all	127	1.78 (1.40-2.25)	30.42 (26.28-35.09)						
11-Deoxycortisol (11S)	Reproductive	norWC	132	0.197 (0.168-0.231)	3.277 (2.681-4.020)	dysWC	52	0.212 (0.168-0.269)	0.667	2.891 (2.153-3.911)	0.568
	Menopausal	norTG	112	0.231 (0.186-0.284)	2.679 (2.334-3.065)	dysTG	10	0.574 (0.370-0.862)	0.003	2.666 (1.919-3.635)	0.981
Cortisol (F)	Reproductive	norWC	132	148.2 (135.3-162.2)	566.1 (521.6-611.8)	dysWC	52	112.7 (94.8-133.5)	0.020	539.6 (465.1-624.9)	0.640
	Menopausal	norTG	112	158.7 (142.3-176.2)	548.9 (511.7-587.8)	dysTG	10	199.3 (137.4-276.9)	0.319	649.7 (510.9-811.0)	0.261
Dihydrotestosterone (DHT)	Reproductive	NW and norHDL	65	0.107 (0.093-0.123)	0.618 (0.513-0.749)	OW/OB or dysHDL	76	0.093 (0.081-0.106)	0.235	0.553 (0.461-0.666)	0.484
						OW/OB only	40	0.087 (0.072-0.107)	0.178	0.553 (0.426-0.727)	0.578
Progesterone (P4)*	Menopausal	all	127	< 0.16	0.28						
17-Hydroxyprogesterone (17OHP4)	Menopausal	norTG	112	0.243 (0.204-0.287)	1.937 (1.711-2.186)	dysTG	10	0.301 (0.150-0.564)	0.613	3.001 (1.862-4.677)	0.141
Estrone (E1)	Menopausal	norWC and norHOMA	50	0.032 (0.027-0.038)	0.162 (0.137-0.192)	dysWC or dysHOMA	54	0.039 (0.033-0.045)	0.201	0.193 (0.164-0.227)	0.220
						dysWC only	29	0.039 (0.031-0.050)	0.279	0.225 (0.177-0.286)	0.066
Estradiol (E2)*	Menopausal	all	127	<0.036 (<0.036)	0.088 (0.037-0.252)						

Table 4. Reference intervals of age-independent steroid circulating levels in the reference sub-cohorts, and comparison with limits in dysmetabolic sub-cohorts.

LRL: lower reference limit; 90CI: 90% confidence interval; URL: upper reference limit; norWC: waist circumference \leq 88 cm; dysWC: waist circumference \geq 88cm; norTG: triglyceride level \leq 1.69 mmol/L; dysTG: triglyceride level \geq 1.69 mmol/L; NW: normal weight; norHDL: high density lipoprotein cholesterol level \geq 1.29 mmol/L; OW/OB: overweight/obese; dysHDL: high density lipoprotein cholesterol level \leq 1.29 mmol/L; OW/OB: overweight/obese; dysHDL: high density lipoprotein cholesterol level \leq 1.29 mmol/L; OW/OB: overweight/obese; dysHDL: high density lipoprotein cholesterol level \leq 1.29 mmol/L; orHOMA: homeostatic model assessment insulin resistance \leq 2.5; dysHOMA: homeostatic model assessment insulin resistance \geq 2.5. LRL and URL were calculated as the mean - 1.96xSD and mean + 1.96xSD of hormone distribution, respectively. * LRL and URL calculated as the 2.5th and 97.5th of hormone distribution, respectively. *P* values refer to Z-test comparison of LRL (*) between dysmetabolic and reference subgroups.

Table 5. Upper and lower reference limits of serum steroid levels varying with the menstrual pha

Steroid hormone	Reference		Early follicular (day 1-6)			Pre-ovulator	ry (day 9-13)	Mid-luteal (day 18-24) [#]		
	cohort features	n	LRL (90CI)	URL (90CI)	n	LRL (90CI)	URL (90CI)	n	LRL (90CI)	URL (90CI)
Progesterone (P4)*	all	31	<0.16	1.41	30	<0.16	7.66	22	13.09	82.14
17-Hydroxyprogesterone (17OHP4)	NW	22	0.405 (0.301-0.538)	2.336 (1.788-3.042)	26	0.447 (0.334-0.608)	5.403 (3.536-8.501)	17	2.855 (2.028-3.693)	7.612 (6.742-8.488)
Estrone (E1)	norTC	20	0.069 (0.057-0.086)	0.367 (0.256-0.556)	25	0.091 (0.021-0.158)	0.516 (0.456-0.576)	17	0.131 (0.088-0.182)	0.523 (0.433-0.622)
Estradiol (E2)	norTG	25	0.046 (0.032-0.065)	0.751 (0.464-1.254)	28	0.074 (0.010 -0.171)	1.273 (1.048-1.513)	21	0.198 (0.132-0.274)	0.827 (0.701-0.961)

LRL: lower reference limit; 90CI: 90% confidence interval; URL: upper reference limit. Data are reported in nmol/L. NW: normal weight; norTC: total cholesterol levels <5.17 mmol/L; norTG:

triglyceride level <5.17 mmol/L. LRL and URL were calculated as the mean - 1.96xSD and mean + 1.96xSD of hormone distribution, respectively. # Women showing P4 levels <10 nmol/L were excluded. * LRL and URL calculated as the 2.5th and 97.5th of hormone distribution, respectively.

Page 25 of 27Accepted Manuscript published as EJE-20-1147.R1. Accepted for publication: 12-Jan-2021Figure 1. Distribution of steroid hormone serum levels by age.



Continuous lines: lower and upper reference limits; dashed lines: 90% confidence intervals.

Steroid analyte	Abbreviation	Sensitivity	ity Intra- Inter- T		Trueness	Accuracy vs	Assay	Ref.
		limit	limit assay a		%	certified		
		(nmol/L)	CV%	CV%		QC*		
17-Hydroxypregnenolone	17OHP5	0.117	3 – 4	7 – 9	89 - 114		2	21
Dehydroepiadrosterone	DHEA	2.71	7 - 8	8 - 10	95 - 102		1	18
Progesterone	P4	0.156	5 - 8	6 – 11		84 - 92	1	18
17-Hydroxyprogesterone	17OHP4	0.236	4 - 5	5 – 9		101 - 104	1	18
11-Deoxycorticosterone	DOC	0.236	5 - 6	6 – 9	100 - 104		1	18
Corticosterone	В	0.903	2 - 6	5 - 10	92 - 98		1	18
11-Deoxycortisol	11S	0.226	3 - 8	2 - 8	99 – 106		1	18
Cortisol	F	0.673	2 - 3	5 - 8		94 - 104	1	18
Androstenedione	A4	0.136	7 - 10	10 - 11	86 - 101		1	18
Testosterone	Т	0.066	3 - 4	4 - 7		97 - 100	1	18
Dihydrotestosterone	DHT	0.134	4 - 6	3 – 9	81 - 112		2	21
Estrone	E1	0.018	2 - 3	3 – 9	83 - 111		2	21
Estradiol	E2	0.036	3 – 6	5 - 7	84 - 113	92 - 108	2	21

Supplemental Table 1. Functional sensitivity limits by the LC-MS/MS assays used in the present study.

* Quality control materials provided the Reference Institute for Bioanalytics.

The absence of interference was evaluated from prednisone, prednisolone, triamcinolone acetonide, methylprednisolone, dexamethasone, betamethasone, cortisone, 21-deoxycortisol, 20α -dihydrocortisone, 20β -dihydrocortisone, epitestosterone, DHEA-sulfate, 16-hydroxyprogesterone, 11-hydroxyprogesterone, 17-hydroxypregnenolone and pregnenolone for assay 1, and from cortisol, DHEA, testosterone, epitestosterone, pregnenolone, progesterone, 17-hydroxyprogesterone, 17-hydroxyprogestero

Supplemental Table 2. Anthropometric, metabolic and hormonal features of women sub-classified according

to the menstrual phase.

	Early follicular (n=31)	Pre-ovulatory (n=30)	Mid-luteal (n=33)	ANOVA, <i>P</i> Value*
Age, years	39.8±8.7	39.2±8.2	38.7±8.3	0.845
Body mass index, kg/m ²	22.8±4.2	22.2±2.4	23.7±6.1	0.217
Waist circumference, cm	80.7±10.4	78.0 ± 8.9	80.5±13.4	0.516
Systolic blood pressure, mmHg	116.3±12.1	112.2±9.0	116.4±19.4	0.497
Diastolic blood pressure, mmHg	77.5±8.0	77.0±7.7	78.0±9.5	0.956
Glucose, mmol/L	4.48±0.83	<mark>4.36±0.74</mark>	4.57±0.51	<mark>0.601</mark>
Insulin, $\mu U/mL$	5.4±2.9	5.0±2.2	6.4±3.6	0.176
HOMA-IR	1.07 ± 0.76	0.99±0.51	$1.34{\pm}0.82$	0.122
Total cholesterol, mmol/L	<mark>4.70±0.89</mark>	<mark>4.34±0.68</mark>	<mark>4.60±0.76</mark>	<u>0.165</u>
HDL-cholesterol, mmol/L	1.51±0.42	1.60±0.28	1.47±0.30	<u>0.453</u>
Triglycerides, mmol/L	0.70±0.38	0.60±0.26	<mark>0.69±0.33</mark>	<u>0.150</u>
17-Hydroxypregnenolone (17OHP5), nmol/L	6.51±9.30	4.89±9.15	6.16±5.52	0.302
Dehydroepiandrosterone (DHEA), nmol/L	18.2±12.2	13.7±12.6	$16.0{\pm}10.1$	0.178
Progesterone (P4), nmol/L	0.36±0.30	0.44±1.65	$11.67 \pm 19.23^{a,b}$	< 0.001
17-Hydroxyprogesterone (17OHP4), nmol/L	0.998 ± 0.661	1.356±1.101	$3.126{\pm}1.967^{a,b}$	< 0.001
11-Deoxycorticosterone (DOC), nmol/L	<0.236 (<0.236)	<0.236 (<0.236)	<0.236 (<0.236)°	0.043#
Corticosterone (B), nmol/L	8.99 ± 9.89	6.19±9.65	8.92±11.59	0.135
11-Deoxycortisol (11S), nmol/L	0.821±0.736	0.591 ± 0.555	0.878 ± 0.774	0.048
Cortisol (F), nmol/L	300.3±101.7	253.6±107.9	292.3±115.6	0.167
Androstenedione (A4), nmol/L	2.25±1.45	2.22±0.73	$2.44{\pm}0.95$	0.585
Testosterone (T), nmol/L	0.691±0.330	0.755 ± 0.279	0.741±0.334	0.659
Dihydrotestosterone (DHT), nmol/L	0.149±0.131	0.147 ± 0.132	0.135±0.113	0.487
Estrone (E1), nmol/L	$0.149{\pm}0.062$	0.264±0.119ª	0.238±0.119ª	< 0.001
Estradiol (E2), nmol/L	0.179 ± 0.180	$0.482{\pm}0.310^{a}$	0.373±0.269ª	< 0.001

Data are presented as the back-transformed means of the transformed variables \pm SD, except 11-deoxycorticosterone which is shown as median (75° centile). Blood was withdrawn between 8:00–10:00 am after overnight fasting. HOMA-IR: homeostatic model assessment insulin resistance; HDL: high density lipoprotein. ^a vs follicular phase, P<0.001; ^b vs pre-ovulatory phase, P<0.050. [#] Kruskal-Wallis test.