ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Occupancy and activity rhythms of the Siberian roe deer

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:
Emiliano Mori, M.C. (2021). Occupancy and activity rhythms of the Siberian roe deer. BIOLOGIA, 76(10), 2991-2999 [10.1007/s11756-021-00790-1].

Availability:
This version is available at: https://hdl.handle.net/11585/851306 since: 2022-02-02

Published:

DOI: http://doi.org/10.1007/s11756-021-00790-1

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/).
When citing, please refer to the published version.

(Article begins on next page)

09 September 2026

This is the final peer-reviewed accepted manuscript of:

Mori, E., Cicero, M., Lovari, S. *et al.* Occupancy and activity rhythms of the Siberian roe deer. *Biologia* 76, 2991–2999 (2021)

The final published version is available online at: ***https://doi.org/10.1007/s11756-021-00790-1***

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/)

When citing, please refer to the published version.

Occupancy and activity rhythms of the Siberian roe deer

Emiliano Mori[1,*], Maila Cicero[2], Sandro Lovari[3,4], Marco Zaccaroni[5], Silvia Salomoni[6], Andrea Vendramin[7], Claudio Augugliaro[2]

1. Consiglio Nazionale delle Ricerche – Istituto di Ricerca sugli Ecosistemi Terrestri – Via Madonna del Piano 10, 50019, Sesto Fiorentino (FI), Italy. ORCID ID: 0000-0001-8108-795.
2. Wildlife Initiative NGO, Bayangol District, 6th Khoroo, Micro District 10, Ulaanbaatar, 210349, Mongolia. Claudio Augugliaro, ORCID: 0000-0001-9917-709X.
3. Unità di Ricerca in Ecologia Comportamentale, Etologia e Gestione della Fauna– Dipartimento di Scienze della Vita, Università di Siena, Via P.A. Mattioli 4, 53100 Siena, Italy.
4. Maremma Natural History Museum, Strada Corsini, 5, 58100 Grosseto, Italy.
5. Department of Biology, University of Florence, Via Madonna del Piano 8, 50019, Sesto Fiorentino (FI), Italy.
6. World Biodiversity Association Onlus c/o Museo Civico di Storia Naturale, Lungadige Porta Vittoria 9, 37129, Verona, Italy.
7. Università degli studi di Udine - DI4A (Dipartimento di Scienze Agroalimentari, 23 ambientali e animali), via Sondrio 2/A, 33100, Udine, Italy.

*, Corresponding Author. Emiliano Mori, Consiglio Nazionale delle Ricerche – Istituto di Ricerca sugli Ecosistemi Terrestri – Via Madonna del Piano 10, 50019, Sesto Fiorentino (FI), Italy. Email: emiliano.mori@cnr.it

Short title: Activity and occupancy of Siberian roe deer

Abstract

The assessment of spatiotemporal behaviour patterns of wild species is pivotal both for conservation and for management, especially when involving rare or elusive species, or species living in delicate ecosystems, e.g. mountains. The Siberian roe deer *Capreolus pygargus* is a native Asian ungulate, whose ecology is still poorly known, especially on mountain ecosystems. In particular, information on its spatial behaviour and temporal patterns of activity is poor. We have assessed its patterns of circadian rhythms in relation to moon phases, with some conclusions on spatial behaviour in respect to potential predation and slope inclination. Data were collected between August and October 2019, with 35 camera-traps deployed over an area of mountain forests alternated to steppe, in Central Mongolia. Camera trap data were analyzed with occupancy models and kernel smoothers, providing a reliable assessment of the presence of Siberian roe deer, with only 1% of false absence and a very high detection probability. This ungulate showed a bimodal temporal behaviour, with activity peaks at dawn and at early night, irrespective from moon phases. The occurrence of Siberian roe deer was negatively influenced by the presence of large carnivores and by increasing slope steepness.

Keywords: *Capreolus pygargus*; moonlight; mountain ecosystems; predation risk; spatiotemporal behaviour.

Introduction

Assessing spatiotemporal behaviour of wildlife is pivotal to design specific management plans, to preserve rare and elusive species and to manage pest species and biological invasions (Srivathsa et al. 2004; Ancillotto et al. 2018; Merson et al. 2019; Mori et al. 2020a). This research field may be challenging, particularly for those species inhabiting impenetrable habitats and remote geographic areas (e.g. De Luca and Rovero 2006; Botts et al. 2020). Camera

53 traps placed for inventories of medium-large mammals in remote study areas (e.g. Augugliaro
54 et al. 2020; Zaccaroni et al. 2020) should provide a wealth of new information (Pamplin 2013;
55 Edward et al. 2018; Mazzamuto et al. 2019), integrating occurrence data very relevant for
56 poorly known, elusive species (Augugliaro et al. 2019; Viviano et al. 2020).
57 The Siberian roe deer *Capreolus pygargus* Pallas, 1771 is a widespread, elusive ungulate
58 species occurring throughout temperate-cold Eurasia, from Baltic countries to the Far East
59 (Argunov 2013; Argunov and Safronov 2013; Lorenzini et al. 2014; Lovari et al. 2016). The
60 current climatic change has promoted its range expansion towards northern latitudes and upper
61 elevations (Argunov and Safronov 2013), although Siberian roe deer tend to select areas at
62 lower elevations with respect to coexisting deer species (Jiang et al. 2006, 2008; Adhikari et al.
63 2016). Little is known on the ecology and behaviour of the Siberian roe deer (Danilkin et al.
64 1995; Danilkin 1996; Jiang et al. 2008, 2009; Adhikari et al. 2016; Lovari et al. 2016). This
65 species inhabits dense forests and forest-steppes up to 3300 m a.s.l. (when permanent ice is
66 absent: Fig. 1); it tends to feed in open habitats where grass cover is abundant, making seasonal
67 movements of up to 200 km between their wintering and breeding ranges in spring, and back
68 in autumn, in groups of up to 500 individuals (Danilkin et al. 1995; Danilkin 1996; Jiang et al.
69 2008; Lovari et al. 2016). However, where larger-sized deer species are present (e.g. *Cervus*
70 *canadensis* Erxleben, 1777, *Cervus elaphus* Linnaeus, 1758, *Alces alces* Linnaeus, 1758), the
71 Siberian roe deer may use sheltered habitats (i.e., forests) in the surroundings of human
72 settlements (Jiang et al. 2008; 2009). Territoriality has only been confirmed in males, before
73 and during the rut, between mid-July and mid-September (Danilkin 1996; Lovari et al. 2016).
74 In the warm months (May-September), females live with their offspring, whereas males are
75 solitary. Conversely in the cold season (October-April), the Siberian roe deer forms mixed
76 groups of up to 20-30 individuals (Argunov and Safronov 2013; Lovari et al. 2016). There is
77 no information on covariates influencing the occupancy of this species. Furthermore, no data

on circadian rhythms of the Siberian roe deer are available, despite being reported as a mainly nocturnal species, with peaks at dawn and dusk in the warm season, while being active mainly at dawn during the cold season (Danilkin 1996; Lovari et al. 2016; Tang et al. 2019). In our work, we have aimed to help filling these gaps, by investigating the patterns of activity rhythms of the Siberian roe deer in Central Mongolia, in the warm months, and the relevant effect of moonlight. We also estimated the probability of occupancy of the Siberian roe deer in this area, and the minimum number of camera-trap nights required to reliably assess its local absence.

Given the biological requirements of this ungulate (Jiang et al. 2008; Lovari et al. 2016), we expected that occupancy by the Siberian roe deer (*i*) would be influenced by the presence of its main predators (i.e., large carnivores and perhaps the red fox *Vulpes vulpes* Linnaeus, 1758 on fawns) and human / livestock presence; (*ii*) would decrease with increasing slopeness, elevation and distance from forest patches, and that (*iii*) ranging movements of this cervid would be reduced to a minimum in bright full moon nights (Monterroso et al. 2013) to reduce predation risk.

Material and Methods

Study area

Our survey was conducted in the Northern Khangai Nuruu National Park (KNNP) (N 46°898′; E 101°346′) in Orkhon district, Arkhangai province, Mongolia. The sampling area was characterized by mountain forests alternated to steppe, bounded by peaks of over 3000 m a.s.l. in the North and in the West. The mean temperature in the study period was 24.4±3.3°C (mean ± SD), whereas monthly precipitations were 4.9±1.5 mm. The vegetation was stratified following an altitudinal range with the steppe system including Asteraceae, Poaceae, Caryophyllaceae and Fabaceae, up to 2100 m a.s.l. A forest-steppe system dominated by *Larix*

sibirica Ledeb. and *Pinus sibirica* Du Tour, as well as shrubs such as *Lonicera altaica* (Pall.) Gladkova, *Ribes altissimum* (Turcz.), and *Rosa acicularis* Lindl., occurs between 2000-2700 m a.s.l. Above the upper treeline, a stony mountain tundra occurred with artic high-mountain species (Klimek and Starkel 1980); mosses and lichens were present in the few peaks over 3000 m a.s.l. (Brzeźniak 1977).

The area hosts over 30 mammalian species (Zaccaroni et al. 2020), e.g. the Siberian roe deer, the Siberian ibex *Capra sibirica* Pallas, 1776 and the wapiti *Cervus canadensis* among ungulates. The large carnivore guild includes the grey wolf *Canis lupus* Linnaeus, 1758, the wolverine *Gulo gulo* (Linnaeus, 1758) and the Eurasian lynx *Lynx lynx* (Linnaeus, 1758). A high number of small and meso-carnivore species also occurs, i.e. the Pallas's cat *Otocolobus manul* (Pallas, 1776), the Asian badger *Meles leucurus* Hodgson, 1847, the stoat *Mustela erminea* Linnaeus, 1758, the Siberian weasel *Mustela sibirica* Pallas, 1773, the least weasel *Mustela nivalis* Linnaeus, 1766, the steppe polecat *Mustela eversmanii* (Lesson, 1827), the beech marten *Martes foina* (Erxleben, 1777), the red fox and the corsac fox *Vulpes corsac* Linnaeus, 1768 (Zaccaroni et al. 2020). Amongst meso-carnivores, only the red fox has been recorded as potential predators of roe deer fawns (Danilkin 1996; Aanes et al. 1998).

Herder families live in sparse groups of 5-10 households (i.e., "*ger*": Narangerel 2010). Sheep, goats, horses and yaks were bred throughout the area. The small-sized livestock was guarded by dogs and herders, and it was enclosed in corrals at night. Yaks were free ranging in the grazing area, and brought back to the surroundings of the herder's camp in the evening, whereas horses were left free-ranging round-the-clock.

Sampling design

We set 35 camera traps in 35 locations, between August and the end of October 2018, with no intermediate control. Camera traps were grouped in two clusters with respectively 17 and 18 cameras, distant approximately 13 km one another between their closest points (Fig. 2). These two areas were selected as they included the highest number of habitat types in the study area and the highest elevation ranges, from woodlands to alpine meadows, to screes. The minimum distance between 2 cameras was 680 ± 30 m (Rovero and Zimmermann 2016).

Cameras were placed on randomly selected cells of a 1 km x 1 km grid created on QGIS 2.18 (QGIS Development Team 2018), including forest patches and areas over the tree-line level, depending on their local accessibility. We arrayed the cameras on video-mode, with an average height of 50 cm above the ground (Tobler et al. 2008), on animal trails and forced passages, at elevations ranging between 2065 and 2944 m a.s.l.

Occupancy models

We counted as one "independent event" all Siberian roe deer videos taken by the same camera trap in less than 30 min; when > 1 roe deer video was recorded by the same camera trap in ≤ 30 min, we kept in our dataset only one record, placed in the mid-time between the first and the last video (Viviano et al. 2021). We filled a binary string (1, detection of Siberian roe deer; 0, non-detection) for all camera-trap checks for each camera-trap site (MacKenzie et al. 2006). Occupancy models were calculated by using the package "unmarked" (Fische and Chandler 2011) for the software R (version 3.6.1., R Foundation for Statistical Computing, Vienna, Austria: cran.r-project.org). We used the "occu" model-fitting function, which fits occurrence models with no linkage between abundance and detection (Fische and Chandler 2011).

As our field work lasted three months, we ran single-season models (MacKenzie et al. 2006). We included the effects of seven environmental variables used as covariates: (*i*) elevation (metres above sea level); (*ii*) slope (degrees); (*iii*) local presence of large carnivores (i.e. the

154 wolverine, the grey wolf and the lynx: 1, detected; 0, not detected); (*iv*) local presence of the
155 red fox, potential predator of roe deer fawns (1, detected; 0, not detected); (*v*) local presence of
156 livestock as a proxy of human pressure and as food competitors (1, detected; 0, not detected);
157 (*vi*) camera trap located in a forest patch (1, yes; 0, no); (*vii*) aspect (i.e. North-Eastern, North-
158 Western, South-Eastern and South-Western oriented camera traps on mountainsides).
159 We started with a "null" model which does not include covariates for both detectability and
160 occupancy (Mori et al. 2020a). Then, we focused first on the effect of all covariates and their
161 interactions on detectability (p) and, afterwards, on occupancy (ψ) and on both p and ψ, as
162 detectability may affect occupancy, following Luzi et al. (2021). The goodness of each model
163 was assessed through a multi-model inference approach; for each obtained model the AIC_C
164 value was computed. Models were considered as informative and reliable when ΔAIC_C (i.e., the
165 difference between the AIC_C of each model and the lowest AIC_C) was lower than 2 ("top
166 models": Burnham and Anderson 2002; Mori et al. 2020a). We estimated the minimum number
167 of camera-trap days of activity to reliably infer the absence of the Siberian roe deer through the
168 formula: $N = \ln(\alpha)/\ln(1-p)$, where α represents the probability of type-I error (i.e. 0.05: Reed
169 1996).
170
171 *Pattern of circadian activity rhythms*
172
173 We defined the "activity" as the cumulate period which roe deer spend while "non-sleeping",
174 regardless of their behaviour (Lashley et al. 2018). For each Siberian roe deer detection, we
175 reported the date and the solar hour of capture directly shown on each file, in a dataset.
176 Circadian rhythms and associated 95% confidence intervals (hereafter, CIs) were computed
177 through the R 3.6.1 package overlap (Meredith and Ridout 2014). We removed from our dataset
178 records occurring at the same site within less than 30 min, to limit pseudo-replication (Meredith

and Ridout 2014). Each detection kept in the final dataset after this filtering was considered arbitrarily as independent. We tested whether moonlight had an effect on the locomotor activity patterns of the Siberian roe deer by classifying moon phases into four clusters: phase (1) from new moon to ¼; phase (2) from ¼ to ½; phase (3) from ½ to ¾ and phase (4) over ¾ and full moon. Then, we performed a chi-squared test on the numbers of detections recorded during each moon phase, to assess if they were uniform throughout the four clusters (Mori et al. 2020a). We performed a Hermans–Rasson test to evaluate whether a random activity pattern was exhibited over the 24 hours (Landler et al. 2019).

Results

Our final dataset included 55 records of Siberian roe deer from a total of 7/35 camera trap sites (Fig. 1), with 23.6% records in forest edges and 76.4% in forest areas: no roe deer was recorded in steppe and on bare rocks beyond the upper tree line.

We obtained 1% of false absence, i.e. the percentage of sites in which the Siberian roe deer was actually found (naïve estimate=0.20) was only 1% smaller than the estimate of proportion of occupied sites ψ (0.21). The top model obtained ($\Delta AIC_C < 2$: Table 1) assumed that the occupancy by the Siberian roe deer was a function of both slope and presence of large carnivores, with a very high value for the parameter w_i (98%: Table 1). The effect of both covariates was negative (coefficient $\beta \pm$ SE, presence of potential predators = -1.38 ± 0.42; slope = -1.59 ± 0.27), thus indicating that sites where potential predators were not recorded and characterised by a lower degree of declivity were more likely to be occupied by the Siberian roe deer, with respect to those where large carnivores were recorded. The effects of all the other environmental variables included in our models were not significant. The detection probability (p) was high for the best selected model (69%). According to our results, four camera-trap days are required to reliably infer the local absence of the Siberian roe deer.

A non-random activity pattern was exhibited round-the-clock, peaking at dawn and early night (Hermans–Rasson test: R = 74.70; p = 0.02: Fig. 3). Activity of Siberian roe deer appeared independent from moon phases (χ^2 = 0.53, df = 3, p = 0.08).

Discussion

Our models, valid for the warm season only, showed a strong negative relationship between presence and detection of Siberian roe deer and detection of large carnivores, thus fulfilling our prediction (*i*). We are aware of the limitations related to our small sample size, and further data are needed to assess the Siberian roe deer behaviour throughout the whole year.

Although the diets of the grey wolf, the wolverine and the Eurasian lynx have not been assessed in this area (where they occur: Zaccaroni et al. 2020), they are predators whose presence may affect the local population density of European (cf. Odden et al. 2006; Van Dijk 2008; Newsome al. 2016) and that of Siberian (Danilkin 1916; Soyumert et al. 2019; Tan et al. 2019; Balayed-Lyngdoh et al. 2020) roe deer. In China, Tang et al. (2019) showed a low encounter probability between Siberian roe deer and the lynx, suggesting it as a potential predator avoidance tactic by this ungulate and providing support to our results. In contrast, the presence of the red fox (Danilkin 1916; Aanes et al. 1998) was not a significant variable affecting the presence of the Siberian roe deer in Central Mongolia. Fawns of Siberian roe deer are 80% larger than those of the European species and this may prevent foxes to prey easily on them (cf. Hewison and Danilkin 2001, but see Danilkin 1996). Furthermore, the red fox is an inferior competitor of the lynx, and presence of larger predators may lead meso-carnivores to avoid areas and hours where/when the superior predator is present, thus coinciding with roe deer habits (cf. Helldin et al. 2006; Tang et al. 2019, but see Ferretti et al. 2021, and Rossa et al. 2021, as to facilitative relationships between the wolf and the red fox). Our models showed also that an increasing degree of declivity limits the occurrence of the Siberian roe deer, in line with

231 our prediction (*ii*). Conversely, forest patches and elevation were not significant variables,
232 possibly because Siberian roe deer are adapting to higher elevation over the treeline as a
233 consequence of climate change (Ross et al. 2021). Most likely, the European roe deer evolved
234 as a glade inhabitant in forested habitats (Lovari et al. 2017), rarely observed beyond the upper
235 treeline (cf. Robin 1975; Mori et al. 2018; Boitani et al. 2003), as their anatomy is not well-
236 adapted to broken terrain or cliffs (Boitani et al. 2003). Accordingly, Siberian roe deer use
237 mountains only in the warm season, i.e. when snow cover is not present, whereas they move to
238 lowlands in the cold months through fixed migration routes (Danilkin et al. 1995; Danilkin
239 1996). Our occupancy models did not show any importance of forest patches for Siberian roe
240 deer occupancy. In North-Eastern China, the Siberian roe deer is a typical species of mature
241 coniferous forests, avoiding sparse scrublands and bare rock areas which do not provide food
242 nor shelter sites (Guo et al. 2017). Adhikari et al. (2016) showed that, in mountainous areas of
243 Jeju Island (South Korea), Siberian roe deer mostly use open areas to feed in the warm months,
244 i.e. areas rich in forbs-climbers and graminoid plants. Although we did not attempt to estimate
245 habitat-selection in our study area, the Siberian roe deer may only use forests (where most
246 photos were taken) as cover to rest, being also active in areas with sparse trees, where their food
247 is most abundant and time required for vigilance is the lowest (see San Josè et al. 1996; Jiang
248 et al. 2008; Adhikari et al. 2016).
249 Both the European roe deer (Bonnot et al. 2019; Ossi et al. 2020; Viviano et al. 2021; Zanni et
250 al. 2021) and the Siberian roe deer are mainly crepuscular, i.e. their activity locomotor patterns
251 significantly peak at dawn and dusk. Our results confirm the results of Tang et al. (2019), which
252 showed that ranging movements of the Siberian roe deer are defined by heliophany (including
253 sunrise and sunset times). The roe deer seems to shape its activity to limit encounters with its
254 main local predators, e.g. the lynx and the wolf (Soyumert et al. 2019; Tang et al. 2019; Bonnot
255 et al. 2020; Mori et al. 2020b). Bonnot et al. (2020) showed a remarkable plasticity in European

roe deer diel activity in relation to spatiotemporal variation of mortality/predation risks. This flexibility was also suggested for populations of Siberian roe deer (Jiang et al. 2008; Tang et al. 2019). This small cervid species shows a limited capacity of the rumen and a fast digestion, requiring frequent feeding bouts (Cederlund 1981; Danilkin 1996). This feature may explain why roe deer activity patterns do not show a single peak, but two peaks of activity/day alternated with resting in cover. An effect of moon phases was also expected, as most prey species tend to avoid bright moonlight nights to limit encounter probabilities with their predators (Lima 1998; Monterroso et al. 2013). Prugh and Golden (2014) reported that the trend in ungulate activity with respect to moon phases is still poorly studied, and it may change with species and adapt to local ecological conditions. Nocturnal activity of mostly-diurnal species peaks in brightest nights (Carnevali et al. 2016; Brisbane and van den Burg 2020), as their visual acuity is fundamental for orientation. Instead, ungulate species with mostly crepuscular and cathemeral activity may decrease predation risk by moving to cover at night, thus limiting their visibility to predators, without avoidance of bright moonlight (Wronski et al. 2006; Prugh and Golden 2014). Consistently, our results showed that, despite the local presence of several large carnivores, the activity rhythms of Siberian roe deer were independent from moon phases, thus not fulfilling our prediction (*iii*). In fact, Pagon et al. (2013) detected no effect of moon phases on the activity rhythms of the European roe deer. Being active at dawn and dusk might thus represent a strategy to avoid both diurnal and nocturnal local predators (cf. Soyumert et al. 2019; Bonnot et al. 2020; Mori et al. 2020b).

Besides individual behaviour (Stache et al. 2013), human activities (e.g hunting pressure and livestock breeding) may shape the spatiotemporal behaviour of wild ungulates (Douglas 1971; De Boer et al. 2004; Scillitani et al. 2010; Bonnot et al. 2013; Zbyryt et al. 2018). Jiang et al. (2008) reported that Siberian roe deer seem to resent human disturbance, much more than coexisting Manchurian wapitis *Cervus canadensis xanthopygus* Milne-Edwards, 1867,

therefore occurring mostly far from human settlements. Ecological studies on European roe deer have shown that hunting pressure increases vigilance levels, with consequent shifts of feeding sites to cover (Benhaiem et al. 2008) and of activity rhythms (Bonnot et al. 2020). Hunting Siberian roe deer is allowed in Mongolia by the National Law on Hunting of May 2000, between the 1st of August and the 1st of December. However, our study area was included in a National Park, where hunting is forbidden (cf. Art. 18, Law on Special Protected Areas of Mongolia), although some poaching may have occurred. In our study area, the temporal behaviour of the Siberian roe deer coincided with that of the European roe in sites where human disturbance is the lowest (Bonnot et al. 2020; Mori et al. 2020b). In our study area, the presence of a single human was detected only once (Zaccaroni et al. 2020).

Our data refer only to roe deer activity in the warm season, i.e. they cover the birth and mating periods: two rather special biological phases of roe deer (Melis et al. 2005; Bongi et al. 2008; Lovari et al. 2008). Information on activity rhythms in the cold season, with a much longer night-duration, could yield different results. In particular, the activity of large predators seems to occur mainly at night (Soyumert et al. 2019; Tang et al. 2019; Bonnot et al. 2020; Mori et al. 2020b), thus leading potentially to an even greater impact on the activity rhythms of Siberian roe deer, in the cold season.

Acknowledgements

An anonymous reviewer and the Associate Editor Peter Kaňuch greatly improved our first draft with their comments.

Author contributions

CA conceived the study; EM and SL conducted literature search and performed the analysis for the review; MC, MZ, SS, AV and CA collected field data. EM, SL and CA led manuscript writing. All authors contributed critically to the draft and gave final approval for submission. Data supporting this study are available in the main text.

Research funding

No fund was received for this research

Conflict of interest statement

No potential conflict of interest was reported by authors.

Compliance with ethical standard

All procedures were conducted without animal handling and with no disturbance to them, in line with all national and international laws on animal and wildlife wellness.

References

Aanes R, Linnell JDC, Perzanowski K, Karlsen J, Odden J (1998) Roe deer as prey. In: Andersen R, Duncan P, Linnell JDC (Eds) The European roe deer the biology of success. Scandinavian University Press, Oslo, Norway, pp. 138–159.

Adhikari P, Park SM, Kim TW, Lee JW, Kim GR, Han SH, Oh H (2016) Seasonal and altitudinal variation in roe deer (*Capreolus pygargus tianschanicus*) diet on Jeju Island, South Korea. J Asia-Pacific Biodiv 9: 422-428. https://doi.org/10.1016/j.japb.2016.09.001

Ancillotto L, Notomista T, Mori E, Bertolino S, Russo D (2018) Assessment of detection methods and vegetation associations for introduced Finlayson's squirrels (*Callosciurus*

finlaysonii) in Italy. Environm Manage 61: 875-883. https://doi.org/10.1007/s00267-018-1013-x

Argunov AV (2013) Formation of the range of the Siberian roe deer (*Capreolus pygargus*, Cervidae) and its present distribution in Yakutia. Biol Bull 40: 692-697. https://doi.org/10.1134/S1062359013080025

Argunov AV, Safronov VM (2013) Demographic structure of Siberian roe deer (*Capreolus pygargus*, Cervidae) population in Central Yakutia. Russ J Ecol 44: 402-407. https://doi.org/10.1134/S1067413613050032

Augugliaro C, Havmøller RW, Monti IE, Havmøller L, Choikhand J, Lkhagvasuren B (2020) Non-volant mammal inventory of western Mongolian-Manchurian Grassland Ecoregion: a biogeographic crossroad worth preserving. Check List 16: 287. https://doi.org/10.15560/16.2.287

Augugliaro C, Paniccia C, Janchivlamdan C, Monti IE, Boldbaatar T, Munkhtsog B (2019) Mammal inventory in the Mongolian Gobi, with the southeasternmost documented record of the Snow Leopard, *Panthera uncia* (Schreber, 1775), in the country. Check List 15: 565–578. https://doi.org/10.15560/15.4.565

Balayed-Lyngdoh S, Habib B, Shrotriya S (2020) Dietary spectrum in Himalayan wolves: comparative analysis of prey choice in conspecifics across high-elevation rangelands of Asia. J Zool 310: 24-33. https://doi.org/10.1111/jzo.12724

Benhaiem S, Delon M, Lourtet B, Cargnelutti B, Aulagnier S, Hewison AJM, Morellet N, Verheyden H (2008) Hunting increases vigilance levels in roe deer and modifies feeding site selection. Anim Behav 76: 611-618. https://doi.org/10.1016/j.anbehav.2008.03.012

Bongi P, Ciuti S, Grignolio S, Del Frate M, Simi S, Gandelli D, Apollonio M (2008) Anti-predator behaviour, space use and habitat selection in female roe deer during the fawning

season in a wolf area. J Zool 276: 242-251. https://doi.org/10.1111/j.1469-7998.2008.00481.x

Boitani L, Lovari S, Vigna-Taglianti A (2003) Fauna d'Italia. Mammalia III – Carnivora, Artiodactyla. Edizioni Calderini, Bologna, Italy.

Bonnot NC, Morellet N. Verheyden H, Cargnelutti B, Lourtet B, Klein F, Hewison AJM (2013) Habitat use under predation risk: hunting, roads and human dwellings influence the spatial behaviour of roe deer. Eur J Wildl Res 59: 185-193. https://doi.org/10.1007/s10344-012-0665-8

Bonnot NC, Couriot O, Berger A, Cagnacci F, Ciuti S, De Groeve JE, Gehr B, Heurich M, Kjellander P, Kröschel M, Morellet N, Sönnischen L, Hewison AJM (2019) Fear of the dark? Contrasting impacts of humans versus lynx on diel activity of roe deer across Europe. J Anim Ecol 89: 132-145. https://doi.org/10.1111/1365-2656.13161

Botts RT, Eppert AA, Wiegman TJ, Rodriguez A, Blankenship SR, Asselin EM, Garley WM, Wagner AP, Ullrich SE, Allen GR, Mooring MS (2020) Circadian activity patterns of mammalian predators and prey in Costa Rica. J Mammal 101: 1313-1331. https://doi.org/10.1111/1365-2656.13161

Brisbane JL, van den Burg MP (2020) No need for artificial light: nocturnal activity by a diurnal reptile under lunar light. Neotrop Biodivers 6: 193-196. https://doi.org/10.1080/23766808.2020.1844993

Brzeźniak E (1977) Thermal Zones on the Southern Slopes of the Khangai. Bull Acad Pol Sci Sci Terre 25: 211-218.

Carnevali L, Lovari S, Monaco A, Mori E (2016) Nocturnal activity of a "diurnal" species, the northern chamois, in a predator-free Alpine area. Behav Processes 126: 101-107. https://doi.org/10.1016/j.beproc.2016.03.013

382 Cederlund G (1981) Daily and seasonal activity patterns of roe deer in a boreal habitat. Viltrevy
383 Swedish Wildl Res 11: 315-353.
384 Danilkin A (1996) Behavioural ecology of Siberian and European roe deer. Chapman & Hall,
385 London, Great Britain.
386 Danilkin AA, Darman IA, Minaev AN, Sempere A (1995) The social organization, behavior
387 and ecological parameters of a migrating population of Siberian roe deer (*Capreolus*
388 *pygargus* Pall.). Izvestiia Akademii Nauk Seriia Biologischeskaia 1: 48-61.
389 De Boer HY, van Breukelen L, Hootsman MJM, van Wieren SE (2004) Flight distance in roe
390 deer *Capreolus capreolus* and fallow deer *Dama dama* related to hunting and other factors.
391 Wildl Biol 10: 35-41. https://doi.org/10.2981/wlb.2004.007
392 De Luca DW, Rovero F (2006) First records in Tanzania of the vulnerable Jackson's mongoose
393 *Bdeogale jacksoni* (Herpestidae). Oryx 40: 468-471.
394 Douglas MJW (1971) Behaviour responses of red deer and chamois to cessation of hunting.
395 New Zeal J Sci 14: 507-518.
396 Edwards S, Cooper S, Uiseb K, Hayward M, Wachter B, Melzheimer J (2018) Making the most
397 of by-catch data: assessing the feasibility of utilising non-target camera trap data for
398 occupancy modelling of a large felid. Afr J Ecol 56: 885-894.
399 https://doi.org/10.1111/aje.12511
400 Ferretti F, Pacini G, Belardi I, ten Cate B, Sensi M, Oliveira R, Rossa M, Burrini L, Lovari S
401 (2021) Recolonizing wolves and opportunistic foxes: interference or facilitation? Biol J Linn
402 Soc 132: 196-210. https://doi.org/10.1093/biolinnean/blaa139
403 Fiske I, Chandler R (2011) Unmarked: a R package for fitting hierarchical models of wildlife
404 occurrence and abundance. J Stat Soft 43: 1-23.

405 Guo K, Liu H, Bao H, Hu J, Wang S, Zhang W, Zhao W, Jiang G (2017) Habitat selection and
406 their interspecific interactions for mammal assemblage in the Greater Khingan Mountains,
407 northeastern China. Wildl Biol 4: 1-8. https://doi.org/10.2981/wlb.00261
408 Helldin JO, Liberg O, Glöersen G (2006) Lynx (*Lynx lynx*) killing red foxes (*Vulpes vulpes*) in
409 Boreal Sweden? Frequency and population effects. J Zool 270: 657-663.
410 https://doi.org/10.1111/j.1469-7998.2006.00172.x
411 Hewison AJM, Danilkin A (2001) Evidence for separate specific status of European (*Capreolus*
412 *capreolus*) and Siberian (*C. pygargus*) roe deer. Mammal Biol 66: 13-21.
413 Jiang G, Jianzhang M, Minghai Z (2006) Spatial distribution of ungulate responses to habitat
414 factors in Wandashan Forest Region, northeastern China. J Wildl Manage 70: 1470-1476.
415 Jiang GS, Zhang MH, Ma JZ (2008) Habitat use and separation between red deer and roe deer
416 in relation to human disturbance in the Wandashan mountains, northeastern China. Wildl
417 Biol 14: 92-100.
418 Jiang G, Ma J, Zhang M, Stott P (2009) Assessing microhabitat use by roe deer and moose in
419 China. Wildl Res 36: 134-142. https://doi.org/10.1071/WR08092
420 Klimek K, Starkel L (1980) Vertical zonality in the Southern Khangai Mountains (Mongolia).
421 Geographical Studies 136, Polish Academy of Sciences, NAUK, Wroclaw, Poland.
422 Landler L, Ruxton GD, Malkemper EP (2019) The Hermans–Rasson test as a powerful
423 alternative to the Rayleigh test for circular statistics in biology. BMC Ecol 19: 1-8.
424 https://doi.org/10.1186/s12898-019-0246-8
425 Lashley MA, Cove MV, Chitwood MC, Penido G, Gardner B, DePerno CS, Moorman CE
426 (2018) Estimating wildlife activity curves: comparison of methods and sample size. Sci Rep
427 8: 4173. https://doi.org/10.1038/s41598-018-22638-6
428 Lima SL (1998) Stress and decision making under the risk of predation: recent developments
429 from behavioural, reproductive, and ecological perspectives. Adv Study Behav 27: 215-290.

Lorenzini R, Garofalo L, Qin X, Voloshina I, Lovari S (2014) Global phylogeography of the genus *Capreolus* (Artiodactyla: Cervidae), a Palaearctic meso-mammal. Zool J Linn Soc 170: 209-221. https://doi.org/10.1111/zoj12091

Lovari S, Bartolommei P, Meschi F, Pezzo F (2008) Going out to mate: excursion behaviour of female roe deer. Ethology 114: 886-896. https://doi.org/10.1111/j.1439-0310.2008.01549.x

Lovari S, Serrao G, Mori E (2017) Woodland features determining home range size of roe deer. Behav Processes 140: 115-120. https://doi.org/10.1016/j.beproc.2017.04.012

Lovari S, Masseti M, Lorenzini R (2016) *Capreolus pygargus*. The IUCN Red List of Threatened Species 2016: https://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T42396A22161884.en. Accessed on 09 October 2020.

Luzi G, Mori E, Puddu G, Zapparoli M (2021) Does the crested porcupine select coppice forest? Habitat preference and activity patterns of a large rodent in the Lago di Vico Natural Reserve. Mammalia https://doi.org/10.1515/mammalia-2020-0143.

MacKenzie DI, Nichols JD, Royle JA, Pollock KH, Bailey LL, Hines JE (2006) Occupancy estimation and modeling. Inferring patterns and dynamics of species occurrence. Elsevier Academic Press Editions, Burlington, Massachusetts, USA.

Mazzamuto MV, Lo Valvo M, Anile S (2019) The value of by-catch data: How species-specific surveys can serve non-target species. Eur J Wildl Res 65: 68. https://doi.org/10.1007/s10344-019-1310-6

Melis C, Cagnacci F, Lovari S (2005) Do male roe deer clump together during the rut? Acta Theriol 50: 253-262.

Meredith M, Ridout M (2014) Overview of the Overlap Package. Available from: http://cran.cs.wwu.edu/web/packages/overlap/vignettes/overlap.pdf. Accessed on 12.12.2020.

454 Merson SD, Dollar LJ, Ta CKW, Macdonald DW (2019) Activity patterns of sympatric living
455 exotic and endemic carnivores (the fosa) in Western Madagascar's deciduous forests. J Zool
456 307: 186-194. https://doi.org/10.1111/jzo.12630
457 Monterroso P, Alves PC, Ferreras P (2013) Catch me if you can: diel activity patterns of
458 mammalian prey and predators. Ethology 119: 1044-1056.
459 https://doi.org/10.1111/eth.12156
460 Mori E, Di Bari P, Coraglia M (2018) Interference between roe deer and Northern chamois in
461 the Italian Alps: are Facebook groups effective data sources? Ethol Ecol Evol 30: 277-284.
462 https://doi.org/10.1080/03949370.2017.1354922
463 Mori E, Sangiovanni G, Corlatti L (2020a) Gimme shelter: the effects of rock and moonlight
464 on occupancy and activity pattern of an endangered rodent, the garden dormouse *Eliomys*
465 *quercinus*. Behav Processes 170: 103999. https://doi.org/10.1016/j.beproc.2019.103999
466 Mori E, Bagnato S, Serroni P, Sangiuliano A, Rotondaro F, Marchianò V, Cascini V, Poerio L,
467 Ferretti F (2020b) Spatiotemporal mechanisms of coexistence in a European mammal
468 community in a protected area of southern Italy. J Zool 310: 232-245.
469 https://doi.org/10.1111/jzo.12743
470 Narangerel Y (2010) Issues in the Policy on Establishing New Communities Among Pastoral
471 Nomadic Herders in Mongolia. MA Thesis. Keio University, Japan.
472 Newsome TM, Boitani L, Chapron G, Ciucci P, Dickman CR, Dellinger JA, Lopez-Bao JV,
473 Peterson RO, Shores CR, Wirsing AJ, Ripple WJ (2016) Food habits of the world's grey
474 wolves. Mammal Rev 46: 255-269. https://doi.org/10.1111/mam.12067
475 Odden J, Linnell JDC, Andersen R (2006) Diet of Eurasian lynx, *Lynx lynx*, in the boreal forest
476 of southeastern Norway: The relative importance of livestock and hares at low roe deer
477 density. Eur J Wildl Res 52: 237-244. https://doi.org/10.1007/s10344-006-0052-4

478 Ossi F, Ranc N, Moorcroft P, Bonanni P, Cagnacci F (2020) Ecological and behavioral drivers
479 of supplemental feeding use by roe deer *Capreolus capreolus* in a peri-urban context.
480 Animals 10: 2088. https://doi.org/10.3390/ani10112088
481 Pagon N, Grignolio S, Pipia A, Bongi A, Bertolucci C, Apollonio M (2013) Seasonal variation
482 of activity patterns in roe deer in a temperate forested area. Chronobiol International 30:
483 772-785. https://doi.org/10.25225/fozo.v66.i4.a9.2017
484 Pamplin FA (2013) Optimising the value of by-catch from *Lynx lynx* camera trap surveys in the
485 Swiss Jura region. MSc thesis in in Applied Ecology and Conservation University of East
486 Anglia, UK.
487 QGIS Development Team (2019) QGIS Geographic Information System. Open Source
488 Geospatial Foundation Project. http://qgis.osgeo.org. Accessed on 20.04.2021.
489 Reed JM (1996) Using statistical probability to increase confidence of inferring species
490 extinction. Conserv Biol 10: 1283-1285. https://doi.org/10.1046/j.1523-
491 1739.1996.10041283.x
492 Robin K (1975) Rauemliche Verscienbungen von markieten Rehen (*Capreolus capreolus* L.)
493 in einem voralpinem Gebiet der Ostchweiz. Z Jagdwiss 21: 145-163.
494 Ross JGB, Peters W, Ossi F, Moorcroft PR, Cordano E, Eccel E, Bianchini F, Ramanzin M,
495 Cagnacci F (2021) Climate change and anthropogenic food manipulation interact in shifting
496 the distribution of a large herbivore at its altitudinal range limit. Sci Rep 11: 1-12.
497 https://doi.org/10.1038/s41598-021-86720-2
498 Rossa M, Lovari S, Ferretti F (2021) Spatiotemporal patterns of wolf, mesocarnivores and prey
499 in a Mediterranean area. Behav Ecol Sociobiol 75:32. https://doi.org/10.1007/s00265-020-
500 02956-4
501 Rovero F, Zimmermann F (2016) Camera trapping for wildlife research. Pelagic Publishing
502 Ltd, Exeter, UK.

503 San Josè C, Lovari S, Ferrari N (1996) Temporal evolution of vigilance in roe deer. Behav
504 Processes 38: 155-159.
505 Scillitani L, Monaco A, Toso A (2010) Do intensive drive hunts affect wild boar (*Sus scrofa*)
506 spatial behaviour in Italy? Some evidences and management implications. Eur J Wildl Res
507 56: 307-318. https://doi.org/10.1007/s10344-009-0314-z
508 Soyumert A, Ertürk A, Tavşanoğlu Ç (2019) The importance of lagomorphs for the Eurasian
509 lynx in Western Asia: results from a large scale camera-trapping survey in Turkey. Mammal
510 Biol 95: 18-25. https://doi.org/10.1016/j.mambio.2019.01.003
511 Srivathsa A, Karanth KK, Jathanna D, Kumar NS, Karanth KU (2014) On a dhole trail:
512 examining ecological and anthropogenic correlates of dhole habitat occupancy in the
513 Western Ghats of India. PloS One 9: e98803. https://doi.org/10.1371/journal.pone.0098803
514 Stache A, Heller E, Hothorn T, Heurich M (2013) Activity patterns of roe deer (*Capreolus*
515 *capreolus*) are strongly influenced by individual behaviour. Folia Zool 62: 67-75.
516 https://doi.org/10.25225/fozo.v62.i1.a10.2013
517 Tan S, Jiang X, Wang F, Zhang G, Zhang J, Men H, Han Y, Bao W (2019). Preliminary analysis
518 on the diet composition of the Eurasian lynx at Saihanwula National Nature Reserve. Chin
519 J Zool 54: 151-158.
520 Tang X, Tang S, Li X, Menghe D, Bao W, Xiang C, Gao F, Bao W (2019) A study of population
521 size and activity patterns and their relationship to the prey species of the Eurasian lynx using
522 a camera trapping approach. Animals 9: 864. https://doi.org/10.3390/ani9110864
523 Tobler MW, Carrillo-Percastegui SE, Leite Pitman R, Mares R, Powell G (2008) Further notes
524 on the analysis of mammal inventory data collected with camera traps. Animal Cons 11:
525 187–189. https://doi:10.1111/j.1469-1795.2008.00181.x

526 Van Dijk J (2008) Wolverine foraging strategies in a multiple-use landscape. Ph.D.
527 Dissertation, Norwegian University of Science and Technology Faculty of Natural Sciences
528 and Technology Department of Biology, Trondheim, Norway.
529 Viviano A, Amori G, Luiselli L, Oebel H, Bahleman F, Mori E (2020) Blessing the rains down
530 in Africa: spatiotemporal behaviour of the crested porcupine *Hystrix cristata* (Mammalia:
531 Rodentia) in the rainy and dry seasons, in the African savannah. Tropical Zool 33: 113-124.
532 https://doi.org/10.4081/tz.2020.80
533 Viviano A, Mori E, Fattorini E, Mazza G, Lazzeri L, Panichi A, Strianese L, Mohamed WF
534 (2021) Spatiotemporal overlap between the European brown hare and its potential predators
535 and competitors. Animals 11: 562. https://doi.org/10.3390/ani11020562
536 Wronski T, Apio A, Plath M (2006) Activity patterns of bushbuck (*Tragelaphus scriptus*) in
537 Queen Elizabeth National Park. Behav Processes 73: 333-341.
538 https://doi.org/10.1016/j.beproc.2006.08.003
539 Yan W, Zeng Z, Gong H, Duan Y, Zhao L, Peng A (2020) Locomotor activity patterns of takin
540 (*Budorcas taxicolor*) in a temperate mountain region. Plos One 15: e0235464.
541 https://doi.org/10.1371/journal.pone.0235464
542 Zaccaroni M, Monti IE, Biliotti N, Munkhtsog B, Comand N, De Donà G, Augugliaro C (2020)
543 A preliminary mammal inventory of the Southernmost Mongolian forest: discovery a
544 mesocarnivores heaven. Species 21: 208-221.
545 Zanni M, Brivio F, Grignolio S, Apollonio M (2021) Estimation of spatial and temporal overlap
546 in three ungulate species in a Mediterranean environment. Mammal Res 66: 149-162.
547 https://doi.org/10.1007/s13364-020-00548-1
548 Zbyryt A, Bubnicki JW, Kuijper DP, Dehnhard M, Churski M, Schmidt K (2018) Do wild
549 ungulates experience higher stress with humans than with large carnivores? Behav Ecol 29:
550 19-30. https://doi.org/10.1093/beheco/arx142

Tables

Table 1. Summary of the five parameter estimates for the occupancy. Models are ranked according to the AIC_C, w_i (Akaike weight), ψ (occupancy probability), p (detectability), SE (standard error), N (sample size estimate). The best selected model is shown in bold.

Model	AIC_C	ΔAIC_C	w_i	ψ (SE)	p (SE)	N
ψ ("slope + presence of large carnivores")p(".")	**307.30**	**0.00**	**0.98**	**0.21 (0.07)**	**0.74 (0.12)**	**4**
ψ ("presence of large carnivores")p("elevation")	317.54	10.24	0.01	0.11 (0.03)	0.32 (0.21)	4
ψ ("slope")p(".")	338.17	12.12	0.01	0.10 (0.03)	0.30 (0.17)	5
ψ (".")p(".")	347.22	15.04	0.01	0.12 (0.02)	0.31 (0.24)	5

560 **Figure Legends**

561

562 **Figure 1.** Camera-trapped Siberian roe deer and environmental features of the study area.

563

564 **Figure 2**. Location of the study area (inset map), location of camera traps and occurrences of
565 Siberian roe deer.

566

567 **Figure 3**. Circadian rhythms of the Siberian roe deer, expressed as Kernel density estimate of
568 activity throughout the 24-h cycle (N = 51 independent records). Blue lines represent
569 bootstrapped estimates of activity patterns; dashed black lines represent 95 % CIs.

570