

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Occupancy and activity rhythms of the Siberian roe deer

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version: Occupancy and activity rhythms of the Siberian roe deer / Emiliano Mori, Maila Cicero, Sandro Lovari, Marco Zaccaroni, Silvia Salomoni, Andrea Vendramin, Claudio Augugliaro. - In: BIOLOGIA. - ISSN 1336-9563. - ELETTRONICO. - 76:10(2021), pp. 2991-2999. [10.1007/s11756-021-00790-1]

This version is available at: https://hdl.handle.net/11585/851306 since: 2022-02-02 Published: DOI: http://doi.org/10.1007/s11756-021-00790-1

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version. This is the final peer-reviewed accepted manuscript of:

Mori, E., Cicero, M., Lovari, S. *et al.* Occupancy and activity rhythms of the Siberian roe deer. *Biologia* 76, 2991–2999 (2021)

The final published version is available online at: <u>https://doi.org/10.1007/s11756-</u> 021-00790-1

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

1	Occupancy and activity rhythms of the Siberian roe deer
2	
3	Emiliano Mori ^{1,*} , Maila Cicero ² , Sandro Lovari ^{3,4} , Marco Zaccaroni ⁵ , Silvia Salomoni ⁶ , Andrea
4	Vendramin ⁷ , Claudio Augugliaro ²
5	
6	1. Consiglio Nazionale delle Ricerche – Istituto di Ricerca sugli Ecosistemi Terrestri – Via
7	Madonna del Piano 10, 50019, Sesto Fiorentino (FI), Italy. ORCID ID: 0000-0001-
8	8108-795.
9	2. Wildlife Initiative NGO, Bayangol District, 6 th Khoroo, Micro District 10, Ulaanbaatar,
10	210349, Mongolia. Claudio Augugliaro, ORCID: 0000-0001-9917-709X.
11	3. Unità di Ricerca in Ecologia Comportamentale, Etologia e Gestione della Fauna-
12	Dipartimento di Scienze della Vita, Università di Siena, Via P.A. Mattioli 4, 53100
13	Siena, Italy.
14	4. Maremma Natural History Museum, Strada Corsini, 5, 58100 Grosseto, Italy.
15	5. Department of Biology, University of Florence, Via Madonna del Piano 8, 50019, Sesto
16	Fiorentino (FI), Italy.
17	6. World Biodiversity Association Onlus c/o Museo Civico di Storia Naturale, Lungadige
18	Porta Vittoria 9, 37129, Verona, Italy.
19	7. Università degli studi di Udine - DI4A (Dipartimento di Scienze Agroalimentari, 23
20	ambientali e animali), via Sondrio 2/A, 33100, Udine, Italy.
21	*, Corresponding Author. Emiliano Mori, Consiglio Nazionale delle Ricerche – Istituto di
22	Ricerca sugli Ecosistemi Terrestri – Via Madonna del Piano 10, 50019, Sesto Fiorentino (FI),
23	Italy. Email: emiliano.mori@cnr.it
24	
25	Short title: Activity and occupancy of Siberian roe deer

- 26 Abstract
- 27

The assessment of spatiotemporal behaviour patterns of wild species is pivotal both for 28 29 conservation and for management, especially when involving rare or elusive species, or species living in delicate ecosystems, e.g. mountains. The Siberian roe deer Capreolus pygargus is a 30 native Asian ungulate, whose ecology is still poorly known, especially on mountain ecosystems. 31 32 In particular, information on its spatial behaviour and temporal patterns of activity is poor. We have assessed its patterns of circadian rhythms in relation to moon phases, with some 33 34 conclusions on spatial behaviour in respect to potential predation and slope inclination. Data 35 were collected between August and October 2019, with 35 camera-traps deployed over an area of mountain forests alternated to steppe, in Central Mongolia. Camera trap data were analyzed 36 37 with occupancy models and kernel smoothers, providing a reliable assessment of the presence of Siberian roe deer, with only 1% of false absence and a very high detection probability. This 38 ungulate showed a bimodal temporal behaviour, with activity peaks at dawn and at early night, 39 irrespective from moon phases. The occurrence of Siberian roe deer was negatively influenced 40 by the presence of large carnivores and by increasing slope steepness. 41

42

43 Keywords: *Capreolus pygargus*; moonlight; mountain ecosystems; predation risk;
44 spatiotemporal behaviour.

45

46 Introduction

47

Assessing spatiotemporal behaviour of wildlife is pivotal to design specific management plans, to preserve rare and elusive species and to manage pest species and biological invasions (Srivathsa et al. 2004; Ancillotto et al. 2018; Merson et al. 2019; Mori et al. 2020a). This research field may be challenging, particularly for those species inhabiting impenetrable habitats and remote geographic areas (e.g. De Luca and Rovero 2006; Botts et al. 2020). Camera traps placed for inventories of medium-large mammals in remote study areas (e.g. Augugliaro
et al. 2020; Zaccaroni et al. 2020) should provide a wealth of new information (Pamplin 2013;
Edward et al. 2018; Mazzamuto et al. 2019), integrating occurrence data very relevant for
poorly known, elusive species (Augugliaro et al. 2019; Viviano et al. 2020).

The Siberian roe deer Capreolus pygargus Pallas, 1771 is a widespread, elusive ungulate 57 species occurring throughout temperate-cold Eurasia, from Baltic countries to the Far East 58 (Argunov 2013; Argunov and Safronov 2013; Lorenzini et al. 2014; Lovari et al. 2016). The 59 current climatic change has promoted its range expansion towards northern latitudes and upper 60 elevations (Argunov and Safronov 2013), although Siberian roe deer tend to select areas at 61 62 lower elevations with respect to coexisting deer species (Jiang et al. 2006, 2008; Adhikari et al. 63 2016). Little is known on the ecology and behaviour of the Siberian roe deer (Danilkin et al. 1995; Danilkin 1996; Jiang et al. 2008, 2009; Adhikari et al. 2016; Lovari et al. 2016). This 64 65 species inhabits dense forests and forest-steppes up to 3300 m a.s.l. (when permanent ice is absent: Fig. 1); it tends to feed in open habitats where grass cover is abundant, making seasonal 66 movements of up to 200 km between their wintering and breeding ranges in spring, and back 67 in autumn, in groups of up to 500 individuals (Danilkin et al. 1995; Danilkin 1996; Jiang et al. 68 2008; Lovari et al. 2016). However, where larger-sized deer species are present (e.g. Cervus 69 70 canadensis Erxleben, 1777, Cervus elaphus Linnaeus, 1758, Alces alces Linnaeus, 1758), the Siberian roe deer may use sheltered habitats (i.e., forests) in the surroundings of human 71 settlements (Jiang et al. 2008; 2009). Territoriality has only been confirmed in males, before 72 73 and during the rut, between mid-July and mid-September (Danilkin 1996; Lovari et al. 2016). In the warm months (May-September), females live with their offspring, whereas males are 74 75 solitary. Conversely in the cold season (October-April), the Siberian roe deer forms mixed groups of up to 20-30 individuals (Argunov and Safronov 2013; Lovari et al. 2016). There is 76 no information on covariates influencing the occupancy of this species. Furthermore, no data 77

on circadian rhythms of the Siberian roe deer are available, despite being reported as a mainly nocturnal species, with peaks at dawn and dusk in the warm season, while being active mainly at dawn during the cold season (Danilkin 1996; Lovari et al. 2016; Tang et al. 2019). In our work, we have aimed to help filling these gaps, by investigating the patterns of activity rhythms of the Siberian roe deer in Central Mongolia, in the warm months, and the relevant effect of moonlight. We also estimated the probability of occupancy of the Siberian roe deer in this area, and the minimum number of camera-trap nights required to reliably assess its local absence.

Given the biological requirements of this ungulate (Jiang et al. 2008; Lovari et al. 2016), we expected that occupancy by the Siberian roe deer (*i*) would be influenced by the presence of its main predators (i.e., large carnivores and perhaps the red fox *Vulpes vulpes* Linnaeus, 1758 on fawns) and human / livestock presence; (*ii*) would decrease with increasing slopeness, elevation and distance from forest patches, and that (*iii*) ranging movements of this cervid would be reduced to a minimum in bright full moon nights (Monterroso et al. 2013) to reduce predation risk.

92

93 Material and Methods

94

96

Our survey was conducted in the Northern Khangai Nuruu National Park (KNNP) (N 46°898';
E 101°346') in Orkhon district, Arkhangai province, Mongolia. The sampling area was
characterized by mountain forests alternated to steppe, bounded by peaks of over 3000 m a.s.l.
in the North and in the West. The mean temperature in the study period was 24.4±3.3°C (mean
± SD), whereas monthly precipitations were 4.9±1.5 mm. The vegetation was stratified
following an altitudinal range with the steppe system including Asteraceae, Poaceae,
Caryophyllaceae and Fabaceae, up to 2100 m a.s.l. A forest-steppe system dominated by *Larix*

⁹⁵ *Study area*

sibirica Ledeb. and *Pinus sibirica* Du Tour, as well as shrubs such as *Lonicera altaica* (Pall.)
Gladkova, *Ribes altissimum* (Turcz.), and *Rosa acicularis* Lindl., occurs between 2000-2700 m
a.s.l. Above the upper treeline, a stony mountain tundra occurred with artic high-mountain
species (Klimek and Starkel 1980); mosses and lichens were present in the few peaks over 3000
m a.s.l. (Brzeźniak 1977).

The area hosts over 30 mammalian species (Zaccaroni et al. 2020), e.g. the Siberian roe deer, 109 the Siberian ibex Capra sibirica Pallas, 1776 and the wapiti Cervus canadensis among 110 ungulates. The large carnivore guild includes the grey wolf *Canis lupus* Linnaeus, 1758, the 111 wolverine Gulo gulo (Linnaeus, 1758) and the Eurasian lynx Lynx lynx (Linnaeus, 1758). A 112 113 high number of small and meso-carnivore species also occurs, i.e. the Pallas's cat Otocolobus 114 manul (Pallas, 1776), the Asian badger Meles leucurus Hodgson, 1847, the stoat Mustela erminea Linnaeus, 1758, the Siberian weasel Mustela sibirica Pallas, 1773, the least weasel 115 Mustela nivalis Linnaeus, 1766, the steppe polecat Mustela eversmanii (Lesson, 1827), the 116 beech marten Martes foina (Erxleben, 1777), the red fox and the corsac fox Vulpes corsac 117 Linnaeus, 1768 (Zaccaroni et al. 2020). Amongst meso-carnivores, only the red fox has been 118 recorded as potential predators of roe deer fawns (Danilkin 1996; Aanes et al. 1998). 119

Herder families live in sparse groups of 5-10 households (i.e., "*ger*": Narangerel 2010). Sheep, goats, horses and yaks were bred throughout the area. The small-sized livestock was guarded by dogs and herders, and it was enclosed in corrals at night. Yaks were free ranging in the grazing area, and brought back to the surroundings of the herder's camp in the evening, whereas horses were left free-ranging round-the-clock.

125

126 Sampling design

We set 35 camera traps in 35 locations, between August and the end of October 2018, with no intermediate control. Camera traps were grouped in two clusters with respectively 17 and 18 cameras, distant approximately 13 km one another between their closest points (Fig. 2). These two areas were selected as they included the highest number of habitat types in the study area and the highest elevation ranges, from woodlands to alpine meadows, to screes. The minimum distance between 2 cameras was 680 ± 30 m (Rovero and Zimmermann 2016).

Cameras were placed on randomly selected cells of a 1 km x 1 km grid created on QGIS 2.18 (QGIS Development Team 2018), including forest patches and areas over the tree-line level, depending on their local accessibility. We arrayed the cameras on video-mode, with an average height of 50 cm above the ground (Tobler et al. 2008), on animal trails and forced passages, at elevations ranging between 2065 and 2944 m a.s.l.

139

140 Occupancy models

141

We counted as one "independent event" all Siberian roe deer videos taken by the same camera 142 trap in less than 30 min; when > 1 roe deer video was recorded by the same camera trap in ≤ 30 143 min, we kept in our dataset only one record, placed in the mid-time between the first and the 144 last video (Viviano et al. 2021). We filled a binary string (1, detection of Siberian roe deer; 0, 145 non-detection) for all camera-trap checks for each camera-trap site (MacKenzie et al. 2006). 146 147 Occupancy models were calculated by using the package "unmarked" (Fische and Chandler 2011) for the software R (version 3.6.1., R Foundation for Statistical Computing, Vienna, 148 Austria: cran.r-project.org). We used the "occu" model-fitting function, which fits occurrence 149 models with no linkage between abundance and detection (Fische and Chandler 2011). 150

As our field work lasted three months, we ran single-season models (MacKenzie et al. 2006).
We included the effects of seven environmental variables used as covariates: (*i*) elevation
(metres above sea level); (*ii*) slope (degrees); (*iii*) local presence of large carnivores (i.e. the

wolverine, the grey wolf and the lynx: 1, detected; 0, not detected); (*iv*) local presence of the
red fox, potential predator of roe deer fawns (1, detected; 0, not detected); (*v*) local presence of
livestock as a proxy of human pressure and as food competitors (1, detected; 0, not detected);
(*vi*) camera trap located in a forest patch (1, yes; 0, no); (*vii*) aspect (i.e. North-Eastern, NorthWestern, South-Eastern and South-Western oriented camera traps on mountainsides).

We started with a "null" model which does not include covariates for both detectability and 159 occupancy (Mori et al. 2020a). Then, we focused first on the effect of all covariates and their 160 interactions on detectability (p) and, afterwards, on occupancy (ψ) and on both p and ψ , as 161 162 detectability may affect occupancy, following Luzi et al. (2021). The goodness of each model was assessed through a multi-model inference approach; for each obtained model the AIC_C 163 value was computed. Models were considered as informative and reliable when ΔAIC_C (i.e., the 164 difference between the AIC_C of each model and the lowest AIC_C) was lower than 2 ("top 165 166 models": Burnham and Anderson 2002; Mori et al. 2020a). We estimated the minimum number of camera-trap days of activity to reliably infer the absence of the Siberian roe deer through the 167 168 formula: N = $\ln(\alpha)/\ln(1-p)$, where α represents the probability of type-I error (i.e. 0.05: Reed 1996). 169

170

171 Pattern of circadian activity rhythms

172

We defined the "activity" as the cumulate period which roe deer spend while "non-sleeping", regardless of their behaviour (Lashley et al. 2018). For each Siberian roe deer detection, we reported the date and the solar hour of capture directly shown on each file, in a dataset. Circadian rhythms and associated 95% confidence intervals (hereafter, CIs) were computed through the R 3.6.1 package overlap (Meredith and Ridout 2014). We removed from our dataset records occurring at the same site within less than 30 min, to limit pseudo-replication (Meredith

and Ridout 2014). Each detection kept in the final dataset after this filtering was considered 179 arbitrarily as independent. We tested whether moonlight had an effect on the locomotor activity 180 patterns of the Siberian roe deer by classifying moon phases into four clusters: phase (1) from 181 new moon to $\frac{1}{4}$; phase (2) from $\frac{1}{4}$ to $\frac{1}{2}$; phase (3) from $\frac{1}{2}$ to $\frac{3}{4}$ and phase (4) over $\frac{3}{4}$ and full 182 moon. Then, we performed a chi-squared test on the numbers of detections recorded during 183 each moon phase, to assess if they were uniform throughout the four clusters (Mori et al. 2020a). 184 We performed a Hermans-Rasson test to evaluate whether a random activity pattern was 185 exhibited over the 24 hours (Landler et al. 2019). 186

- 187
- 188 Results
- 189

Our final dataset included 55 records of Siberian roe deer from a total of 7/35 camera trap sites
(Fig. 1), with 23.6% records in forest edges and 76.4% in forest areas: no roe deer was recorded
in steppe and on bare rocks beyond the upper tree line.

193 We obtained 1% of false absence, i.e. the percentage of sites in which the Siberian roe deer was 194 actually found (naïve estimate=0.20) was only 1% smaller than the estimate of proportion of 195 occupied sites ψ (0.21). The top model obtained ($\Delta AIC_C < 2$: Table 1) assumed that the 196 occupancy by the Siberian roe deer was a function of both slope and presence of large 197 carnivores, with a very high value for the parameter w_i (98%: Table 1). The effect of both covariates was negative (coefficient $\beta \pm SE$, presence of potential predators = -1.38 \pm 0.42; 198 slope = -1.59 ± 0.27), thus indicating that sites where potential predators were not recorded and 199 200 characterised by a lower degree of declivity were more likely to be occupied by the Siberian roe deer, with respect to those where large carnivores were recorded. The effects of all the other 201 environmental variables included in our models were not significant. The detection probability 202 203 (p) was high for the best selected model (69%). According to our results, four camera-trap days are required to reliably infer the local absence of the Siberian roe deer. 204

A non-random activity pattern was exhibited round-the-clock, peaking at dawn and early night (Hermans–Rasson test: R = 74.70; p = 0.02: Fig. 3). Activity of Siberian roe deer appeared independent from moon phases ($\chi^2 = 0.53$, df = 3, p = 0.08).

208

209 Discussion

210

Our models, valid for the warm season only, showed a strong negative relationship between presence and detection of Siberian roe deer and detection of large carnivores, thus fulfilling our prediction (*i*). We are aware of the limitations related to our small sample size, and further data are needed to assess the Siberian roe deer behaviour throughout the whole year.

Although the diets of the grey wolf, the wolverine and the Eurasian lynx have not been assessed 215 216 in this area (where they occur: Zaccaroni et al. 2020), they are predators whose presence may affect the local population density of European (cf. Odden et al. 2006; Van Dijk 2008; 217 Newsome al. 2016) and that of Siberian (Danilkin 1916; Soyumert et al. 2019; Tan et al. 2019; 218 Balayed-Lyngdoh et al. 2020) roe deer. In China, Tang et al. (2019) showed a low encounter 219 probability between Siberian roe deer and the lynx, suggesting it as a potential predator 220 avoidance tactic by this ungulate and providing support to our results. In contrast, the presence 221 222 of the red fox (Danilkin 1916; Aanes et al. 1998) was not a significant variable affecting the 223 presence of the Siberian roe deer in Central Mongolia. Fawns of Siberian roe deer are 80% larger than those of the European species and this may prevent foxes to prey easily on them (cf. 224 Hewison and Danilkin 2001, but see Danilkin 1996). Furthermore, the red fox is an inferior 225 226 competitor of the lynx, and presence of larger predators may lead meso-carnivores to avoid areas and hours where/when the superior predator is present, thus coinciding with roe deer 227 habits (cf. Helldin et al. 2006; Tang et al. 2019, but see Ferretti et al. 2021, and Rossa et al. 228 229 2021, as to facilitative relationships between the wolf and the red fox). Our models showed also that an increasing degree of declivity limits the occurrence of the Siberian roe deer, in line with 230

our prediction (ii). Conversely, forest patches and elevation were not significant variables, 231 232 possibly because Siberian roe deer are adapting to higher elevation over the treeline as a consequence of climate change (Ross et al. 2021). Most likely, the European roe deer evolved 233 as a glade inhabitant in forested habitats (Lovari et al. 2017), rarely observed beyond the upper 234 treeline (cf. Robin 1975; Mori et al. 2018; Boitani et al. 2003), as their anatomy is not well-235 adapted to broken terrain or cliffs (Boitani et al. 2003). Accordingly, Siberian roe deer use 236 mountains only in the warm season, i.e. when snow cover is not present, whereas they move to 237 lowlands in the cold months through fixed migration routes (Danilkin et al. 1995; Danilkin 238 1996). Our occupancy models did not show any importance of forest patches for Siberian roe 239 240 deer occupancy. In North-Eastern China, the Siberian roe deer is a typical species of mature 241 coniferous forests, avoiding sparse scrublands and bare rock areas which do not provide food nor shelter sites (Guo et al. 2017). Adhikari et al. (2016) showed that, in mountainous areas of 242 Jeju Island (South Korea), Siberian roe deer mostly use open areas to feed in the warm months, 243 i.e. areas rich in forbs-climbers and graminoid plants. Although we did not attempt to estimate 244 habitat-selection in our study area, the Siberian roe deer may only use forests (where most 245 photos were taken) as cover to rest, being also active in areas with sparse trees, where their food 246 247 is most abundant and time required for vigilance is the lowest (see San Josè et al. 1996; Jiang 248 et al. 2008; Adhikari et al. 2016).

Both the European roe deer (Bonnot et al. 2019; Ossi et al. 2020; Viviano et al. 2021; Zanni et al. 2021) and the Siberian roe deer are mainly crepuscular, i.e. their activity locomotor patterns significantly peak at dawn and dusk. Our results confirm the results of Tang et al. (2019), which showed that ranging movements of the Siberian roe deer are defined by heliophany (including sunrise and sunset times). The roe deer seems to shape its activity to limit encounters with its main local predators, e.g. the lynx and the wolf (Soyumert et al. 2019; Tang et al. 2019; Bonnot et al. 2020; Mori et al. 2020b). Bonnot et al. (2020) showed a remarkable plasticity in European

roe deer diel activity in relation to spatiotemporal variation of mortality/predation risks. This 256 257 flexibility was also suggested for populations of Siberian roe deer (Jiang et al. 2008; Tang et al. 2019). This small cervid species shows a limited capacity of the rumen and a fast digestion, 258 requiring frequent feeding bouts (Cederlund 1981; Danilkin 1996). This feature may explain 259 why roe deer activity patterns do not show a single peak, but two peaks of activity/day 260 alternated with resting in cover. An effect of moon phases was also expected, as most prey 261 species tend to avoid bright moonlight nights to limit encounter probabilities with their 262 predators (Lima 1998; Monterroso et al. 2013). Prugh and Golden (2014) reported that the trend 263 in ungulate activity with respect to moon phases is still poorly studied, and it may change with 264 265 species and adapt to local ecological conditions. Nocturnal activity of mostly-diurnal species 266 peaks in brightest nights (Carnevali et al. 2016; Brisbane and van den Burg 2020), as their visual acuity is fundamental for orientation. Instead, ungulate species with mostly crepuscular 267 268 and cathemeral activity may decrease predation risk by moving to cover at night, thus limiting their visibility to predators, without avoidance of bright moonlight (Wronski et al. 2006; Prugh 269 and Golden 2014). Consistently, our results showed that, despite the local presence of several 270 large carnivores, the activity rhythms of Siberian roe deer were independent from moon phases, 271 272 thus not fulfilling our prediction (iii). In fact, Pagon et al. (2013) detected no effect of moon 273 phases on the activity rhythms of the European roe deer. Being active at dawn and dusk might 274 thus represent a strategy to avoid both diurnal and nocturnal local predators (cf. Soyumert et al. 2019; Bonnot et al. 2020; Mori et al. 2020b). 275

Besides individual behaviour (Stache et al. 2013), human activities (e.g hunting pressure and
livestock breeding) may shape the spatiotemporal behaviour of wild ungulates (Douglas 1971;
De Boer et al. 2004; Scillitani et al. 2010; Bonnot et al. 2013; Zbyryt et al. 2018). Jiang et al.
(2008) reported that Siberian roe deer seem to resent human disturbance, much more than
coexisting Manchurian wapitis *Cervus canadensis xanthopygus* Milne-Edwards, 1867,

therefore occurring mostly far from human settlements. Ecological studies on European roe deer have shown that hunting pressure increases vigilance levels, with consequent shifts of feeding sites to cover (Benhaiem et al. 2008) and of activity rhythms (Bonnot et al. 2020).

Hunting Siberian roe deer is allowed in Mongolia by the National Law on Hunting of May 2000, between the 1st of August and the 1st of December. However, our study area was included in a National Park, where hunting is forbidden (cf. Art. 18, Law on Special Protected Areas of Mongolia), although some poaching may have occurred. In our study area, the temporal behaviour of the Siberian roe deer coincided with that of the European roe in sites where human disturbance is the lowest (Bonnot et al. 2020; Mori et al. 2020b). In our study area, the presence of a single human was detected only once (Zaccaroni et al. 2020).

Our data refer only to roe deer activity in the warm season, i.e. they cover the birth and mating periods: two rather special biological phases of roe deer (Melis et al. 2005; Bongi et al. 2008; Lovari et al. 2008). Information on activity rhythms in the cold season, with a much longer night-duration, could yield different results. In particular, the activity of large predators seems to occur mainly at night (Soyumert et al. 2019; Tang et al. 2019; Bonnot et al. 2020; Mori et al. 2020b), thus leading potentially to an even greater impact on the activity rhythms of Siberian roe deer, in the cold season.

298

299 Acknowledgements

300

An anonymous reviewer and the Associate Editor Peter Kaňuch greatly improved our first draftwith their comments.

303

304 Author contributions

306	CA conceived the study; EM and SL conducted literature search and performed the analysis for
307	the review; MC, MZ, SS, AV and CA collected field data. EM, SL and CA led manuscript
308	writing. All authors contributed critically to the draft and gave final approval for submission.
309	Data supporting this study are available in the main text.
310	
311	Research funding
312	
313	No fund was received for this research
314	
315	Conflict of interest statement
316	
317	No potential conflict of interest was reported by authors.
318	
319	Compliance with ethical standard
320	
321	All procedures were conducted without animal handling and with no disturbance to them, in
322	line with all national and international laws on animal and wildlife wellness.
323	
324	References
325	
326	Aanes R, Linnell JDC, Perzanowski K, Karlsen J, Odden J (1998) Roe deer as prey. In:
327	Andersen R, Duncan P, Linnell JDC (Eds) The European roe deer the biology of success.
328	Scandinavian University Press, Oslo, Norway, pp. 138–159.
329	Adhikari P, Park SM, Kim TW, Lee JW, Kim GR, Han SH, Oh H (2016) Seasonal and
330	altitudinal variation in roe deer (Capreolus pygargus tianschanicus) diet on Jeju Island,
331	South Korea. J Asia-Pacific Biodiv 9: 422-428. https://doi.org/10.1016/j.japb.2016.09.001
332	Ancillotto L, Notomista T, Mori E, Bertolino S, Russo D (2018) Assessment of detection
333	methods and vegetation associations for introduced Finlayson's squirrels (Callosciurus

- *finlaysonii*) in Italy. Environm Manage 61: 875-883. https://doi.org/10.1007/s00267-018 1013-x
- Argunov AV (2013) Formation of the range of the Siberian roe deer (*Capreolus pygargus*,
 Cervidae) and its present distribution in Yakutia. Biol Bull 40: 692-697.
 https://doi.org/10.1134/S1062359013080025
- Argunov AV, Safronov VM (2013) Demographic structure of Siberian roe deer (*Capreolus pygargus*, Cervidae) population in Central Yakutia. Russ J Ecol 44: 402407. https://doi.org/10.1134/S1067413613050032
- Augugliaro C, Havmøller RW, Monti IE, Havmøller L, Choikhand J, Lkhagvasuren B (2020)
 Non-volant mammal inventory of western Mongolian-Manchurian Grassland Ecoregion: a
 biogeographic crossroad worth preserving. Check List 16: 287.
 https://doi.org/10.15560/16.2.287
- Augugliaro C, Paniccia C, Janchivlamdan C, Monti IE, Boldbaatar T, Munkhtsog B (2019)
 Mammal inventory in the Mongolian Gobi, with the southeasternmost documented record
- of the Snow Leopard, *Panthera uncia* (Schreber, 1775), in the country. Check List 15: 565–
- 349 578. https://doi.org/10.15560/15.4.565
- Balayed-Lyngdoh S, Habib B, Shrotriya S (2020) Dietary spectrum in Himalayan wolves:
- 351 comparative analysis of prey choice in conspecifics across high-elevation rangelands of
 352 Asia. J Zool 310: 24-33. https://doi.org/10.1111/jzo.12724
- Benhaiem S, Delon M, Lourtet B, Cargnelutti B, Aulagnier S, Hewison AJM, Morellet N,
- 354 Verheyden H (2008) Hunting increases vigilance levels in roe deer and modifies feeding site
- selection. Anim Behav 76: 611-618. https://doi.org/10.1016/j.anbehav.2008.03.012
- Bongi P, Ciuti S, Grignolio S, Del Frate M, Simi S, Gandelli D, Apollonio M (2008) Antipredator behaviour, space use and habitat selection in female roe deer during the fawning

- 358 season in a wolf area. J Zool 276: 242-251. https://doi.org/10.1111/j.1469359 7998.2008.00481.x
- Boitani L, Lovari S, Vigna-Taglianti A (2003) Fauna d'Italia. Mammalia III Carnivora,
 Artiodactyla. Edizioni Calderini, Bologna, Italy.
- Bonnot NC, Morellet N. Verheyden H, Cargnelutti B, Lourtet B, Klein F, Hewison AJM (2013)
- 363 Habitat use under predation risk: hunting, roads and human dwellings influence the spatial
- behaviour of roe deer. Eur J Wildl Res 59: 185-193. https://doi.org/10.1007/s10344-0120665-8
- Bonnot NC, Couriot O, Berger A, Cagnacci F, Ciuti S, De Groeve JE, Gehr B, Heurich M,
- 367 Kjellander P, Kröschel M, Morellet N, Sönnischen L, Hewison AJM (2019) Fear of the dark?
- 368 Contrasting impacts of humans versus lynx on diel activity of roe deer across Europe. J Anim

369 Ecol 89: 132-145. https://doi.org/10.1111/1365-2656.13161

- Botts RT, Eppert AA, Wiegman TJ, Rodriguez A, Blankenship SR, Asselin EM, Garley WM,
- Wagner AP, Ullrich SE, Allen GR, Mooring MS (2020) Circadian activity patterns of
- mammalian predators and prey in Costa Rica. J Mammal 101: 1313-1331.
- 373 https://doi.org/10.1111/1365-2656.13161
- Brisbane JL, van den Burg MP (2020) No need for artificial light: nocturnal activity by a diurnal
 reptile under lunar light. Neotrop Biodivers 6: 193-196.
 https://doi.org/10.1080/23766808.2020.1844993
- Brzeźniak E (1977) Thermal Zones on the Southern Slopes of the Khangai. Bull Acad Pol Sci
 Sci Terre 25: 211-218.
- 379 Carnevali L, Lovari S, Monaco A, Mori E (2016) Nocturnal activity of a "diurnal" species, the
- 380 northern chamois, in a predator-free Alpine area. Behav Processes 126: 101-107.
- 381 https://doi.org/10.1016/j.beproc.2016.03.013

- Cederlund G (1981) Daily and seasonal activity patterns of roe deer in a boreal habitat. Viltrevy
 Swedish Wildl Res 11: 315-353.
- Danilkin A (1996) Behavioural ecology of Siberian and European roe deer. Chapman & Hall,
 London, Great Britain.
- 386 Danilkin AA, Darman IA, Minaev AN, Sempere A (1995) The social organization, behavior
- and ecological parameters of a migrating population of Siberian roe deer (*Capreolus pygargus* Pall.). Izvestiia Akademii Nauk Seriia Biologischeskaia 1: 48-61.
- 389 De Boer HY, van Breukelen L, Hootsman MJM, van Wieren SE (2004) Flight distance in roe
- deer *Capreolus capreolus* and fallow deer *Dama dama* related to hunting and other factors.
- 391 Wildl Biol 10: 35-41. https://doi.org/10.2981/wlb.2004.007
- 392 De Luca DW, Rovero F (2006) First records in Tanzania of the vulnerable Jackson's mongoose
 393 *Bdeogale jacksoni* (Herpestidae). Oryx 40: 468-471.
- 394 Douglas MJW (1971) Behaviour responses of red deer and chamois to cessation of hunting.
 395 New Zeal J Sci 14: 507-518.
- 396 Edwards S, Cooper S, Uiseb K, Hayward M, Wachter B, Melzheimer J (2018) Making the most
- 397 of by-catch data: assessing the feasibility of utilising non-target camera trap data for
- 398 occupancy modelling of a large felid. Afr J Ecol 56: 885-894.
 399 https://doi.org/10.1111/aje.12511
- 400 Ferretti F, Pacini G, Belardi I, ten Cate B, Sensi M, Oliveira R, Rossa M, Burrini L, Lovari S
- 401 (2021) Recolonizing wolves and opportunistic foxes: interference or facilitation? Biol J Linn
- 402 Soc 132: 196-210. https://doi.org/10.1093/biolinnean/blaa139
- 403 Fiske I, Chandler R (2011) Unmarked: a R package for fitting hierarchical models of wildlife
 404 occurrence and abundance. J Stat Soft 43: 1-23.

- Guo K, Liu H, Bao H, Hu J, Wang S, Zhang W, Zhao W, Jiang G (2017) Habitat selection and
 their interspecific interactions for mammal assemblage in the Greater Khingan Mountains,
 northeastern China. Wildl Biol 4: 1-8. https://doi.org/10.2981/wlb.00261
- 408 Helldin JO, Liberg O, Glöersen G (2006) Lynx (*Lynx lynx*) killing red foxes (*Vulpes vulpes*) in
- 409 Boreal Sweden? Frequency and population effects. J Zool 270: 657-663.
 410 https://doi.org/10.1111/j.1469-7998.2006.00172.x
- Hewison AJM, Danilkin A (2001) Evidence for separate specific status of European (*Capreolus*) *capreolus*) and Siberian (*C. pygargus*) roe deer. Mammal Biol 66: 13-21.
- Jiang G, Jianzhang M, Minghai Z (2006) Spatial distribution of ungulate responses to habitat
- factors in Wandashan Forest Region, northeastern China. J Wildl Manage 70: 1470-1476.

Jiang GS, Zhang MH, Ma JZ (2008) Habitat use and separation between red deer and roe deer

- 416 in relation to human disturbance in the Wandashan mountains, northeastern China. Wildl417 Biol 14: 92-100.
- Jiang G, Ma J, Zhang M, Stott P (2009) Assessing microhabitat use by roe deer and moose in
 China. Wildl Res 36: 134-142. https://doi.org/10.1071/WR08092
- 420 Klimek K, Starkel L (1980) Vertical zonality in the Southern Khangai Mountains (Mongolia).
- 421 Geographical Studies 136, Polish Academy of Sciences, NAUK, Wroclaw, Poland.
- 422 Landler L, Ruxton GD, Malkemper EP (2019) The Hermans-Rasson test as a powerful
- 423 alternative to the Rayleigh test for circular statistics in biology. BMC Ecol 19: 1-8.
- 424 https://doi.org/10.1186/s12898-019-0246-8
- 425 Lashley MA, Cove MV, Chitwood MC, Penido G, Gardner B, DePerno CS, Moorman CE
- 426 (2018) Estimating wildlife activity curves: comparison of methods and sample size. Sci Rep
- 427 8: 4173. https://doi.org/10.1038/s41598-018-22638-6
- 428 Lima SL (1998) Stress and decision making under the risk of predation: recent developments
- from behavioural, reproductive, and ecological perspectives. Adv Study Behav 27: 215-290.

- 430 Lorenzini R, Garofalo L, Qin X, Voloshina I, Lovari S (2014) Global phylogeography of the
- 431 genus *Capreolus* (Artiodactyla: Cervidae), a Palaearctic meso-mammal. Zool J Linn Soc
 432 170: 209-221. https://doi.org/10.1111/zoj12091
- 433 Lovari S, Bartolommei P, Meschi F, Pezzo F (2008) Going out to mate: excursion behaviour of
- 434 female roe deer. Ethology 114: 886-896. https://doi.org/10.1111/j.1439-0310.2008.01549.x
- 435 Lovari S, Serrao G, Mori E (2017) Woodland features determining home range size of roe deer.

436 Behav Processes 140: 115-120. https://doi.org/10.1016/j.beproc.2017.04.012

- Lovari S, Masseti M, Lorenzini R (2016) *Capreolus pygargus*. The IUCN Red List of
 Threatened Species 2016: https://dx.doi.org/10.2305/IUCN.UK.20161.RLTS.T42396A22161884.en. Accessed on 09 October 2020.
- 440 Luzi G, Mori E, Puddu G, Zapparoli M (2021) Does the crested porcupine select coppice forest?
- Habitat preference and activity patterns of a large rodent in the Lago di Vico Natural
 Reserve. Mammalia https://doi.org/10.1515/mammalia-2020-0143.
- 443 MacKenzie DI, Nichols JD, Royle JA, Pollock KH, Bailey LL, Hines JE (2006) Occupancy
- estimation and modeling. Inferring patterns and dynamics of species occurrence. Elsevier
 Academic Press Editions, Burlington, Massachusetts, USA.
- 446 Mazzamuto MV, Lo Valvo M, Anile S (2019) The value of by-catch data: How species-specific
- 447 surveys can serve non-target species. Eur J Wildl Res 65: 68.
 448 https://doi.org/10.1007/s10344-019-1310-6
- Melis C, Cagnacci F, Lovari S (2005) Do male roe deer clump together during the rut? Acta
 Theriol 50: 253-262.
- 451 Meredith M, Ridout M (2014) Overview of the Overlap Package. Available from:
 452 http://cran.cs.wwu.edu/web/packages/overlap/vignettes/overlap.pdf. Accessed on
 453 12.12.2020.

- Merson SD, Dollar LJ, Ta CKW, Macdonald DW (2019) Activity patterns of sympatric living
 exotic and endemic carnivores (the fosa) in Western Madagascar's deciduous forests. J Zool
 307: 186-194. https://doi.org/10.1111/jzo.12630
- Monterroso P, Alves PC, Ferreras P (2013) Catch me if you can: diel activity patterns of
 mammalian prey and predators. Ethology 119: 1044-1056.
 https://doi.org/10.1111/eth.12156
- 460 Mori E, Di Bari P, Coraglia M (2018) Interference between roe deer and Northern chamois in
- the Italian Alps: are Facebook groups effective data sources? Ethol Ecol Evol 30: 277-284.

462 https://doi.org/10.1080/03949370.2017.1354922

- Mori E, Sangiovanni G, Corlatti L (2020a) Gimme shelter: the effects of rock and moonlight
 on occupancy and activity pattern of an endangered rodent, the garden dormouse *Eliomys quercinus*. Behav Processes 170: 103999. https://doi.org/10.1016/j.beproc.2019.103999
- 466 Mori E, Bagnato S, Serroni P, Sangiuliano A, Rotondaro F, Marchianò V, Cascini V, Poerio L,
- 467 Ferretti F (2020b) Spatiotemporal mechanisms of coexistence in a European mammal
 468 community in a protected area of southern Italy. J Zool 310: 232-245.
 469 https://doi.org/10.1111/jzo.12743
- 470 Narangerel Y (2010) Issues in the Policy on Establishing New Communities Among Pastoral
 471 Nomadic Herders in Mongolia. MA Thesis. Keio University, Japan.
- 472 Newsome TM, Boitani L, Chapron G, Ciucci P, Dickman CR, Dellinger JA, Lopez-Bao JV,
- 473 Peterson RO, Shores CR, Wirsing AJ, Ripple WJ (2016) Food habits of the world's grey
- 474 wolves. Mammal Rev 46: 255-269. https://doi.org/10.1111/mam.12067
- 475 Odden J, Linnell JDC, Andersen R (2006) Diet of Eurasian lynx, *Lynx lynx*, in the boreal forest
- 476 of southeastern Norway: The relative importance of livestock and hares at low roe deer
- 477 density. Eur J Wildl Res 52: 237-244. https://doi.org/10.1007/s10344-006-0052-4

- 478 Ossi F, Ranc N, Moorcroft P, Bonanni P, Cagnacci F (2020) Ecological and behavioral drivers
 479 of supplemental feeding use by roe deer *Capreolus capreolus* in a peri-urban context.
 480 Animals 10: 2088. https://doi.org/10.3390/ani10112088
- 481 Pagon N, Grignolio S, Pipia A, Bongi A, Bertolucci C, Apollonio M (2013) Seasonal variation
- 482 of activity patterns in roe deer in a temperate forested area. Chronobiol International 30:
- 483 772-785. https://doi.org/10.25225/fozo.v66.i4.a9.2017
- Pamplin FA (2013) Optimising the value of by-catch from *Lynx lynx* camera trap surveys in the
 Swiss Jura region. MSc thesis in Applied Ecology and Conservation University of East
 Anglia, UK.
- 487 QGIS Development Team (2019) QGIS Geographic Information System. Open Source
 488 Geospatial Foundation Project. http://qgis.osgeo.org. Accessed on 20.04.2021.
- Reed JM (1996) Using statistical probability to increase confidence of inferring species
 extinction. Conserv Biol 10: 1283-1285. https://doi.org/10.1046/j.15231739.1996.10041283.x
- 492Robin K (1975) Rauemliche Verscienbungen von markieten Rehen (Capreolus capreolus L.)
- in einem voralpinem Gebiet der Ostchweiz. Z Jagdwiss 21: 145-163.
- 494 Ross JGB, Peters W, Ossi F, Moorcroft PR, Cordano E, Eccel E, Bianchini F, Ramanzin M,
- 495 Cagnacci F (2021) Climate change and anthropogenic food manipulation interact in shifting
- the distribution of a large herbivore at its altitudinal range limit. Sci Rep 11: 1-12.
 https://doi.org/10.1038/s41598-021-86720-2
- 498 Rossa M, Lovari S, Ferretti F (2021) Spatiotemporal patterns of wolf, mesocarnivores and prey
 499 in a Mediterranean area. Behav Ecol Sociobiol 75:32. https://doi.org/10.1007/s00265-020-
- 500 02956-4
- Rovero F, Zimmermann F (2016) Camera trapping for wildlife research. Pelagic Publishing
 Ltd, Exeter, UK.

- San Josè C, Lovari S, Ferrari N (1996) Temporal evolution of vigilance in roe deer. Behav
 Processes 38: 155-159.
- Scillitani L, Monaco A, Toso A (2010) Do intensive drive hunts affect wild boar (*Sus scrofa*)
 spatial behaviour in Italy? Some evidences and management implications. Eur J Wildl Res
- 507 56: 307-318. https://doi.org/10.1007/s10344-009-0314-z
- 508 Soyumert A, Ertürk A, Tavşanoğlu Ç (2019) The importance of lagomorphs for the Eurasian
- 509 lynx in Western Asia: results from a large scale camera-trapping survey in Turkey. Mammal

510 Biol 95: 18-25. https://doi.org/10.1016/j.mambio.2019.01.003

- 511 Srivathsa A, Karanth KK, Jathanna D, Kumar NS, Karanth KU (2014) On a dhole trail:
- 512 examining ecological and anthropogenic correlates of dhole habitat occupancy in the
- 513 Western Ghats of India. PloS One 9: e98803. https://doi.org/10.1371/journal.pone.0098803
- Stache A, Heller E, Hothorn T, Heurich M (2013) Activity patterns of roe deer (*Capreolus*) *capreolus*) are strongly influenced by individual behaviour. Folia Zool 62: 67-75.
 https://doi.org/10.25225/fozo.v62.i1.a10.2013
- Tan S, Jiang X, Wang F, Zhang G, Zhang J, Men H, Han Y, Bao W (2019). Preliminary analysis
 on the diet composition of the Eurasian lynx at Saihanwula National Nature Reserve. Chin
- 519 J Zool 54: 151-158.
- Tang X, Tang S, Li X, Menghe D, Bao W, Xiang C, Gao F, Bao W (2019) A study of population
- size and activity patterns and their relationship to the prey species of the Eurasian lynx using
- a camera trapping approach. Animals 9: 864. https://doi.org/10.3390/ani9110864
- 523 Tobler MW, Carrillo-Percastegui SE, Leite Pitman R, Mares R, Powell G (2008) Further notes
- 524 on the analysis of mammal inventory data collected with camera traps. Animal Cons 11:
- 525 187–189. https://doi:10.1111/j.1469-1795.2008.00181.x

- 526 Van Dijk J (2008) Wolverine foraging strategies in a multiple-use landscape. Ph.D.
 527 Dissertation, Norwegian University of Science and Technology Faculty of Natural Sciences
 528 and Technology Department of Biology, Trondheim, Norway.
- 529 Viviano A, Amori G, Luiselli L, Oebel H, Bahleman F, Mori E (2020) Blessing the rains down
- 530 in Africa: spatiotemporal behaviour of the crested porcupine *Hystrix cristata* (Mammalia:
- Rodentia) in the rainy and dry seasons, in the African savannah. Tropical Zool 33: 113-124.
- 532 https://doi.org/10.4081/tz.2020.80
- 533 Viviano A, Mori E, Fattorini E, Mazza G, Lazzeri L, Panichi A, Strianese L, Mohamed WF
- 534 (2021) Spatiotemporal overlap between the European brown hare and its potential predators
- and competitors. Animals 11: 562. https://doi.org/10.3390/ani11020562
- Wronski T, Apio A, Plath M (2006) Activity patterns of bushbuck (*Tragelaphus scriptus*) in
 Queen Elizabeth National Park. Behav Processes 73: 333-341.
 https://doi.org/10.1016/j.beproc.2006.08.003
- Yan W, Zeng Z, Gong H, Duan Y, Zhao L, Peng A (2020) Locomotor activity patterns of takin
 (*Budorcas taxicolor*) in a temperate mountain region. Plos One 15: e0235464.
 https://doi.org/10.1371/journal.pone.0235464
- 542 Zaccaroni M, Monti IE, Biliotti N, Munkhtsog B, Comand N, De Donà G, Augugliaro C (2020)
- A preliminary mammal inventory of the Southernmost Mongolian forest: discovery a
 mesocarnivores heaven. Species 21: 208-221.
- 545 Zanni M, Brivio F, Grignolio S, Apollonio M (2021) Estimation of spatial and temporal overlap
 546 in three ungulate species in a Mediterranean environment. Mammal Res 66: 149-162.
- 547 https://doi.org/10.1007/s13364-020-00548-1
- 548 Zbyryt A, Bubnicki JW, Kuijper DP, Dehnhard M, Churski M, Schmidt K (2018) Do wild
- ungulates experience higher stress with humans than with large carnivores? Behav Ecol 29:
- 550 19-30. https://doi.org/10.1093/beheco/arx142

551 Tables

552

553

Table 1. Summary of the five parameter estimates for the occupancy. Models are ranked according to the AIC_C, w_i (Akaike weight), ψ (occupancy probability), p (detectability), SE (standard error), N (sample size estimate). The best selected model is shown in bold.

Model	AIC _C	ΔAIC_{C}	\mathbf{W}_i	ψ (SE)	p (SE)	Ν
ψ ("slope + presence of large carnivores")p(".")	307.30	0.00	0.98	0.21 (0.07)	0.74 (0.12)	4
ψ ("presence of large carnivores")p("elevation")	317.54	10.24	0.01	0.11 (0.03)	0.32 (0.21)	4
ψ ("slope")p(".")	338.17	12.12	0.01	0.10 (0.03)	0.30 (0.17)	5
ψ(".")p(".")	347.22	15.04	0.01	0.12 (0.02)	0.31 (0.24)	5

558

560	Figure	Legends
	0	

561

Figure 1. Camera-trapped Siberian roe deer and environmental features of the study area.

563

Figure 2. Location of the study area (inset map), location of camera traps and occurrences ofSiberian roe deer.

566

Figure 3. Circadian rhythms of the Siberian roe deer, expressed as Kernel density estimate of activity throughout the 24-h cycle (N = 51 independent records). Blue lines represent bootstrapped estimates of activity patterns; dashed black lines represent 95 % CIs.