

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Normalized Difference Vegetation Index versus Dark Green Colour Index to estimate nitrogen status on bermudagrass hybrid and tall fescue

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Caturegli, L., Gaetani, M., Volterrani, M., Magni, S., Minelli, A., Baldi, A., et al. (2020). Normalized Difference Vegetation Index versus Dark Green Colour Index to estimate nitrogen status on bermudagrass hybrid and tall fescue. *INTERNATIONAL JOURNAL OF REMOTE SENSING*, 41(2), 455-470 [10.1080/01431161.2019.1641762].

Availability:

This version is available at: <https://hdl.handle.net/11585/716429> since: 2021-11-25

Published:

DOI: <http://doi.org/10.1080/01431161.2019.1641762>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **Title** Normalized Difference Vegetation Index versus Dark Green Color Index to estimate nitrogen
2 status on bermudagrass hybrid and tall fescue.

3 **Short title:** NDVI vs DGCI to estimate N on two turfgrass species
4

5 **Authors:** Lisa Caturegli^{*1}, Monica Gaetani¹, Marco Volterrani¹, Simone Magni¹, Alberto Minelli²,
6 Ada Baldi³, Giada Brandani³, Marco Mancini³, Anna Lenzi³, Simone Orlandini³, Filippo Lulli⁴,
7 Claudia de Bertoldi⁴, Marco Dubbini⁵, Nicola Grossi¹
8

9 ¹ Department of Agriculture, Food and Environment, University of Pisa, Pisa, Italy,

10 ² Department of Agricultural and Food Sciences, University of Bologna, Bologna, Italy

11 ³ Department of Agriculture, Food, Environment and Forestry, University of Florence, Florence,
12 Italy

13 ⁴ Turf Europe Srl, Livorno, Italy

14 ⁵ Department of History, Cultures and Civilizations, University of Bologna, Bologna, Italy;

15 *Corresponding author Email: lisa.caturegli@gmail.com
16

17 **Abbreviations:**

18 DGCI Dark Green Color Index

19 GPS Global Positioning System

20 HSB Hue Saturation Brightness

21 NDVI Normalized Difference Vegetation Index

22 PA Precision Agriculture

23 PTM Precision Turfgrass Management

24 RGB Red Green Blue

25 UAS Unmanned Aerial Systems

26 UAV Unmanned Aerial Vehicle

27 **Keywords:**

28 *Cynodon dactylon x transvaalensis*, *Schedonorus phoenix*, Turfgrass, Color, Quality, Unmanned

29 Aerial Vehicle.

30

31 **Abstract**

32 In recent years digital sensors have been successfully integrated on board Unmanned Aerial
33 Vehicles (UAV) to assess crop vigor, vegetation coverage, and to quantify the “greenness” of
34 foliage as indirect measurements of crop nitrogen status. The classical approach of precision
35 agriculture has involved the use of multispectral sensors onboard UAV and the development of
36 numerous vegetation indices associated with vegetation parameters, such as the mostly used
37 Normalized Difference Vegetation Index (NDVI). However, the main negative issue when dealing
38 with multi and hyper-spectral reflectance measuring tools is their high cost and complexity from the
39 operational point of view. As a low-cost alternative, vegetation indices derived from Red Green
40 Blue (RGB) cameras have been employed for remote sensing assessment, providing data on
41 different stress conditions and species. Digital images record information as amounts of RGB light
42 emitted for each pixel of the image; however, the intensity of red and blue will often alter how
43 green an image appears. To simplify the interpretation of digital color data, recent studies have
44 suggested converting RGB values to the more intuitive Hue, Saturation, and Brightness (HSB) color
45 spectrum, and then into a single measure of dark green color, the Dark Green Color Index (DGCI).
46 In this study NDVI acquired by a ground-based handheld crop sensor and by a multispectral camera
47 mounted on board a UAV have been compared with DGCI calculated from images taken with a
48 commercial digital camera on board a UAV, trying to quantify the color of turfgrass that had
49 received different nitrogen (N) rates.

50 The objectives of the trial were to study an affordable easy-to-use tool evaluating the relationship
51 among NDVI, DGCI and leaf nitrogen content on turfgrass.

52

53 **Introduction**

54 Nitrogen fertilization on turfgrasses is one of the factors that most influence physiological and
55 aesthetic aspects (Volterrani et al. 2005; Perry and Davenport 2007; Samborski, Tremblay, and
56 Fallon 2009; Caturegli et al., “Monitoring turfgrass”, 2014; Caturegli et al., “Turfgrass spectral
57 reflectance”, 2014; Grossi et al. 2016). Thus, nitrogen (N) represents an important nutrient that
58 contributes to maintain green color, density, recovery from drought diseases, and a general good
59 turfgrass quality (Walters and Bingham 2007; Dordas 2008; Magni et al. 2014; Caturegli et al.
60 2016).

61 However, the excessive fertilization of N wastes fertilizers and leads to pollution of ground and
62 surface water, not improving the quality of the turf (Bell and Xiong 2008; Bremer et al. 2011;
63 Rhezali and Lahlali 2017). To avoid over-fertilization, site-specific nutrition management brought
64 significant environmental and economic benefits (Huang et al. 2008). Indeed, a precise analysis of
65 the plant nitrogen status is important to determine the amount of nitrogen fertilizer the plant really
66 needs (Corwin and Lesch 2005; Li et al. 2015).

67 Previous studies have focused on implementation of indirect sensing tools (chlorophyll meters,
68 reflectance measurements, color analysis) to try to obtain an almost optimal quality by reducing the
69 N inputs and the loss N to a minimum (Rorie, Purcell, Mozaffari et al. 2011; Caturegli, Casucci et
70 al. 2015; Caturegli, Grossi et al. 2015; Caturegli et al. 2016).

71 These concepts are the basis of Precision Agriculture (PA), which aims to obtain detailed site-
72 specific information by mapping the variation in important soil and plant properties in order to
73 allow better site-specific management. Inputs such as water, fertilizers and pesticides are applied
74 only where, when and in the amount needed by plant (Caturegli et al., “Turfgrass spectral
75 reflectance”, 2014). Related to PA is Precision Turfgrass Management (PTM) that is useful to
76 monitor pests, fertilization, salinity stress and irrigation deficiency on turfgrass (Carrow et al. 2010;
77 Krum, Carrow, and Karnok 2010). The approach of PA implied the combined use of multispectral

78 sensors and vegetation indices associated with vegetation parameters (Trenholm, Carrow, and
79 Duncan 1999; Jiang and Carrow 2007; Vergara-Díaz et al. 2016). Thus, vegetation indices were
80 calculated by combining various reflectance bands of the spectrum and correlated with relevant
81 turfgrass canopy parameters. Among the indices, the Normalized Difference Vegetation Index
82 (NDVI) is the most widely used as reflectance-based plant stress indicator (Hansen and Schjoerring
83 2003; Johnsen et al. 2009; Aguilar et al. 2012; Barton 2012; Fensholt and Proud 2012; Rhezali and
84 Lahlali 2017). It is based on the relationship between the absorption of visible light and resilient
85 reflectance of near-infrared light to the chlorophyll in vegetation (Bell et al. 2004; Caturegli,
86 Casucci et al. 2015). The NDVI value ranges from -1 to 1, with higher values indicating greater
87 plant health, and correlates positively with turfgrass quality (Trenholm, Carrow, and Duncan 1999;
88 Fitz-Rodriguez and Choi 2002; Leinauer et al. 2014). This index is also influenced by differences in
89 species, environmental stresses, fertilization and pest injuries (Xiong et al. 2007; Bremer et al.
90 2011; Caturegli, Grossi et al. 2015). It can be obtained with hand-held ground-based instruments
91 (Graeff and Claupein 2003; Ma, Morrison, and Dwyer 1996) and aerial vehicle-mounted sensors
92 (Bausch and Duke 1996; Blackmer et al. 1996; Scharf and Lory 2009; Rorie, Purcell, Karcher et al.
93 2011). In recent years, digital sensors have been successfully integrated on board Unmanned Aerial
94 Vehicles (UAV) to assess crop vigor, vegetation coverage, and to quantify the “greenness” of
95 foliage as indirect measurements of crop N status (White et al. 2012; Andrade-Sanchez et al. 2014).
96 Furthermore, small commercial Unmanned Aerial Systems (UAS) (< 50 kg) (Laliberte and Rango
97 2011) have been available for PA for environmental and agricultural applications (Gupta et al.
98 2013; Zhang and Kovacs 2012; Caturegli et al. 2016). However, the main negative issue when it
99 comes to multi and hyper-spectral reflectance measuring tools is their high cost and complexity.
100 Vegetation indices derived from Red-Green-Blue (RGB) cameras have been employed for remote
101 sensing assessment, as a low-cost alternative (Vergara-Díaz et al. 2016). This method may provide
102 data on different stress conditions in different crops (Casadesús et al. 2007; Casadesus and Villegas
103 2014; Zhou et al. 2015) and turfgrass (Karcher and Richardson 2003; Karcher and Richardson

104 2013). Digital images are composed by pixels that record information as amounts of RGB light
105 emitted. However, the greenness of an image can be often altered by the intensity of red and blue.
106 To simplify the interpretation of data, Karcher and Richardson (2003) suggested converting RGB
107 values to the more intuitive Hue, Saturation, and Brightness (HSB) based on human perception of
108 color. Working with quality of turfgrass in response to N fertilizer, Karcher and Richardson (2003)
109 processed HSB values into a single measure of dark green color, the Dark Green Color Index
110 (DGCI).

111 This method proposed by (Karcher and Richardson 2003) may represent a proper alternative to the
112 spectroradiometric approaches that involves the use of NDVI from aerial platforms and from
113 ground-based measurements (Vergara-Díaz et al. 2016). To facilitate the DGCI acquisition,
114 recently, also a smartphone application called FieldScout GreenIndex+ Turf (Spectrum
115 Technologies, Inc., Aurora, IL, USA) (Spectrum Technologies, Inc. 2018) has been developed and
116 tested (O'Brien 2017; Xiang et al. 2017; Xiang et al. 2018) The application (APP) captures images
117 with a smartphone or tablet, calculates the DGCI, and shows a turfgrass quality visual rating
118 (Karcher and Richardson 2003).

119 The aim of this research was to study an affordable easy-to-use tool evaluating the relationship
120 among NDVI, DGCI and leaf nitrogen content on turfgrass. Trying to quantify the color of turfgrass
121 that had received different N rates, NDVI acquired by a ground-based handheld crop sensor and by
122 a multispectral camera mounted on board a UAV have been compared with DGCI calculated from
123 images taken with a commercial digital camera on board a UAV.

124

125 **Materials and Methods**

126 The trial was carried out in July 2017 in S. Piero a Grado, Pisa, at the Centre for Research on
127 Turfgrass for the Environment and Sports (CeRTES) of the Department of Agriculture, Food and
128 Environment of the University of Pisa (43°40'N, 10°19'E, 6 metres above sea level (m. a. s. l.).

129 The turfgrasses selected for the study were a mature turfgrass stands of the warm-season
130 bermudagrass hybrid (*Cynodon dactylon* [L.] Pers. (Linnaeus Persoon) variety *dactylon* x *Cynodon*
131 *transvaalensis* Burt-Davy) cultivar (cv) 'Patriot' and the cool-season tall fescue (*Schedonorus*
132 *phoenix* [Scop.] (Scopoli) Holub) cv 'Grande'.

133 The swards were all established on a calcareous fluvisoil (Coarse-silty, mixed, thermic, Typic
134 Xerofluvents) with pH 7.8 and 18 g kg⁻¹ of organic matter.

135 No fertilizer had been applied to the turfgrass before the trial started. In order to create a linear
136 nitrogen gradient, on June 2017 fertilization was carried out applying ammonium sulphate (21-0-0)
137 with a rotary spreader (ICL Specialty Fertilizers AccuPro 2000, Ipswich, UK).

138 The experimental designs were:

139 a) For tall fescue 8 nitrogen rates were applied, from 0 to 210 kg ha⁻¹ of N with increases of 30 kg
140 ha⁻¹ (0 kg ha⁻¹, 30 kg ha⁻¹, 60 kg ha⁻¹, 90 kg ha⁻¹, 120 kg ha⁻¹, 150 kg ha⁻¹, 180 kg ha⁻¹, 210 kg ha⁻¹
141 of N). The plot size was 3 m × 3 m, with 3 replications.

142 b) For bermudagrass hybrid, which tolerates higher doses of fertilizer, 11 nitrogen rates were
143 applied, from 0 to 300 kg ha⁻¹ of N with increases of 30 kg ha⁻¹ (0 kg ha⁻¹, 30 kg ha⁻¹, 60 kg ha⁻¹,
144 90 kg ha⁻¹, 120 kg ha⁻¹, 150 kg ha⁻¹, 180 kg ha⁻¹, 210 kg ha⁻¹, 240 kg ha⁻¹, 270 kg ha⁻¹, 300 kg ha⁻¹
145 of N). The plot size was 3 m × 3 m, with 3 replications.

146 Extreme N rates were applied in order to reach the nitrogen saturation level for both species,
147 regardless of the agronomic drawbacks to the turfgrasses.

148 After the fertilization, an irrigation of 5 mm was applied. During the trial period a turf height of 2.0
149 cm was maintained by mowing with a walk-behind reel mower (John Deere 20SR7, Moline IL,

150 USA) with clippings removal. In the entire experimental area, in order to evaluate nitrogen
151 fertilization as the only variability source, identical and maintenance practices were applied.
152 Irrigation was applied as needed to avoid wilt, in order to maintain the soil moisture constant and
153 equal in all areas. During the trial no weed or pest control was necessary.

154 On each of the two experimental areas proximity and remote sensed readings were acquired starting
155 from the unfertilized control to the highest nitrogen rate in each plot.

156 The ground-based instrument used to acquire NDVI values was a Handheld Crop Sensor (HCS)
157 (GreenSeeker, Model HSC-100, Trimble Navigation Unlimited, Sunnyvale, CA) while the remote
158 sensed readings were collected with a UAV which was a VTOL (Vertical Take Off and Landing)
159 DJI s900 hexacopter (DJI, Shenzhen, China) equipped with a digital commercial camera Sony Nex
160 5 (Sony, Surrey, United Kingdom) and a lightweight multispectral sensor MAIA S2 (SAL
161 Engineering, Modena Italy; EOPTIS, Trento, Italy). Spectral measurements (proximity and aerial)
162 were taken on 6 July 2017 between 11:30 AM (ante meridiem) and 1:30 PM (post meridiem) (local
163 time), in complete absence of clouds. The weather parameters of July 2017 were as follows:
164 average air temperature 25 °C, average relative humidity 60%; July average of the noon
165 Photosynthetic Photon Flux Density 1,482 $\mu\text{mol m}^{-2} \text{s}^{-1}$; average wind speed 6 km h⁻¹. Each ground-
166 based measurement was geo-referenced to sub-meter accuracy with a Global Positioning System
167 (GPS) receiver Leica 1200 in Real Time Kinematic, in order to find the exact position on the UAV
168 images and to compare data acquired with the two systems (Caturegli, Casucci et al. 2015;
169 Caturegli, Grossi et al. 2015; Caturegli et al. 2016).

170 **Ground-based measurements**

171 Proximity sensed measurements of spectral reflectance were acquired with a HCS at a height
172 of 110 cm from the ground, thus monitoring a surface of about 2,000 cm² ($\varnothing = 50$ cm). The HCS
173 has an active light source that makes readings unaffected by sunlight (Bell, Kruse, and Krum 2013).
174 Reflectance was measured in the red region at 660 nm, and in the near infrared region of the

175 spectrum at 780 nm. The output is directly provided as NDVI, which is calculated using the
176 equation:

$$177 \quad \text{NDVI} = ((\text{NIR}) - R) / ((\text{NIR}) + R) \quad (1)$$

178 where R is the reflectance in the red band and NIR is the reflectance in the near-infrared band.

179 In the same day and in the same area also the following parameters were studied:

- 180 - Color intensity (1 = very light green; 6 = acceptable green; 9 = very dark green): visual
181 assessments (Morris and Shearman 2008);
- 182 - Turfgrass Quality: (1 = poor; 6 = acceptable; 9 = excellent): visual assessments (Morris and
183 Shearman 2008);
- 184 - Total N content of leaves: samples of clippings were collected on each sampling area with a
185 walk-behind reel mower from a surface of 0.5 m² (1.0 m × 0.5 m). Fresh clippings were put
186 in a ventilated stove at 70 °C, dried to constant weight, and the total N was determined by
187 the micro-Kjeldahl method (Bremner 1965);
- 188 - Plant water content (PWC): calculated as follows:

$$189 \quad \text{PWC (\%)} = \frac{190 \quad (\text{FW}) - (\text{DW})}{(\text{FW})} \times 100 \quad (2)$$

191 where FW is the leaf fresh weight and DW the leaf dry weight. Leaves were cut and quickly
192 put into a plastic bag with hermetic closure. The bags were refrigerated and kept in the dark
193 until arrival to the laboratory, where they have been weighed.

194 **UAV flight and analysis of UAV derived imagery**

195 The UAV system used for surveying was a DJI s900 hexacopter (Figure 1 (a)) with Global
196 Navigation Satellite System (GNSS), with L1 code solution and a 3 axis accelerometer based
197 stabilization system. The hexacopter was equipped with a digital commercial camera Sony Nex 5
198 (Sony, Surrey, United Kingdom) and a lightweight multispectral camera MAIA S2 (SAL
199 Engineering, Modena Italy; EOPTIS, Trento, Italy) (Figure 1 (b)). The images were acquired at 90

200 m of altitude to guarantee a GSD (Ground Sample Distance) of less than 5 cm and a FOV (Field Of
201 View) of about 58 m × 43 m. The direction and altitude of the aircraft were controlled by the
202 rotation speed or by the direction of the propellers (Li et al. 2015). Real-time images, and other
203 information such as altitude and battery voltage, were transmitted to a ground monitor through a
204 radio link.

205 **Please insert Figure 1 near here**

206

207 **UAS derived imagery NDVI**

208 The UAS derived imagery NDVI was obtained using the UAV cited above equipped with a
209 multispectral camera MAIA S2 (SAL Engineering, Modena Italy; EOPTIS, Trento, Italy), which
210 features an array of nine sensors with 1.2 Megapixel resolution: specifically, one RGB color and
211 eight monochrome sensors are available for analysis of the visible and near infra-red (VIS-NIR)
212 spectrum from 390 nm to 950 nm, operating with a frame rate of 5 Hz per sensor. Each of the eight
213 sensors is provided with a band-pass filter (Table 1). Global shutter technology is so such that all of
214 the pixels in each sensor start to collect charge simultaneously, allowing images to be scanned in
215 “one shot” for synchronized multiband measurements. The extremely fast exposure times of the
216 nine global shutter complementary metal-oxide semiconductor (CMOS) sensors (up to 10^{-4} s) and
217 the low travel speed ($< 0.5 \text{ m s}^{-1}$) guarantee the absence of the blur effect. The images obtained
218 were geometrically corrected with calibrated optics, and radiometrically corrected with the
219 acquisition of the reflectance values of the incident light through a calibrated white panel. After the
220 corrections, the 9 images for each shot were registered using the proprietary MAIA software based
221 on photogrammetric method (Dubini et al. 2017). Every pixel of the image contained coordinates
222 and an NDVI value that was extracted using Quantum GIS (Geographic Information System) 2.18
223 software.

224 **Please insert Table 1 near here**

225

226 **Dark Green Color Index (DGCI)**

227 A common digital camera, Sony Nex 5 (Sony, Surrey, United Kingdom) was used to capture
228 RGB images of the selected area. The Sony Nex-5 is a mirrorless interchangeable-lens camera, with
229 the Advanced Photo System-Classic (APS-C) Exmor CMOS sensor and a maximum image
230 resolution of $4,912 \times 3,264$ and a pixel size of $5 \mu\text{m}$ in both x and y directions (Remondino and
231 Fraser 2006; Fryskowska et al. 2016). To reduce the effect of vibration during the flight and capture

232 clear images, the camera was mounted on a pan-tilt set which keeps the lens horizontal. In the same
233 day as the ground NDVI readings and the NDVI by the multispectral camera, the digital camera
234 recorded UAS derived imagery RGB images above the interested area, always in a zenithal plane.
235 Images were taken with auto-focus, auto-white balance and an automatic exposure, and they were
236 saved in Joint Photographic Experts Group (JPEG) format. Subsequently, images were analyzed
237 with the open source Quantum GIS 2.18 software to extract the RGB values of the pixels where the
238 NDVI values by the ground and by the multispectral camera were calculated. To simplify the
239 interpretation of data, RGB values were converted into HSB values, using the method suggested by
240 (Karcher and Richardson 2003), to finally calculate the DGCI. DGCI value is on a scale from 0
241 (very yellow) to 1 (dark green) (Rhezali and Lahlali 2017). DGCI was calculated as:

$$242 \quad \text{DGCI} = [((\text{Hue}) - 60)/60 + (1 - (\text{Saturation})) + (1 - (\text{Brightness}))]/3 \quad (3)$$

243

244 **Statistical analysis**

245 The correlations between the two different NDVI reading methods (ground-based sensing
246 with a HCS and remote sensing with UAV) and DGCI were studied using CoStat software (CoHort,
247 Monterey, CA, USA) and Pearson's correlation coefficients (r) were calculated in order to verify
248 whether: (a) NDVI-ground data and NDVI-UAV were suitably correlated with DGCI obtained from
249 RGB images captured by the digital camera on board a UAV; (b) UAV imagery with a low cost
250 digital camera could be a diagnostic tool to identify variation in N status of turfgrass, comparable to
251 a more expensive multispectral camera. Linear relationships were studied for the correlations
252 showing statistically significant coefficients.

253 **Results and discussion**

254 **Relationship between DGCI, NDVI and observed parameters**

255 In bermudagrass hybrid, considering r among NDVI values obtained with the two different
256 instruments (proximity sensed with the HCS GreenSeeker and remotely sensed with the
257 multispectral camera MAIA mounted on board a UAV), and the measured parameters, the r values
258 were highly significant. The r values ranged between 0.92 for PWC-NDVI of both the instruments
259 and 0.97 for turfgrass quality-NDVI GreenSeeker. Comparing DGCI and all the measured
260 parameters, the index was significantly correlated with color intensity, turfgrass quality and plant
261 water content with r values ranging between 0.83 for color intensity and 0.84 for turfgrass quality
262 and PWC (Table 2).

263 In tall fescue the correlations between NDVI obtained with GreenSeeker and with UAV and color
264 intensity ($r = 0.96$ and $r = 0.95$) has showed higher r values than the same in bermudagrass hybrid
265 ($r = 0.94$). Also PWC-NDVI (GreenSeeker and UAV) showed a degree of association significantly
266 higher in tall fescue ($r = 0.98$) than bermudagrass hybrid ($r = 0.92$).

267 Furthermore, observing the correlations, the DGCI was highly correlated with all the measured
268 parameters with r values ranging between 0.92 for DGCI-Quality and 0.98 for DGCI-PWC. These
269 relationships were all significantly higher in tall fescue than bermudagrass hybrid (Table 2) also in
270 the case of DGCI-turfgrass color ($r = 0.95$). Previous reports by Zhang and Kovacs (2012), and
271 Leinauer et al. (2014) also indicated this trend of values between DGCI and turfgrass quality and
272 turfgrass color. As in our study also in the report by Leinauer et al. (2014), the association between
273 DGCI and turfgrass quality in tall fescue showed higher r values than the same association in
274 bermudagrass hybrid. As for the turfgrass color, Zhang and Kovacs (2012) also studied the
275 relationship between visual color rating and DGCI, with higher Pearson correlation coefficient in
276 tall fescue than bermudagrass hybrid. Previous reports by Karcher and Richardson (2003) also

277 confirm that visual ratings can be used to separate treatment effects on turf color. Frequently raters
278 ranked the turf plots similarly although differences in color existed. Therefore, visual color rating
279 remains a valid evaluation tool if data are not compared across raters. However, the accuracy of
280 DGCI, as demonstrated in previous studies, enables researchers to record reflected turfgrass color
281 on a standardized scale rather than using arbitrary rating values.

282 **Please insert Table 2 near here**

283

284 **Relationship between DGCI and NDVI**

285 Both in bermudagrass hybrid (Figure 2 (a)) and tall fescue (Figure 2 (b)) DCGI significantly
286 related to the average NDVI values measured with a HCS (GreenSeeker) and with the multispectral
287 camera MAIA mounted on board a UAV, although data have been collected by instruments that
288 measure at different heights and spatial resolutions. In fact DGCI has been collected only with RGB
289 camera mounted on board a UAV, while NDVI has been measured by a multispectral camera on
290 board a UAV and also by a ground based HCS.

291 As shown in the Figure 2, DGCI values were linearly associated with NDVI, as also demonstrated
292 by Leinauer et al. 2014. In Figure 2 (a) bermudagrass hybrid has performed a higher degree of
293 association in NDVI GreenSeeker-DGCI ($r = 0.91$) than with UAV ($r = 0.85$), while in the case of
294 tall fescue the degree of association was statistically the same (Figure 2 (b)). Comparing the two
295 species, it was interesting to note that in tall fescue the correlation coefficients (Table 2) between
296 both NDVI (GreenSeeker and UAV) and DGCI were higher than in Bermudagrass (Table 2; Figures
297 2 (a) - (b)).

298

299 **Please insert Figure 2 near here**

300

301

302 **Relationship between DGCI and clipping nitrogen content**

303 Figure 3 showed the linear relationship between DGCI and clipping nitrogen content
304 percentage in bermudagrass hybrid (a) and tall fescue (b) and it was of interest to note that the
305 coefficients were high for both the species. However, DGCI in tall fescue showed a higher degree of
306 association with clipping N content ($r = 0.95$), than in bermudagrass hybrid ($r = 0.86$) (Figures 3 (a)
307 - (b)). DGCI values were linearly associated with clipping nitrogen content, as also demonstrated in
308 other crops (Rorie, Purcell, Mozaffari et al. 2011; Vergara-Díaz et al. 2016). Thus, DGCI values
309 could predict the average nitrogen concentrations of tall fescue and bermudagrass hybrid clippings
310 in different plots and with different application rates.

311 The close association between DGCI and leaf nitrogen therefore provided an additional tool for the
312 assessment of leaf nitrogen content. Our research was consistent with previous work by Karcher and
313 Richardson (2003) who found that DGCI values were able to differentiate among turfgrass cultivars
314 receiving various N treatments.

315

316 **Please insert Figure 3 near here**

317

318

319 **Conclusions**

320 DGCI values were highly correlated with the nitrogen clipping content and NDVI with a
321 highly significant degree of association. The results suggested that UAS derived imagery RGB
322 photography by UAVs had a great potential in supporting decisions. Thus, DGCI could be a
323 promising remote-sensing tool for mapping the crop nitrogen status or NDVI at large scale with
324 high precision and low cost (Li et al. 2015). This method could be used by farmers operating in
325 large-scale farms to precisely manage the application of fertilizers, although the farmers especially
326 in the developing and underdeveloped counties, they do not have enough knowledge to operate the
327 UAV and manage the technology. As turfgrass, especially in the most developed countries, this
328 method could allow golf course superintendents and turf management specialists to make critical
329 decisions in real time without high up-front costs. Differences in camera quality and settings and
330 lighting conditions could affect DGCI and limit their utility in diagnosing N deficiencies.
331 Furthermore, disease, water status, nutritional deficiencies other than N, or different uniformity,
332 texture and growth habit may affect greenness regardless of N status as suggested also by Rorie,
333 Purcell, Karcher et al. 2011 and by Leinauer et al. 2014. More research is required on this
334 technology and on the Smartphone APP FieldScout GreenIndex+ Turf (Spectrum Technologies,
335 Inc., Aurora, IL, USA) (Spectrum Technologies, Inc. 2018) to study and overcome possible
336 discrepancies between the APP and the Smartphone camera. Although the accuracy of a
337 Smartphone camera is not comparable to a digital camera, the precision of a Smartphone camera
338 could still help to detect minor changes in turf greenness over time and-or relative to other areas of
339 the golf course or sports field. In fact, if the imagery was conveyed quickly to the user, a broader
340 usage of this technology could allow golf course superintendents and turf management specialists to
341 make critical decisions in real time without high up-front costs, in small areas. To use efficiently
342 this technology on large scale, DGCI could be use directly on board an UAV and could serve as an
343 indicator of N deficiency on turfgrass, thus increasing turfgrass nitrogen fertilization efficiency.

344 Indeed, applications installed in drones could be good solutions for farmers or golf course
345 superintendents and turf management specialists so they can adopt and benefit from DGCI
346 technology.

347 **Acknowledgements**

348 Authors wish to thank Mar. Project Srl, Geographike Srl, Geomind Srl and E.T.G. Srl staff for
349 the cooperation in research.

350

351 The present research was carried out within the Project "TurfUp_Remote monitoring and
352 management of turfgrass" co-financed under Tuscany ERDF ROP 2014-2020 Research
353 development and innovation Call 2.

354

355 **References**

- 356 Aguilar, C., J. C. Zinnert, M. J. Polo, and D. R. Young. 2012. “NDVI as an indicator for
357 changes in water availability to woody vegetation”. *Ecological Indicators* 23: 290–300.
- 358 Andrade-Sanchez, P., M. A. Gore, J. T. Heun, K. R. Thorp, A. E. Carmo-Silva, A. N. French,
359 M. E. Salvucci, and J. White. 2014. “Development and evaluation of a field-based high-
360 throughput phenotyping platform”. *Functional Plant Biology* 41 (1): 68-79.
- 361 Barton, C. V. 2012. “Advances in remote sensing of plant stress”. *Plant and Soil*; 354: 41–44.
- 362 Bausch, W. C., and H. R. Duke. 1996. “Remote sensing of plant nitrogen status in corn”.
363 *Transactions of the ASAE* 39: 1869-1875.
- 364 Bell, G. E., and X. Xiong. 2008. “The history, role, and potential of optical sensing for practical
365 turf management”. In P. Pessaraki (Ed.), *Handbook of turfgrass management and physiology*
366 (pp. 641–658). Boca Raton, FL: CRC Press.
- 367 Bell, G. E., B. M. Howell, G. V. Johnson, W. R. Raun, J. B. Solie, and M. L. Stone. 2004.
368 “Optical sensing of turfgrass chlorophyll content and tissue nitrogen”. *HortScience* 39(5):
369 1130–1132.
- 370 Bell, G. E., J. K. Kruse, and J. M. Krum. 2013. “The evolution of spectral sensing and advances
371 in precision turfgrass management”. In *Turfgrass: Biology, use and management*, edited by
372 J. C. Stier, B. P. Horgan, and S. A. Bonos, 1151–1188. Madison, WI: American Society of
373 Agronomy.
- 374 Blackmer, T. M., J. S. Schepers, G. E. Varvel, and G. E. Meyer. 1996 “Analysis of aerial
375 photography for nitrogen stress within corn fields”. *Agronomy Journal* 88 (5): 729-733.
- 376 Bremer, D. J., H. Lee, K. Su, and S. J. Keeley. 2011, “Relationships between normalized
377 difference vegetation index and visual quality in cool-season turfgrass: II. Factors affecting
378 NDVI and its component reflectances”. *Crop Science* 51(5), 2219-2227.

379 Bremner, J. M. 1965 “Total Nitrogen”. In *Methods of Soil Analysis*, Part 2. Agron. Monogr. no.
380 9, edited by C. A. Black, 1149–1178. Madison, WI: American Society of Agronomy.

381 Carrow, R. N., J. M. Krum, I. Flitcroft, and V. Cline. 2010. “Precision turfgrass management:
382 challenges and field applications for mapping turfgrass soil and stress”. *Precision*
383 *Agriculture* 11: 115–134.

384 Casadesus, J., and D. Villegas. 2014. “Conventional digital cameras as a tool for assessing leaf
385 area index and biomass for cereal breeding”. *Journal of integrative plant biology* 56 (1): 7-
386 14.

387 Casadesús, J., Y. Kaya, J. Bort, M. M. Nachit, J. L. Araus, S. Amor, G. Ferrazzano, et al. 2007.
388 “Using vegetation indices derived from conventional digital cameras as selection criteria for
389 wheat breeding in water-limited environments”. *Annals of Applied Biology* 150 (2): 227-
390 236.

391 Caturegli L., F. Lulli, L. Foschi, L. Guglielminetti, E. Bonari, and M. Volterrani. 2014.
392 “Monitoring turfgrass species and cultivars by spectral reflectance”. *European Journal of*
393 *Horticultural Science* 79 (3): 97–107.

394 Caturegli L., F. Lulli, L. Foschi, L. Guglielminetti, E. Bonari, and M. Volterrani. 2014.
395 “Turfgrass spectral reflectance: simulating satellite monitoring of spectral signatures of main
396 C3 and C4 species”. *Precision. Agriculture*. 6: 297–310.

397 Caturegli L., M. Corniglia, M. Gaetani, N. Grossi, S. Magni, M. Migliazzi, L. Angelini, M. et
398 al. 2016. “Unmanned Aerial Vehicle to estimate nitrogen status of turfgrasses”. *PLOS ONE*.
399 11(6), e0158268. doi:10.1371/journal.pone.0158268.

400 Caturegli, L., M. Casucci, F. Lulli, N. Grossi, M. Gaetani, S. Magni, E. Bonari, and M.
401 Volterrani. 2015. “GeoEye-1 satellite versus groundbased multispectral data for estimating
402 nitrogen status of turfgrasses”. *International Journal of Remote Sensing* 36: 2238–2251.

403 Caturegli, L., N. Grossi, M. Saltari, M. Gaetani, S. Magni, A. E. Nikolopoulou, E. Bonari, and
404 M. Volterrani. 2015. “Spectral reflectance of tall fescue (*Festuca Arundinacea* Schreb.)

405 under different irrigation and nitrogen conditions”. *Agriculture and Agricultural Science*
406 *Procedia* 4: 59–67.

407 Corwin, D. L., and S. M Lesch. 2005. “Apparent soil electrical conductivity measurements in
408 agriculture”. *Computers and electronics in agriculture* 46(1), 11-43.
409 doi:10.1016/j.compag.2004.10.005.

410 Dordas, C. 2008. “Role of nutrients in controlling plant diseases in sustainable agriculture: A
411 review”. *Agronomy for Sustainable Development* 28: 33–46. doi:10.1051/agro:2007051..

412 Dubbini, M. , A. Pezzuolo, M. D. Giglio, M. Gattelli, L. Curzio, D. Covi, T. Yezekyan, and F.
413 Marinello. 2017. “Last generation instrument for agriculture multispectral data collection”.
414 *Agricultural Engineering International: CIGR Journal* 19 (1): 87–93.

415 Fensholt, R., and S. R. Proud. 2012. “Evaluation of earth observation based global long term
416 vegetation trends-Comparing GIMMS and MODIS global NDVI time series”. *Remote*
417 *Sensing of Environment* 119: 131–147.

418 Fitz-Rodriguez, E., and C. Y. Choi. 2002. “Monitoring turfgrass quality using multispectral
419 radiometry”. *Transactions of the ASAE* 45 (3):865-867.

420 Fryskowska, A., M. Kedzierski, A. Grochala, and A. Braula. 2016. “Calibration of low cost
421 RGB and NIR UAV cameras”. *International Archives of Photogrammetry, Remote Sensing*
422 *and Spatial Information Sciences* 41: 817-821.

423 Graeff, S., and W. Claupein. 2003. “Quantifying nitrogen status of corn (*Zea mays* L.) in the
424 field by reflectance measurements”. *European Journal of Agronomy* 19 (4): 611–618.

425 Grossi N., M. Fontanelli, E. Garramone, A. Peruzzi, M. Raffaelli, M. Pirchio, L. Martelloni, et
426 al. 2016. “Autonomous mower saves energy and improves quality of tall fescue
427 lawn”. *HortTechnology*, 26 (6): 825-830.

428 Gupta, S. G., M. M. Ghonge, and P. M. Jawandhiya. 2013. “Review of unmanned aircraft
429 system (UAS)”. *International Journal of Advanced Research in Computer Engineering &*
430 *Technology* 2 (4): 1646–1658.

431 Hansen, P. M., and J. K. Schjoerring. 2003. "Reflectance measurement of canopy biomass and
432 nitrogen status in wheat crops using normalized difference vegetation indices and partial
433 least squares regression". *Remote Sensing of Environment* 86: 542–553.

434 Huang, J, F. He, K. Cui, R. J. Buresh, B. Xu, W. Gong, and S. Peng. 2008. "Determination of
435 optimal nitrogen rate for rice varieties using a chlorophyll meter". *Field Crops Research*
436 105: 70–80.

437 Jiang, Y, and R. N. Carrow. 2007. "Broadband spectral reflectance models of turfgrass species
438 and cultivars to drought stress". *Crop Science* 47: 1611–1618.

439 Johnsen, A. R., B. P. Horgan, B. S. Hulke, and V. Cline. 2009. "Evaluation of remote sensing to
440 measure plant stress in Creeping Bentgrass (L.) fairways". *Crop Science* 49: 2261–2274.

441 Karcher, D. E., and M. D. Richardson. 2003. "Quantifying turfgrass color using digital image
442 analysis". *Crop Science* 43: 943-951.

443 Karcher, D. E., and M. D. Richardson. 2013. "Digital image analysis in turfgrass research". In
444 *Turfgrass: Biology, use and management*, edited by J. C. Stier, B. P. Horgan, and S. A.
445 Bonos, 1133–1149. Madison, WI: American Society of Agronomy.

446 Krum, J. M., R. N. Carrow, and K. Karnok. 2010. "Spatial mapping of complex turfgrass sites:
447 Site-specific management units and protocols". *Crop Science* 50: 301–315

448 Laliberte, A. S., and A. Rango. 2011. "Image processing and classification procedures for
449 analysis of sub-decimeter imagery acquired with an unmanned aircraft over arid
450 rangelands". *GIScience & Remote Sensing* 48 (1): 4–23.

451 Leinauer, B., D. M. VanLeeuwen, M. Serena, M. Schiavon, and E. Sevostianova. 2014. "Digital
452 image analysis and spectral reflectance to determine turfgrass quality". *Agronomy Journal*
453 106 (5): 1787-1794.

454 Li, J., F. Zhang, X. Qian, Y. Zhu, and G. Shen. 2015. "Quantification of rice canopy nitrogen
455 balance index with digital imagery from unmanned aerial vehicle". *Remote Sensing Letters*
456 6: 183–189.

457 Ma, B. L., M. J. Morrison, and L. M Dwyer. 1996. "Canopy light reflectance and field
458 greenness to assess nitrogen fertilization and yield of maize". *Agronomy Journal* 88: 915–
459 920.

460 Magni S., M. Gaetani, N. Grossi, L. Caturegli, S. La Bella, C. Leto, G. Virga, T. Tuttolomondo,
461 F. Lulli and M. Volterrani. 2014. "Bermudagrass adaptation in the Mediterranean climate:
462 phenotypic traits of 44 accessions". *Advances in Horticultural Science* 28 (1): 29-34.

463 Morris, K. N., and R. C. Shearman. 2008. "NTEP turfgrass evaluation guidelines". In NTEP
464 turfgrass evaluation workshop, 1–5. Beltsville, MD. Available:
465 <http://www.ntep.org/cooperator.html>. Accessed August 2018.

466 O'Brien, D. 2017. "New technologies for evaluating putting green surface characteristics". Ph.D
467 dissertation, University of Arkansas, AR, USA. Available from
468 <https://scholarworks.uark.edu/etd/2614/>

469 Perry, E. M., and J. R. Davenport. 2007. "Spectral and spatial differences in response of
470 vegetation indices to nitrogen treatments on apple". *Computer and Electronics in*
471 *Agriculture* 59 (1-2): 56–65. doi:10.1016/j.compag.2007.05.002.

472 Quantum GIS 2.18 software. Accessed July, 2018. <https://www.qgis.org>

473 Remondino, F., and C. Fraser. 2006. "Digital camera calibration methods: considerations and
474 comparisons". *International Archives of Photogrammetry, Remote Sensing and Spatial*
475 *Information Sciences* 36 (5): 266-272.

476 Rhezali, A., and R. Lahlali. 2017. "Nitrogen (N) mineral nutrition and imaging sensors for
477 determining N status and requirements of maize". *Journal of Imaging* 3(4): 51.

478 Rorie, R. L., L. C. Purcell, D. E. Karcher, and C. A. King. 2011. "The assessment of leaf
479 nitrogen in corn from digital images". *Crop Science* 51 (5): 2174-2180.

480 Rorie, R. L., L. C. Purcell, M. Mozaffari, D. E. Karcher, C. A. King, M. C. Marsh, and D. E.
481 Longer. 2011. "Association of "greenness" in corn with yield and leaf nitrogen
482 concentration". *Agronomy Journal* 103(2): 529-535.

483 SAL engineering MAIA S2 (Modena Italy; EOPTIS, Trento, Italy). Accessed July, 2018.
484 <https://www.salengineering.it/public/en/p/maia.asp>

485 Samborski, S. M., N. Tremblay, and E. Fallon. 2009. “Strategies to make use of plant sensors-
486 based diagnostic information for nitrogen recommendations”. *Agronomy Journal* 101 (4):
487 800–816.

488 Scharf, PC, and J. A. Lory. 2009. “Calibrating reflectance measurements to predict optimal
489 sidedress nitrogen rate for corn”. *Agronomy Journal* 101 (3): 615-625.

490 Spectrum Technologies, Inc. 2018. Product manual, Item no. 2910TA, 2910T.
491 Available:http://www.specmeters.com/assets/1/22/2910TA_2910T_GreenIndex__Turfl.pdf
492 Accessed 11 Sept. 2018.

493 Trenholm, L. E., R. N. Carrow, and R. R. Duncan. 1999. Relationship of multispectral
494 radiometry data to qualitative data in turfgrass research. *Crop Science* 39: 763–769.

495 Vergara-Díaz, O., M. A. Zaman-Allah, B. Masuka, A. Hornero, P. Zarco-Tejada, B. M.
496 Prasanna, J. E. Cairns and J. L. Araus. 2016. “A novel remote sensing approach for
497 prediction of maize yield under different conditions of nitrogen fertilization”. *Frontiers in*
498 *Plant Science* 7:666. doi: 10.3389/fpls.2016.00666.

499 Volterrani, M., N. Grossi, L. Foschi, and S. Miele. 2005. “Effects of nitrogen nutrition on
500 bermudagrass spectral reflectance”. *International Turfgrass Society Research Journal* 10:
501 1005–1014.

502 Walters, D. R., and I. J. Bingham. 2007. “Influence of nutrition on disease development caused
503 by fungal pathogens: implications for plant disease control.” *Annals of Applied Biology* 151
504 (3): 307–324. doi:10.1111/aab.2007.151.issue-3.

505 White, J. W., P. Andrade-Sanchez, M. A. Gore, K. F. Bronson, T. A. Coffelt, M. M. Conley, K.
506 A. Feldmann, et al. 2012. “Field-based phenomics for plant genetics research”. *Field Crops*
507 *Research* 133: 101-112.

508 Xiang, M., J. Q. Moss, D. L. Martin, and Y. Wu. 2018. “The salinity tolerance of seeded-type
509 common bermudagrass cultivars and experimental selections”. *HortTechnology* 28(3): 276-
510 283.

511 Xiang, M., J. Q. Moss, D. L. Martin, K. Su, B. L. Dunn, and Y. Wu. 2017. “Evaluating the
512 salinity tolerance of clonal-type bermudagrass cultivars and an experimental selection”.
513 *HortScience* 52 (1): 185-191.

514 Xiong, X., G. E. Bell, J. B. Solie, M. W. Smith, and B. Martin. 2007. “Bermudagrass seasonal
515 responses to nitrogen fertilization and irrigation detected using optical sensing”. *Crop*
516 *Science* 47 (4): 1603–1610.

517 Zhang, C., and J. M. Kovacs. 2012. “The application of small unmanned aerial systems for
518 precision agriculture: a review”. *Precision Agriculture* 13: 693–712.

519 Zhou, B., A. Elazab, J. Bort, O. Vergara, M. D. Serret, and J. L. Araus. 2015. “Low-cost
520 assessment of wheat resistance to yellow rust through conventional RGB
521 images”. *Computers and electronics in agriculture* 116: 20-29.

522

523 **Captions**

524 **Table 1.** Instrument monochrome sensors with relative band-pass filters of the multispectral camera
525 MAIA.

526 **Table 2.** Pearson product-moment correlation coefficients (*r*) among clipping nitrogen content,
527 color intensity, turfgrass quality, plant water content (PWC), NDVI measured with a handheld crop
528 sensor (GreenSeeker) and NDVI measured with multispectral camera mounted on an unmanned
529 aerial vehicle (UAV) and dark green color index (DGCI) on a) bermudagrass hybrid; b) tall fescue.
530 For each species correlation coefficients are calculated across all entries.

531 All values are significant at the 0.010 level, except for DGCI color intensity, quality and PWC for
532 bermudagrass hybrid and DGCI quality for tall fescue, whose values are significant at the 0.001
533 level.

534 **Figure 1.** (a) UAV during flight operations (6 July 2017; Pisa, Italy; 43°40'N, 10° 19'E, 6 m. a. s.
535 1.); (b) The multispectral camera MAIA mounted on the UAV.

536 **Figure 2.** Linear relationship between NDVI measured with a handheld crop sensor (GreenSeeker)
537 and NDVI measured with a multispectral camera mounted on UAV and DGCI on (a) bermudagrass
538 hybrid; and (b) tall fescue. Values represented the 3 replications.

539 **Figure 3.** Linear relationship between DGCI and the clipping nitrogen content (%) on (a)
540 bermudagrass hybrid; and (b) tall fescue. Values represented the average of 3 replications.

541

542

543 **Tables**

544 **Table 1.** Instrument monochrome sensors with relative band-pass filters of the multispectral camera
 545 MAIA.

Wavelength (nm)		
Start	Central	Stop
395.0	422.5	450.0
455.0	487.5	520.0
525.0	550.0	575.0
580.0	602.5	625.0
630.0	660.0	690.0
705.0	725.0	745.0
750.0	785.0	820.0
825.0	887.5	950.0

546
 547 **Table 2.** Pearson product-moment correlation coefficients (*r*) among clipping nitrogen content,
 548 color intensity, turfgrass quality, plant water content (PWC), NDVI measured with a handheld crop
 549 sensor (GreenSeeker) and NDVI measured with multispectral camera mounted on an unmanned
 550 aerial vehicle (UAV) and dark green color index (DGCI) on a) bermudagrass hybrid; b) tall fescue.
 551 For each species correlation coefficients are calculated across all entries.
 552 All values are significant at the 0.010 level, except for DGCI color intensity, quality and PWC for
 553 bermudagrass hybrid and DGCI quality for tall fescue, whose values are significant at the 0.001
 554 level.

<i>r</i>	Color intensity	Quality	PWC	NDVI GreenSeeker	NDVI UAV	DGCI
a) Bermudagrass hybrid						
N clipping (%)	0.97	0.97	0.95	0.94	0.92	0.86
Color intensity (1-9)	N/A	0.94	0.99	0.94	0.94	0.83
Quality (1-9)	N/A	N/A	0.97	0.97	0.94	0.84
PWC (%)	N/A	N/A	N/A	0.92	0.92	0.84
NDVI GreenSeeker (780,660)	N/A	N/A	N/A	N/A	0.96	0.91
NDVI UAV (830,660)	N/A	N/A	N/A	N/A	N/A	0.85
b) Tall fescue						
N clipping (%)	0.99	0.99	0.99	0.95	0.94	0.95
Color intensity (1-9)	N/A	0.99	0.99	0.96	0.95	0.95
Quality (1-9)	N/A	N/A	0.98	0.94	0.93	0.92
PWC (%)	N/A	N/A	N/A	0.98	0.98	0.98
NDVI GreenSeeker (780,660)	N/A	N/A	N/A	N/A	0.99	0.95
NDVI UAV (830,660)	N/A	N/A	N/A	N/A	N/A	0.96

555

556