



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Environmental correlates of growth patterns in Neolithic Liguria (northwestern Italy)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Environmental correlates of growth patterns in Neolithic Liguria (northwestern Italy) / Dori I., Varalli A., Seghi F., Moggi-Cecchi J., Sparacello V.S.. - In: INTERNATIONAL JOURNAL OF PALEOPATHOLOGY. - ISSN 1879-9817. - ELETTRONICO. - 28:(2020), pp. 112-122. [10.1016/j.ijpp.2019.12.002]

Availability:

This version is available at: <https://hdl.handle.net/11585/850072> since: 2023-08-25

Published:

DOI: <http://doi.org/10.1016/j.ijpp.2019.12.002>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Dori, I., Varalli, A., Seghi, F., Moggi-Cecchi, J. & Sparacello, V. S. Environmental correlates of growth patterns in Neolithic Liguria (northwestern Italy). *International Journal of Paleopathology* 28, 112–122 (2020).

The final published version is available online at:
<https://doi.org/10.1016/j.ijpp.2019.12.002>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Environmental correlates of growth patterns in Neolithic Liguria (northwestern Italy)

Dori I^{1,2,3}, Varalli A⁴, Seghi F², Moggi-Cecchi J², Sparacello VS^{1*}.

¹Univ. Bordeaux, CNRS, MC, PACEA, UMR 5199, 33615 Pessac, France

² Department of Biology, Laboratory of Anthropology, University of Florence, via del Proconsolo 12, 50122 Florence, Italy

³ Soprintendenza Archeologia, Belle Arti e Paesaggio per le province di Verona, Rovigo e Vicenza, Piazza S. Fermo 3, 37121 Verona, Italy

⁴ Department of Archaeology, Durham University, UK

* Corresponding Author: Vitale Stefano Sparacello. PACEA - UMR 5199 Université de Bordeaux, Bâtiment B8 Allée Geoffroy Saint Hilaire, CS 50023 33615 PESSAC CEDEX. E-mail: vitale.sparacello@u-bordeaux.fr; vitosparacello@gmail.com; grifis1979@gmail.com.

Abstract

Objective: This study evaluates patterns of human growth in the Neolithic to make inferences about environmental correlates of developmental disturbances.

Materials: 33 children/adolescents from the Neolithic of Liguria (Italy), 29 of which date between 4,800-4,400 cal BCE.

Methods: Neolithic patterns of growth are compared with a modern sample (the Denver Growth Study; DGS). Dental development was used to determine age at death. Proxies for postcranial maturation are femoral length, and proportion of mean adult femoral length attained.

Results: Ligurian children show growth faltering compared to DGS, especially between 4 and 9 years of age. Between 1 and 2 years, and in later childhood and adolescence, values are more similar, or higher than DGS, when using the proportion of adult femoral length attained.

Conclusions: The pattern of growth in Ligurian Neolithic children may reflect a deprived and highly-infectious environment: three individuals show skeletal lesions suggestive of tuberculosis. The relatively faster growth in infancy may result from the buffering provided by maternal milk. Older children and adolescents may exhibit catch-up growth.

Significance: This study contributes to our understanding of Neolithic selective pressures and possible biocultural adaptive strategies.

Limitations: The cross-sectional nature of the data, and the small sample size, make unclear whether the observed pattern is representative of the growth patterns in the living population. The possibility that adults are stunted undermines the interpretation of optimal growth in the first years.

Suggestions for Further Research: Refine age estimates, increase sample size through the study of other bone elements.

Keywords

Growth disturbances; stunting; Neolithic Transition; life history; tuberculosis.

1. Introduction

Developing individuals subjected to high levels of environmental stress, such as poor nutrition (Johnston et al., 1976; Martorell, 1985), infectious disease and parasites (Solomons et al., 1993; Stephenson, 1987), or even psycho-social stress (Powell et al., 1967), may exhibit slower rates of growth, delayed maturation, prolonged growth, and smaller final adult size (Bogin, 1988, 1999; Ulijaszek et al., 1998). Although genetic factors interact with the environment in the expression of body size and rate of growth (e.g. Eveleth and Tanner, 1990; Johnston et al., 1976), studying growth trajectories at a population level provides insights about variation in environmental correlates of growth disturbances, such as warfare, famines, and socioeconomic parameters (Powell et al., 1967; Ulijaszek et al., 1998).

In bioarchaeology, the study of growth trajectories and their interpretation, including comparisons with modern reference samples, faces a number of problems (reviews in Lewis, 2007; Saunders, 2008). Problems related to archaeological assemblages include the differential preservation of juvenile remains (Bello et al., 2006), the cross-sectional rather than longitudinal nature of data (Humphrey, 2003), the representativeness of data due to the osteological paradox (Goodman, 1993; Saunders and Hoppa, 1993; Wood et al., 1992), and the temporal resolution of the data (Albanese, 2002, 2009). Methodological problems mainly relate to age estimation based on dental development (Saunders, 2008). Nevertheless, numerous studies since the early work by Johnston (1962) have explored patterns of growth in prehistoric and more recent skeletal assemblages (reviews in Humphrey, 2000, 2003; Larsen, 2015; Lewis, 2007; Mays, 2018; Saunders, 2008), and made inferences on environmental

correlates of developmental disturbances, such as malnutrition and disease, as well as the effect of migration, colonialism, and changing subsistence patterns.

The Neolithic Transition, i.e. is the adoption of a production economy based on the domestication of plants and animals, is “*one of the fundamental structural processes of human history*” (Bocquet-Appel, 2011a, 2011b), and dramatically changed several aspects of the human experience. Various studies suggest that the Neolithic Transition was accompanied by a worsening in health status and well-being, resulting in an increase of osteological markers of stress (e.g. *cribra orbitalia* and enamel hypoplasia; Armelagos et al., 2005; Cohen and Armelagos, 1984; Cohen and Crane-Kramer, 2007; Eshed et al., 2010; Gleń-Haduch et al., 1997; Larsen, 1995; Temple, 2010). Mortality rates increased, especially in infants (Armelagos et al., 2009; Bocquet-Appel 2002, 2009, 2011a, 2011b; Page et al., 2016; Pérez-Losada and Fort, 2010), possibly due to unsanitary and deprived conditions and a more infectious environment (Armelagos et al., 1991, 1996, 2005). In addition, it has been suggested that certain Neolithic infant feeding practices may have had a negative impact on children survivorship (Pearson et al., 2010). Exploring patterns of growth in Neolithic children can integrate data coming from paleopathological and paleodemographic sources, and can contribute to the debate on biocultural Neolithic adaptive strategies.

The prehistoric skeletal series from Liguria (northwestern Italy) represent an important source of information on the past peopling of the northwestern Mediterranean from the Upper Paleolithic to the Metal Ages (Del Lucchese, 1997; Formicola et al., 2005; Maggi, 1997; Sparacello et al., 2018). Evidence of Neolithic occupation comes from several caves and rock shelters opening in the Finalese area, where renowned sites such as Arene Candide have yielded detailed stratigraphic successions (Arobba et al., 2017; Maggi, 1997; Tiné, 1999). About 200 burials and an undefined number of scattered human remains have been reported from these sites (e.g. Del Lucchese, 1997; Delfino, 1981; Issel, 1908; Panelli and Rossi, 2015, 2017; Parenti and Messeri, 1962; Richard, 1942; Sparacello et al., 2018, 2019). However, a large portion of the skeletal series had been excavated since the mid-19th century, and was accompanied by little information about the depositional context (De Pascale, 2007, 2008; Rossi et al., 2014). Only recently, a large-scale campaign of direct dating on Ligurian human remains has been conducted (project BUR.P.P.H., PI VSS and DEN.P.H., PI ID), and allowed for the chronological characterization of most burials and individuals reconstructed from the scattered remains (Sparacello et al., 2018, 2019, and in review).

This study explores patterns of growth in Ligurian Neolithic children through the analysis of femoral length in individuals whose age at death was independently estimated from dental development. The comparison with a reference sample of modern, healthy, and well-nourished children (the Denver Growth Study; Maresh, 1943, 1955, 1970) will allow for an evaluation of well-being among these early agriculturalists, as done in previous studies (e.g. Harrington and Pfeiffer, 2008; Humphrey 2000, 2003; Pfeiffer and Harrington, 2010). Subsistence of Neolithic Ligurian people was based on a variety of domesticated plants (Arobba et al., 2017; Nisbet, 2008), and on livestock breeding, especially sheep (Macphail et al., 1997; Rowley-Conwy, 1992, 1997, 1998), in a highly mountainous environment, resulting in strenuous physical activity and high logistic mobility levels (Marchi et al., 2006, 2011; Sparacello and Marchi, 2008; Sparacello et al., 2011, 2014). Environmental stress and poor health conditions are suggested by a high prevalence of enamel hypoplasia (Formicola, 1987; Orellana-Gonzales et al., in review), and by several cases of osteoarticular tuberculosis (Canci et al., 1996; Formicola et al., 1987; Sparacello et al., 2017; Sparacello et al., 2018, and unpublished data, see below), a highly infectious, debilitating, and growth-impairing disease (Sparacello et al., 2016). We expect that environmental hardships experienced by Neolithic Ligurian people will result in a pattern of development showing growth faltering when compared to a modern industrialized sample.

2. Materials and Methods

The sub-adults included in this study consist of 33 individuals spanning birth to late adolescence, chronologically belonging to the Neolithic of Liguria (c. 5800-3800 cal. BCE). We included only the individuals that were directly dated (Table 1; Sparacello et al., in review), and the vast majority (29/33) chronologically overlap with the Square Mouthed Pottery Culture (SMP; c. 5000-4300 cal. BCE; Binder and Sénépart, 2010; Del Lucchese and Starnini, 2015). Three individuals from Arma dell'Aquila chronologically overlap with the earlier Impresso-Cardial Complex (ICC; Binder et al., 2017; Sparacello et al., 2019), and one from Grotta Pollera with the later Chasséen (4300-3700 BCE; Crepaldi, 2001; Maggi, 1997). All remains were unearthed from six caves (Arene Candide Cave, Arma dell'Aquila, Grotta marina di Bergeggi, Grotta dei Pipistrelli, Grotta Pollera, Arma Strapatente), situated within a radius of 5 km in the Finalese area (Figure 1). Data was collected at the Museo di Archeologia Ligure, Genova, at the Museo Archeologico del Finale, Finale Ligure, and at the Museo di Storia Naturale – Sezione di Antropologia e Etnologia, Università degli Studi di Firenze. All

the available remains belonging to the Neolithic of Liguria were surveyed, with the exception of the four subadults housed at the Museo delle Civiltà in Rome.

[Figure 1 about here]

For all children below the age of 12, age at death estimates are based on dental mineralization following AlQahtani et al. (2010). Dental formation shows considerably less populational variation than eruption, and is less influenced by environmental factors (Demirjian, 1986; review in Saunders, 2008). However, the accuracy of age at death determination is influenced by the number of teeth that could be examined. Dental eruption following AlQahtani et al. (2010) was used when mineralization could not be assessed due to impossibility of examining the root (two adolescent individuals). All age estimates and information on tooth development is available in Supplementary Information S1. Due to the small sample size for each age class, we decided to report the results using the midpoint of the age estimate for each individual.

Femoral maximum length was measured following the standards in the field (e.g. Schaefer et al., 2009) using a digital caliper and an osteometric table, as appropriate. Length was measured without epiphyses in children, and with the epiphysis in adolescents above 12 years of age (Ruff, 2007). Unfortunately, in two adolescents (e.g. Pollera 1 PE and Pollera 34 PE) one or both epiphyses were plastered to the metaphysis, and a slight overestimation of the maximum length cannot be excluded (Table 1; Pollera 34 PE was not included in the analysis due to absence of the distal epiphysis, and its measurements are in Supplementary Information S2).

[Table 1 about here]

The growth pattern in our sample was compared with the standards derived from the Denver Growth Study, which consists of longitudinal bone length data from healthy children, deriving from radiographs collected in the United States between 1935-1967 (Maresh, 1943, 1955, 1970), and commonly used in comparisons with prehistoric populations (review in Mays, 2018). The mean of male and female diaphyseal length was used, due to the absence of sex estimation for most of our immature remains.

Following previous research (e.g. Harrington and Pfeiffer, 2008; Humphrey 2000, 2003; Ives and Humphrey, 2017; Pfeiffer and Harrington, 2010), we compared absolute length for age, and relative length as a percentage of the mean adult length (pooled sexes) for the same population (for the Neolithic Ligurian sample: $n = 25$; males = 13; females: 12; average

femur M1 = 410.5 mm; for the Denver Growth Study: $n = 68$; femur M1 = 489 mm; Harrington and Pfeiffer, 2008).

It should be noted that the values of absolute length for age are biased by radiographic magnification, proportionally to bone size (Feldesman, 1992; Humphrey, 2003). Ruff (2007) proposed two regression equations correcting the radiographic measurements made on the original x-rays of the Denver Growth Study, and applied it to a smaller sample of 20 individuals (10 males and 10 females). The data points obtained for the Ligurian sample were therefore plotted against: 1) the mean and $\pm 1/2$ SD absolute length between birth and 12 years of age of the Denver Growth Study (Maresh, 1955, 1970), corrected for radiographic magnification using the formulae proposed by Ruff (Ruff, 2007:702); 2) the mean and $\pm 1/2$ SD absolute length between 1 and 17 years of age of the subsample studied by Ruff, based on the data presented in his Table 1 (Ruff, 2007:702).

The proportion of adult age attained at a given age is only minimally influenced by radiographic magnification. Ligurian data were plotted against the mean and $\pm 1/2$ SD proportion of adult size attained between birth and 17 years of age of 1) the Denver Growth Study (Maresh, 1955, 1970; see also Humphrey, 2000; Pfeiffer and Harrington, 2010), and 2) the subsample studied by Ruff, based on the data presented in his Table 1 (Ruff, 2007:702), and correcting the adult length of the Denver Growth Study using the regression for bones longer than 217 mm (adjusted = $0.949 \times \text{original length} + 5.63$).

In addition, we calculated a residual relative to the proportion of adult femur length achieved, when comparing the Ligurian Neolithic people with the Denver Growth Study (cf. Harrington and Pfeiffer, 2008; Humphrey 2000, 2003; Pfeiffer and Harrington, 2010). The proportion of adult femur length achieved was estimated graphically for the age points of the Ligurian Neolithic children (Supplementary Information S2), and the residuals were plotted on the mean and $\pm 1/2$ SD proportion of adult size attained by the Denver sample between birth and 12 years of age, redrawn with the mean as a horizontal line (e.g. Humphrey, 2000, 2003).

3. Results

Figure 2 shows the absolute maximum length of the femur in the Ligurian Neolithic sample, when compared with the data of the Denver Growth Study, corrected for radiographic magnification following Ruff (2007), and the subsample from the Denver Growth Study selected and corrected by Ruff (2007). In both graphs, the Ligurian Neolithic growth pattern tends to fall consistently two standard deviations below the mean.

[Figure 2 about here]

Figure 3 shows the pattern of growth expressed as proportion of mean adult length attained in the Ligurian Neolithic sample, when compared with the data of the Denver Growth Study, and the sample from the Denver Growth Study selected by Ruff (2007). The Ligurian Neolithic sample now falls close to the mean, or above, of the Denver Growth Study up to the age of 2.5, and consistently below between the ages of 4.5 and 9. Most of older children and adolescents fall close or above the mean of the Denver Growth Study, except for one who is more than two standard deviations below the mean. The slight differences observed between the whole Denver sample and the results by Ruff (2007) are mostly due to the greater standard deviations in the latter study, which is based on a smaller sample size ($n = 20$ compared to $n = 70-80$ in Maresh, 1970; Humphrey, 2000).

[Figure 3 about here]

The pattern from Figure 3 is visible more clearly in Figure 4, showing the residual relative to the proportion of adult femur length achieved, when comparing the Ligurian Neolithic sample (only measurements without epiphyses) with a) the Denver Growth Study, and b) the sample from the Denver Growth Study selected by Ruff (2007). Neolithic Ligurian individuals tend to fall consistently below the mean of the Denver Growth Study between 4.5 and 8.5 years of age, and in three cases more than two standard deviations below the mean (Figure 4a). Note that Ruff (2007) did not present data for individuals below 1 years of age. Also, the larger standard deviation using data from Ruff's (2007) study is probably influenced by the smaller sample size (see above).

[Figure 4 about here]

4. Discussion

Through the comparison of dental maturation and skeletal development in a bioarchaeological sample of children and juveniles, this study aimed at the evaluation of Neolithic growth disturbances in Liguria (Italy), and at discerning their possible environmental correlates. This study has the advantage of a very narrow geographical focus, being all sites in a radius of a few kilometers, and most likely belonging to the same agropastoral system (Rowley-Conwy, 1992). In addition, most individuals belong to a precise chronological phase of human occupation, with dates primarily spanning c. 4800-4400 BCE at 2σ , when the Square Mouthed Pottery culture was attested in the region (Maggi et al., 1997). The few individuals

belonging to earlier or later chrono-cultural phases do not seem to deviate significantly from the general pattern observed. Although important caveats should be taken into account in any bioarchaeological study, especially when patterns of development are involved (review in Saunders, 2008), the Ligurian sample can be considered representative of a Neolithic population, with presumably a well-defined set of biocultural adaptations to specific environmental challenges.

When compared to the reference sample of the Denver Growth Study, the Ligurian Neolithic people tend to have shorter femora during growth, especially when considering absolute femoral length, and less markedly when length is expressed as a proportion of mean adult size. In addition, using the latter method, a majority of Ligurian individuals falls above the reference sample in early life (between birth and c. 2.5 years), indicating a longer femur for their age compared to Denver, followed by a clear downward deflection between 4.5-9 years. Considering that the Denver sample is composed by healthy and well-nourished modern children, which are assumed to have attained their full growth potential, the pattern observed in the Ligurian Neolithic sample would indicate optimal development in early life, followed by growth faltering.

In previous studies, some prehistoric populations showed a similar pattern: Humphrey (2003) observed that the three Native North American samples included in her review (Knoll, Libben, and San Cristóbal samples; Johnston, 1962; Lovejoy et al., 1990; Ryan, 1976) exhibited relatively longer femoral length than Denver during infancy (until about 1.5 years), followed by a dramatic reversal in the growth trajectory, and discussed the possibility of a genetic component influencing rates of skeletal growth and/or dental maturation. However, the different timing of the growth rate reversal in the three groups suggested that additional environmental factors, such as infant early feeding practices and the weaning process, might have contributed to varying growth trajectories (Humphrey, 2000, 2003).

Maternal milk supplies passive immunity, and weaning increases the pathogen load, requiring a sudden energetic investment into immune defense by the child (McDade, 2003; McDade and Worthmann, 1999). Therefore, various studies have associated the onset of deficits in growth with the cessation of breast feeding (e.g. Mays, 2010; Humphrey, 2000). For example, in the 18th and 19th century London, a widespread practice of early breast feeding cessation was introduced, due to social and cultural factors (Fildes, 1986; 1995; Nitsch et al., 2011), and resulted in growth deficits (beginning around 8 months of age, and becoming marked by 15 months; Humphrey, 2000; Ives and Humphrey, 2017), high prevalence of enamel defects in

teeth (King et al., 2002), and high infant mortality (Humphrey et al., 2012). An even earlier onset of growth faltering was attributed to deprived nutritional status of the mother, or poor quality of early supplementary foods (Humphrey, 2003). Mays (2010), noted how the cessation of breast feeding in a medieval sample between 1-2 years of age (estimated via isotopic analysis) “marks the start of a general pattern of deficient growth” (Mays, 2010:69; see also Mays, 2007). Although the link between breast feeding cessation and growth faltering is speculative, given the small sample size, the pattern observed in the Ligurian Neolithic sample, with growth retardation apparent only after c. 2.5 years of age, seems to be compatible with the direct estimation of breast feeding duration in two SMP Neolithic individuals (including the adolescent individual Arene Candide V BB studied here; Goude et al., in review). The isotopic profiles ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) suggest that breast feeding extended into the third year of life, as observed also in other Neolithic groups (Cienkosz-Stepanczak et al., 2017; Fernández-Crespo et al. 2018; Howcroft, 2013; Howcroft et al., 2014; Pearson et al., 2010, 2015; Scharlotta et al., 2018). The presence of a period of metabolic stress around 2.5-3.5 years of age is also supported by a significant increase in linear enamel hypoplasia prevalence in the same Ligurian SMP Neolithic sample (Orellana-Gonzales et al., in review). These multiple signals of developmental disturbances – in correspondence with the estimated end of passive immunity – may be due to poor nutrition and/or increased pathogen load. Although it is difficult to quantitatively assess the caloric intake of SMP Neolithic people, their diet included a significant component of animal protein (Le Bras-Goude et al., 2006; Goude et al., 2014), which was also probably used as a weaning food (Goude et al., *in review*). In this context, the effect of disease may have been relatively more important, as we discuss further below.

It could be argued that higher relative femoral length in early infancy may be influenced by a combination of two factors: relatively low variation in neonatal size (e.g. Leary et al., 2006), and markedly small adult size between the Denver sample and prehistoric populations (mean maximum length of the femur in the pooled sex Ligurian sample is 74 mm smaller). If differences in adult size were not entirely due to genetic differences, but also to stunting due to later metabolic stress, the pattern observed here in children below 2.5 years may not be a reflection of rapid growth, and in general absolute femoral length patterns may be more informative. Although it is difficult to determine the degree to which Ligurian adults attained their full growth potential, body proportions in the European Neolithic were markedly different than in modern times (e.g. Ruff et al., 2006), and Ligurian Neolithic adults do not

appear to be significantly smaller than other contemporary Mediterranean Neolithic populations (e.g. Rosenstock et al., 2019). The use of percent of adult size attained is generally advised in these contexts (Hoppa and Fitzgerald, 1999; Humphrey, 2003), but further research is necessary to independently assess the nutritional and developmental status of Ligurian Neolithic infants.

Regardless of the method used to compare Ligurian children with the Denver sample, it is clear that, by the age of 4.5 years, Ligurian Neolithic children are experiencing growth faltering. All individuals fall below the mean of the Denver sample, the majority being between -1 and -2 standard deviations (percent of adult size attained) or well below -2 standard deviations (absolute femoral length). Previous bioarchaeological studies comparing growing individuals from prehistoric groups with the Denver sample almost invariably show a growth deficit in the former (reviews in Humphrey, 2000, 2003; Larsen, 2015; Lewis, 2007; Mays, 2018; Saunders, 2008). However, the variation in the prehistoric patterns can be quite marked, and is assumed to reflect different social and environmental condition, in addition to genetic differences, as happens in contemporary groups (Bogin, 1988; Eveleth and Tanner, 1990). The more apparent growth deficit in agriculturalists when compared to hunter-gatherers has been attributed to their lesser reliance on animal protein (Cook, 1979, 1984; Goodman, 1998; Larsen et al., 2002). Within agriculturalists, the overreliance on staple foods with poor nutritional properties (e.g. maize), seem to coincide with poor growth (Cook, 1984; Goodman et al., 1984). In medieval Croatia, agriculturalist groups from the inland show lower long bone length at the same age than pastoralist communities from the coast (Pinhasi et al., 2014). Archaeological evidence suggests that the subsistence of Ligurian Neolithic people had a strong pastoral component (Macphail et al., 1997; Rowley-Conwy, 1992, 1997, 1998), which led to the consumption of animal protein since early life, as suggested by isotopic studies (Le Bras-Goude et al., 2006; Goude et al., 2014, and in review). The nutritional status of Ligurian Neolithic people may therefore have been relatively good for a prehistoric group, yet the pattern of growth faltering appears to be among the most relevant when compared with other bioarchaeological populations (cf. Humphrey 2003). Few studies on growth patterns have been conducted on Neolithic skeletal series from western Eurasia, probably due to lack of reasonably numerous samples of children. Pinhasi et al. (2011) found similar growth patterns between the lower limbs of the Denver sample and a small sample of Greek Neolithic children. The larger skeletal series from Çatalhöyük (Turkey), which is comparable to the Ligurian one in terms of chronology, diet, and subsistence (e.g. Pearson et al., 2015; Richards,

2003), shows growth patterns during childhood that are in line with the Denver sample (Ruff et al., 2013), and in general appears to have had a good health status (Hillson et al., 2013; Larsen et al., 2019).

In fact, in addition to dietary deprivation, health status and infectious load are considered a major influence on growth patterns in modern and prehistoric populations (Bogin, 1988; Larsen, 2015; Mays, 2018; Stephenson, 1999; Stinson, 2000). In previous bioarchaeological studies, disease burden was inferred based on increased sedentism or contact with European-introduced infectious disease (e.g. Jantz and Owsley, 1984; Lovejoy et al., 1990). In the Neolithic Ligurian sample, several individuals show osteoarticular lesions compatible with tuberculosis, including two children and one adolescent included in this study. Pollera 21, c. 5 years of age, shows multiple cystic lesions and bone erosions in the vertebral column, shoulder joint, and pelvis (Sparacello et al., 2017). At the age of 8.5 years, the most stunted individual (Arene Candide 6730.3+6623.1+6625.2) displays lesions suggestive of tuberculosis in the thoracic and sacral vertebral bodies (Figure 5; Sparacello, unpublished data, forthcoming). Arene Candide V (excavations Bernabò Brea – Cardini), c. 15 years old, suffered from Pott's spine, a collapse of the vertebral column which is considered pathognomonic for TB (Formicola et al., 1987). These individuals add to the growing evidence for this disease in both adults and children in Neolithic Liguria (e.g. Canci et al., 1996; Sparacello et al., 2018, and unpublished data, forthcoming), and mark a sharp contrast with the aforementioned site of Çatalhöyük, where no evidence of skeletal tuberculosis has been found despite the analysis a large skeletal series spanning over a millennium (Larsen et al., 2019).

[Figure 5 about here]

Indeed, evidence of tuberculosis is rare in the bioarchaeological record (Roberts and Buikstra, 2003); finding three individuals with lesions compatible with osteoarticular tuberculosis in our small sample of developing individuals suggests a high prevalence in the Ligurian Neolithic population. This hypothesis is supported by the fact that skeletal lesions manifest only in a small percentage of the affected individuals (estimates ranging from 1% to 3–5%; Turgut, 2001; Vigorita, 1999). In addition, the 5-10 years age class is the one with the lower risk of contracting the disease in modern epidemiological studies (Seddon and Shingadia, 2014). Active tuberculosis is a debilitating disease that impairs skeletal development (Mansukoski and Sparacello, 2018; Sparacello et al., 2016), but also the more common latent and sub-clinical states require a constant investment in immune defenses (Ulrichs et al., 2005;

Lin and Flynn, 2010), possibly diverting energy from growth (Ganmaa et al., 2012; see also McDade et al., 2008). We propose that significant infectious burden due to a high prevalence of tuberculosis in the Neolithic of Liguria may contribute to explain the pattern of growth faltering observed in this study. However, further research is necessary to investigate the paleoepidemiology of tuberculosis among Ligurian Neolithic people, by cross-referencing demographic data with new differential diagnoses.

Although the sample size is small, older children and early adolescents appear to have attained a proportion of the adult femoral length similar or higher than the reference Denver sample, with the exception of the adolescent with tuberculosis. This may indicate growth retardation followed by catch-up growth, resulting from an adaptation or acclimatization to environmental hardships (e.g. Beaton, 1989; Lewis, 2007:67; Stini, 1975). Indeed, during adolescence genetic influence on growth is expressed more strongly than during childhood, and environmental factors are relatively less important (Bogin, 1999). However, when considering absolute femoral length, Ligurian Neolithic adolescents are still well below the Denver sample. As discussed above, the relevance of the two methods to infer optimal growth patterns ultimately depends on whether differences in final stature between the Ligurian and Denver sample are due to genetic factors or failure to attain the full growth potential.

As in all bioarchaeological studies, there are numerous caveats that should be taken into account when interpreting the above results, which are not limited to the small sample size. Growth data from prehistoric populations are cross-sectional instead of longitudinal, and represent a cross-section of non-survivors. Cross-sectional data may not accurately describe the developmental patterns of a population, since growth events are not synchronized between individuals, resulting into a smoothing in the slope of growth curves (Humphrey, 2003). Furthermore, growth of non-survivors may be not representative of the normal development of the living population (which is part of the “osteological paradox”; Wood et al., 1992; Wright and Yoder, 2003), especially if they died of long-term, debilitating diseases (Goodman, 1993; Saunders and Hoppa, 1993; Sundick, 1978). Saunders and Hoppa (1993) noted that the linear growth of survivors is usually greater than that of non-survivors, but they concluded that the effect of this bias is relatively minor. However, given the widespread evidence of tuberculosis, the possibility that non survivors suffered from long-term developmental disturbances may be particularly relevant in the Ligurian sample. The individuals with tuberculous lesions appear stunted, but also most of the individuals between c 4.5 and 8.5 years of age. Ideally, it would be necessary to verify whether children dying of

different causes at the same age had similar dimensions, which is problematic given the small sample size of bioarchaeological samples, and especially given the uncertainties in the assessment of causes of death. Luckily, in our small sample, two individuals, Arene Candide VIII (excavations Bernabò Brea – Cardini), age midpoint 4.5 years, and Arene Candide 3 (excavations Tiné), age midpoint 8.5 years, show clear signs of perimortem trauma (Figure 6; Sparacello, unpublished data). Paradoxically, their violent and presumably sudden death makes them more likely to be representative of the population of survivors, although we cannot exclude that they suffered of a long-term disease which did not leave obvious traces in the skeleton. Nevertheless, their proportion of adult femoral length attained falls, like for most individuals between 4.5 and 8.5, between 1 and 2 standard deviation below the mean of the Denver sample. Although the small sample makes any inference tentative at best, this would suggest that growth disturbances during childhood were widespread in the Neolithic of Liguria.

[Figure 6 about here]

5. Conclusions

This study is part of a renewed, multidisciplinary attempt to characterize population health and well-being, subsistence patterns, and biocultural adaptive strategies in a chronologically and spatially well-defined window of European Neolithic variability (e.g. Orellana-Gonzales et al., in review; Goude et al., in review; Sparacello et al., 2017, 2018, 2019, in review), adding to the current debate on the competitive advantages and disadvantages of a Neolithic lifestyle.

When considering absolute femoral length, the Ligurian skeletal series is significantly smaller than the modern Denver sample throughout development. However, when considering the proportion of adult size attained at a given age, results suggests optimal development among Ligurian Neolithic people in early infancy, until the age of c. 2.5-3 years. Between 4.5 and 8.5 years, growth faltering is apparent, while later children and adolescents may show catch-up growth.

Although the sample size is small, the onset of growth faltering pattern corresponds with the estimated timing of weaning in the same Neolithic sample, reconstructed via isotopic analysis (Goude et al., in review), and with developmental disturbances observed in enamel mineralization (Orellana-Gonzales, in review). We propose that the growth pattern observed may relate more to disease load than nutritional factors. Early in life, optimal growth may be

due to infant feeding practices attempting to favor both growth, via the early introduction of animal protein (Goode et al., in review), and immune protection, by delaying the termination of breast feeding into the third year of life. Evidence of growth faltering between 4.5-8.5 years of age may reflect, among other environmental hardships, the stunting effect of debilitating diseases such as tuberculosis (which is manifest in three individuals) as well as the significant metabolic investment required into the immune system at the expense of growth in areas with significant infectious load.

In addition to small sample size, several methodological and theoretical caveats suggest caution when interpreting these results, and further research is necessary to test the above hypotheses. New differential diagnoses on individuals with suspect tuberculous osteoarticular lesions, coupled with demographic data, will contribute to the assessment of the paleoepidemiology of this disease among Ligurian Neolithic people. Demographic studies on the complete skeletal series of dated Neolithic Ligurian children will evaluate pattern of child mortality in order to verify whether the results observed here may be influenced by differential survivorship and frailty (Goodman, 1993; Saunders and Hoppa, 1993; Wood et al., 1992). The integration of developmental and demographic data will also inform the current debate on the Neolithic Demographic transition and its possible determinants (Bocquet-Appel 2002, 2009, 2011a, 2011b; Page et al., 2016).

The possibility of adaptation/acclimatization to slow growth among Ligurian Neolithic people, followed by catch up growth with adolescence, should be explored with a larger sample of subadults, which can be attained only by expanding the regional focus of the research, or by conducting further excavations in the Finalese area. Additionally, the use of other long bones from this same skeletal series may provide further confirmation of the results found here (e.g. Goode et al., 1993).

Other limitations of this study are the use of the midpoint of the age estimate based on available information about tooth development (Supplementary Information S1). A refined age estimation will be possible with the advancement of non-invasive analyses of tooth microstructure (e.g. Smith et al., 2015). Furthermore, error may be introduced by the lack of sex determination, given the well-known differences in developmental trajectories between sexes (e.g. Schaefer et al., 2009), which may be explored in the future using the amelogenin analysis of the enamel (Stewart et al., 2017).

Acknowledgements

The authors thank the Soprintendenza Archeologia, Belle Arti e Paesaggio per la città metropolitana di Genova e le provincie di Imperia, La Spezia e Savona, for granting access to the skeletal collections, especially the Superintendent Vincenzo Tiné and the Officers Elisabetta Starnini, Marta Conventi, Nico Radi, and Stefano Costa. Thanks to the Director of the Museo di Archeologia Ligure, Patrizia Garibaldi, for granting access to the collections owned by the Comune di Genova. We are grateful to the directors and curators of the museums where the skeletal collections are preserved, for continuous assistance during the data collection: Monica Zavattaro (Museo di Storia Naturale – Sezione di Antropologia e Etnologia, Università degli Studi di Firenze), Patrizia Garibaldi, Guido Rossi, Irene Molinari (Museo di Archeologia Ligure, Genova), Daniele Arobba, and Andrea De Pascale (Museo Archeologico del Finale, Finale Ligure). Thanks to Chiara Panelli, Stefano Rossi, Roberto Maggi, Paolo Biagi, Giovanni Murialdo, Elisa Bianchi, Simona Mordeglia, Walter Siciliano, Gwenaëlle Goude, Kate McGrath, Sacha Kacki, Eric Pubert, Alain Queffelec, Giovanna Stefania, Luca Bachechi, Chiara Bullo, and Brunetto Chiarelli for assistance during data collection and for their scientific input. A special thank goes to Mario.

We are grateful to the editor and the two reviewers who significantly improved this manuscript with their comments and suggestions.

The project DEN.P.H.: Dental anthropology at the Pleistocene-Holocene transition – insights on lifestyle and funerary behaviour from Neolithic Liguria (Italy) (ID) is funded by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 752626. The project BUR.P.P.H: Burial practices at the Pleistocene - Holocene transition: the changing role of pathology, violence, and “exceptional events” (VSS) has received financial support from the French State in the framework of the “Investments for the future” Program, IdEx Bordeaux, reference ANR-10-IDEX-03-02.

Literature Cited

Albanese, J., 2002. The use of skeletal data for the study of secular change: methodological implications of combining data from different sources. *Am. J. Phys. Anthropol. Suppl.* 34, 36.

Albanese, J., 2009. A Critical Review of the Methodology for the Study of Secular Change Using Skeletal Data. Ontario Archaeology No. 85-88/London Chapter OAS Occasional Publication No. 9, 139-155.

AlQahtani, S.J., Hector, M.P., Liversidge, H.M., 2010. Brief communication: The London atlas of human tooth and eruption. *Am. J. Phys. Anthropol.* 142, 481-490. <https://doi.org/10.1002/ajpa.21258>.

Armelagos, G.J., Barnes, K.C., Lin, J., 1996. Disease in human evolution: the re-emergence of infectious disease in the third epidemiological transition. *AnthroNotes* 18, 1-7.

Armelagos, G.J., Brown, P.J., Turner, B., 2005. Evolutionary, historical and political economic perspectives on health and disease. *Soc. Sci. Med.* 61, 755-765. <https://doi.org/10.1016/j.socscimed.2004.08.066>

Armelagos, G.J., Goodman, A.H., Harper, K.N., & Blakey, M.L., 2009. Enamel hypoplasia and early mortality: bioarcheological support for the Barker hypothesis. *Evol. Anthropol.* 18, 261-271. <https://doi.org/10.1002/evan.20239>.

Armelagos, G.J., Goodman, A.H., Jacobs, K.H., 1991. The origins of agriculture: population growth during a period of declining health. *Popul. Environ.* 13, 9-22. <https://doi.org/10.1007/BF01256568>

Arobba, D., Panelli, C., Caramiello, R., Gabriele, M., Maggi, R. 2017. Cereal remains, plant impressions and 14C direct dating from the Neolithic pottery of Arene Candide Cave (Finale Ligure, NW Italy). *J. Archaeol. Sci. Rep.* 12, 395-404. <https://doi.org/10.1016/j.jasrep.2017.02.015>

Beaton, G.H. 1989. Small but healthy? Are we asking the right question? *Eur. J. Clin. Nutr.* 43, 863-875.

Bello, S.M., Thomann, A., Signoli, M., Dutour, O., Andrews, P. 2006. Age and sex bias in the reconstruction of past population structures. *Am. J. Phys. Anthropol.* 129, 24-38. <https://doi.org/10.1002/ajpa.20243>

Binder, D., Sénépart, I. 2010. La séquence de l'Impresso-Cardial de l'abri Pendimoun et l'évolution des assemblages céramiques en Provence. *Mémoire LI de la Société Préhistorique française*, 51, 149-167.

Binder, D., Battentier, J, Delhon, C., Sénépart, I. 2017. In pursuit of a missing transition: the Mesolithic and Neolithic radiocarbon chronology at La Font-aux-Pigeons rockshelter. *Antiquity*, 91, 605-620. <https://doi.org/10.15184/aqy.2017.65>

Bocquet-Appel, J.-P. 2002. Paleoanthropological traces of a Neolithic demographic transition. *Curr. Anthropol.* 43, 637-650. <https://doi.org/10.1086/342429>

Bocquet-Appel, J.-P. 2009. The demographic impact of the agricultural system in human history. *Curr. Anthropol.* 50, 657-660. <https://doi.org/10.1086/605552>

Bocquet-Appel, J-P. 2011a. When the world's population took off: the springboard of the Neolithic Demographic Transition. *Science* 333, 560-561. <https://doi.org/10.1126/science.1208880>

Bocquet-Appel, J-P. 2011b. The agricultural demographic transition during and after the agriculture inventions. *Curr. Anthropol.* 52, S497-S510. <https://www.jstor.org/stable/10.1086/659243>

Bogin, B. 1988. *Patterns of human growth*. Cambridge: Cambridge University Press.

Bogin, B. 1999. Evolutionary perspective on human growth. *Annu. Rev. Anthropol.* 28, 109-153.

Canci, A., Minozzi, S., Borgognini Tarli, S. 1996. New evidence of tuberculous spondylitis from Neolithic Liguria (Italy). *Int. J. Osteoarchaeol.* 6, 497-501. [https://doi.org/10.1002/\(SICI\)1099-1212\(199612\)6:5<497::AID-OA291>3.0.CO;2-O](https://doi.org/10.1002/(SICI)1099-1212(199612)6:5<497::AID-OA291>3.0.CO;2-O)

Cienkosz-Stepanczak, B., Lisowak-Gaczorek, A., Haduch, E., Ellam, R., Cook, G., Kruk, J., Milisauskas, S., Koziel, S., Szostek, K. 2017. Nitrogen and strontium isotopes as tools for reconstruction of breastfeeding practices and human behavior – a Neolithic collective grave in Bronocice (Poland). *Coll. Anthropol.* 41, 191-199.

Cohen, M.N., Armelagos, G.J. 1984. *Paleopathology at the origins of agriculture*. Orlando, FL: Academic Press.

Cohen, M.N., Crane-Kramer, G.M.M. (Eds.). 2007. *Ancient health: skeletal indicators of agricultural and economic intensification*. Gainesville, FL: University Press.

Cook, D.C. 1979. Health and differential survival in prehistoric populations: prenatal dental defects. *Am. J. Phys. Anthropol.* 51:649-664.

Cook, D.C. 1984. Subsistence and health in the Lower Illinois Valley: osteological evidence, in: Cohen, M.N., Armelagos, G.J. (Eds.), *Paleopathology at the Origins of Agriculture*. New York: Academic Press, pp. 235-269.

Crepaldi, F., 2001. Le Chasséen en Ligurie. *Bulletin de La Société Préhistorique Française* 98(3), 485–494.

Delfino, E. 1981. *Liguria preistorica: sepolture dal Paleolitico Superiore all'Età del Ferro in Liguria e nell'area ligure*. Savona, Italy: Sabatelli Editore.

Del Lucchese, A. 1997. The Neolithic burials from Arene Candide cave – the Bernabò Brea-Cardini excavations, in: Maggi, R. (Ed.), *Arene Candide: a functional and environmental assessment of the Holocene sequence (excavations Bernabò Brea-Cardini 1940-50)*. *Mem. Ist. Ital. Paleontol. Um.* 5, pp. 605-611.

Del Lucchese, A., Starnini, E. 2015. Aggiornamenti sulla fase antica della Cultura dei Vasi a Bocca Quadrata in Liguria da una revisione dei materiali ceramici in corso, in: Conventi, M., Del Lucchese, A., Gardini, A. (Eds.), *Archeologia in Liguria, n.s., Vol. V, 2012-2013*. Genova, Italy: Sagep Editrice, pp. 23-37

De Pascale, A. 2007. Spunti e riflessioni per una storia delle prime ricerche paleontologiche nel Finalese. *Rivista di Scienze Preistoriche*, LVII, 379-398.

De Pascale, A. 2008. Le prime esplorazioni nelle caverne ossifere del Finalese: tracce, ipotesi e scoperte ad opera di Issel, Perrando, Morelli, Rovereto, Rossi, Amerano, in: De Pascale, A., Del Lucchese, A., Raggio, O. (Eds.), *La nascita della Paleontologia in Liguria: personaggi, scoperte e collezioni tra XIX e XX secolo*. Bordighera, Italy: Istituto Internazionale di studi Liguri, Atti del Convegno (Finale Ligure Borgo, 22-23 settembre 2006), pp. 223-248.

Demirjian, A. 1986. Dentition, in: Falkner, F., Tanner, J.M. (Eds.), *Human growth - a comprehensive treatise*. New York, Plenum Press, pp. 198-269

Eshed, V., Gopher, A., Pinhasi, R., Hershkovitz, I. 2010. Paleopathology and the origin of agriculture in the Levant. *Am. J. Phys. Anthropol.* 143, 121-133.
<https://doi.org/10.1002/ajpa.21301>

Eveleth, P.B., Tanner, J.M. 1990. *Worldwide variation in human growth*, 2nd edition. Cambridge: Cambridge University Press.

Feldesman, M.R. 1992. Femur/stature ratio and estimates of stature in children. *Am. J. of Phys. Anthropol.* 87, 447-412

Fernández-Crespo, T., Czermak, A., Lee-Thorp, J.A., Schulting, R.J. 2018. Infant and childhood diet at the passage tomb of Alto de la Huesera (north-central Iberia) from bone

collagen and sequential dentine isotope composition. *Int. J. Osteoarchaeol.* 28, 542-551.
<https://doi.org/10.1002/oa.2659>

Fildes VA. 1986. *Breasts, bottles and babies: a history of infant feeding*. Edinburgh: Edinburgh University Press.

Fildes, V.A. 1995. The culture and biology of breastfeeding: an historical review of Western Europe, in: Dettwyler, K.A., Stuart-Macadams, P. (Eds.), *Breastfeeding: biocultural perspectives*. New York: Aldine de Gruyter, pp. 101-126.

Formicola, V. 1987. Neolithic transition and dental changes: the case of an Italian site. *J. Hum. Evol.* 16, 231-239. [https://doi.org/10.1016/0047-2484\(87\)90078-9](https://doi.org/10.1016/0047-2484(87)90078-9)

Formicola, V., Milanesi, Q., Scarsini, C. 1987. Evidence of spinal tuberculosis at the beginning of the fourth millenium BC from Arene Candide cave (Liguria, Italy). *Am. J. Phys. Anthropol.* 72, 1-6. <https://doi.org/10.1002/ajpa.1330720102>

Ganmaa, D., Giovannucci, E., Bloom, B.R., Fawzi, W., Burr, W., Batbaatar, D., Sumberzul, N., Holick, M.F., Willett, W.C. 2012. Vitamin D, tuberculin skin test conversion, and latent tuberculosis in Mongolian school-age children: a randomized, double-blind, placebo-controlled feasibility trial. *Am. J. Clin. Nutr.* 96, 391-396.

Gleń-Haduch, E., Szostez, K., Głąb, H. 1997. Cribra orbitalia and trace element content in human teeth from Neolithic and Early Bronze Age graves in southern Poland. *Am. J. Phys. Anthropol.* 103, 201-207. [https://doi.org/10.1002/\(SICI\)1096-8644\(199706\)103:2<201::AIDAJPA5>3.0.CO;2-W](https://doi.org/10.1002/(SICI)1096-8644(199706)103:2<201::AIDAJPA5>3.0.CO;2-W)

Goode, H., Waldron, T., Rogers, J. 1993. Bone growth in juveniles: a methodological note. *Int. J. Osteoarchaeol.* 3, 321-323. <https://doi.org/10.1002/oa.1390030411>

Goodman, A.H. 1993. On the interpretation of health from skeletal remains. *Curr. Anthropol.* 34, 281-288. <https://doi.org/10.1086/204170>

Goodman, A.H. 1998. Skeletal growth and time of agricultural intensification, in: Ulijaszek, S.J., Johnson, F.E., Preece, M.A. (Eds.). *The Cambridge encyclopaedia of human growth and development*. Cambridge, Cambridge University Press, pp. 387-389.

Goodman, A.H., Lallo, J., Armelagos, G.J., Rose, J.C. 1984. Health changes at Dickson Moundsm, Illinois (A.D. 950-1300), in: Cohen, M.N., Armelagos, G.J. (Eds.), *Paleopathology at the Origins of Agriculture*. New York: Academic Press, pp. 271-301.

Goude, G., Binder, D., Del Lucchese, A. 2014. Alimentation et modes de vie néolithiques en Ligurie, in: Bernabo Brea, M., Maggi, R., Manfredini, A., (Eds.). *Il Pieno Neolitico in Italia (8-10 June, Finale Ligure 2009)*. *Riv. Stud. Lig.* 77, 371-387.

Goude, G., Dori, I., Sparacello, V.S., Starnini, E., Varalli, A. In Review. Multi-proxy dentine microsections analyses reveal diachronic changes in life history adaptations, mobility, and tuberculosis-induced wasting in prehistoric Liguria (Finale Ligure, Italy, northwestern Mediterranean). *Int. J. Paleopath.*

Harrington, L., Pfeiffer, S. 2008. Juvenile mortality in southern African archaeological contexts. *South African Archaeological Bulletin* 63, 95-101.

<https://doi.org/10.2307/20475004>

Hillson, S.W., Larsen, C.S., Boz, B., Pilloud, M.A., Sadvari, J.W., Agarwal, S C., Glencross, B., Beauchesne, P., Pearson, J., Ruff, C.B., Garofalo, E.M., Hager, L., Scott, D.H. 2013. The human remains I: Interpreting community structure, health, and diet in Neolithic Çatalhöyük., in: Hodder, I. (Ed.), *Humans and landscapes of Çatalhöyük: Reports from the 2000–2008 seasons*. Los Angeles: Cotsen Institute of Archaeology Press, pp. 339–396.

Hoppa, R.D., Fitzgerald, C.M. 1999. From head to toe: integrating studies from bone and teeth in biological anthropology, in: Hoppa, R.D., Fitzgerald, C.M. (Eds.), *Human growth in the past: studies from bones and teeth*. Cambridge, Cambridge University Press, pp. 1-31.

Howcroft, R. 2013. Weaned upon a time. *Studies of the infant diet in prehistory*. Ph.D. Thesis. Stockholm University, Sweden.

Howcroft, R., Eriksson, G., Lidén, K. 2014. Infant feeding practices at the Pitted Ware Culture site of Ajvide, Gotland. *J. Anthropol. Archaeol.* 34, 42-53.

<https://doi.org/10.1016/j.jaa.2014.01.001>

Humphrey, L.T. 2000. Growth studies of past populations: An overview and an example, in: Cox, M., Mays, S. (Eds.), *Human Osteology in Archaeology and Forensic Science*. Cambridge: Cambridge University Press, pp. 23-38.

Humphrey, L.T. 2003. Linear growth variation in the archaeological record, in: Thompson, J.L., Krovitz, G.E., Nelson, A.J. (Eds.), *Patterns of Growth in the Genus Homo*. Cambridge: Cambridge University Press, pp. 144-169.

Humphrey, L.T., Bello, S., Rousham, E. 2012. Sex differences in infant mortality in Spitalfields, London, 1750–1839. *J. Biosoc. Sci.* 44, 95-119.

Issel, A. 1908. *Liguria preistorica*. Genova: Società Ligure di Storia Patria.

Ives, R., Humphrey, L. 2017. Patterns of long bone growth in a mid-19th century documented sample of the urban poor from Bethnal Green, London, UK. *Am. J. Phys. Anthropol.* 163, 173-186. DOI: 10.1002/ajpa.23198

Jantz, R.L., Owsley, D.W. 1984. Long bone growth variation among Arikara skeletal populations. *Am. J. Phys. Anthropol.* 63, 13-20.

Johnston, F.E. 1962. Growth of the long bones of infants and young children at Indian Knoll. *Am. J. Phys. Anthropol.* 20, 249-254. <https://doi.org/10.1002/ajpa.1330200309>

Johnston, F.E., Wainer, B.E., Thissen, D., MacVean, R. 1976. Hereditary and environment determinants of growth in height in a longitudinal sample of children and youth of Guatemalan and European ancestry. *Am. J. Phys. Anthropol.* 44, 469-476. <https://doi.org/10.1002/ajpa.1330440310>

King, T., Humphrey, L.T., Hillson, S.W. 2005. Linear enamel hypoplasia as indicator of systemic physiological stress: evidence from two known age-at-death and sex populations from postmedieval London. *Am. J. Phys. Anthropol.* 128, 547-559. <https://doi.org/10.1002/ajpa.20232>

Larsen, C.S. 1995. Biological changes in human populations with agriculture. *Annu. Rev. Anthropol.* 24, 185-213. <https://doi.org/10.1146/annurev.an.24.100195.001153>

Larsen, C.S. 2015. *Bioarchaeology. Interpreting behavior from the human skeleton*, 2nd edition. Cambridge: Cambridge University Press.

Larsen, C.S., Crosby, A.W., Griffin, M.C., Hutchinson, D.L., Ruff, C.B., Russel, K.F., Schoeninger, M.J., Sering, L.E., Simpson, S.W., Takács, J.L., Teaford, M.F. 2002. A biohistory of health and behavior in the Georgia Bight: the agricultural transition and the impact of European contact, in: Steckel, R.H., Rose, J.C. (Eds.). *The backbone of history: health and nutrition in the western hemisphere*. Cambridge, Cambridge University Press, pp. 406-439.

Larsen, C.S., Knüsel, C.J., Haddow, S.D., Pilloud, M.A., Milella, M., Sadvari, J.W., Pearson, J, Ruff, C.B., Garofalo, E.M., Bocaege, E., Betz, B.J., Dori, I., Glencross, B. 2019. Bioarchaeology of Neolithic Çatalhöyük reveals fundamental transitions in health, mobility, and lifestyle in early farmers. *PNAS* 116(26), 12615-12623.

Le Bras-Goude, G., Binder, D., Formicola, V., Duday, H., Couture, C., Hublin, J.-J., Richards, M. 2006. Stratégies de subsistance et analyse culturelle de populations Néolithiques de Ligurie: approche par l'étude isotopique ($\delta^{13}\text{C}$ et $\delta^{15}\text{N}$) des restes osseux. *Bull. Mém. Soc. Anthropol. Paris* 18, 43-53.

Leary, S., Fall, C., Osmond, C., Lovel, H., Campbell, D., Eriksson, J., Forrester, T., Godfrey, K., Hill, J., Jie, Mi., Law, C., Newby, R., Robinson, S., Yajnik, C. 2006. Geographical variation in neonatal phenotype. *Acta Obstet. Gynecol. Scand.*, 85(9), 1080–1089.

Lewis, M.E. 2007. *The bioarchaeology of children*. Cambridge: Cambridge University Press.

- Lin, P.L., Flynn, J.L. 2010. Understanding latent tuberculosis: a moving target. *J. Immunol.* 185(1), 15-22.
- Lovejoy, C.O., Russell, K.F., Harrison, M.L. 1990. Long bone growth velocity in the Libben population. *Am. J. Hum. Biol.* 2, 533-542. <https://doi.org/10.1002/ajhb.1310020509>
- Macphail, R.I., Courty, M.A., Hater, J., Watez, J. 1997. The soil micromorphological evidence of domestic occupation and stabling activities, in: Maggi, R. (Ed.), *Arene Candide: a functional and environmental assessment of the Holocene sequence (excavations Bernabò Brea-Cardini 1940-50)*. *Mem. Ist. Ital. Paleontol. Um.* 5, pp. 31-52.
- Maggi, R., 1997. The radiocarbon chronology, in: Maggi, R., Starnini, E., Voytek, B.A., (Eds.), *Arene Candide: a functional and environmental assessment of the Holocene sequence (excavations Bernabò Brea-Cardini 1940-1950)*. *Mem. Ist. Ital. Paleontol. Um.* 5, pp. 31-52.
- Mansukoski, L., Sparacello, V.S. 2018. Smaller long bone cross-sectional size in people who died of tuberculosis: insights on frailty factors from a 19th and early 20th century Finnish population. *Int. J. Paleopathol.* 20, 38-44. <https://doi.org/10.1016/j.ijpp.2017.12.005>
- Marchi, D., Sparacello, V.S., Holt, B.M., Formicola, V. 2006. Biomechanical approach to the reconstruction of activity patterns in Neolithic Western Liguria, Italy. *Am. J. Phys. Anthropol.* 131, 447-455. <https://doi.org/10.1002/ajpa.20449>
- Marchi, D., Sparacello, V.S., Shaw, C.N. 2011. Mobility and lower limb robusticity of a pastoralist Neolithic population from North-Western Italy, in: Pinhasi, R., Stock, J. (Eds.), *Human Bioarchaeology of the Transition to Agriculture*. New York, NY: Wiley-Liss, pp. 317-346.
- Maresh, M.M. 1955. Linear growth of long bones of extremities from infancy through adolescence; continuing studies. *Am. J. Dis. Child.*, 89, 725-742.
- Maresh, M.M. 1943. Growth of major long bones in healthy children: a preliminary report on successive roentgenograms of the extremities from early infancy to twelve years of age. *Am. J. Dis. Child.*, 6, 227-257.
- Maresh, M.M. 1970. Measurements from roentgenograms, in: McCammon, R.W. (Ed.) *Human Growth and Development*. Springfield IL: C.C. Thomas, pp. 157-200.
- Martorell, R. 1985. Child growth retardation: a discussion of its causes and its relationship to health, in: Blaxter, K., Waterlow, J.C. (Eds.), *Nutritional adaptation in man*. London, John Libby, pp. 13-30.
- Mays, S. 2007. The human remains, in: Mays, S., Harding, C., Heighway, C. (Eds.), *Wharram: a study of settlement on the Yorkshire Wolds, XI the churchyard*. York University Archaeological Publications 13. York, University of York, pp. 77-192, 337-397.

Mays, S. 2010. The Effects of Infant Feeding Practices on Infant and Maternal Health in a Medieval Community. *Childhood in the Past*, 3:1, 63-78. DOI: 10.1179/cip.2010.3.1.63

Mays, S. 2018. The study of growth in skeletal populations, in: Crawford, S.C., Hadley D.M., Shepherd, G. (Eds.), *The Oxford Handbook of the Archaeology of Childhood*. Oxford: Oxford University Press, pp. 71-89.

McDade, T.W. 2003. Life history theory and the immune system: steps toward a human ecological immunology. *Yearb. Phys. Anthropol.* 46, 100-125.

McDade, T.W., Reyes-Garcia, V., Tanner, S., Huanca, T., Leonard, W.R. 2008. Maintenance versus growth: investigating the costs of immune activation among children in lowland Bolivia. *Am. J. Phys. Anthropol.* 136, 478-484. <https://doi.org/10.1002/ajpa.20831>

McDade, T.W., Worthman, C.M. 1998. The weanling's dilemma reconsidered: a biocultural analysis of breast-feeding ecology. *J. Dev. Behav. Pediatr.* 19, 286-299.

Nisbet, R. 2008. Environment and agriculture in the early Neolithic of the Arene Candide (Liguria), in: Fiorentino, G., Magri, D. (Eds.), *Charcoals from the Past: Cultural and Palaeoenvironmental Implications. Proceedings of the Third International Meeting of Anthracology, Cavallino- Lecce (Italy), June 28th- July 1st 2004*, BAR International Series, 1807, pp. 193-198.

Nitsch, E.K., Humphrey, L.T., Hedges, R.E.M. 2011. Using stable isotope analysis to examine the effect of economic change on breastfeeding practices in Spitalfields, London, UK. *Am. J. Phys. Anthropol.* 146, 619-628. <https://doi.org/10.1002/ajpa.21623>

Orellana González, E., Sparacello, V.S., Bocaage, E., Varalli, A., Moggi-Cecchi, J., Dori, I. n.d. Insights on patterns of developmental disturbances from analysis of linear enamel hypoplasia in a Neolithic sample from Liguria (northwestern Italy). *Int. J. Paleopathol.* (in review).

Page, A.E., Viguier, S., Dyble, M., Smith, D., Chaudhary, N., Salali, G.D., Thompson, J., Vinicius, L., Mace, R., Migliano, A.B. 2016. Reproductive trade-offs in extant hunter-gatherers suggest adaptive mechanism for the Neolithic expansion. *PNAS*, 113, 4694-4699. <https://doi.org/10.1073/pnas.152403111>

Panelli, C., Rossi, S. 2015. Alfred J. Wall “medico inglese dell’esercito delle Indie Orientali” e gli scavi nella Caverna delle Arene Candide e della Grotta Pollera (Finale Ligure). *Archeologia in Liguria*, 5, 300-302.

Panelli, C., Rossi, S. 2017. Desenterrando el pasado (reciente): las excavaciones de Alfred John Wall en las Cuevas de Arene Candide y Pollera (Finale Ligure, Savona – Italia),

in: Ayarzagüena Sanz, M., Mora, G., Salas Álvarez, J. (Eds.), 150 Años de Historia de la Arqueología: Teoría y método de una disciplina. Madrid, pp. 125-142.

Parenti, R., Messeri, P. 1962. I resti scheletrici umani del Neolitico Ligure. *Palaeontographia Italica*, 50, 5-165.

Pearson, J.A., Haddow, S.D., Hillson, S.W., Knüsel, C.J., Larsen, C.S., Sadvari, J.W. 2015. Stable carbon and nitrogen isotope analysis and dietary reconstruction through the life course at Neolithic Çatalhöyük, Turkey. *J. Soc. Archaeol.* 15, 210-232.
<https://doi.org/10.1177/1469605315582983>

Pearson, J.A., Hedges, R.E.M., Molleson, T.I., Özbek, M. 2010. Exploring the relationship between weaning and infant mortality: An isotope case study from Aşikli Höyük and Cayönü Tepesi. *Am. J. Phys. Anthropol.* 143, 448-457.
<https://doi.org/10.1002/ajpa.21335>

Pérez-Losada, J., Fort, J. 2010. Age-dependent mortality, fecundity and mobility effects on front speeds: theory and application to the Neolithic transition. *J. Stat. Mech.: Theory Exp.* P11006. <https://doi.org/10.1088/1742-5468/2010/11/P11006>

Pfeiffer, S., Harrington, L. 2010. Child growth among southern African foragers, in: Moffat, T., Prowse, T. (Eds.) *Human diet and nutrition in biocultural perspective. Studies of the Biosocial Society vol. 5*, Berghahn, Oxford, pp. 35–56.

Pinhasi, R., Stefanovic, S., Papathanasiou, A., Stock, J.T. 2011. Variability in long bone growth patterns and limb proportions within and amongst Mesolithic and Neolithic populations from southeast Europe, in: Pinhasi, R., Stock, J.T. (Eds.), *Human bioarchaeology of the Transition to Agriculture*. Chichester, Wiley-Blackwell, pp. 177-202.

Pinhasi, R., Timpson, A., Thomas, M., Šlaus, M. 2014. Bone growth, limb proportions and non-specific stress in archaeological populations from Croatia. *Ann. Hum. Biol.* 41(2), 127-137.

Powell, G.F., Brasel, J.A., Raiti, S., Blizzard, R.M. 1967. Emotional deprivation and growth retardation simulating idiopathic hypopituitarism: Clinical evaluation of the syndrome. *N. Eng. J. Med.* 276, 1271-1278. <https://doi.org/10.1056/NEJM196706082762301>

Richard, C. 1942. Scavi nell'Arma dell'Aquila a Finale Ligure: prima relazione. *Bullettino di Paleontologia Italiana, Nuova Serie, V-VI*, 43-100.

Richards, M.P., Pearson, J.A., Molleson, T.I., Russell, N., Martin, L. 2003. Stable Isotope evidence of diet at Neolithic Çatalhöyük. *J. Archaeol. Sci.* 30, 67-76.

Roberts, C.A., Buikstra, J.E. 2003. *The bioarchaeology of tuberculosis*. Gainesville: University Press of Florida.

Rosenstock, E., Ebert, J., Martin, R., Hicketier, A., Walter, P., Groß, M. 2019. Human stature in the Near East and Europe ca. 10,000-1000 BC: its spatiotemporal development in a Bayesian errors-in-variables model. *Archaeol. Anthropol. Sci.* doi.org/10.1007/s12520-019-00850-3

Rowley-Conwy, P. 1992. Arene Candide: a Small Part of a Larger Pastoral System?, in: Maggi, R., Nisbet, R., Barker, G. (Eds.) *Archeologia della pastorizia nell'Europa meridionale. Rivista di Studi Liguri*, LVII (1-4), 95-116.

Rowley-Conwy, P. 1997. The animal bones from Arene Candide (Holocene sequence): final report, in: Maggi, R. (Ed.), *Arene Candide: a functional and environmental assessment of the Holocene sequence (Bernabò Brea-Cardini 1940–50)*. *Mem. Ist. Ital. Paleontol. Um.* 5, pp. 153-279.

Rowley-Conwy, P. 1998. Improved separation of Neolithic metapodials of sheep (*Ovis*) and goat (*Capra*) from Arene Candide cave, Liguria, Italy. *J. Archaeol. Sci.* 25, 251-258. <https://doi.org/10.1006/jasc.1997.0204>

Rossi, S., Panelli, C., De Pascale, A., Maggi, R. 2014. “Di una caverna ossifera di Finale”: evidenze di archeologia ottocentesca nella Caverna delle Arene Candide, in: Guidi, A. (Ed.), *150 anni di Preistoria e Protostoria in Italia. Il contributo della Preistoria e della Protostoria alla formazione dello Stato unitario (Atti della XLVI Riunione Scientifica IIPP, Roma, 23-26 Novembre 2011)*. *Studi di Preistoria e Protostoria* 1, Firenze, pp. 237-244.

Ruff, C.B. 2007. Body size prediction from juvenile skeletal remains. *Am. J. Phys. Anthropol.* 133, 698-716. <https://doi.org/10.1002/ajpa.20568>

Ruff, C.B., Garofalo, E., Holmes, M.A. 2013. Interpreting skeletal growth in the past from a functional and physiological perspective. *Am. J. Phys. Anthropol.* 150, 29-37.

Ruff, C.B., Holt, B.M., Sladek, V., Berner, M., Murphy, W.A.Jr., zur Nedden, D., Seidler, H., Recheis, W. 2006. Body size, body proportions, and mobility in the Tyrolean "Iceman". *J. Hum. Evol.* 51, 91-101.

Ryan, A. S. 1976. Long bone growth in a prehistoric population from San Cristóbal, New Mexico. *Michigan Discussions in Anthropology*, 2, 55-75.

Saunders, S.R. 2008. Juvenile skeletons and growth-related studies, in: Katzemberg, M.A., Saunders, S.R. (Eds.), *Biological anthropology of the human skeleton*, second edition. New York: John Wiley & Sons, Inc., pp. 117-147.

Saunders, S.R., Hoppa, R.D. 1993. Growth deficit in survivors and non-survivors: Biological mortality bias in subadult skeletal samples. *Yearb. Phys. Anthropol.* 36, 127-151. <https://doi.org/10.1002/ajpa.1330360608>

Schaefer, M., Black, S., Scheuer, L. 2009. Juvenile osteology – a laboratory and field manual. New York, NY: Academic Press.

Scharlotta, I., Goude, G., Herrscher, E., Bazaliiskii, V.I., Weber, A.W. 2018. Shifting weaning practices in Early Neolithic Cis-Baikal, Siberia: New insights from stable isotope analysis of molar micro-samples. *Int. J. Osteoarchaeol.* 28(5), 579-598.

Seddon, J.A., Shingadia, D. 2014. Epidemiology and disease burden of tuberculosis in children: a global perspective. *Infect. Drug Resist.* 7, 153-165.
<https://doi.org/10.2147/IDR.S45090>

Smith, T.M., Tafforeau, P., Le Cabec, A., Bonnin, A., Houssaye, A., Pouech, J., Moggi-Cecchi, J., Manthi, F., Ward, C., Makaremi, M., Menter, C.G. 2015. Dental ontogeny in Pliocene and early Pleistocene hominins. *PLoS ONE* 10(2): e0118118.
<https://doi.org/10.1371/journal.pone.0118118>

Solomons, N.W., Mazariegos, M., Brown, K.H., Klasing, K. 1993. The underprivileged, developing country child: environmental contamination and growth failure revisited. *Nutr. Rev.* 51, 327-332. <https://doi.org/10.1111/j.1753-4887.1993.tb03758.x>

Sparacello, V.S., Marchi, D. 2008. Mobility and subsistence economy: a diachronic comparison between two groups settled in the same geographical area (Liguria, Italy). *Am. J. Phys. Anthropol.* 136, 485-495. <https://doi.org/10.1002/ajpa.20832>

Sparacello, V.S., Marchi, D., Shaw, C.N. 2014. The importance of considering fibular robusticity when inferring the mobility patterns of past populations, in: Carlson, K.J., Marchi, D. (Eds.), *Reconstructing Mobility: Environmental, Behavioral, and Morphological Determinants*. New York, NY: Springer, pp. 91-110. https://doi.org/10.1007/978-1-4899-7460-0_6

Sparacello VS, Roberts CA, Canci A, Moggi-Cecchi J, Marchi D. 2016. Insights on the paleoepidemiology of ancient tuberculosis from the structural analysis of postcranial remains from the Ligurian Neolithic (northwestern Italy). *Int J Paleopath* 15:50-64.

Sparacello, V.S., Roberts, C.A., Kerudin, A., Müller, R. 2017. A 6,500-year-old Middle Neolithic child from Pollera Cave (Liguria, Italy) with probable multifocal osteoarticular tuberculosis. *Int. J. Paleopath.* 17, 67-74. <https://doi.org/10.1016/j.ijpp.2017.01.004>

Sparacello, V.S., Panelli, C., Rossi, S., Dori, I., Varalli, A., Goude, G., Kacki, S., Partiot, C., Roberts, C.A., Moggi-Cecchi, J. 2018. Chapter 9: archaeoethanatology and palaeobiology of the burials and “scattered human remains” from Arma dell’Aquila (Finale Ligure, Savona), in Biagi, P., Starnini E. (Eds.), *Gli scavi all’Arma dell’Aquila (Finale Ligure, Savona): le Ricerche e i Materiali degli Scavi del Novecento*. Trieste, Italy: Società

per la Preistoria e Protostoria della Regione Friuli-Venezia Giulia, Quaderno 15, pp. 143-181.

Sparacello, V.S., Panelli, C., Rossi, S., Dori, I., Varalli, A., Goude, G., Starnini, E., Biagi, P. 2019. The re-discovery of Arma dell'Aquila (Finale Ligure, Italy): new insights on Neolithic funerary behavior from the sixth millennium BCE in the north-western Mediterranean. *Quat. Int.* 512, 67-81. doi.org/10.1016/j.quaint.2019.02.003.

Sparacello, V.S., Pearson, O.M., Coppa, A., Marchi, D. 2011. Changes in skeletal robusticity in an Iron Age agropastoral group: the Samnites from the Alfedena necropolis (Abruzzo, Central Italy). *Am. J. Phys. Anthropol.* 144, 119-130. <https://doi.org/10.1002/ajpa.21377>

Sparacello, V.S., Varalli, A., Rossi, S., Panelli, C., Goude, G., Palstra, S.W.L., Conventi, M., Del Lucchese, A., Arobba, D., De Pascale, A., Zavattaro, M., Garibaldi, P., Rossi, G., Molinari, I., Maggi, R., Moggi Cecchi, J., Starnini, E., Biagi, P., Dori, I. In Review. The funerary use of caves in Liguria (northwestern Italy) from the Neolithic to historic times: results from a large-scale AMS dating campaign on human skeletal series. *Quat. Int.*

Stephenson, C. B. 1999. Burden of infection on growth failure. *J. Nutr.* S129, 534-538.

Stephenson, L.S. 1987. Impact of helminth infections on human nutrition. London: Taylor & Francis.

Stewart, N.A., Gerlach, R.F., Gowland, R.L., Gron, K.J., Montgomery, J. 2017. Sex determination of human remains from peptides in tooth enamel. *PNAS*, 114, 13649-13654. <https://doi.org/10.1073/pnas.1714926115>

Stini, W.A. 1975. Adaptive strategies of human populations under nutritional stress, in: Watts, E.S., Johnston, F.E., Lasker, G.W. (Eds.), *Biosocial Interrelations in Population Adaptation*. Paris: Mouton, pp. 19-41.

Stinson, S. 2000. Growth variation: biological and cultural factors, in: Stinson, S., Bogin, B, Huss-Ashmore, R., O'Rourke, D. (Eds.), *Human biology: an evolutionary and biocultural perspective*. Chinchester: Wiley-Liss, pp. 423-462.

Sundick, R.I. 1978. Human skeletal growth and age determination. *Homo*, 29, 228-249.

Temple, D.H. 2010. Patterns of systematic stress during the agricultural transition in prehistoric Japan. *Am. J. Phys. Anthropol.* 142, 112-124. <https://doi.org/10.1002/ajpa.21208>

Tiné, V. 1999. Transizione tra Neolitico antico e Neolitico medio. Le ceramiche dello stile Pollera. *Strato* 13, in: Tinè, S. (Ed.), *Il Neolitico nella Caverna delle Arene Candide (scavi 1972-1977)*. Bordighera, Italy: Istituto Internazionale di studi Liguri, Collezione di Monografie Preistoriche ed Archeologiche, X, pp. 142-180.

Turgut, M. 2001. Spinal tuberculosis (Pott's disease): its clinical presentation, surgical management, and outcome. A survey study on 694 patients. *Neurosurg. Rev.* 24, 8-13.

Ulijaszek, S.J., Johnston, F.E., Preece, M.A. 1998. *The Cambridge encyclopedia of human growth and development*. Cambridge, UK: Cambridge University Press.

Ulrichs, T., Kosmiadi, G.A., Jörg, S., Pradl, L., Titukhina, M., Mishenko, V., Gushina, N., Kaufmann, S.H.E. Differential organization of the local immune response in patients with active cavitary tuberculosis or with nonprogressive tuberculosis. *J. Infect. Dis.* 192, 89-97.

Vigorita, V.J. 2008. *Orthopaedic pathology*. Philadelphia: Lippincott Williams & Wilkins.

Wood, J.W., Milner, G.R., Harpending, H.C., Weiss, K.M. 1992. The osteological paradox: problems of inferring prehistoric health from skeletal samples. *Curr. Anthropol.* 3, 343-370. <https://doi.org/10.1086/204084>

Wright, L.E., Yoder, C.J. 2003. Recent progress in bioarchaeology: approaches to the osteological paradox. *J. Archaeol. Res.* 11, 43-70. <https://doi.org/10.1023/A:1021200925063>

Figure 1 – Geographical location of the sites included in this study. Top: the red square indicates the Finalese area within the Liguria region (highlighted in yellow) in northwestern Italy. Bottom: the cave sites analyzed in this study in the municipality of Finale Ligure: 1) Arene Candide Cave; 2) Grotta Pollera; 3) Arma dell’Aquila; 4) Grotta dei Pipistrelli; 5) Arma Strapatente; 6) Grotta marina di Bergeggi. Modified from ArcGIS and Google Maps.

Figure 2 - Absolute maximum length of the femur by age in the Ligurian Neolithic sample, when compared with: Left) the data of the Denver Growth Study, corrected for radiographic magnification using the formulae proposed by Ruff (Ruff, 2007:702), and Right) the sample from the Denver Growth Study selected and corrected by Ruff (2007). The solid lines indicate the mean, the two dotted lines indicate ± 1 and ± 2 standard deviations. In Ruff (2007) the lengths without epiphyses are reported until the age of 12, and with epiphyses from the age of 11. Solid circles: ICC; empty squares: SMP; stars: Chassean.



Figure 3 – Length of the femur expressed as percentage of mean adult length attained by age in the Ligurian Neolithic sample, when compared with Left) the data of the Denver Growth Study, and Right) the sample from the Denver Growth Study selected and corrected by Ruff (2007). The solid lines indicate the mean, the two dotted lines indicate ± 1 and ± 2 standard deviations. The lengths without epiphyses are reported until the age of 12, and from the age of 10 (11 in Ruff, 2007) the lengths with epiphyses are reported. Solid circles: ICC; empty squares: SMP; stars: Chassean.



Figure 4 – The residuals of the length of the femur expressed as percentage of mean adult length attained by age in the Ligurian Neolithic sample, with respect to Left) the data of the Denver Growth Study, and Right) the sample from the Denver Growth Study selected and corrected by Ruff (2007). The solid lines indicate the mean, the two dotted lines indicate ± 1 and ± 2 standard deviations. Solid circles: ICC; empty squares: SMP; stars: Chassean.



Figure 5 – A vertebral body of an individual from Arene Candide (AC 6623.1, catalogue number of the Museo di Storia Naturale – Sezione di Antropologia e Etnologia, Università degli Studi di Firenze; possibly burial n°6 from the excavations Morelli 1884-87) showing an erosion of a thoracic vertebral body compatible with osteoarticular tuberculosis (cf. Sparacello et al., 2017).

Figure 6 – Signs of perimortem trauma in two Neolithic children from Liguria included in this analysis. Left: mandibular fracture in Arene Candide VIII excavations Bernabò Brea 1940-50. Right: fracture of the neural arch of a thoracic vertebra in Arene Candide 3 excavations Tiné 1973-76.











a.



b.



Individual Id	Dental Age	Age Midpoint	Femur R (mm)	Femur L (mm)	AMS date cal. BCE (95.4%) ¹	Chrono-cultural attribution ¹
Arma dell'Aquila 7 Richard	perinatal	0.0	70.4	70.2	5657-5533	ICC
Arma dell'Aquila 8 Richard	perinatal	0.0		75.5	5646-5527	ICC
Pollera 6663.1	perinatal	0.0	74	73	4701-4548	SMP
Pollera 6664.1	perinatal	0.0		75	4536-4373	SMP
Pollera 6665.1+6670.1	perinatal	0.0		73	3946-3775	CH
Arene Candide 6633.1	4.5-7.5 months	0.5	96.3		4726-4557	SMP
Arene Candide 6629.1	7.5-10.5 months	0.8	98.2	98.3	4767-4586	SMP
Pollera 6675.2+6676.1	7.5-12 months	0.8	126	129	4719-4557	SMP
Arene Candide 6630.1	10.5-18 months	1.2	124.5	124.7	4682-4502	SMP
Arene Candide 6628.1+6625.4	1-1.5	1.3	123	122	4768-4592	SMP
Pollera 6666.2	1-1.5	1.3		119	4712-4556	SMP
Pollera 6669.1+6671.1	1-1.5	1.3	134	133	4712-4587	SMP
Pollera 6680.1	1-2	1.5		128	4836-4717	SMP
Pollera 6670.2	1.5-2.5	1.8	147	148	4682-4502	SMP
Arene Candide 6631.1	1.5-2.5	2.0	140.4	139.9	4691-4545	SMP
Arma dell'Aquila 6 Richard	1.5-2.5	2.0		153	5657-5538	ICC
Arene Candide 6632.1+6623.2	2-3	2.5		161.6	4726-4557	SMP
Arene Candide VIII BB	3.5-5.5	4.5	184	190	4800-4619	SMP
Bergeggi S2 01178PE, 6894-5FI	4.5-5.5	5.0		193	5047-4857	SMP
Pollera 21PE	4-6	5.0	190		4779-4587	SMP
Pollera 6678.1	5-6	5.5	206	209	4794-4686	SMP
Arma dell'Aquila Zambelli JUV B	6-7	6.5	228		4727-4546	SMP
Pipistrelli 1_dep. 23.I.2_JUV	6-7	6.5		222.5	4703-4545	SMP
Arene Candide 6627.1	6-8	7.0		214	4690-4544	SMP
Pollera 20PE	6-8	7.0		230	4715-4556	SMP
Pollera 6682.1	7-9	8.0		231	4707-4555	SMP
Strapatente I	7.5-8.5	8.0		255	4531-4369	SMP
Arene Candide 3 Tinè	7.5-9.5	8.5	246	249	4782-4502	SMP
Arene Candide 6730.3+6623.1+6625.2	7-10	8.5		243	4779-4608	SMP
Pollera 1PE Issel-Morelli	10-12	11.0		373 ²	4783-4620	SMP
Arene Candide V BB	14-16	15.0	348	351	4720-4557	SMP
Arene Candide 6621.1	13-19	16.0		400.5	4726-4557	SMP
Arene Candide 1 Tinè	15-18	16.5		404	4704-4374	SMP

Table 1 – Subadult individuals from the Neolithic of Liguria included in this study. Estimate of age at death based on dental development is in years unless otherwise indicated. ICC: Impresso-Cardial Complex; SMP: Square Mouthed Pottery; CH: Chassean. ¹ Details on the direct AMS dates and chrono-cultural attributions are provided in Sparacello et al., in review. ² Epiphyses were plastered to the diaphysis, the length may be slightly overestimated.

Legend:

- Table legend: upper: upper dentition, lower: lower dentition; n/e: dentition non-erupted, e: dentition partially or fully erupted; R: right, L: left.
- Teeth legend: I, i: incisor; C, c: canine; P: premolar; M, m: molar; U: upper; L: lower; R: right; L: left; d: deciduous; capital letters: indicate the maxilla or mandible tooth and the permanent tooth (e.g. URI1: upper right first incisor), lower case letters: indicate the deciduous tooth (e.g. URdi1: upper right first deciduous incisors). See AlQahtani et al. (2010) for the description used to identify tooth developmental stages of single and multirooted teeth.

Arma dell'Aquila 7 Richard: deciduous dentition not fully mineralized.

		<i>upper</i>																							
R	n/e									i2		i1					m1	m2					n/e	L	
	e																							e	
		<i>lower</i>																							
	e																							e	
	n/e									m1		i2	i1		i1	i2		m1					n/e		

- ULdi1: Cr $\frac{3}{4}$
- URdi2: Cr $\frac{3}{4}$
- LRdi1-LLdi1: Cr $\frac{3}{4}$
- LRdi2-LLdi2: Cr $\frac{1}{2}$ - Cr $\frac{3}{4}$
- LRdm1-LLdm1: Coc
 - Age at death: perinatal/new born

Arma dell'Aquila 8 Richard: no teeth present. The estimation of the age at death has been made with the analysis of cranial and post-cranial bones. The skeleton has been compared with the other newborns of the Neolithic Ligurian sample, particularly Arma dell'Aquila 7 Richard.

- Age at death: perinatal/new born

Pollera 6663.1: deciduous dentition not fully mineralized.

		<i>upper</i>																							
R	n/e																							n/e	L
	e																							e	
		<i>lower</i>																							
	e																							e	
	n/e																							n/e	

- ULdi1: Crc
- ULdi2: Cr $\frac{3}{4}$
- URdm1: Coc
- URdm2-ULdm2: Cco - Ci
- LRdi1-LLdi1: Crc
- LLdi2: Cr $\frac{3}{4}$
- LRdm1-LLdm1: Coc
- LRdm2-LLdm1: Cco - Ci
 - Age at death: perinatal/new born

Arene Candide 6629.1: deciduous dentition not fully mineralized.

		<i>upper</i>																	
R	n/e				m2		c	i2	i1	i1	i2	c	m1					n/e	L
	e																	e	
		<i>lower</i>																	
	e																	e	
	n/e				m2	m1	c	i2	i1	i1	i2	c	m1	m2				n/e	

- ULdi1: R ¼
- URdi2: Ri – R ¼
- ULdi2: Ri – R ¼
- LLdm1: CrC
- ULdm1: Ri
- URdm2: Cr ¾

➤ Age at death: c. 7.5-10.5 months

Pollera 6675.2+6676.1: deciduous dentition not fully mineralized. Partially erupted some deciduous teeth.

		<i>upper</i>																	
R	n/e				m2		c					c		m2				n/e	L
	e					m1			i1		i2		m1					e	
		<i>lower</i>																	
	e					m1				i1			m1					e	
	n/e			M1	m2		c	i2			i2	c		m2	M1			n/e	

- URdm2: R ¼
- LLdi2: R ½

➤ Age at death: c. 7.5 months-1 year (< 1 yr)

Arene Candide 6630.1+CR NNPE Infans: deciduous dentition not fully mineralized. Partially erupted some deciduous teeth.

		<i>upper</i>																	
R	n/e						c					c						n/e	L
	e							i2	i1	i1	i2							e	
		<i>lower</i>																	
	e				m2	m1		i2	i1	i1	i2		m1	m2				e	
	n/e			M1			c					c			M1			n/e	

- ULdi1: R ½ - R ¾
- ULdm1: R ½

➤ Age at death: c. 10.5-18 months

Arene Candide 6628.1+6625.4: two isolated teeth, one deciduous and one permanent, not fully mineralized.

- URdi2: R ½ - R ¾
- URM1: Cr ½

➤ Age at death: c. 1-1.5 years

Pollera 6666.2: deciduous dentition not fully mineralized and erupted. It is possible to observe the presence of the lower first mandibular molar. Maxillary teeth not present.

		<i>upper</i>																	
R	n/e																	n/e	L
	e																	e	
		<i>lower</i>																	
	e							i1		i2		m1						e	
	n/e						c					m2	M1				n/e		

- LRdi1: R ½
- LLdi2: R ½
- LRdc: Ri
- LLdm1: R ½
- Age at death: c. 1-1.5 years

Pollera 6669.1+6671.1: deciduous dentition not fully mineralized and erupted. It is possible to observe the presence of the lower first permanent molar.

		<i>upper</i>																	
R	n/e				m2		c					c		m2				n/e	L
	e				m1		i2	i1	i1	i2		m1					e		
		<i>lower</i>																	
	e					m1		i2	i1	i1	i2		m1					e	
	n/e				M1	m2		c				c		m2	M1			n/e	

- URdm2-ULdm2: Ri - R ¼
- LRdc: R ¼
- Age at death: c. 1-1.5 years

Pollera 6680.1: deciduous dentition not fully erupted and mineralized. It is possible to observe the presence of the lower first and second permanent incisors. Maxillary teeth not present.

		<i>upper</i>																	
R	n/e																	n/e	L
	e																	e	
		<i>lower</i>																	
	e					m1							m1					e	
	n/e				m2		c	I2	I1	I1		c		m2	M1			n/e	

- LRdm2: Ri
- LLM1: Cr ½
- Age at death: c. 1-2 years

Pollera 6670.2: deciduous dentition erupted but not fully mineralized. It is possible to observe the degree of mineralization of upper first permanent molar.

		<i>upper</i>																	
R	n/e																M1	n/e	L
	e				m2	m1	c	i2		i1		c	m1	m2				e	
		<i>lower</i>																	
	e				m2	m1												e	
	n/e				M1													n/e	

- URdc: R ¾

URdm2: R ½

ULM1: Crc

- Age at death: c. 1.5-2.5 years

Arene Candide 6631.1: deciduous teeth erupted, permanent teeth not fully mineralized.

		<i>upper</i>																		
R	n/e																	n/e	L	
	e				m2			c							m2			e		
		<i>lower</i>																		
	e				m2										m1	m2			e	
	n/e			M1									I2	C			M1		n/e	

LRM1: Cr ¾ - Crc

LLI2: Cr ¾

LLC: Cr ¾

- Age at death: c. 1.5-2.5 years

Arma dell'Aquila 6 Richard: deciduous dentition erupted but not fully mineralized. It is possible to observe the presence of the first permanent molars of both jaws.

		<i>upper</i>																		
R	n/e			M1													M1		n/e	L
	e				m2	m1							c	m1	m2				e	
		<i>lower</i>																		
	e				m2	m1	c		i1	i1	i2	c	m1	m2					e	
	n/e			M1													M1		n/e	

LRdc: R ½ - R ¾

- Age at death: c. 1.5-2.5 years

Arene Candide 6632.1+6623.2: deciduous dentition erupted but not fully mineralized. Permanent dentition not fully mineralized.

		<i>upper</i>																		
R	n/e			M1					I2										n/e	L
	e				m2	m1	c												e	
		<i>lower</i>																		
	e					m1								m1					e	
	n/e			M1	m2		C	I2	I1	I1	I2	C			m2	M1			n/e	

URdm2: R ½ - R ¾

URI2: Cr ¾

URM1: Crc

LLdm2: R ¼ - R ½

- Age at death: c. 2-3 years

Arene Candide VIII BB: deciduous dentition erupted. Permanent dentition not fully mineralized.

		<i>upper</i>																	
R	n/e			M1						I1		C	P1	P2	M1	M2		n/e	L
	e				m2	m1	c	i2	i1			c						e	
		<i>lower</i>																	
	e				m2	m1	c	i2	i1	i1	i2	c	m1	m2				e	
	n/e			M2	M1						I2	C				M1	M2	n/e	

ULdc: Ac
 LL12: Ri
 LLC: Crc

➤ Age at death: c. 3.5-5.5 years (c. 4-5 yr)

Berggegi S2 01178 PE, 6894-5-FI: deciduous dentition erupted. Permanent dentition not fully mineralized. For many permanent teeth, it is not possible to determine the degree of mineralization.

		<i>upper</i>																		
R	n/e		M2	M1					I1	I1						M1	M2		n/e	L
	e				m2	dm1	c	i2	i1	i1	i2	c	m1	m2					e	
		<i>lower</i>																		
	e				m2	m1	c			i1	i2	c	m1	m2					e	
	n/e			M1									P1			M1	M2		n/e	

URM2-ULM2: Cr ½

➤ Age at death: c. 4.5-5.5 years

Pollera 21PE: deciduous dentition fully erupted. Permanent dentition not fully mineralized.

		<i>upper</i>																		
R	n/e			M1												M1			n/e	L
	e				m2	m1	c	i2	i1	i1	i2	c	m1	m2					e	
		<i>lower</i>																		
	e				m2	m1	c	i2			i2	c	m1	m2					e	
	n/e								I1	I1						M1	M2		n/e	

LLM1: R ¼

LLM2: Cr ½

➤ Age at death: c. 4-6 years

Pollera 6678.1: deciduous dentition fully erupted. Permanent dentition not fully mineralized. For many permanent teeth, it is not possible to determine the degree of mineralization.

		<i>upper</i>																		
R	n/e			M1												M1			n/e	L
	e				m2	m1	c	i2	i1			c	m1	m2					e	
		<i>lower</i>																		
	e				m2	m1						c	m1	m2					e	
	n/e		M2	M1												M1	M2		n/e	

LLM2-LRM2: Coc

➤ Age at death: c. 3.5-6.5 years

Arma dell'Aquila Zambelli JUV B: three isolated permanent teeth.

URM1-ULM1: R ½

LLM1: R ½

➤ Age at death: c. 6-7 years

Pipistrelli 1_dep23.I.2_JUV: mixed dentition, deciduous and permanent teeth are erupted. Permanent dentition not fully mineralized.

R	n/e							I2			I2						n/e	L
	e			M1	m2	m1							m1	m2	M1		e	
<i>lower</i>																		
	e			M1	m2	m1			I1	I1		c	m1	m2	M1		e	
	n/e							I2			I2					M2	n/e	

URI2-ULI2: R ¼
 URM1-ULM1: R ¾
 LRI1-LLI1: R ¾
 LRI2-LLI2: R ¾
 LRM1-LLM1: R ¾
 LLM2: Ri

➤ Age at death: c. 6-7 years

Arene Candide 6627.1: mixed dentition, deciduous and permanent teeth are erupted. Permanent dentition not fully mineralized.

R	n/e										C	P1	P2				n/e	L
	e							I1	I1				m2	M1			e	
<i>lower</i>																		
	e			M1	m2	m1	c	I2	I1	I1	I2	c	m1	m2	M1		e	
	n/e		M2					C				C				M2	n/e	

ULdm2: Res ¼
 LRdc-LLdc: Res ¼
 ULC: R ½
 ULP1: R ¼ - R ½
 ULP2: R ¼
 LLI1: Rc
 LRI2-LLI2: Rc - A ½
 LLC: R ½
 LLM1: R ¾ - Rc

➤ Age at death: c. 6-8 years

Pollera 20PE: mixed dentition, deciduous and permanent teeth are erupted. Permanent dentition not fully mineralized.

R	n/e		M2					I2			I2					M2	n/e	L
	e			M1	m2	m1	c		I1	I1		c	m1	m2	M1		e	
<i>lower</i>																		
	e			M1	m2	m1	c	i2	I1	I1	i2	c	m1	m2	M1		e	
	n/e		M2													M2	n/e	

URM2-ULM2: Crc - Ri
 LLM2: Crc

➤ Age at death: c. 6-8 years

Pollera 6682.1: mixed dentition. All teeth are isolated (no maxilla and/or mandible bones). Permanent dentition not fully mineralized.

		<i>upper</i>																		
R	n/e		M2				P1	C											n/e	L
	e			M1	m2	m1	c	I2	I1	I1	I2								e	
		<i>lower</i>																		
	e					m1	c	I2	I1	I1	I2								e	
	n/e											C	P1						n/e	

URdm1: Res ½

URdm2: Res ¼

URC: R ½

URP1: R ¼

URM2: R ¼

LLdm1: Res ½

LRI1-LLI1: A ½

LRI2-LLI2: Rc

LLC: R ½

LLP1: R ¼

➤ Age at death: c. 7-9 years

Strapatente I: mixed dentition, deciduous and permanent teeth are erupted. Permanent dentition not fully mineralized.

		<i>upper</i>																			
R	n/e		M2																	n/e	L
	e			M1	m2	m1	c	I2	I1	I1	I2	c	m1	m2	M1				e		
		<i>lower</i>																			
	e			M1	m2	m1	c	i2	I1	I1		c	m1	m2	M1				e		
	n/e		M2								I2								n/e		

URI1-ULI1: Rc

URI2-ULI2: R ¾

URM1-ULM1: R ¾ - Rc

URM2-ULM2: R ¼

➤ Age at death: c. 7.5-8.5 years

Arene Candide 3 Tinè: mixed dentition, deciduous and permanent teeth are erupted. Permanent dentition not fully mineralized.

		<i>upper</i>																			
R	n/e		M2																	n/e	L
	e			M1	m2	m1			I1		I2		m1	m2	M1				e		
		<i>lower</i>																			
	e			M1	m2	m1	c	I2	I1	I1	I2	c	m1	m2	M1				e		
	n/e	M3																	n/e		

URM2-ULM2: Crc

LRdc: Res ¼

LLI2: A ½

➤ Age at death: c. 7.5-9.5 years

Arene Candide 6730.3+6623.1+6625.2: permanent teeth not fully mineralized. It is possible to observe in the mandible the presence of the alveoli of deciduous teeth that are not preserved (probably mixed dentition). Maxillary teeth not present.

		<i>upper</i>																				
R	n/e																			n/e	L	
	e																			e		
		<i>lower</i>																				
	e																	M1		e		
	n/e																P1		M2	M3	n/e	

LLM3: Coc
 ➤ Age at death: c. 7-10 years

Pollera 1PE Issel-Morelli: permanent dentition not fully mineralized and erupted.

		<i>upper</i>																				
R	n/e	M3															P2			M3	n/e	L
	e		M2	M1	P2	P1	C	I2	I1	I1	I2	C	P1					M1	M2		e	
		<i>lower</i>																				
	e		M2	M1	P2	P1	C	I2	I1	I1	I2	C					P2	M1	M2		e	
	n/e	M3																		M3	n/e	

URM3- ULM3: Crc - Cr ½
 LRM3: Cr ½
 ➤ Age at death: c. 10-12 years

Arene Candide V BB: permanent dentition erupted. It is not possible to determine the degree of teeth mineralization.

		<i>upper</i>																				
R	n/e	M3																			n/e	L
	e		M2	M1	P2	P1	C	I2	I1	I1	I2	C	P1	P2	M1	M2					e	
		<i>lower</i>																				
	e		M2	M1																	e	
	n/e																				n/e	

➤ Age at death: c. 14-16 years

Arene Candide 6621.1: isolated permanent teeth not fully mineralized (no maxilla and/or mandible bones). Mandibular teeth not present.

		<i>upper</i>																				
R	n/e																			M3	n/e	L
	e		M2							I1		I2		P1	P2						e	
		<i>lower</i>																				
	e																				e	
	n/e																				n/e	

ULM3: R ½
 ➤ Age at death: c. 13-19 years

Arene Candide 1 Tinè: permanent dentition not fully erupted. It is not possible to determine the degree of teeth mineralization.

R	<i>n/e</i>	<i>upper</i>														<i>n/e</i>	L
		M3								I1	I2	C	P1	P2	M1		
	<i>e</i>		M2	M1	P2	P1	C	I2	I1	I2	C	P1	P2	M1	M2		<i>e</i>
		<i>lower</i>															
	<i>e</i>		M2	M1	P2	P1	C	I2	I1	I2	C	P1	P2	M1	M2		<i>e</i>
	<i>n/e</i>	M3														M3	<i>n/e</i>

➤ Age at death: c. 15-18 years