

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

The bianchetto truffle (*Tuber borchii*) a lead-resistant ectomycorrhizal fungus increases *Quercus cerris* phytoremediation potential

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Sisti, D., Donati Zeppa, S., Amicucci, A., Vittori Antisari, L., Vianello, G., Puliga, F., et al. (2022). The bianchetto truffle (*Tuber borchii*) a lead-resistant ectomycorrhizal fungus increases *Quercus cerris* phytoremediation potential. ENVIRONMENTAL MICROBIOLOGY, 24(12), 6439-6452 [10.1111/1462-2920.16273].

Availability:

This version is available at: <https://hdl.handle.net/11585/906424> since: 2022-11-23

Published:

DOI: <http://doi.org/10.1111/1462-2920.16273>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **The bianchetto truffle (*Tuber borchii*) a lead-resistant ectomycorrhizal fungus**
2 **increases *Quercus cerris* phytoremediation potential**

3

4 Davide Sisti¹, Sabrina Donati Zeppa¹, Antonella Amicucci¹, Livia Vittori Antisari²,
5 Gilmo Vianello², Federico Puliga², Pamela Leonardi², Mirco Iotti³, Alessandra
6 Zambonelli^{2*}

7

8 ¹Department of Biomolecular Sciences, University of Urbino Carlo Bo, 61029 Urbino, Italy

9 ²Department of Agricultural and Food Sciences, University of Bologna, Viale G. Fanin 44, 40127
10 Bologna, Italy

11 ³Department of Life, Health and Environmental Science, University of L'Aquila, via Vetoio,
12 67100, Coppito, L'Aquila, Italy

13

14 *Corresponding Author: alessandr.zambonelli@unibo.it, tel. + 39 051 2096572

15 Running title: *Tuber borchii-Quercus cerris* Pb tolerance

16

17 *Originality-significance statement*

18 This work firstly demonstrates the high tolerance of *Tuber borchii* mycelium and of *Quercus cerris*
19 seedlings mycorrhized with *T. borchii* to high Pb concentrations.

20 *Tuber borchii* mycorrhization as well as Pb treatment influence the uptake and translocation of Pb
21 and other elements within the host plant.

22 The results suggest that *T. borchii* - *Q. cerris* mycorrhized plants could have great potential for
23 practical application in bioremediation.

24

25

26 **Abstract**

27 *Tuber borchii* is a European edible truffle which forms ectomycorrhizas with several soft- and
28 hardwood plants. In this paper the effects of high level of Pb on the in vitro growth of five *T.*
29 *borchii* strains and the molecular mechanisms involved in Pb tolerance were studied. Moreover,
30 the effects of the Pb treatment on *T. borchii* ectomycorrhizas and on the growth, element uptake
31 and distribution in different organs of *Quercus cerris* seedlings were investigated. The results
32 showed an extraordinary tolerance of *T. borchii* mycelium to Pb: all the tested strains were able to
33 grow at Pb concentration over 4000 mg L⁻¹. The mechanisms of tolerance seem related to Pb
34 sequestration in the vacuole and its immobilization as crystal of Pb oxalate outside the hyphae
35 rather than detoxification processes, considering the low expression of glutaredoxin and
36 thioredoxin genes.

37 *T. borchii* - *Q. cerris* mycorrhizas tolerate a soil concentration of Pb from 1,869 to 4,030 mg kg⁻¹
38 although at these Pb concentrations *T. borchii* showed a reduced ability to colonize roots. *T. borchii*

39 mycorrhization increased the uptake of Pb by *Q. cerris*. Mycorrhization and Pb treatment also
40 significantly influenced the uptake and translocation in the plant of other elements.

41

42 **Key words:** lead nitrate, mycelium, detoxification mechanisms, ectomycorrhizas,
43 bioconcentration factor, translocation factor, element uptake

44

45 **Introduction**

46 Contamination of groundwater and soil with heavy metals is now one of the major problems of
47 pollution of natural environments and pose risks to human health as they enter the food chain via
48 agricultural products or contaminated drinking water (Cerbasi and Yetis, 2001; Zhang, 2020).

49 Acidification of forest soils in highly industrialized areas accompanied by high concentrations of
50 heavy metals determine the impoverishment of the fungal community and, a strong reduction of
51 biodiversity in different ecosystems (Fellner and Pešková, 1995; Lenart-Boroń and Boroń, 2014;

52 Passarini *et al.*, 2022). Among soil organisms, fungi are highly resistant to heavy metal pollution
53 and play important roles in element cycling, mineral transformation, plant stress tolerance and soil
54 bioremediation (Massaccesi *et al.*, 2002; Gadd, 2007; Văcar *et al.*, 2021). In particular,

55 mycorrhizal fungi have a high tolerance to heavy metals and are able to colonize areas that have
56 high concentrations of Pb^{2+} , Cu^{2+} , Zn^{2+} , Cd^{2+} , and Ni^{2+} (Hachani *et al.*, 2020; Yu *et al.*, 2020). The
57 biochemical and genetic mechanisms underlying heavy metal uptake and translocation in the plant

58 by mycorrhizal fungi are not completely known at this time, and their understanding is the basis
59 for developing new strategies for myco-remediation (Akhtar and Mannanab, 2020; Robinson *et*
60 *al.*, 2021). To fill this knowledge gap, specific *in vitro* and in greenhouse studies on the tolerance

61 and the accumulation capacities of mycorrhizal fungi to heavy metals are powerful strategies.

62 Among heavy metals, lead is one of the most harmful to plant and animal health. In the plants lead
63 toxicity causes disturbs in plant water and nutritional relations and causes oxidative damages; as
64 consequence, it inhibits the plant growth from seed germination (Zulfiqar *et al.*, 2019). The
65 oxidative damages caused by lead exposure have also detrimental effects on animals and humans
66 causing a range of cardiovascular, renal, neurologic, and hematologic dysfunctions with serious
67 health problems which might be permanent and lead to fatality (Assi *et al.*, 2016).

68 The diffusion of lead into the environment is primarily due to disposal of metals, the use of paints
69 based on Pb and the use of pesticides and fertilizers. Lead is also the main pollutant produced by
70 the combustion of leaded petrol (Page and Gange, 1970), and it was accumulated in intense traffic
71 areas for a long time (Wang and Zhang, 2018).

72 Truffles are hypogeous ectomycorrhizal ascomycetes belonging to the order of Pezizales. Some
73 species in the genus *Tuber* are highly appreciated for the organoleptic properties (i.e., taste and
74 flavor) of their edible ascomata (Mello *et al.*, 2006). *Tuber borchii* Vittad. is one of the most
75 common truffles in Europe being found from Finland to Sicily and from Spain to Iran, in different
76 soil conditions (Hall *et al.*, 2007; Puliga *et al.*, 2021) and, in contrast to other *Tuber* species, also
77 tolerates acidic soils (Gardin, 2005; Lancellotti and Franceschini, 2013) where heavy metals are
78 more available (Sintorini *et al.*, 2021). It can form mycorrhizas with many soft- and hard-wood
79 plants (Hall *et al.*, 2007). It is also the unique *Tuber* species which mycelium can growth easily in
80 vitro and, for this reason, it is considered a model species for studying the truffle biology and
81 genetics (Leonardi *et al.*, 2019). Previous *in vitro* studies carried out on agarized substrate have
82 shown that *T. borchii* increases the tolerance of *Cistus creticus* L. to zinc, lead and chromium
83 (Sabella *et al.*, 2016) but the molecular mechanisms involved in the tolerance to heavy metals of
84 a truffle species have never been investigated so far. Among the favorite hosts, *Quercus cerris* L.

85 is a pollution-tolerant tree native to southern Europe and Asia Minor which dominates in the mixed
86 forests of the Mediterranean basin (Najib *et al.*, 2021). This tree is often used as host plant in truffle
87 cultivation being able to form ectomycorrhizas with all the European edible *Tuber* species (Hall *et*
88 *al.*, 2007).

89 In this paper the effects of high level of Pb on *T. borchii* mycelium and the molecular mechanisms
90 related to its tolerance have been investigated by *in vitro* growth tests and gene expression
91 analyses. Moreover, the effects of Pb on *T. borchii* root colonization as well as the growth and
92 nutrient uptake of *Q. cerris* and the distribution of Pb and other elements in the different plant
93 organs (roots, stems and leaves) have been also tested. The potential of *Q. cerris* mycorrhized
94 plants for phytoremediation was also discussed.

95

96 **Results**

97 *In vitro* assessment of Pb on *T. borchii* mycelial growth

98 All *T. borchii* strains analyzed showed a dry weight not significantly dependent from Pb
99 concentration ($p=0.60$), whilst strains showed a different overall mean weight ($p<0.001$) (Fig. 1
100 and Tables S3). The oldest strain (1BO in culture from 1987) had the lowest dry weight at all the
101 tested Pb concentration and in the control (Fig. 1). However, all strains mostly responded to the
102 presence of Pb by increasing their diameter growth, in particular at the highest Pb concentrations
103 (Table S4, Fig. S1). Light microscope images of the Tbo5118 mycelium showed that the Pb
104 absorbed was sequestered into the vacuoles of mycelia grown at concentrations up to 2.56 mM
105 (Fig. 2a) while at higher Pb concentrations (up to 20.48 mM), a large number of needle- and star-
106 shaped crystals were formed on the cell wall surface of hyphae (Fig. 2b).

107

108 *Gene expression*

109

110 Changes in gene expression of mycelia treated or untreated with lead was verified by analyzing
111 the following five genes: 1) thioredoxin (TbThio), 2) glutaredoxin (TbGlut), 3) a putative
112 Na(+)/Li(+)-exporting P-type ATPase (TbNL), 4) TbRhoGdi, 5) TbCdc42. TbThio and TbGlut
113 were used to evaluate the possible activation of cell detoxification mechanisms, TbRhoGdi and
114 TbCdc42 are related to hyphal apical growth (Menotta et al., 2007; Picceri et al., 2018) and TbNL
115 was selected as possible gene involved in metal transport.

116 The expression of the genes *TbCdc42*, *TbRhoGdi* and *TbNL* was significantly higher in pure
117 culture of the strain Tbo5118 grown in presence of lead with respect to the controls (Fig. 3).
118 Instead, the expression analysis of *TbGlu* and *TbThio*, genes did not reveal significant differences.

119

120 *Effects of Pb on plant growth and on mycorrhizal infection*

121 Both mycorrhizal colonization with *T. borchii* and Pb treatment did not influence the height of the
122 plants one year after inoculation and eight months after the first Pb treatment, whereas the collar
123 diameter was only significantly higher in mycorrhizal plants (Table S5). No foliar symptoms were
124 observed in any plant treated with Pb (mycorrhized and non-mycorrhized: 1mic-1Pb and 0mic-
125 1Pb). At the second check of mycorrhizal infection two plants, one treated with Pb (1mic-1Pb) and
126 one not treated with Pb (1mic-0Pb), were found damaged and therefore were not considered in the
127 subsequent analyses. No contaminations with other fungal species were found. The plants
128 increased their mycorrhization degree with *T. borchii* 8 months after the first Pb treatment.
129 However, the mycorrhizas increased significantly more ($p < 0.05$) in the plants not treated with Pb
130 (+ 30% for 1mic-0Pb and + 5% for 1mic-1Pb, Fig. 4a). Microscope analyses of ectomycorrhizas

131 showed the presence of Pb in vacuoles of the extraradical mycelium (Fig. 4c) and needle-like Pb
132 crystals on the surface on some hyphae and cystidia (Fig. 4d, f). Inside the cross and longitudinal
133 sections of the mycorrhizas, lead was found to accumulate in the inner part of the mantle and in
134 the Hartig net (Fig. 4e,g).

135

136 *Phytoextraction efficiency*

137 At the end of the experiment, the mean soil Pb concentration was 2,006 and 3,594 mg kg⁻¹ in 1mic-
138 1Pb and 0mic-1Pb pots, respectively and reached 4,000 mg kg⁻¹ in two 0mic-1Pb pots. It was
139 significantly higher in the 0mic-1Pb pots than in the 1mic-1Pb pots (Fig. 5b). The mean Pb
140 concentration of untreated soils was 12.98 and 22.95 mg kg⁻¹ in 1mic0Pb and 0mic-0Pb pots,
141 respectively and no significant differences were found between them. The presence of *T. borchii*
142 mycorrhizas as well as the Pb treatment significantly increase the BFC (Fig. 5a). The Pb was
143 significantly more concentrated in roots and stem than in the leaves (Table 1). The Pb quantity was
144 significantly higher in all the organs of treated plants, whereas *T. borchii* mycorrhization
145 significantly increases the Pb content only in the stems (Table 1).

146

147 The treatment with Pb and the mycorrhization with *T. borchii* influenced also the concentration of
148 several elements in plant organs. In particular, the leaf concentration of K and Zn significantly
149 increases in 1Pb and 1mic. The Pb treatment also increases the leaf concentration of Ca, S, Li, Ni,
150 Sr in the leaf, whereas decreases the P concentration. In contrast, the mycorrhization with *T.*
151 *borchii* increased the P, Cu and Na concentration in the leaf and reduced the Se and Mn
152 concentration.

153 The concentration of K, Cu, P and Zn was also significantly higher in the stem of both 1mic and
154 1Pb plants. In contrast the Ba concentration was significantly lower in both 1mic and 1Pb plants.
155 The Pb treatment significantly increased Mo concentration in the stem and decreased the B content.
156 *Tuber borchii* mycorrhization decreased Ca concentration in the stem.
157 In the roots Ca, K, P and Sr concentration was significantly higher of both 1mic and 1Pb plants.
158 The Pb treatment also significantly increased the Na, S, Ba, Cu and Se concentration in the roots.
159
160 The translocation factors (TFs and TFI) reported in the Fig. 6 show how the elements are
161 translocated from the roots into the other organs. Most of heavy metals such as Pb, Mo, Cu, Cr, Fe
162 and Al are scarcely translocated in the leaf and in the stem. In contrast Ca, Mn and Ba are
163 translocated from the root to the stem and the leaf. P, Mg, Li, B, Na, S and K are principally
164 translocated in the leaf. There was no obvious effect of Pb treatment and mycorrhization on TFs
165 and TFI.

166

167 **Discussion**

168 In the last 25 years, *T. borchii* has been frequently used as a model species within the genus *Tuber*
169 for transcriptomic and functional studies and to shed light on issues of the truffle life cycle
170 (Zambonelli *et al.*, 2021). The responses of its mycelium and/or mycorrhizas to different stress
171 conditions such as high temperatures, nitrogen starvation or heavy metals have been also
172 investigated (Montanini *et al.*, 2006; Leonardi *et al.*, 2017; Sabella *et al.*, 2016).

173 In this work we adopted a multidisciplinary approach to analyses the effect of lead on mycelium
174 and mycorrhizas of *T. borchii* and the detoxification strategies implemented by the fungus as well
175 as the effect on plant nutrition.

176 In our study, *T. borchii* showed a very high tolerance to Pb concentration in the substrate. All the
177 tested strains tolerate a Pb concentration up to 4,200 mg L⁻¹ although they were isolated in non-
178 contaminated soils. This Pb concentration is higher than those previously tested for most of the
179 other mycorrhizal fungi (Table S6). Only the mycelia of fungal species isolated from contaminated
180 soils showed a similar tolerance to Pb (Liquat *et al.*, 2020). The mycelial biomass was little
181 affected by high Pb concentration in the substrate whereas the colony expansion was even slightly
182 stimulated by high Pb concentration in the substrate. In this regard, we evaluated the expression
183 of genes related to apical growth in fungi, such as *TbCdc42* and *TbRhoGdi* in the tested fungal
184 strains. The greater expression of these two genes in mycelia grown in the presence of lead
185 compared to the controls, demonstrated that *T. borchii* is stimulated in apical growth in the
186 presence of Pb. This behavior, in our opinion, can be interpreted as a defensive strategy adopted
187 by the fungus to colonize new parts of the substrate in search of unpolluted soil.

188 The microscopic analyses showed that the fungus is able to perform two different active
189 mechanisms in order to avoid the cellular toxic effects of the lead. As shown by the specific
190 coloration with sodium rhodizonate (Tung and Temple, 1996), Pb at low concentration is
191 sequestered in the vacuole of the fungal cell. This type of behavior would allow the fungus to
192 continue to perform its vital functions without suffering the toxic element. In fact, it has been
193 reported that ECM fungi are able to chelate metal by glutathione and to compartment in the vacuole
194 the metal-glutathione complex (Smith and Read, 2010).

195 At higher concentrations of Pb, up to 20.48 mM, we also noted the presence of a large number of
196 crystals of Pb on the hyphal cell wall. This led us to speculate that beyond a certain threshold
197 concentration, the Pb is inactivated, in crystalline form, and accumulated outside the fungal cells.
198 In this way the fungus would prevent the intracellular accumulation of potentially lethal amounts

199 of Pb. The formation of crystals was also observed on the mycorrhizal mantle as well as in the
200 cystidia and the extra-radical mycelium.

201 The precipitation of extracellular Pb could be enhanced by an active secretion of fungal exudates
202 in the wall, such as oxalic acid which was found to contribute to the 'outer defense line' to resist
203 Pb (Tian *et al.*, 2019). In fact, oxalic acid can react with Pb cations to form lead oxalate crystals
204 (Ceci *et al.*, 2015) which were easily recognized under microscopy due to their characteristic shape
205 and red color after staining with sodium rhodizonate.

206 These data are strictly correlated to the evidence highlighted by gene expression analyses, that
207 showed a higher expression of the *TbNL* gene in the presence of lead in the growth substrate. We
208 can assume that this gene may be involved in the compartmentalization in the vacuoles of Pb,
209 favoring an outward transport, to compensate for the low level of protective detoxification
210 enzymes. In fact, the lack of upregulation of thioredoxin and glutaredoxin genes in mycelia grown
211 in substrate added with 5.12 mM $\text{Pb}(\text{NO}_3)_2$ indicates that the cellular mechanisms of detoxification
212 were not activated.

213 The treatments with $\text{Pb}(\text{NO}_3)_2$ in the pots where mycorrhized and non mycorrhized plants were
214 grown allowed to reach a final Pb concentration in the soil variable from 1869 to 4030 mg/kg.
215 These concentrations are similar to those found in the highly polluted soil around mining areas in
216 China (Shi *et al.*, 2021). At so high Pb concentration no symptoms were observed in neither control
217 nor mycorrhized plants treated with lead. The plant development was also not affected by Pb
218 treatment. *Tuber borchii* ectomycorrhizas were not visibly damaged by the presence of Pb but the
219 total degree of mycorrhization at the end of the experiment increased significantly more in non-
220 treated plants. The lower level of root colonization in plants treated with Pb could be interpreted
221 as conflicting with the results obtained from mycelial growth tests. However, the presence of Pb

222 in soil may interfere with the symbiosis establishment, hindering the plant-fungus recognition or
223 slowing down the root colonization. The negative effect of Pb and other heavy metals on
224 abundance of ectomycorrhizas was already demonstrated by different authors (Chappelka *et al.*,
225 1991, Bojarczuk and Kieliszewska-Rokicka, 2010; Ouatiki *et al.*, 2021).

226 Nevertheless, the presence of *T. borchii* mycorrhizas significantly increase the uptake of Pb from
227 soil and its accumulation in plant tissues. These results are in contrast with the experiment carried
228 out on poplar microcuttings mycorrhized with *Paxillus involutus* where the presence of
229 ectomycorrhizas significantly reduced the bioaccumulation factor of Pb (Szuba *et al.*, 2021) but it
230 is coherent with Sun *et al.* (2022) who found that *Suillus luteus* enhanced Cd accumulation in oak
231 seedlings.

232 After absorption, Pb concentration in stems and roots of mycorrhized plants was, respectively, 13
233 and 15 times higher than leaves while this gap was reduced to about 4.5 times in plants without *T.*
234 *borchii* ectomycorrhizas. Our result is consistent with the reports of many similar studies that
235 demonstrate that Pb is mainly accumulated in the roots (Xiong, 1997; Escobar and Dussan, 2016
236 Szuba *et al.*, 2021) and partially translocated in the shoots. The leaf showed a low concentration
237 of Pb that was the probable reason why no foliar symptoms were detected during the four months
238 of experiments. As reported in literature foliar tissues are the final frontier for metal(oid) uptake
239 via the roots; thus, part of a plant's plan is to limit the entrance or translocation of metallic elements
240 into photosynthetic tissues (Clemens and Ma, 2016; Angulo-Bejarano, 2021).

241 The uptake and translocation of the elements in the plant was differentially affected by Pb
242 treatment and/or the symbiosis with *T. borchii*. Only 6 out of 21 analyzed elements (Fe, Al, Cr,
243 Mg, Sb, V) did not significantly changed their concentration in roots, stems or leaves regardless
244 the treatment. The effect of mycorrhizas was limited to 10 elements of which 6 increased and 3

245 decreased their concentration in one or more plant tissues while Ca was differentially translocated
246 in stems and roots. Pb treatment significantly increased the concentration of 13 different elements
247 in at least one of the plant tissues and reduced the concentration of only three elements in leaves
248 (P) or stems (B and Ba). In general, our results are in contrast with those reported by other authors
249 who report that Pb can disrupt the assimilation of some fertilization elements such as Ca, Fe, and
250 Mg and limit their absorption and/or transport to leaves (Houda *et al.*, 2016). In our work, the Pb
251 treatment increased the quantity of Ca as well as of other metals such as K, Li, Zn, Ni and Sr in
252 the leaves. On the other hand, the treatment with Pb increases the concentration of heavy metals
253 (such as Cu and Sr) in the roots of the plants treated with Pb especially in the mycorrhized plants.
254 As reported in literature, most of the Pb is translocated to the shoots after its absorption into the
255 roots using passive mechanisms that rely on H⁺/ATPase systems (Angulo-Bejarano, 2021). The
256 increased expression of *TbNL* in *T. borchii* mycelium in presence of Pb (this work) as well in
257 truffle mycorrhizas (Martin *et al.*, 2010) would explain the greater Pb and other metal uptake by
258 the mycorrhizal structure and their transport into the plant body. In fact, the protein encoded by
259 this gene has a homology with Na⁽⁺⁾/Li⁽⁺⁾-exporting P-type ATPase, a P-type sodium transporter
260 involved in several processes, including cellular response to glucose starvation, hyperosmotic
261 response and metal ion transport. Moreover, it has also a lower homology with a metal transporter
262 ATPases (OSNPB_060665800 Smith *et al.*, 2014, Amari *et al.*, 2017).

263 On the other hand, the high presence of Pb could have also induced changes in the activity of the
264 plant plasma membrane H⁺-ATPase and increased their transport into the host plant.

265 The Pb absorbed by *T. borchii* hyphae seems not to have accumulated in the mantle as suggested
266 by some works (Chot and Reddy, 2022) but transferred towards the Hartig net where Pb was found
267 more visible by staining with sodium rhodizonate.

268 Another hypothesis to explain the increasing Pb uptake in mycorrhizal plants is that the Pb uptake
269 of roots follows water flow, which is facilitated by the aquaporins as suggested by Zhang *et al.*
270 (2021) in plants of *Medicago truncatula* colonized with the arbuscular mycorrhizal fungus
271 *Rhizophagus irregularis*.

272 P concentration in the leaf was remarkably reduced in the plants treated with Pb. This could be
273 attributed to Pb phytotoxicity on major metabolic processes such as reducing water and nutrients
274 uptake and transport, chlorophyll formation with induced oxidative stress via production of
275 reactive oxygen species (Zulfiquar *et al.*, 2019). *Tuber borchii* mycorrhization promoted P uptake
276 and translocation compared to non-inoculated plants in the soil contaminated with Pb, similarly to
277 those observed with soybean plants inoculated with arbuscular mycorrhizal fungi (Adeyemi *et al.*,
278 2021). Similarly, a previous study on the heavy metal Cd showed that the mycorrhization of
279 *Quercus acutissima* seedlings even doubled the P content in the roots although only at low Cd
280 concentrations (0.1 mg/kg) (Sun *et al.*, 2022). It is supposed that soil exploration type (Agerer
281 2001) of the ectomycorrhizal fungi is one of the most important factor in P uptake. ECM fungal
282 species with long exploration type via abundant extraradical mycelium may have a competitive
283 advantage over species with contact exploration in the search for water and P (Köhler *et al.*, 2018).
284 Even though the mycorrhizas formed by *Tuber* spp. are considered of contact exploration type
285 (Agerer, 2006), they were able to significantly increase P uptake. On the other hand, *T. borchii* as
286 other ectomycorrhizal fungi seems to secrete organic acids and probably phosphatases which
287 improve the availability of soil P, or increase mycelium inputs to facilitate plant acquisition of P.
288 In the acidification process, organic acids secreted by *T. borchii* could reduce pH in the rhizosphere
289 soil and increase the exchange capacity of soluble and exchangeable cations (e.g., K⁺, and Ca²⁺),

290 thereby liberating these mineral elements increasing their uptake (Liu *et al.*, 2020) as shown by
291 our results.

292 The discovery of an ectomycorrhizal fungal species (*T. borchii*) highly tolerant to Pb and its ability
293 to increase Pb uptake by the host plant is of fundamental importance for phytoremediation.
294 *Quercus cerris* plants mycorrhized with *T. borchii* could have great potential for practical
295 application in phytoremediation due to the high biomass of its aerial parts and the low
296 environmental requirements which are considered important factors in plant bioremediation
297 (Escobar and Dussan 2016; Houda *et al.*, 2016; Kalubi *et al.* 2016; Yang *et al.* 2015; Mleczek *et*
298 *al.*, 2017). This work also contributed to a knowledge on *Q. cerris* accumulation and distribution
299 of trace elements, macronutrients and heavy metals which seem to reflect a specific growth pattern
300 in plant species (Subramanian *et al.*, 2022). Further detailed biochemical and molecular studies
301 are needed to decipher the physiological and molecular mechanisms underlying variation in
302 nutrient uptake and translocation in response to Pb contamination and *T. borchii* mycorrhization.

303

304 **Experimental procedures**

305 *Mycelial strains*

306 Five mycelial strains designated 1BO (ATCC 96540), 10RA, 17BO, 43BO and Tbo5118 isolated
307 from *T. borchii* fruiting bodies harvested in central Italy were used. These soils contain less than
308 30 mg/kg of Pb (Amorosi *et al.*, 2012) (Table S1). Identity of the strains was molecularly verified
309 by sequencing their ITS region (Bonuso *et al.*, 2006, 2010; in this work for Tbo5118). They were
310 maintained by subculturing on half strength Potato dextrose agar (PDA, Difco).

311

312 *In vitro assessment of Pb on T. borchii mycelial growth*

313 The response of *T. borchii* strains to various concentration of Pb (0.04, 0.08, 0.16, 0.32, 0.64, 1.28,
314 2.56, 5.12, 10.24, 20.48 mM) was assessed. Before adding the lead to the growth medium it was
315 melted in the form of $\text{Pb}(\text{NO}_3)_2$, in a solution with a chelating agent (EDTA sodium salt) at the
316 same molar concentration of $\text{Pb}(\text{NO}_3)_2$ as described by Vassil *et al.* (1998).

317 The isolates were grown in the dark at 24°C, with no agitation, in 100 L-flasks containing 70 ml
318 of a modified Murashige-Skoog liquid medium (MS/2) (Sisti *et al.*, 1998). Each flask was
319 inoculated with one plug (0.7 cm in diameter) from 30-d old cultures grown on PDA. Colony
320 diameter of each strain was measured every five days for a month in cultures grown in Petri dishes
321 (9 cm in diameter) on agarized MS/2 at the same conditions applied for liquid cultures. The pH of
322 each medium was adjusted to 5.6-5.8 after the addition of chelated lead and before autoclaving
323 them < 120 °C for 20 min.

324

325 *Gene expression assay*

326 RNA isolation

327 Total RNA was isolated from 1-month-old liquid cultures of Tbo5118 (the strain then used for
328 plant inoculation) added with 5.12 mM $\text{Pb}(\text{NO}_3)_2$ or without lead addition (control), using RNeasy
329 Plant-mini kit from QIAGEN, following the manufacturer's instructions. The final concentration
330 and quality of RNA samples were estimated either spectrophotometrically by a NanoDrop®ND-
331 1000 (Celbio) or by agarose gel electrophoresis, staining with ethidium bromide. Total RNA was
332 treated with DnaseI, DNA-free™ kit (Ambion), according to the manufacturer's instructions. Three
333 biological replicates were processed for each treatment.

334

335 Real Time PCR analysis

336 The reverse transcription reactions were performed in 20 µl volume reactions. Firstly, 1 µg of
337 DNase-treated total RNAs from *T. borchii* strain grown with or without Pb addition and 1 µl of
338 random hexamers (12 µM, Promega) were combined and incubated at 65° C for 2 min; 1 µl RT-
339 buffer, 0.5 mM dNTPs, 1 U RNase inhibitor and 4 U of MMLV Reverse Transcriptase (QIAGEN)
340 (for a total of 6 µl) were then added. The reactions were incubated at 37° C for 1 h and then at 72°
341 C for 5 min. Finally, the cDNAs were diluted 1:2 for the subsequent PCR reaction.

342 Suitable primer pairs for each selected gene with a high melting temperature (> 60 °C) were
343 designed. The primers used to amplify *TbThio* (BM56For 5'-CTTCCATCACACATCCATCAA-
344 3', BM56Rev 5'-AATCAGTTTGCAGGGACCAC-3'), *TbGlut* (BM55For 5'-
345 ACCCCGTTGCTTATCTTTTCC-3', BM55Rev 5'-CTCCTTGAGAGCAGCCTGG-3'), *TbNL*
346 (AF23For 5'-TGCCTGGTGACATTATCGAA-3', AF23Rev 5'-
347 CCCCAGTCAAAAGTGCTTCA-3') genes were designed on *TbThio* (BM266256), *TbGlut*
348 (BM266155, Lacourt *et al.*, 2002) sequences and AF23 Na(+)/Li(+)-exporting P-type ATPase EST
349 sequences (AF487323 Zeppa *et al.*, 2002), using the software Primer 3.0
350 (<http://www.bioinformatics.nl/cgi-bin/primer3plus.cgi>). Moreover *TbRhoGdi* and *TbCdc42*
351 primer pairs were used to amplify *TbRhoGdi* and *TbCdc42* genes, as reported in Menotta *et al.*
352 (2007). The 18S rRNA gene from *T. borchii* was selected as a reference (18S RT F 5'-
353 TGGTCCGGTCGGATCTT-3', 18S RT R 5'-CATTACGGCGGTCCTAGAAA-3') (Menotta *et*
354 *al.*, 2008). To avoid amplification of genomic DNA, all the primers used, were designed including
355 a splice junction.

356 Quantitative real time PCR (qPCR) was performed in a Bio-Rad iCycler iQ Multi-Color Real Time
357 PCR Detection System (Bio-Rad) using, for all genes, the following thermal parameters: 95° C for
358 10 min, followed by 50 cycles of 95° C for 30 s and 60° C for 30 s.

359 Each sample was analysed in triplicate in 25 µl reaction consisting of 1 µl diluted cDNA, 2X
360 QuantiTect SYBR Green PCR kit, 300 nM of primers and 0.6 U of Hot Rescue Real Time DNA
361 polymerase (Diatheva). The specificity of the amplification products was confirmed by examining
362 thermal denaturation plots and by sample separation in a 3% DNA agarose gel. The amount of
363 each target transcript was related to that of the reference gene using the method described by Pfaffl
364 (2001). The Pfaffl method is a comparative method calibrated on the single gene efficiency. Three
365 independent replicates of amplification products were used to calculate the means and standard
366 errors.

367

368 *Greenhouse experiments*

369 *Quercus cerris* acorns were germinated in sterile sand in a greenhouse. Six-week-old seedlings
370 were transplanted individually in plastic pots (Bamaplast, Italy), 7 cm diameter and 18 cm deep,
371 filled with 500 ml of a calcareous loam soil (pH in water, 7.79; total CaCO₃, 9.9%; organic matter,
372 2.18%; electrical conductivity, 0.33 dS m⁻¹; texture: sand 34.8%, silt 47.0%, clay 18.2%) mixed
373 with vermiculite and sand (8:1:1). The substrate was autoclaved at 120 °C for 2 h. Fifteen seedlings
374 were inoculated with *T. borchii*, using the mycelium of the strain Tbo5118 as inoculum, before
375 transplanting. This strain was chosen because it was isolated more recently than the other strains
376 and thus it is more infective (Iotti *et al.*, 2012). Mycelial inoculum was prepared following the
377 instruction reported in the Italian patent application nr. 102021000023342 filed on September 9,
378 2021. Fifteen uninoculated seedlings were also prepared and run in parallel as controls.

379 Six months after inoculation the high of the stem and the collar diameter of each seedling were
380 measured and all seedlings were removed from their pots to assess *T. borchii* mycorrhiza formation
381 and the presence of contaminants. The identity of the obtained *T. borchii* mycorrhizas was

382 morphologically (Zambonelli *et al.*, 1993) and molecularly confirmed under a dissecting
383 microscope ($\times 20$) and by direct PCR strategy (Iotti and Zambonelli, 2006) using *T. borchii*
384 species-specific primers (Amicucci *et al.*, 1998), respectively. The degree of mycorrhization was
385 measured by counting the number of colonized and non-colonized root tips of 3-4 root fragments
386 randomly selected roots from the top, middle and bottom sector of the root system. The results
387 were expressed as a percentage of colonized tips out of the total number of tips.

388 Ten seedlings mycorrhized with *T. borchii* (1mic) and ten control seedlings completely free of
389 ectomycorrhizal contaminants (0mic) were selected for the following experiments. An half of these
390 seedlings (five 1mic and five 0mic) were treated during 6 months with an increasing dose of
391 $\text{Pb}(\text{NO}_3)_2$ (1Pb) (Table S2) whereas the other seedlings (five 1mic and five 0mic) were not treated
392 with Pb (0Pb). EDTA sodium salt was added to the seedlings of the first two treatments at the
393 same molar concentration of $\text{Pb}(\text{NO}_3)_2$. In total each treated seedling received around 3.5 g of
394 $\text{Pb}(\text{NO}_3)_2$. Each pot was watered taking care to minimize water leakage and, in treated pots, Pb
395 leaching.

396 Two months after the last treatment the high of the stem and the collar diameter of each seedling
397 were measured; all the seedlings were removed from their pots and, after accurately washing the
398 roots system under tap water, the degree of ectomycorrhizal infection was assessed as previously
399 described. Then, each seedling was separated into roots, stems and leaves which were dried in an
400 oven at 60°C until constant weight for Pb content analyses.

401

402 *Morphological analyses of mycelium and ectomycorrhizas*

403 Before morphological observation, the mycelium from liquid cultures was gently washed in
404 distilled water. Morphological features of ectomycorrhizas were firstly observed under a dissecting

405 microscope (20x) and then hand-made cross sections of fresh unramified ends were carried out by
406 using a razor blade. Sodium rhodizonate was applied to mycelia and ectomycorrhizal sections to
407 visualize the lead particles forming a pink-colored lead–rhodizonate complex (Glater and
408 Hernandez 1972; Tung and Temple, 1996). All the microscopic observations were carried out
409 under an Eclipse TE 2000-E microscope (1000 X) (Nikon) and images captured with a DXM1200F
410 digital camera (Nikon).

411

412 *Pb content measurements in soil and plant tissues*

413 The different dried organs (root, stem and leaves) were ground in a titanium ball mill. 250 mg of
414 dried plant tissues were mineralized with 6 ml of HNO₃ (Suprapur, Merck, Kenilworth) and 2 ml
415 of H₂O₂ (Carlo Erba, MI, Italy) using a microwave oven (Milestone 2100, Sorisone). After
416 digestion, the sample solution was made up to 20 ml with Milli-Q ultrapure distilled water and
417 filtered with Whatman 42 filter paper. Pb concentration was determined in mineralized samples
418 by Inductively Coupled Plasma Optical Emission Spectrometry (ICP-OES, Spectro, Ametek, MI,
419 Italy), after appropriate calibration. The quality of the data analysis was assessed using an
420 International Standard (CRM 482, Community Bureaux of Reference, BCR).

421 Soils of the four treatments (0Pb-0mic, 1Pb-0mic, 0Pb-1mic and 1Pb-0mic) were air-dried and
422 ground (less than 2 mm in size) with a titanium ball mill (Vittori Antisari *et al.*, 2012, 2013, 2014).
423 Briefly, 250 mg of soil samples was mineralized with aqua regia solution (2 ml 65% HNO₃ plus 6
424 ml 37% HCl, suprapur grade, Carlo Erba) using a microwave oven (Milestone 2100, Sorisone).
425 After digestion, the soil sample solution was made up to 20 ml with Milli-Q ultrapure distilled
426 water and filtered with Whatman 42 filter paper. Pb concentration was determined by ICP-OES;

427 each soil sample was analysed three time and the data calibrated using International Reference
428 Materials (BCR) and internal laboratory standards (Ferronato *et al.*, 2021).

429

430 *Calculation of phytoextraction efficiency*

431 Capacity of mycorrhized and non-mycorrhized oak seedlings for Pb phytoextraction was estimated
432 based on the lead bioconcentration factor in roots (BFC) [BFC = Pb concentration in the roots (mg
433 kg^{-1}) / Pb concentration in the soil (mg kg^{-1})]. The translocation capacity of Pb and other analyzed
434 elements from roots to stems and to leaves was calculated as translocation factor (TF) [TFs = Pb
435 concentration stem (mg kg^{-1}) / Pb concentration in the root (mg kg^{-1}) and TF_l = Pb concentration
436 leaves (mg kg^{-1}) / Pb concentration in the roots (mg kg^{-1})] (Placek *et al.*, 2016). These parameters
437 were expressed as percentage values.

438

439 *Statistical analyses*

440 The dry weight of the mycelia as a function of the different strains (considered as a random factor),
441 and as a function of the Pb concentration (considered as covariate) was compared with a univariate
442 GLM model.

443 The data of plant growth (height and stem diameter), soil Pb content and BFC were compared
444 using a two-way ANOVA with Tukey's post hoc test. To establish the effects of Pb on *T. borchii*
445 mycorrhizal colonization a Before-After-Control-Impact (BACI) design (Smith, 2002) was
446 applied. BACI provides a way of comparing data obtained before treatment with data obtained
447 after treatment, as repeated measurement analysis of variance ANOVA with Tukey's post-hoc test.

448 The effects of Pb treatment and mycorrhizal symbiosis on the element uptake and distribution in
449 roots, shoots and leaves was analyzed by MANOVA analysis; mycorrhization and Pb treatment

450 were both binary predictive factors, while dependent variables were all the elements measured
451 (Table 1); significance of mycorrhization, Pb treatment and their interaction have been calculated.
452 RT-PCR expression signals were analyzed by Wilcoxon test for one sample, using Bonferroni
453 correction for multiple comparison. All elaborations were performed using SPSS statistical
454 package version 22.0 or XLSTAT.

455

456 **ACKNOWLEDGEMENTS**

457 We thank E. Vecchi for its help in laboratory analyses.

458

459 **CONFLICT OF INTEREST**

460 The authors declare that there is no conflict of interest.

461

462 **DATA AVAILABILITY**

463 Some of the data that support the findings of this study are available in the supplementary material
464 of this article. Any additional information will be made available upon request.

465

466 **References**

467 Adeyemi, N.O., Atayese, M.O., Sakariyawo, O.S., Azeez, J.O., and Ridwan, M. (2021) Arbuscular
468 mycorrhizal fungi species differentially regulate plant growth, phosphorus uptake and stress
469 tolerance of soybean in lead contaminated soil. *J Plant Nutr* 44 (11): 1633-1648.

470 <https://doi.org/10.1080/01904167.2021.1871748>

471 Agerer, R. (2001) Exploration types of ectomycorrhizae: a proposal to classify ectomycorrhizal
472 mycelial systems according to their patterns of differentiation and putative ecological importance.
473 Mycorrhiza 11: 107–114. <https://doi.org/10.1007/s005720100108>

474 Agerer, R. (2006) Fungal relationships and structural identity of their ectomycorrhizae. Mycol
475 Progress 5: 67–107. <https://doi.org/10.1007/s11557-006-0505-x>

476 Akhtar, N. and Mannanab M.A. (2020) Mycoremediation: expunging environmental pollutants.
477 Biotech Rep 26: e00452. <https://doi.org/10.1016/j.btre.2020.e00452>

478 Amari, T., Ghnaya, T. and C. Abdelly, C. (2017) Nickel, cadmium and lead phytotoxicity and
479 potential of halophytic plants in heavy metal extraction. S Afr J Bot 111: 99-110.
480 <https://doi.org/10.1016/j.sajb.2017.03.011>

481 Amicucci, A., Zambonelli, A., Giomaro, G., Potenza, L., and Stocchi, V. (1998) Identification of
482 ectomycorrhizal fungi of the genus *Tuber* by species-specific ITS primers. Mol Ecol 7: 273–277.
483 <https://doi.org/10.1046/j.1365-294X.1998.00357.x>

484 [Amorosi, A., Guermendi, M., Marchi, N. and Sammartino, I.\(2012\) The Pedogeochemical Map of](https://doi.org/10.1016/j.sajb.2017.03.011)
485 [the Emilia-Romagna plain \(1:250,000 scale\). Explanatory Notes. https://ambiente.regione.emilia-](https://doi.org/10.1016/j.sajb.2017.03.011)
486 [romagna.it/en/geologia/archivio_pdf/soil/pedogeochemical-map-1-250-000-explanatory-notes](https://doi.org/10.1016/j.sajb.2017.03.011)

487 Angulo-Bejarano, P. I., Puente-Rivera, J., and Cruz-Ortega, R. (2021) Metal and metalloid toxicity
488 in plants: An overview on molecular Aspects. Plants 10(4): 635.
489 <https://doi.org/10.3390/plants10040635>

490 Assi,M.A., Hezme, M,N., Haron, A.W., Sabri, M.Y.,and Rajion, M.A. (2016) The detrimental
491 effects of lead on human and animal health. Vet World 9(6): 660-671.
492 <https://doi.org/10.14202/vetworld.2016.660-671>

493 Bojarczuk, K., and Kieliszewska-Rokicka, B. (2010) Effect of ectomycorrhiza on Cu and Pb
494 accumulation in leaves and roots of silver birch (*Betula pendula* Roth.) seedlings grown in metal-
495 contaminated soil. *Water Air Soil Pollut* 207:227–240. [https://doi.org/10.1007/s11270-009-0131-](https://doi.org/10.1007/s11270-009-0131-8)
496 [8](https://doi.org/10.1007/s11270-009-0131-8)

497 Bonuso, E., Iotti, M., Macrì, A., and Zambonelli A. (2006) Approccio innovativo per
498 l'identificazione molecolare di funghi filamentosi. *Micologia Italiana* 35: 32 – 40.

499 Bonuso, E., Zambonelli, A., Bergemann, S., Iotti, M., and Garbelotto, M. (2010) Multilocus
500 phylogenetic and coalescent analyses identify two cryptic species in the Italian bianchetto truffle,
501 *Tuber borchii* Vittad. *Conserv Genet* 11: 1453-1466. <https://doi.org/10.1007/s10592-009-9972-3>

502 Ceci, A., Rhee, Y. J., Kierans, M., Hillier, S., Pendrowski, H., Gray, N., *et al.* (2015)
503 Transformation of vanadinite [Pb₅(VO₄)₃Cl] by fungi. *Environ Microbiol* 17: 2018–2034.
504 <https://doi.org/10.1111/1462-2920.12612>

505 Cerbasi, I.H., and Yetis, U. (2001) Biosorption of Ni (ii) and Pb (ii) by *Phanerochaete*
506 *chrysosporium* from binary metal system-kinetics. *Water Res* 27: 15–20.
507 <https://doi.org/10.4314/wsa.v27i1.5004>

508 Chappelka, A.H., Kush, J.S., Runion, G.B., Meier, S., and Kelley, W.D. (1991) Effects of soil-
509 applied lead on seedling growth and ectomycorrhizal colonization of loblolly pine. *Environment*
510 *Pollut* 72(4): 307-316. [https://doi.org/10.1016/0269-7491\(91\)90004-G](https://doi.org/10.1016/0269-7491(91)90004-G)

511 Chot, E., and Reddy, M.S. (2022) Role of ectomycorrhizal symbiosis behind the host plants
512 ameliorated tolerance against heavy metal stress. *Front Microbiol* 13:855473.
513 <https://doi.org/10.3389/fmicb.2022.855473>

514 Clemens, S., and Ma, J.F. (2016) Toxic heavy metal and metalloid accumulation in crop plants
515 and foods. *Annu. Rev Plant Biol* 67: 489–512. [https://doi.org/10.1146/annurev-arplant-043015-](https://doi.org/10.1146/annurev-arplant-043015-112301)
516 [112301](https://doi.org/10.1146/annurev-arplant-043015-112301)

517 Escobar, M.P., and Dussan, J. (2016) Phytoremediation potential of chromium and lead by *Alnus*
518 *acuminata* subsp. *acuminata*. *Environ Prog Sustain Energy* 35: 942–948.
519 <https://doi.org/10.1002/ep.12297>

520 Fellner, R., and Pešková, V. (1995) Effects of industrial pollutants on ectomycorrhizal
521 relationships in temperate forests. *Can J Bot* 73: S1310-S1315. <https://doi.org/10.1139/b95-392>

522 Ferronato, C., Vianello, G., De Feudis, M. and Vittori Antisari, L. (2021) Technosols development
523 in an abandoned mining area and environmental risk assessment. *Appl Sci* 11(15): 6982.
524 <https://doi.org/10.3390/app11156982>

525 Gadd, G.M. (2007) Fungi and industrial pollutants. In: *The Mycota, Volume IV: Environmental*
526 *and Microbial Relationships*. Kubicek, C.P., and Druzhinina, I.S. (eds). Berlin, Germany:
527 Springer-Verlag, pp. 68–84.

528 Gardin, L (2005) I tartufi minori in Toscana. Gli ambienti di crescita dei tartufi marzuolo e
529 scorzone. *Quaderno ARSIA*, January 2005. <https://doi.org/10.13140/RG.2.1.2509.5844>

530 Glater, R.A.B. and Hernandez, L. Jr. (1972) Lead Detection in Living Plant Tissue Using a New
531 Histochemical Method. *J Air Pollut Control Assoc* 22 (6): 463-467.
532 <https://doi.org/10.1080/00022470.1972.10469663>

533 Hachani, C., Lamhamedi, M.S., Cameselle, C., Gouveia, S., Zine El Abidine, A., Khasa, D.P.,
534 and Béjaoui, Z. (2020) Effects of ectomycorrhizal fungi and heavy metals (Pb, Zn, and Cd) on
535 growth and mineral nutrition of *Pinus halepensis* seedlings in North Africa. *Microorganisms* 8:
536 2033. <https://doi.org/10.3390/microorganisms8122033>

537 Hall, I, Brown, G, and Zambonelli A (2007) Taming the Truffle. The History, Lore, and Science
538 of the Ultimate Mushroom. Portland, OR: Timber Press.

539 Houda, Z., Bejaoui, Z., Albouchi, A., Gupta, D.K., and Corpas F.J. (2016) Comparative study of
540 plant growth of two poplar tree species irrigated with treated wastewater, with particular reference
541 to accumulation of heavy metals (Cd, Pb, As, and Ni). Environ Monit Assess 188: 99.
542 <https://doi.org/10.1007/s10661-016-5102-0>

543 Iotti, M. and Zambonelli, A. (2006) A quick and precise technique for identifying ectomycorrhizas
544 by PCR. Mycol Res 110: 60-65. <https://doi.org/10.1016/j.mycres.2005.09.010>

545 Iotti, M., Piattoni, F. and Zambonelli A. (2012) Techniques for host plant inoculation with truffles
546 and other edible ectomycorrhizal mushrooms. In Edible ectomycorrhizal mushrooms, current
547 knowledge and future prospects. Zambonelli, A., and Bonito, G.M. (eds). Soil Biology 34. Berlin:
548 Springer-Verlag, pp. 145 – 161. https://doi.org/10.1007/978-3-642-33823-6_9

549 Kalubi, K.N., Mehes-Smith, M., and Omri, A. (2016) Comparative analysis of metal translocation
550 in red maple (*Acer rubrum*) and trembling aspen (*Populus tremuloides*) populations from stressed
551 ecosystems contaminated with metals. Chem Ecol 32: 312–323.
552 <https://doi.org/10.1080/02757540.2016.1142978>

553 Köhler, J., Yang, N., Pena, R., Raghavan, V., Polle, A., and Meier, I.C. (2018) Ectomycorrhizal
554 fungal diversity increases phosphorus uptake efficiency of European beech. New Phytol 220 (4):
555 1200-1210. <https://doi.org/10.1111/nph.15208>

556 Lacourt, I., Duplessis, S., Abbà, S., Bonfante, P., and F. Martin, F. (2002) Isolation and
557 characterization of differentially expressed genes in the mycelium and fruit body of *Tuber borchii*.
558 Appl Environ Microbiol 68: 4574-4582. <https://doi.org/10.1128/AEM.68.9.4574-4582.2002>

559 Lancellotti, E., and Franceschini, A. (2013) Studies on the ectomycorrhizal community in a
560 declining *Quercus suber* L. stand. *Mycorrhiza* 23:533-542. [https://doi.org/10.1007/s00572-013-](https://doi.org/10.1007/s00572-013-0493-z)
561 [0493-z](https://doi.org/10.1007/s00572-013-0493-z)

562 Lenart-Boroń, A., and Boroń, P. (2014) The effect of industrial heavy metal pollution on microbial
563 abundance and diversity in soils – A review. In *Environmental risk assessment of soil*
564 *contamination*. Hernandez-Soriano, M.C. (ed). London: IntechOpen, pp.759-783.
565 <https://doi.org/10.5772/57406>

566 Leonardi, P., Iotti, M., Donati Zeppa, S., Lancellotti, E., Amicucci, A., and Zambonelli, A. (2017)
567 Morphological and functional changes in mycelium and mycorrhizas of *Tuber borchii* due to heat
568 stress. *Fungal Biol* 29: 20-29. <http://dx.doi.org/10.1016/j.funeco.2017.05.003>

569 Leonardi, P., Murat, C., Puliga, F, Iotti, M., and Zambonelli A. (2019) Ascoma genotyping and
570 mating type analyses of mycorrhizas and soil mycelia of *Tuber borchii* in a truffle orchard
571 established by mycelial inoculated plants. *Environ Microbiol.* 22(3) 964-975.
572 <https://doi.org/10.1111/1462-2920.14777>

573 Liaquat, F., Munis, M.F.H., Haroon, U., Arif, S., Saqib, S., Zaman, W., *et al.*, (2020) Evaluation
574 of metal tolerance of fungal strains isolated from contaminated mining soil of Nanjing, China.
575 *Biology (Basel)*. 9(12): 469. <https://doi.org/10.3390/biology9120469>

576 Liu, Y., Li, X., and Kou, Y. (2020) Ectomycorrhizal fungi: participation in nutrient turnover and
577 community assembly pattern in forest ecosystems. *Forests* 11: 453.
578 <https://doi.org/10.3390/f11040453>

579 Martin, F., Kohler, A., Murat, C., Balestrini, R., Coutinho, P.M., Jaillon O., *et al.* (2010). Périgord
580 black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* 464:
581 1033-1038. <https://doi.org/10.1038/nature08867>

582 Massacesi, G., Romero, M.C., Cazau, M.C. and Bucsinszky, A.M. (2002) Cadmium removal
583 capacities of filamentous soil fungi isolated from industrially polluted sediments, in La Plata
584 (Argentina). *World J Microbiol Biotechnol* 18: 817-820.
585 <https://doi.org/10.1023/A:1021282718440>

586 [Mello, A., Murat, C., and Bonfante P. \(2006\) Truffles: much more than a prized and local fungal
587 delicacy. *FEMS Microbiol Lett* 260\(1\): 1-8. <https://doi.org/10.1111/j.1574-6968.2006.00252.x>](#)

588 Menotta, M., Amicucci, A., Basili, G., Rivero, F., Polidori, E., Sisti, D., and Stocchi V. (2007)
589 Molecular characterisation of the small GTPase CDC42 in the ectomycorrhizal fungus *Tuber*
590 *borchii* Vittad. *Protoplasma* 231: 227–237. <https://doi.org/10.1007/s00709-007-0254-y>

591 Menotta, M., Amicucci, A., Basili, G., Polidori, E., Stocchi V., and Rivero, F. (2008) Molecular
592 and functional characterization of a Rho GDP dissociation inhibitor in the filamentous fungus
593 *Tuber borchii*. *BMC Microbiol* 8: 57. <https://doi.org/10.1186/1471-2180-8-57>

594 Mleczek, M., Goliński, P., Krzesłowska, M., Gąsecka, M., Magdziak, Z., Rutkowski, P., *et al.*
595 (2017) Phytoextraction of potentially toxic elements by six tree species growing on hazardous
596 mining sludge. *Environ Sci Pollut Res Int* 24(28): 22183-22195. [https://doi.org/10.1007/s11356-
597 017-9842-3](https://doi.org/10.1007/s11356-017-9842-3)

598 Montanini, B., Gabella, S., Abbà, S., Peter, M., Kohler, A., Bonfante, P., *et al.* (2006) Gene
599 expression profiling of the nitrogen starvation stress response in the mycorrhizal ascomycete *Tuber*
600 *borchii*. *Fungal Genet Biol* 43(9): 630-641. <https://doi.org/10.1016/j.fgb.2006.04.001>

601 Najib, R., Hour, T., Khairallah, Y. and Khalil, M. (2021) *Quercus cerris* L.: An Overview. –
602 *Forestry Studies* 74:1–9, <https://doi.org/10.2478/fsmu-2021-0001>

603 Ouatiki, E., Tounsi, A., Amir, S., Midhat, L., Radi, M., and Ouahmane, L. (2021) Inoculation of
604 *Pinus halepensis* with the ectomycorrhizal fungi *Scleroderma* helps in phytoremediation of soil

605 polymetallic pollution. Pol J Environ Stud 30(6): 5669-5680.
606 <https://doi.org/10.15244/pjoes/131979>

607 Page, A.L., and Gange, T.J. (1970). Accumulation of lead in soils for regions of high and low
608 motor vehicle traffic density. Environ Sci Technol 4: 140-142.
609 <https://doi.org/10.1021/es60037a001>

610 Passarini, M.R.Z., Ottoni, J.R., Costa, P.E.dS., Hissa, D.C., Falcão, R.M., Melo, V.M.M, *et al.*
611 (2022) Fungal community diversity of heavy metal contaminated soils revealed by metagenomics.
612 Arch Microbiol 204: 255. <https://doi.org/10.1007/s00203-022-02860-7>

613 Pfaffl, M.W. (2001) A New Mathematical Model for Relative Quantification in Real-Time RT-
614 PCR. Nucleic Acids Res 29: e45. <http://dx.doi.org/10.1093/nar/29.9.e45>

615 Picceri, G.G., Leonardi, P., Iotti, M., Gallo, M. Baldo, F., Zambonelli, A. et al. (2018) Bacteria-
616 produced ferric exopolysaccharide nanoparticles as iron delivery system for truffles (*Tuber*
617 *borchii*). Appl Microbiol Biotechnol 102: 1429–1441. [https://doi.org/10.1007/s00253-017-8615-](https://doi.org/10.1007/s00253-017-8615-8)
618 [8](https://doi.org/10.1007/s00253-017-8615-8)

619 Placek, A., Grobelak, A., and Kacprzak , M. (2016) Improving the phytoremediation of heavy
620 metals contaminated soil by use of sewage sludge. Int J Phytoremediation 18(6): 605-618.
621 <https://doi.org/10.1080/15226514.2015.1086308>

622 Puliga, F., Illice, M., Iotti, M., Leonardi, P., Baghdadi, A., Mozafari, A.A., and Zambonelli, A.
623 (2021) True truffle diversity in Iran. It J Mycol 50: 52-62. [https://doi.org/10.6092/issn.2531-](https://doi.org/10.6092/issn.2531-7342/12822)
624 [7342/12822](https://doi.org/10.6092/issn.2531-7342/12822)

625 Robinson, J.R., Isikhuemhen, O.S. and Anike, F.N. (2021) Fungal–metal interactions: A Review
626 of Toxicity and Homeostasis. J. Fungi 7: 225. <https://doi.org/10.3390/jof7030225>

627 Sabella, E., Nutricati, E., Aprile, A., Miceli, A., Negro, C., Rampino, P. *et al.* (2016) *Tuber borchii*
628 Vitt. mycorrhiza protects *Cistus creticus* L. from heavy metal toxicity. *Environ Exp Bot* 130: 181-
629 188. <https://doi.org/10.1016/j.envexpbot.2016.06.007>

630 Shi, J., Du, P., Luo, H., Chen, J., Zhang, Y., Wu, M., and Xu, G. (2021). Characteristics and risk
631 assessment of soil polluted by lead around various metal mines in China. *Int J Environ Res Public*
632 *Health* 18(9): 4598. <https://doi.org/10.3390/ijerph18094598>

633 Sintorini M. M., Widyatmoko H., Sinaga E., and Aliyah N. (2021) Effect of pH on metal mobility
634 in the soil. *IOP Conf Ser: Earth Environ Sci* 737: 012071. <https://doi.org/10.1088/1755-1315/737/1/012071>

635

636 Sisti, D., Zambonelli, A., Giomaro, G., Rossi, I., Ceccaroli, P., Citterio, B., *et al.* (1998) In vitro
637 mycorrhizal synthesis of micropropagated *Tilia platyphyllos* Scop. plantlets with *Tuber borchii*
638 Vittad. mycelium in pure culture. *Acta Horti* 457 379–387.
639 <https://doi.org/10.17660/ActaHortic.1998.457.47>

640 Smith, A.T., Smith, K.P. and Rosenzweig, A.C. (2014) Diversity of the metal-transporting P1B-
641 type ATPases. *J Biol Inorg Chem* 19, 947–960. <https://doi.org/10.1007%2Fs00775-014-1129-2>

642 Smith, E. (2002) BACI design. In *Encyclopedia of Environmetrics* El-Shaarawi, A.H., and
643 Piegorsch, W.W. (eds). Chichester: John Wiley & Sons Ltd, vol.1, pp. 141e148.

644 Smith, S. E., and Read, D. J. (2010) *Mycorrhizal Symbiosis*. London: Academic press.

645 Subramanian, D., Subha, R., and Murugesan, A.K. (2022) Accumulation and translocation of trace
646 elements and macronutrients in different plant species across five study sites. *Ecol Ind* 135:
647 108522. <https://doi.org/10.1016/j.ecolind.2021.108522>

648 Sun, W., Yang, B., Zhu, Y., Wang, H., Qin, G., and Yang, H. (2022) Ectomycorrhizal fungi
649 enhance the tolerance of phytotoxicity and cadmium accumulation in oak (*Quercus acutissima*

650 Carruth.) seedlings: modulation of growth properties and the antioxidant defense responses.
651 Environ Sci Pollut Res 29(5): 6526-6537. [https://doi-org.ezproxy.unibo.it/10.1007/s11356-021-](https://doi-org.ezproxy.unibo.it/10.1007/s11356-021-16169-3)
652 [16169-3](https://doi-org.ezproxy.unibo.it/10.1007/s11356-021-16169-3)

653 Szuba, A., Marczak, Ł., and Kozłowski, R. (2021) Pb stress and ectomycorrhizas: strong protective
654 proteomic responses in poplar roots inoculated with *Paxillus involutus* isolate and characterized
655 by low root colonization intensity. Int J Mol Sci 22: 4300. <https://doi.org/10.3390/ijms22094300>

656 Tian, D., Jiang, , Z., Jiang, L., Su, M. and Hu S. (2019) A new insight into lead (II) tolerance of
657 environmental fungi based on a study of *Aspergillus niger* and *Penicillium oxalicum*. Environ
658 Microbiol 21: 471-479. <https://doi.org/10.1111/1462-2920.14478>

659 Tung, G., and Temple, P. J. (1996) Histochemical detection of lead in plant tissues. Environ
660 Toxicol Chem 15(6): 906-914. <https://doi.org/10.1002/etc.5620150612>

661 Văcar, C. L., Covaci, E., Chakraborty, S., Li, B., Weindorf, D. C., Frențiu, T., Pârvu, M., and
662 Podar, D. (2021) Heavy metal-resistant filamentous fungi as potential mercury bioremediators. J
663 Fungi (Basel, Switzerland) 7(5): 386. <https://doi.org/10.3390/jof7050386>

664 Vassil, A.D., Kapulnik, Y., Raskin, I., and Salt, D.E. (1998) The role of EDTA in lead transport
665 and accumulation by Indian Mustard. Plant Physiol 117: 447-453.
666 <https://doi.org/10.1104/pp.117.2.447>

667 Vittori Antisari, L., Carbone, S., Ferronato, C., Simoni, A., and Vianello, G. (2012) Leaf washing
668 as an assessment tool to characterize dry atmospheric deposition. Int J Environ Qual 9:37–50.
669 <https://doi.org/10.6092/issn.2281-4485/3737>

670 Vittori Antisari, L., Cremonini, S., Desantis, P., Calastri, C., and Vianello, G. (2013) Chemical
671 characterisation of anthro-technosols from Bronze to Middle Age in Bologna (Italy). J Archaeol
672 Sci 40(10): 3660-3671. <https://doi.org/10.1016/j.jas.2013.04.023>

673 Vittori Antisari, L., Bianchini, G., Dinelli, E., Falsone, G., Gardini, A., Simoni, A., *et al.* (2014).
674 Critical evaluation of an intercalibration project focused on the definition of new multi-element
675 soil reference materials (AMS-MO1 and AMS-ML1). *EQA - J Environ Qual* 15(15): 41–64.
676 <https://doi.org/10.6092/issn.2281-4485/4553>

677 Wang, M., and Zhang, H. (2018) Accumulation of heavy metals in roadside soil in urban area and
678 the related impacting factors. *Int J Environ Res Public Health*, 15(6): 1064.
679 <https://doi.org/10.3390/ijerph15061064>

680 Xiong, Z. T. (1997) Bioaccumulation and physiological effects of excess lead in a roadside pioneer
681 species *Sonchus oleraceus* L. *Environ Pollut* 97: 275–279. [https://doi.org/10.1016/S0269-](https://doi.org/10.1016/S0269-7491(97)00086-9)
682 [7491\(97\)00086-9](https://doi.org/10.1016/S0269-7491(97)00086-9)

683 Yang, J., Li, K., Zheng, W., Zhang, H., Cao, X., Lan, Y., *et al.* (2015) Characterization of early
684 transcriptional responses to cadmium in the root and leaf of Cd-resistant *Salix matsudana* Koidz.
685 *BMC Genom* 16:705. <https://doi.org/10.1186/s12864-015-1923-4>

686 Yu, P., Sun, Y., Huang, Z., Zhu, F., Sun, Y., and Jiang, L. (2020) The effects of ectomycorrhizal
687 fungi on heavy metals' transport in *Pinus massoniana* and bacteria community in rhizosphere soil
688 in mine tailing area. *J Hazard Mater* 381: 121203. <https://doi.org/10.1016/j.jhazmat.2019.121203>

689 Zambonelli, A., Iotti, M., Puliga, F., and Hall, I.R. (2021) Enhancing white truffle (*Tuber*
690 *magnatum* Picco and *T. borchii* Vittad.) cultivation through biotechnology innovation. In:
691 *Advances in Plant Breeding Strategies: Vegetable Crops*. Al-Khayri, J.M., Jaim, S.M., and
692 Johnson, D.V.(eds). Cham: Springer, pp505-532. https://doi.org/10.1007/978-3-030-66969-0_14

693 Zambonelli, A., Salomoni, S. and Pisi A. (1993) Caratterizzazione anatomo-morfologica e
694 micromorfologica delle micorrize di *Tuber* spp. su *Quercus pubescens* Willd. *Micologia Italiana* 3:
695 73-90.

696 Zeppa, S., Guidi, C., Zambonelli, A., Potenza, L., Vallorani, L., Pierleoni R., *et al.* (2002)
697 Identification of putative genes involved in the development of *Tuber borchii* fruit body by mRNA
698 differential display in agarose gel. *Curr Genet* 42: 161–168. [https://doi.org/10.1007/s00294-002-](https://doi.org/10.1007/s00294-002-0343-6)
699 [0343-6](https://doi.org/10.1007/s00294-002-0343-6)

700 Zhang, H., Ren, W., Zheng, Y., Li, Y., Zhu, M. and Tang, M. (2021) Arbuscular mycorrhizal fungi
701 Increase Pb uptake of colonized and non-colonized *Medicago truncatula* root and deliver extra Pb
702 to colonized root segment. *Microorganisms* 9 :
703 1203. <https://doi.org/10.3390/microorganisms9061203>

704 Zhang, Y., O'Connor, D., Xu, W., and Hou, D. (2020). Blood lead levels among Chinese children:
705 the shifting influence of industry, traffic, and e-waste over three decades, *Environ Int* 135 :105379.
706 <https://doi.org/10.1016/j.envint.2019.105379>

707 Zulfiqar, U., Farooq M., Hussain, S., Maqsood, M., Hussain, M., Ishfaq, M. *et al.*, (2019) Lead
708 toxicity in plants: impacts and remediation. *J Environ Manag* 250: 109557.
709 <https://doi.org/10.1016/j.jenvman.2019.109557>

710

711

712

713 **Legends of the figures**

714 Fig. 1 - Dry weights (mean and standard deviation) of the 5 *Tuber borchii* strains, cultivated at
715 different Pb concentration. The complete dataset is reported in Table S3.

716 Fig. 2 - *Tuber borchii* hyphae treated with Pb. The pink-colored elements correspond to the lead–
717 rhodizonate complex inside the vacuoles (a), or to oxalate crystals outside the hyphae (b).

718 Fig. 3 – Relative expression of *TbCdc42*, *TbRhoGdi*, *TbGlu*, *TbThio*, *TbNL* genes in *T. borchii*
719 strain Tbo5118 grown on MS/2 liquid medium treated with lead compared to the controls (dotted
720 line). Data are mean \pm standard error of at least three independent experiments each performed in
721 triplicate. The expression data were expressed as fold expression versus C. Asterisks indicate
722 significant differences with the control (** $p < 0.01$, *** $p < 0.001$).

723 Fig. 4 - Difference between the percentages of mycorrhizal colonization of the control plants (0Pb,
724 green column) and those treated with Pb (Pb1) before the first Pb treatment and two months after
725 the last Pb treatment (a). *Tuber borchii* mycorrhizas untreated with Pb (0Pb) (b) and treated with
726 Pb (1Pb) (c,d, e, f and g). The pink-colored elements correspond to the lead–rhodizonate complex
727 inside the vacuoles of extra-radical hyphae (c), to the oxalate crystals on the cystidia surface (d
728 and f), lead–rhodizonate complex in the hyphae forming the Hartig net (e and g) and in the
729 innermost part of the mantle (g).

730 Fig. 5 - Lead concentration in the pot soil (a) and BFC (b) of the plants treated (1Pb) or untreated
731 with Pb (0Pb), mycorrhized (1mic) or not mycorrhized (0mic) with *T. borchii*. Bars indicate
732 standard error. p values determined by two-way ANOVA were the following: $p(\text{Mic}) = 0.000$,
733 $p(\text{Pb}) = 0.0001$, $p(\text{Mic} \times \text{Pb}) = 0.002$ for the soil lead concentration (a); $p(\text{Mic}) = 0.038$,
734 $p(\text{Pb}) = 0.003$, $p(\text{Mic} \times \text{Pb}) = 0.05$ for the BFC (b).

735 Fig. 6 - TFs (black bar) and TFI (grey bar) values (log₁₀ scale) of the principal element measured;
736 in ordinate symbols of element and status of plants analyzed (0mic-0Pb: non mycorrhizal plant,
737 without Pb; 1mic-0Pb: mycorrhizal plant, without Pb; 0mic-1Pb: non mycorrhizal plant, with Pb;
738 1mic-1Pb: mycorrhizal plant, with Pb). For figure clarity, some elements are omitted (Ni, V, Zn,
739 and Sr).

740

741 **Supporting Information**

742 Additional Supporting Information may be found in the online version of this article at the
743 publisher's web-site:

744 **Table S1** - Metadata of the *Tuber borchii* strains selected in this work. Herbarium numbers
745 are of dried specimens in the Mycology Center, University of Bologna, Italy (CMI-Unibo).
746 The accession number is referred to the ITS sequences deposited in GenBank.

747 **Table S2** - Timetable of the Pb treatments

748 **Table S3** – Dry weight (mean \pm standard deviation) of the mycelia of different *T. borchii* strains
749 grown on liquid MS/2 untreated (Control) or treated with different Pb concentrations.

750 **Table S4** – Colony diameter (mean \pm standard deviation) of the different *T. borchii* strains grow
751 on agarized MS/2 untreated (Control) or treated with different Pb concentrations.

752 **Table S5** – Height and collar diameter of the plants at the end of the experiment (one year after
753 inoculation and eight months after the first Pb treatment).

754 **Fig. S1** – Grow curve of the different *T. borchii* strains grown on agarized MS/2 untreated or
755 treated with 2.53, 10.24 and 20.48 mM of Pb. The data obtained at all Pb concentrations are
756 reported in Table S4.

757

758

759

760

761

762

763

764

TABLE 1 Element concentration in the organs of the plants treated (1Pb) or untreated with Pb (0Pb), mycorrhized (1mic) and non-mycorrhized (0mic) with *Tuber borchii*. In bold the significant values of *p*

	0mic-0Pb (ppm)	0mic-1Pb (ppm)	1mic-0Pb (ppm)	1mic-1Pb (ppm)	<i>p</i> (mic)	<i>p</i> (Pb)	<i>p</i> (mic × Pb)
Leaf							
Al	0.07 ± 0.02	0.07 ± 0.09	0.11 ± 0.04	0.08 ± 0.02	0.429	0.602	0.505
Ca	10.4 ± 0.85	15.8 ± 4.07	11.8 ± 5.87	16.7 ± 3.04	0.556	0.018	0.898
Fe	0.15 ± 0.03	0.14 ± 0.10	0.19 ± 0.07	0.13 ± 0.03	0.638	0.335	0.409
K	4.86 ± 0.88	6.41 ± 1.29	5.98 ± 0.99	9.02 ± 0.84	0.003	0.001	0.167
Mg	3.26 ± 0.50	2.93 ± 0.53	2.60 ± 0.26	3.17 ± 0.42	0.345	0.594	0.062
Mn	0.31 ± 0.06	0.26 ± 0.02	0.17 ± 0.04	0.19 ± 0.02	<0.001	0.544	0.088
Na	2.36 ± 0.82	1.97 ± 1.31	3.83 ± 1.55	4.09 ± 1.54	0.017	0.923	0.632
P	0.51 ± 0.10	0.43 ± 0.04	0.93 ± 0.16	0.62 ± 0.07	<0.001	0.001	0.026
S	0.94 ± 0.14	1.21 ± 0.08	0.99 ± 0.10	1.23 ± 0.03	0.508	<0.001	0.696
B	54.63 ± 14.6	59.8 ± 23.4	65.9 ± 23.5	77.8 ± 14.7	0.154	0.392	0.734
Ba	37.6 ± 4.16	43.1 ± 14.0	40.5 ± 23.7	37.3 ± 9.25	0.840	0.873	0.553
Cr	2.31 ± 1.35	2.72 ± 3.00	4.12 ± 3.32	1.95 ± 1.26	0.673	0.478	0.304
Cu	4.92 ± 0.48	6.40 ± 0.52	8.81 ± 1.24	8.84 ± 0.93	<0.001	0.086	0.099
Li	3.27 ± 0.60	5.96 ± 1.07	4.24 ± 1.81	6.42 ± 1.44	0.277	0.002	0.696
Mo	0.43 ± 0.16	0.34 ± 0.10	0.42 ± 0.14	0.35 ± 0.01	0.952	0.165	0.911
Ni	0.28 ± 0.30	1.36 ± 0.78	0.54 ± 0.41	1.02 ± 0.42	0.895	0.010	0.271
Pb	0.71 ± 0.26	295 ± 215	0.28 ± 0.24	167 ± 30.0	0.290	0.002	0.293
Sb	0.36 ± 0.03	0.35 ± 0.07	0.69 ± 0.61	0.39 ± 0.09	0.226	0.313	0.316
Se	0.36 ± 0.10	0.26 ± 0.06	0.18 ± 0.17	0.19 ± 0.06	0.030	0.423	0.330
Sr	45.7 ± 3.76	84.0 ± 26.3	55.7 ± 27.1	89.1 ± 21.6	0.498	0.006	0.824
V	0.03 ± 0.02	0.08 ± 0.16	0.06 ± 0.10	0.01 ± 0.01	0.740	0.929	0.356
Zn	15.5 ± 1.63	18.9 ± 5.10	22.2 ± 1.86	49.5 ± 26.7	0.013	0.033	0.085
Stem							
Al	0.15 ± 0.05	0.23 ± 0.11	0.18 ± 0.07	0.13 ± 0.01	0.304	0.618	0.069
Ca	14.1 ± 1.34	14.5 ± 1.47	10.4 ± 0.77	12.1 ± 1.40	<0.001	0.116	0.301
Fe	0.21 ± 0.05	0.29 ± 0.12	0.21 ± 0.04	0.14 ± 0.01	0.058	0.892	0.066
K	2.35 ± 0.14	3.00 ± 0.44	2.65 ± 0.54	3.94 ± 0.52	0.013	0.001	0.157
Mg	1.19 ± 0.24	0.93 ± 0.14	0.98 ± 0.16	0.94 ± 0.14	0.236	0.098	0.224
Mn	0.25 ± 0.13	0.15 ± 0.02	0.17 ± 0.06	0.14 ± 0.06	0.259	0.109	0.346
Na	0.74 ± 0.09	1.09 ± 0.61	1.00 ± 0.47	1.32 ± 0.43	0.298	0.160	0.923
P	0.23 ± 0.05	0.24 ± 0.02	0.49 ± 0.08	0.60 ± 0.06	<0.001	0.035	0.094
S	0.37 ± 0.03	0.46 ± 0.04	0.44 ± 0.17	0.55 ± 0.11	0.137	0.064	0.931
B	18.3 ± 1.80	12.7 ± 2.67	17.4 ± 2.55	12.1 ± 2.31	0.533	<0.001	0.911
Ba	58.7 ± 9.90	40.9 ± 6.24	32.8 ± 2.34	28.0 ± 5.67	<0.001	0.004	0.064
Cr	2.00 ± 1.06	3.55 ± 1.58	2.88 ± 1.56	1.50 ± 0.44	0.366	0.890	0.035
Cu	4.14 ± 0.40	5.70 ± 1.33	6.14 ± 0.74	9.01 ± 1.07	<0.001	<0.001	0.197
Li	1.22 ± 0.07	1.39 ± 0.14	1.24 ± 0.18	1.11 ± 0.14	0.086	0.778	0.044
Mo	0.29 ± 0.05	0.45 ± 0.08	0.26 ± 0.08	0.37 ± 0.02	0.125	0.001	0.416
Ni	1.14 ± 0.60	0.80 ± 0.37	0.84 ± 0.21	0.44 ± 0.17	0.096	0.067	0.869
Pb	0.58 ± 0.62	1334 ± 324	0.46 ± 0.52	2185 ± 212	0.001	<0.001	0.001
Sb	0.49 ± 0.13	0.51 ± 0.08	0.44 ± 0.04	0.51 ± 0.09	0.602	0.372	0.609
Se	0.15 ± 0.08	0.20 ± 0.13	0.13 ± 0.07	0.12 ± 0.11	0.323	0.739	0.606
Sr	96.2 ± 8.47	98.2 ± 13.8	91.3 ± 8.51	95.9 ± 10.9	0.511	0.544	0.808
V	0.48 ± 0.19	0.63 ± 0.30	0.63 ± 0.26	0.34 ± 0.11	0.548	0.540	0.076
Zn	9.05 ± 1.62	9.59 ± 1.37	12.0 ± 0.94	14.4 ± 0.92	<0.001	0.032	0.153

(Continues)

TABLE 1 (Continued)

	0mic-0Pb (ppm)	0mic-1Pb (ppm)	1mic-0Pb (ppm)	1mic-1Pb (ppm)	$\rho(\text{mic})$	$\rho(\text{Pb})$	$\rho(\text{mic} \times \text{Pb})$
Root							
Al	0.56 ± 0.41	0.66 ± 0.11	0.74 ± 0.32	0.94 ± 0.56	0.225	0.406	0.792
Ca	7.51 ± 1.26	9.09 ± 0.69	8.58 ± 0.70	10.7 ± 1.74	0.035	0.006	0.666
Fe	0.69 ± 0.43	0.77 ± 0.12	0.84 ± 0.37	0.97 ± 0.66	0.400	0.612	0.915
K	2.81 ± 0.31	3.31 ± 0.40	3.33 ± 0.47	3.96 ± 0.56	0.017	0.021	0.756
Mg	1.22 ± 0.50	1.31 ± 0.19	1.20 ± 0.15	1.49 ± 0.40	0.647	0.261	0.544
Mn	0.05 ± 0.02	0.06 ± 0.01	0.05 ± 0.01	0.05 ± 0.02	0.723	0.501	0.501
Na	0.79 ± 0.29	1.22 ± 0.41	0.86 ± 0.29	1.37 ± 0.31	0.533	0.013	0.825
P	0.23 ± 0.03	0.26 ± 0.03	0.54 ± 0.10	0.71 ± 0.14	<0.001	0.024	0.119
S	0.46 ± 0.04	0.55 ± 0.07	0.47 ± 0.08	0.62 ± 0.10	0.323	0.007	0.426
B	15.1 ± 1.26	15.3 ± 1.77	16.25 ± 2.04	12.9 ± 3.24	0.571	0.157	0.114
Ba	18.7 ± 6.40	23.1 ± 2.84	20.2 ± 1.38	29.1 ± 8.62	0.178	0.026	0.411
Cr	9.28 ± 8.66	9.00 ± 4.18	8.61 ± 4.21	10.7 ± 7.15	0.868	0.770	0.703
Cu	8.08 ± 2.07	24.2 ± 7.09	12.4 ± 1.68	24.9 ± 3.72	0.271	<0.001	0.428
Li	1.48 ± 0.40	1.61 ± 0.16	1.73 ± 0.37	2.03 ± 0.81	0.180	0.372	0.718
Mo	0.62 ± 0.13	0.62 ± 0.07	0.62 ± 0.08	0.67 ± 0.20	0.670	0.652	0.758
Ni	3.25 ± 1.81	2.87 ± 0.78	3.47 ± 1.14	3.47 ± 1.91	0.571	0.792	0.800
Pb	1.14 ± 0.66	1383 ± 1025	1.20 ± 0.38	2461 ± 1467	0.244	0.001	0.244
Sb	0.60 ± 0.15	0.58 ± 0.07	0.61 ± 0.09	0.60 ± 0.15	0.787	0.787	0.919
Se	0.10 ± 0.12	0.29 ± 0.08	0.22 ± 0.03	0.35 ± 0.17	0.126	0.014	0.643
Sr	51.2 ± 8.48	65.0 ± 6.82	61.8 ± 6.40	78.4 ± 12.5	0.015	0.003	0.754
V	2.31 ± 0.85	2.18 ± 0.06	2.63 ± 0.49	2.47 ± 1.34	0.457	0.719	0.971
Zn	18.5 ± 14.2	15.2 ± 2.04	14.0 ± 2.26	22.5 ± 9.34	0.742	0.524	0.168




FIGURE 1 Dry weights (mean and SD) of the 5 *Tuber borchii* strains, cultivated at different Pb concentration. The complete dataset is reported in Table S3.




FIGURE 2 *Tuber borchii* hyphae treated with Pb. The pink-coloured elements correspond to the lead-rhodizonate complex inside the vacuoles (A), or to oxalate crystals outside the hyphae (B).




FIGURE 3 Relative expression of *TbCdc42*, *TbRhoGdi*, *TbGlu*, *TbThio*, *TbNL* genes in *Tuber borchii* strain Tbo5118 grown on MS/2 liquid medium treated with lead compared to the controls (dotted line). Data are mean \pm SE of at least three independent experiments each performed in triplicate. The expression data were expressed as fold expression versus C. Asterisks indicate significant differences with the control (** $p < 0.01$, *** $p < 0.001$).




FIGURE 4 Difference between the percentages of mycorrhizal colonization of the control plants (0Pb, green column) and those treated with Pb (Pb1) before the first Pb treatment and 2 months after the last Pb treatment (A). *Tuber borchii* mycorrhizas untreated with Pb (0Pb) (B) and treated with Pb (1Pb) (C–G). The pink-coloured elements correspond to the lead–rhodizonate complex inside the vacuoles of extra-radical hyphae (C), to the oxalate crystals on the cystidia surface (D and F), lead–rhodizonate complex in the hyphae forming the Hartig net (E and G) and in the innermost part of the mantle (G).






FIGURE 5 Lead concentration in the pot soil (A) and BFC (B) of the plants treated (1Pb) or untreated with Pb (0Pb), mycorrhized (1mic) or not mycorrhized (0mic) with *Tuber borchii*. Bars indicate standard error. p -Values determined by two-way ANOVA were the following: p (mic) = 0.000, p (Pb) = 0.0001, p (mic \times Pb) = 0.002 for the soil lead concentration (A); p (mic) = 0.038, p (Pb) = 0.003, p (mic \times Pb) = 0.05 for the BFC (B).

Translocation factor 0mic



Translocation factor 1mic




FIGURE 6 Translocation factor (TFs) (grey bar) and TFI (black bar) values (log₁₀ scale) of the principal elements measured in not mycorrhized (0mic) and in mycorrhized plants (1mic); in ordinate symbols of element and status of plants analysed (0Pb: untreated plants, 1Pb: treated Pb plants).