



ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Perfect Strategies for Non-Local Games

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Perfect Strategies for Non-Local Games / Lupini M; Mancinska L; Paulsen VI; Roberson DE; Scarpa G; Severini S; Todorov IG; Winter A. - In: MATHEMATICAL PHYSICS ANALYSIS AND GEOMETRY. - ISSN 1385-0172. - STAMPA. - 23:1(2020), pp. 7.1-7.31. [10.1007/s11040-020-9331-7]

This version is available at: <https://hdl.handle.net/11585/914658> since: 2023-02-10

Published:

DOI: <http://doi.org/10.1007/s11040-020-9331-7>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

This is the final peer-reviewed accepted manuscript of:

Lupini, M., Mančinska, L., Paulsen, V.I. et al. Perfect Strategies for Non-Local Games. *Math Phys Anal Geom* 23, 7 (2020)

The final published version is available online at <https://doi.org/10.1007/s11040-020-9331-7>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

PERFECT STRATEGIES FOR NON-LOCAL GAMES

M. LUPINI, L. MANČINSKA, V. I. PAULSEN, D. E. ROBERSON, G. SCARPA,
S. SEVERINI, I. G. TODOROV, AND A. WINTER

ABSTRACT. We introduce a new class of non-local games, called imitation games, in which the players display linked behaviour, and which contain as subclasses the classes of variable assignment games, binary constraint system games, synchronous games, many games based on graphs, and unique games. We associate a C*-algebra $C^*(\mathcal{G})$ to any imitation game \mathcal{G} , and show that the existence of perfect quantum commuting (resp. quantum, local) strategies of \mathcal{G} can be characterised in terms of properties of this C*-algebra. We single out a subclass of imitation games, which we call mirror games, and provide a characterisation of their quantum commuting strategies that has an algebraic flavour, showing in addition that their approximately quantum perfect strategies arise from amenable traces on the encoding C*-algebra. We describe the main classes of non-signalling bipartite correlations in terms of states on operator system tensor products. This leads to the introduction of another new class of games, called reflexive games, which are characterised as the hardest non-local games that can be won using a given set of strategies. We provide a characterisation of their perfect strategies in terms of operator system quotients.

1. INTRODUCTION

The study of correlated behaviour between two parties obeying the laws of Quantum Mechanics has been central in the subject since its initiation. The celebrated Bell Theorem demonstrates that the set \mathcal{C}_q of quantum correlations, arising from shared finite dimensional entanglement, is strictly larger than the set of classical (or local) correlations, arising from shared randomness. A third natural class of correlations, \mathcal{C}_{qc} , arising from Algebraic Quantum Field Theory, corresponds to the *commuting model* of Quantum Mechanics. According to it, the measurements of the two parties act on a single (infinite dimensional) Hilbert space; this setting was studied in [19] and subsequently in [29], where the author showed that every correlation from \mathcal{C}_{qc} can be approximated by ones from \mathcal{C}_q if and only if the Connes Embedding Problem in operator algebra theory [7] has an affirmative answer. Deep results about the inequality between those classes of correlations, answering questions left open by Tsirelson (see [40] and [41]) were recently obtained by Slofstra in [37] and [38] when the number of inputs is large and in [11] similar results are shown for a small number of inputs. The relevance

Date: 7 November 2018.

of operator algebraic techniques in the study of correlation sets became also apparent through [13] and, subsequently, [33], where operator systems and their tensor products were used to describe some correlation classes.

Non-local, or two-player one-round, games have attracted substantial attention in theoretical physics, mathematics and computer science (see e.g. [26], [36], [30] and [39]). For some games, quantum or other non-signalling strategies, have been successfully used to obtain strategies for such games that outperform the classical ones. A prominent such example is the graph colouring game defined in [3], where it was demonstrated that the quantum chromatic number of a graph, arising from the set \mathcal{C}_q , can be strictly smaller than its classical counterpart. In [33], the authors defined a corresponding commuting version of the chromatic number, using the set \mathcal{C}_{qc} of quantum commuting correlations, in addition to other related analogues. The graph homomorphism game was introduced in [25, 27] and subsequently studied [28] as a generalisation of the graph colouring game. An even more general class – that of synchronous games – was considered in [32], where perfect strategies from the classes \mathcal{C}_q and \mathcal{C}_{qc} were described in terms of traces on a canonical C^* -algebra, associated with the input-output sets of the game.

Another prominent class of games of importance in Quantum Information Theory, that was at the heart of Slofstra’s work [37], is the class of binary constraint system (BCS) games [6]. In the present paper, we observe that there is a unifying feature of all aforementioned non-local games, which leads us to the introduction and study of the class of *imitation games*. In addition to the games discussed previously, it includes the class of variable assignment games and the class of unique games [36]. One of our main objectives is to develop the formalism of these games and exhibit the interconnections between its various subclasses.

The paper is organised as follows. After collecting some necessary preliminary material in Section 2, we describe, in Section 3, the perfect non-signalling strategies of a general non-local game that belong to a given class, in terms of states of operator system tensor products. These descriptions have two important advantages as compared to the existing use of C^* -algebras, encountered more commonly in the literature. Firstly, the operator systems involved are finite dimensional and hence easier to handle than their infinite dimensional C^* -algebraic counterparts. Secondly, the tensor theory of operator systems is genuinely richer than that of C^* -algebras [21], and this allows to express, via operator system tensor products, correlations pertinent to generalised probabilistic theories as opposed to ones appearing in Quantum Mechanics alone (for a background on generalised probabilistic theories, we refer the reader to [24]).

We further define the new class of *reflexive games*; a reflexive game can be thought of as the hardest game that can be won using a family of strategies from a given class. We show that the perfect strategies of reflexive games are in a one-to-one correspondence to states on certain operator system quotients. In fact, we introduce, more generally, *reflexive covers* of games,

and exhibit several examples where the reflexive cover of a game can be strictly harder than the original game.

In Sections 4 and 5, we study the class of imitation games. With every imitation game \mathcal{G} , we associate a canonical C^* -algebra $C^*(\mathcal{G})$, and provide an explicit description of $C^*(\mathcal{G})$ in the case \mathcal{G} is a variable assignment game. If \mathcal{G} is a linear BCS game, we relate $C^*(\mathcal{G})$ to the group C^* -algebra of the group of \mathcal{G} that was introduced in [5] and used extensively in [37]. We show that the perfect quantum commuting strategies of an imitation game \mathcal{G} correspond to traces on $C^*(\mathcal{G})$, while the perfect quantum strategies of \mathcal{G} correspond to finite dimensional representations of $C^*(\mathcal{G})$. We prove the equality of several classes of correlations, namely the quantum ones, the quantum spatial ones and the maximally entangled quantum ones.

In Section 6, we consider a subclass of the class of imitation games, which we call *mirror games*, and provide an algebraic, Hilbert-space free, approach, to their perfect strategies. As a consequence, we give a different representation of their perfect quantum commuting strategies, using traces on canonical input-output C^* -algebras. We show that the *quantum approximate* perfect strategies of these games correspond precisely to amenable traces on these C^* -algebras, extending a recent result from [23].

2. PRELIMINARIES

We fix finite sets X, Y, A and B . A collection of scalars

$$p = \{p(a, b|x, y) : (x, y) \in X \times Y, (a, b) \in A \times B\}$$

is called *non-signalling* if

$$(1) \quad \sum_{b \in B} p(a, b|x, y) = \sum_{b \in B} p(a, b|x, y'), \quad x \in X, y, y' \in Y, a \in A$$

and

$$(2) \quad \sum_{a \in A} p(a, b|x, y) = \sum_{a \in A} p(a, b|x', y), \quad x, x' \in X, y \in Y, b \in B.$$

If, in addition, $p(\cdot, \cdot|x, y)$ is a probability distribution for every $(x, y) \in X \times Y$, then p is called a *non-signalling correlation* on (X, Y, A, B) . The set \mathcal{C}_{ns} of all non-signalling correlations is canonically endowed with a compact metrisable topology and a convex structure by regarding it as a subset of \mathbb{R}^m where m is the cardinality of $X \times Y \times A \times B$. If $p \in \mathcal{C}_{\text{ns}}$, we set $p(a|x)$ (resp. $p(b|y)$) to be equal to the sum in (1) (resp. (2)), for any choice of $y \in Y$ (resp. $x \in X$).

We recall the definition of several sets of non-signalling correlations. The set \mathcal{C}_{det} of *deterministic* correlations consists of all correlations p for which there exist functions $f : X \rightarrow A$ and $g : Y \rightarrow B$ such that $p(a, b|x, y) = 1$ if and only if $a = f(x)$ and $b = g(y)$.

The set \mathcal{C}_{loc} of *local correlations* is the convex hull of correlations of the form $p(a, b|x, y) = p^1(a|x)p^2(b|y)$ where $p^1 : A \times X \rightarrow [0, 1]$ and $p^2 :$

$B \times Y \rightarrow [0, 1]$ satisfy $\sum_a p^1(a|x) = \sum_b p^2(b|y) = 1$ for every $x \in X$ and every $y \in Y$.

In order to define the rest of the classes, known as non-classical, we recall that a *positive operator-valued measure (POVM)* on a Hilbert space \mathcal{H} is a tuple (A_1, \dots, A_n) of positive operators on \mathcal{H} summing up to the identity operator. A *projection-valued measure (PVM)* is a POVM consisting of projection operators.

In each of the definitions given below, we use PVM's, but results of [16] (see also [33]) show that the sets of correlations that we obtain are the same if we replace PVM's with POVM's in each definition.

The set \mathcal{C}_q of *quantum correlations* consists of the correlations of the form

$$p(a, b|x, y) = \langle \xi | P_{x,a} \otimes Q_{b,y} | \xi \rangle$$

where \mathcal{H} is a *finite-dimensional* Hilbert space, $|\xi\rangle \in \mathcal{H} \otimes \mathcal{H}$ is a unit vector, and for every $x \in X$ and $y \in Y$, $(P_{x,a})_{a \in A}$ and $(Q_{y,b})_{b \in B}$ are PVM's on \mathcal{H} .

Suppose that \mathcal{H} is a finite-dimensional Hilbert space with fixed basis $|i\rangle$, $i = 1, 2, \dots, d$. The *maximally entangled vector* in $\mathcal{H} \otimes \mathcal{H}$ is the unit vector $|\eta\rangle = d^{-1/2} \sum_{i=1}^d |i\rangle \otimes |i\rangle$. The set \mathcal{C}_{qm} is the *convex hull* of the quantum correlations of the form

$$p(a, b|x, y) = \langle \eta | P_{x,a} \otimes Q_{b,y} | \eta \rangle,$$

where $(P_{x,a})_{a \in A}$ (resp. $(Q_{y,b})_{b \in B}$) is a PVM on \mathcal{H} for every $x \in X$ (resp. $y \in Y$).

The set \mathcal{C}_{qs} of *quantum spatial* correlations is defined similarly to the set of quantum correlations, but the restriction that \mathcal{H} be finite dimensional is dropped. The set \mathcal{C}_{qa} of *quantum approximate* correlations is defined to be the closure of \mathcal{C}_q . Finally, the set \mathcal{C}_{qc} of *quantum commuting* correlations consists of the correlations of the form

$$p(a, b|x, y) = \langle \xi | P_{x,a} Q_{y,b} | \xi \rangle,$$

where \mathcal{H} is a Hilbert space, $|\xi\rangle \in \mathcal{H}$ is a unit vector and $(P_{x,a})_{a \in A}$ and $(Q_{y,b})_{b \in B}$, $x \in X$, $y \in Y$ are PVMs on \mathcal{H} such that $P_{x,a} Q_{y,b} = Q_{y,b} P_{x,a}$ for all $x \in X$, $y \in Y$, $a \in A$ and $b \in B$. A separability argument shows that considering *separable* Hilbert spaces in the definition of the sets \mathcal{C}_{qs} and \mathcal{C}_{qc} yields equivalent definitions. It is clear from the definition (after observing that \mathcal{C}_{qc} is closed) that we have the following inclusions between these sets of correlations

$$\mathcal{C}_{\det} \subseteq \mathcal{C}_{\text{loc}} \subseteq \mathcal{C}_{qm} \subseteq \mathcal{C}_q \subseteq \mathcal{C}_{qs} \subseteq \mathcal{C}_{qa} \subseteq \mathcal{C}_{qc} \subseteq \mathcal{C}_{\text{ns}}.$$

We now recall the connection between non-signalling correlations and perfect strategies for non-local games. A *non-local game* is a tuple $\mathcal{G} = (X, Y, A, B, \lambda_{\mathcal{G}})$, where X, Y, A and B are finite sets and $\lambda_{\mathcal{G}} : X \times Y \times A \times B \rightarrow \{0, 1\}$ is a function. We think of X and Y as sets of possible *inputs* or *questions* for two players (Alice and Bob) of a two-party single-round game, and of A and B as sets of possible *outputs* or *answers* for Alice and Bob, respectively. The function $\lambda_{\mathcal{G}}$ is called the *payoff*, or *rule, function* of \mathcal{G} ,

assigning value 1 to (x, y, a, b) if a, b are *acceptable answers* to the pair (x, y) of questions, and 0 otherwise. When there is no risk of confusion, we write $\lambda = \lambda_{\mathcal{G}}$. Notice that in this subclass of non-local games, we do not consider probability distributions on the input sets and we restrict our attention to two players. Correspondingly, we are interested in the perfect strategies for these games, which are automatically winning strategies for any given probability distribution on the direct product of the input sets of the game. More precisely, we call a non-signalling correlation p on (X, Y, A, B) a *perfect strategy* for \mathcal{G} if

$$\lambda(x, y, a, b) = 0 \implies p(a, b|x, y) = 0.$$

We let $\mathcal{C}_{\text{ns}}(\mathcal{G})$ (or $\mathcal{C}_{\text{ns}}(\lambda)$) be the set of all such correlations.

More specifically, if $\lambda : X \times Y \times A \times B \rightarrow \{0, 1\}$ and $p \in \mathcal{C}_{\text{ns}}$, write

$$N(\lambda) = \{(x, y, a, b) \in X \times Y \times A \times B : \lambda(x, y, a, b) = 0\}$$

and

$$N(p) = \{(x, y, a, b) \in X \times Y \times A \times B : p(a, b|x, y) = 0\}.$$

Setting

$$\mathcal{C}_x(\lambda) := \{p \in \mathcal{C}_x : N(\lambda) \subseteq N(p)\} = \mathcal{C}_x \cap \mathcal{C}_{\text{ns}}(\mathcal{G}),$$

we obtain a corresponding chain

$$\mathcal{C}_{\text{det}}(\lambda) \subseteq \mathcal{C}_{\text{loc}}(\lambda) \subseteq \mathcal{C}_{\text{qm}}(\lambda) \subseteq \mathcal{C}_{\text{q}}(\lambda) \subseteq \mathcal{C}_{\text{qs}}(\lambda) \subseteq \mathcal{C}_{\text{qa}}(\lambda) \subseteq \mathcal{C}_{\text{qc}}(\lambda) \subseteq \mathcal{C}_{\text{ns}}(\lambda).$$

A *perfect x-strategy* for the game \mathcal{G} is an element of $\mathcal{C}_x(\lambda)$.

In this paper, we will arrive at characterisations of such sets of correlations in terms of states on operator systems. We refer the reader to [31] for an introduction to the basic notions of non-commutative functional analysis (see also [12, 34]). Let \mathcal{S} be an operator system, that is, a subspace of a unital C^* -algebra \mathcal{A} such that $1 \in \mathcal{S}$ and $x \in \mathcal{S} \implies x^* \in \mathcal{S}$. Then $M_n(\mathcal{S}) \subseteq M_n(\mathcal{A})$, and we let $M_n(\mathcal{S})^+$ be the cone of all elements of $M_n(\mathcal{S})$ that are positive in the C^* -algebra $M_n(\mathcal{A})$.

Let \mathcal{S} and \mathcal{T} be operator systems. Given a linear map $\phi : \mathcal{S} \rightarrow \mathcal{T}$, let $\phi^{(n)} : M_n(\mathcal{S}) \rightarrow M_n(\mathcal{T})$ be the map given by $\phi^{(n)}((x_{i,j})_{i,j}) = (\phi(x_{i,j}))_{i,j}$. The map ϕ is called *positive* if $\phi(\mathcal{S}^+) \subseteq \mathcal{T}^+$, and *completely positive* if $\phi^{(n)}(M_n(\mathcal{S})^+) \subseteq M_n(\mathcal{T})^+$ for every $n \in \mathbb{N}$. A state on \mathcal{S} is a positive linear map $s : \mathcal{S} \rightarrow \mathbb{C}$ with $s(1) = 1$.

We say that two operator systems are completely order isomorphic if there exists a bijective completely positive map with completely positive inverse between them. We write $\mathcal{S} \subseteq_{\text{c.o.i.}} \mathcal{T}$ when, as linear spaces, $\mathcal{S} \subseteq \mathcal{T}$, and $M_n(\mathcal{S})^+ = M_n(\mathcal{T})^+ \cap M_n(\mathcal{S})$ for each $n \in \mathbb{N}$. We denote by $\mathcal{S} \oplus^1 \mathcal{T}$ the coproduct of \mathcal{S} and \mathcal{T} in the category of operator systems and unital completely positive maps; it is characterised by the following universal property: $\mathcal{S} \oplus^1 \mathcal{T}$ is generated as a linear space by \mathcal{S} and \mathcal{T} , its unit is also the unit of \mathcal{S} and of \mathcal{T} , and whenever \mathcal{R} is an operator system and $\phi : \mathcal{S} \rightarrow \mathcal{R}$ and $\psi : \mathcal{T} \rightarrow \mathcal{R}$ are unital completely positive maps then there exists a unique (unital) completely positive map $\theta : \mathcal{S} \oplus^1 \mathcal{T} \rightarrow \mathcal{R}$ extending ϕ and ψ . We

refer the reader to [16, Section 3] and [20, Section 8] for further properties of the operator system coproduct. We further let $\mathcal{S} \oplus^\infty \mathcal{T}$ be the product of \mathcal{S} and \mathcal{T} and, for a finite set A , define $\ell^\infty(A)$ to be the direct sum of $|A|$ copies of the operator system \mathbb{C} , indexed by A .

In the sequel, we will make use of the tensor theory of operator systems developed in [21]. If \mathcal{S} and \mathcal{T} are operator systems, we denote by $\mathcal{S} \otimes_{\min} \mathcal{T}$ (resp. $\mathcal{S} \otimes_{\mathbb{C}} \mathcal{T}$, $\mathcal{S} \otimes_{\max} \mathcal{T}$) the minimal (resp. commuting, maximal) tensor product of \mathcal{S} and \mathcal{T} introduced therein. We note that, if $\mathcal{S} \subseteq_{\text{c.o.i.}} \mathcal{A}$ and $\mathcal{T} \subseteq_{\text{c.o.i.}} \mathcal{B}$ for C^* -algebras \mathcal{A} and \mathcal{B} , then $\mathcal{S} \otimes_{\min} \mathcal{T} \subseteq_{\text{c.o.i.}} \mathcal{A} \otimes_{\min} \mathcal{B}$, where $\mathcal{A} \otimes_{\min} \mathcal{B}$ is the spatial tensor product of \mathcal{A} and \mathcal{B} . By their definition, the tensor product $\mathcal{S} \otimes_{\mathbb{C}} \mathcal{T}$ linearises pairs of unital completely positive maps $\phi : \mathcal{S} \rightarrow \mathcal{A}$ and $\psi : \mathcal{T} \rightarrow \mathcal{A}$ with commuting ranges, while the maximal tensor product $\mathcal{S} \otimes_{\max} \mathcal{T}$ linearises jointly completely positive maps $\theta : \mathcal{S} \times \mathcal{T} \rightarrow \mathcal{A}$ (here \mathcal{A} is an arbitrary C^* -algebra).

3. CORRELATIONS AS PERFECT STRATEGIES

Let X , Y , A and B be finite sets. Following [33], we let $\mathcal{S}_{X,A}$ be the coproduct of $|X|$ copies of $\ell^\infty(A)$, indexed by X . Let $\mathcal{A}(X, A) = \ell^\infty(A) *_1 \cdots *_1 \ell^\infty(A)$ be the C^* -algebra free product, amalgamated over the unit, of $|X|$ copies of $\ell^\infty(A)$, and $\mathbb{F}(X, A) = \mathbb{Z}_{|A|} * \cdots * \mathbb{Z}_{|A|}$ be the free product of $|X|$ copies of the cyclic group with $|A|$ elements. The Fourier transform implements a $*$ -isomorphism between the C^* -algebra $C^*(\mathbb{Z}_{|A|})$ of the group $\mathbb{Z}_{|A|}$ and $\ell^\infty(A)$; thus, we have a canonical $*$ -isomorphism $\mathcal{A}(X, A) \cong C^*(\mathbb{F}(X, A))$. Letting $(e_{x,a})_{a=1}^{|A|}$ be the canonical basis of the x -th copy of $\ell^\infty(A)$, we have that

$$\mathcal{S}_{X,A} = \text{span}\{e_{x,a} : x \in X, a \in A\}$$

within $\mathcal{A}(X, A)$. Note the relations

$$\sum_{a \in A} e_{x,a} = 1, \quad x \in X.$$

Set $\mathcal{A}_{\min}(X, A) = \ell^\infty(A) \otimes \cdots \otimes \ell^\infty(A)$ ($|X|$ copies) and note that the C^* -algebra $\mathcal{A}_{\min}(X, A)$ is $*$ -isomorphic to $\ell^\infty(\Delta_{X,A})$, where $\Delta_{X,A} = A^X$. Let

$$\mathcal{S}_{X,A}^{\min} = \text{span}\{e'_{x,a} : x \in X, a \in A\} \subseteq \mathcal{A}_{\min}(X, A),$$

where $e'_{x,a}(x', a') = 0$ if $x = x'$ and $a \neq a'$, and $e'_{x,a}(x', a') = 1$ otherwise.

For conceptual convenience, we will denote the canonical generators of $\mathcal{S}_{Y,B}$ (resp. $\mathcal{S}_{Y,B}^{\min}$) by $f_{y,b}$ (resp. $f'_{y,b}$). For an element s of the dual vector space $(\mathcal{S}_{X,A} \otimes \mathcal{S}_{Y,B})^d$ of $\mathcal{S}_{X,A} \otimes \mathcal{S}_{Y,B}$, write

$$p_s(a, b|x, y) = s(e_{x,a} \otimes f_{y,b}), \quad (x, y) \in X \times Y, (a, b) \in A \times B,$$

and

$$p_s = \{p_s(a, b|x, y) : (x, y) \in X \times Y, (a, b) \in A \times B\}.$$

Clearly, the collection p_s is non-signalling. Conversely, given a non-signalling collection of scalars p , let $s_p \in (\mathcal{S}_{X,A} \otimes \mathcal{S}_{Y,B})^d$ be the (well-defined and unique linear) functional satisfying

$$s_p(e_{x,a} \otimes f_{y,b}) = p(a, b|x, y), \quad (x, y) \in X \times Y, (a, b) \in A \times B.$$

It is clear that $p \rightarrow s_p$ is a bijective correspondence between $(\mathcal{S}_{X,A} \otimes \mathcal{S}_{Y,B})^d$ and the set of all non-signalling collections on (X, Y, A, B) .

Let $\lambda : X \times Y \times A \times B \rightarrow \{0, 1\}$. Set

$$J(\lambda) = \text{span}\{e_{x,a} \otimes f_{y,b} : \lambda(x, y, a, b) = 0\};$$

thus, $J(\lambda)$ is a linear subspace of $\mathcal{S}_{X,A} \otimes \mathcal{S}_{Y,B}$.

If τ is any of the tensor products max, c or min and $J \subseteq \mathcal{S}_{X,A} \otimes \mathcal{S}_{Y,B}$, let

$$\mathcal{P}_\tau(J) = \{s \in (\mathcal{S}_{X,A} \otimes_\tau \mathcal{S}_{Y,B})^d : s \text{ is a state with } J \subseteq \ker(s)\}.$$

Let also

$$\mathcal{P}_{\text{omin}}(J) = \{s \in (\mathcal{S}_{X,A}^{\text{min}} \otimes_{\text{min}} \mathcal{S}_{Y,B}^{\text{min}})^d : s \text{ is a state with } J \subseteq \ker(s)\}.$$

We write $\mathcal{P}_\tau = \mathcal{P}_\tau(\{0\})$.

Theorem 3.1. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a non-local game and*

$$p = \{(p(a, b|x, y)) : (x, y) \in X \times Y, (a, b) \in A \times B\}$$

be a non-signalling collection of scalars. The map $p \rightarrow s_p$ is a continuous affine isomorphism between

- (i) $\mathcal{C}_{\text{ns}}(\lambda)$ and $\mathcal{P}_{\text{max}}(J(\lambda))$;
- (ii) $\mathcal{C}_{\text{qc}}(\lambda)$ and $\mathcal{P}_{\text{c}}(J(\lambda))$;
- (iii) $\mathcal{C}_{\text{qa}}(\lambda)$ and $\mathcal{P}_{\text{min}}(J(\lambda))$;
- (iv) $\mathcal{C}_{\text{loc}}(\lambda)$ and $\mathcal{P}_{\text{omin}}(J(\lambda))$,

and a bijection between

- (v) $\mathcal{C}_{\text{det}}(\lambda)$ and the extreme points of the set $\mathcal{P}_{\text{omin}}(J(\lambda))$.

Proof. (i) Set

$$\mathcal{R}_{X,A} = \left\{ (z_{x,a})_{x \in X, a \in A} : \sum_{a \in A} z_{x,a} = \sum_{a \in A} z_{x',a}, \text{ for all } x, x' \in X \right\},$$

viewed as an operator subsystem of $\ell^\infty(X \times A)$. By [14, Theorem 5.9], the dual operator system $\mathcal{R}_{X,A}^d$ of $\mathcal{R}_{X,A}$, which is again an operator system by finite dimensionality [4, Theorem 4.4], is completely order isomorphic to $\mathcal{S}_{X,A}$, the duality being given by

$$\langle (z_{x,a}), e_{x',a'} \rangle = z_{x',a'}.$$

By [22, Proposition 6.2] and the fact that $\mathcal{S}_{X,A}$ is finite-dimensional, $\mathcal{S}_{X,A}^d \cong \mathcal{R}_{X,A}$. By [15, Proposition 1.9],

$$(\mathcal{S}_{X,A} \otimes_{\text{max}} \mathcal{S}_{Y,B})^d \cong \mathcal{R}_{X,A} \otimes_{\text{min}} \mathcal{R}_{X,A}.$$

By the injectivity of the minimal tensor product, $\mathcal{R}_{X,A} \otimes_{\text{min}} \mathcal{R}_{X,A}$ is (completely order isomorphic to) an operator subsystem of $\ell^\infty(X \times A \times Y \times$

B), and it is straightforward to check that the image of the state space of $\mathcal{S}_{X,A} \otimes_{\max} \mathcal{S}_{Y,B}$ under this isomorphism is precisely the set of all non-signalling correlations. Statement (i) is now clear.

(ii) By [33, Lemma 2.6], $\mathcal{S}_{X,A} \otimes_c \mathcal{S}_{Y,B} \subseteq_{\text{c.o.i.}} \mathcal{A}(X,A) \otimes_{\max} \mathcal{A}(Y,B)$ and hence every state on $\mathcal{S}_{X,A} \otimes_c \mathcal{S}_{Y,B}$ extends to a state on $\mathcal{A}(X,A) \otimes_{\max} \mathcal{A}(Y,B)$. The proof now follows the arguments in [33, Theorem 2.8].

(iii) Here the proof follows the one of [33, Theorem 2.9], using the fact that $\mathcal{S}_{X,A} \otimes_{\min} \mathcal{S}_{Y,B} \subseteq_{\text{c.o.i.}} \mathcal{A}(X,A) \otimes_{\min} \mathcal{A}(Y,B)$.

(iv) Note that

$$(3) \quad \mathcal{S}_{X,A}^{\min} \otimes_{\min} \mathcal{S}_{Y,B}^{\min} \subseteq_{\text{c.o.i.}} \ell^\infty(\Delta_{X,A} \times \Delta_{Y,B}).$$

Suppose that $p_x^1 = (p^1(a|x))_{a \in A}$ is a probability distribution on A ; then p_x^1 gives rise to a state s_x^1 on $\ell^\infty(A)$, $x \in X$. The product state $s^1 = \otimes_{x \in X} s_x^1$ on $\ell^\infty(X \times A)$ is given by $s_1 \left(\sum_{x,a} \lambda_{x,a} e'_{x,a} \right) = \sum_{x,a} p^1(a|x) \lambda_{x,a}$. If, similarly, $p_y^2 = (p^2(b|y))_{b \in B}$ is a probability distribution on B , $y \in Y$, and s^2 is the associated state on $\ell^\infty(Y \times B)$, then the product state $s^1 \otimes s^2$ on $\ell^\infty(X \times A \times Y \times B)$ arises from the families of product distributions $p_x^1 \otimes p_y^2$, $x \in X$, $y \in Y$. It now follows that, if $p \in \mathcal{C}_{\text{loc}}$ then s_p is a state on $\mathcal{S}_{X,A}^{\min} \otimes_{\min} \mathcal{S}_{Y,B}^{\min}$.

Conversely, suppose that s is a state on $\mathcal{S}_{X,A}^{\min} \otimes_{\min} \mathcal{S}_{Y,B}^{\min}$. In view of (3), s has an extension, which we denote in the same way, to a state on $\ell^\infty(\Delta_{X,A} \times \Delta_{Y,B})$. It is thus a convex combination of pure states. On the other hand, if s_0 is a pure state of $\ell^\infty(\Delta_{X,A} \times \Delta_{Y,B})$ then there exists $(t_1, t_2) \in \Delta_{X,A} \times \Delta_{Y,B}$ such that $s_0(f) = f(t_1, t_2)$, $f \in \ell^\infty(\Delta_{X,A} \times \Delta_{Y,B})$. Thus, $s_0 = s_1 \otimes s_2$, where s_1 (resp. s_2) is the state on $\ell^\infty(\Delta_{X,A})$ (resp. $\ell^\infty(\Delta_{Y,B})$) of evaluation on t_1 (resp. t_2), and it follows that p_s is a local correlation. Statement (iv) is now immediate.

(v) Suppose that $p \in \mathcal{C}_{\text{det}}(\lambda)$. Clearly, p is an extreme point of $\mathcal{C}_{\text{loc}}(\lambda)$; by (iv), s_p is an extreme point of $\mathcal{P}_{\text{omin}}(J(\lambda))$. Conversely, suppose that s_p is an extreme point of $\mathcal{P}_{\text{omin}}(J(\lambda))$. By (iv), p is an extreme point of $\mathcal{C}_{\text{loc}}(\lambda)$. Suppose that $p = tp_1 + (1-t)p_2$, where $0 \leq t \leq 1$ and $p_1, p_2 \in \mathcal{C}_{\text{loc}}$. It follows that $p_1(a, b|x, y) = p_2(a, b|x, y) = 0$ whenever $\lambda(x, y, a, b) = 0$, and hence $p = p_1 = p_2$. Thus, p is an extreme point of \mathcal{C}_{loc} and therefore belongs to \mathcal{C}_{det} . Thus, $p \in \mathcal{C}_{\text{det}}(\lambda)$. \square

We record, in the following two statements, some additional descriptions of the sets of quantum commuting and quantum approximate correlations that will be used in the sequel.

Corollary 3.2. *Let p be a non-signalling correlation on $X \times Y \times A \times B$. The following statements are equivalent:*

- (i) $p \in \mathcal{C}_{\text{qc}}$;
- (ii) $p(a, b|x, y) = s(e_{x,a} \otimes f_{y,b})$ for some state s on $\mathcal{A}(X,A) \otimes_{\max} \mathcal{A}(Y,B)$;
- (iii) $p(a, b|x, y) = s(e_{x,a} \otimes f_{y,b})$ for some state s on $\mathcal{S}_{X,A} \otimes_c \mathcal{S}_{Y,B}$.

Proof. (i) \Rightarrow (iii) follows from Theorem 3.1 (ii).

(iii) \Rightarrow (ii) follows from Krein's Theorem and the fact that $\mathcal{S}_{X,A} \otimes_c \mathcal{S}_{Y,B} \subseteq_{\text{c.o.i.}} \mathcal{A}(X, A) \otimes_{\max} \mathcal{A}(Y, B)$ [33, Lemma 2.7].

(ii) \Rightarrow (i) is similar to the arguments in the proof of [33, Lemma 2.7]. \square

Corollary 3.3. *Let p be a non-signalling correlation on $X \times Y \times A \times B$. The following statements are equivalent:*

- (i) $p \in \mathcal{C}_{\text{qa}}$;
- (ii) $p(a, b|x, y) = s(e_{x,a} \otimes f_{y,b})$ for some state s on $\mathcal{A}(X, A) \otimes_{\min} \mathcal{A}(Y, B)$;
- (iii) $p(a, b|x, y) = s(e_{x,a} \otimes f_{y,b})$ for some state s on $\mathcal{S}_{X,A} \otimes_{\min} \mathcal{S}_{Y,B}$.

Proof. (i) \Rightarrow (iii) follows from Theorem 3.1 (iii).

(iii) \Rightarrow (ii) follows from Krein's Theorem and the fact that, by the injectivity of the minimal tensor product, $\mathcal{S}_{X,A} \otimes_{\min} \mathcal{S}_{Y,B} \subseteq_{\text{c.o.i.}} \mathcal{A}(X, A) \otimes_{\min} \mathcal{A}(Y, B)$.

(ii) \Rightarrow (i) follows by the arguments in the proof of [33, Theorem 2.9]. \square

In this rest of this section, we give a more precise description of perfect strategies for a special class of games, which we now introduce. If $\mathcal{G}_1 = (X, Y, A, B, \lambda_1)$ and $\mathcal{G}_2 = (X, Y, A, B, \lambda_2)$ are games, we write $\mathcal{G}_1 \leq \mathcal{G}_2$ if $\lambda_1 \leq \lambda_2$, and in this case say that \mathcal{G}_1 is *harder* (or *smaller*) than \mathcal{G}_2 .

Let $x \in \{\text{det}, \text{loc}, \text{qm}, \text{q}, \text{qs}, \text{qa}, \text{qc}, \text{ns}\}$. For $\Sigma \subseteq \mathcal{C}_x$, let $\lambda_\Sigma : X \times Y \times A \times B \rightarrow \{0, 1\}$ be the function defined by the equality

$$N(\lambda_\Sigma) = \bigcap_{p \in \Sigma} N(p).$$

Clearly, λ_Σ is the rule function of the hardest game for which every element of Σ is a perfect strategy.

Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a game. We set $\lambda_x = \lambda_{\mathcal{C}_x(\lambda)}$. Thus,

$$\lambda_x(x, y, a, b) = 0 \iff p(a, b|x, y) = 0 \text{ for every } p \in \mathcal{C}_x(\lambda).$$

Set $\text{Ref}_x(\mathcal{G}) = (X, Y, A, B, \lambda_x)$ and call it the *reflexive x-cover* of \mathcal{G} . We call \mathcal{G} *x-reflexive* if $\text{Ref}_x(\mathcal{G}) = \mathcal{G}$.

Note the inequalities

$$\lambda_{\text{det}} \leq \lambda_{\text{loc}} \leq \lambda_{\text{qm}} \leq \lambda_{\text{q}} \leq \lambda_{\text{qs}} \leq \lambda_{\text{qa}} \leq \lambda_{\text{qc}} \leq \lambda_{\text{ns}} \leq \lambda.$$

Example 3.4. Consider the graph colouring game for the graph G with vertex set $X = \{1, 2, 3, 4\}$ and edge set $\{(1, 2), (2, 3), (3, 4)\}$. Then every deterministic 2-colouring of G is also a deterministic 2-colouring of the 4-cycle. This shows that the reflexive covers of a game can be strictly harder than the original game.

Proposition 3.5. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a game and $x \in \{\text{det}, \text{loc}, \text{qm}, \text{q}, \text{qs}, \text{qa}, \text{qc}, \text{ns}\}$. The following hold:*

- (i) $\text{Ref}_x(\text{Ref}_x(\mathcal{G})) = \text{Ref}_x(\mathcal{G})$.
- (ii) \mathcal{G} is *x-reflexive* if and only if there exists a set $\Sigma \subseteq \mathcal{C}_x$ such that $\lambda = \lambda_\Sigma$.

Proof. (i) Clearly, $N(\lambda) \subseteq N(\lambda_x)$, and so $\mathcal{C}_x(\lambda_x) \subseteq \mathcal{C}_x(\lambda)$. Suppose that $p \in \mathcal{C}_x(\lambda)$ and $(x, y, a, b) \in N(\lambda_x)$. By the definition of λ_x , we have that $p(a, b|x, y) = 0$. Thus, $p \in \mathcal{C}_x(\lambda_x)$ and so $\mathcal{C}_x(\lambda) = \mathcal{C}_x(\lambda_x)$. It now follows that $\lambda_x = \lambda_{xx}$, that is, $\text{Ref}_x(\mathcal{G}) = \text{Ref}_x(\text{Ref}_x(\mathcal{G}))$.

(ii) If \mathcal{G} is reflexive then we can take $\Sigma = \mathcal{C}_x(\lambda)$. Conversely, suppose that $\lambda = \lambda_\Sigma$ for some $\Sigma \subseteq \mathcal{C}_x$. The inclusion $N(\lambda) \subseteq N(\lambda_x)$ follows by the definition of λ_x . On the other hand, since $\Sigma \subseteq \mathcal{C}_x(\lambda)$, we have

$$\cap\{N(p) : p \in \mathcal{C}_x(\lambda)\} \subseteq \cap\{N(p) : p \in \Sigma\},$$

that is, $N(\lambda_x) \subseteq N(\lambda)$. Thus, $N(\lambda_x) = N(\lambda)$ and \mathcal{G} is x -reflexive. \square

A *kernel* in an operator system \mathcal{S} [22] is a subspace $J \subseteq \mathcal{S}$ for which there exist an operator system \mathcal{T} and a completely positive map $\phi : \mathcal{S} \rightarrow \mathcal{T}$ such that $J = \ker(\phi)$. If $J \subseteq \mathcal{S}$ is a kernel, then the quotient linear space \mathcal{S}/J can be equipped with a (unique) operator system structure with the property that, whenever \mathcal{T} is an operator system and $\phi : \mathcal{S} \rightarrow \mathcal{T}$ is a completely positive map with $J \subseteq \ker(\phi)$, the induced map $\hat{\phi} : \mathcal{S}/J \rightarrow \mathcal{T}$ is completely positive. For a fixed game $\mathcal{G} = (X, Y, A, B, \lambda)$ and $\tau \in \{\max, c, \min, \text{omin}\}$, write

$$J_\tau(\lambda) = \bigcap \{\ker(s) : s \in \mathcal{P}_\tau(J(\lambda))\}.$$

By [22, Proposition 3.1], $J_\tau(\lambda)$ is a kernel in $\mathcal{S}_{X,A} \otimes_\tau \mathcal{S}_{Y,B}$ in the case $\tau \in \{\max, c, \min\}$, and in $\mathcal{S}_{X,A}^{\min} \otimes_{\min} \mathcal{S}_{Y,B}^{\min}$ in the case $\tau = \text{omin}$.

Theorem 3.6. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a game. Then*

$$(4) \quad \lambda_{\text{ns}}(x, y, a, b) = 0 \iff e_{x,a} \otimes e_{y,b} \in J_{\max}(\lambda);$$

$$(5) \quad \lambda_{\text{qc}}(x, y, a, b) = 0 \iff e_{x,a} \otimes e_{y,b} \in J_c(\lambda);$$

$$(6) \quad \lambda_{\text{qa}}(x, y, a, b) = 0 \iff e_{x,a} \otimes e_{y,b} \in J_{\min}(\lambda);$$

$$(7) \quad \lambda_{\text{det}}(x, y, a, b) = 0 \iff e_{x,a} \otimes e_{y,b} \in J_{\text{omin}}(\lambda).$$

Moreover, the map $p \rightarrow s_p$ defines a one-to-one correspondence between

- (i) the perfect non-signalling strategies for $\text{Ref}_{\text{ns}}(\mathcal{G})$ and the states on $(\mathcal{S}_{X,A} \otimes_{\max} \mathcal{S}_{Y,B})/J_{\max}(\lambda)$;
- (ii) the perfect non-signalling strategies for $\text{Ref}_{\text{qc}}(\mathcal{G})$ and the states on $(\mathcal{S}_{X,A} \otimes_c \mathcal{S}_{Y,B})/J_c(\lambda)$;
- (iii) the perfect non-signalling strategies for $\text{Ref}_{\text{qa}}(\mathcal{G})$ and the states on $(\mathcal{S}_{X,A} \otimes_{\min} \mathcal{S}_{Y,B})/J_{\min}(\lambda)$;
- (iv) the perfect non-signalling strategies for $\text{Ref}_{\text{loc}}(\mathcal{G})$ and the states on $(\mathcal{S}_{X,A}^{\min} \otimes_{\min} \mathcal{S}_{Y,B}^{\min})/J_{\text{omin}}(\lambda)$. Also, $\lambda_{\text{det}} = \lambda_{\text{loc}}$.

Proof. The equivalences (4)-(7) follow from Theorem 3.1.

(i) We have that p is a perfect non-signalling strategy for $\text{Ref}_{\text{ns}}(\mathcal{G})$ if and only if p is a perfect non-signalling strategy for \mathcal{G} , if and only if s_p annihilates $J(\lambda)$, if and only if s_p annihilates $J_{\max}(\lambda)$, if and only if s_p induces a state on the quotient operator system $(\mathcal{S}_{X,A} \otimes_{\max} \mathcal{S}_{Y,B})/J_{\max}(\lambda)$.

Now suppose that $p \neq p'$ are two perfect non-signaling strategies for $\text{Ref}_{\text{ns}}(\mathcal{G})$, then because their corresponding states are well-defined on the quotient, we have that for some a, b, x, y ,

$$s_p(e_{x,a} \otimes f_{y,b} + J_{\max}(\lambda)) = p(a, b|x, y) \neq p'(a, b|x, y) = s_{p'}(e_{x,a} \otimes f_{y,b} + J_{\max}(\lambda)),$$

which shows that the correspondence is one-to-one. The proofs of the correspondence for (ii), (iii), and (iv) are similar to (i).

Finally, to see that $\lambda_{\text{det}} = \lambda_{\text{loc}}$, let $p \in \mathcal{C}_{\text{loc}}$. By Theorem 3.1, there exist $l \in \mathbb{N}$, $p_k \in \mathcal{C}_{\text{det}}$, and $t_k \in (0, 1]$, $k = 1, \dots, l$, such that $p = \sum_{k=1}^l t_k p_k$. It follows that $N(p) = \cap_{k=1}^l N(p_k)$. Thus,

$$\cap\{N(p) : p \in \mathcal{C}_{\text{loc}}(\lambda)\} = \cap\{N(p) : p \in \mathcal{C}_{\text{det}}(\lambda)\},$$

and hence $\lambda_{\text{loc}} = \lambda_{\text{det}}$. \square

In view of Example 3.4, it is natural to consider chromatic non-signalling covers of graphs. Let G be a graph and $x \in \{\text{det}, \text{loc}, \text{qm}, \text{q}, \text{qs}, \text{qa}, \text{qc}, \text{ns}\}$. As customary, let $\chi_x(G)$ be the x -chromatic number of G (see [33]). The *chromatic x -cover* $\text{Chrom}_x(G)$ of G is the largest supergraph that is coloured by every $\chi_x(G)$ -colouring of G . We write $\text{Chrom}(G) = \text{Chrom}_{\text{det}}(G)$. Example 3.4 shows that $\text{Chrom}(G)$ can be strictly larger than $\chi(G)$.

Corollary 3.7. *For any graph G , we have that $\text{Chrom}(G) = \text{Chrom}_{\text{loc}}(G)$.*

Example 3.8. Let G and H be graphs. If u and v are adjacent vertices, we will write $u \sim v$. A *graph homomorphism* from G to H is a function $\varphi : V(G) \rightarrow V(H)$ that preserves adjacency, i.e., such that if $u, v \in V(G)$ and $u \sim v$, then $\varphi(u) \sim \varphi(v) \in V(H)$. The (G, H) -*homomorphism game* (see [25, 27]) has input sets $X = Y = V(G)$, output sets $A = B = V(H)$, and rule function λ is given by

$$\lambda(x, y, a, b) = \begin{cases} 0 & \text{if } (x = y \ \& \ a \neq b) \text{ or } (x \sim y \ \& \ a \not\sim b) \\ 1 & \text{otherwise.} \end{cases}$$

In other words, the two players are given vertices of G and must respond with vertices of H . If they both receive the same vertex of G , then they must answer with the same vertex of H . If they are given adjacent vertices of G , they must respond with adjacent vertices of H . It is easy to notice that the deterministic strategies for the (G, H) -homomorphism game are in one-to-one correspondence with the homomorphisms from G to H .

A *walk* of length ℓ in a graph G is a sequence u_0, u_1, \dots, u_ℓ of vertices of G such that $u_{i-1} \sim u_i$ for all $i \in \{1, \dots, \ell\}$. We say that this is a walk from u_0 to u_ℓ , and call u_0 and u_ℓ the endpoints of the walk. Suppose that φ is a homomorphism from G to H . Clearly, if u_0, \dots, u_ℓ is a walk of length ℓ in G , then $\varphi(u_0), \dots, \varphi(u_\ell)$ is a walk of length ℓ in H . Therefore, if $u, v \in V(G)$ are the endpoints of a walk of length ℓ in G , then $\varphi(u), \varphi(v) \in V(H)$ are the endpoints of a walk of length ℓ in H . It follows that $\lambda_{\text{loc}}(x, y, a, b) = 0$ when there is a walk of length ℓ with endpoints $x, y \in V(G)$, but there is no walk of length ℓ with endpoints $a, b \in V(H)$. It follows from this that, for some

choices of G and H , the (G, H) -homomorphism game is not loc-reflexive. In fact, a special case of this is given in Example 3.4. There, the graph G is the path on four vertices, and H is the complete graph on two vertices. In G , vertices 1 and 4 have a walk of length 3 between them. In H , there is no walk of length 3 between a vertex and itself, and so vertices 1 and 4 must be mapped to distinct vertices of H .

Surprisingly, it was shown in [25] that a similar fact holds even for quantum strategies for the homomorphism game. More precisely, if $p \in \mathcal{C}_q$, is a perfect correlation for the (G, H) -homomorphism game, and $x, y \in V(G)$ are the endpoints of a walk of length ℓ in G , then $p(a, b|x, y) = 0$ unless $a, b \in V(H)$ are the endpoints of a walk of length ℓ in H . Therefore, as in the deterministic/local case, we have that $\lambda_q(x, y, a, b) = 0$ whenever there is a walk of length ℓ in G with endpoints $x, y \in V(G)$, but there is no walk of length ℓ in H with endpoints $a, b \in V(H)$. This can be used to show that, for certain graphs G and H , the (G, H) -homomorphism game is not q-reflexive.

Let us consider the special case where $x = y$ and there is a walk of length ℓ beginning and ending at $x \in V(G)$ (this is called a *closed* walk). In this case, if p is a perfect q-strategy, we have that $p(a, b|x, x) = 0$ unless $a = b$ (this is by definition) and a is contained in a closed walk of length ℓ . Furthermore, since $p(a, b|x, x) = 0$ whenever $b \neq a$, we have that, if $p(a|x)$ is the corresponding marginal probability (given by $p(a|x) = \sum_{b \in V(H)} p(a, b|x, x)$) then $p(a|x) = p(a, a|x, x)$. Therefore, for any perfect q-strategy p of the (G, H) -homomorphism game, if $x \in V(G)$ is contained in a closed walk of length ℓ , then we have that the marginal $p(a|x)$ is equal to zero unless $a \in V(H)$ is also contained in a closed walk of length ℓ . Thus, if Alice (or Bob) receive a vertex contained in a closed walk of length ℓ , then they must respond with a vertex contained in a closed walk of length ℓ if they are employing a perfect q-strategy. This implies that $\lambda_q(a, b|x, y) = 0$ if $x \in V(G)$ is contained in a closed walk of length ℓ , but $a \in V(H)$ is not, regardless of the values of y and b . This is noteworthy because it is an example where λ_q has additional zeros that depend only on the input and output of a single party, even though λ has no such zeros.

The above remarks show that there are many homomorphism games which are not q-reflexive. However, the proof of the above fact about walks that was given for q-strategies in [25] works just as well for qc-strategies. So these homomorphism games will also not be qc-reflexive. Next we will see an example of a game that is not even ns-reflexive.

Example 3.9. The (G, H) -*isomorphism game* is similar to the homomorphism game described in Example 3.8 but, in it, both adjacency and non-adjacency must be preserved. More precisely, in the (G, H) -isomorphism game, the input and output sets for both players are all equal to $V(G) \cup V(H)$, where the vertex sets of the two graphs are assumed to be disjoint. If Alice (Bob) receives a vertex from G , she (he) must respond with a vertex

from H , and vice versa. If this first condition of the game is met, then Alice either receives as input or sends as output a vertex g_A of G . Similarly, Alice sends or receives a vertex h_A of H . We can define g_B and h_B analogously for Bob. The remaining rule of the (G, H) -isomorphism game is that these four vertices must satisfy $\text{rel}(g_A, g_B) = \text{rel}(h_A, h_B)$, where rel is a function determining whether two vertices are equal, adjacent, or distinct and non-adjacent. All of these conditions are encoded in the rule function λ for the (G, H) -isomorphism game.

A partition P_1, \dots, P_k of the vertex set $V(G)$ of a graph G is said to be equitable if there exist integers c_{ij} for $i, j \in \{1, \dots, k\}$ such that each vertex in P_i is adjacent to precisely c_{ij} vertices in P_j . We refer to the numbers c_{ij} as the *partition numbers* of the partition P_1, \dots, P_k . For example, the trivial partition into singletons is always equitable for any graph. Also, the single element partition is equitable for a graph G if and only if every vertex of G has the same number of neighbors, i.e. if and only if G is *regular*. In general, given two equitable partitions of a graph G , their *join* will be an equitable partition of G . It follows that any graph has a unique (up to permutation of the parts) coarsest equitable partition. Suppose that P_1, \dots, P_k and Q_1, \dots, Q_ℓ are the coarsest equitable partitions of G and H with partition numbers c_{ij} for $i, j \in \{1, \dots, k\}$, and d_{ij} for $i, j \in \{1, \dots, \ell\}$, respectively. We say that G and H have *common coarsest equitable partitions* if $k = \ell$, and (up to some permutation of the indices) $|P_i| = |Q_i|$ for all $i = 1, \dots, k$, and $c_{ij} = d_{ij}$ for all $i, j \in \{1, \dots, k\}$. In [1], it was shown that there exists a perfect ns-strategy for the (G, H) -isomorphism game if and only if the graphs have *common coarsest equitable partitions*.

Now suppose that G and H are graphs with common coarsest equitable partitions P_1, \dots, P_k and Q_1, \dots, Q_k respectively. Though it is not explicitly stated in [1], it follows from the proof of Lemma 4.2 therein and the proof of [35, Theorem 2.2], that in any perfect ns-strategy p for the (G, H) -isomorphism game, the marginal probability $p(h|g) = p(h, h|g, g)$ vanishes unless $g \in P_i$ and $h \in Q_i$ for some $i \in \{1, \dots, k\}$. Therefore, we obtain that $\lambda_{\text{ns}}(g, g', h', h') = 0$ if $g \in P_i$ and $h \in Q_j$ with $i \neq j$, regardless of the values of g' and h' . Thus, if $k \geq 2$, then the (G, H) -isomorphism game is not ns-reflexive. This occurs if and only if the graphs G and H are not regular.

4. IMITATION GAMES

In this section, we introduce a new class of games that we call *imitation games*, give examples, and establish some first properties.

Definition 4.1. A game $\mathcal{G} = (X, Y, A, B, \lambda)$ will be called an *imitation game* if

- (a) for every $x \in X$ and $a, a' \in A$ with $a \neq a'$, there exists $y \in Y$ such that

$$\sum_{b \in B} \lambda(a, b|x, y) \lambda(a', b|x, y) = 0,$$

and

- (b) for every $y \in Y$ and $b, b' \in B$ with $b \neq b'$, there exists $x \in X$ such that

$$\sum_{a \in A} \lambda(a, b|x, y) \lambda(a, b'|x, y) = 0.$$

Conditions (a) and (b) in Definition 4.1 assert that, in some sense, in a perfect strategy for \mathcal{G} , Alice's answers are completely determined by Bob's answers and vice versa.

Given an imitation game \mathcal{G} , we wish to define a corresponding *-algebra $A^*(\mathcal{G})$ and C*-algebra $C^*(\mathcal{G})$. First recall that given any set of generators, $\{a_t\}_{t \in T}$ then the free unital *-algebra on these generators is just the complex linear span of words in the generators and the unit 1, with multiplication defined by concatenation of words and with the *-operation given by reversing the order in words and taking the complex conjugates of the coefficients. When we impose a *relation* on these words, such as, $a_t^2 = a_t$, we mean that we are forming the quotient of the free *-algebra by the smallest 2-sided ideal J containing $a_t^2 - a_t$ and satisfying, $x \in J \implies x^* \in J$ (this condition is called **-closed*). More generally given a set of generators and relations, by the *free *-algebra given by generators and relations*, we mean the quotient of the free *-algebra on those generators by the smallest 2-sided *-closed ideal containing all of the relations. Of course, it is possible that when we impose relations that the ideal J is the entire free algebra, equivalently, that $1 \in J$, in which case this algebra is (0).

Given a unital *-algebra \mathcal{A} , its *C*-completion* \mathcal{B} is the C*-algebra obtained by first defining a seminorm on \mathcal{A} by setting

$$\|a\| = \sup\{\|\pi(a)\| : \pi : \mathcal{A} \rightarrow B(\mathcal{H})\},$$

where the supremum is taken over all Hilbert spaces \mathcal{H} and all unital *-homomorphisms $\pi : \mathcal{A} \rightarrow B(\mathcal{H})$, and then completing. In the case that there are no unital *-homomorphisms, we say that the C*-completion is (0).

Definition 4.2. Let \mathcal{G} be an imitation game. The *-algebra $A^*(\mathcal{G})$ of the imitation game \mathcal{G} is the unital free *-algebra given by generators $(p_{x,a})_{x \in X, a \in A} \cup (q_{y,b})_{y \in Y, b \in B}$ and relations:

- (a) $p_{x,a}^2 = p_{x,a}^* = p_{x,a}$, $\forall x \in X, a \in A$ and $q_{y,b} = q_{y,b}^* = q_{y,b}$, $\forall y \in Y, b \in B$.
- (b) $\sum_{a \in A} p_{x,a} = 1$ and $p_{x,a} p_{x,c} = 0$ for every $x \in X, a \neq c$;
- (c) $\sum_{b \in B} q_{y,b} = 1$ and $q_{y,b} q_{y,c} = 0$ for every $y \in Y, b \neq c$;
- (d) $\lambda_{\mathcal{G}}(x, y, a, b) = 0$ implies $p_{x,a} q_{y,b} = 0$, for any $(x, y, a, b) \in X \times Y \times A \times B$.

The C*-algebra of the game $C^*(\mathcal{G})$ is the C*-completion of $A^*(\mathcal{G})$.

Of course it is possible that either $A^*(\mathcal{G})$ or $C^*(\mathcal{G})$ is (0).

Example 4.3. Suppose that $A = B$ and $X = Y$. The *synchronicity game* \mathcal{G}^s is obtained by setting $\lambda_{\mathcal{G}^s}(a, b, x, y) = 0$ if and only if $x = y$ and $a \neq b$.

The correlations in $\mathcal{C}_{\text{ns}}(\mathcal{G}^{\text{s}})$ are called *synchronous correlations* and denoted by \mathcal{C}^{s} .

Example 4.4. Any game \mathcal{G} , with $X = Y$ and $A = B$, harder than the synchronicity game, is called *synchronous*. It is clear that any synchronous game is an imitation game. In this case, the C*-algebra of the game \mathcal{G} as in Definition 4.2 coincides with the C*-algebra of the game as defined in [17]. The class of synchronous games contains in particular the graph coloring games and the graph homomorphism games introduced and studied in [3, 25, 27, 32, 33, 28]. In the case of the graph homomorphism game, the game C*-algebra as in Definition 4.2 recovers the C*-algebra considered in [28, Section 4].

Example 4.5. Suppose that Y is a finite set of *variables* $\{v_1, \dots, v_n\}$. A *binary constraint* in v_1, \dots, v_n is an expression of the form $f((v_j)_{j \in V}) = 1$ for some $V \subseteq \{1, 2, \dots, n\}$, where $f : \{-1, 1\}^V \rightarrow \{-1, 1\}$ is a function. A *binary constraint system* \mathcal{S} is a set X of such binary constraints with the property that every variable appears in some constraint. In [6], a game $\mathcal{G}_{\mathcal{S}}$ has been associated with a binary constraint system, as follows. The input sets are X and Y as above, while the output sets are $B = \{+1, -1\}$ and $A = \{(a_i)_{i \in V} : V \subseteq \{1, 2, \dots, n\}, a_i \in \{+1, -1\}\}$. The payoff function is defined by $\lambda(x, y, (a_i)_{i \in V}, b) = 1$ if and only if x is a constraint of the form $f((v_k)_{k \in V}) = 1$, the tuple $(a_k)_{k \in V}$ satisfies the constraint x , $j \in V$, and $b = a_j$. A particular example of a binary constraint system game is the Mermin-Peres magic square game [18].

Example 4.6. Suppose that C is a finite set of possible *values* for some *variables* v_1, \dots, v_n . As before, we let X and Y represent the set of possible questions for Alice and Bob, respectively. To each $z \in X \cup Y$ we assigns a subset V_z of $\{1, 2, \dots, n\}$, in such a way that, for any $i \in \{1, 2, \dots, n\}$, there exist $x \in X$ and $y \in Y$ such that $i \in V_x \cap V_y$. We let the sets A and B be both equal to the set of tuples $(c_i)_{i \in V}$ for some $V \subseteq \{1, 2, \dots, n\}$, which we can think of as a valuation of the variables v_i for $i \in V$. A *variable assignment game* \mathcal{G} on (X, Y, A, B) is any game whose payoff function $\lambda_{\mathcal{G}} : X \times Y \times A \times B \rightarrow \{0, 1\}$ satisfies the conditions

- (a) $\lambda(x, y, (a_i)_{i \in V}, (b_j)_{j \in W}) = 1$ implies $V = V_x$ and $W = V_y$, and
- (b) $a_i = b_i$ for every $i \in V_x \cap V_y$.

It is clear that any binary constraint system game is a variable assignment game.

We show that every variable assignment game is an imitation game. Suppose that $x \in X$ and $a, a' \in A$ with $a \neq a'$, say $a = (a_i)_{i \in V}$ and $a' = (a'_i)_{i \in V'}$, where V and V' are subsets of $\{1, \dots, n\}$. If $V \neq V'$ then either $V \neq V_x$, in which case choosing an arbitrary $y \in Y$ we have that $\lambda(x, y, a, b) = 0$ for all $b \in B$, or $V' \neq V_x$, in which case choosing an arbitrary $y \in Y$ we have that $\lambda(x, y, a', b) = 0$ for all $b \in B$. Suppose that $V = V'$, and choose $i \in V$ such that $a_i \neq a'_i$. Let $y \in Y$ be such that $i \in V_x \cap V_y$. Let $b = (b_j)_{j \in W} \in B$. If

$W \neq V_y$ then clearly $\lambda(x, y, a, b) = 0$. If, on the other hand, $W = V_y$ then $\lambda(x, y, a, b)\lambda(x, y, a', b) = 0$. By symmetry, \mathcal{G} is an imitation game.

Example 4.7. We call a non-local game $\mathcal{G} = (X, Y, A, B, \lambda)$ a *mirror game* if there exist functions $\xi : X \rightarrow Y$ and $\eta : Y \rightarrow X$ such that

$$\lambda(x, \xi(x), a, b)\lambda(x, \xi(x), a', b) = 0, \quad x \in X, \quad a, a' \in A, \quad b \in B, \quad a \neq a'$$

and

$$\lambda(\eta(y), y, a, b)\lambda(\eta(y), y, a, b') = 0, \quad y \in Y, \quad a \in A, \quad b, b' \in B, \quad b \neq b'.$$

Clearly, every mirror game is an imitation game, and the difference between the two classes consists in that, given $x \in X$ (resp. $y \in Y$) and $a, a' \in A$ (resp. $b, b' \in B$) with $a \neq a'$ (resp. $b \neq b'$), the element $y \in Y$ (resp. $x \in X$) satisfying condition (a) (resp. (b)) in Definition 4.1 depends on a and a' (resp. b and b') for general imitation games, and is independent of them for mirror games.

Example 4.8. Recall that a non-local game $\mathcal{G} = (X, Y, A, B, \lambda)$ is called *unique* (see e.g. [36]) if for every $(x, y) \in X \times Y$ there exists a bijection $\phi_{x,y} : A \rightarrow B$ such that

$$\lambda(x, y, a, b) = 1 \iff b = \phi_{x,y}(a).$$

Every unique game is a mirror game; indeed, given $x \in X$, any element y of Y satisfies (a) in Definition 4.1 for any choice of $a, a' \in A$ with $a \neq a'$; the claim follows by symmetry.

Most known examples of games admitting a perfect quantum but not local strategy fall within the class of imitation games. However, there do exist examples that do not fulfill the requirements set out in the definition of imitation games. One family of such examples can be obtained by considering the non-local game associated with one-shot entanglement-assisted zero-error codes [9]. In the setting of zero-error communication Alice uses a classical noisy channel \mathcal{N} , characterized by its confusability graph $G = G(\mathcal{N})$, to communicate one out of $m \in \mathbb{N}$ messages to Bob with zero chance of confusion. The idea for turning this communication scenario into a non-local game is to think of Alice's answer as the input to the channel and the question for Bob as output of the channel. Specifically, we let $X = \{1, \dots, m\}$, $A = V(G)$, $Y = E(G)$, and $B = X$. The payoff function $\lambda(x, y, a, b)$ is defined to be 1, if either

- (1) edge y is not incident to vertex a (Bob's question cannot arise as an output of \mathcal{N} upon inputting Alice's answer a) or
- (2) y is incident to a and $x = b$ (Bob correctly decoded the intended message).

If the one-shot entanglement-assisted zero-error capacity, $c_{SE}(\mathcal{N})$, of \mathcal{N} exceeds its classical counterpart, then the above game has a perfect quantum but not local strategy when $m = c_{SE}(\mathcal{N})$. However, the above game is not

an imitation game, since $\lambda(a, b|x, y)\lambda(a', b|x, y) = 1$ if we pick b to be equal to x (this violates condition (a) in Definition 4.1).

Proposition 4.9. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a variable assignment game with n variables and set of variable values C . Its game C^* -algebra $C^*(\mathcal{G})$ is $*$ -isomorphic to the universal C^* -algebra generated by a family $\{e_{i,c} : i = 1, \dots, n, c \in C\}$ of projections subject to the relations*

- (a) $\sum_{c \in C} e_{i,c} = 1$, $i = 1, 2, \dots, n$;
- (b) $e_{i,c}e_{j,d} = e_{j,d}e_{i,c}$ for all $c, d \in C$, whenever there exists $z \in X \cup Y$ such that $i, j \in V_z$;
- (c) the projections $\prod_{i \in V_x} e_{i,a_i}$ and $\prod_{j \in V_y} e_{j,b_j}$ are orthogonal whenever $\lambda(x, y, (a_i)_{i \in V_x}, (b_i)_{i \in V_y}) = 0$.

Proof. Let \mathcal{U} be the universal C^* -algebra as in the statement, and fix a faithful $*$ -representation $C^*(\mathcal{G}) \subseteq \mathcal{B}(\mathcal{H})$. By the universal property of $C^*(\mathcal{G})$, there exists a canonical $*$ -homomorphism $\pi : C^*(\mathcal{G}) \rightarrow \mathcal{U}$ such that

$$\pi(p_{x,(a_i)_{i \in V}}) = \begin{cases} \prod_{i \in V_x} e_{i,a_i} & \text{if } V = V_x, \\ 0 & \text{otherwise} \end{cases}$$

and

$$\pi(q_{y,(b_i)_{i \in V}}) = \begin{cases} \prod_{i \in V_y} e_{i,a_i} & \text{if } V = V_y, \\ 0 & \text{otherwise} \end{cases}.$$

We now define a $*$ -homomorphism $\rho : \mathcal{U} \rightarrow C^*(\mathcal{G})$ which is the inverse of π . If $x \in X$, $i \in V_x$, and $c \in C$, let

$$g_{i,c}^x = \sum \{p_{x,a} : a \in C^{V_x}, a_i = c\}.$$

Similarly if $y \in Y$, $i \in V_y$, and $c \in C$, let

$$h_{i,c}^y = \sum \{q_{y,b} : b \in C^{V_y}, b_i = c\}.$$

It is clear that $g_{i,c}^x$ and $h_{i,c}^y$ are projections. Fix $x \in X$ and $y \in Y$. Observe that, for every $i \in V_x$ and $j \in V_y$,

$$(8) \quad \sum_{c \in C} g_{i,c}^x = \sum_{c \in C} h_{j,c}^y = 1.$$

Suppose that $i \in V_x \cap V_y$. Since $p_{x,a}q_{y,b} = 0$ whenever $a_i \neq b_i$, we have

$$(9) \quad 1 = \left(\sum_{c \in C} g_{i,c}^x \right) \left(\sum_{d \in C} h_{j,d}^y \right) = \sum_{c \in C} g_{i,c}^x h_{i,c}^y.$$

Let $|\xi\rangle \in \mathcal{H}$ be a unit vector, $|\alpha\rangle = \left(g_{i,c}^x |\xi\rangle \right)_{c \in C}$ and $|\beta\rangle = \left(h_{i,c}^y |\xi\rangle \right)_{c \in C}$. By (8), $|\alpha\rangle$ and $|\beta\rangle$ are unit vectors in $\oplus_{c \in C} \mathcal{H}$ while, by (9), they satisfy the relation $\langle \alpha | \beta \rangle = 1$. Thus $\alpha = \beta$, that is, $g_{i,c}^x |\xi\rangle = h_{i,c}^y |\xi\rangle$ for every $|\xi\rangle \in \mathcal{H}$ and hence $g_{i,c}^x = h_{i,c}^y$. Therefore $g_{i,c}^x = h_{i,c}^y$ for every $i \in V_x \cap V_y$ and every $c \in C$. We conclude that $g_{i,c}^x = g_{i,c}^{x'} := g_{i,c}$ for any $x, x' \in X$, $i \in V_x \cap V_{x'}$, and $c \in C$. Since $(p_{x,a})_{a \in A}$ and $(q_{y,b})_{b \in B}$ are PVM's for all

$x \in X$ and $y \in Y$, the family $\{g_{i,c} : i = 1, \dots, n, c \in C\}$ satisfies condition (b). Condition (c) is straightforward. Therefore the map $e_{i,c} \mapsto g_{i,c}$ extends to a *-homomorphism $\rho : \mathcal{U} \rightarrow C^*(\mathcal{G})$, which is easily seen to be the inverse of π . \square

A particular instance of a binary constraint system is a *linear* binary constraint system as considered in [5]. This is a binary constraint system \mathcal{S} where the functions $f : \{-1, 1\}^V \rightarrow \{-1, 1\}$ are of the form $f((\lambda_i)_{i \in V}) = (-1)^\rho \prod_{i \in V} \lambda_i$ for some $\rho \in \{0, 1\}$. The *solution group* $\Gamma(\mathcal{S})$ associated to such a linear binary constraint system as in [5, 37] is defined to be the group generated by involutions u_1, \dots, u_n, J subject to the following relations: J commutes with u_1, \dots, u_n , and u_i, u_j commute whenever a constraint of the form $(-1)^\rho \prod_{k \in V} \lambda_k = 1$, with $i, j \in V$, is an element of the input set X , in which case $J^\rho \prod_{k \in V} u_k = 1$.

We specialise Example 4.6 to the case of a binary constraint system \mathcal{S} and describe the canonical non-local game $\mathcal{G}_\mathcal{S}$ associated with it. We have that X is the set of all constraints and $C = \{+1, -1\}$. For $x \in X$, V_x is the set of indices of variables that appear in the constraint x . We set $Y = \{1, 2, \dots, n\}$ and $V_y = \{y\}$ for $y \in Y$. The payoff function λ is obtained by letting, for $x \in X$ and $j \in V_x$, where x is the constraint $(-1)^{\rho_x} \prod_{i \in V_x} \lambda_i = 1$, $\lambda((a_i)_{i \in V_x}, a_j, x, j) = 1$ if and only if $(-1)^{\rho_x} \prod_{i \in V_x} a_x = 1$.

We denote by $C^*(\Gamma(\mathcal{S}))$ the full group C^* -algebra of the solution group $\Gamma(\mathcal{S})$ and identify $\Gamma(\mathcal{S})$ with a subgroup of the unitary group of $C^*(\Gamma(\mathcal{S}))$. A natural question that arises, and is addressed in the next proposition, is what the relation between $C^*(\mathcal{G}_\mathcal{S})$ and $C^*(\Gamma(\mathcal{S}))$ is.

Proposition 4.10. *Let \mathcal{S} be a linear binary constraint system with a corresponding non-local game $\mathcal{G}_\mathcal{S}$ and solution group $\Gamma(\mathcal{S})$. The game C^* -algebra $C^*(\mathcal{G}_\mathcal{S})$ is *-isomorphic to the quotient of the full group C^* -algebra $C^*(\Gamma(\mathcal{S}))$ by the relation $J + 1 = 0$.*

Proof. Let J be the closed two sided ideal of $C^*(\Gamma(\mathcal{S}))$ generated by $J + 1$ and $\mathcal{A} = C^*(\Gamma(\mathcal{S}))/J$. We denote by $\hat{u}_1, \dots, \hat{u}_n$ the images inside \mathcal{A} of the canonical generators of $\Gamma(\mathcal{S})$ under the canonical quotient mapping $C^*(\Gamma(\mathcal{S})) \rightarrow \mathcal{A}$. We consider the description of $C^*(\mathcal{G})$ as in Proposition 4.9. (Recall that a binary constraint system game is, in particular, a variable assignment game in the sense of Example 4.6, and therefore Proposition 4.9 applies.) Consider the assignment $u_i \mapsto e_{i,+1} - e_{i,-1}$ and $J \mapsto -1$. It is easy to verify that this defines a unitary representation of $\Gamma(\mathcal{S})$, and hence it extends to a *-homomorphism $\pi : C^*(\Gamma(\mathcal{S})) \rightarrow C^*(\mathcal{G})$. Since such a *-homomorphism maps J to -1 , it induces a *-homomorphism $\mathcal{A} \rightarrow C^*(\mathcal{G})$. Conversely, the assignment $e_{i,+1} \mapsto \hat{u}_i^+$ and $e_{i,-1} \mapsto \hat{u}_i^-$ defines a *-homomorphism $C^*(\mathcal{G}) \rightarrow \mathcal{A}$, where \hat{u}_i^+ and \hat{u}_i^- denote the spectral subspaces of the selfadjoint unitary \hat{u}_i associated with $+1$ and -1 , respectively. It is clear from the definition that the *-homomorphisms π and ρ are inverses of each other. \square

5. PERFECT STRATEGIES FOR IMITATION GAMES

In [17] the existence of various types of perfect strategies for synchronous games was characterised in terms of types of traces on the game C^* -algebra. In this section, we extend those results to imitation games. Given an imitation game \mathcal{G} , we characterise the elements of the set $\mathcal{C}_{\text{qc}}(\mathcal{G})$ of perfect quantum commuting strategies, and of the set $\mathcal{C}_{\text{q}}(\mathcal{G})$ of perfect quantum strategies, for the game \mathcal{G} , in terms of traces on the game C^* -algebra $C^*(\mathcal{G})$.

Theorem 5.1. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be an imitation game and p be a non-signalling correlation on (X, Y, A, B) . The following statements are equivalent:*

- (a) $p \in \mathcal{C}_{\text{qc}}(\mathcal{G})$;
- (b) *the C^* -algebra of the game $C^*(\mathcal{G})$ is nonzero, and there exists a tracial state τ on $C^*(\mathcal{G})$ such that $p(a, b|x, y) = \tau(p_{x,a}q_{y,b})$.*

Proof. (i) \Rightarrow (ii) Fix $p \in \mathcal{C}_{\text{qc}}(\mathcal{G})$. By Corollary 3.2, there exists a separable Hilbert spaces \mathcal{H} , a unit vector $|\xi\rangle \in \mathcal{H} \otimes \mathcal{H}$, and PVMs $(P_{x,a})_{a \in A}$ and $(Q_{y,b})_{b \in B}$ on \mathcal{H} for $x \in X$ and $y \in Y$ such that

$$p(a, b|x, y) = \langle \xi | P_{x,a} Q_{y,b} | \xi \rangle, \quad (x, y, a, b) \in X \times Y \times A \times B.$$

Let \mathcal{A} (resp. \mathcal{B}) be the C^* -algebra generated by $\{P_{x,a} : x \in X, a \in A\}$ (resp. $\{Q_{y,b} : y \in Y, b \in B\}$). Let \mathcal{M} (resp. \mathcal{N}) be the WOT-closure of \mathcal{A} (resp. \mathcal{B}). Observe that $\mathcal{M} \subseteq \mathcal{N}'$. For $x \in X$, $y \in Y$ and $b \in B$, set

$$\Pi_{y,b}^x = \sum_{a \in A, \lambda(x,y,a,b)=1} P_{x,a}.$$

Clearly, $\Pi_{y,b}^x$ is a projection in \mathcal{A} . We have

$$\begin{aligned} \langle \xi | Q_{y,b} | \xi \rangle &= \sum_{a \in A} \langle \xi | P_{x,a} Q_{y,b} | \xi \rangle = \sum_{a \in A, \lambda(x,y,a,b)=1} \langle \xi | P_{x,a} Q_{y,b} | \xi \rangle \\ &= \langle \xi | \Pi_{y,b}^x Q_{y,b} | \xi \rangle. \end{aligned}$$

Since $(I - \Pi_{y,b}^x)Q_{y,b}$ is an idempotent, this shows that

$$\|(I - \Pi_{y,b}^x)Q_{y,b} | \xi \rangle\|^2 = \langle \xi | (I - \Pi_{y,b}^x)Q_{y,b} | \xi \rangle = 0,$$

that is,

$$(10) \quad \Pi_{y,b}^x Q_{y,b} | \xi \rangle = Q_{y,b} | \xi \rangle, \quad x \in X, y \in Y, b \in B.$$

By assumption, for $b \neq b'$ and $y \in Y$ there exists $x \in X$ such that $\sum_{a \in A} \lambda(x, y, a, b) \lambda(x, y, a, b') = 0$. For such a choice of x, y, b, b' we have

$$\langle \xi | \Pi_{y,b}^x Q_{y,b'} | \xi \rangle = \sum_{a \in A, \lambda(x,y,a,b)=1} \langle \xi | P_{x,a} Q_{y,b'} | \xi \rangle = 0;$$

henceforth $\Pi_{y,b}^x Q_{y,b'} | \xi \rangle = 0$.

Let $\Pi_{y,b}$ be the projection onto the intersection of the ranges of $\Pi_{y,b}^x$ for $x \in X$. Since the projections in \mathcal{M} form a complete sublattice of the lattice of projections in $\mathcal{B}(\mathcal{H})$, we have that $\Pi_{y,b} \in \mathcal{M}$. By (10),

$$(11) \quad \Pi_{y,b} Q_{y,b} |\xi\rangle = Q_{y,b} |\xi\rangle.$$

Suppose now that $b' \in B$ and $b' \neq b$. By the preceding paragraph, there exists $x \in X$ such that $\Pi_{y,b}^x Q_{y,b'} |\xi\rangle = 0$. Thus $\Pi_{y,b} Q_{y,b'} |\xi\rangle = 0$ whenever $b' \neq b$. Thus, using (11), we obtain

$$(12) \quad \Pi_{y,b} |\xi\rangle = \Pi_{y,b} \left(\sum_{b' \in B} Q_{y,b'} \right) |\xi\rangle = \Pi_{y,b} Q_{y,b} |\xi\rangle = Q_{y,b} |\xi\rangle.$$

Similarly, define

$$\Xi_{x,a}^y = \sum_{b \in B, \lambda(a,b,x,y)=1} Q_{y,b},$$

let $\Xi_{x,a} \in \mathcal{N}$ be the projection onto the intersection of the ranges of $\Xi_{x,a}^y$ for $y \in Y$, and show that

$$(13) \quad \Xi_{x,a} |\xi\rangle = P_{x,a} |\xi\rangle, \quad x \in X, a \in A.$$

Let $\mathcal{K}_{\mathcal{M}}$ (resp. $\mathcal{K}_{\mathcal{N}}$) be the closure of the span of $\{A |\xi\rangle : A \in \mathcal{M}\}$ (resp. $\{B |\xi\rangle : B \in \mathcal{N}\}$). Suppose that $x_1, \dots, x_n \in X$ and $a_1, \dots, a_n \in A$. Then

$$P_{x_1, a_1} \cdots P_{x_n, a_n} |\xi\rangle = \Xi_{x_n, a_n} \cdots \Xi_{x_1, a_1} |\xi\rangle \in \mathcal{K}_{\mathcal{N}};$$

therefore $\mathcal{K}_{\mathcal{M}} \subseteq \mathcal{K}_{\mathcal{N}}$. Similarly, $\mathcal{K}_{\mathcal{N}} \subseteq \mathcal{K}_{\mathcal{M}}$. Set $\mathcal{K} := \mathcal{K}_{\mathcal{M}} = \mathcal{K}_{\mathcal{N}}$. Clearly, \mathcal{K} is invariant under the projections $P_{x,a}$ and $Q_{y,b}$; after replacing $P_{x,a}$ and $Q_{y,b}$ with their restrictions to \mathcal{K} , we can thus assume that $\mathcal{K} = \mathcal{H}$.

Fix $y \in Y$ and $b, b' \in B$ with $b \neq b'$. By assumption, there exists $x \in X$ such that $\sum_{a \in A} \lambda(x, y, a, b) \lambda(x, y, a, b') = 0$. We thus have that $\Pi_{y,b}^x \Pi_{y,b'}^x = 0$. Since $\Pi_{y,b} \leq \Pi_{y,b}^x$ and $\Pi_{y,b'} \leq \Pi_{y,b'}^x$, we conclude that $\Pi_{y,b} \Pi_{y,b'} = 0$. On the other hand, if $Z \in \mathcal{N}$ then, using (12), we have

$$\sum_{b \in B} \Pi_{y,b} Z |\xi\rangle = \sum_{b \in B} Z \Pi_{y,b} |\xi\rangle = Z \sum_{b \in B} Q_{y,b} |\xi\rangle = Z |\xi\rangle.$$

Since this is true for every $Z \in \mathcal{N}$ we conclude that $\sum_{b \in B} \Pi_{y,b} = I$.

We show that the vector state $\tau : \mathcal{A} \rightarrow \mathbb{C}$, given by $\tau(A) = \langle \xi | A |\xi\rangle$, is tracial. Indeed, for $x_1, \dots, x_n, x'_1, \dots, x'_m \in X$ and $a_1, \dots, a_n, a'_1, \dots, a'_m \in A$

\mathcal{A} , using (13), we have

$$\begin{aligned}
 & \left\langle \xi \mid (P_{x_1, a_1} \cdots P_{x_n, a_n}) (P_{x'_1, a'_1} \cdots P_{x'_m, a'_m}) \mid \xi \right\rangle \\
 &= \left\langle \xi \mid \Xi_{x'_m, a'_m} (P_{x_1, a_1} \cdots P_{x_n, a_n}) (P_{x'_1, a'_1} \cdots P_{x'_{m-1}, a'_{m-1}}) \mid \xi \right\rangle \\
 &= \left\langle \Xi_{x'_m, a'_m} \xi \mid (P_{x_1, a_1} \cdots P_{x_n, a_n}) (P_{x'_1, a'_1} \cdots P_{x'_{m-1}, a'_{m-1}}) \mid \xi \right\rangle \\
 &= \left\langle P_{x'_m, a'_m} \xi \mid (P_{x_1, a_1} \cdots P_{x_n, a_n}) (P_{x'_1, a'_1} \cdots P_{x'_{m-1}, a'_{m-1}}) \mid \xi \right\rangle \\
 &= \left\langle P_{x'_m, a'_m} \xi \mid \Xi_{x'_{m-1}, a'_{m-1}} (P_{x_1, a_1} \cdots P_{x_n, a_n}) (P_{x'_1, a'_1} \cdots P_{x'_{m-2}, a'_{m-2}}) \mid \xi \right\rangle \\
 &= \left\langle \xi \mid \Xi_{x'_{m-1}, a'_{m-1}} P_{x'_m, a'_m} (P_{x_1, a_1} \cdots P_{x_n, a_n}) (P_{x'_1, a'_1} \cdots P_{x'_{m-2}, a'_{m-2}}) \mid \xi \right\rangle \\
 &= \cdots = \left\langle \xi \mid (P_{x'_1, a'_1} \cdots P_{x'_m, a'_m}) (P_{x_1, a_1} \cdots P_{x_n, a_n}) \mid \xi \right\rangle.
 \end{aligned}$$

This shows that τ is a tracial state on \mathcal{A} . A similar argument shows that τ is a tracial state on \mathcal{B} .

We next show that τ is faithful on \mathcal{A} . Suppose that $C \in \mathcal{A}$ is positive and assume that $\tau(C) = 0$. Then, for every $T \in \mathcal{A}$, we have that

$$\begin{aligned}
 |\langle \xi \mid T^* C^2 T \mid \xi \rangle| &= \tau(T^* C^2 T) = \tau(CTT^*C) \\
 &\leq \|T\|^2 \tau(C^{\frac{1}{2}} C C^{\frac{1}{2}}) \leq \|T\|^2 \|C\| \tau(C) = 0.
 \end{aligned}$$

Therefore $CT \mid \xi \rangle = 0$ for every $T \in \mathcal{A}$, and hence $C = 0$. A similar argument shows that τ is faithful on \mathcal{B} .

Suppose now that $\lambda(x, y, a, b) = 0$. Then

$$0 = \langle \xi \mid P_{x,a} Q_{y,b} \mid \xi \rangle = \langle \xi \mid P_{x,a} \Pi_{y,b} \mid \xi \rangle = \tau(P_{x,a} \Pi_{y,b});$$

therefore, $P_{x,a} \Pi_{y,b} = 0$.

This shows that the C^* -algebra of the game $C^*(\mathcal{G})$ is nonzero, and the assignment $p_{x,a} \mapsto P_{x,a}$, $q_{y,b} \mapsto \Pi_{y,b}$ defines a unital $*$ -homomorphism $\pi : C^*(\mathcal{G}) \rightarrow \mathcal{A}$. Since τ is a tracial state on \mathcal{A} , we conclude that $\tau \circ \pi$ is a tracial state on $C^*(\mathcal{G})$. For $(x, y, a, b) \in X \times Y \times A \times B$, we have

$$(\tau \circ \pi)(p_{x,a} q_{y,b}) = \tau(P_{x,a} \Pi_{y,b}) = \langle \xi \mid P_{x,a} Q_{y,b} \mid \xi \rangle = p(a, b \mid x, y).$$

This concludes the proof of the implication.

(ii) \Rightarrow (i) Suppose that there exists a tracial state τ on $C^*(\mathcal{G})$ such that

$$\tau(p_{x,a} q_{y,b}) = p(a, b \mid x, y), \quad (x, y, a, b) \in X \times Y \times A \times B.$$

Consider the left regular representation π_τ of $C^*(\mathcal{G})$ associated with τ , and the right regular representation π_τ^{op} of $C^*(\mathcal{G})^{\text{op}}$ associated with τ ; recall that they act on the Hilbert space $\mathcal{H} = L^2(C^*(\mathcal{G}), \tau)$ produced via τ through the GNS construction and are defined by setting $\pi_\tau(z) \mid w \rangle = \mid zw \rangle$ and $\pi_\tau^{\text{op}}(z) \mid w \rangle = \mid wz \rangle$ for $z, w \in C^*(\mathcal{G})$ (see [2]). Set $P_{x,a} = \pi(p_{x,a})$ and $Q_{y,b} = \rho(q_{y,b})$, $x \in X$, $y \in Y$, $a \in A$, $b \in B$. Observe that $(P_{x,a})_{a \in A}$ and $(Q_{y,b})_{b \in B}$ are PVMs and

$$\langle 1 \mid P_{x,a} Q_{y,b} \mid 1 \rangle_{L^2(C^*(\mathcal{G}), \tau)} = \langle p_{x,a} \mid q_{y,b} \rangle = \tau(p_{x,a} q_{y,b})$$

for any $(x, y, a, b) \in X \times Y \times A \times B$. \square

Corollary 5.2. *Suppose that \mathcal{G} is an imitation game. The following assertions are equivalent:*

- (1) \mathcal{G} has a perfect quantum commuting strategy;
- (2) the C^* -algebra of the game $C^*(\mathcal{G})$ is nonzero, and it has a tracial state.

In view of Proposition 4.10, Corollary 5.2 recovers [5, Theorem 4] as a particular case.

We now turn our attention to other quantum strategies. Given an imitation game \mathcal{G} , in the next theorem, we characterise the elements of the set $\mathcal{C}_q(\mathcal{G})$ in terms of traces on the game C^* -algebra in a fashion analogous to Theorem 5.1.

Theorem 5.3. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be an imitation game and p be a non-signalling correlation on (X, Y, A, B) . The following statements are equivalent:*

- (i) $p \in \mathcal{C}_{\text{qs}}(\mathcal{G})$;
- (ii) $p \in \mathcal{C}_q(\mathcal{G})$;
- (iii) $p \in \mathcal{C}_{\text{qm}}(\mathcal{G})$;
- (iv) the game C^* -algebra $C^*(\mathcal{G})$ is nonzero, and there exists a finite-dimensional C^* -algebra \mathcal{F} with a tracial state τ and a unital $*$ -homomorphism $\pi : C^*(\mathcal{G}) \rightarrow \mathcal{F}$ such that

$$p(a, b|x, y) = (\tau \circ \pi)(p_{xa}q_{yb}), \quad (x, y, a, b) \in X \times Y \times A \times B.$$

Proof. (i) \Rightarrow (iv) Suppose that $p \in \mathcal{C}_{\text{qs}}(\mathcal{G})$. There exist separable Hilbert spaces \mathcal{H}_A and \mathcal{H}_B , a unit vector $|\xi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$, and PVMs $(P_{x,a})_{a \in A}$ (resp. $(Q_{y,b})_{b \in B}$) on \mathcal{H}_A (resp. \mathcal{H}_B) for $x \in X$ (resp. $y \in Y$), such that

$$p(a, b|x, y) = \langle \xi | P_{x,a} \otimes Q_{y,b} | \xi \rangle, \quad (x, y, a, b) \in X \times Y \times A \times B.$$

Following the proof of Theorem 5.1, for $y \in Y, b \in B$ and $x \in X$, set

$$\Pi_{y,b}^x = \sum_{a \in A, \lambda(x,y,a,b)=1} P_{x,a}.$$

Similarly, for $x \in X, a \in A$ and $y \in Y$, set

$$\Xi_{x,a}^y = \sum_{b \in B, \lambda(x,y,a,b)=1} Q_{y,b}.$$

Clearly, $(\Pi_{y,b}^x)_{b \in B}$ (resp. $(\Xi_{x,a}^y)_{a \in A}$) is a family of projections in $\mathcal{B}(\mathcal{H}_A)$ (resp. $\mathcal{B}(\mathcal{H}_B)$). Let $\Pi_{y,b}$ (resp. $\Xi_{x,a}$) be the projection onto the intersection of the ranges of $\Pi_{y,b}^x$ (resp. $\Xi_{x,a}^y$) for $x \in X$ (resp. $y \in Y$).

It follows from the proof of the implication (i) \Rightarrow (ii) in Theorem 5.1 that $(\Pi_{y,b})_{b \in B}$ (resp. $(\Xi_{x,a})_{a \in A}$) is a PVM on \mathcal{H}_A (resp. \mathcal{H}_B),

$$(14) \quad (\Pi_{y,b} \otimes I) |\xi\rangle = (I \otimes Q_{y,b}) |\xi\rangle$$

and

$$(15) \quad (I \otimes \Xi_{x,a}) |\xi\rangle = (P_{x,a} \otimes I) |\xi\rangle.$$

We now follow the arguments in the proof of [6, Theorem 1]. Consider the Schmidt decomposition

$$|\xi\rangle = \sum_{i \in I} \alpha_i (|\phi_i\rangle \otimes |\psi_i\rangle)$$

for the unit vector $|\xi\rangle$. By (14) and (15),

$$(16) \quad \sum_{i \in I} \alpha_i (\Pi_{y,b} |\phi_i\rangle \otimes |\psi_i\rangle) = \sum_{i \in I} \alpha_i (|\phi_i\rangle \otimes Q_{y,b} |\psi_i\rangle)$$

and

$$(17) \quad \sum_{i \in I} \alpha_i (|\phi_i\rangle \otimes \Xi_{x,a} |\psi_i\rangle) = \sum_{i \in I} \alpha_i (P_{x,a} |\phi_i\rangle \otimes |\psi_i\rangle).$$

Let $(\mu_b)_{b \in B}$ be unimodular scalars such that the operators

$$U_y = \sum_{b \in B} \mu_b \Pi_{y,b} \quad \text{and} \quad S_y := \sum_{b \in B} \mu_b Q_{y,b}$$

are unitary. Similarly, let $(\nu_a)_{a \in A}$ be unimodular scalars such that the operators

$$V_x := \sum_{a \in A} \nu_a \Xi_{x,a} \quad \text{and} \quad T_x := \sum_{a \in A} \nu_a P_{x,a}$$

are unitary. Equations (16) and (17) imply

$$(18) \quad \sum_{i \in I} \alpha_i (U_y |\phi_i\rangle \otimes |\psi_i\rangle) = \sum_{i \in I} \alpha_i (|\phi_i\rangle \otimes S_y |\psi_i\rangle)$$

and

$$(19) \quad \sum_{i \in I} \alpha_i (|\phi_i\rangle \otimes V_x |\psi_i\rangle) = \sum_{i \in I} \alpha_i (T_x |\phi_i\rangle \otimes |\psi_i\rangle).$$

Fix a Schmidt coefficient α for $|\xi\rangle$ and set $I^\alpha = \{i \in I : \alpha_i = \alpha\}$. Define $\mathcal{K}_A^\alpha := \text{span}\{|\phi_i\rangle : i \in I^\alpha\}$ and $\mathcal{K}_B^\alpha := \text{span}\{|\psi_i\rangle : i \in I^\alpha\}$. By the uniqueness of the Schmidt decomposition, (16) and (17), we deduce that

$$\text{span}\{U_y |\phi_i\rangle : i \in I^\alpha\} = \text{span}\{T_x |\phi_i\rangle : i \in I^\alpha\} = \mathcal{K}_A^\alpha$$

and

$$\text{span}\{U_x |\psi_i\rangle : i \in I^\alpha\} = \text{span}\{S_y |\psi_i\rangle : i \in I^\alpha\} = \mathcal{K}_B^\alpha.$$

Therefore, \mathcal{K}_A^α is $\Pi_{y,b}$ -invariant and $P_{x,a}$ -invariant for every $x \in X, y \in Y, b \in B, a \in A$. Similarly, \mathcal{K}_B^α is $\Xi_{x,a}$ -invariant and $Q_{y,b}$ -invariant for every $x \in X, y \in Y, b \in B, a \in A$.

Let $P_{x,a}^\alpha$ and $\Pi_{y,b}^\alpha$ be the restriction of $P_{x,a}$ and $\Pi_{y,b}$, respectively, to \mathcal{K}_A^α , which we views as operators on \mathcal{K}_A^α . Define $Q_{y,b}^\alpha, \Xi_{x,a}^\alpha \in \mathcal{B}(\mathcal{K}_B^\alpha)$ similarly. Let also

$$|\xi^\alpha\rangle = \frac{1}{\sqrt{|I^\alpha|}} \sum_{i \in I^\alpha} |\phi_i\rangle \otimes |\psi_i\rangle$$

be the maximally entangled vector in $\mathcal{K}_A^\alpha \otimes \mathcal{K}_B^\alpha$. We have that

$$(\Pi_{y,b}^\alpha \otimes I) |\xi^\alpha\rangle = (I \otimes Q_{y,b}^\alpha) |\xi\rangle, \quad y \in Y, b \in B.$$

Clearly, $\sum_{b \in B} \Pi_{y,b}^\alpha$ is the identity of \mathcal{K}_A^α . If $\lambda(x, y, a, b) = 0$ then

$$0 = (P_{x,a}^\alpha \otimes Q_{y,b}^\alpha) |\xi^\alpha\rangle = (P_{x,a}^\alpha \Pi_{y,b}^\alpha \otimes I) |\xi^\alpha\rangle = (P_{x,a}^\alpha \Pi_{y,b}^\alpha \otimes I) |\xi^\alpha\rangle$$

and hence $P_{x,a}^\alpha \Pi_{y,b}^\alpha = 0$ by [6, Lemma 2]. This shows that the assignment $p_{x,a} \mapsto P_{x,a}^\alpha$ and $q_{y,b} \mapsto \Pi_{y,b}^\alpha$ defines an unital *-homomorphism $\pi^\alpha : C^*(\mathcal{G}) \rightarrow \mathcal{B}(\mathcal{K}_A^\alpha)$. Define τ^α to be the canonical tracial state of $\mathcal{B}(\mathcal{K}_A^\alpha)$, and observe that $\tau^\alpha(T) = \langle \xi^\alpha | T \otimes I | \xi^\alpha \rangle$ for $T \in \mathcal{B}(\mathcal{K}_A^\alpha)$.

Let (α_n) be an enumeration of the Schmidt coefficients of $|\xi\rangle$. We have

$$\begin{aligned} p(a, b|x, y) &= \langle \xi | P_{x,a} \otimes Q_{y,b} | \xi \rangle \\ &= \sum_{i,j} \alpha_i \bar{\alpha}_j (\langle \phi_j | \otimes \langle \psi_j |) (P_{x,a} \otimes Q_{y,b}) (|\phi_i\rangle \otimes |\psi_i\rangle) \\ &= \sum_n |\alpha_n|^2 \langle \xi^{\alpha_n} | P_{x,a}^{\alpha_n} \otimes Q_{y,b}^{\alpha_n} | \xi^{\alpha_n} \rangle \\ &= \sum_n |\alpha_n|^2 \langle \xi^{\alpha_n} | P_{x,a}^{\alpha_n} \Pi_{y,b}^{\alpha_n} \otimes I | \xi^{\alpha_n} \rangle \\ &= \sum_n |\alpha_n|^2 (\tau^{\alpha_n} \circ \pi^{\alpha_n})(p_{x,a} q_{y,b}). \end{aligned}$$

By [8], we can replace the infinite convex combination above with a finite convex combination. This concludes the proof.

(iv) \Rightarrow (iii) Suppose that \mathcal{F} is a finite-dimensional C*-algebra, τ is a tracial state on \mathcal{F} , and $\pi : C^*(\mathcal{G}) \rightarrow \mathcal{F}$ is a unital *-homomorphism such that $p(a, b|x, y) = (\tau \circ \pi)(p_{x,a} q_{y,b})$ for $(a, b, x, y) \in A \times B \times X \times Y$. After observing that τ is a convex combination of canonical tracial states on matrix algebras, one can proceed as in the proof of (ii) \Rightarrow (i) in Theorem 5.1 to show that $p \in C_{\text{qm}}$.

(iii) \Rightarrow (ii) \Rightarrow (i) hold trivially. \square

Corollary 5.4. *Let \mathcal{G} be an imitation game. The following assertions are equivalent:*

- (i) \mathcal{G} has a perfect spatial quantum strategy;
- (ii) \mathcal{G} has a perfect quantum strategy;
- (iii) \mathcal{G} has a perfect quantum strategy using a maximally entangled vector;
- (iv) the C*-algebra of the game $C^*(\mathcal{G})$ is nonzero, and it has a nondegenerate finite-dimensional representation.

In view of Proposition 4.10, Corollary 5.2 recovers [5, Theorem 5]—see also [6, Section 4]—as a particular case.

We also obtain the following result of [23].

Corollary 5.5. *The set $\mathcal{C}_q^s = \mathcal{C}_q \cap \mathcal{C}^s$ of synchronous quantum correlations is equal to the set $\mathcal{C}_{\text{qs}}^s = \mathcal{C}_{\text{qs}} \cap \mathcal{C}^s$ of synchronous quantum spatial correlations,*

as well as to the convex hull $\mathcal{C}_{\text{qm}}^{\text{s}} = \mathcal{C}_{\text{qm}} \cap \mathcal{C}^{\text{s}}$ of the synchronous quantum correlation defined using a maximally entangled vector.

One can similarly characterise the local perfect strategies for \mathcal{G} in terms of traces on $C^*(\mathcal{G})$. We omit the proof since it follows closely the ideas in the proof of Theorems 5.1 and 5.3.

Theorem 5.6. *Suppose that \mathcal{G} is an imitation game with input sets X, Y and output sets A, B . Let $p : A \times B \times X \times Y \rightarrow [0, 1]$ be a correlation. The following statements are equivalent:*

- (i) $p \in \mathcal{C}_{\text{loc}}(\mathcal{G})$;
- (ii) *the C^* -algebra of the game $C^*(\mathcal{G})$ is nonzero, and there exists a finite-dimensional abelian C^* -algebra \mathcal{F} with a tracial state τ and a unital $*$ -homomorphism $\pi : C^*(\mathcal{G}) \rightarrow \mathcal{F}$ such that $p(a, b|x, y) = (\tau \circ \pi)(p_{xa}q_{yb})$.*

Corollary 5.7. *Let \mathcal{G} be an imitation game. Then \mathcal{G} has a perfect local strategy if and only if there exists a unital $*$ -homomorphism $\pi : C^*(\mathcal{G}) \rightarrow \mathbb{C}$.*

6. MIRROR GAMES

In this section, we consider the subclass of mirror games and provide a different kind of representation of their perfect quantum commuting strategies in terms of traces. The approach is Hilbert-space-free and allows us to characterise the perfect quantum approximate strategies for these games as ones arising from amenable traces, extending significantly the corresponding result for synchronous games from [23].

Let X, Y, A and B be finite sets. Recall that $\mathcal{A}(X, A) = \ell^\infty(A) *_1 \cdots *_1 \ell^\infty(A)$, where the free product is taken $|X|$ times, $(e_{x,a})_{a=1}^{|A|}$ is the canonical basis of x -th copy of $\ell^\infty(A)$, and $\mathcal{S}_{X,A} = \text{span}\{e_{x,a} : x \in X, a \in A\}$. The canonical generators of $\mathcal{A}(Y, B)$ are denoted by $f_{y,b}$, and their span is denoted by $\mathcal{S}_{Y,B}$.

Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a non-local game. For $x \in X, y \in Y, a \in A$ and $b \in B$, set

$$E_{x,y}^a = \{b \in B : \lambda(x, y, a, b) = 1\} \quad \text{and} \quad E_{x,y}^b = \{a \in A : \lambda(x, y, a, b) = 1\}.$$

Recall that \mathcal{G} is a *mirror game* if there exist functions $\xi : X \rightarrow Y$ and $\eta : Y \rightarrow X$ such that, of every $x \in X$, we have

$$E_{x,\xi(x)}^a \cap E_{x,\xi(x)}^{a'} = \emptyset, \quad a, a' \in A, \quad a \neq a',$$

and

$$E_{\eta(y),y}^b \cap E_{\eta(y),y}^{b'} = \emptyset, \quad b, b' \in B, \quad b \neq b'.$$

In the statement of the following theorem, we use the correspondence $s \rightarrow p_s$ between states and non-signalling families highlighted in Section 3.

Theorem 6.1. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a mirror game, $p \in \mathcal{C}_{\text{qc}}(\lambda)$ and $s \in S(\mathcal{A}(X, A) \otimes_{\text{max}} \mathcal{A}(Y, B))$ be such that $p = p_s$. Then*

(i) the functional $\tau : \mathcal{A}(X, A) \rightarrow \mathbb{C}$ given by $\tau(z) = s(z \otimes 1)$, $z \in \mathcal{A}(X, A)$, is a trace;

(ii) there exists a unital $*$ -homomorphism $\rho : \mathcal{A}(Y, B) \rightarrow \mathcal{A}(X, A)$ such that

$$(20) \quad p(a, b|x, y) = \tau(e_{x,a}\rho(f_{y,b})), \quad x \in X, y \in Y, a \in A, b \in B,$$

and

$$(21) \quad s(z \otimes f_{y_1, b_1} \cdots f_{y_k, b_k}) = \tau(z\rho(f_{y_k, b_k} \cdots f_{y_1, b_1})),$$

for all $z \in \mathcal{A}(A, X)$, $k \in \mathbb{N}$, $y_i \in Y, b_i \in B$, $i = 1, \dots, k$.

Proof. (i) We assume first that

$$(22) \quad \cup_{a \in A} E_{x, \xi(x)}^a = B \text{ and } \cup_{b \in B} E_{\eta(y), y}^b = A, \quad x \in X, y \in Y.$$

For $x \in X, y \in Y, a \in A$ and $b \in B$, let

$$p_{x,a} = \sum_{b \in E_{x, \xi(x)}^a} f_{\xi(x), b}, \quad q_{y,b} = \sum_{a \in E_{\eta(y), y}^b} e_{\eta(y), a}.$$

By (22), $\sum_{b \in B} q_{y,b} = 1$, for all $y \in Y$, and the universal property of $\mathcal{A}(Y, B)$ implies that the assignment $f_{y,b} \rightarrow q_{y,b}$, $y \in Y, b \in B$, extends to a unital $*$ -homomorphism $\rho : \mathcal{A}(Y, B) \rightarrow \mathcal{A}(X, A)$.

If $u_1, u_2 \in \mathcal{A}(X, A) \otimes_{\max} \mathcal{A}(Y, B)$, write $u_1 \sim u_2$ if $s(u_1 - u_2) = 0$. Clearly, \sim is an equivalence relation. Fix $x \in X$ and $a \in A$. Then

$$s(e_{x,a} \otimes 1) = \sum_{b \in B} s(e_{x,a} \otimes f_{\xi(x), b}) = \sum_{b \in E_{x, \xi(x)}^a} s(e_{x,a} \otimes f_{\xi(x), b}) = s(e_{x,a} \otimes p_{x,a}).$$

On the other hand, if $a' \neq a$ then $E_{x, \xi(x)}^{a'} \cap E_{x, \xi(x)}^a = \emptyset$ and so $s(e_{x,a'} \otimes f_{\xi(x), b}) = 0$ whenever $b \in E_{x, \xi(x)}^a$, implying

$$s(e_{x,a'} \otimes p_{x,a}) = \sum_{b \in E_{x, \xi(x)}^a} s(e_{x,a'} \otimes f_{\xi(x), b}) = 0.$$

It follows that

$$s(1 \otimes p_{x,a}) = \sum_{a' \in A} s(e_{x,a'} \otimes p_{x,a}) = s(e_{x,a} \otimes p_{x,a}).$$

Thus,

$$e_{x,a} \otimes 1 \sim e_{x,a} \otimes p_{x,a} \sim 1 \otimes p_{x,a}, \quad x \in X, a \in A.$$

Write $h_{x,a} = e_{x,a} \otimes 1 - 1 \otimes p_{x,a}$. Clearly, $h_{x,a}$ is selfadjoint and

$$h_{x,a}^2 = e_{x,a} \otimes 1 - e_{x,a} \otimes p_{x,a} - e_{x,a} \otimes p_{x,a} + 1 \otimes p_{x,a};$$

thus, $h_{x,a}^2 \sim 0$. The Cauchy-Schwarz inequality now implies

$$(23) \quad uh_{x,a} \sim 0 \text{ and } h_{x,a}u \sim 0, \quad x \in X, a \in A, u \in \mathcal{A}(X, A) \otimes_{\max} \mathcal{A}(Y, B).$$

In particular,

$$(24) \quad ze_{x,a} \otimes 1 \sim z \otimes p_{x,a} \sim e_{x,a}z \otimes 1, \quad x \in X, a \in A, z \in \mathcal{A}(X, A).$$

Similarly, setting $h_{y,b} = q_{y,b} \otimes 1 - 1 \otimes f_{y,b}$, where $y \in Y$ and $b \in B$, we obtain $h_{y,b}^2 \sim 0$, and therefore

$$(25) \quad zq_{y,b} \otimes 1 \sim z \otimes f_{y,b} \sim q_{y,b}z \otimes 1, \quad y \in Y, b \in B, z \in \mathcal{A}(X, A),$$

and

$$(26) \quad zq_{y,b} \otimes w \sim z \otimes wf_{y,b}, \quad y \in Y, b \in B, z \in \mathcal{A}(X, A), w \in \mathcal{A}(Y, B).$$

It is clear that τ is a state on $\mathcal{A}(X, A)$. Let z and w be words on the set $\mathcal{E} := \{e_{x,a} : x \in X, a \in A\}$. We show by induction on the length $|w|$ of w that

$$(27) \quad zw \otimes 1 \sim wz \otimes 1.$$

In the case $|w| = 1$, the claim reduces to (24). Suppose (27) holds if $|w| \leq n - 1$. Let $|w| = n$ and write $w = w'e$, where $e \in \mathcal{E}$. Then, using (24), we have

$$zw \otimes 1 = zw'e \otimes 1 \sim ezw' \otimes 1 \sim w'ez \otimes 1 = wz \otimes 1.$$

From (27) and the fact that the set of all linear combinations of words on \mathcal{E} is dense in \mathcal{A} , we conclude that τ is a trace on $\mathcal{A}(X, A)$.

Now assume that the conditions from (22) are not fulfilled. Choose $a_0 \in A$ and $b_0 \in B$, and define $p_{x,a}$, $a \neq a_0$, $q_{y,b}$, $b \neq b_0$, as in (22). Set $E_{y,0} = A \setminus \left(\cup_{b \in B} E_{\eta(y),y}^b \right)$, $F_{x,0} = B \setminus \left(\cup_{a \in A} E_{x,\xi(x)}^a \right)$, and let

$$p_{x,a_0} = \sum_{b \in F_{x,0}} f_{\xi(x),b}, \quad q_{y,b_0} = \sum_{a \in E_{y,0}} e_{\eta(y),a}.$$

The proof thereafter proceeds as before.

(ii) By (26),

$$p(a, b|x, y) = s(e_{x,a} \otimes f_{y,b}) = s(e_{x,a}q_{y,b} \otimes 1) = \tau(e_{x,a}q_{y,b}) = \tau(e_{x,a}\rho(f_{y,b})),$$

for all $x \in X, y \in Y, a \in A, b \in B$. We show (21) by induction on k . If $k = 1$, the claim follows from (26). Assuming validity for up to $k - 1$ terms, using (26) we have

$$\begin{aligned} s(z \otimes f_{y_1,b_1} \cdots f_{y_k,b_k}) &= s(zq_{y_k,b_k} \otimes f_{y_1,b_1} \cdots f_{y_{k-1},b_{k-1}}) \\ &= \tau(z\rho(f_{y_k,b_k})\rho(f_{y_{k-1},b_{k-1}} \cdots f_{y_1,b_1})) \\ &= \tau(z\rho(f_{y_k,b_k}f_{y_{k-1},b_{k-1}} \cdots f_{y_1,b_1})). \end{aligned}$$

□

Remark. Theorem 6.1 is a generalisation of [32, Theorem 5.5], which was concerned with synchronous games, in two directions: it specifies concretely the C*-algebra involved in the strategy representation, and it extends this representation to the wider class of mirror games.

Recall that, if \mathcal{A} is a C*-algebra, its opposite C*-algebra \mathcal{A}^{op} is defined to be the same involutive normed space, whose elements are denoted by a^{op} , $a \in \mathcal{A}$, but equipped with the product given by $a^{\text{op}}b^{\text{op}} = (ba)^{\text{op}}$. The following lemma was established in [23].

Lemma 6.2. *There exists a *-isomorphism $\gamma : \mathcal{A}(X, A) \rightarrow \mathcal{A}(X, A)^{\text{op}}$ such that*

$$\gamma(e_{x_1, a_1} \cdots e_{x_k, a_k}) = (e_{x_k, a_k} \cdots e_{x_1, a_1})^{\text{op}}, \quad x_i \in X, a_i \in A, i = 1, \dots, k, k \in \mathbb{N}.$$

Theorem 6.3. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a mirror game and $p \in \mathcal{C}_{\text{ns}}(\lambda)$. The following are equivalent:*

- (i) $p \in \mathcal{C}_{\text{qc}}(\lambda)$;
- (ii) *there exist a trace $\tau : \mathcal{A}(X, A) \rightarrow \mathbb{C}$ and a unital *-homomorphism $\rho : \mathcal{A}(Y, B) \rightarrow \mathcal{A}(X, A)$ with $\rho(\mathcal{S}_{X, A}) \subseteq \mathcal{S}_{Y, B}$ such that*

$$p(a, b|x, y) = \tau(e_{x, a} \rho(f_{y, b})), \quad x \in X, y \in Y, a \in A, b \in B.$$

Proof. (i) \Rightarrow (ii) follows from Theorem 6.1 and its proof, using the fact that the projections $q_{y, b}$ defined therein lie in $\mathcal{S}_{X, A}$.

(ii) \Rightarrow (i) Let $\phi : \mathcal{A}(X, A) \otimes \mathcal{A}(X, A)^{\text{op}} \rightarrow \mathbb{C}$ be the bilinear form given by

$$\phi(z \otimes w^{\text{op}}) = \tau(zw), \quad z, w \in \mathcal{A}(X, A).$$

It is well-known that ϕ extends to a state on $\mathcal{A}(X, A) \otimes_{\text{max}} \mathcal{A}(X, A)^{\text{op}}$ (this can be seen, for example, by noting that ϕ is unital and jointly completely positive, and using results from [21]). Let $s : \mathcal{A}(X, A) \otimes_{\text{max}} \mathcal{A}(Y, B) \rightarrow \mathbb{C}$ be the linear functional defined by

$$s = \phi \circ (\text{id} \otimes \gamma) \circ (\text{id} \otimes \rho).$$

Then s is a state on $\mathcal{A}(X, A) \otimes_{\text{max}} \mathcal{A}(Y, B)$, and

$$s(e_{x, a} \otimes f_{y, b}) = \phi(e_{x, a} \otimes \gamma(\rho(f_{y, b}))) = \phi(e_{x, a} \otimes \rho(f_{y, b})^{\text{op}}) = \tau(e_{x, a} \rho(f_{y, b})).$$

By Corollary 3.2, $p \in \mathcal{C}_{\text{qc}}(\lambda)$. \square

For the next theorem, we will need the notion of an amenable trace; we refer the reader to [2] for its many equivalent definitions and properties. The definition that we shall use is that a trace τ on a C*-algebra \mathcal{A} is *amenable* if and only if the map on the algebraic tensor product $\phi : \mathcal{A} \otimes \mathcal{A}^{\text{op}} \rightarrow \mathbb{C}$ given by $\phi(a \otimes b^{\text{op}}) = \tau(ab)$ is bounded with respect to the minimal C*-tensor norm.

Theorem 6.4. *Let $\mathcal{G} = (X, Y, A, B, \lambda)$ be a mirror game and $p \in \mathcal{C}_{\text{ns}}(\lambda)$.*

The following are equivalent:

- (i) $p \in \mathcal{C}_{\text{qa}}(\lambda)$;
- (ii) *there exist an amenable trace $\tau : \mathcal{A}(X, A) \rightarrow \mathbb{C}$ and a unital *-homomorphism $\rho : \mathcal{A}(Y, B) \rightarrow \mathcal{A}(X, A)$ with $\rho(\mathcal{S}_{X, A}) \subseteq \mathcal{S}_{Y, B}$ such that*

$$p(a, b|x, y) = \tau(e_{x, a} \rho(f_{y, b})), \quad x \in X, y \in Y, a \in A, b \in B.$$

Proof. (ii) \Rightarrow (i) Let $\phi : \mathcal{A}(X, A) \otimes \mathcal{A}(X, A)^{\text{op}} \rightarrow \mathbb{C}$ be the bilinear form given by

$$\phi(z \otimes w^{\text{op}}) = \tau(zw), \quad z, w \in \mathcal{A}(X, A).$$

Since τ is amenable, ϕ extends to a state on $\mathcal{A}(X, A) \otimes_{\text{min}} \mathcal{A}(X, A)^{\text{op}}$ [2, Theorem 6.2.7]. Define a state s on $\mathcal{A}(X, A) \otimes_{\text{min}} \mathcal{A}(Y, B) \rightarrow \mathbb{C}$ by letting

$$s = \phi \circ (\text{id} \otimes \gamma) \circ (\text{id} \otimes \rho).$$

As in the proof of Theorem 6.3,

$$p(a, b|x, y) = s(e_{x,a} \otimes f_{y,b}), \quad x \in X, y \in Y, a \in A, b \in B;$$

by Corollary 3.3, $p \in \mathcal{C}_{\text{qa}}(\lambda)$.

(i) \Rightarrow (ii) Since $p \in \mathcal{C}_{\text{qa}}(\lambda)$, by Corollary 3.3, there exists a state s on $\mathcal{A}(X, A) \otimes_{\min} \mathcal{A}(Y, B)$ such that

$$p(a, b|x, y) = s(e_{x,a} \otimes f_{y,b}), \quad x \in X, y \in Y, a \in A, b \in B.$$

Arguing as in the proof of Theorem 6.1 and using (24), we may obtain a unital *-homomorphism $\rho : \mathcal{A}(Y, B) \rightarrow \mathcal{A}(X, A)$ with $\rho(\mathcal{S}_{Y,B}) \subseteq \mathcal{S}_{X,A}$ and a unital *-homomorphism $\pi : \mathcal{A}(X, A) \rightarrow \mathcal{A}(Y, B)$ with $\pi(\mathcal{S}_{X,A}) \subseteq \mathcal{S}_{Y,B}$ such that:

- (a) the functional $\tau : \mathcal{A}(X, A) \rightarrow \mathbb{C}$, given by $\tau(z) = s(z \otimes 1)$, is a trace;
- (b) $s(z \otimes \pi(e_{x_1, a_1} \cdots e_{x_k, a_k})) = \tau(ze_{x_k, a_k} \cdots e_{x_1, a_1})$, for $z \in \mathcal{A}(A, X)$, $x_i \in X$, $a_i \in A$, $i = 1, \dots, k$, and
- (c) $p(a, b|x, y) = \tau(e_{x,a} \rho(f_{y,b}))$, for $x \in X$, $y \in Y$, $a \in A$, $b \in B$.

Let $\phi : \mathcal{A}(X, A) \otimes_{\min} \mathcal{A}(X, A)^{\text{op}} \rightarrow \mathbb{C}$ be the state defined by letting

$$\phi = s \circ (\text{id} \otimes \pi) \circ (\text{id} \otimes \gamma^{-1}).$$

Let $z \in \mathcal{A}(X, A)$ and $w = e_{x_1, a_1} \cdots e_{x_k, a_k}$, for some $x_i \in X$, $a_i \in A$, $i = 1, \dots, k$. Set $\bar{w} := \gamma^{-1}(w^{\text{op}}) = e_{x_k, a_k} \cdots e_{x_1, a_1}$. Thus, using (b) we have

$$\begin{aligned} \phi(z \otimes w^{\text{op}}) &= s(z \otimes \pi(\bar{w})) = s(z \otimes \pi(e_{x_k, a_k} \cdots e_{x_1, a_1})) \\ &= \tau(ze_{x_1, a_1} \cdots e_{x_k, a_k}) = \tau(zw). \end{aligned}$$

By linearity and continuity,

$$\phi(z \otimes w^{\text{op}}) = \tau(zw), \quad z, w \in \mathcal{A}(X, A).$$

By [2, Theorem 6.2.7], τ is amenable. \square

Acknowledgement. Part of this research was conducted during two Focused Research Meetings, funded by the Heilbronn Institute, and hosted at Queen's University Belfast in October 2016 and March 2017. M. Lupini was partially supported by the NSF grant DMS-1600186. G. Scarpa acknowledges the support of MTM2014-54240-P (MINECO), QUITEMAD+-CM Reference: S2013/ICE-2801 (Comunidad de Madrid), ICMAT Severo Ochoa project SEV-2015-0554 (MIN-ECO), and grant 48322 from the John Templeton Foundation. The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation.

REFERENCES

- [1] A. ATSERIAS, L. MANČINSKA, D. E. ROBERSON, R. ŠÁMAL, S. SEVERINI AND A. VARVITSIOTIS, *Quantum and non-signalling graph isomorphisms*, preprint (2016), arXiv:1611.09837v3.
- [2] N. P. BROWN AND N. OZAWA, *C*-algebras and finite-dimensional approximations*, American Mathematical Society, 2008.

- [3] P. J. CAMERON, A. MONTANARO, M. W. NEWMAN, S. SEVERINI AND A. WINTER, *On the quantum chromatic number of a graph*, Electronic J. Combinatorics 14 (2007), no. 1, Paper 81, 15 pp.
- [4] M.D. CHOI AND E.G. EFFROS, *Injectivity and operator spaces*, J. Funct. Anal. 24 (1977), 156-209.
- [5] R. CLEVE, L. LIU AND W. SLOFSTRA, *Perfect commuting-operator strategies for linear system games*, preprint (2016), arXiv:1606.02278.
- [6] R. CLEVE AND R. MITTAL, *Characterization of binary constraint system games*, Automata, languages, and programming. Part I, Lecture Notes in Comput. Sci., vol. 8572, Springer (2014), 320-331.
- [7] A. CONNES, *Classification of injective factors*, Ann. of Math. (2) 104 (1976), 73-115.
- [8] W. D. COOK AND R. J. WEBSTER, *Carathéodory's theorem*, Canad. Math. Bull. 15 (1972), 293-293.
- [9] T. S. CUBITT, D. LEUNG, W. MATTHEWS, AND A. WINTER, *Improving zero-error classical communication with entanglement*, Phys. Rev. Lett. **104** (2010), 230503.
- [10] K. DYKEMA, V. I. PAULSEN, J. PRAKASH, *The Delta game*, preprint (2017), arXiv:1707.06186.
- [11] K. DYKEMA, V. I. PAULSEN, J. PRAKASH, *Non-closure of the set of quantum correlations via graphs*, preprint (2017), arXiv:1709.05032.
- [12] E. G. EFFROS AND ZH.-J. RUAN, *Operator spaces*, Oxford University Press, 2000.
- [13] D. FARENICK, A. KAVRUK, V. I. PAULSEN AND I. G. TODOROV, *Operator systems from discrete groups*, Comm. Math. Phys. 329 (2014), 207-238.
- [14] D. FARENICK, A. KAVRUK, V. I. PAULSEN AND I. G. TODOROV, *Characterizations of the weak expectation property*, New York J. Math., to appear.
- [15] D. FARENICK AND V. I. PAULSEN, *Operator system quotients of matrix algebras and their tensor products*, Math. Scand. 111 (2012), 210-243.
- [16] T. FRITZ, *Operator system structures on the unital direct sum of C^* -algebras*, Rocky Mountain J. Math. 44 (2014), no. 3, 913-936.
- [17] J. W. HELTON, K. P. MEYER, V. I. PAULSEN AND M. SATRIANO, *Algebras, synchronous games, and chromatic numbers of graphs*, preprint (2017), arXiv:1703.00960.
- [18] A. S. HOLEVO, *Quantum systems, channels, information*, De Gruyter, 2013.
- [19] M. JUNGE, M. NAVASCUES, C. PALAZUELOS, D. PEREZ-GARCIA, V. B. SCHOLTZ AND R. F. WERNER, *Connes' embedding problem and Tsirelson's problem*, J. Math. Physics 52 (2011), 012102.
- [20] A. S. KAVRUK, *Nuclearity related properties in operator systems*, J. Operator Theory 71 (2014), no. 1, 95-156.
- [21] A. KAVRUK, V. I. PAULSEN, I. G. TODOROV AND M. TOMFORDE, *Tensor products of operator systems*, J. Funct. Anal. 261 (2011), no. 2, 267-299.
- [22] A. S. KAVRUK, V. I. PAULSEN, I. G. TODOROV, AND M. TOMFORDE, *Quotients, exactness, and nuclearity in the operator system category*, Adv. Math. 235 (2013), 321-360.
- [23] S.-J. KIM, V. I. PAULSEN AND C. SCHAFHAUSER, *A synchronous game for binary constraint systems*, preprint (2017), arXiv:1707.01016.
- [24] L. LAMI, *Non-classical correlations in quantum mechanics and beyond*, PhD thesis, arXiv:1803.02902.
- [25] L. MANČINSKA AND D. E. ROBERSON, *Quantum homomorphisms*, J. Combin. Theory Ser. B, 118 (2016), 228-267.
- [26] L. MANČINSKA AND T. VIDICK, *Unbounded entanglement in nonlocal games*, Quantum Inf. Comput. 15 (2015), no. 15-16, 1317-1332.
- [27] D. E. ROBERSON, *Variations on a theme: Graph homomorphisms*, Ph.D. thesis, University of Waterloo, 2013.
- [28] C. M. ORTIZ AND V. I. PAULSEN, *Quantum graph homomorphisms via operator systems*, Linear Algebra App. 497 (2016), 23-43.

- [29] N. OZAWA, *About the Connes' embedding problem—algebraic approaches*, Japan. J. Math. 8 (2013), no. 1, 147-183.
- [30] C. PALAZUELOS AND T. VIDICK, *Survey on nonlocal games and operator space theory*, J. Math. Phys. 57 (2016), no. 1, 015220.
- [31] V. I. PAULSEN, *Completely bounded maps and operator algebras*, Cambridge University Press, 2002.
- [32] V. I. PAULSEN, S. SEVERINI, D. STAHLKE, I. G. TODOROV AND A. WINTER, *Estimating quantum chromatic numbers*, J. Funct. Anal. 270 (2016), no. 6, 2188-2222.
- [33] V. I. PAULSEN AND I. G. TODOROV, *Quantum chromatic numbers via operator systems*, Q. J. Math. 66 (2015), no. 2, 677-692.
- [34] G. PISIER, *Introduction to operator space theory*, Cambridge University Press, 2003.
- [35] M. V. RAMANA, E. R. SCHEINERMAN AND D. ÜLLMAN, *Fractional isomorphism of graphs*, Discrete Math. 132 (1994), no. 1-3, 247-265.
- [36] A. RAO, *Parallel repetition in projection games and a concentration bound*, SIAM J. Comput. 40 (2011), no. 6, 1871-1891.
- [37] W. SLOFSTRA, *Tsirelson's problem and an embedding theorem for groups arising from non-local games*, preprint (2016), arXiv:1606.03140.
- [38] W. SLOFSTRA, *The set of quantum correlations is not closed*, preprint (2017), arXiv:1703.08618.
- [39] W. SLOFSTRA AND T. VIDICK, *Entanglement in non-local games and the hyperlinear profile of groups*, preprint (2017), arXiv:1711.10676.
- [40] B. S. TSIRELSON, *Quantum generalizations of Bell's inequality*, Lett. Math. Phys. 4 (1980), no. 4, 93-100.
- [41] B. S. TSIRELSON, *Some results and problems on quantum Bell-type inequalities*, Hadronic J. Suppl. 8 (1993), no. 4, 329-345.

MATHEMATICS DEPARTMENT, CALIFORNIA INSTITUTE OF TECHNOLOGY, 1200 EAST CALIFORNIA BOULEVARD, MAIL CODE 253-37, PASADENA, CA 91125

Current address: School of Mathematics and Statistics, Victoria University of Wellington, PO Box 600, Wellington 6140, New Zealand

E-mail address: martino.lupini@vuw.ac.nz

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN 5, 2100 COPENHAGEN, DENMARK

E-mail address: mancinska@math.ku.dk

INSTITUTE FOR QUANTUM COMPUTING AND DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, CANADA

E-mail address: vpaulsen@uwaterloo.ca

DEPARTMENT OF PHYSICS, TECHNICAL UNIVERSITY OF DENMARK, RICHARD PETERSENS PLADS, COPENHAGEN, DENMARK

E-mail address: davidroberson@gmail.com

FACULTY OF MATHEMATICAL SCIENCES, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA DE LAS CIENCIAS, 3, DPCHO. 489, 28040 MADRID, SPAIN

E-mail address: giannicolascarpa@gmail.com

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON WC1E 6BT, UNITED KINGDOM, AND INSTITUTE OF NATURAL SCIENCES, SHANGHAI JIAO TONG UNIVERSITY, 200240 SHANGHAI, CHINA

E-mail address: s.severini@ucl.ac.uk

MATHEMATICAL SCIENCES RESEARCH CENTRE, QUEEN'S UNIVERSITY BELFAST, BELFAST BT7 1NN, UNITED KINGDOM, AND SCHOOL OF MATHEMATICAL SCIENCES, NANKAI UNIVERSITY, 300071 TIANJIN, CHINA

E-mail address: i.todorov@qub.ac.uk

ICREA AND FÍSICA TEÒRICA: INFORMACIÓ I FENOMENS QUÀNTICS, UNIVERSITAT AUTÒNOMA DE BARCELONA, ES-08193 BELLATERRA, BARCELONA, SPAIN

E-mail address: andreas.winter@uab.cat