

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

The value of dermoscopy of the nail plate free edge and hyponychium

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

lorizzo, M., Starace, M., Di Altobrando, A., Alessandrini, A., Veneziano, L., Piraccini, B.M. (2021). The value of dermoscopy of the nail plate free edge and hyponychium. JOURNAL OF THE EUROPEAN ACADEMY OF DERMATOLOGY AND VENEREOLOGY, 35(12), 2361-2366 [10.1111/jdv.17521].

Availability:

This version is available at: https://hdl.handle.net/11585/871187 since: 2022-02-27

Published:

DOI: http://doi.org/10.1111/jdv.17521

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

DR. MICHELA STARACE (Orcid ID : 0000-0002-3981-1527)
DR. AMBRA DI ALTOBRANDO (Orcid ID : 0000-0003-3236-8822)
DR. AURORA ALESSANDRINI (Orcid ID : 0000-0003-2895-4861)
DR. BIANCA MARIA PIRACCINI (Orcid ID : 0000-0001-6537-9689)

Article type : Review Article

#### JEADV-2021-0208

# The value of dermoscopy of the nail plate free edge and hyponychium Matilde Iorizzo<sup>1</sup>, Michela Starace<sup>2</sup>, Ambra di Altobrando<sup>2</sup>, Aurora Alessandrini<sup>2</sup>, Leonardo Veneziano<sup>2</sup>, Bianca Maria Piraccini<sup>2</sup>

Private Dermatology Practice, Bellinzona, Switzerland.

<sup>2</sup> Dermatology- IRCCS Policlinico Sant'Orsola, Department of Experimental, Diagnostic Specialty Medicine (DIMES), *Alma Mater Studiorum*, University of Bologna, Italy

Manuscript word count: 1744 Abstract word count: 102 References: 27 Tables: 1 Figures: 1 Keywords: dermoscopy, onychoscopy, nail plate free edge, hyponychium, melanonychia, onycholysis, onychomycosis, psoriasis, capillaries, subungual mass.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the <u>Version of Record</u>. Please cite this article as <u>doi:</u> <u>10.1111/jdv.17521</u>

Funding sources: None declared.

Conflict of interests: None declared.

Acknowledgement: The patients in this manuscript have given written informed consent to the publication of their case details

#### **Corresponding Author:**

Michela Starace, MD, PhD Dermatology Unit, Policlinico Sant'Orsola Department of Specialized, Experimental and Diagnostic Medicine, University of Bologna Via Massarenti, 1, 40138 Bologna, Italy e-mail: michela.starace2@ unibo.it phone: +390512143677 fax: +39051214486

# ABSTRACT

The non-invasive examination of the nail unit using a dermoscope is known as onychoscopy. This technique has become increasingly appreciated to facilitate the clinical diagnosis of nail disorders, opening up a valuable second front with a potential to avoid invasive diagnostic procedures. During a nail consultation, the nail unit should always be examined with

the aid of a dermatoscope in all its components. The aim of this paper review is to provide practical information about onychoscopy of the nail plate free edge and hyponychium, two components of the nail unit difficult to evaluate at naked eye and often forgotten, but of paramount importance.

### **INTRODUCTION**

The free edge, the thickest part of the nail plate, grows beyond the fingertip, protecting it. Frontally, it appears as a slightly curve and flexible transverse band, approximately 0.5 - 1 mm thick, of a white yellowish color and made up of three layers corresponding to the dorsal nail plate (produced by the proximal portion of the nail matrix), the intermediate nail plate (produced by the distal matrix) and the ventral nail plate (produced by the nail bed)<sup>1</sup>.

Under the free edge lies the hyponychium, a 0.1 - 0.2 mm cutaneous margin that continues the nail bed and corresponds to the site where the nail plate detaches from the bed. The hyponychium is proximally margined by the onychodermal band and distally by the distal groove, a cutaneous ridge demarcating the border between subungual structures and the finger pulp<sup>2-4</sup>. Together with the free edge, the hyponychium forms a crevice, well known reservoir for different microbes often responsible for nail plate secondary changes<sup>5</sup>. Both the hyponychium and the free edge can be however hit by different primary dermatological conditions, often overlooked because this anatomic area is difficult to evaluate by naked eye. Things may change with the use of a dermoscope.

Onychoscopy has become increasingly appreciated in the last years to facilitate the clinical diagnosis of nail disorders, opening up a valuable second front with a potential to avoid invasive diagnostic methods<sup>6-8</sup>. Enabling easier and closer observation of nail features not visible to naked eye, onychoscopy is a valuable aid not only in revealing cryptic details of diagnostic value, but also in monitoring the evolution of a disease and response to treatment through stored images<sup>9</sup>. The aim of this paper is then to provide practical information about onychoscopy of the nail plate free edge and hyponychium, two components of the nail unit often forgotten, but of paramount importance. A practical table with the most common findings will also be provided according to authors' experience, supported and confirmed by the existing literature listed as references of this paper.

#### METODOLOGY

The unique anatomy of the nail apparatus makes onychoscopy a difficult technique to be performed, in every site. The nail plate, in fact, is a convex structure that doesn't allow to a flat and rigid lens to adhere perfectly. When observing the free edge of the nail plate and hyponychium, the lens should be placed on the nail plate free edge, pushing back the fingertip. The dermoscope should then be moved up and down and from side to side. An initial dry observation (dry dermoscopy technique) can be followed by the application of an interface medium (wet dermoscopy technique). Given the nail plate small size, convexity and hardness, which prevent a complete adherence of the lens to the surface, the dermoscope needs to be tilted and an interface medium with high viscosity should be used, in order not to roll off/slide off the plate. Water-based gels (e.g., ultrasound gel), antiseptic gels or oils serve this purpose very well. However, ultrasound gel is probably the best option because it doesn't flow<sup>10</sup>. The lowest magnifications, i.e. 10X to 30X, are the ones to start with: the nail plate free edge, subungual contents and digital creases are easily seen. Higher magnifications (50X-200X) detect

instead the detailed capillary architecture of the hyponychium.

The patient undergoing the exam must initially be seated comfortably with the hand or foot to be examined placed on a flat surface, since the whole procedure may take 15-20 minutes. A grey or black backdrop is always preferrable because the colour absorbs the excess of light without reflecting it <sup>4</sup>. Lighter backdrop is instead better for darker phototypes.

Regarding the choice of the device, videodermoscopes are versatile devices capable of providing high magnifications, though resolution may be suboptimal. On the other hand, handheld dermoscopes provide much lower magnification, but clarity. Nonmore contact (polarized) handheld dermoscopes may also be used, proving very useful in visualizing deep structures and capillary architecture without pressing or blanching vessels<sup>11,12</sup>. Both devices, non-polarized and polarized, should however be used when examining the nail plate free edge and hyponychium to perform a complete evaluation of the disease.

#### **INTERPRETATION**

As stated, specific disorders affecting these areas require the use of the dermatoscope as mandatory.

Considering that these nail structures are able to respond only with a limited number of reaction patterns to the number of disorders affecting them, collecting diagnostic clues can be extremely challenging. Knowledge of nail anatomy and the most common nail disorders are mandatory

before starting onychoscopy because it is very important to know where and what to look at. Being the plate a product of the matrix, for example, the free edge can also reveal alterations coming from the matrix.

Lower dermoscopy magnifications are usually enough to visualize the nail plate free edge that, in normal conditions, appears as a white - yellowish upside-down "U", more or less curved, with a thickness regular in all its length. The hyponychium is instead not always visible, as frequently covered by the free edge, but it is seen as a "cul de sac" below the free edge when the nail plate is cut too short or in nail biters. Higher magnifications are needed to visualize the capillary network of this area that appears as a regular pool of red dots: the appearance as dots is due to the perpendicular arrangement of the dermal capillaries to the skin (each dot represents the top of one loop)<sup>13</sup>.

Different conditions may alter the aspect of these areas and dermoscopic observation can often give diagnostic hints to clinical examination that might avoid biopsy or tell us which is the right place to biopsy. Table I reports the most common findings seen at dermoscopy when examining the nail plate free edge and hyponychium affected by a nail disorder. Figure 1 reports the most common disorders affecting these areas.

The free edge may be locally or diffusely thickened or thinned, it may be splitted, fissured or detached and the color may vary. The curvature may also be affected totally or in small areas. When onycholysis is present, the hyponychium becomes more evident. Scales of different thickness and colors may be present under the nail plate and the capillary network might be changed and more visible. A subungual mass of different shape, size and consistency may also be present.

#### **CLINICAL RELEVANCE**

Dermoscopic features of the nail unit should always be observed in all its parts, including the distal nail plate free edge and the hyponychium, because they can reveal important clues for the correct diagnosis. The most important data that the nail plate free edge is able to provide are probably those collected during the evaluation of a longitudinal melanonychia. The color of the nail plate free edge permits first to distinguish between a melanic and a non-melanic pigmentation: the first one is located within the nail plate that is then grey to brown to black, the second is located below or above the nail plate leaving it free from pigment <sup>14</sup>. Once we are dealing with melanonychia, the origin of the pigment is very important because biopsy the distal matrix will

result in a limited risk of scarring, and this is a useful information we can provide to patients to further encourage them before surgery. If the pigment is located in the upper portion of the free edge, the source is likely to be the proximal portion of the matrix; if the pigment is found in the lower portion, it favors a more distal matrix location. Moreover, a dermoscopic evaluation of the hyponychium can reveal a micro-Hutchinson sign, the periungual melanic pigmentation due to tumor spread into adjacent tissues, not visible at naked eye<sup>15-17</sup>. Detecting the micro Hutchinson sign is particularly useful in presence of an ulcerated mass of the nail bed and hyponychium because it might be a sign of amelanotic melanoma. Hyponychial pigmentation can be however observed also in congenital nevi, where it just suggests presence of nest of melanocytes in the periungual skin.

The distal nail edge evaluation is also important in case of single white longitudinal bands of the nail plate. Presence of multiple sharply demarcated honeycomb holes within a locally thickened nail plate is diagnostic for onychomatricoma<sup>18</sup>. The holes correspond to the longitudinal digitations of the tumor excavated within the nail plate.

Enhancing vision of a subungual mass is also important to define the diagnosis and avoid unnecessary procedures. Onychopapilloma is, for instance, almost always diagnosed through dermoscopy and clearly differentiated from a wart or a longitudinal melanonychia in case of pigmentation. Onychopapilloma grows as a focal gray filiform keratotic plug underneath the

free edge of the nail plate, and it is never surrounded by a collarette<sup>19</sup>. A collarette characterizes subungual warts, which also present dotted dilated and intermingled thrombosed vessels (dark redblackish hemorrhagic dots), some of which surrounded by a white halo<sup>20</sup>.

Fibroma and Fibrokeratomas appears as skin-colored papules with a central pale yellowish area more or less hyperkeratotic<sup>21</sup>. Exostosis are hard nodules with possible vascular ectasia or ulceration<sup>22</sup>.

The presence of subungual hyperkeratosis and scales can instead suggest an inflammatory disorder like psoriasis or eczema, but if the scales are yellow-orange or multicolored, with a ruinlike appearance, a diagnosis of onychomycosis should be always taken into account<sup>23,24</sup>. The rare form of onychomycosis called endonyx, not characterized by subungual scales, can instead be easily identified observing the nail plate free edge that shows shiny white areas confined to the nail plate<sup>25</sup>. Usually, subungual hyperkeratosis and scales are associated with onycholysis, but when this presents alone a traumatic etiology can be suspected. The presence of pustules in the hyponychial area can also be detected, facilitating the diagnosis of pustular psoriasis (Hallopeau's acrodermatitis), but differential diagnosis should include herpes infection, which can be ruled out by clinical observation, history and Tzanck smear test.

Evaluation of the vascular architecture is finally very important. The presence of irregularly distributed, dilated, tortuous, and elongated capillaries can be a clue for the definitive diagnosis of nail psoriasis when the nail plate signs are unspecific<sup>26</sup>. Abnormal capillaries are absent in allergic contact dermatitis even when it presents with psoriasiform nail changes<sup>27</sup>. Capillary density is also correlated with nail psoriasis severity and can be a marker of response to treatment.

### CONCLUSIONS

Performing onychoscopy is not easy like skin dermoscopy or trichoscopy, but due to the small and often hidden nail components this technique is always advisable in any patient during a nail consultation and should become a routine to reinforce presumptive clinical diagnoses.

# REFERENCES

- 1. Garson JC, Baltenneck F, Leroy F, et al. Histological structure of human nail as studied by synichroton x-ray microdiffration. *Cell Mol Biol* 2000; **46**:1025-1034.
- 2. Terry RB. The onychodermal band in health and disease. *Lancet* 1955; i:179–181.
- 3. Perrin C. Peculiar zone of the distal nail unit: the nail isthmus. *Am J Dermatopathol* 2007; **29:**108–109.
- 4. de Berker D, Ruben BS, Baran R. The science of the nail apparatus. In: Baran R, de Berker D, Holzberg M, Piraccini BM; Richert B, Thomas L. Baran & Dawber's diseases of the nails and their management (5<sup>th</sup> Edition). Wiley Blackwell, Oxford, 2019:1-58.
- 5. Canavan TN, Graham LV, Elewski BE. Subungual Space: The Next Frontier. *Skin Appendage Disord* 2018; **5**:50-51.
- Piraccini BM, Bruni F, Starace M. Dermoscopy of non-skin cancer nail disorders. *Dermatol Ther* 2012; 25:594-602.
- Alessandrini A, Starace M, Piraccini BM. Dermoscopy in the evaluation of nail disorders. *Skin Appendage Disord* 2017;3:70-82.
- Piraccini BM, Alessandrini A, Starace M. Onychoscopy: Dermoscopy of the Nails. *Dermatol Clin* 2018; 36:431-438.

- 9. Grover C, Jakhar D. Onychoscopy: A practical guide. *Indian J Dermatol Venereol Leprol* 2017;**83:**536-549.
- 10. Gewirtzman AJ, Saurat JH, Braun RP. An evaluation of dermoscopy fluids and application techniques. *Br J Dermatol* 2003; **149:**59-63.
- Benvenuto-Andrade C, Dusza SW, Agero AL, et al. Differences between polarized light dermoscopy and immersion contact dermoscopy for the evaluation of skin lesions. *Arch Dermatol* 2007; 143:329-338.
- Pan Y, Gareau DS, Scope A, Rajadhyaksha M, Mullani NA, Marghoob AA. Polarized and nonpolarized dermoscopy: the explanation for the observed differences. *Arch Dermatol* 2008; 144:828-829.
- Humbert P, Sainthillier JM, Mac-Mary S, et al. Capillaroscopy and videocapillaroscopy assessment of skin microcirculation: dermatologic and cosmetic approaches. *J Cosmet Dermatol* 2005; 4:153-62.
- 14. Braun RP, Baran R, Saurat JH, Thomas L. Surgical Pearl: Dermoscopy of the free edge of the nail to determine the level of nail plate pigmentation and the location of its probable origin in the proximal or distal nail matrix. J Am Acad Dermatol 2006; 55:512-513.
- 15. Ronger S, Touzet S, Ligeron C, et al. Dermoscopic examination of nail pigmentation. *Arch Dermatol* 2002; **138**:1327-1333.
- 16. Starace M, Dika E, Fanti PA, et al. Nail apparatus melanoma: dermoscopic and histopathologic correlations on a series of 23 patients from a single centre. *J Eur Acad Dermatol Venereol* 2018; **32**:164-173.
- Starace M, Alessandrini A, Brandi N, Piraccini BM. Use of Nail Dermoscopy in the Management of Melanonychia: Review. *Dermatol Pract Concept* 2019; 9:38-43.
- 18. Di Chiacchio N, Tavares GT, Tosti A, et al. Onychomatricoma: epidemiological and clinical findings in a large series of 30 cases. *Br J Dermatol* 2015;**173:**1305-1307.
- Tosti A, Schneider SL, Ramirez-Quizon MN, et al. Clinical, dermoscopic, and pathologic features of onychopapilloma: A review of 47 cases. J Am Acad Dermatol 2016;74:521-526.
- 20. Subhadarshani S, Sarangi J, Verma KK. Dermoscopy of Subungual Wart. *Dermatol Pract Concept* 2019; **9:**22-23.

- 21. Ehara Y, Yoshida Y, Ishizu S, Shiomi T, Yamamoto O. Acquired subungual fibrokeratoma. *J Dermatol* 2017; **44:** e140-e141.
- 22. Piccolo V, Argenziano G, Alessandrini AM, et al. Dermoscopy of Subungual Exostosis: A Retrospective Study of 10 Patients. *Dermatology* 2017; **233**:80-85.
- 23. Piraccini BM, Balestri R, Starace M, Rech G. Nail digital dermoscopy (onychoscopy) in the diagnosis of onychomycosis. *J Eur Acad Dermatol Venereol* 2013; **27:**509-513.
- Yorulmaz A, Yalcin B. Dermoscopy as a first step in the diagnosis of onychomycosis. *Postepy Dermatol Alergol* 2018; 35:251-258.
- 25. Jakhar D, Grover C, Kaur I. Dendritic pattern: Unique onychoscopic feature in endonyx. *J Am Acad Dermatol* 2019; **81**:e89-e90.
- 26. Iorizzo M, Dahdah M, Vincenzi C, Tosti A. Videodermoscopy of the hyponychium in nail bed psoriasis. *J Am Acad Dermatol* 2008; **58:**714-715.
- 27. Piccolo V, Piraccini BM, Argenziano G, Russo T, Alessandrini A, Starace M. Onychoscopy of allergic contact dermatitis caused by artificial nails: A double-center retrospective study on 34 patients. *J Am Acad Dermatol* 2020; 83:1485-1487.

# **FIGURE LEGENDS**

**Figure 1**: Dermoscopy of the nail plate free edge and hyponychium: (A) Hutchinson's sign in nail melanoma; (B) onychomatricoma; (C) allergic contact dermatitis; (D) subungual exostosis; (E) longitudinal melanonychia - proliferation – with the pigment coming from the distal matrix; (F) pincer nail; (G) traumatic onycholysis; (H) psoriasis; (I) psoriasis; (L) onyychomycosis; (M) onychopapilloma; (N) wart.

 Table 1: Most common dermoscopic findings observed in nail plate free edge and hyponychium.

	NAIL PLATE FREE EDGE	HYPONYCHIUM
	Thickness: normal	Scales: absent
Longitudinal Melanonychia	Shape: normal	Color: possible presence of Hutchinson sign (also
(proliferation)	Color: pigmentation of the upper portion (PM	micro) and/or vascular nodule with irregular vascular
	involvement)	pattern (dots & irregular lines) in case of Melanoma
	pigmentation of the lower portion (DM involvement)	Capillaries: normal
	pigmentation of the full thickness (PM+DM inv.)	
	Margin: possible splitting/dystrophy in case of	
	Melanoma	
	frequent splitting in case of pediatric nevi	
	Subungual scales/mass: absent	
	Thickness: normal	Scales: absent
Longitudinal Melanonychia	Shape: normal	Color: normal
(activation)	Color: normal to light gray	Capillaries: normal
	Margin: normal	
	Subungual scales/mass: absent	

Table 1

	Thickness: partial or full-length thickening	Scales: absent
Onychomatricoma	Shape: transverse overcurvature	Color: normal
	Color: yellowish	Capillaries: normal
	Margin: honeycomb cavities (sharply demarcated	
	holes)	
	Subungual scales/mass: absent	
	Thickness: Whole length thickening	Scales: absent
Pincer Nails	Shape: overcurvature	Color: normal
	Color: yellowish	Capillaries: normal
	Margin: normal	
	Subungual scales/mass: absent	
	Thickness: Whole length thinning	Scales: absent
Koylonychia	Shape: concave shape	Color: normal
	Color: normal	Capillaries: normal
	Margin: normal	
	Subungual scales/mass: absent	
	Thickness: normal	Scales: absent
Traumatic Onycholisis	Shape: normal	Color: normal

Accepted Article

(acute / chronic)	Color: normal	Capillaries: normal
	Margin: normal	
	Subungual scales/mass: absent	
	Thickness: may be diffusely thickened	Scales: may be present
Psoriasis	Shape: normal	Color: normal
	Color: normal or yellowish	Capillaries: irregularly distributed, dilated, tortuous,
	Margin: normal	and elongated capillaries (best seen at 50x)
	Subungual scales/mass: absent	
	Thickness: normal	Scales: present with skin fissures
Allergic Contact Dermatitis	Shape: normal	Color: erythema, hemorrhages
	Color: normal or yellowish	Capillaries: normal
	Margin: normal	
	Subungual scales/mass: absent	
	<b>Thickness:</b> slightly increased (especially in DLSO)	Scales: irregularly distributed hyperkeratosis with
Onychomycosis	Shape: normal	yellow-orange scales,
	<b>Color:</b> normal, but shiny white in case of endonyx	Color: possible black/multicolored pigment and
	onychomycosis	indentations ("ruin appearance")
	Margin: may be irregular	Capillaries: normal

	Subungual scales/mass: ruin-shaped yellow-white
	orange scales
	Thickness: normal
	Shape: normal
Warts	Color: normal
	Margin: normal
	Subungual scales/mass: keratotic mass in a mosai
	pattern with dotted vessels; presence of a collarette
	limiting the lesion
	Thickness: normal
Onychopapilloma	Shape: normal
	Color: normal
	Margin: fissured
	Subungual scales/mass: absent localized pink-
	brownish filiform keratotic plug
	Of the proximal nail fold:

Scales: may be present

Capillaries: normal

**Color:** normal

Scales: absent

Color: normal

Scales: absent

Capillaries: normal

Fibroma / Fibrokeratoma	Thickness: localized tinning	Color: normal
	Shape: localized concave shape	Capillaries: normal
	Color: normal	
	Margin: fissured	
	Subungual scales/mass: skin colored papule with a	
	central pale yellowish area ( $\pm$ hyperkeratotic),	
	surrounded by a scaly collarette	
	Subungual localization:	
	Thickness: normal	
	Shape: normal	
	Color: normal	
	Margin: normal	
	Subungual scales/mass: skin colored papule with a	
	central pale yellowish area ( $\pm$ hyperkeratotic),	
	surrounded by a scaly collarette	
	Thickness: normal	Scales: absent
Exostosis	Shape: normal	Color: normal
	Color: normal	Capillaries: normal

 Margin: normal

 Subungual scales/mass: hard nodule together with

 hyperkeratosis and ulceration, possible vascular

 ectasia

PM: proximal matrix; DM: distal matrix; NB: nail bed; NP: nail plate; DLSO: distal lateral subungual onychomycosis



jdv\_17521\_f1.png