

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Perivascular adipose tissue as a source of therapeutic targets and clinical biomarkers A clinical consensus statement from the European Society of Cardiology Working Group on Coronary Pathophysiology and Micro-circulation

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Antoniades, C., Tousoulis, D., Vavlukis, M., Fleming, I., Duncker, D.j., Eringa, E., et al. (2023). Perivascular adipose tissue as a source of therapeutic targets and clinical biomarkers A clinical consensus statement from the European Society of Cardiology Working Group on Coronary Pathophysiology and Micro-circulation. EUROPEAN HEART JOURNAL, 44(38), 3827-3844 [10.1093/eurheartj/ehad484].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/945934> since: 2023-10-23

*Published:*

DOI: <http://doi.org/10.1093/eurheartj/ehad484>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

**Perivascular adipose tissue as a source of  
therapeutic targets and clinical biomarkers**  
***A Clinical Consensus Statement from the ESC Working Group  
on Coronary Pathophysiology and Microcirculation***

Charalambos Antoniades<sup>1</sup>, Dimitris Tousoulis<sup>2</sup>, Marija Vavlukis<sup>3</sup>, Ingrid Fleming<sup>4</sup>, Dirk J. Duncker<sup>5</sup>, Etto Eringa<sup>6</sup>, Olivia Manfrini<sup>7</sup>, Alexios S. Antonopoulos<sup>1,2</sup>, Evangelos Oikonomou<sup>2</sup>, Teresa Padró<sup>8,15</sup>, Danijela Trifunovic-Zamaklar<sup>9</sup>, Giuseppe De Luca<sup>10</sup>, Tomasz Guzik<sup>12</sup>, Edina Cenko<sup>7</sup>, Ana Djordjevic-Dikic<sup>13</sup>, Filippo Crea<sup>14</sup>

<sup>1</sup> Acute Multidisciplinary Imaging and Interventional Centre, RDM Division of Cardiovascular Medicine, University of Oxford, UK.

<sup>2</sup> 1<sup>st</sup> Cardiology Department, National and Kapodistrian University of Athens, Greece.

<sup>3</sup> University Clinic for Cardiology, Medical Faculty, University Ss' Cyril and Methodius in Skopje, Skopje, North Macedonia

<sup>4</sup> Institute for Vascular Signalling, Centre of Molecular Medicine, Goethe University, Frankfurt, Germany.

<sup>5</sup> Department of Cardiology, Thorax center, Erasmus University Medical Center, Rotterdam, the Netherlands.

<sup>6</sup> Cardiovascular-Program ICCC, Research Institute—Hospital Santa Creu i Sant Pau, IIB-Sant Pau, Barcelona, Spain.

<sup>7</sup> Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy.

<sup>8</sup> Cardiovascular Program-ICCC, Institut d'Investigació Biomèdica Sant Pau (IIB SANT PAU)

<sup>9</sup> Cardiology clinic, University clinical Center, MedicalClinical, University of Belgrade, Serbia

<sup>10</sup> Division of Clinical and Experimental Cardiology, AOU Sassari, Sassari, Italy.

<sup>11</sup> Cardiologia Ospedaliera, Nuovo Galeazzi-Sant'Ambrogio, Milan Italy.

<sup>12</sup> Institute of Cardiovascular Medical Sciences, BHF Glasgow Cardiovascular Research Centre, UK and Department of Medicine, Jagiellonian University, Collegium Medicum, Krakow, Poland

<sup>13</sup> Cardiology Clinic, University clinical Center, Medical Faculty, University of Belgrade, Serbia.

<sup>14</sup> Department of Cardiology and Pulmonary Sciences, Catholic University of the Sacred Heart, Rome, Italy.

<sup>15</sup> CiberCV, Institute Carlos III, Madrid, Spain.

**Total word count (main text): 6027, Figures: 6, Tables: 4**

**Clinical Consensus Statement from the ESC working group on coronary pathophysiology and microcirculation**

**Corresponding author**

Prof Charalambos Antoniades

BHF Chair of Cardiovascular Medicine University of Oxford

Acute Multidisciplinary Imaging & Interventional Centre

Radcliffe Department of Medicine University of Oxford,

AMIIC, L2 John Radcliffe Hospital, Headley Way, Headington OX39DU

Oxford, United Kingdom

Email: [Charalambos.antoniades@cardiov.ox.ac.uk](mailto:Charalambos.antoniades@cardiov.ox.ac.uk)

## **Content Page**

Abstract	Page 3
Introduction	Page 4
Defining the different adipose tissue depots: Consensus terminology	Page 5
Biological differences between adipose tissue depots	Page 6
Understanding the cross-talk between PVAT and the vascular wall	Page 7
Bi-directional cross-talk between PVAT and the vascular wall	Page 9
PVAT as a source of therapeutic targets in cardiovascular disease	Page 9
Evaluating PVAT biology in metabolically healthy vs unhealthy obesity	Page 10
PVAT in diabetes and insulin resistance	Page 11
PVAT in arterial hypertension	Page 12
PVAT and sex differences	Page 13
Pharmacological targeting of PVAT	Page 14
PVAT as a source of diagnostic and prognostic biomarkers	Page 15
Imaging PVAT: Quantity vs Quality	Page 15
PVAT as sensor of disease signals from the vascular wall	Page 16
Using CT imaging of PVAT to measure vascular inflammation	Page 16
Using PVAT imaging for risk prediction: Is it prime time for its use in clinical practice?	Page 20
PVAT imaging vs circulating biomarkers of inflammation	Page 24
Future Perspectives and Remaining Challenges	Page 25
Conclusions	Page 26

## **Abstract**

Obesity is a modifiable cardiovascular risk factor, but adipose tissue (AT) depots in humans are heterogenous anatomically, histologically, and functionally. For example, visceral AT is a pro-atherogenic secretory AT depot, while subcutaneous AT represents a more classical energy storage depot. Perivascular adipose tissue (PVAT) regulates vascular biology via paracrine crosstalk signals. In this position paper we review the state-of-the-art knowledge of various adipose tissue depots and provide a consensus definition of PVAT around the coronary arteries, as the AT surrounding the artery up to a distance from its outer wall equal to the luminal diameter of the adjacent artery. We focus on the interactions between PVAT and the vascular wall, that render PVAT as a potential therapeutic target in cardiovascular disease. This Clinical Consensus Statement also discusses the role of PVAT as a clinically relevant source of diagnostic and prognostic biomarkers of vascular function, which may guide precision medicine in atherosclerosis, hypertension, heart failure and other cardiovascular diseases (CVD). We also highlight its role as a “biosensor” of vascular inflammation and describe recent imaging technologies that visualise PVAT in clinical practice, allowing non-invasive quantification of coronary inflammation and the related residual cardiovascular inflammatory risk, guiding deployment of therapeutic interventions. Finally, we review the current and future clinical applicability of artificial intelligence and machine learning technologies that integrate PVAT information into prognostic models to provide clinically meaningful information in primary and secondary prevention.

## Introduction

Obesity is a widely accepted risk factor for cardiovascular disease, as increased body mass index (BMI) has been associated not only with classic cardiovascular risk factors like type 2 diabetes mellitus, hypertension and hyperlipidemia, but also with coronary artery disease, heart failure, atrial fibrillation and stroke, independently from the other risk factors.<sup>1-3</sup> This was initially believed to be due to the detrimental effects of adipose tissue (AT)-derived molecules (e.g. adipokines) on the human arterial wall and heart.<sup>1</sup> However, during the last decade, it became clear that the association between obesity and cardiovascular disease is much more complex than originally anticipated. Epidemiological studies have shown a U-shape relationship between BMI and survival, with higher short- and mid- term survival rates observed among overweight individuals (BMI 25-30 Kg/m<sup>2</sup>), while “normal” body weight (BMI 20-24.9 kg/m<sup>2</sup>) only related with better long-term survival (>15 years).<sup>2</sup> This paradoxically “protective” effect of obesity became more evident in individuals with chronic diseases, including heart failure, renal failure or cancer.<sup>4</sup> This “obesity paradox” was originally explained by the inability of BMI to capture fat distribution in the body, since visceral obesity is the one that we should target, while gluteal/subcutaneous fat has a neutral effect on cardiovascular risk prediction. Visceral obesity was therefore defined as metabolically unhealthy obesity, while gluteal obesity was defined as metabolically healthy obesity.<sup>5</sup> However, the most important lesson learned by studying the “obesity paradox” was that AT may actually sense systemic signals of cachexia related with chronic diseases, activating lipolysis, a phenomenon driving weight-loss in chronic conditions like cancer or heart failure, leading to misinterpretations of the relationship between body fat mass and survival.<sup>4</sup> This could mean that AT can be used as a biosensor of the severity of chronic diseases, turning it into a

potential source of diagnostic or prognostic biomarkers, beyond its role as a therapeutic target. Therefore, understanding the biological nature of the cross-talk between AT and the cardiovascular system, is an unmet need, that will lead to the discovery of new therapeutic targets and the development of novel diagnostic biomarkers in cardiovascular medicine.

### **Defining the different adipose tissue depots: Consensus terminology**

Human AT is biologically classified into three main types: *white (WAT*, responsible for energy storage), *brown (BAT, responsible for ‘shivering thermogenesis’)* and beige (that includes adipocytes with intermediate phenotype between that of white and brown), distributed in the human body as shown in Figure 1.<sup>6,7</sup>

Anatomically, WAT can be subdivided into *visceral (VAT)* and *subcutaneous (SAT)*. SAT is located within the reticular dermis of the skin; in males, SAT is mainly located over the trunk, whilst in females is predominantly distributed around the hips and lower limbs.<sup>7</sup> The abdominal SAT is further distinguished into the superficial and deep subcutaneous AT layers, the latter being phenotypically closer to VAT.<sup>7</sup>

VAT includes the *abdominal* and *thoracic* AT. Abdominal VAT is located within the abdominal cavity, surrounding the visceral organs, and it is further subdivided into mesenteric peritoneal and retroperitoneal (perirenal) AT. Thoracic VAT includes the epicardial AT (EAT) which is enclosed between the cardiac surface and the visceral pericardium, the pericardial AT (external to the pericardium and surrounding the cardiac silhouette) as well as the non-pericardial thoracic AT (located anywhere inside the thoracic cavity, but outside the pericardium; this includes the pericardial depot) (**Figure 1**).<sup>8</sup> Perivascular AT (PVAT)<sup>9-11</sup> around the coronary

arteries has distinct biological properties compared to the rest of EAT, far from the arterial wall,<sup>12</sup> but as there is gradual transition between PVAT and non-PVAT within EAT, without any anatomical structure separating the two, the definitions need to be done by consensus. Indeed, in the recent literature,<sup>13-19</sup> PVAT is defined as the layer of AT located within a distance equal to the luminal diameter of the adjacent artery, a definition adopted by the working group.<sup>10</sup> This definition applies to the PVAT surrounding any artery in the human body up to a luminal diameter of 2cm (see **Figure 1**). For arteries with luminal diameter >2cm (such as the aorta), PVAT extends up to 2cm from the outer surface of the vessel.

### **Biological differences between adipose tissue depots**

The nomenclature of human AT depots does not serve only the purposes of the anatomical classification as the anatomical location is a critical determinant of AT phenotype and its biology (**Table 1**). For example, VAT contains smaller, less differentiated adipocytes with significantly higher infiltration by macrophages compared to the subcutaneous depot.<sup>20</sup> VAT is highly active in terms of adipokine secretion, and its secretome closely correlates with systemic metabolic status and insulin resistance. In contrast SAT contains larger, well-differentiated adipocytes, and is less active in terms of adipokine secretion.<sup>21</sup> PVAT has a unique role in the physiology of the cardiovascular system given its anatomical proximity to the vascular wall (**Table 2**). In this article we will focus on the role of PVAT in cardiovascular physiology and will discuss its role as a therapeutic target and/or as a source of diagnostic biomarkers in clinical practice.

## Understanding the cross-talk between PVAT and the vascular wall

AT secretes bioactive molecules, collectively called adipokines, that can affect cardiovascular physiology in an endocrine manner (via their release in systemic circulation). Not all depots affect the cardiovascular system in a similar way. The secretome of the highly vascular VAT is released into the bloodstream, exerting primarily detrimental effects on the cardiovascular system, in an endocrine manner. Conversely, the less active SAT has a neutral relationship with cardiovascular risk, while the accumulation of gluteal AT may even be a marker of lower cardiovascular risk, as it secretes largely anti-atherogenic adipokines like adiponectin. Beyond its endocrine effects on the cardiovascular system, AT can also affect the heart and the vascular wall directly, in a paracrine (i.e., by diffusion) or vasocrine (i.e., mediated by the local microcirculation) way, as shown in **Figure 2A**.<sup>22,23</sup>

PVAT was originally thought to serve only as a structural support tissue providing physical protection to the blood vessels. However, in the past 20 years its role in regulating vessel function has become increasingly recognized, as it releases numerous vasoactive factors (adipokines) with paracrine (outside-in) effects on the vascular wall, in addition to its ability to also sense signals released from the vessel wall (inside-out) that determine its biological behavior, as part of a cross-talk between the two.<sup>24,25</sup>

PVAT includes a range of cell types including adipocytes, neuronal cells (autonomic nerves), immune cells and the stromal vascular fraction that consists of pre-adipocytes, fibroblasts, endothelial and mesenchymal stem cells derived from the local microcirculation.<sup>26</sup> In clinical studies, PVAT is generally considered part of WAT, but PVAT's adipocytes are generally smaller in size, with less lipid droplet accumulation and less differentiated adipocytes, with gene



expression patterns often similar to those of adipocytes in BAT, particularly in animal models.<sup>27,28</sup> PVAT also houses a community of innate and adaptive immune cells consisting of lymphocyte (T and B cells), macrophages, eosinophils, NK-cells and mast cells. Under healthy conditions, macrophages, and T cells form the dominant immune cell types in PVAT, contributing to the anti-inflammatory effects and, together with the eosinophils to the PVAT's anti-constrictive effects on the vessel wall.<sup>27,29</sup>

PVAT, releases bioactive molecules such as adipokines, anti-inflammatory and pro-inflammatory factors, miRNAs and others (e.g. hydrogen sulfide, reactive oxygen species, fatty acid metabolites, etc.) and contributes to the regulation of vascular homeostasis.<sup>30</sup> Under physiological conditions, PVAT exerts a net vasodilatory, anti-oxidant and anti-inflammatory effects on the vasculature.<sup>24</sup> The vasodilatory effects are due to the secretion of adipocyte-derived relaxing factor, adiponectin, apelin, leptin and omentin, which induce vasodilation either by acting directly on vascular smooth muscle cells (e.g. adipocyte-derived relaxing factor, leptin and adiponectin) and/or indirectly by increasing the endothelium-derived vasodilators such as nitric oxide and endothelium-derived hyperpolarizing factor (e.g. apelin, omentin, leptin and adiponectin).<sup>1</sup> The paracrine anti-oxidant and anti-inflammatory actions by adipokines released from PVAT under physiological conditions likely involve 5' adenosine monophosphate-activated protein kinase (AMPK) and NO signaling<sup>31 32</sup> within the adjacent vessel.<sup>33-35</sup> **Table 3** summarizes the cellular sources and the vascular effects of key PVAT-derived adipokines.

### ***Bi-directional cross-talk between PVAT and the vascular wall***

PVAT is involved in bi-directional interactions with the vascular wall. Indeed, PVAT responds to paracrine signals from vascular cells that facilitate phenotypic changes of PVAT adipocytes, modifying the composition of their secretome that exerts paracrine effects back on to the vascular wall. This crosstalk is different in health and disease. In the presence of high vascular oxidative stress, the vascular wall secretes lipid peroxidation products such as 4-hydroxynonenal (4-HNE), which are diffused to the surrounding PVAT, activating PPAR- $\gamma$  signaling in the perivascular adipocytes; this leads to up-regulation of adiponectin expression and secretion from PVAT which then suppress vascular oxidative stress as a local defense mechanism against vascular oxidative damage.<sup>36-38</sup> In a similar way, PVAT adipocytes sense inflammatory molecules secreted from the vascular wall (e.g. IL-6, TNF $\alpha$ , IFN $\gamma$  etc), and they change their phenotype from storage cells to secretory cells, activating lipolysis and inhibiting adipogenesis.<sup>12</sup> In summary, the bi-directional communication (**Figure 2B**) between vascular cells and PVAT plays a role in regulating vascular function. Better understanding of PVAT specific secretome changes in response to vascular signals, including adipokines, cytokines and lipid peroxidation products, is essential to identify new diagnostic biomarkers<sup>39</sup> and therapeutic targets in cardiovascular medicine.<sup>40</sup>

### **PVAT as a source of therapeutic targets in cardiovascular disease**

The paracrine effects of PVAT, particularly on vascular tone and vascular inflammation, are significantly altered in diseases such as (pre)diabetes, hypertension, and cardiac diseases. Indeed, in pathological conditions, cytokines and classical adipokines dysregulate the expression of

critical genes involved in vascular redox balance, such as NADPH oxidases (Nox) or endothelial nitric oxide synthase (eNOS), leading to endothelial dysfunction. For example, while leptin directly induces vasorelaxation, prolonged exposure of endothelial cells to leptin leads to increased oxidative stress, activation of NADPH oxidases and endothelial dysfunction.<sup>41</sup> Resistin and visfatin have also been shown to induce prooxidant and proinflammatory phenotypes in the vasculature.<sup>42</sup>

It has recently been shown that PVAT secretes also adipokines such as secreted frizzled-related protein 5 (SFRP5) and ligands regulating non-canonical Wnt-signalling (e.g. wnt5a), playing a major role in controlling vascular redox state.<sup>43</sup> At the same time, PVAT's metabolic and secretory function is regulated by signals originating from vascular cells. For example, it has been shown that cytokines released either from the atherosclerotic plaques or by the inflamed vascular wall in the absence of atherosclerotic plaques (such as IL-6 or TNF- $\alpha$ ) leads to adipocyte dedifferentiation, decreased lipid storage and increased ability to further promote perivascular inflammation by inducing the expression of chemokines such as RANTES, that drive further development of perivascular inflammation.<sup>44</sup> Similarly, in hypertensive mice, eosinophilic granulocytes within PVAT determine vascular resistance and blood pressure.<sup>29</sup>

### **Evaluating PVAT biology in metabolically healthy vs unhealthy obesity**

Obesity does not always result in metabolic dysregulation, and there are metabolically “healthy” individuals with obesity.<sup>45</sup> In general, the metabolically “healthy” individual with obesity does carry an expanded total AT mass in the absence of metabolic diseases such as type 2 diabetes mellitus, dyslipidemia, or hypertension (such as sumo wrestlers). While the local accumulation of

PVAT has been consistently associated with the development of cardiometabolic complications in obesity,<sup>46,47</sup> there is evidence that metabolically healthy obesity is accompanied by a more favorable PVAT's inflammatory status than metabolically unhealthy obesity.<sup>11</sup> The “metabolically healthy” obesity state is more often observed in young, physically active individuals with a good nutritional status and low levels of ectopic and visceral fat storage.<sup>48</sup>

### **PVAT in diabetes and insulin resistance**

PVAT in skeletal muscles controls insulin-stimulated perfusion (via insulin-mediated microvascular recruitment), mitochondrial protein expression and glucose uptake.<sup>9</sup> Insulin-mediated microvascular recruitment and related vasodilatation, expands the endothelial surface area in direct contact with blood, facilitating extraction of glucose and insulin into the muscle interstitium. This vasodilatory insulin effect strongly depends on locally (in PVAT) produced adiponectin which penetrates the vascular wall either by diffusion (paracrine effect) or possibly through the adipomuscular arterioles (vasocrine effect),<sup>9</sup> although the later remains largely speculative. PVAT is also important for insulin regulation of muscle glucose metabolism, controlling heat shock protein expression (HSP 90AB1 that is involved in the regulation of glucose and fatty acid metabolism) and expression of the mitochondrial protein complexes engaged in the respiratory chain.<sup>49</sup>

PVAT is dysfunctional in individuals with insulin resistance and type 2 diabetes mellitus,<sup>50</sup> as it loses its ability to mediate insulin-induced vasodilatation and to antagonize sympathetic tone, predominately due to impaired adiponectin secretion.<sup>50</sup> In mice, inflammation of PVAT impairs insulin-induced vasodilatation during weight gain, contributing to impaired muscle blood flow

and insulin resistance.<sup>51</sup> In addition, removal of PVAT (in vivo animal experiments) causes decreased expression of mitochondrial electron transport chain components, a characteristic of muscle insulin resistance in diabetes.<sup>9</sup> Paracrine signaling by intramuscular PVAT strongly modulates muscle insulin sensitivity and plays an important role in the pathogenesis of type 2 diabetes. Impaired cross-talk between PVAT and microvascular endothelium also predisposes to insulin resistance and type 2 diabetes.<sup>35</sup> The mechanisms by which PVAT biology shifts towards its pro-insulin resistance phenotype is not fully understood. Under diabetic conditions the biosynthetic activity of para-aortic PVAT shifts towards a pro-inflammatory (increased CRP, CCL2, CD36), pro-oxidant (increased aldose reductase, and reduced antioxidant defence enzymes) and vasoconstrictive state.<sup>52</sup> Pro-inflammatory phenotype of PVAT can be induced by high-fat feeding.<sup>53</sup> Since the skeletal muscles account for the greatest proportion of insulin resistance in diabetes, it might be speculated that PVAT dysfunction is one of the key steps for the development of insulin resistance and type 2 diabetes, although further data is needed to document this role.

### **PVAT in arterial hypertension**

Inflammation is essential in the pathogenesis of hypertension<sup>54,55</sup> Immune cell activation and its infiltration into the target organs including the vasculature, increases blood pressure<sup>56</sup> and causes target organ damage.<sup>57,58</sup> In the vasculature, PVAT is the primary site of immune cell infiltration in hypertension. Angiotensin II, Salt, as well as initial prehypertensive increases of blood pressure induce chemokine release from PVAT which mediates perivascular inflammation (PMID). This includes chemokines such as RANTES, IP-10, MCP-1 attracting T cells as well as

macrophages into PVAT.<sup>57,59</sup> These cells release effector cytokines implicated in hypertension, such as IFN-gamma, IL-17, TNF-a, IL-6, that cause vascular dysfunction, oxidative stress and vascular stiffness, all of which are critical processes in hypertension.<sup>54,57</sup> The role of PVAT inflammation in the regulation of blood pressure, provides a further mechanistic link between obesity, atherosclerosis and hypertension.<sup>60,61</sup>

### **PVAT and sex differences**

Women exhibit significantly higher pericardial and epicardial fat volumes with declining oestrogen levels following menopause.<sup>1</sup> This observation highlights a potential role for ectopic fat depots in sex differences in cardiovascular risk.<sup>1,8</sup> Although there is a paucity of human data, healthy female pigs show greater PVAT-derived relaxing factors such as adiponectin and lower levels of adipose-derived contracting factor such as thromboxane A<sub>2</sub>, compared to healthy males.<sup>62,63</sup> Additionally, PVAT releases nitric oxide (NO) and cyclooxygenase metabolites to induce coronary vasodilation in young female but not male pigs, suggesting a sex-specific effect of PVAT on vascular tone.<sup>62</sup> Nevertheless, reduction of oestrogen levels following menopause is associated with a change from prevalent release of vasorelaxing factors towards vasoconstrictive as well as pro-inflammatory adipokines.<sup>64</sup> Ovariectomy increased the vasoconstrictor endothelial dependent responses by reducing endothelial NO and increasing TXA<sub>2</sub>/ cyclooxygenase (COX) signalling and attenuated the vasodilatory effects of PVAT, in female rats as compared with sham rats suggesting that oestrogen regulates PVAT-mediated microvascular tone.<sup>65</sup> Oestrogen receptors are expressed in AT, but their potential role in PVAT remains unknown, while the role of oestrogen-receptor signalling as a therapeutic target within the human PVAT is unclear.

## Pharmacological targeting of PVAT

Given its well-established role in the pathophysiology of vascular disease, PVAT forms a rational therapeutic target in cardiovascular medicine. Current anti-diabetic treatments, partly exert their beneficial cardiovascular effects by acting on AT. There is evidence that PVAT of animals and humans phenotype changes quickly in response to changes in diet<sup>66,67</sup> or physical activity,<sup>68</sup> while pharmacological anti-diabetic treatments may also affect PVAT's function. Glucagon-like peptide-1 receptor (GLP-1R) expression levels in adipocytes are associated with insulin resistance, and GLP-1R agonists promote the differentiation of adipocytes and restore adipocyte health in cell culture models.<sup>69</sup> In humans, treatment of patients with obesity with the GLP-1 analog liraglutide has beneficial effects on plasma lipids;<sup>70</sup> while maintaining weight loss, liraglutide also lowers apolipoprotein B plasma levels which may help reducing CVD risk;<sup>71</sup> for example, fasting GLP-1 levels are associated with decreased carotid intima-media thickness;<sup>71</sup> these effects of liraglutide may be partly mediated suppresses plasma C16:0-ceramide species, AT-derived sphingolipids that induce vascular dysfunction and drive cardiovascular risk.<sup>72,73</sup> Sodium–glucose cotransporter-2 inhibitors (SGLT2-i) also exert beneficial effects on human AT. For example, the SGLT2-i empagliflozin promotes AT browning in white AT and activates residing M2 macrophages attenuating obesity-induced inflammation and insulin resistance in animal models.<sup>74</sup> In humans, empagliflozin treatment reduces the volume of EAT, which is closely linked to cardiometabolic risk.<sup>75</sup> Whether such beneficial effects are also extended to PVAT biology and PVAT-vascular wall interactions remains to be seen.<sup>76</sup> Statins lower vascular inflammation and this is reflected in respective changes in PVAT phenotype after an acute coronary syndrome (ACS) and statin-treatment initiation.<sup>12</sup> Statins increase H<sub>2</sub>S bioavailability in PVAT and this may contribute to the

anticontractile vascular effects of atorvastatin.<sup>77</sup> Other potential strategies to restore PVAT health, include rosiglitazone (albeit with known adverse effects in congestive heart failure) and cannabinoid CB1 receptor agonists to promote H<sub>2</sub>S release from PVAT, or inhibition of PVAT inflammation with melatonin or cytokine antagonists, or enhancers of adiponectin expression.<sup>78</sup> Although part of the effectiveness of these drugs may be mediated by their effects on PVAT, this is difficult to prove. There is an unmet need to test PVAT-specific drug-delivery systems, to test the effectiveness of PVAT as a direct therapeutic target in cardiovascular disease.

## **PVAT as a source of diagnostic and prognostic biomarkers**

### **Imaging PVAT: Quantity vs Quality**

The development of new imaging modalities and the incorporation of computational systems in the analysis and interpretation of medical images has opened new horizons in measuring the volume of AT (e.g., EAT as a surrogate marker of metabolically unhealthy obesity which predicts non-CV mortality)<sup>79</sup> or even its “quality”, by using complex image post-processing. The amount and type of information provided by each imaging modality is different but Computed Tomography (CT) is considered to be the gold standard in visualizing and characterizing PVAT, due to its very high resolution, and the distinct attenuation signals of AT, falling typically between -30 and -190 Hounsfield Units (HU). Other imaging modalities still have major limitations; for example, echocardiography cannot be used for tissue characterization or for coronary artery imaging, positron emission tomography (PET) suffers from high noise/signal ratio coming from vessels and/or underlying myocardium, while MRI still lacks the anatomical resolution for the



imaging of coronary arteries. **Table 4** summarizes the value of different imaging modalities in assessing PVAT phenotype.

### **PVAT as sensor of disease signals from the vascular wall**

As discussed earlier, in disease states, the human arterial wall secretes various mediators such as oxidation products (e.g. 4-HNE) which diffuse to PVAT, triggering “re-programming” of adipocytes from quiet lipid-storage cells to active biosynthetic cells secreting more antioxidant adipokines such as adiponectin, back to the vascular wall as a “defense mechanism” against vascular oxidative damage.<sup>6,36,38</sup> Inflammatory molecules from the vascular wall diffuse into PVAT<sup>12</sup> and stall the differentiation of pre-adipocytes into mature adipocytes in PVAT.<sup>12</sup> Moreover, vessel-derived inflammatory molecules induce perivascular lipolysis and lead to a gradient of adipocyte size around the inflamed artery, with smaller and low fat- containing adipocytes close to the vessel, transiting into larger and fat-filled adipocytes further away from the outer vascular surface.<sup>6,12</sup> This gradient in adipocyte size within PVAT surrounding the inflamed artery, results into higher lipid/water ratio in PVAT’s layers close to the inflamed vascular wall.<sup>10,12</sup> This gradient of PVAT’s structure and composition around inflamed arteries may serve as an internal “thermometer” or vascular inflammation, if it can be visualized and quantified non-invasively. The creation of 3D gradients of adipocyte size around inflamed arteries is demonstrated in **Figure 3**.

### Using CT imaging of PVAT to measure vascular inflammation

CT can be used to extract not only quantitative (volumetric) information about PVAT, but also qualitative information about its structure and composition (including its 3D texture).<sup>13</sup> Indeed, the gradient in lipid accumulation and adipocyte size around inflamed coronary arteries can be visualized and quantified from routine coronary CT angiograms (CCTA).<sup>12</sup> This continuum of morphological changes in PVAT can be detected as a gradient in the CT signal attenuation in the perivascular space (within the window of -30 to -190 HU).<sup>12</sup> A metric developed to quantify these three-dimensional attenuation gradients in CCTA imaging of PVAT is the perivascular Fat Attenuation Index (FAI).<sup>12</sup> This measurement incorporates corrections for technical scan parameters that affect the attenuation readings around human arteries, in a non-linear way. Perivascular FAI has been biologically and clinically validated and it is now considered as the main imaging biomarker of coronary inflammation extracted from CCTA (**Figure 3**).<sup>80</sup> Perivascular FAI was originally measured around the proximal segment of the right coronary artery (RCA) over a 40mm-segment (10-50mm from RCA origin) in a radial distance from the arterial wall equal to the diameter of the underlying artery) as it was technically easier to validate the method in that vessel.<sup>12</sup> However, further validation studies have led to the development of appropriate algorithms that calculate perivascular FAI around the proximal segments of the left circumflex artery (LCx) as well as the left anterior descending artery (LAD).<sup>39,81</sup> Analysis of the PVAT around the proximal segments of epicardial coronary arteries can be used a measure of the overall background inflammatory burden of the respective coronary artery.<sup>10</sup> Standardised FAI measurement could take into account any side branches, subtracting the PVAT volume of the side branch from the volume included into the calculation of FAI around the main epicardial artery.

This becomes very important when attempting to measure FAI in mid-/distal segments of the coronary arteries.

A per-lesion analysis of PVAT may be also applied around individual coronary plaques to detect inflammatory signals. Perivascular FAI increases significantly around vulnerable / ruptured plaques in patients with an ACS.<sup>10,12</sup> The average pericoronary AT attenuation is higher around atherosclerotic coronary segments compared to coronary segments without disease,<sup>82</sup> predicts non-calcified plaque progression,<sup>83</sup> while there are significant differences in average pericoronary fat attenuation around culprit vs. non-culprit lesions of patients with ACS.<sup>19,84,85</sup> The measurement of pericoronary fat attenuation is also nearly co-linear to <sup>18</sup>FNaF coronary uptake, measured using PET/CT, which is considered the gold standard to measure microcalcification in atherosclerotic plaques, an indirect metric of inflammation in vivo (**Figure 4A**).<sup>86</sup> Albeit higher pericoronary attenuation has been reported around lesions of intermediate luminal severity with a low fractional flow reserve (FFR),<sup>87,88</sup> this findings should be interpreted with caution; FAI is a metric of vascular inflammation and any association with FFR could be explained by unaccounted confounders (e.g., atheroma burden). The two methods provide complementary results, as they capture different biology (local vascular inflammation vs hemodynamic compromise due to local atheroma, inflamed or not), and associations between them are likely to be indirect.

#### *Using FAI measurements in clinical practice: challenges and opportunities*

Measurement of pericoronary FAI is derived from the post-processing of standard CCTA images, and therefore its calculation involves all typical requirements for acquiring a CCTA scan including medication (beta-blockers, nitroglycerine), ECG-gating, and radiation exposure as per

Society of Cardiovascular Computed Tomography guidelines,<sup>89</sup> while it suffers from the same technical limitations of CCTA interpretation (motion artefacts, arrhythmias etc). Nonetheless in the CRISP-CT study only 5.6% of the scans were not analyzable for FAI because of poor scan quality (e.g. motion artefacts) and these were also clinically non-diagnostic. Pericoronary FAI is a crude method to allow assessment of coronary inflammation, and it performs well as a research tool. However, translating this into a clinically meaningful reading of coronary inflammation, which will make sense for individual patient's management is challenging. Indeed, the measurement of attenuation on CCTA is influenced by various technical factors (e.g. tube voltage etc),<sup>90</sup> biological factors (e.g. the background adipocyte size is driving the physiological range of attenuation measurements to more negative values, resulting into respective shifts of the expected normal values), and anatomical factors (e.g. the segment of the epicardial coronary tree where the measurements are performed affects the expected attenuation values etc).<sup>12</sup> A recent study has presented a new algorithm (FAI-Score) which corrects the FAI values for a range of technical, biological and anatomical factors, and provides a standardized metric of the degree of background coronary inflammation for each of the 3 epicardial coronary arteries separately. Indeed, FAI-score is expressed in arbitrary units (AU), and to be clinically meaningful it needs to be interpreted on age- and sex- specific nomograms (**Figure 3**).<sup>39</sup> This artery-specific measurement is currently the only regulatory cleared metric of coronary inflammation derived from CCTA in Europe, and it is measured separately for each epicardial coronary artery. However, the local measurement of perivascular FAI around specific atherosclerotic plaques (plaque-specific inflammatory burden, **Figure 4B**) remains a promising research tool to detect the

inflamed plaque, but it is neither validated clinically nor regulatory cleared, so it can not be used in clinical practice yet.<sup>13</sup>

#### *Perivascular FAI and responsiveness to anti-inflammatory treatments*

Evidence suggests that perivascular FAI changes significantly in response to anti-inflammatory treatments.<sup>15,91</sup> A recent study has demonstrated significant reduction in pericoronary FAI around non-calcified and mixed coronary plaques (but not calcified plaques) at one year after initiation of statin treatment,<sup>29</sup> confirming the antiinflammatory effect of statins. In a prospective clinical study involving patients with psoriasis, a condition which is associated with increased levels of vascular inflammation, biologic anti-inflammatory agents (i.e., anti-TNF- $\alpha$  or anti-IL 12/23 and anti-IL-17 therapies) reduced perivascular FAI within 1-year of treatment (**Figure 4C**).<sup>91</sup> In ACS, the culprit lesion inflammatory burden (calculated as the area under the curve of FAI-changes across the atherosclerotic plaque, above the baseline value defined by a reference segment proximal to the plaque- **Figure 4B**) is elevated in culprit lesions during ACS, and it is returning to a steady state within 6 months from the event, in at least 80% of the patients.<sup>10,12,13</sup> The remaining ~20% of the patients whose atherosclerotic plaques do not “cool down” within 6 months, may be more likely to develop recurrent ACS, and could represent a target population for anti-inflammatory agents like colchicine or novel therapeutics targeting coronary inflammation.<sup>10</sup> Further randomized clinical trials are needed, to quantify the effect size of various anti-inflammatory treatments on coronary inflammation, measured using this method.

### Using PVAT imaging for risk prediction: Is it prime time for its use in clinical practice?

The CRISP-CT (Cardiovascular RISK Prediction using CT) study explored the prognostic value of pericoronary FAI in two independent cohorts in a total of 3,912 patients, followed up for up to a decade post-CCTA.<sup>81,92</sup> Perivascular FAI around the three epicardial coronary arteries was strongly prognostic of both cardiac mortality (HR 5.6 and 9 in the two cohorts, above vs below a FAI threshold of -70.1 HU, after correction for all risk factors, high risk plaque (HRP), degree of coronary stenosis etc.) and non-fatal myocardial infarction (HR 5.0 using the same FAI threshold and following the same corrections mentioned above)- **Figure 4D-E**.<sup>81</sup> In the SCOTHEART study,<sup>93</sup> uncorrected measurement of pericoronary attenuation above and below the -70HU cut-off, showed a highly significant prognostic value for non-fatal cardiac events but with a smaller effect size compared to the weighted FAI used in CRISP-CT (**Figure 4F**), consistent with previous findings suggesting a significantly lower prognostic value of uncorrected pericoronary attenuation vs the weighted FAI measurement (**Figure 4G**).<sup>94</sup> Moreover, the FAI cut-off used in the CRISP-CT study (-70.1HU) was derived from cox-regression models, and it cannot be used as a meaningful cut-off in clinical practice. As mentioned earlier, the absolute FAI values are influenced by various technical, biological and anatomical factors, and corrected measures like the FAI-Score projected on nomograms, are more appropriate for standardized measurements of coronary inflammation.<sup>95</sup> Indeed, FAI-Score above the 50<sup>th</sup> percentile in the RCA and the LAD is related with 2-fold increase of relative risk for fatal cardiac events, while increase over the 75<sup>th</sup> and 95<sup>th</sup> percentiles is related with 2.4 and 5 fold increase of the relative risk respectively (**Figure 4H**).<sup>95</sup> Interestingly, for the LCx, a FAI-Score above the 95<sup>th</sup> percentile is needed to achieve ~2.4 times increase of the relative risk for fatal cardiac events, possibly because the LCX is often a small

vessel and it is responsible for fewer fatal cardiac events compared to the other two main coronary branches.<sup>95</sup>

#### *Using pericoronary fat imaging in clinical practice*

The information provided by FAI-Score is complementary to existing risk factors as well as the atherosclerotic plaque volume (calcified or non-calcified plaque burden) and/or the characteristics of vulnerable plaque detected in CCTA (e.g. high-risk plaque features like low-attenuation plaque, positive remodelling, spotty calcification and napkin ring sign).<sup>96</sup> A prognostic model that includes FAI-Score, atherosclerotic plaque burden as well as the clinical risk factors was trained in the US population of CRISP-CT study and validated in the European population against hard endpoints, reclassifying ~16% of the patients to a higher and ~20% to a lower risk category for cardiac mortality.<sup>39</sup> Indeed, such a model has been proposed for use in conjunction with the ESC SCORE, providing more sophisticated and accurate risk prediction in those patients with CCTA information available, and has formed part of a regulatory cleared medical device used in clinical practice. This revised absolute risk calculator has been proposed as a tool to treat these patients according to the ESC Guidelines on cardiovascular disease prevention in clinical practice.<sup>97,98</sup> It has also been proposed that in patients with at least ~2.4-fold increase of their relative risk for a fatal cardiac event (i.e. when FAI-Score in LAD or RCA >75<sup>th</sup> percentile), intense statin- or other anti-inflammatory treatments may be able to reduce long-term risk.<sup>39</sup>

Although the current risk-factors-based risk scores (like the ESC-SCORE<sup>97</sup>) work well in primary prevention, their value could be massively increased if they are combined with information from imaging, like PVAT phenotyping and plaque characteristics. Indeed, combining perivascular FAI analysis with HRP features, identifies a group of patients at very high risk for

future cardiac events (i.e. those with both high peri-coronary FAI and high-risk plaque, with ~7 fold increase of the 10 years risk of fatal cardiac events) and up to 11-fold increase of the risk of non-fatal myocardial infarction.<sup>93</sup> This approach also identifies a group of intermediate/high risk patients, with high peri-coronary FAI and no HRP (whose relative risk for fatal cardiac event over the decade is ~5).<sup>17</sup> Whether a FAI-based approach is superior to functional tests of myocardial ischaemia for risk stratification remains to be seen as studies have published both negative<sup>14</sup> and positive<sup>99</sup> results. However, there is strong evidence that adding FAI into a model that includes risk factors, high risk plaque and calcium score increased the prognostic performance of the model;<sup>81</sup> similarly, a risk prediction model with perivascular FAI performed much better compared in predicting cardiac risk.<sup>100</sup> It is therefore important to come up with prognostic models that combine classic risk factors with information on plaque and perivascular fat phenotyping, which can be used in clinical practice to calculate the absolute risk of a patient for a cardiac event. Because such models combine information from imaging as well as prognostic modelling, they need to meet the appropriate regulatory standards as medical devices and be cleared by the respective regulatory authorities (i.e. CE mark under the European Medical Device Regulations, Food and Drugs Administration clearance for the USA etc), before they are implemented in clinical practice.<sup>101</sup> The performance of such a model is expected to be significantly superior to the risk-factors-only based risk stratification models, but they need to be trained against appropriate clinical cohorts with significant follow-up after their CCTA. Currently, such models can only be created for patients having CCTA as part of routine clinical care, because they can be trained only in such cohorts of patients linked with prospective 10-years outcomes (e.g., in CRISP-CT study<sup>81</sup>). Studies like SCAPIS,<sup>102</sup> linking CCTA information from



primary prevention populations with outcomes data, will allow the development of such prognostic models for asymptomatic individuals, enabling community screening.

*Using artificial intelligence for radiotranscriptomic phenotyping of PVAT in risk prediction:*

Extraction of PVAT radiomic features and application of machine learning algorithms can generate more sophisticated biomarkers for the deep phenotyping of PVAT.<sup>13</sup> Different types of vascular inflammation can give different texture changes in PVAT, driven by variable degrees of perivascular lipolysis, adipogenesis, oedema, fibrosis and angiogenesis.<sup>6</sup> By using tissue biopsies and basic science tools (like RNA sequencing or histology) to generate the “ground truth” for these changes, one can train radiomic signatures of PVAT from CCTA images to best describe the type of vascular inflammation of interest. The term “radiotranscriptomic” has been introduced in cardiovascular imaging to describe the process of training radiomic signatures against the transcriptomic profile of the tissue.<sup>13</sup> Indeed, by using AT biopsies from patients undergoing cardiac surgery and available coupled CCTA scans, machine learning algorithms were trained to identify AT inflammation, fibrosis and vascularity from the radiomic phenotype of fat and generate Fat Radiomic Profile (FRP, **Figure 5**). FRP was externally tested in a cohort of 5,487 participants from the CRISP-CT<sup>81</sup> & SCOT-HEART studies<sup>103</sup> and was able to independently predict MACEs beyond traditional risk factors, coronary calcium score, coronary stenosis, and HRP features on CCTA.<sup>13</sup> Such approach was tested with similar success in identifying unstable coronary plaques and predicting outcomes, in other cohorts.<sup>104-106</sup>

Recently, a radiotranscriptomic signature of acute cytokine-driven vascular inflammation was trained using radiomic features of PVAT around the internal mammary arteries against RNA sequencing data extracted from the same arteries.<sup>107</sup> That signature was then tested in patients

with COVID-19, and it was found to change significantly during acute COVID-19. This radiotranscriptomic signature had striking prognostic value for in-hospital mortality in acute COVID-19, even when applied in non-gated CT angiograms of the pulmonary artery. Texture radiotranscriptomics can also be used to capture and quantify microcirculation in the perivascular space, in addition to lipolysis/adipogenesis, fibrosis and edema, providing incremental prognostic value over FAI for cardiac events.<sup>13,107</sup> Such machine learning/radiotranscriptomic approaches are expected to revolutionize our capacity to use PVAT as a window into vascular biology.

### **PVAT imaging vs circulating biomarkers of inflammation**

The non-invasive detection of vascular inflammation was hailed as the “Holy Grail” in the field of cardiovascular medicine.<sup>39</sup> The detection of residual inflammatory risk is particularly relevant in the light of recent clinical evidence supporting the reduction of CAD risk by anti-inflammatory treatments.<sup>108-110</sup> Therefore, the use of inflammatory biomarkers has been proposed as a way to guide the deployment anti-inflammatory treatments in primary and secondary prevention.<sup>108,111</sup>

Plasma biomarkers of inflammation, such as C-reactive protein (CRP) or IL-6, are not specific for vascular inflammation; for example it is estimated that approximately 50% of patients in secondary prevention, have high inflammatory risk based on serum CRP levels.<sup>112</sup> Although HRP characteristics on CCTA provide indirect information about the potential of a plaque to be inflamed, the combination of HRP with newer metrics of inflammation such as the perivascular FAI-Score, allows for a further refinement of the risk for cardiovascular events.<sup>17,113</sup> A recent meta-analysis has shown that CCTA-based biomarkers such as HRP or pericoronary FAI provide

much higher added prognostic value on top of clinical risk profile and atherosclerosis extent for MACEs, compared to plasma biomarkers (**Figure 6**).<sup>114</sup>

### **Other imaging techniques to assess PVAT: advantages, disadvantages, and future perspectives**

Other non-invasive imaging modalities like MRI, PET or even ultrasound could potentially add value in assessing PVAT. Indeed, MRI can quantify PVAT volume around large arteries like the aorta and this is independently associated with measures of subclinical atherosclerosis.<sup>115</sup> However, MRI does not allow accurate visualization of PVAT around the coronaries, while another major limitation is the standardized image quality to allow assessment of adipose tissue quality, able to provide between-patient comparisons. However, crude measurements of the total epicardial and pericardial adipose tissue volume provides some useful information about the predictive value of visceral obesity in the general population, as demonstrated in the UK biobank recently,<sup>116</sup> which is however significantly lower compared to the volumetric quantification of EAT using CT.<sup>79</sup>

PET is another promising imaging modality that provides functional information regarding adipose tissue metabolic activity.<sup>18</sup> Indeed, FDG uptake in large adipose tissue depots provides useful information regarding the inflammatory and metabolic activity of adipose tissue.<sup>117</sup> When PET is coupled with CT imaging anatomical information (PET-CT), it allows better understanding of the anatomical distribution of the radiotracer uptake, particularly around large vessels like the aorta.<sup>118</sup> However, the low spatial resolution of PET, as well as the uptake of FDG by the myocardium, does not allow reliable assessment of peri-coronary adipose tissue.<sup>118</sup> Future improvements in the spatial resolution of the method as well as the development of novel

radiotracers that would target specific metabolic or inflammatory pathways in the AT (adipocytes of AT-infiltrating inflammatory cells), could potentially transform this method into a valuable tool to assess PVAT biology, with possible clinical implications.

Other imaging techniques like ultrasound may also be of some value in obtaining surrogate measurements of EAT<sup>20</sup> but its limited ability to differentiate adipose tissue from other structures, as well as its low penetration and the operator-dependent nature of the technique, limit its value in the assessment of PVAT around the coronaries. Improvements in the post-processing of ultrasound images, may change the ability of this method to assess PVAT in the future.

Finally, the advent of the revolutionary technology of Photon Counting CT with its superior image resolution and capacity for tissue characterization by using subtraction techniques, is also expected to open up new possibilities in the field of PVAT imaging.<sup>119</sup> For now, there is a significant volume of technical work that needs to be done for the calibration of this technology. Indeed, the established definitions of PVAT using contemporary CT (quantitative e.g., EAT or qualitative e.g., FAI Score) will need to be re-defined in an environment of changing energies (keV) and higher spatial resolution, while the behaviour of radiomic features in this environment will also need to be further evaluated. Validation studies against tissue biopsies<sup>120</sup> as well as against contemporary CT analysis of PVAT<sup>121</sup> are ongoing, and are expected to be reported in the coming months. Until then, photon counting CT images are incompatible with the existing clinical tools used for the qualitative assessment of PVAT.

## Future Perspectives and Remaining Challenges

The potential value of PVAT as a therapeutic target is still unclear. Interventions that could enhance the production of adiponectin from PVAT could be useful against vascular disease.<sup>36,38</sup> although there are complex interactions between the vascular wall and PVAT, with multiple messengers involved in their cross-talk, temporally and spatially regulated by vascular disease development. Identifying those bioactive molecules or signaling pathways amenable to treatment remains an open challenge.

Coronary PVAT has been in the spotlight of research over the last years, but the role of PVAT in other vascular beds remains less well studied. The study of peri-aortic<sup>122</sup> or peri-femoral AT<sup>123</sup> may provide new insights into the development, prevention and treatment of aortic aneurysms or peripheral arterial disease respectively.

Therapeutic interventions could be potentially deployed either by targeting PVAT itself or by using PVAT imaging as a tool of precision medicine. Indeed, peri-coronary FAI-Score provides a regulatory clear metric of coronary inflammation and it can be used for personalized absolute risk prediction.<sup>39</sup> It has therefore been suggested that patients with high residual inflammatory risk identified by this approach, could be candidates for anti-inflammatory strategies targeting vascular inflammation, like statins, colchicine or other novel therapeutics.<sup>39,124</sup>

The pipeline for the validation of new imaging biomarkers must be rigorous before their clinical use. First and foremost is the technical validation of any new imaging biomarker i.e., the repeatability and accuracy of its measurement as well as a steady diagnostic performance across platforms or acquisition parameters.<sup>79</sup> Equally important are the issues of biological and clinical

validation of any new biomarker, and the confirmation of its generalizability, and diagnostic/prognostic value in large clinical datasets.<sup>95,125</sup>

All these issues are even more pressing in the case of AI-derived imaging biomarkers, which are typically derived out of deep learning techniques and the mining of high-dimensional data, in a 'black-box' process.<sup>125</sup> The use of AI biomarkers in clinical practice requires rigorous validation and the use of very large datasets to ensure worldwide applicability, as mandated by the Medical Device Regulations in Europe and the Food and Drugs Administration in the USA.

## **Conclusions**

This Clinical Consensus Statement presents the optimum terminology for the definition of the various AT depots, and particularly consensus definitions of perivascular AT. It provides an overview of the range of adipokines secreted by PVAT, with either protective or detrimental effects on the cardiovascular system, which can be used as therapeutic targets for the prevention and treatment of cardiovascular diseases. Although existing pharmacotherapies affect the biology of AT, targeting PVAT specifically is more challenging, as it requires local delivery of the intervention. This position paper also discusses the role of PVAT as a "biosensor" of vascular inflammation and presents recent imaging technologies that visualise PVAT around the coronary arteries and quantify indirectly the degree of coronary inflammation from routine CCTA scans, predicting future cardiovascular events. We also present the current clinical applicability of these imaging biomarkers, and highlight the importance of regulatory clearance for any medical device that provides these measurements.

In conclusion, PVAT is currently used as a source of imaging biomarkers directly applicable in clinical practice, and this role will be further enhanced with the advance of AI-powered image analysis in the near future. However, further research is needed before we can use PVAT as a therapeutic target in cardiovascular diseases.

## REFERENCES

1. Badimon L, Bugiardini R, Cenko E, Cubedo J, Dorobantu M, Duncker DJ, *et al.* Position paper of the European Society of Cardiology-working group of coronary pathophysiology and microcirculation: obesity and heart disease. *Eur Heart J* 2017;**38**:1951-1958. doi: 10.1093/eurheartj/ehx181
2. Aune D, Sen A, Prasad M, Norat T, Janszky I, Tonstad S, *et al.* BMI and all cause mortality: systematic review and non-linear dose-response meta-analysis of 230 cohort studies with 3.74 million deaths among 30.3 million participants. *BMJ* 2016;**353**:i2156. doi: 10.1136/bmj.i2156
3. Aguilar M, Rose RA, Takawale A, Nattel S, Reilly S. New aspects of endocrine control of atrial fibrillation and possibilities for clinical translation. *Cardiovasc Res* 2021;**117**:1645-1661. doi: 10.1093/cvr/cvab080
4. Antonopoulos AS, Tousoulis D. The molecular mechanisms of obesity paradox. *Cardiovasc Res* 2017;**113**:1074-1086. doi: 10.1093/cvr/cvx106
5. Collaboration NCDRF. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 2017;**390**:2627-2642. doi: 10.1016/S0140-6736(17)32129-3
6. Oikonomou EK, Antoniadou C. The role of adipose tissue in cardiovascular health and disease. *Nat Rev Cardiol* 2019;**16**:83-99. doi: 10.1038/s41569-018-0097-6
7. Rosen ED, Spiegelman BM. What we talk about when we talk about fat. *Cell* 2014;**156**:20-44. doi: 10.1016/j.cell.2013.12.012
8. Antonopoulos AS, Antoniadou C. The role of epicardial adipose tissue in cardiac biology: classic concepts and emerging roles. *J Physiol* 2017;**595**:3907-3917. doi: 10.1113/JP273049
9. Turaihi AH, Serne EH, Molthoff CFM, Koning JJ, Knol J, Niessen HW, *et al.* Perivascular Adipose Tissue Controls Insulin-Stimulated Perfusion, Mitochondrial Protein Expression, and Glucose Uptake in Muscle Through Adipomuscular Arterioles. *Diabetes* 2020;**69**:603-613. doi: 10.2337/db18-1066
10. Antoniadou C, Antonopoulos AS, Deanfield J. Imaging residual inflammatory cardiovascular risk. *Eur Heart J* 2020;**41**:748-758. doi: 10.1093/eurheartj/ehz474
11. Costa RM, Neves KB, Tostes RC, Lobato NS. Perivascular Adipose Tissue as a Relevant Fat Depot for Cardiovascular Risk in Obesity. *Front Physiol* 2018;**9**:253. doi: 10.3389/fphys.2018.00253
12. Antonopoulos AS, Sanna F, Sabharwal N, Thomas S, Oikonomou EK, Herdman L, *et al.* Detecting human coronary inflammation by imaging perivascular fat. *Sci Transl Med* 2017;**9**. doi: 10.1126/scitranslmed.aal2658
13. Oikonomou EK, Williams MC, Kotanidis CP, Desai MY, Marwan M, Antonopoulos AS, *et al.* A novel machine learning-derived radiotranscriptomic signature of perivascular fat improves cardiac risk prediction using coronary CT angiography. *Eur Heart J* 2019;**40**:3529-3543. doi: 10.1093/eurheartj/ehz592
14. Bengs S, Haider A, Warnock GI, Fiechter M, Pargaetzi Y, Rampidis G, *et al.* Quantification of perivascular inflammation does not provide incremental prognostic value over myocardial perfusion imaging and calcium scoring. *Eur J Nucl Med Mol Imaging* 2021;**48**:1806-1812. doi: 10.1007/s00259-020-05106-0



15. Dai X, Yu L, Lu Z, Shen C, Tao X, Zhang J. Serial change of perivascular fat attenuation index after statin treatment: Insights from a coronary CT angiography follow-up study. *Int J Cardiol* 2020;**319**:144-149. doi: 10.1016/j.ijcard.2020.06.008
16. Antonopoulos AS, Antoniades C. Perivascular Fat Attenuation Index by Computed Tomography as a Metric of Coronary Inflammation. *J Am Coll Cardiol* 2018;**71**:2708-2709. doi: 10.1016/j.jacc.2018.03.511
17. Oikonomou EK, Desai MY, Marwan M, Kotanidis CP, Antonopoulos AS, Schottlander D, *et al.* Perivascular Fat Attenuation Index Stratifies Cardiac Risk Associated With High-Risk Plaques in the CRISP-CT Study. *J Am Coll Cardiol* 2020;**76**:755-757. doi: 10.1016/j.jacc.2020.05.078
18. Mazurek T, Kobylecka M, Zielenkiewicz M, Kurek A, Kochman J, Filipiak KJ, *et al.* PET/CT evaluation of F-FDG uptake in pericoronary adipose tissue in patients with stable coronary artery disease: Independent predictor of atherosclerotic lesions' formation? *J Nucl Cardiol* 2016. doi: 10.1007/s12350-015-0370-6
19. Goeller M, Achenbach S, Cadet S, Kwan AC, Commandeur F, Slomka PJ, *et al.* Pericoronary Adipose Tissue Computed Tomography Attenuation and High-Risk Plaque Characteristics in Acute Coronary Syndrome Compared With Stable Coronary Artery Disease. *JAMA Cardiol* 2018;**3**:858-863. doi: 10.1001/jamacardio.2018.1997
20. Iacobellis G. Epicardial adipose tissue in contemporary cardiology. *Nat Rev Cardiol* 2022. doi: 10.1038/s41569-022-00679-9
21. Antonopoulos AS, Margaritis M, Coutinho P, Digby J, Patel R, Psarros C, *et al.* Reciprocal effects of systemic inflammation and brain natriuretic peptide on adiponectin biosynthesis in adipose tissue of patients with ischemic heart disease. *Arterioscler Thromb Vasc Biol* 2014;**34**:2151-2159. doi: 10.1161/ATVBAHA.114.303828
22. Lu C, Su LY, Lee RM, Gao YJ. Alterations in perivascular adipose tissue structure and function in hypertension. *Eur J Pharmacol* 2011;**656**:68-73. doi: 10.1016/j.ejphar.2011.01.023
23. Antonopoulos AS, Sanna F, Sabharwal N, Thomas S, Oikonomou EK, Herdman L, *et al.* Detecting human coronary inflammation by imaging perivascular fat. 2017;**9**. doi:
24. Kim HW, Shi H, Winkler MA, Lee R, Weintraub NL. Perivascular Adipose Tissue and Vascular Perturbation/Atherosclerosis. *Arterioscler Thromb Vasc Biol* 2020;**40**:2569-2576. doi: 10.1161/ATVBAHA.120.312470
25. Mancio J, Oikonomou EK, Antoniades C. Perivascular adipose tissue and coronary atherosclerosis. *Heart* 2018;**104**:1654-1662. doi: 10.1136/heartjnl-2017-312324
26. Kumar RK, Jin Y, Watts SW, Rockwell CE. Naive, Regulatory, Activated, and Memory Immune Cells Co-exist in PVATs That Are Comparable in Density to Non-PVAT Fats in Health. *Front Physiol* 2020;**11**:58. doi: 10.3389/fphys.2020.00058
27. Gil-Ortega M, Somoza B, Huang Y, Gollasch M, Fernandez-Alfonso MS. Regional differences in perivascular adipose tissue impacting vascular homeostasis. *Trends Endocrinol Metab* 2015;**26**:367-375. doi: 10.1016/j.tem.2015.04.003
28. Nosalski R, Guzik TJ. Perivascular adipose tissue inflammation in vascular disease. *Br J Pharmacol* 2017;**174**:3496-3513. doi: 10.1111/bph.13705
29. Withers SB, Forman R, Meza-Perez S, Sorobetea D, Sitnik K, Hopwood T, *et al.* Eosinophils are key regulators of perivascular adipose tissue and vascular functionality. *Sci Rep* 2017;**7**:44571. doi: 10.1038/srep44571
30. Szasz T, Bomfim GF, Webb RC. The influence of perivascular adipose tissue on vascular homeostasis. *Vasc Health Risk Manag* 2013;**9**:105-116. doi: 10.2147/VHRM.S33760

31. Yamawaki H, Kuramoto J, Kameshima S, Usui T, Okada M, Hara Y. Omentin, a novel adipocytokine inhibits TNF-induced vascular inflammation in human endothelial cells. *Biochem Biophys Res Commun* 2011;**408**:339-343. doi: 10.1016/j.bbrc.2011.04.039
32. Chun HJ, Ali ZA, Kojima Y, Kundu RK, Sheikh AY, Agrawal R, *et al.* Apelin signaling antagonizes Ang II effects in mouse models of atherosclerosis. *J Clin Invest* 2008;**118**:3343-3354. doi: 10.1172/JCI34871
33. Kotanidis CP, Antoniades C. Perivascular fat imaging by computed tomography (CT): a virtual guide. *Br J Pharmacol* 2021;**178**:4270-4290. doi: 10.1111/bph.15634
34. Li X, Ma Z, Zhu YZ. Regional Heterogeneity of Perivascular Adipose Tissue: Morphology, Origin, and Secretome. *Front Pharmacol* 2021;**12**:697720. doi: 10.3389/fphar.2021.697720
35. Saxton SN, Clark BJ, Withers SB, Eringa EC, Heagerty AM. Mechanistic Links Between Obesity, Diabetes, and Blood Pressure: Role of Perivascular Adipose Tissue. *Physiol Rev* 2019;**99**:1701-1763. doi: 10.1152/physrev.00034.2018
36. Antonopoulos AS, Margaritis M, Coutinho P, Shirodaria C, Psarros C, Herdman L, *et al.* Adiponectin as a link between type 2 diabetes and vascular NADPH oxidase activity in the human arterial wall: the regulatory role of perivascular adipose tissue. *Diabetes* 2015;**64**:2207-2219. doi: 10.2337/db14-1011
37. Antonopoulos AS, Margaritis M, Verheule S, Recalde A, Sanna F, Herdman L, *et al.* Mutual Regulation of Epicardial Adipose Tissue and Myocardial Redox State by PPAR-gamma/Adiponectin Signalling. *Circ Res* 2016;**118**:842-855. doi: 10.1161/CIRCRESAHA.115.307856
38. Margaritis M, Antonopoulos AS, Digby J, Lee R, Reilly S, Coutinho P, *et al.* Interactions between vascular wall and perivascular adipose tissue reveal novel roles for adiponectin in the regulation of endothelial nitric oxide synthase function in human vessels. *Circulation* 2013;**127**:2209-2221. doi: 10.1161/CIRCULATIONAHA.112.001133
39. Oikonomou EK, Antonopoulos AS, Schottlander D, Marwan M, Mathers C, Tomlins P, *et al.* Standardized measurement of coronary inflammation using cardiovascular computed tomography: integration in clinical care as a prognostic medical device. *Cardiovasc Res* 2021;**117**:2677-2690. doi: 10.1093/cvr/cvab286
40. Murray EC, Nosalski R, MacRitchie N, Tomaszewski M, Maffia P, Harrison DG, *et al.* Therapeutic targeting of inflammation in hypertension: from novel mechanisms to translational perspective. *Cardiovasc Res* 2021;**117**:2589-2609. doi: 10.1093/cvr/cvab330
41. Chen JX, Stinnett A. Critical role of the NADPH oxidase subunit p47phox on vascular TLR expression and neointimal lesion formation in high-fat diet-induced obesity. *Lab Invest* 2008;**88**:1316-1328. doi: 10.1038/labinvest.2008.92
42. Guzik TJ, Skiba DS, Touyz RM, Harrison DG. The role of infiltrating immune cells in dysfunctional adipose tissue. *Cardiovasc Res* 2017;**113**:1009-1023. doi: 10.1093/cvr/cvx108
43. Akoumianakis I, Sanna F, Margaritis M, Badi I, Akawi N, Herdman L, *et al.* Adipose tissue-derived WNT5A regulates vascular redox signaling in obesity via USP17/RAC1-mediated activation of NADPH oxidases. *Sci Transl Med* 2019;**11**. doi: 10.1126/scitranslmed.aav5055
44. Wu H, Ghosh S, Perrard XD, Feng L, Garcia GE, Perrard JL, *et al.* T-cell accumulation and regulated on activation, normal T cell expressed and secreted upregulation in adipose tissue in obesity. *Circulation* 2007;**115**:1029-1038. doi: 10.1161/CIRCULATIONAHA.106.638379

45. Iacobini C, Pugliese G, Blasetti Fantauzzi C, Federici M, Menini S. Metabolically healthy versus metabolically unhealthy obesity. *Metabolism* 2019;**92**:51-60. doi: 10.1016/j.metabol.2018.11.009
46. Chang L, Milton H, Eitzman DT, Chen YE. Paradoxical roles of perivascular adipose tissue in atherosclerosis and hypertension. *Circ J* 2013;**77**:11-18. doi: 10.1253/circj.cj-12-1393
47. Lim S, Meigs JB. Links between ectopic fat and vascular disease in humans. *Arterioscler Thromb Vasc Biol* 2014;**34**:1820-1826. doi: 10.1161/ATVBAHA.114.303035
48. Despres JP. Body fat distribution and risk of cardiovascular disease: an update. *Circulation* 2012;**126**:1301-1313. doi: 10.1161/CIRCULATIONAHA.111.067264
49. Koentges C, Konig A, Pfeil K, Holscher ME, Schnick T, Wende AR, *et al.* Myocardial mitochondrial dysfunction in mice lacking adiponectin receptor 1. *Basic Res Cardiol* 2015;**110**:37. doi: 10.1007/s00395-015-0495-4
50. Greenstein AS, Khavandi K, Withers SB, Sonoyama K, Clancy O, Jeziorska M, *et al.* Local inflammation and hypoxia abolish the protective anticontractile properties of perivascular fat in obese patients. *Circulation* 2009;**119**:1661-1670. doi: 10.1161/CIRCULATIONAHA.108.821181
51. Meijer RI, Hoevenaars FPM, Serne EH, Yudkin JS, Kokhuis TJA, Weijers EM, *et al.* JNK2 in myeloid cells impairs insulin's vasodilator effects in muscle during early obesity development through perivascular adipose tissue dysfunction. *Am J Physiol Heart Circ Physiol* 2019;**317**:H364-h374. doi: 10.1152/ajpheart.00663.2018
52. Azul L, Leandro A, Boroumand P, Klip A, Seica R, Sena CM. Increased inflammation, oxidative stress and a reduction in antioxidant defense enzymes in perivascular adipose tissue contribute to vascular dysfunction in type 2 diabetes. *Free Radic Biol Med* 2020;**146**:264-274. doi: 10.1016/j.freeradbiomed.2019.11.002
53. Chatterjee TK, Stoll LL, Denning GM, Harrelson A, Blomkalns AL, Idelman G, *et al.* Proinflammatory phenotype of perivascular adipocytes: influence of high-fat feeding. *Circ Res* 2009;**104**:541-549. doi: 10.1161/CIRCRESAHA.108.182998
54. Drummond GR, Vinh A, Guzik TJ, Sobey CG. Immune mechanisms of hypertension. *Nat Rev Immunol* 2019;**19**:517-532. doi: 10.1038/s41577-019-0160-5
55. Zhu X, Wang Y, Zhu L, Zhu Y, Zhang K, Wang L, *et al.* Class A1 scavenger receptor prevents obesity-associated blood pressure elevation through suppressing overproduction of vascular endothelial growth factor B in macrophages. *Cardiovasc Res* 2021;**117**:547-560. doi: 10.1093/cvr/cvaa030
56. Guzik TJ, Hoch NE, Brown KA, McCann LA, Rahman A, Dikalov S, *et al.* Role of the T cell in the genesis of angiotensin II induced hypertension and vascular dysfunction. *J Exp Med* 2007;**204**:2449-2460. doi: 10.1084/jem.20070657
57. Nosalski R, Siedlinski M, Denby L, McGinnigle E, Nowak M, Cat AND, *et al.* T-Cell-Derived miRNA-214 Mediates Perivascular Fibrosis in Hypertension. *Circ Res* 2020;**126**:988-1003. doi: 10.1161/CIRCRESAHA.119.315428
58. Frangogiannis NG. Cardiac fibrosis. *Cardiovasc Res* 2021;**117**:1450-1488. doi: 10.1093/cvr/cvaa324
59. Mikolajczyk TP, Nosalski R, Szczepaniak P, Budzyn K, Osmenda G, Skiba D, *et al.* Role of chemokine RANTES in the regulation of perivascular inflammation, T-cell accumulation, and vascular dysfunction in hypertension. *FASEB J* 2016;**30**:1987-1999. doi: 10.1096/fj.201500088R
60. Nosalski R, Guzik TJ. IL-15 and IL-7: keys to dysregulated inflammation in acute coronary syndromes. *Cardiovasc Res* 2021;**117**:1806-1808. doi: 10.1093/cvr/cvab189

61. Hall JE, Mouton AJ, da Silva AA, Omoto ACM, Wang Z, Li X, *et al.* Obesity, kidney dysfunction, and inflammation: interactions in hypertension. *Cardiovasc Res* 2021;**117**:1859-1876. doi: 10.1093/cvr/cvaa336
62. Ahmad AA, Randall MD, Roberts RE. Sex differences in the role of phospholipase A2 - dependent arachidonic acid pathway in the perivascular adipose tissue function in pigs. *J Physiol* 2017;**595**:6623-6634. doi: 10.1113/JP274831
63. Ahmad AA, Randall MD, Roberts RE. Sex differences in the regulation of porcine coronary artery tone by perivascular adipose tissue: a role of adiponectin? *Br J Pharmacol* 2017;**174**:2773-2783. doi: 10.1111/bph.13902
64. Meyer MR, Barton M. Role of Perivascular Adipose Tissue for Sex Differences in Coronary Artery Disease and Spontaneous Coronary Artery Dissection (SCAD). *Endocrine and Metabolic Science* 2021;**2**:100068. doi: <https://doi.org/10.1016/j.endmts.2020.100068>
65. Wang D, Wang C, Wu X, Zheng W, Sandberg K, Ji H, *et al.* Endothelial dysfunction and enhanced contractility in microvessels from ovariectomized rats: roles of oxidative stress and perivascular adipose tissue. *Hypertension* 2014;**63**:1063-1069. doi: 10.1161/HYPERTENSIONAHA.113.02284
66. Chatterjee TK, Stoll LL, Denning GM, Harrelson A, Blomkalns AL, Idelman G, *et al.* Proinflammatory Phenotype of Perivascular Adipocytes: Influence of High-Fat Feeding. *Circulation Research* 2009;**104**:541-549. doi: 10.1161/ATVBAHA.120.314129
67. Emanuel AL, Meijer RI, Woerdeman J, Raalte DHv, Diamant M, Kramer MHH, *et al.* Effects of a Hypercaloric and Hypocaloric Diet on Insulin-Induced Microvascular Recruitment, Glucose Uptake, and Lipolysis in Healthy Lean Men. *Arteriosclerosis, Thrombosis, and Vascular Biology* 2020;**40**:1695-1704. doi: doi:10.1161/ATVBAHA.120.314129
68. Withers SB, gabiti-Rosei C, Livingstone DM, Little MC, Aslam R, Malik RA, *et al.* Macrophage activation is responsible for loss of anticontractile function in inflamed perivascular fat. *Arterioscler. Thromb. Vasc. Biol* 2011;**31**:908-913. doi: 10.1161/ATVBAHA.111.22284
69. Vendrell J, El Bekay R, Peral B, Garcia-Fuentes E, Megia A, Macias-Gonzalez M, *et al.* Study of the potential association of adipose tissue GLP-1 receptor with obesity and insulin resistance. *Endocrinology* 2011;**152**:4072-4079. doi: 10.1210/en.2011-1070
70. Engelbrechtsen L, Lundgren J, Wewer Albrechtsen NJ, Mahendran Y, Iepsen EW, Finocchietto P, *et al.* Treatment with liraglutide may improve markers of CVD reflected by reduced levels of apoB. *Obes Sci Pract* 2017;**3**:425-433. doi: 10.1002/osp4.133
71. Jujic A, Nilsson PM, Atabaki-Pasdar N, Dieden A, Tuomi T, Franks PW, *et al.* Glucose-Dependent Insulinotropic Peptide in the High-Normal Range Is Associated With Increased Carotid Intima-Media Thickness. *Diabetes Care* 2021;**44**:224-230. doi: 10.2337/dc20-1318
72. Akawi N, Checa A, Antonopoulos AS, Akoumianakis I, Daskalaki E, Kotanidis CP, *et al.* Fat-Secreted Ceramides Regulate Vascular Redox State and Influence Outcomes in Patients With Cardiovascular Disease. *J Am Coll Cardiol* 2021;**77**:2494-2513. doi: 10.1016/j.jacc.2021.03.314
73. Lundgren JR, Janus C, Jensen SBK, Juhl CR, Olsen LM, Christensen RM, *et al.* Healthy Weight Loss Maintenance with Exercise, Liraglutide, or Both Combined. *N Engl J Med* 2021;**384**:1719-1730. doi: 10.1056/NEJMoa2028198
74. Xu L, Ota T. Emerging roles of SGLT2 inhibitors in obesity and insulin resistance: Focus on fat browning and macrophage polarization. *Adipocyte* 2018;**7**:121-128. doi: 10.1080/21623945.2017.1413516

75. Mullens W, Martens P. Empagliflozin-Induced Changes in Epicardial Fat. 2021;**9**:590-593. doi: doi:10.1016/j.jchf.2021.05.006
76. Kagota S, Maruyama-Fumoto K, McGuire JJ, Shinozuka K. A Sodium Glucose Cotransporter 2 Inhibitor Fails to Improve Perivascular Adipose Tissue-Mediated Modulation of Vasodilation and Cardiac Function in Rats With Metabolic Syndrome. *J Cardiovasc Pharmacol Ther* 2021;**26**:480-489. doi: 10.1177/10742484211001853
77. Wojcicka G, Jamroz-Wisniewska A, Atanasova P, Chaldakov GN, Chylinska-Kula B, Beltowski J. Differential effects of statins on endogenous H<sub>2</sub>S formation in perivascular adipose tissue. *Pharmacol Res* 2011;**63**:68-76. doi: S1043-6618(10)00203-3 [pii] 10.1016/j.phrs.2010.10.011
78. Xia N, Li H. The role of perivascular adipose tissue in obesity-induced vascular dysfunction. *Br J Pharmacol* 2017;**174**:3425-3442. doi: 10.1111/bph.13650
79. West H, Siddique M, Williams MC, Volpe L, Desai R, Lyasheva M, *et al.* Deep-Learning for Epicardial Adipose Tissue Assessment with Computed Tomography: Implications for Cardiovascular Risk Prediction. *JACC Cardiovasc Imaging* 2022 (*accepted*). doi:
80. In: P. L, ed. Braunwald's Heart Disease, Elsevier; 12th edition (November 29, 2021), Chapter 20 - Cardiac Computed Tomography.
81. Oikonomou EK, Marwan M, Desai MY, Mancio J, Alashi A, Hutt Centeno E, *et al.* Non-invasive detection of coronary inflammation using computed tomography and prediction of residual cardiovascular risk (the CRISP CT study): a post-hoc analysis of prospective outcome data. *Lancet* 2018;**392**:929-939. doi: 10.1016/S0140-6736(18)31114-0
82. Marwan M, Hell M, Schuhback A, Gauss S, Bittner D, Pflederer T, *et al.* CT Attenuation of Pericoronary Adipose Tissue in Normal Versus Atherosclerotic Coronary Segments as Defined by Intravascular Ultrasound. *J Comput Assist Tomogr* 2017;**41**:762-767. doi: 10.1097/RCT.0000000000000589
83. Lee SE, Sung JM, Andreini D, Al-Mallah MH, Budoff MJ, Cademartiri F, *et al.* Association Between Changes in Perivascular Adipose Tissue Density and Plaque Progression. *JACC Cardiovasc Imaging* 2022;**15**:1760-1767. doi: 10.1016/j.jcmg.2022.04.016
84. Gaibazzi N, Martini C, Botti A, Pinazzi A, Bottazzi B, Palumbo AA. Coronary Inflammation by Computed Tomography Pericoronary Fat Attenuation in MINOCA and Tako-Tsubo Syndrome. *J Am Heart Assoc* 2019;**8**:e013235. doi: 10.1161/JAHA.119.013235
85. Sugiyama T, Kanaji Y, Hoshino M, Yamaguchi M, Hada M, Ohya H, *et al.* Determinants of Pericoronary Adipose Tissue Attenuation on Computed Tomography Angiography in Coronary Artery Disease. *J Am Heart Assoc* 2020;**9**:e016202. doi: 10.1161/JAHA.120.016202
86. Kwiecinski J, Dey D, Cadet S, Lee SE, Otaki Y, Huynh PT, *et al.* Peri-Coronary Adipose Tissue Density Is Associated With (18)F-Sodium Fluoride Coronary Uptake in Stable Patients With High-Risk Plaques. *JACC Cardiovasc Imaging* 2019;**12**:2000-2010. doi: 10.1016/j.jcmg.2018.11.032
87. Hoshino M, Yang S, Sugiyama T, Zhang J, Kanaji Y, Yamaguchi M, *et al.* Peri-coronary inflammation is associated with findings on coronary computed tomography angiography and fractional flow reserve. *J Cardiovasc Comput Tomogr* 2020;**14**:483-489. doi: 10.1016/j.jcct.2020.02.002
88. Yan H, Zhao N, Geng W, Hou Z, Gao Y, Lu B. Pericoronary fat attenuation index and coronary plaque quantified from coronary computed tomography angiography identify ischemia-causing lesions. *Int J Cardiol* 2022;**357**:8-13. doi: 10.1016/j.ijcard.2022.03.033

89. Leipsic J, Abbara S, Achenbach S, Cury R, Earls JP, Mancini GJ, *et al.* SCCT guidelines for the interpretation and reporting of coronary CT angiography: a report of the Society of Cardiovascular Computed Tomography Guidelines Committee. *J Cardiovasc Comput Tomogr* 2014;**8**:342-358. doi: 10.1016/j.jcct.2014.07.003
90. Boussoussou M, Vattay B, Szilveszter B, Simon J, Lin A, Vecsey-Nagy M, *et al.* The effect of patient and imaging characteristics on coronary CT angiography assessed pericoronary adipose tissue attenuation and gradient. *J Cardiovasc Comput Tomogr* 2022. doi: 10.1016/j.jcct.2022.09.006
91. Elnabawi Y, Oikonomou E, Dey A, Mancio J, Rodante J, Gelfand J, *et al.* Treatment with biologic therapy in psoriasis is associated with a reduction in coronary artery inflammation, assessed by perivascular Fat Attenuation Index. *J Am Coll Cardiol* 2019;**73**:87. doi: 10.1016/S0735-1097(19)30695-3 %J Journal of the American College of Cardiology
92. Desai MY. Noninvasive detection of perivascular inflammation by coronary computed tomography in the CRISP-CT study and its implications for residual cardiovascular risk. *Cardiovasc Res* 2019;**115**:e3-e4. doi: 10.1093/cvr/cvy282
93. Tzolos E, Williams MC, McElhinney P, Lin A, Grodecki K, Flores Tomasino G, *et al.* Pericoronary Adipose Tissue Attenuation, Low-Attenuation Plaque Burden, and 5-Year Risk of Myocardial Infarction. *JACC Cardiovasc Imaging* 2022. doi: 10.1016/j.jcmg.2022.02.004
94. Antoniadis C, Kotanidis CP, Berman DS. State-of-the-art review article. Atherosclerosis affecting fat: What can we learn by imaging perivascular adipose tissue? *J Cardiovasc Comput Tomogr* 2019;**13**:288-296. doi: 10.1016/j.jcct.2019.03.006
95. Oikonomou EK, Siddique M, Antoniadis C. Artificial intelligence in medical imaging: A radiomic guide to precision phenotyping of cardiovascular disease. *Cardiovasc Res* 2020;**116**:2040-2054. doi: 10.1093/cvr/cvaa021
96. Mahabadi AA, Bamberg F, Toepker M, Schlett CL, Rogers IS, Nagurney JT, *et al.* Association of aortic valve calcification to the presence, extent, and composition of coronary artery plaque burden: from the Rule Out Myocardial Infarction using Computer Assisted Tomography (ROMICAT) trial. *Am Heart J* 2009;**158**:562-568. doi: 10.1016/j.ahj.2009.07.027
97. Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Back M, *et al.* 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2021;**42**:3227-3337. doi: 10.1093/eurheartj/ehab484
98. Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, *et al.* 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts)Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J* 2016;**37**:2315-2381. doi: 10.1093/eurheartj/ehw106
99. Dai X, Hou Y, Tang C, Lu Z, Shen C, Zhang L, *et al.* Long-term prognostic value of the serial changes of CT-derived fractional flow reserve and perivascular fat attenuation index. *Quant Imaging Med Surg* 2022;**12**:752-765. doi: 10.21037/qims-21-424
100. Oikonomou EK, Antonopoulos AS, Schottlander D, Marwan M, Mathers C, Tomlins P, *et al.* Standardised measurement of coronary inflammation using cardiovascular CT: integration in clinical care as a prognostic medical device. *Cardiovasc Res* 2021. doi: 10.1093/cvr/cvab286
101. Antoniadis C, Shirodaria C. Detecting Coronary Inflammation With Perivascular Fat Attenuation Imaging. 2019;**12**:2011-2014. doi: doi:10.1016/j.jcmg.2018.12.024

102. Bergstrom G, Berglund G, Blomberg A, Brandberg J, Engstrom G, Engvall J, *et al.* The Swedish CardioPulmonary BioImage Study: objectives and design. *J Intern Med* 2015;**278**:645-659. doi: 10.1111/joim.12384
103. investigators S-H. CT coronary angiography in patients with suspected angina due to coronary heart disease (SCOT-HEART): an open-label, parallel-group, multicentre trial. *Lancet* 2015;**385**:2383-2391. doi: 10.1016/S0140-6736(15)60291-4
104. Kolossvary M, Karady J, Szilveszter B, Kitslaar P, Hoffmann U, Merkely B, *et al.* Radiomic Features Are Superior to Conventional Quantitative Computed Tomographic Metrics to Identify Coronary Plaques With Napkin-Ring Sign. *Circ Cardiovasc Imaging* 2017;**10**. doi: 10.1161/CIRCIMAGING.117.006843
105. Shang J, Ma S, Guo Y, Yang L, Zhang Q, Xie F, *et al.* Prediction of acute coronary syndrome within 3 years using radiomics signature of pericoronary adipose tissue based on coronary computed tomography angiography. *Eur Radiol* 2022;**32**:1256-1266. doi: 10.1007/s00330-021-08109-z
106. Lin A, Kolossvary M, Cadet S, McElhinney P, Goeller M, Han D, *et al.* Radiomics-Based Precision Phenotyping Identifies Unstable Coronary Plaques From Computed Tomography Angiography. **0**. doi: doi:10.1016/j.jcmg.2021.11.016
107. Kotanidis CP, Xie C, Alexander D, Rodrigues JCL, Burnham K, Mentzer A, *et al.* Constructing custom-made radiotranscriptomic signatures of vascular inflammation from routine CT angiograms: a prospective outcomes validation study in COVID-19. *Lancet Digit Health* 2022;**4**:e705-e716. doi: 10.1016/S2589-7500(22)00132-7
108. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, *et al.* Antiinflammatory Therapy with Canakinumab for Atherosclerotic Disease. *N Engl J Med* 2017. doi: 10.1056/NEJMoa1707914
109. Nidorf SM, Fiolet ATL, Mosterd A, Eikelboom JW, Schut A, Opstal TSJ, *et al.* Colchicine in Patients with Chronic Coronary Disease. *N Engl J Med* 2020. doi: 10.1056/NEJMoa2021372
110. Tardif JC, Kouz S, Waters DD, Bertrand OF, Diaz R, Maggioni AP, *et al.* Efficacy and Safety of Low-Dose Colchicine after Myocardial Infarction. *N Engl J Med* 2019;**381**:2497-2505. doi: 10.1056/NEJMoa1912388
111. Elliott P, Cowie MR, Franke J, Ziegler A, Antoniades C, Bax J, *et al.* Development, validation, and implementation of biomarker testing in cardiovascular medicine state-of-the-art: proceedings of the European Society of Cardiology-Cardiovascular Round Table. *Cardiovasc Res* 2021;**117**:1248-1256. doi: 10.1093/cvr/cvaa272
112. Ridker PM. How Common Is Residual Inflammatory Risk? *Circ Res* 2017;**120**:617-619. doi: 10.1161/CIRCRESAHA.116.310527
113. Antoniades C, West HW. Coronary CT angiography as an 'one-stop shop' to detect the high-risk plaque and the vulnerable patient. *Eur Heart J* 2021;**42**:3853-3855. doi: 10.1093/eurheartj/ehab538
114. Antonopoulos AS, Angelopoulos A, Papanikolaou P, Simantiris S, Oikonomou EK, Vamvakaris K, *et al.* Biomarkers of Vascular Inflammation for Cardiovascular Risk Prognostication: A Meta-Analysis. *JACC Cardiovasc Imaging* 2021. doi: 10.1016/j.jcmg.2021.09.014
115. Randrianarisoa E, Stefan N, Fritsche A, Reis-Damaschk N, Hieronimus A, Balletshofer B, *et al.* Periaortic Adipose Tissue Compared With Peribrachial Adipose Tissue Mass as Markers

- and Possible Modulators of Cardiometabolic Risk. *Angiology* 2018;**69**:854-860. doi: 10.1177/0003319718755581
116. Ardissino M, McCracken C, Bard A, Antoniadis C, Neubauer S, Harvey NC, *et al.* Pericardial adiposity is independently linked to adverse cardiovascular phenotypes: a CMR study of 42 598 UK Biobank participants. *Eur Heart J Cardiovasc Imaging* 2022;**23**:1471-1481. doi: 10.1093/ehjci/jeac101
117. Toutouzas K, Antonopoulos AS, Koutagiar I, Skoumas I, Benetos G, Kafouris P, *et al.* Visceral adipose tissue phenotype and hypoadiponectinemia are associated with aortic Fluorine-18 fluorodeoxyglucose uptake in patients with familial dyslipidemias. *J Nucl Cardiol* 2022;**29**:1405-1414. doi: 10.1007/s12350-020-02472-y
118. Antoniadis C, Antonopoulos AS, Deanfield J. Imaging residual inflammatory cardiovascular risk. *Eur Heart J* 2019. doi: 10.1093/eurheartj/ehz474
119. Risch F, Schwarz F, Braun F, Bette S, Becker J, Scheurig-Muenkler C, *et al.* Assessment of epicardial adipose tissue on virtual non-contrast images derived from photon-counting detector coronary CTA datasets. *Eur Radiol* 2022. doi: 10.1007/s00330-022-09257-6
120. The AdipoRedOx study - Interactions between adipose tissue, vascular wall and myocardium in human atherosclerosis. <https://oxhvf.com/adiporedox/overview/>. (20/01/2023)
121. The VIP study - Validation study for introduction of novel computed tomography Imaging biomarkers in clinical Practice. <https://oxhvf.com/vip/overview/>. (20/01/2023)
122. Piacentini L, Werba JP, Bono E, Saccu C, Tremoli E, Spirito R, *et al.* Genome-Wide Expression Profiling Unveils Autoimmune Response Signatures in the Perivascular Adipose Tissue of Abdominal Aortic Aneurysm. *Arterioscler Thromb Vasc Biol* 2019;**39**:237-249. doi: 10.1161/ATVBAHA.118.311803
123. Piacentini L, Saccu C, Bono E, Tremoli E, Spirito R, Colombo GI, *et al.* Gene-expression profiles of abdominal perivascular adipose tissue distinguish aortic occlusive from stenotic atherosclerotic lesions and denote different pathogenetic pathways. *Sci Rep* 2020;**10**:6245. doi: 10.1038/s41598-020-63361-5
124. Antoniadis C, West HW. ESC CVD Prevention Guidelines 2021: improvements, controversies, and opportunities. *Cardiovasc Res* 2022;**118**:e17-e19. doi: 10.1093/cvr/cvab371
125. Antoniadis C, Oikonomou EK. Artificial intelligence in cardiovascular imaging-principles, expectations, and limitations. *Eur Heart J* 2021. doi: 10.1093/eurheartj/ehab678
126. Antoniadis C, Shirodaria C. Detecting Coronary Inflammation With Perivascular Fat Attenuation Imaging: Making Sense From Perivascular Attenuation Maps. *JACC Cardiovasc Imaging* 2019;**12**:2011-2014. doi: 10.1016/j.jcmg.2018.12.024



**Table 1. Definitions of adipose tissue depots: Clinical Consensus Statement**

**Subcutaneous adipose tissue (ScAT):** in the hypodermis (i.e., just below the skin); may include the subcutaneous thoracic, abdominal, gluteal or femoral adipose tissue.

**Visceral adipose tissue (VAT):** the adipose tissue located inside the abdominal or thoracic cavity, between the viscera; subclassified into thoracic adipose tissue (pericardial, non-pericardial, epicardial and perivascular) and abdominal adipose tissue (intraperitoneal, retroperitoneal).

**Thoracic adipose tissue (ThAT):** the visceral adipose tissue of the chest comprising of pericardial & non-pericardial thoracic adipose tissue depots, epicardial adipose and perivascular adipose tissue (around any vessel like the coronaries, the aorta, the internal mammary arteries etc).





**Epicardial adipose tissue (EpAT):** the adipose tissue between the visceral pericardium and the myocardium. That includes the perivascular adipose tissue around the coronary arteries.

**Pericardial adipose tissue:** the adipose tissue of the thorax, surrounding the heart located outside the pericardial sac (does not include peri-aortic adipose tissue)

**Paracardial adipose tissue:** a term that has been used to refer to adipose tissue close to the heart. Should be abandoned since it has a less clear meaning.

**Perivascular adipose tissue (PVAT):** the adipose tissue that lies around a vessel in a radial distance from the vascular wall equal to one luminal diameter of the adjacent vessel. For arteries with diameter > 2cm, perivascular adipose tissue is defined by convention as the adipose tissue within a distance of 2cm from the outer vessel wall.

**Table 2.** Morphological and functional differences between adipose tissue depots in the chest.

				
	<b>Subcutaneous adipose tissue (ScAT)</b>	<b>Pericardial adipose tissue (PAT)</b>	<b>Epicardial adipose tissue (EAT)</b>	<b>Perivascular adipose tissue (PVAT)</b>
<b>Relation with sex</b>	more in females	abdominal obesity in males	no difference	no difference
<b>Relation with age</b>	less with age	increasing with age	increasing with age	unclear
<b>Adipocyte size</b>	Large	Small	Comparable to PAT	Gradient in adipocyte size (smaller closer to the vascular wall)
<b>Adipocyte differentiation status</b>	Well differentiated	Poorly differentiated	Comparable to PAT	Dependent on vascular biology
<b>Vascular-stromal cell content</b>	Less rich vs. VAT	Rich vasculature High immune-cell content	Comparable to PAT	Dependent on vascular biology
<b>Biological profile</b>	Less metabolically active Thermogenic potential (beiging)	High metabolic activity Inflammatory profile Lipolytic potential	High metabolic activity Inflammatory profile Lipolytic potential	High secretory profile with non adipocytes (pre-adipocytes and inflammatory cells) driving its' secretome
<b>Association with metabolic status</b>	Neutral / protective	Strong. Negatively associated with IR)-biology driven by systemic stimuli	Strong (less than PAT)- biology largely driven by myocardial signals	Less than EAT-biology largely driven by vascular signals
<b>Association with cardiovascular risk</b>	Neutral	Strong	Strong	Strong

**Table 3.** Main bioactive components in PVAT: Cell origin and biological functions in the vascular bed

Bioactive component	Main cell origin	Vasomotor effects	Inflammation	Other biological effects
<b>Adipokine</b>				
Adiponectin	Adipocyte	eNOS-dependent vasodilatation	anti-inflammatory	Adenosine 5'-monophosphate-activated protein kinase (AMPK) activation
Leptin	Adipocyte	eNOS dependent Vasodilatation ET-1-mediated vasoconstriction	pro-inflammatory	Induce oxidative stress
Apelin	Adipocyte	Unclear: possibly vasodilator in small arteries and vasoconstrictor in veins (9,10)	Anti-inflammatory	Increase glucose utilization (insulin-independent effect) (12,13)
Omentin	Adipocyte, stromal cells	Anti-contractile effects	Anti-inflammatory	Increases NO bioavailability Anti-oxidant effects
Chemerin	Adipocyte	Vasoconstriction	Pro-inflammatory	Induce adhesion molecules in EC - Induces VSMC-proliferation
Visfatin	Stromal cells, Inflammatory cells (macrophage),	Vasoconstriction	Pro-inflammatory	Proliferative response in VSMC
Resistin	Inflammatory cells (macrophage), adipocyte	Vasoconstriction	Pro-inflammatory	Impairs insulin-stimulated glucose uptake
SFRP5	Adipocytes	Vasodilation		Reduces oxidative stress
Wnt5a	Adipocytes	Vasoconstriction		Mediates oxidative stress (NADPH oxidase activity)
<b>Cytokines &amp; Growth Factors</b>				





**Table 3.** Main bioactive components in PVAT: Cell origin and biological functions in the vascular bed

<b>Bioactive component</b>	<b>Main cell origin</b>	<b>Vasomotor effects</b>	<b>Inflammation</b>	<b>Other biological effects</b>
Interleukin-6 (IL-6)	Inflammatory cells (T-cells, macrophage); adipocyte	Vasoconstriction	Pro-inflammatory	Induce oxidative stress in EC and VSMC
Interleukin-10 (IL-10)	Inflammatory cells (T-cells, macrophage)	Unclear	anti-inflammatory	Reduce oxidative stress
IFN-gamma	Inflammatory cells (T-cells, NK-cells), stromal cells	Vasoconstriction	Pro-inflammatory	Impairs endothelium-dependent relaxation oxidative-stress
TNF-alpha	Inflammatory cells; stromal cells; adipocytes	-	Pro-inflammatory	ROS-production
MCP-1	Inflammatory cells, adipocytes	-	Pro-inflammatory	-
VEGF	Adipocytes	-	-	VSMC proliferation
<b>Other bioactive components</b>				
Hydrogen sulphide	Adipocytes, stroma cells (EC)	Vasodilatation		Anti-oxidant effects at physiological levels. At high levels lead to generation of free radicals
Nitric oxide	Endothelial cells (via eNOS) & Sympathetic nerves (via nNOS)	Vasorelaxation	Anti-inflammatory	-
Palmitic acid methyl ester	Adipocytes	Vasodilatation	Pro-inflammatory	Induces ICAM-1 expression
Angiotensin (1-7)	Adipocytes	Vasodilatation		Reduces oxidative stress
Ceramides (C16/C24)	Adipocytes	Vasoconstriction		Oxidative stress

eNOS: endothelial nitric oxide synthase; ET-1: endothelin-1; ICAM-1: Intercellular adhesion molecule-1; IFN: interferon; MCP: monocyte chemoattractant protein-1; NO: nitric oxide; ROS: reactive oxygen


species; TNF-alpha: tumor necrosis factor-alpha; VEGF: vascular endothelial growth factor; VSMC: vascular smooth muscle cell.

**Table 4.** Imaging modalities to assess adipose tissue


<b>Modality</b>	<b>Measurements Characteristics</b>	<b>Strengths</b>	<b>Limitations</b>
<b>Echocardiography</b> 	EAT thickness	<ul style="list-style-type: none"> <li>✓ Clinical availability</li> <li>✓ Low Cost</li> <li>✓ Lack of Radiation exposure</li> <li>✓ Ease of use</li> <li>✓ Safe</li> </ul>	<ul style="list-style-type: none"> <li>- No PVAT assessment</li> <li>- Lack of reproducibility</li> <li>- Low/intermediate spatial resolution</li> <li>- No 3D analysis / volumetric data</li> <li>- No qualitative analysis</li> <li>- Can image only epicardial or subcutaneous fat but not PVAT</li> <li>- Lack of experience</li> <li>- Image quality is dependent on subject characteristics</li> </ul>
<b>Computed Tomography</b> 	EAT thickness, area, volume, attenuation & quality i.e, Perivascular FAI, Perivascular Fat Radiomic Profile (FRP)	<ul style="list-style-type: none"> <li>✓ PVAT assessment</li> <li>✓ Clinical availability</li> <li>✓ Reproducibility</li> <li>✓ Excellent spatial resolution</li> <li>✓ Easy to perform</li> <li>✓ Quantitative and qualitative data</li> <li>✓ 3D volume data</li> <li>✓ Incorporation of AI and Machine learning algorithms</li> <li>✓ Simultaneous assessment of coronary calcium and coronary plaque</li> </ul>	<ul style="list-style-type: none"> <li>- Intermediate cost</li> <li>- Iodine contrast use</li> <li>- Radiation exposure</li> </ul>
<b>Magnetic Resonance Imaging</b> 	EAT thickness, area, volume, proton density fat fraction	<ul style="list-style-type: none"> <li>✓ Lack of radiation exposure</li> <li>✓ Reproducibility</li> <li>✓ Quantitative and qualitative data</li> <li>✓ 3D volume data</li> <li>✓ Simultaneous assessment of myocardial structure and function</li> </ul>	<ul style="list-style-type: none"> <li>- PVAT assessment for large arteries like the aorta/carotids only, not the coronaries</li> <li>- Lack of availability</li> <li>- Intermediate/high cost</li> <li>- Intermediate spatial resolution (inferior to CT)</li> <li>- Time consuming</li> </ul>
<b>Positron Emission Tomography</b> 	SUV, TBR	<ul style="list-style-type: none"> <li>✓ Gold standard to assess tissue inflammation by imaging</li> </ul>	<ul style="list-style-type: none"> <li>- Cannot be used to image EAT or PVAT</li> <li>- High cost</li> <li>- Lack of availability</li> <li>- Radiation exposure</li> </ul>

3D: Three dimensional; <sup>18</sup>F-fluorodeoxyglycose positron emission tomography; CT: Computed tomography; EAT: Epicardial adipose tissue; FAI: Fat Attenuation Index; PVAT: Perivascular adipose tissue; SUV: Standardized Uptake Values; TBR: target-to-background ratio


## Figure legends



**Figure 1.** Imaging of human adipose tissue depots by computed tomography. Axial (A), coronal (B) and sagittal (C) views of the chest showing subcutaneous, visceral abdominal, thoracic (including the pericardial and epicardial) adipose tissue. Reconstruction of perivascular adipose tissue (PVAT) around an epicardial coronary artery (D) and thoracic aorta (E). PVAT is defined as the adipose tissue lying within a radial distance from the outer vessel wall equal to the vessel diameter (or at a maximum distance of 2cm in the case of large vessels with diameter  $>2\text{cm}$ , like the aorta).




**Figure 2. (A)** Routes via which remote and local adipose tissue depots affect the cardiovascular system. **(B)** Illustration of the major components of the bidirectional interplay between the vascular wall and fat in the perivascular space. EC, endothelial cell; eNOS, endothelial NOS; PVAT, perivascular adipose tissue; VSMC, vascular smooth muscle cell (Reused with permission by Kotanidis & Antoniadis Br J Pharmacol. 2021;178:4270–4290). An animated video [web link](#)<sup>33</sup> presents how the cross-talk between PVAT and the vascular wall leads to changes in PVAT’s texture and composition, visible with computed tomography.




**Figure 3.** Schematic representation of the biology underlying the detection of coronary inflammation by imaging perivascular adipose tissue (PVAT). Illustration of a healthy artery and surrounding PVAT (bottom) and the resulting PVAT phenotype in states of high vascular wall inflammation (top). Biopsies of PVAT surrounding inflamed vessels demonstrates high macrophage infiltration and contains small adipocytes that when cultured in the presence of inflammatory cytokines, do not store intracellular lipids. Indeed, in the presence of vascular inflammation there is a steep change in adipocyte size and lipid:water content with increasing distance from the vascular wall due to the paracrine effects of vascular inflammation on surrounding perivascular adipose tissue (PVAT). These changes in water:lipid can be detected by coronary computed tomography angiography (CCTA), visualised by the Fat Attenuation Index (FAI) mapping of PVAT and quantified using FAI-Score. The latter is interpreted clinically by using age- and gender- nomograms, and when it is used in prognostic models with plaque and clinical risk factors, it provides a powerful way to calculate the patient's specific risk for cardiovascular events. An example of a patient with high FAI-Score at baseline who reduced vascular inflammation after 1 year's treatment with atorvastatin 40mg/od. Images obtained from Antonopoulos et al Science Transl Med 2017 (with permission). The animated images on the process by which vascular inflammation drives changes to PVAT visible by CCTA, have been obtained with permission from this animated video [web link](#).<sup>33</sup>






**Figure 4.** Pericoronary adipose tissue attenuation for detecting vascular inflammation. **(A)** Pericoronary Fat Attenuation Index (FAI) is increased around culprit lesions in acute coronary syndrome patients with evidence of inflammation as assessed by  $^{18}\text{F}$ -sodium fluoride ( $^{18}\text{F}$ -NaF) by positron emission

tomography/computed tomography (adapted *from* <sup>86</sup>). **(B)** Changes in FAI of perivascular adipose tissue (PVAT) around ruptured (culprit) atherosclerotic lesions of acute MI patients, non-culprit lesions of the same patients, or lesions in stable CAD patients.<sup>12</sup> **(C)** Temporal changes in FAI around a culprit lesion in patients with acute ST elevation myocardial infarction (n= 10). **(D)** Prognostic value of FAI for cardiac mortality and **(E)** cardiac mortality or nonfatal myocardial infarction in CRISP-CT study in a cohort of 2,040 patients undergoing diagnostic CCTA in Cleveland Clinic, US.<sup>81</sup> **(F)** Prognostic value of unadjusted coronary PVAT attenuation values in SCOT-HEART trial in 1,697 evaluable participants.<sup>81</sup> **(G)** The superior performance of fully weighted FAI vs. unadjusted PVAT density; unadjusted PVAT density had borderline predictive value for cardiac mortality, whereas fully weighted FAI led to improved risk prediction by 13.6% in CRISP-CT study.<sup>126</sup> **(H)** Standardization of coronary perivascular FAI and nomograms for age (FAI-Score) for left anterior descending artery (LAD) right coronary artery (RCA) and left circumflex coronary artery (LCx). FAI-Score informs on the coronary vessel-specific inflammation burden and the associated relative risk for a fatal cardiac event compared to the age group of reference. For instance, a young individual with no traditional risk factors may be at low absolute risk for a fatal cardiac event; however, a high FAI-Score may indicate increased relative risk for cardiac events in the long-term as a result of subclinical vascular inflammation.<sup>39</sup> All Figure panels were reproduced with permission from the authors/publishers.



**Figure 5.** Radiomic phenotyping of coronary perivascular adipose tissue (PVAT). (A) Coronary PVAT imaging features can be used to extract a high number of shape-, attenuation-, and texture-related statistics (i.e, radiomics). (B) Heatmap of scaled radiomic features (right) of all 1391 stable radiomic features in the SCOT-HEART population (n= 1575 patients). (C) Extracted radiomic features can be tested against the transcriptome profile of PVAT to identify features informing on distinct biological processes such as adipose tissue inflammation, vascularity, and fibrosis. (D) Selected radiomic features can form distinct radiomic signatures of PVAT, in this case Fat Radiomic Profile adipose tissue inflammation, vascularity, and fibrosis, which had strong independent predictive value for major adverse cardiac events (MACE) in the SCOT-HEART population (obtained from Oikonomou EO et al, Eur Heart J 2019, with permission).



**Figure 6.** Prognostic value of vascular inflammation biomarkers for major adverse cardiovascular events.

The added prognostic value of imaging biomarkers on top of patient risk profile and coronary atherosclerosis extent is greater than that of plasma biomarkers according to a meta-analysis of available published evidence from clinical studies (n=351,628 individuals). Perivascular adipose tissue (PVAT) imaging by CT was associated with the maximum added prognostic information among the studied vascular inflammation biomarkers. Obtained with permission from Antonopoulos AS et al; J Am Coll Cardiol Cardiovasc Imaging 2022.