

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Contribution of cover crop residue decomposition to peach tree nitrogen nutrition

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Tassinari A., da Silva L.O.S., Drescher G.L., de Oliveira R.A., Baldi E., de Melo G.W.B., et al. (2021). Contribution of cover crop residue decomposition to peach tree nitrogen nutrition. JOURNAL OF SOIL SCIENCE AND PLANT NUTRITION, 21(3), 2124-2136 [10.1007/s42729-021-00508-x].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/849636> since: 2022-01-31

*Published:*

DOI: <http://doi.org/10.1007/s42729-021-00508-x>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

## Contribution of cover crop residue decomposition to peach trees nitrogen nutrition

Tassinari<sup>a</sup>, Adriele; Silva<sup>a</sup>, Lincon Oliveira Stefanello da; Drescher<sup>b\*</sup>, Gerson Laerson; Oliveira<sup>c</sup>, Rodolfo Assis de; Baldi<sup>d</sup>, Elena; Melo<sup>e</sup>, George Wellington Bastos de; Zalamena<sup>f</sup>, Jovani; Mayer<sup>g</sup>, Newton Alex; Giacomini<sup>a</sup>, Sandro José; Carranca<sup>h</sup>, Corina Luisa de Abreu Fernandes; Ferreira<sup>i</sup>, Paulo Ademar Avelar; Paula<sup>a</sup>, Betania Vahl de; Loss<sup>j</sup>, Arcângelo; Toselli<sup>d</sup>, Moreno; Brunetto<sup>a</sup>, Gustavo

<sup>a</sup> Department of Soil Science, Federal University of Santa Maria, 1000 Roraima Ave, Santa Maria - RS 97105900, Brazil

<sup>b</sup> Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, 1366 West Altheimer Drive, Fayetteville - AR 72704, United States

<sup>c</sup> Department of Agricultural Sciences, Federal University of Santa Catarina, 1346 Admar Gonzaga Highway, Florianópolis - SC 88034000, Brazil

<sup>d</sup> Department of Agriculture and Food Sciences, University of Bologna. 44 Giuseppe Fanin St, Bologna 40127, Italy

<sup>e</sup> Brazilian Agricultural Research Corporation - Embrapa Grape and Wine, 515 Livramento St, Bento Gonçalves – RS 95701008, Brazil

<sup>f</sup> Federal Institute of Education, Science and Technology of Rio Grande do Sul, IFRS – Campus Restinga, 285 Alberto Hoffmann St, Porto Alegre - RS 91791508, Brazil

<sup>g</sup> Brazilian Agricultural Research Corporation - Embrapa Temperate Climate, BR 392 Highway Km 78, Pelotas – RS 96010971, Brazil

<sup>h</sup> National Institute of Agricultural and Veterinary Research, Republic Ave, Lisboa 2780157, Portugal

<sup>i</sup> Federal University of Santa Maria - Campus Cachoeira do Sul, 1345 Ernesto Barros St, Cachoeira do Sul - RS 96506-310, Brazil

<sup>j</sup> Department of Rural Engineering, Federal University of Santa Catarina, 1346 Admar Gonzaga Highway, Florianópolis - SC 88034000, Brazil

\*Corresponding author. *E-mail*: [gersondrescher@gmail.com](mailto:gersondrescher@gmail.com). Orcid ID: 0000-0002-3843-7908

**Acknowledgments** The authors gratefully acknowledge all students and staff for their contributions in the development of this research.

**Funding** This study was financed (grant and scholarships) in part by the Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Embrapa Clima Temperado - Pelotas, and Embrapa Uva e Vinho.

# Contribution of cover crop residue decomposition to peach tree nitrogen nutrition

## Abstract

*Purpose:* Cover crop nitrogen (N) cycling has an important role in agricultural production and contributes to peach (*Prunus persica* (L.) Batsch.) N nutrition. This study evaluated black oat (*Avena strigosa* Schreb) and ryegrass (*Lolium multiflorum* L.) residue decomposition dynamics, N recovery from cover crop residues, and N compartmentalization in peach tree organs.

*Methods:* A two-year field trial was developed with labeled (5 atom%  $^{15}\text{N}$  excess) cover crop shoot biomass application in a 5-year old peach orchard. The region's climate is warm temperate (Cfb), and the soil is classified as a Typic Hapludalf. Litter bags with unlabeled shoot residues were also deposited in the orchard to assess biomass, carbon (C), N, lignin, cellulose, and non-structural biomass decomposition dynamics.

*Results:* After 13 months, the leaves, trunk, and roots showed the greatest proportion of N derived from residues (Ndfr) (35.4, 25.1, and 22.4%, respectively) while the greatest concentrations of  $^{15}\text{N}$  and Ndfr occurred in roots <2 mm (0.0376 and 0.94%, respectively). The N derived from cover crop shoots in the second production cycle was similar among tree organs. Ryegrass residues presented the highest decomposition constant (k) values for dry matter, total organic carbon (TOC), cellulose, and lignin. Hence, black oat residues presented a higher half-life ( $t^{1/2}$ ) for dry matter, TOC, total N, cellulose, and lignin.

*Conclusions:* The N derived from black oat and ryegrass residues in mature trees was expressively low (<1%) and similar between species. Within organs, the highest Ndfr occurred in peach leaves during the flowering stage, when the greatest residue decomposition rate also occurred. Soil N and plant internal N reserves are the major N sources for newly formed organs, but greater contributions to tree N nutrition may occur with long-term cover crop residue deposition and different plant species.

**Keywords:** *Avena strigosa*, *Lolium multiflorum*, N cycling;  $^{15}\text{N}$  recovery, *Prunus persica* (L.) Batsch.

## 1 Introduction

Nutrient management in orchards has a great impact on plant growth, yields, and fruit quality. Sustainable soil fertility management in orchards requires not only a fine-tuning of fertilizer rates, but also a higher use efficiency of nutrients already present in soil (Tagliavini 2012). There is a multitude of factors, including the cycling of plant residues, that dictate nutrient availability in the soil system. A key step to improving nutrient use efficiency is understanding the nutrient's fate during leaf-litter decomposition on the soil surface and the ability of that nutrient to become available for plant uptake (Tagliavini and Scandellari 2013).

Nitrogen (N) is a nutrient with a great impact on growth and reproductive development in fruit trees, such as peach [*Prunus persica* (L.) Batsch.], as it directly affects the flowering quality, fruit set, fruit quality, and yield potential. However, mature peach trees tend to absorb only small amounts of the N derived from fertilization (Policarpo et al. 2002) as the majority of the N used by the new vegetative organs comes from internal N reserves. Part of the N-fertilizer applied to the orchard soils can be lost by volatilization (especially when using urea as N source) (Roccuzzo et al. 2017), leaching (greater loss potential in sandy soils with low organic matter content) (Lorensini et al. 2012), and surface runoff (especially in areas with steep grades) (Martínez et al. 2006). Moreover, greater losses can occur when the N-fertilizers are not applied at the right time and rate to meet trees' nutritional demand.

Cover crops, such as black oat (*Avena strigosa* Schreb.) and ryegrass (*Lolium multiflorum* L.), are used in the inter-row space of orchards to dissipate the kinetic energy of rainfall and reduce soil erosion. Throughout tree development, cover crop shoots are frequently cut and deposited on the soil surface contributing to increasing soil organic matter (SOM)

content and nutrient cycling (Brunetto et al. 2017; Ferreira et al. 2014).

During the decomposition of cover crop residues, part of the carbon (C) present in the plant tissue may remain in the soil, especially from residues with lower lability. However, the majority of the C will probably return to the atmosphere as CO<sub>2</sub>, depending on the residue composition and mineralization rate (Oliveira BS et al. 2016; Reichert et al. 2015). Part of the plant tissue N can increase soil available N (nitrate or ammonium), which can be absorbed by mature trees. The decomposition of plant material and the release of N from cover crop residues depends on the residue biochemical composition, especially the cellulose and lignin content, but also the C/N, lignin/N, and cellulose/lignin ratios (Carranca et al. 2009). Residues with low cellulose content, high lignin content, and high C/N ratio typically have a low decomposition rate and may even temporarily immobilize soil N (Bonanomi et al. 2013). On the other hand, more labile residues with higher cellulose content, lower lignin content, and lower C/N and cellulose/lignin ratios can mineralize N to the soil, increasing the plant-available N forms (Cabrera et al. 2005). The residue decomposition and nutrient release is also dependent on edaphoclimatic characteristics, especially soil texture, moisture, aeration, temperature, and nutrient availability, which directly affect soil microbial activity (Brunetto et al. 2011; Cabrera et al. 2005; Ferreira et al. 2014).

Studies investigating the N derived from cover crop residues can be accurately performed using the <sup>15</sup>N isotope dilution technique (Brunetto et al. 2014; Neto et al. 2008; Tagliavini et al. 2007). Annual tree organs such as leaves, twigs of the year, and fruits are major sinks for the absorbed N derived from in-season fertilization (Brunetto et al. 2017), SOM mineralization, and decomposing cover crop residues. Part of the N stored in perennial organs is remobilized to the tree meristematic tissues in the subsequent growth cycle, reducing the tree N needs from N fertilizers at this vegetative phase. Even though the contribution of cover crops for plant nutrition has been evidenced for different crops and production systems (Amossé et

al. 2014; Brunetto et al. 2011, 2014; Oliveira RA et al. 2016; Ovalle et al. 2010), information on how the N derived from cover crop residues is redistributed in different peach organs is still scarce.

We hypothesize that a) ryegrass and black oat residues will present similar decomposition dynamics, and b) the N released will be used by trees and preferentially redistributed to newly formed organs and trunk. Thus, this study aims to evaluate a) the recovery of N derived from the decomposition of black oat and ryegrass shoot residues in mature trees, b) the N absorption dynamics and the N compartmentalization in different tree organs, and c) to assess black oat and ryegrass shoot residue decomposition dynamics in a mature peach orchard.

## **2 Materials and methods**

### **2.1 Cover crop cultivation**

Soil for this experiment was collected from the 0-0.20 m depth of a soil classified as a Rhodic Paleudalf (Soil Survey Staff 2014) in Southern Brazil (29°42'54" S, 53°42'25" W). The soil was air-dried, ground to pass through a sieve with 2-mm openings, and reserved for analysis. Selected soil chemical and physical characteristics are shown in Table 1. Polypropylene pots (20.0 cm x 25.0 cm) with four kilograms of soil were used to cover crop cultivation.

Seeds of black oat and ryegrass were pre-germinated in plastic germination boxes (Gerbox®) by placing the seeds on paper towels and moistening them with distilled water amounting to 2.5 times the weight of the dry paper. The seed boxes were placed in a BOD (Biochemical Oxygen Demand) chamber at a constant temperature of 25°C and a positive photoperiod of 8 h. A germination test was carried out from 01 May to 13 May 2014, when 25 germinated seeds of each plant species were sown in the polypropylene pots. Seven days after

emergence, the less developed seedlings were thinned, leaving 20 black oat or 20 ryegrass plants in each pot.

Urea containing 46.6% N, labeled with 5 atom%  $^{15}\text{N}$ , was applied to 15 pots containing black oat and 15 pots containing ryegrass at a rate of 10 g of N  $\text{m}^{-2}$ . Unlabeled urea with 45% N was applied to an additional 15 pots grown with black oat and 15 pots grown with ryegrass (control plots). Urea, with and without  $^{15}\text{N}$ , was diluted in deionized water and applied to the soil, thereafter pots were irrigated daily with deionized water to maintain the soil at 70% field capacity (Casaroli and Jong van Lier 2008) during the experimental period. The N-fertilizer solution application was split six times throughout the cultivation period: 10, 17, 24, 31, 38, and 45 days after thinning. At 18 and 41 days after thinning, 50 mL of a nutrient solution containing 500, 502, 48, 19, 9, and 14000 mg  $\text{L}^{-1}$  of B, Mn, Zn, Cu, Mo, and  $\text{KH}_2\text{PO}_4$ , respectively, were applied to each pot. Throughout the crop cycle, weeds were removed by hand from the pots.

Sixty-three days after sowing (*i.e.*, 17 July 2014), the shoots of black oat and ryegrass grown with and without  $^{15}\text{N}$  applications were cut close to the soil surface. The cover crop shoot residues were washed and oven-dried at 65 °C until constant weight. The cover crop  $^{15}\text{N}$ -labeled shoot residues were reserved for Experiment 1, while the unlabeled shoot residues were reserved for Experiment 2. Subsamples of both  $^{15}\text{N}$ -labeled and unlabeled residues were ground with a Wiley mill and separated for chemical characterization and  $^{15}\text{N}$  analysis.

## **2.2 Experiment 1 - Mature trees N recovery from black oat and ryegrass shoot residue decomposition**

### **2.2.1 Experimental layout**

The experiment was carried out in a mature peach orchard of ‘Chimarrita’ [*Prunus persica* (L.) Batsch] (Raseira et al. 2014) grafted on ‘Capdeboscq’ [*Prunus persica* (L.) Batsch]

rootstock (Mayer et al. 2014). The orchard was established in 2009 in the experimental area of Embrapa Uva e Vinho, located in the city of Bento Gonçalves, Southern Brazil (29°09'44" S, 51°31'50" W). Before the installation of the orchard, 30 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (as triple superphosphate) and 40 kg K<sub>2</sub>O ha<sup>-1</sup> (as potassium chloride) were applied to increase soil-test P and K to optimum levels. Subsequently, the orchard received annual fertilizer rates of 90 kg N ha<sup>-1</sup> (as urea), 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (as triple superphosphate), and 80 kg K<sub>2</sub>O ha<sup>-1</sup> (as potassium chloride) to maintain adequate nutrient levels according to the regional production guidelines (CQFS - RS/SC, 2004). After the installation of the experimental area, the adjacent plants received no fertilization. The spacing between tree rows and between plants within rows was 4.0 and 1.5 m, respectively, totaling 1667 plants per hectare. The Y-shaped pruning system was used for trees. The climate of the region is warm temperate (Cfb), according to the Köppen-Geiger classification system, with an average air temperature of 17.1°C and an average annual rainfall of 1755 mm (Fig. 1). The soil is classified as a Typic Hapludalf (Soil Survey Staff 2014) and its selected physical and chemical characteristics are presented in Table 1. The local landscape is gently sloping.

Before the experiment set up, the orchard included cover crops, mainly grasses (*e.g.*, black oat and ryegrass). In the evaluation area, however, all vegetation, including roots, was removed, so that no plants were competing with the trees. On 17 July 2014, <sup>15</sup>N-labeled shoot residues of black oat and ryegrass (sampled 63 days after sowing) were deposited at the soil surface to an area of 0.96 m<sup>2</sup> (0.8 m x 1.2 m) surrounding the tree trunk. Each plot was composed of three trees that each received 0.13 kg dry matter of either black oat or ryegrass shoot residue, corresponding to 1354 kg of dry matter per hectare. A nylon net (2 mm mesh) covered the residue and was fixed to the soil with metal clamps to prevent residue displacement by the wind, rain, and animals. For the duration of the experiment, non-residual herbicide was applied to the area of residue deposition to manage weeds and avoid <sup>15</sup>N absorption by weeds.



### 2.2.2 $^{15}\text{N}$ recovery by mature trees

To follow the pattern of  $^{15}\text{N}$  uptake during the mature tree development, the first peach leaf sampling was carried out on 17 August 2014 (30 days after cover crop shoot residue deposition), while subsequent leaf sampling was performed monthly from September 2014 to April 2015. At each leaf sampling time, twenty mature leaves per plant were collected from the middle position on twigs of the year, surrounding the tree canopy. The leaves were oven-dried with forced air at 65 °C until constant weight, ground, and reserved for total N and  $^{15}\text{N}$  analysis.

On 17 December 2014, 115 days after cover crop shoot residue deposition, a PVC pipe (250 mm diameter) was inserted into the soil to collect a soil core sample from the 0-0.20 m depth of each plot. Three soil cores were collected in the central position at one side of the trees row, where  $^{15}\text{N}$ -labeled black oat and ryegrass shoot residues were deposited. Each soil core was separated into 0.05 m layers (*i.e.*, 0-0.05, 0.05-0.10, 0.10-0.15, and 0.15-0.20 m depths) and combined for a composite sample of each depth. Thereafter, the soil samples were air-dried, ground to pass a 2-mm sieve, and reserved for total N and  $^{15}\text{N}$  analysis.

On 23 November 2015 (15 months after cover crop shoot residue deposition), the trunks of trees that received cover crop residue were cut close to the soil surface and separated into the stem, twigs of the year, 2-year-old branches, and leaves. The fruits were collected from each tree and weighed to determine fruit yield. Thereafter, ten fruits per tree were randomly selected to determine total N content and  $^{15}\text{N}$  enrichment in the pulp (*i.e.*, mesocarp plus exocarp). The tree roots were removed from the soil, washed with distilled water, and separated into three diameter classes (< 2 mm, 2 - 5 mm, > 5 mm). Root material was reserved for analysis of total N and  $^{15}\text{N}$  enrichment. The same stratified soil sampling procedure and analysis, described above, was repeated before peach roots were removed from the soil.

## 2.3 Experiment 2 - Black oat and ryegrass shoot residue decomposition and N release in

## **a mature peach orchard**

### **2.3.1 Experimental layout**

Experiment 2 was laid out in the same orchard used for Experiment 1. Unlabeled black oat and ryegrass shoot residues were placed in nylon litter bags (2 mm mesh) covering an area of 0.16 m<sup>2</sup> (0.4 m x 0.4 m) and deposited on 22 July 2014 on the soil surface of planting rows. Each plot was composed of four trees that received either black oat or ryegrass shoot residue. Each tree received 64 g of dry matter from black oat or ryegrass shoot residue, corresponding to 4,000 kg dry matter per hectare. The bags were fixed to the soil with a metal clamp to increase the area of contact with the soil surface and prevent displacement by the wind. The litter bags were collected at 30-day intervals (*i.e.*, 0, 30, 60, 90, 120, 150, and 180 days after deposition), and at each sampling time, four litter bags of each cover crop were removed from the field. In the laboratory, the residues were washed to remove the adhering soil particles and then oven-dried at 65 °C, ground, sieved through 2-mm openings, and reserved for further analysis.

### **2.4 Plant and soil analysis**

Before cover crop residue deposition in the experiments, a subsample was subjected to sulfuric acid digestion to determine total N, phosphorus (P), potassium (K), calcium (Ca), and magnesium (Mg) contents (Tedesco et al. 1995). Total N was determined by steam distillation (TE-0364, Tecnal, Brazil). Total P was determined in a spectrophotometer (Bell Photonics, 1105, Brazil) at 882 nm (Murphy and Riley 1962). Total K was determined in a flame photometer (Digimed, BM-62, Brazil). Calcium and Mg were determined in an atomic absorption spectrophotometer (PerkinElmer, AAnalyst 200, USA). Total organic carbon (TOC) was determined by wet combustion (Yeomans and Bremner 1988). The determination of lignin, cellulose, and non-structural biomass was performed according to the methodology described by Aber and Martin (1999). Total N and <sup>15</sup>N were analyzed in an elemental analyzer (Thermo

Scientific, Flash EA 1112, Milan, Italy) and by isotope-ratio mass spectrometry (Thermo Scientific, Delta V Advantage, Bremen, Germany), respectively. Cover crop shoot residue chemical characteristics are presented in Table 2.

The tree organs' tissue and soil samples from Experiment 1 were also analyzed for total N and %<sup>15</sup>N following the above-mentioned procedures. The remaining cover crop residues from litter bags in Experiment 2 were also analyzed for C content, lignin, cellulose, and non-structural biomass following the above-mentioned methodologies. The decomposition of residue dry matter and the release of C, N, cellulose, lignin, and non-structural biomass were estimated by subtracting the initial content from the amount determined after each sampling time.

## 2.5 Calculations and statistical analysis

Atom% <sup>15</sup>N excess in soil and plant tissue samples was calculated based on the natural <sup>15</sup>N abundance (Mariotti 1983). The N derived from residue (Ndfr) and the N derived from soil (Ndfs) was calculated by the following equations:

$$^{15}\text{N excess in sample (\%)} = \% ^{15}\text{N in sample} - 0.3663\% \quad (\text{Eq. 1})$$

$$\text{Ndfr (\%)} = (\% ^{15}\text{N excess in sample} / \% ^{15}\text{N excess in residue}) \times 100 \quad (\text{Eq. 2})$$

$$\text{Ndfs(\%)} = 100 - \text{Ndfr} \quad (\text{Eq. 3})$$

The residual percentage of each variable (total C and N, cellulose, lignin, and non-structural biomass) were adjusted by the exponential mathematical model described by Wieder and Lang (1982):

$$X = X_0^{(-kt)} \quad (\text{Eq. 4})$$

Where: X is the amount of dry matter or nutrient remaining in the residue after a period t (days); X<sub>0</sub> is the initial amount of dry matter or nutrient in the residue; k is the decomposition constant.

The half-life ( $t^{1/2}$ ) was calculated with the value of k (Paul and Clark 1996) (Equation 5). The  $t^{1/2}$  expresses the time required for half of the residue to decompose and half of the nutrients contained in the residue to be released.

$$t^{1/2} = 0.693/k \quad (\text{Eq. 5})$$

All data were submitted to normality and homogeneity of variance by the Lilliefors and Shapiro-Wilk tests prior to the analysis of variance (ANOVA). The dry matter, atom% <sup>15</sup>N, total N (mg tree<sup>-1</sup>), and total soil N (Experiment 1) variables were transformed [log<sub>10</sub>(x)] to fit a normal distribution before running the ANOVA. Experiment 1 had a completely randomized block design with five replicates. ANOVA was performed to determine the influence of black oat and ryegrass shoot residues on leaf total N, atom% <sup>15</sup>N excess, and N<sub>dfr</sub> for each leaf's sampling time. ANOVA was also conducted to determine the influence of black oat and ryegrass shoot residue, tree organs, and their interaction on dry matter, total N, atom% <sup>15</sup>N excess, N<sub>dfr</sub>, and N<sub>dfs</sub>. Experiment 2 had a completely randomized block design with five replicates. ANOVA was performed to determine the influence of black oat and ryegrass shoot residues on the remaining dry matter, TOC, total N, cellulose, lignin, and non-structural biomass decomposition constant rate (k) and half-life ( $t^{1/2}$ ). Means were compared by the Tukey-test ( $p < 0.05$ ). When no difference was observed among cover crop species, the mean

value of the treatments was compared. ANOVA was also conducted to determine the influence of black oat and ryegrass shoot residues, residue deposition time, and their interaction on C/N ratio, dry matter, TOC, and total N. When the interaction was significant, two times the standard error of means (SEM) was used as the minimum difference between means statistically different for  $p \leq 0.05$ .

### 3 Results

#### 3.1 Experiment 1 - Mature trees N recovery from black oat and ryegrass shoot residue decomposition

Following the pattern of leaf response of mature trees cultivated in soils with black oat and ryegrass  $^{15}\text{N}$ -labeled shoot residue deposition, the highest N concentration was found in trees cultivated with ryegrass shoot residue, especially in November and December 2014 (Fig. 2a). However, the highest leaf N concentration occurred in August 2014, shortly after the deposition of both cover crop residues on the soil surface. Thereafter, leaf N concentration decreased over time in all trees. The highest atom%  $^{15}\text{N}$  and the Ndf<sub>r</sub> were observed in September in leaves of trees that received the deposition of black oat residue (Fig. 2b and 2c). In subsequent evaluations, there were no differences for leaf concentration of total N, atom %  $^{15}\text{N}$ , and % Ndf<sub>r</sub> between trees cultivated under both cover crop treatments.

After 75 weeks (*i.e.*, December 2015), the dry matter, atom %  $^{15}\text{N}$ , and Ndf<sub>r</sub> of mature peach organs did not differ between the two treatments, but significant differences were observed within tree organs (Table 3). The highest total N concentration occurred in mature tree leaves. The highest dry matter yield was observed in the peach trunk, followed by roots > 5 mm and leaves (7677.6, 2243.1, and 1212.2 g tree<sup>-1</sup>, respectively). Mature trees leaves and trunk presented the highest Ndf<sub>r</sub> (mg tree<sup>-1</sup>) and Ndf<sub>s</sub> (mg tree<sup>-1</sup>) while the highest atom %  $^{15}\text{N}$  excess and % Ndf<sub>r</sub> were observed in roots < 2 mm.

Soil characteristics were not affected by the type of cover crop residue, but differences were observed among the soil layers (Table 4). For the mean effect of crop residues, the soil samples collected at 23 and 75 weeks (*i.e.*, 115 and 375 days, respectively) after cover crop residue deposition showed the highest atom%  $^{15}\text{N}$  and Ndf<sub>r</sub> at the 0-0.05 m depth (Table 4), with little variation among other soil depths. It should be noted that 23 weeks after the deposition of black oat and ryegrass shoot residues, atom%  $^{15}\text{N}$  and Ndf<sub>r</sub> values were higher than the values observed 75 weeks after the residue deposition. Overall, the percentage of soil Ndf<sub>r</sub> below the 0.05-m depth was low.

### **3.2 Experiment 2 - Black oat and ryegrass shoot residue decomposition and N release in a mature peach orchard**

The temporal dynamics of shoot residue dry matter, TOC, N, lignin, cellulose, and non-structural biomass contents were explained by the exponential decay model (Eq. 4). Cover crop residue dry matter and TOC, N, cellulose, lignin, and non-structural biomass contents decreased rapidly during the experimental period, with slight differences between cover crop species (Fig. 3). At the end of the trial (*i.e.*, 150 days after the residue deposition), only approximately 15% of the cover crop residue dry matter and TOC remained in the soil surface (Fig. 3a and 3b, respectively). There was an 82 and 79% decrease in the N content of black oat and ryegrass remaining residues, respectively (Fig. 3c). Lignin and cellulose presented comparable decomposition dynamics, with remaining percentages close to zero after 150 days of residue decomposition. The remaining non-structural biomass followed a similar trend as dry matter, TOC, and total N, with slightly higher values (20 and 30% for black oat and ryegrass shoot residues, respectively) at the end of the experiment (Fig. 3f). The highest  $k$  values for dry matter, TOC, cellulose, and lignin were observed for ryegrass shoot residue (Table 5). Black oat residue presented the highest  $t^{1/2}$  values for dry matter, TOC, total N, cellulose, and lignin

(Table 5). There was no statistical difference for  $k$  and  $t^{1/2}$  values of the non-structural biomass in both cover crop residues. The C/N ratio of litter bag residues over time was greater in black oat than in ryegrass, except at 90 and 180 days after deposition, when the two cover crops presented similar values (Table 6). The C/N ratio constantly decreased until 90 days of deposition, then remained stable for a month and slightly increased at the end of the trial. Black oat shoot residue showed greater dry matter than ryegrass, with values decreasing over time in both cover crops. Total organic C decreased across time and was different between cover crop species. Total organic C of black oat shoot residue was higher than ryegrass from 30 to 120 days after deposition; at the beginning of the experiment, ryegrass showed higher values than black oat, while at the end of the experiment (150 and 180 days after deposition), the values were similar between the two cover crops species. Black oat total N content was higher than ryegrass at 0, 30, 90, and 120 days after deposition; in other sampling times, the situation was inverted. The remaining residue N content decreased over time, the only exception was black oat that showed a slight increase 90 days after deposition and then again decreased until the end of the trial (Table 6).

#### **4 Discussion**

Different from annual crops, perennial plants, such as fruit trees, use two main sources of N for their vegetative growth and reproduction: the root N uptake and the internal N cycling (Carranca et al. 2018). This behavior can be observed in our study by the highest total N concentration in mature tree leaves at the first sampling time, which is probably related to the plant's internal N reserves. Likewise, the highest N derived from cover crop residues measured in leaves at 30 and 60 days after black oats and ryegrass shoot deposition, respectively, indicates that the cover crop residues were rapidly decomposing and releasing N to the soil, which was taken up by trees during an active absorption phase. This timing coincided with the flowering

stage when plants increase young root growth, which is responsible for the absorption of water and nutrients, including soil N (Bravo et al. 2012). The trees' superficial roots, which were in proximity with the cover crops' decomposing residues, may have contributed to a greater extent to the N absorption at this stage. Furthermore, this behavior indicates that synchronizing cover crop termination (*e.g.*, mowing) so that a greater decomposition rate occurs in a period of high tree N demand can contribute to enhancing N recovery.

A higher soil N availability favors plant dry matter production by increasing the formation and growth of new vegetative organs such as leaves and twigs, which have a high nutritional N demand due to intense cell division and elongation (Nario et al. 2003; Roccuzzo et al. 2017). However, the highest leaf N content in mature trees at different phenological stages was mostly derived from other N forms than cover crop shoot residue decomposition. This is because the mineral N fertilization that the orchard received over the years may have had a residual effect on our study. The low N recovery from the cover crop shoot residue decomposition by the peach organs might also be related to N losses from the plant-soil system, especially by denitrification, runoff, and leaching (Carranca et al. 2018; Roccuzzo et al. 2017) as only a small proportion of N<sub>dfr</sub> (below 0.8%) remained in the soil. On the other hand, the high N content in newly formed organs may also be a response to internal N remobilization.

The similar behavior observed for the black oat and ryegrass shoot residue decomposition dynamics, dry matter, organic C, N, cellulose, lignin, and non-structural biomass contents is related to the residue initial lignin concentration and C/N ratio (Table 2). The decrease of dry matter, organic C, N, cellulose, lignin, and non-structural biomass that occurred over time is mediated by the activity of soil fauna and the degradation by the microbial population (Carranca et al. 2009; Nguyen and Marschner 2017; Oliveira RA et al. 2016). During the decomposition period, cover crop residues had a C/N ratio below 20 (Table 6) at all sampling times, which facilitates the colonization and mineralization by microbial population



(Ferreira et al. 2014; Oliveira RA et al. 2016). Consequently, the cover crop shoot residue dry matter reduction over the 150 days of the trial contributed to the decrease of soil cover (this if there is no deposition of residues in short intervals of time) and the potential to dissipate the kinetic energy of the raindrops, leaving the soil more susceptible to wind and water erosion (Ferreira et al. 2014; Oliveira RA et al. 2016), and therefore N loss by runoff. This process was probably even more intense in the surrounding area of the tree, without cover crop residues. Frequent rainfall and warm temperatures during spring and summer are known to increase soil microbial activity and therefore intensify residue decomposition rate (Chen et al. 2020). Hence, continuous cover crop residue deposition in orchards is paramount to increase both soil cover and SOM content, while reducing nutrient losses over time, which ultimately will positively affect N use efficiency.

Using cover crops in orchards is economically feasible since, in addition to the protection against soil erosion, the residue deposition can contribute to the cycling of different nutrients, increasing soil health and potentially reducing the costs associated with fertilizer application. On average, during the experimental period, black oat and ryegrass residues released an equivalent of 120 kg N ha<sup>-1</sup> and 136 kg N ha<sup>-1</sup>, corresponding to 261 and 296 kg urea ha<sup>-1</sup>, respectively. However, this is a substantial amount of N released in 150 days for the present climatic conditions, and therefore a major part of the N derived from cover crop residue was probably lost from the soil-plant system, which is supported by the low N recovery by mature trees. The greater N recovery by mature trees during the first 60 days was related to the 50% N released from both cover crops during this period (see Table 6 and Fig. 3), which corresponds to the flowering phase where new roots are active and are responsible for water and nutrient uptake (Bravo et al. 2012). Furthermore, during this phase the trees present an intensive leaf growth, which increases leaf area and consequently surface area for transpiration, resulting in a greater accumulation of minerals within the plant body. These findings agree with

those observed in other fruit studies, such as the grapevines, in soils amended with other types of residue (Brunetto et al. 2011, 2014).

The overall contribution of cover crop residue to peach nutrition in our study was low (below 1%). However, a greater contribution might occur with different cover crop species and residue management that synchronizes the residue N release with trees nutrient demand. It is also important to highlight that even with a low contribution to N nutrition in a short term, cover crops play an important role in cycling other nutrients such as P, K, and S, increasing SOM content and reducing soil and water loss in the system, which is paramount for sustainable food production.

## **5 Conclusions**

The highest percentage of N derived from the decomposition of black oat and ryegrass shoot residues occurred in leaves at the flowering stage, *i.e.*, about 30 to 60 days after the shoot residues deposition on the soil surface. The majority of peach leaf N and the N in other young tree organs was derived from other N sources than the cover crop residues, such as soil available N, organic matter, plant internal N reserves, or even the residual effect of mineral fertilizers applied to the trees adjacent to the study area. This is partially explained by the shoot residue decomposition rate on the soil surface, which was reduced to 25-50% during the flowering period.

Most of the N contained in leaves in the first evaluation cycle and annual and perennial organs in the second evaluation cycle was also derived from other N sources than cover crop decomposing shoot residue. It is expected that cover crop grasses with a similar C/N ratio to black oat and ryegrass will have comparable residue decomposition dynamics and contribution to tree N nutrition.

Further studies need to be performed to evaluate other cover crop species and residue

management to synchronize cover crop N mineralization with the stages of greater plant N demand to avoid N losses and increase the N recovery and plant nutritional status.

**Conflict of Interest** The authors declare that they have no conflict of interest.

## 6 References

Aber JD, Martin M (1999) Leaf Chemistry, 1992-1993 (ACCP). [Leaf Chemistry, 1992-1993 (Accelerated Canopy Chemistry Program)].

Amossé C, Jeuffroy MH, Mary B, David C (2014) Contribution of relay intercropping with legume cover crops on nitrogen dynamics in organic grain systems. *Nutr. Cycl. Agroecosyst.* 98:1–14.

Bonanomi G, Incerti G, Giannino F, Mingo A, Lanzotti V, Mazzoleni S (2013) Litter quality assessed by solid state<sup>13</sup>C NMR spectroscopy predicts decay rate better than C/N and Lignin/N ratios. *Soil Biol Biochem.* 56:40-48.

Bravo K, Moreno T, Baldi E, Marcolini G, Sorrenti G, Quartieri M, Marangoni B (2012) Effect of organic fertilization on carbon assimilation and partitioning in bearing nectarine trees. *Sci. Horticult.* 137:100-106.

Brunetto G, Ventura M, Scandellari F, Ceretta CA, Kaminski J, Melo GWB, Tagliavini M (2011) Nutrients release during the decomposition of mowed perennial ryegrass and white clover and its contribution to nitrogen nutrition of grapevine. *Nutr. Cycl. Agroecosyst.* 90:299-308.

Brunetto G, Ceretta CA, Melo GWB, Kaminski J, Trentin G, Girotto E, Ferreira PAA, Miotto A, Trivelin PCO (2014) Contribution of nitrogen from agricultural residues of rye to ‘*Niagara Rosada*’ grape nutrition. *Sci. Horticult.* 169:66-70.

Brunetto G, Lorensini F, Ceretta CA, Ferreira PAA, Couto RR, De Conti L, Ciotta MN,

427 Kulmann M, Schneider RO, Somavilla LM, et al. (2017) Contribution of mineral N to  
 428 young grapevine in the presence or absence of cover crops. *J. Soil Sci. Plant Nutri.*  
 429 17(3):570-580.

430 Cabrera ML, Kissel DE, Vigil MF (2005) Nitrogen mineralization from organic residues:  
 431 research opportunities. *J. Environ. Qual.* 34:75-79.

432 Carranca C, Rocha I, Varennes A, Oliveira A, Pampulha ME, Torres MO (2009) Effect of  
 433 tillage and temperature on nitrogen mineralization and microbial activity and microbial  
 434 numbers of lupine amended soil. *Agrochimica.* 53(3):183-195.

435 Carranca C, Brunetto G, Tagliavini M (2018) Nitrogen nutrition of fruit trees to reconcile  
 436 productivity and environmental concerns. *Plants.* 7(4):1-12.

437 Casaroli D, Jong van Lier Q (2008). Critérios para determinação da capacidade de vaso.  
 438 [Criteria for pot capacity determination]. *Rev. Bras. Ciênc. Solo*, 32(1):59-66. Portuguese.

439 Chen Q, Niu B, Hu Y, Luo T, Zhang G (2020) Warming and increased precipitation indirectly  
 440 affect the composition and turnover of labile-fraction soil organic matter by directly  
 441 affecting vegetation and microorganisms. *Sci. Total Environ.* 714 (136787): 1-9.

442 Comissão de Química e Fertilidade do Solo (2004) Manual de calagem e adubação para os  
 443 Estados de Rio Grande do Sul e de Santa Catarina. [Liming and fertilization manual for the  
 444 states of Rio Grande do Sul and Santa Catarina]. Porto Alegre, Sociedade Brasileira de  
 445 Ciência do Solo. 400p. Portuguese.

446 Ferreira PAA, Girotto E, Trentin G, Miotto A, Melo GWB, Ceretta CA, Kaminski J, Del Frari  
 447 BK, Marchezan C, Silva LOS, et al. (2014) Biomass decomposition and nutrient release  
 448 from black oat and hairy vetch residues deposited in a vineyard. *Rev. Bras. Ciênc. Solo.*  
 449 38:1621-1632.

450 Lorensini F, Ceretta CA, Girotto E, Cerini JB, Lourenzi CR, De Conti L, Trindade MM, Melo  
 451 GWB, Brunetto G (2012) Lixiviação e volatilização de nitrogênio em um Argissolo

452 cultivado com videira submetida à adubação nitrogenada. [Lixiviation and volatilization of  
 453 nitrogen in Sandy Typic Hapludalf soil cultivated with grapevine submitted to the nitrogen  
 454 fertilization]. Ciênc. Rural. 42:1173-1179. Portuguese.

455 Martínez JRF, Zuazo VHD, Raya AM (2006) Environmental impact from mountainous olive  
 456 orchards under different soil-management systems (SE Spain). Sci. Total Environ. 358:  
 457 46– 60.

458 Mariotti A (1983) Atmospheric nitrogen is a reliable standard for natural  $^{15}\text{N}$  abundance  
 459 measurements. Nature, 303:685-7.

460 Mayer NA, Bianchi VJ, Castro LAS (2014) Porta-enxertos. [Rootstocks]. In: Raseira MCB,  
 461 Pereira JFM, Carvalho FLC, editors. Pessegueiro [Peach tree]. Brasília: Embrapa; p. 173-  
 462 223. Portuguese.

463 Murphy J, Riley JP (1962) A modified single solution method for the determination of  
 464 phosphate in natural waters. Anal. Chim. Acta. 27:31-36.

465 Nario A, Pino I, Zapata F, Albornoz MP, Baherl P (2003) Nitrogen ( $^{15}\text{N}$ ) fertiliser use efficiency  
 466 in peach (*Prunus persica* L.) cv. Goldencrest trees in Chile. Sci. Hortic. 97:279–287.

467 Neto C, Carranca C, Clemente J, De Varennes A (2008) Nitrogen distribution, remobilization  
 468 and re-cycling in young orchard of non-bearing ‘Rocha’ pear trees. Sci. Hortic. 118:299-  
 469 307.

470 Nguyen TT, Marschner P (2017) Soil respiration, microbial biomass and nutrient availability  
 471 in soil after addition of residues with adjusted N and P concentrations. Pedosphere, 27:76-  
 472 85.

473 Oliveira BS, Ambrosini VG, Trapp T, Santos MA, Sete PB, Lovato PE, Loss A, Comin JJ,  
 474 Lourenzi CR, Rosa RC, et al. (2016) Nutrition, productivity and soil chemical properties  
 475 in an apple orchard under weed management. Nutr. Cycl. Agroecosys. 104:247-258.

476 Oliveira RA, Brunetto G, Loss A, Gatiboni LC, Kürtz C, Müller Júnior V, Lovato PE, Oliveira

477 BS, Souza M, Comin JJ (2016) Cover crops effects on soil chemical properties and onion  
 478 yield. Rev. Bras. Ciênc. Solo. 40:1-17.

479 Ovalle C, del Pozo A, Peoples MB, Lavín A (2010) Estimating the contribution of nitrogen  
 480 from legume cover crops to the nitrogen nutrition of grapevines using a  $^{15}\text{N}$  dilution  
 481 technique. Plant Soil. 334:247–259.

482 Paul EA, Clark FE (1996) Dynamics of residue decomposition and soil organic matter turnover.  
 483 In: Paul EA, Clark FE, editors. Soil microbiology and biochemistry. 2nd ed. San Diego:  
 484 Academic, (1996). pp. 158-79.

485 Policarpo M, Di Marco L, Caruso T, Gioacchini P, Tagliavini M (2002) Dynamics of nitrogen  
 486 uptake and partitioning in early and late fruit ripening peach (*Prunus persica*) tree  
 487 genotypes under a Mediterranean climate. Plant Soil. 239:207–214.

488 Raseira MCB, Nakasu BH, Barbosa W (2014) Cultivares: descrição e recomendação.  
 489 [Cultivars: description and recommendation] In: Raseira MCB, Pereira JFM, Carvalho  
 490 FLC, editors. Pessequeiro [Peach tree]. Brasília: Embrapa; p. 73-141. Portuguese.

491 Reichert JM, Rodrigues MF, Bervald CM, Brunetto G, Kato OR, Schumacher MV (2015)  
 492 Fragmentation, fiber separation, decomposition, and nutrient release of secondary-forest  
 493 biomass, mechanically chopped-and-mulched, and cassava production in the Amazon.  
 494 Agric. Ecosyst. Environ. 204:8-16.

495 Roberts TL, Norman RJ, Slaton NA, Wilson CE Jr, Ross WJ, Bushong JT. 2009a. Direct steam  
 496 distillation as an alternative to the Illinois soil nitrogen test. Soil Sci Soc Am J. 73:1268–  
 497 1275.

498 Roccuzzo G, Scandellari F, Allegra M, Torrisi B, Stagno F, Mimmo T, Zanutelli D, Gioacchini  
 499 P, Millard P, Tagliavini M (2017) Seasonal dynamics of root uptake and spring  
 500 remobilisation of nitrogen in field grown orange trees. Sci. Hortic. 226:223–230.

501 Soil Survey Staff (2014) Soil taxonomy: a basic system of soil classification for making and

502 interpreting soil surveys. 12th ed. Washington, DC: United States Department of  
 503 Agriculture, Natural Resources Conservation Service.

504 Tagliavini M, Tonon G, Scandellari F, Quinônes A, Palmieri S, Menarbin G, Gioacchini P,  
 505 Masia M (2007) Nutrient recycling during the decomposition of apple leaves (*Malus*  
 506 *domestica*) and mowed grasses in orchard. *Agric. Ecosyst. Environ.* 118:191-200.

507 Tagliavini M (2012) Nutrient fluxes in orchard ecosystems. In: 3rd Slovenian Fruit Congress  
 508 with international participation, Krško, 21-23. November 2012 / [organizers] Professional  
 509 Fruit Society of Slovenia [in] University of Ljubljana, Biotechnical Faculty, Department  
 510 of Agronomy, Department of Fruit Growing, Viticulture and horticulture; [editor of the  
 511 collection Metka Hudina]. - Ljubljana: Professional Fruit Society of Slovenia, 2012.

512 Tagliavini M, Scandellari F (2013) Methodologies and Concepts in the Study of Nutrient  
 513 Uptake Requirements and Partitioning in Fruit Trees. *Acta Horticulturae*, (984), 47–56 \_  
 514 10.17660\_ActaHortic.2013.984.3.

515 Tedesco MJ, Gianello C, Bissani CA, Bohnen H, Volkweiss SJ (1995) Análise de solo, plantas  
 516 e outros materiais. [Analysis of soil, plants and other materials]. Porto Alegre:  
 517 Departamento de Solos/UFRGS. Portuguese.

518 Walkley A, Black IA (1934) An examination of the Degtjareff method for determining soil  
 519 organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci*  
 520 37(1):29–38.

521 Wieder RK, Lang GE (1982) A critique of the analytical methods used examining de-  
 522 composition data obtained from litter bags. *Ecology*. 63(6):1642-1663.

523 Yeomans JC, Bremmer JM (1988) A rapid and precise method for routine determination of  
 524 organic carbon in soil. *Comm. Soil Sci. Plant Anal.* 19:1467-1476.

525 **Table 1** Selected physical and chemical characteristics of experimental soils at the 0-0.20 m  
526 depth

Soil properties	Unit	Experiment1 and 2
		0-0.20 m layer
Clay (pipette method)	g kg <sup>-1</sup>	333
Silt (pipette method)	g kg <sup>-1</sup>	405
Sand (pipette method)	g kg <sup>-1</sup>	262
Organic C (Walkley and Black 1934)	g kg <sup>-1</sup>	17.3
pH in H <sub>2</sub> O (1:1)	-	5.2
Total N (Kjeldahl method)	g kg <sup>-1</sup>	2.0
NO <sub>3</sub> <sup>-</sup> -N (extracted by KCl 1 mol L <sup>-1</sup> )	mg kg <sup>-1</sup>	34
NH <sub>4</sub> <sup>+</sup> -N (extracted by KCl 1 mol L <sup>-1</sup> )	mg kg <sup>-1</sup>	56
Alkaline hydrolizable N (Roberts et al. 2009)	mg kg <sup>-1</sup>	183
Aluminum (exchangeable) (extracted by KCl 1 mol L <sup>-1</sup> )	mg kg <sup>-1</sup>	15
Magnesium (exchangeable) (extracted by KCl 1 mol L <sup>-1</sup> )	mg kg <sup>-1</sup>	150
Calcium (exchangeable) (extracted by KCl 1 mol L <sup>-1</sup> )	mg kg <sup>-1</sup>	760
Phosphorus (available) (extracted by Mehlich 1)	mg kg <sup>-1</sup>	15
Potassium (available) (extracted by Mehlich 1)	mg kg <sup>-1</sup>	100

527



**Table 2** Chemical characterization of black oat and ryegrass shoot residues at the beginning of the experiment, and the amount of dry matter and nutrients added to the soil for residues decomposition (0-20 cm depth).

Variable	Black oat	Ryegrass
----- g kg <sup>-1</sup> -----		
TOC <sup>a</sup>	463.2±3.5 <sup>k</sup>	431.9±2.0
Total N <sup>b</sup>	42.9±0.7	47.1±0.6
P <sup>c</sup>	4.3±0.2	4.0±0.2
K <sup>d</sup>	24.6±0.4	27.9±1.7
Ca <sup>e</sup>	4.9±0.2	5.4±0.1
Mg <sup>f</sup>	5.8±0.1	5.6±0.1
Cel <sup>g</sup>	388.9±0.6	444.7±1.2
Lig <sup>h</sup>	141.4±0.3	140.5±1.8
Bio <sup>i</sup>	569.7±0.3	414.8±0.8
C/N	10.8±0.2	9.2±0.1
Lig/N	1.0±0.01	3.0±0.04
C/P	109.0±5.9	108.3±5.7
Cel/Lig	9.4±0.8	3.2±0.6
At % N <sup>15</sup>	40.1±0.1	36.0±0.04
Amount of residue and nutrients added to the soil surface (kg ha <sup>-1</sup> )		
DM <sup>j</sup>	5176.8	5607.4
TOC	220.1	228.9
N	13.6	14.9
P	2.4	2.2
K	20.1	22.0
Ca	2.0	2.5
Mg	2.1	2.5

<sup>a</sup>Total organic carbon; <sup>b</sup>Total nitrogen; <sup>c</sup>Total phosphorus; <sup>d</sup>Total potassium; <sup>e</sup>Total calcium; <sup>f</sup>Total magnesium; <sup>g</sup>Cellulose; <sup>h</sup>Lignin; <sup>i</sup>Non-structural biomass; <sup>j</sup>Dry matter; <sup>k</sup>mean standard error (n = 3)

**Table 3** Dry matter, total N, atom  $^{15}\text{N}$  excess,  $^{15}\text{N}$  derived from residues (Ndfr), and N derived from other sources (Ndfs) in mature peach trees organs after 375 days of black oat and ryegrass shoot residues deposition on the soil surface (Experiment 1)

Cover crop	Peach tree organs								CV (%)
	Pulp	Leaves	Twigs of the year	Branches of the year	Trunk	Roots > 5 mm	Roots 2 - 5 mm	Roots < 2 mm	
Dry matter (g tree <sup>-1</sup> )									
Black Oat	54.43	1243.18	523.14	1163.23	6387.33	1891.52	137.58	145.95	6.12 <sup>(2)</sup>
Ryegrass	55.33	1181.28	556.66	985.35	8967.83	2594.65	111.17	98.33	
Average	54.88 <sup>f(1)</sup>	1212.23 <sup>bc</sup>	539.90 <sup>d</sup>	1074.29 <sup>c</sup>	7677.58 <sup>a</sup>	2243.08 <sup>b</sup>	124.37 <sup>e</sup>	122.14 <sup>e</sup>	
Total N (g kg <sup>-1</sup> )									
Black Oat	11.16	26.56	7.44	4.07	2.82	5.90	8.04	8.65	7.36
Ryegrass	12.18	27.87	7.69	4.47	3.12	8.83	11.22	10.54	
Average	11.67 <sup>b</sup>	27.22 <sup>a</sup>	7.60 <sup>d</sup>	4.27 <sup>e</sup>	2.98 <sup>f</sup>	7.37 <sup>d</sup>	9.63 <sup>c</sup>	9.60 <sup>c</sup>	
Total N (mg tree <sup>-1</sup> )									
Black Oat	575.39	33012.08	3839.95	4733.8	18244.51	11110.02	1109.17	1257.05	5.5
Ryegrass	671.81	32957.52	4288.27	4420.87	28052.09	22870.01	1228.49	1027.03	
Average	623.60 <sup>e</sup>	32984.80 <sup>a</sup>	4064.11 <sup>c</sup>	4577.33 <sup>c</sup>	23148.30 <sup>ab</sup>	16990.01 <sup>b</sup>	1168.83 <sup>d</sup>	1142.04 <sup>d</sup>	
atom <sup>15</sup> N excess (%)									
Black Oat	0.019	0.0138	0.0164	0.0211	0.0156	0.0216	0.0279	0.0387	0.0376 <sup>a</sup>
Ryegrass	0.017	0.0170	0.0154	0.0190	0.0151	0.0172	0.0266	0.0364	
Average	0.0180 <sup>c</sup>	0.0154 <sup>c</sup>	0.0159 <sup>c</sup>	0.0200 <sup>bc</sup>	0.0153 <sup>c</sup>	0.0194 <sup>bc</sup>	0.0272 <sup>b</sup>		
<sup>15</sup> N derived from shoot residues (Ndfr) (% total N in the organ)									
Black Oat	0.47	0.35	0.41	0.53	0.39	0.54	0.7	0.96	0.94 <sup>a</sup>
Ryegrass	0.42	0.42	0.38	0.47	0.37	0.43	0.66	0.91	
Average	0.45 <sup>bc</sup>	0.38 <sup>c</sup>	0.39 <sup>c</sup>	0.50 <sup>bc</sup>	0.38 <sup>c</sup>	0.48 <sup>bc</sup>	0.67 <sup>ab</sup>		
<sup>15</sup> N derived from shoot residues (Ndfr) (mg tree <sup>-1</sup> )									
Black Oat	2.69	110.88	15.99	25.17	73.86	60.21	7.98	12.01	21.1
Ryegrass	2.84	140.2	16.46	21.09	104.39	98.99	8.4	8.92	
Average	2.76 <sup>e</sup>	125.54 <sup>a</sup>	16.22 <sup>bc</sup>	23.13 <sup>b</sup>	89.12 <sup>a</sup>	79.60 <sup>a</sup>	8.19 <sup>cd</sup>	10.46 <sup>bc</sup>	
<sup>15</sup> N derived from other sources (Ndfs) (% total N in the organ)									
Black Oat	99.53	99.65	99.59	99.47	99.61	99.46	99.3	99.04	99.06 <sup>c</sup>
Ryegrass	99.58	99.58	99.62	99.53	99.63	99.57	99.34	99.09	
Average	99.56 <sup>ab</sup>	99.62 <sup>a</sup>	99.60 <sup>a</sup>	99.50 <sup>ab</sup>	99.62 <sup>a</sup>	99.52 <sup>ab</sup>	99.32 <sup>b</sup>		
<sup>15</sup> N derived from other sources (Ndfs) (mg tree <sup>-1</sup> )									
Black Oat	572.71	32901.20	3823.96	4708.63	18170.65	11049.81	1101.19	1245.04	5.49
Ryegrass	668.97	32817.32	4271.81	4399.78	27947.71	22771.01	1220.09	1018.12	
Average	620.84 <sup>e</sup>	32859.26 <sup>a</sup>	4047.89 <sup>c</sup>	4554.20 <sup>c</sup>	23059.18 <sup>ab</sup>	16910.41 <sup>b</sup>	1160.64 <sup>d</sup>	1131.58 <sup>d</sup>	

<sup>(1)</sup>Means (n = 5) followed by the same lowercase letter in the column, for each plant organ, do not differ by the Tukey-test ( $p > 0.05$ ).<sup>(2)</sup>Coefficient of variation (CV) of cover crop shoot residue error 1

**Table 4** Soil total nitrogen (TN) concentration, atom  $^{15}\text{N}$  excess, and  $^{15}\text{N}$  derived from residues (Ndfr) in the 0-0.20 m depth of a mature peach tree orchard with black oat and ryegrass shoot residues deposition (Experiment 1)

Cover crop	Soil depth (m)				CV (%)
	0-0.05	0.05-0.10	0.10-0.15	0.15-0.20	
23 Weeks (i.e., 115 days)					
	Total N (g kg <sup>-1</sup> )				
Black Oat	4.1	3.3	2.0	1.8	
Ryegrass	4.7	2.6	2.0	1.3	
<i>Average</i>	<i>4.4 a<sup>(1)</sup></i>	<i>2.9 b</i>	<i>2.0 c</i>	<i>1.6 c</i>	<i>6.22<sup>(2)</sup></i>
	<sup>15</sup> N (atom% <sup>15</sup> N excess)				
Black Oat	0.0224	0.0097	0.0056	0.0051	
Ryegrass	0.0406	0.0098	0.0057	0.0050	
<i>Average</i>	<i>0.0215 a</i>	<i>0.0097 b</i>	<i>0.0056 b</i>	<i>0.0050b</i>	<i>0.40</i>
	Ndfr (%)				
Black Oat	0.56	0.24	0.14	0.13	
Ryegrass	1.01	0.24	0.14	0.13	
<i>Average</i>	<i>0.79 a</i>	<i>0.24 b</i>	<i>0.13 b</i>	<i>0.13 b</i>	<i>5.92</i>
75 Weeks (i.e., 375 days)					
	Total N (g kg <sup>-1</sup> )				
Black Oat	4.6	3.4	2.4	1.9	
Ryegrass	4.0	3.0	2.3	1.9	
<i>Average</i>	<i>4.3 a</i>	<i>3.2 b</i>	<i>2.3 c</i>	<i>1.9 c</i>	<i>5.66</i>
	<sup>15</sup> N (atom% <sup>15</sup> N excess)				
Black Oat	0.0074	0.0041	0.0052	0.0257	
Ryegrass	0.0263	0.0058	0.0051	0.0044	
<i>Average</i>	<i>0.0169 a</i>	<i>0.0050 b</i>	<i>0.0051 b</i>	<i>0.0051 b</i>	<i>0.64</i>
	Ndfr (%)				
Black Oat	0.18	0.10	0.13	0.14	
Ryegrass	0.66	0.15	0.13	0.11	
<i>Average</i>	<i>0.42 a</i>	<i>0.12 b</i>	<i>0.13 b</i>	<i>0.13 b</i>	<i>9.83</i>

<sup>(1)</sup>Means (n = 5) followed by the same lowercase letter within columns not differ by Tukey test ( $p < 0.05$ ).

<sup>(2)</sup>Coefficient of variation (CV) of cover crop residue error 1

**Table 5** Dynamics of black oat and ryegrass shoot residues decomposition in a mature peach orchard. Remaining dry matter, total organic carbon, nitrogen, cellulose, lignin, and non-structural biomass were adjusted to the model  $X = X_0 e^{-kt}$ ; where X is the amount of dry matter or nutrient remaining in the residue after a period t (days),  $X_0$  is the initial amount of dry matter or nutrient in the residue, k is the decomposition constant rate and  $t_{1/2}$  is the half-life for each compartment (Experiment 2)


Treatment	k	$t_{1/2}$	$R^{2(3)}$	k	$t_{1/2}$	$R^2$
	$g\ g^{-1}$	days	-	$g\ g^{-1}$	days	-
Remaining dry matter			Remaining cellulose			
Black Oat	0.0164 b <sup>(1)</sup>	42 a	0.98*	0.0278 b	25 a	0.95*
Ryegrass	0.0208 a	33 b	0.98*	0.0356 a	19 b	0.98*
CV (%) <sup>(2)</sup>	2.61	2.72		9.05	5.34	
Remaining total organic carbon			Remaining lignin			
Black Oat	0.0170 b	41 a	0.97*	0.0325 b	23 a	0.88*
Ryegrass	0.0211 a	33 b	0.97*	0.0614 a	12 b	0.98*
CV (%)	2.26	2.33		13.62	14.16	
Remaining nitrogen			Remaining non-structural biomass			
Black Oat	0.0118 a	54 a	0.94*	0.0117 a	59 a	0.91*
Ryegrass	0.0129 a	59 a	0.97*	0.0459 a	50 a	0.93*
CV (%)	6.06	11.38		41.36	29.48	

<sup>(1)</sup>Means (n = 5) followed by the same lowercase letter do not differ by the Tukey test ( $p < 0.05$ ). <sup>(2)</sup> Coefficient of variation (CV) of the decomposition constant rate (k) and the half-life for each compartment ( $t_{1/2}$ ). <sup>(3)</sup> Coefficient of determination ( $R^2$ ) of the residue's decomposition dynamics model. \* significant at  $p < 0.05$

**Table 6** Carbon/nitrogen (C/N) ratio, dry matter (DM), total organic carbon (TOC), and nitrogen (N) in the remaining shoot residues of black oat and ryegrass deposited on the soil surface in mature peach trees planting rows (Experiment 2)


Treatments	Day after litter bags deposition on the soil surface						
	0	30	60	90	120	150	180
C/N ratio							
Black oat shoot	16.3	8.7	7.0	5.5	5.8	6.7	6.1
Ryegrass shoot	15.7	6.6	4.3	5.1	5.0	7.6	5.9
<i>Significance</i> <sup>(1)</sup>	2 SEM = 0.479						
CV (%)	8.20	15.3	23.9	24.4	23.0	40.8	10.1
DM (kg ha <sup>-1</sup> )							
Black oat shoot	5177	2989	1573	1277	1120	745	161
Ryegrass shoot	5607	2702	1277	1078	838	825	142
<i>Significance</i>	2 SEM = 61.0						
CV (%)	3.79	7.13	16.1	9.72	12.1	20.3	26.0
TOC (kg ha <sup>-1</sup> )							
Black oat shoot	2201	1126	562	512	443	290	44.7
Ryegrass shoot	2289	976	420	419	327	280	69.5
<i>Significance</i>	2 SEM = 35.8						
CV (%)	16.80	7.62	16.81	9.79	11.56	23.53	3.11
N (kg ha <sup>-1</sup> )							
Black oat shoot	136	131	74.8	94.9	77.2	42.7	9.90
Ryegrass shoot	149	150	88.8	83.6	61.1	47.2	12.5
<i>Significance</i>	2 SEM = 4.12						
CV (%)	9.85	13.2	7.34	18.4	20.6	9.00	23.5

<sup>(1)</sup> Values differing by 2 standard error of means (SEM) are statistically different




556

557 **Fig. 1** Average air temperature and cumulative monthly rainfall after the deposition of litter  
 558 bags in the experimental area



Leaf sampling time after cover crop shoot residue deposition (days)

**Fig. 2** Total N a), atom% <sup>15</sup>N excess b), and <sup>15</sup>N derived from shoot residues c) in mature peach trees leaves grown with black oat and ryegrass shoot residues deposition on the soil surface (Experiment 1). Vertical bars indicate the standard error of the mean (n = 3). Lowercase letters compare black oat and ryegrass residues within each sampling time by the Tukey-test ( $p < 0.05$ ), ns = not significant



**Fig. 3** Remaining percentage of dry matter (DM) a), total organic carbon (TOC) b), nitrogen (N) c), cellulose (Cel) d), lignin (Lig) e), and non-structural biomass (Bio) f) in black oat and ryegrass shoot residues deposited on the planting rows of a mature peach tree orchard (Experiment 2)