



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Smoothing effect for time-degenerate Schrödinger operators

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Smoothing effect for time-degenerate Schrödinger operators / Federico, Serena; Staffilani, Gigliola. - In: JOURNAL OF DIFFERENTIAL EQUATIONS. - ISSN 0022-0396. - ELETTRONICO. - 298:(2021), pp. 205-247. [10.1016/j.jde.2021.07.006]

Availability:

This version is available at: <https://hdl.handle.net/11585/842243> since: 2021-12-18

Published:

DOI: <http://doi.org/10.1016/j.jde.2021.07.006>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Serena Federico, Gigliola Staffilani, Smoothing effect for time-degenerate Schrödinger operators, Journal of Differential Equations, Volume 298, 2021, Pages 205-247

The final published version is available online at
<https://dx.doi.org/10.1016/j.jde.2021.07.006>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

SMOOTHING EFFECT FOR TIME-DEGENERATE SCHRÖDINGER OPERATORS

SERENA FEDERICO AND GIGLIOLA STAFFILANI

ABSTRACT. In this work we consider an example of a linear time degenerate Schrödinger operator. We show that with the appropriate assumptions the operator satisfies a Kato smoothing effect. We also show that the solutions to the nonlinear initial value problems involving this operator and polynomial derivative nonlinearities are locally well-posed and their solutions also satisfy the same smoothing estimates as the linear solutions.

1. INTRODUCTION

In this paper we study the smoothing effect for time-degenerate Schrödinger operators of the form

$$(1) \quad \mathcal{L}_\alpha = i\partial_t + t^\alpha \Delta_x + b(t, x) \cdot \nabla_x,$$

where the coefficients $b(t, x) = (b_1(t, x), \dots, b_n(t, x))$ are *complex valued* and satisfy suitable decay assumptions, and $\alpha > 0$.

In the case $b(t, x) \equiv 0$, or when $b(t, x) = ct^\alpha$, where c is a complex vector, we show below that standard Fourier analysis arguments can be applied, and the results will follow by application of more classical techniques.

In the more general situation we deal with space-time variable coefficients, at least in the first order part, and we need to replace the standard use of the Fourier transform with the use of pseudo-differential calculus. This will allow us to obtain smoothing estimates for the linear operator (1) and its non homogenous counterpart. Smoothing estimates for an operator such as (1) where $\alpha = 0$ are by now classical results, see for example [1, 2, 4, 6].

Although the central part of this work is dedicated to linear smoothing estimates, in the second part of the paper we also address local well-posedness of the related Cauchy problem with derivative nonlinearities. As in the case of nondegenerate space variable coefficients Schrödinger equations (see [7, 8, 11] and references therein), also in our case local well-posedness relies heavily on the smoothing estimates proved in the first part of this work.

In proving smoothing estimates for the operator (1), the main problems are given by the presence of the time degeneracy in the second order term and by the presence of the first order term $b(t, x) \cdot \nabla_x$. In particular, the time degeneracy has to be managed in

This project has received funding from the European Unions Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 838661 and No 777822.

G.S. was partially founded by the NSF grant No DMS-1764403.

order to apply a method similar to that of Mizohata [12] and Doi [3] to absorb the space-time variable coefficients first order term through the application of the Sharp Gårding inequality. The presence of the term $b(t, x) \cdot \nabla_x$ affects the applicability of the Sharp Gårding inequality as well, and, in addition, determines, in general, a loss of derivatives.

Due to these considerations it is clear that conditions on $b(t, x)$ are necessary to control the behavior of the operator, and, specifically, conditions relating the coefficient t^α and the coefficients $b_j(t, x)$. It is also well known that even in the case $\alpha = 0$ and $b(t, x) = b(x)$ conditions that are necessary (and sufficient) need to be imposed on $b(x)$ even for local well-posedness to hold, see [12].

Our approach to overcome these problems in proving the linear and non homogeneous smoothing effect estimates is inspired by techniques similar to those in [7, 8], which themselves are based on a method proposed by Mizohata [12] and Doi [3]. The key point is the construction of a pseudo-differential operator K possessing good (specific) properties with respect to the second order operator $t^\alpha \Delta_x$ and that permits to control the first order term $b(t, x) \cdot \nabla_x$. Since the leading coefficient depends only on the time-variable t , we can perform our analysis by keeping t as a parameter provided that b satisfies suitable conditions with respect to t as well. This simplifies considerably the problem, since, this way, we can choose the operator K related to Δ_x instead of $t^\alpha \Delta_x$.

The way the operator K is introduced in the argument is by defining a new norm equivalent to the H_x^s -norm, and in terms of K itself. Thanks to this new norm we will perform useful commutator estimates giving the smoothing effect with a gain of one derivative for the non homogeneous term with respect to the regularity of the initial data (which is necessary to deal with derivative nonlinearities).

In order to extend the smoothing effect estimates we prove in the first part of this work to the nonlinear problem with derivative nonlinearities we follow the work of Kenig, Ponce and Vega who studied smoothing effect and local well-posedness of the nonlinear Cauchy problem both for space-variable nondegenerate Schrödinger operators and for space-variable nondegenerate ultrahyperbolic Schrödinger operators (see [7, 8], and also [13, 11] and references therein).

In the context of time degenerate Schrödinger operators, at least in relation to Cauchy problems, to the best of our knowledge, the only result currently available is due to Cicognani and Reissig in [1], in which they study the local well-posedness of the homogeneous Cauchy problem for time degenerate Schrödinger operators of the same form as (1) (or even more general) in Sobolev and Gevrey spaces. However, our approach is different from that used in [1], in which neither the analysis of the smoothing effect nor the local well-posedness of the derivative nonlinear Cauchy problem is considered.

In this paper we study the following nonlinear prototypes associated with (1):

$$(2) \quad \begin{cases} \mathcal{L}_\alpha u = \pm u |u|^{2k} \\ u(0, x) = u_0(x), \end{cases}$$

and

$$(3) \quad \begin{cases} \mathcal{L}_\alpha u = \pm t^\beta \nabla u \cdot u^{2k}, & \beta \geq \alpha > 0, \\ u(0, x) = u_0(x). \end{cases}$$

We remark immediately that although in this paper we explicitly prove results for (3), as mentioned again in Section 7, the same techniques we use here work when the nonlinearity is a polynomial $t^\beta P(u, \bar{u}, \nabla u, \nabla \bar{u})$. In the case (2), the proofs when¹ $b \equiv 0$ and when $b \not\equiv 0$ are treated separately. This is done in order to show how classical technics can be applied to obtain the local well-posedness in the first case, while, in the more general case $b \not\equiv 0$, the use of pseudo-differential calculus will be needed to get the result.

In the IVP (3), we use a nonlinearity where the time factor appears. This choice is dictated by the fact that it allows for the application of the weighted smoothing estimates at our disposal directly. In addition this type of nonlinearity does not give any limitation on the exponent α , which will be any nonnegative real number. If $\beta > \alpha$ then a contraction mapping theorem based on the fact that in the analysis of the nonlinear term a power $t^{\beta-\alpha}$ appears makes the analysis pretty straightforward. On the other hand when $\beta = \alpha$ more care needs to be used. The result follows from the combination of the technique used here for the IVP (3) when $\beta > \alpha$, and that used in [7, 8] for the nondegenerate case (compare with the problem (1.3) in [8]) in order to remove the smallness of the initial data. The main point consists in fact in modifying the functional space to which the solution belongs in a way that permits to obtain, via a mean value theorem, a time factor needed to apply the contraction argument. The norm to be used is the one used in [7, 8] where the time derivative of the solution is taken into account. Since the proof of this result follows the lines described above we shall omit the proof which is left to the reader.

We want to stress, once again, that the local solutions of the nonlinear problems (2) and (3) above satisfy the weighted smoothing estimates stated below.

Moreover, results concerning more general degenerate Schrödinger operators can be obtained following the procedure described above. In fact, one might consider

$$\mathcal{L}_\alpha = i\partial_t + \sum_{i,j=1}^n D_{x_i} a_{ij}(t, x) D_{x_j} + b(t, x) \cdot \nabla_x,$$

where $a_{ij}(t, x)$ is such that $a_{ij}(t, x) \sim t^\alpha a'_{i,j}(x)$, and, for all $i, j = 1, \dots, n$, $a'_{i,j} \in C_b^\infty(\mathbb{R}^n)$, $\{a'_{i,j}(x)\}_{i,j=1,\dots,n}$ is real valued and positive definite, and $|\partial_x^\gamma a_{i,j}(t, x)| \lesssim t^\alpha \langle x \rangle^{-|\gamma|-\sigma+1}$ for any multiindex $\gamma \in \mathbb{N}^n$, where σ is dictated by the conditions satisfied by $b(t, x)$ (see (10) below). By combining our technique and that in [8] one should be able to prove the weighted smoothing effect, and, as a consequence, that the associated IVP (both linear and nonlinear similar to the ones above) is locally well-posed.

Notations. We recall here, briefly, some notations used throughout the paper.

We use the notation $A \lesssim B$ to indicate that there exists an absolute constant $c > 0$ such that $A \leq cB$. We shall denote by Λ^s the pseudo-differential operator of order s whose symbol is given by $\Lambda^s(\xi) = \langle \xi \rangle^s = (1 + |\xi|^2)^{s/2}$.

Below we often use mixed norm spaces. For example $L_x^p L_t^q([0, T] \times \mathbb{R}^n)$, $1 \leq p, q \leq \infty$ is the space of functions $f(x, t)$ that are in L^q in time on the interval $[0, T]$ and are L^p in

¹As mentioned above the case when $b(t, x) = ct^\alpha$ and c a complex vector, can be treated like the case $b \equiv 0$.

space. The norm is taken in the right to left order. In a similar manner we define the spaces $L^p([0, T]; H^s(\mathbb{R}^n))$, $1 \leq p \leq \infty$ of functions that are L^p in time and in the Sobolev space $H^s(\mathbb{R}^n)$ in space. As remarked in [7], also in this case if the vector $b(t, x) = b(x)$ is a real smooth enough function, then standard energy method gives well-posedness. On the other hand in this paper we are not just concerned with well-posedness, but more importantly with the proof of smoothing estimates.

We shall now summarize the main results of the paper.

Case $b \equiv 0$. We state here the results when $b \equiv 0$. We will remark later that the same results hold when $b = ct^\alpha$, where c is an imaginary vector.

Smoothing effect estimates. Let $W_\alpha(t, s)$ be the operator defined as in (14).

Theorem 1.1. *Let $W_\alpha(t) := W_\alpha(t, 0)$, with $\alpha > 0$, then*

If $n = 1$ for all $f \in L^2(\mathbb{R})$,

$$(4) \quad \sup_x \|t^{\alpha/2} D_x^{1/2} W_\alpha(t) f\|_{L_t^2([0, T])}^2 \lesssim \|f\|_{L^2(\mathbb{R})}^2;$$

If $n \geq 2$, on denoting by $\{Q_\beta\}_{\beta \in \mathbb{Z}^n}$ the family of non overlapping cubes of unit size such that $\mathbb{R}^n = \bigcup_{\beta \in \mathbb{Z}^n} Q_\beta$, then for all $f \in L_x^2(\mathbb{R}^n)$,

$$(5) \quad \sup_{\beta \in \mathbb{Z}^n} \left(\int_{Q_\beta} \int_0^T |t^{\alpha/2} D_x^{1/2} W_\alpha(t) f(x)|^2 dt dx \right)^{1/2} \lesssim \|f\|_{L^2(\mathbb{R}^n)},$$

where $D_x^\gamma f(x) = (|\xi|^\gamma \widehat{f}(\xi))^\vee(x)$.

Theorem 1.2. *Let $n = 1$ and $g \in L_x^1 L_t^2([0, T] \times \mathbb{R})$, then*

$$(6) \quad \|D_x^{1/2} \int_{\mathbb{R}_+} t^{\alpha/2} W_\alpha(0, t) g(t) dt\|_{L_x^2(\mathbb{R})} \lesssim \|g\|_{L_x^1 L_t^2(\mathbb{R} \times [0, T])},$$

and, for all $g \in L_t^1 L_x^2([0, T] \times \mathbb{R})$,

$$(7) \quad \|t^{\alpha/2} D_x^{1/2} \int_0^t W_\alpha(t, \tau) g(\tau) d\tau\|_{L_x^\infty(\mathbb{R}) L_t^2([0, T])} \lesssim \|g\|_{L_t^1 L_x^2([0, T] \times \mathbb{R})}.$$

If $n \geq 2$, denoting by $\{Q_\beta\}_{\beta \in \mathbb{Z}^n}$ a family of non overlapping cubes of unit size such that $\mathbb{R}^n = \bigcup_{\beta \in \mathbb{Z}^n} Q_\beta$, then, for all $g \in L_t^1 L_x^2([0, T] \times \mathbb{R}^n)$,

$$(8) \quad \sup_{\beta \in \mathbb{Z}^n} \left(\int_{Q_\beta} \left\| t^{\alpha/2} D_x^{1/2} \int_0^t W_\alpha(t, \tau) g(\tau) d\tau \right\|_{L_t^2([0, T])}^2 dx \right)^{1/2} \lesssim \|g\|_{L_t^1 L_x^2([0, T] \times \mathbb{R}^n)},$$

Local well-posedness.

Theorem 1.3. *Let $k \geq 1$, then (2) is locally well-posed in H^s for $s > n/2$ and its solution satisfies smoothing estimates.*

Case $b \neq 0$. Let us consider the IVP

$$(9) \quad \begin{cases} \partial_t u = it^\alpha \Delta_x u + ib(t, x) \cdot \nabla_x u + f(t, x) \\ u(0, x) = u_0(x). \end{cases}$$

The results true in this case are true in general, that is, also when $b \equiv 0$.

Smoothing effect estimates.

Theorem 1.4. *Let $u_0 \in H^s(\mathbb{R}^n)$, $s \in \mathbb{R}$. Assume that, for all $j = 1, \dots, n$, b_j is such that $b_j \in C([0, T], C_b^\infty(\mathbb{R}^n))$ and there exists $\sigma > 1$ such that*

$$(10) \quad |\operatorname{Im} \partial_x^\gamma b_j(t, x)|, |\operatorname{Re} \partial_x^\gamma b_j(t, x)| \lesssim t^\alpha \langle x \rangle^{-\sigma-|\gamma|}, \quad x \in \mathbb{R}^n,$$

and denote by $\lambda(|x|) := \langle x \rangle^{-\sigma}$.

Then

- (i) *If $f \in L^1([0, T]; H^s(\mathbb{R}^n))$ then the IVP (9) has a unique solution $u \in C([0, T]; H^s(\mathbb{R}^n))$ and there exist positive constants C_1, C_2 such that*

$$\sup_{0 \leq t \leq T} \|u(t)\|_s \leq C_1 e^{C_2(\frac{T^{\alpha+1}}{\alpha+1} + T)} \left(\|u_0\|_s + \int_0^T \|f(t)\|_s dt \right);$$

- (ii) *If $f \in (L^2[0, T]; H^s(\mathbb{R}^n))$ then the IVP (9) has a unique solution $u \in C([0, T]; H^s(\mathbb{R}^n))$ and there exist two positive constants C_1, C_2 such that*

$$\sup_{0 \leq t \leq T} \|u(t)\|_s^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \left| \Lambda^{s+1/2} u \right|^2 \lambda(|x|) dx dt \leq C_1 e^{C_2(\frac{T^{\alpha+1}}{\alpha+1} + T)} \left(\|u_0\|_s^2 + \int_0^T \|f(t)\|_s^2 dt \right);$$

- (iii) *If $\Lambda^{s-1/2} f \in L^2([0, T] \times \mathbb{R}^n; t^{-\alpha} \lambda(|x|)^{-1} dt dx)$ then the IVP (9) has a unique solution $u \in C([0, T]; H^s(\mathbb{R}^n))$ and there exist positive constants C_1, C_2 such that*

$$\begin{aligned} & \sup_{0 \leq t \leq T} \|u(t)\|_s^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \left| \Lambda^{s+1/2} u \right|^2 \lambda(|x|) dx dt \\ & \leq C_1 e^{C_2 \frac{T^{\alpha+1}}{\alpha+1}} \left(\|u_0\|_s^2 + \int_0^T \int_{\mathbb{R}^n} t^{-\alpha} \lambda(|x|)^{-1} \left| \Lambda^{s-1/2} f \right|^2 dx dt \right). \end{aligned}$$

Above we abbreviated the norm $\|f\|_{H^s(\mathbb{R}^n)} =: \|f\|_s$.

Local well-posedness.

Theorem 1.5. *Let \mathcal{L}_α be such that condition (10) is satisfied. Then the IVP (2) is locally well posed in H^s for $s > n/2$.*

Theorem 1.6. *Let \mathcal{L}_α be such that condition (10) is satisfied with $\sigma = 2N$ (thus $\lambda(|x|) = \langle x \rangle^{-2N}$) for some $N \geq 1$, and $s > n + 4N + 3$ such that $s - 1/2 \in 2\mathbb{N}$. Let $H_\lambda^s := \{u_0 \in H^s(\mathbb{R}^n); \lambda(|x|)u_0 \in H^s(\mathbb{R}^n)\}$, then the IVP (3) with $\beta \geq \alpha > 0$, is locally well posed in H_λ^s .*

We remark that in all our local well-posedness results we do not focus on the optimal index s for the H^s regularity of the initial data. Instead our goal in these theorems is to show that the nonlinear solutions enjoy the same type of smoothing estimates as the linear ones.

We conclude this introduction by giving the plan of the paper. In Section 2 we start the analysis of the case $b \equiv 0$, and we remark how $b = ct^\alpha$, with c a constant imaginary vector, can be analyzed in a similar manner. First we show that the solution (which is explicit) to the linear initial value problem involving \mathcal{L}_α can be written via the use of a two-parameter family of unitary operators (similarly to the standard case). Afterwards, by using standard Fourier analysis methods we derive homogeneous time-weighted smoothing estimates (where the weight depends on the degeneracy) by reducing the case $\alpha \neq 0$ (degenerate case) to the nondegenerate case $\alpha = 0$. This is the first part of Theorem 1.1.

In Section 3 we show that the Duhamel's principle still applies in our context and prove inhomogeneous time-weighted smoothing estimates with a gain of $1/2$ derivative with respect to the initial data. This is Theorem 1.2.

In Section 4 we state the local well-posedness result for the nonlinear Cauchy problem in the case $b \equiv 0$ in which we consider a nonlinearity of the form $u|u|^{2k}$ with $k \geq 1$, and we prove that smoothing estimates propagate to these nonlinear solutions. This is Theorem 1.3.

In Section 5 we start the analysis of the more general case $b \neq 0$ (but the results are true when $b \equiv 0$ as well). We state the hypothesis on the coefficients b_j and prove the local smoothing effect estimates in weighted Sobolev spaces. Additionally, we prove the local well-posedness of the linear Cauchy problem by means of the smoothing estimates. This is Theorem 1.4.

In Section 6 we analyze the local well-posedness of the nonlinear Cauchy problem in presence of two different nonlinearities. First we consider nonlinearities of the form $u|u|^{2k}$, $k \geq 1$. This is Theorem 1.5. Afterwards, we consider derivative nonlinearities of the form $t^\beta \nabla u \cdot u^{2k}$, where $\nabla u := \text{div}(u)$. This is Theorem 1.6.

Finally Section 7 contains some final remarks.

2. THE CASE $\mathcal{L}_\alpha = i\partial_t + t^\alpha \Delta_x$. HOMOGENEOUS SMOOTHING PROPERTIES

We start with the analysis of the homogeneous Cauchy problem

$$(11) \quad \begin{cases} \partial_t u = it^\alpha \Delta_x u \\ u(s, x) = u_s(x), \end{cases}$$

where $0 \leq s < t \leq T$, $x \in \mathbb{R}^n$ and u_0 is at least in $L^2(\mathbb{R}^n)$.

Observe that, by application of the Fourier transform with respect to the space variable, we get

$$\begin{cases} \partial_t \widehat{u}(t, \xi) = -it^\alpha |\xi|^2 \widehat{u}(t, \xi) \\ \widehat{u}(s, \xi) = \widehat{u}_s(\xi), \end{cases}$$

whose solution at time $t \leq T$ is given by

$$\widehat{u}(t, \xi) = e^{-i \frac{t^{\alpha+1} - s^{\alpha+1}}{\alpha+1} |\xi|^2} \widehat{u}_s(\xi),$$

and finally, by Fourier inversion formula,

$$(12) \quad u(t, x) = \int_{\mathbb{R}^n} e^{-i \left(\frac{t^{\alpha+1} - s^{\alpha+1}}{\alpha+1} |\xi|^2 - x \cdot \xi \right)} \widehat{u}_s(\xi) d\xi.$$

Formula (12), giving the solution of the homogeneous problem at $0 < t \leq T$ starting at time $s < t$, can be written as

$$(13) \quad u(t, x) = W_\alpha(t, s) u_s(x) := e^{i \frac{t^{\alpha+1} - s^{\alpha+1}}{\alpha+1} \Delta_x} u_s(x) := \frac{(\alpha+1)^{n/2}}{(i(t^{\alpha+1} - s^{\alpha+1}))^{n/2}} e^{i(\alpha+1) \frac{|\cdot|^2}{t^{\alpha+1} - s^{\alpha+1}}} * u_s(x).$$

Therefore throughout the paper we shall use the notation

$$(14) \quad W_\alpha(t, s) := e^{i \frac{t^{\alpha+1} - s^{\alpha+1}}{\alpha+1} \Delta_x}, \quad \forall s, t \in [0, T],$$

to indicate the operator defined as in (13). Note that $\{W_\alpha(t, s)\}_{s, t \in [0, T]}$ is a two-parameter family of unitary operators and that, for any given (t, s) , $W_\alpha(t, s)$ is the "solution operator" of the IVP (11), that is, $u(t, x) = W_\alpha(t, s) u_s(x)$ is the solution at time t of (11) starting at time s . Moreover the following properties hold:

- (i): $W_\alpha(t, t) = I$;
- (ii): $W_\alpha(t, s) = W_\alpha(t, r) W_\alpha(r, s)$ for every $s, t, r \in [0, T]$;
- (iii): $W_\alpha(t, s) \Delta_x u = \Delta_x W_\alpha(t, s) u$.

In particular we are interested in solutions starting at time $s = 0$ (where the operator is degenerate), that, by (14), will be given by $u(t, x) = W_\alpha(t, 0) u_0(x)$. In what follows we shall often use the notation $W_\alpha(t, 0) =: W_\alpha(t)$.

Note that, if $\alpha = 0$ and $t > 0$,

$$(15) \quad W_0(t) = e^{it\Delta_x},$$

which is the standard Schrödinger semigroup.

By (14) we can easily see the first property of W_α , namely

$$(16) \quad \|W_\alpha(t, s) u_s\|_{H_x^s} = \|u_s\|_{H_x^s}.$$

The second property of the operator W_α is given by the local smoothing result of Theorem 1.1 that we prove below.

Proof of Theorem 1.1. First note that (4) and (5) are true when $\alpha = 0$ in (11), that is, when $W_\alpha(t) = W_0(t) = e^{it\Delta_x}$ is the standard Schrödinger semigroup (see, for instance, [6]). Then it suffices to prove that

$$(17) \quad \|t^{\alpha/2} D_x^{1/2} W_\alpha(t) f\|_{L_t^2([0, T])}^2 = C_\alpha \|D_x^{1/2} W_0(t) f\|_{L_t^2([0, T'])}^2,$$

where C_α is a positive constant depending on α and $T' > 0$, since then the result will follow directly from the standard case $\alpha = 0$. We then reduce the proof to the proof of (17).

We have that

$$(18) \quad \|t^{\alpha/2} D_x^{1/2} W_\alpha(t) f\|_{L_t^2([0, T])}^2 = \int_0^T \left| t^{\alpha/2} \int_{\mathbb{R}^n} e^{-i(t^{\alpha+1}|\xi|^2/(\alpha+1) - x \cdot \xi)} |\xi|^{1/2} \widehat{f}(\xi) d\xi \right|^2 dt,$$

then we apply the change of variables $t^{\alpha+1}/(\alpha+1) = s$, $t = c_\alpha s^{1/(\alpha+1)}$, $c_\alpha = (\alpha+1)^{1/(\alpha+1)}$, $dt = c'_\alpha s^{-\alpha/(\alpha+1)} ds$ with $c'_\alpha = c_\alpha/(\alpha+1)$, and get

$$\begin{aligned} (18) &= \int_0^{\frac{T^{\alpha+1}}{\alpha+1}} c'_\alpha s^{-\alpha/(\alpha+1)} \left| \int_{\mathbb{R}^n} e^{-i(s|\xi|^2 - x \cdot \xi)} |\xi|^{1/2} \widehat{f}(\xi) d\xi \right|^2 c'_\alpha s^{-\alpha/(\alpha+1)} ds \\ &= c_\alpha^\alpha c'_\alpha \int_0^{\frac{T^{\alpha+1}}{\alpha+1}} \left| \int_{\mathbb{R}^n} e^{-i(t|\xi|^2 - x \cdot \xi)} |\xi|^{1/2} \widehat{f}(\xi) d\xi \right|^2 dt \\ &= C_\alpha \int_0^{\frac{T^{\alpha+1}}{\alpha+1}} |D_x^{1/2} W_0(t) f(x)|^2 dt \\ &= C_\alpha \|D_x^{1/2} W_0(t) f\|_{L_t^2([0, T^{\alpha+1}/(\alpha+1)])}^2, \end{aligned}$$

which gives the result. Finally, by application of the smoothing estimates for $W_0(t) = e^{it\Delta_x}$, we get (4) and (5) (see [6], Corollary 2.2). \square

Remark 2.1. Now we consider the case $b(x, t) = ct^\alpha$, where c is imaginary. We note that in his case

$$(19) \quad W_\alpha(t, s) := e^{i\frac{t^{\alpha+1} - s^{\alpha+1}}{\alpha+1}(\Delta_x + c \cdot \nabla_x)} s, t \in [0, T],$$

and

$$W_\alpha(\widehat{t}, 0)u_0(\xi) = e^{-i\frac{\widehat{t}^{\alpha+1}}{\alpha+1}(|\xi|^2 - ic \cdot \xi)},$$

and the argument proceeds exactly as above.

3. THE CASE $-i\partial_t + t^\alpha \Delta_x$. INHOMOGENEOUS SMOOTHING PROPERTIES

We consider here the inhomogeneous Cauchy problem

$$(20) \quad \begin{cases} \partial_t u = it^\alpha \Delta_x u + f(t, x) \\ u(0, x) = u_0(x), \end{cases}$$

where $(t, x) \in [0, T] \times \mathbb{R}^n$, u_0 is at least in $L^2(\mathbb{R}^n)$ and f is at least in $L^2([0, T] \times \mathbb{R}^n)$, and we prove that Duhamel's formula still applies in this case.

Proposition 3.1. *Let $u_0 \in L_x^2(\mathbb{R}^n)$, then the solution at time $t > 0$ of the IVP (initial value problem) (20) is given by*

$$(21) \quad u(t, x) = W_\alpha(t)u_0(x) + \int_0^t W_\alpha(t, \tau)f(\tau, x)d\tau.$$

Proof. To prove (21) we will prove that the solution of the IVP (20) is given by

$$(22) \quad u(t, x) = u_1(t, x) + \int_0^t u_2(\tau, x)d\tau,$$

where u_1 and u_2 are the solutions of

$$\text{IVP1} = \begin{cases} \partial_t u_1 = it^\alpha \Delta_x u \\ u_1(0, x) = u_0(x), \end{cases} \quad \text{IVP2} = \begin{cases} \partial_t u_2 = it^\alpha \Delta_x u \\ u_2(\tau, x) = f(\tau, x), \end{cases}$$

respectively.

By (12) we have that, for t fixed and $\tau < t$,

$$u_1(t, x) = W_\alpha(t)u_0(x),$$

and

$$u_2(t, x) = W_\alpha(t, \tau)f(\tau, x) := e^{i\left(\frac{t^{\alpha+1}-\tau^{\alpha+1}}{\alpha+1}\right)\Delta_x} f(\tau, x).$$

We now suppose that $u(t, x)$ is the solution at time t of (20), and show that u is exactly given by (22). Note that, by using the fact that $W_\alpha(t, s)$ commutes with Δ_x , then if u solves (20) we get

$$\begin{aligned} \frac{d}{dt}W_\alpha(0, t)u(t, x) &= \frac{d}{dt} \left(e^{-i\frac{t^{\alpha+1}}{\alpha+1}\Delta_x} u(t, x) \right) \\ &= -it^\alpha \Delta_x W_\alpha(0, t)u(t, x) + W_\alpha(0, t) \frac{d}{dt}u(t, x) \\ &= -it^\alpha \Delta_x W_\alpha(0, t)u(t, x) + W_\alpha(0, t) (it^\alpha \Delta_x u(t, x) + f(t, x)) \\ &= \cancel{-it^\alpha \Delta_x W_\alpha(0, t)u(t, x)} + \cancel{it^\alpha W_\alpha(0, t)\Delta_x u(t, x)} + W_\alpha(0, t)f(t, x), \end{aligned}$$

which gives

$$\frac{d}{dt}W_\alpha(0, t)u(t, x) = W_\alpha(0, t)f(t, x).$$

We then integrate the last equality

$$\int_0^t \frac{d}{ds}W_\alpha(0, s)u(s, x)ds = \int_0^t W_\alpha(0, \tau)f(\tau, x)d\tau$$

and find

$$W_\alpha(0, t)u(t, x) = W_\alpha(0, 0)u_0(x) + \int_0^t W_\alpha(0, \tau)f(\tau, x)d\tau,$$

which gives, by applying $W_\alpha(t) := W_\alpha(t, 0)$ on both sides, and recalling that $W_\alpha(t, t) = I$,

$$u(t, x) = W_\alpha(t)u_0(x) + W_\alpha(t) \int_0^t W_\alpha(0, \tau)f(\tau, x)d\tau$$

$$\begin{aligned}
&= W_\alpha(t)u_0(x) + \int_0^t W_\alpha(t, \tau)f(\tau, x)d\tau \\
&= u_1(t, x) + \int_0^t u_2(\tau, x)d\tau.
\end{aligned}$$

Therefore, if u solves (20) it is of the form (22), that is, u is given by formula (21).

We conclude the proof by verifying that u given by (21) satisfies (20).

We have

$$\begin{aligned}
\frac{d}{dt}u(t, x) &\stackrel{(21)}{=} it^\alpha \Delta_x W_\alpha(t)u_0(x) + \int_0^t it^\alpha \Delta_x W_\alpha(t, \tau)f(\tau, x) d\tau + f(t, x) \\
&= it^\alpha \Delta_x \left(W_\alpha(t)u_0(x) + \int_0^t W_\alpha(t, \tau)f(\tau, x) dt \right) + f(t, x) \\
&= it^\alpha \Delta_x u(t, x) + f(t, x),
\end{aligned}$$

thus u satisfies (20). □

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2. The first inequality (6) follows directly from (4) by duality.

As regards (7), denoting $L_x^p := L_x^p(\mathbb{R}^n)$, we have

$$\begin{aligned}
&\|t^{\alpha/2} D_x^{1/2} \int_0^t W_\alpha(t, \tau)g(\tau)d\tau\|_{L_x^\infty L_t^2([0, T])} \\
&\leq \left\| \left(\int_0^T \left| \int_0^T |t^{\alpha/2} D_x^{1/2} W_\alpha(t, \tau)g(\tau)| d\tau \right|^2 dt \right)^{1/2} \right\|_{L_x^\infty} \\
&\stackrel{\text{Minkowski}}{\leq} \left\| \int_0^T \left(\int_0^T |t^{\alpha/2} D_x^{1/2} W_\alpha(t, \tau)g(\tau)|^2 dt \right)^{1/2} d\tau \right\|_{L_x^\infty} \\
&= \int_0^T \left\| t^{\alpha/2} D_x^{1/2} W_\alpha(t, 0)(W_\alpha(0, \tau)g(\tau)) \right\|_{L_x^\infty L_t^2([0, T])} d\tau \\
&\stackrel{\text{by (4)}}{\leq} \int_0^T \|W_\alpha(0, \tau)g(\tau)\|_{L_x^2} d\tau = \|g\|_{L_t^1([0, T])L_x^2},
\end{aligned}$$

which gives (7).

As for (8) we first observe that

$$\begin{aligned}
&\left\| t^{\alpha/2} D_x^{1/2} \int_0^t W_\alpha(t, \tau)g(\tau)d\tau \right\|_{L_t^2([0, T])} \\
&\leq \left\| \int_0^T |t^{\alpha/2} D_x^{1/2} W_\alpha(t, \tau)g(\tau)| d\tau \right\|_{L_t^2([0, T])} \\
&\stackrel{\text{Minkowski}}{\leq} \int_0^T \|t^{\alpha/2} D_x^{1/2} W_\alpha(t, 0)W_\alpha(0, \tau)g(\tau)\|_{L_t^2([0, T])} d\tau,
\end{aligned}$$

therefore

$$\begin{aligned}
& \left(\int_{Q_\beta} \left\| t^{\alpha/2} D_x^{1/2} \int_0^t W_\alpha(t, \tau) g(\tau) d\tau \right\|_{L_t^2([0, T])}^2 dx \right)^{1/2} \\
& \leq \left[\int_{Q_\beta} \left(\int_0^T \| t^{\alpha/2} D_x^{1/2} W_\alpha(t, \tau) g(\tau) \|_{L_t^2([0, T])}^2 d\tau \right) dx \right]^{1/2} \\
& \stackrel{\text{Minkowski}}{\leq} \int_0^T \left(\int_{Q_\beta} \| t^{\alpha/2} D_x^{1/2} W_\alpha(t, 0) W_\alpha(0, \tau) g(\tau) \|_{L_t^2([0, T])}^2 dx \right)^{1/2} d\tau.
\end{aligned}$$

We then apply the $\sup_{\beta \in \mathbb{Z}^n}$ on both the RHS and the LHS of the latter inequality and get

$$\begin{aligned}
& \sup_{\beta \in \mathbb{Z}^n} \left(\int_{Q_\beta} \left\| t^{\alpha/2} D_x^{1/2} \int_0^t W_\alpha(t, \tau) g(\tau) d\tau \right\|_{L_t^2([0, T])}^2 dx \right)^{1/2} \\
& \leq \sup_{\beta \in \mathbb{Z}^n} \left(\int_0^T \left(\int_{Q_\beta} \| t^{\alpha/2} D_x^{1/2} W_\alpha(t, 0) W_\alpha(0, \tau) g(\tau) \|_{L_t^2([0, T])}^2 dx \right) d\tau \right)^{1/2} \\
& \leq \int_0^T \sup_{\beta \in \mathbb{Z}^n} \left(\int_{Q_\beta} \| t^{\alpha/2} D_x^{1/2} W_\alpha(t, 0) W_\alpha(0, \tau) g(\tau) \|_{L_t^2([0, T])}^2 dx \right)^{1/2} d\tau \\
& \stackrel{\text{by (5)}}{\leq} \int_0^T \| W_\alpha(0, \tau) g(\tau) \|_{L_x^2(\mathbb{R}^n)} d\tau \\
& = \int_0^T \| g(\tau) \|_{L_x^2(\mathbb{R}^n)} d\tau = \| g \|_{L_t^1([0, T]) L_x^2(\mathbb{R}^n)}
\end{aligned}$$

which concludes the proof. \square

As remarked for the homogeneous problem, also in the inhomogeneous one we considered above we can take $b(x, t) = t^\alpha c$ where c is an imaginary vector and obtain the same results.

4. LOCAL WELL-POSEDNESS FOR THE NONLINEAR CAUCHY PROBLEM

Let us now consider the nonlinear initial value problem

$$(23) \quad \begin{cases} \partial_t u = it^\alpha \Delta_x u \pm u|u|^{2k} \\ u(0, x) = u_0(x), \end{cases}$$

where $k \geq 1$ is a positive integer. We shall prove that (23) is locally well posed in $H^s(\mathbb{R}^n)$, for $s > n/2$.

In the sequel we shall use the notation $W_\alpha(t, 0) =: W_\alpha(t)$.

Definition 4.1. We say that the IVP (23) is locally well-posed (l.w.p) in $H^s(\mathbb{R}^n)$ if for any ball B in the space $H^s(\mathbb{R}^n)$ there exist a time T and a Banach space of functions $X \subset L^\infty([0, T], H^s(\mathbb{R}^n))$ such that for each initial datum $u_0 \in B$ there exists a unique solution $u \in X \subset C([0, T], H^s(\mathbb{R}^n))$ for the integral equation

$$u(x, t) = W_\alpha(t)u_0 + \int_0^t W_\alpha(t, \tau)|u|^{2k}u(\tau)d\tau.$$

Furthermore the map $u_0 \mapsto u$ is continuous as a map from $H^s(\mathbb{R}^n)$ into $C([0, T], H^s(\mathbb{R}^n))$.

To prove the local well-posedness we shall use the following result.

Lemma 4.0.1. Let $g(u) = u|u|^{2k}$ and s, l positive integers with $l \leq s$ and $s > n/2$. Then

$$(24) \quad \|g(u)\|_{H^s} \lesssim \|u\|_{H^s}^{2k+1},$$

$$(25) \quad \|g(u) - g(v)\|_{L^2} \lesssim (\|u\|_{H^s}^{2k} + \|v\|_{H^s}^{2k})\|u - v\|_{L^2},$$

$$(26) \quad \|g^{(l)}(u) - g^{(l)}(v)\|_{L^\infty} \lesssim (\|u\|_{H^s}^{2k-l} + \|v\|_{H^s}^{2k-l})\|u - v\|_{H^s},$$

$$(27) \quad \|g(u) - g(v)\|_{H^s} \lesssim (\|u\|_{H^s}^{2k} + \|v\|_{H^s}^{2k})\|u - v\|_{H^s}.$$

We are now ready to prove Theorem 1.3.

Proof of Theorem 1.3. The proof is based on the standard fixed point argument. For convenience we shall assume that the nonlinear term is given by $+u|u|^{2k}$ but the proof applies with no modification in the focusing case. We assume first that $n = 1$. Let X be the following metric space

$$X = \{u : [0, T] \times \mathbb{R} \rightarrow \mathbb{C}; \|t^{\alpha/2}D_x^{1/2+s}u\|_{L_x^\infty L_t^2([0, T])} < \infty, \|u\|_{L_t^\infty([0, T])H_x^s} < \infty\},$$

equipped with the distance

$$d(u, v) = \|t^{\alpha/2}D_x^{1/2+s}(u - v)\|_{L_x^\infty L_t^2([0, T])} + \|u - v\|_{L_t^\infty([0, T])\dot{H}_x^s} + \|u - v\|_{L_t^\infty([0, T])L_x^2}$$

in which \dot{H}_x^s stands for the homogeneous Sobolev space, and consider

$$\Phi : X \rightarrow X, \quad \Phi(u) = W_\alpha(t)u_0 + \int_0^t W_\alpha(t, \tau)u|u|^{2k}(\tau)d\tau.$$

We now prove that Φ is a contraction, since then the result follows by application of the fixed point theorem.

We have that

$$\|\Phi(u)\|_X \leq \|W_\alpha(t)u_0\|_X + \left\| \int_0^t W_\alpha(t, \tau)u|u|^{2k}(\tau)d\tau \right\|_X,$$

and consider the two terms on the RHS separately.

For the homogeneous term we have

$$\|W_\alpha(t)u_0\|_X = \|t^{\alpha/2}D_x^{1/2+s}W_\alpha(t)u_0\|_{L_x^\infty L_t^2([0, T])} + \|W_\alpha(t)u_0\|_{L_t^\infty([0, T])\dot{H}_x^s} + \|W_\alpha(t)u_0\|_{L_t^\infty([0, T])L_x^2}$$

$$\begin{aligned} &\stackrel{\text{by(4)}}{\leq} \|D_x^s u_0\|_{L_x^2} + 2\|u_0\|_{H_x^s} \\ &\leq 3\|u_0\|_{H^s}. \end{aligned}$$

For the nonlinear term we get

$$\begin{aligned} &\left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_X = \|t^{\alpha/2} D_x^{1/2+s} \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau\|_{L_t^\infty L_x^2} \\ &+ \left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_{L_t^\infty([0, T]) \dot{H}_x^s} + \left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_{L_t^\infty([0, T]) L_x^2} \\ &\stackrel{(7)}{\leq} \|D_x^s u |u|^{2k}\|_{L_t^1([0, T]) L_x^2} + \left\| \int_0^t W_\alpha(t, \tau) D_x^s u |u|^{2k}(\tau) d\tau \right\|_{L_t^\infty([0, T]) L_x^2} \\ &\quad + \left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_{L_t^\infty([0, T]) L_x^2}. \end{aligned}$$

Note that

$$\begin{aligned} &\|D_x^s u |u|^{2k}\|_{L_t^1([0, T]) L_x^2} \\ &\leq T \|u |u|^{2k}\|_{L_t^\infty([0, T]) H_x^s} \\ &\stackrel{\text{by (24)}}{\leq} CT \|u\|_{L_t^\infty([0, T]) H_x^s}^{2k+1} \\ &\leq CT \|u\|_X^{2k+1}, \end{aligned}$$

and that

$$\begin{aligned} &\left\| \int_0^t W_\alpha(t, \tau) D_x^s u |u|^{2k}(\tau) d\tau \right\|_{L_t^\infty([0, T]) L_x^2} = \|W_\alpha(t) \int_0^t W_\alpha(0, \tau) D_x^s u |u|^{2k}(\tau) d\tau\|_{L_t^\infty([0, T]) L_x^2} \\ &\leq \left(\int_{\mathbb{R}} \left| \int_0^T |W_\alpha(0, \tau) D_x^s u |u|^{2k}(\tau)|^2 d\tau \right| dx \right)^{1/2} \\ &\leq \int_0^T \left(\int_{\mathbb{R}} |W_\alpha(0, \tau) D_x^s u |u|^{2k}(\tau)|^2 dx \right)^{1/2} d\tau \\ &= \|D_x^s u |u|^{2k}\|_{L_t^1([0, T]) L_x^2} \\ &\leq T \|u |u|^{2k}\|_{L_t^\infty([0, T]) H_x^s} \\ &\leq CT \|u\|_{L_t^\infty([0, T]) H_x^s}^{2k+1} \\ &\leq CT \|u\|_X^{2k+1}. \end{aligned}$$

Similarly

$$\begin{aligned} &\left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_{L_t^\infty([0, T]) L_x^2} \\ &\leq \|D_x^s u |u|^{2k}\|_{L_t^1([0, T]) L_x^2} \\ &\leq CT \|u\|_X^{2k+1}, \end{aligned}$$

therefore, by application of the previous inequalities we get

$$\left\| \int_0^t W_\alpha(t, \tau) D_x^s u |u|^{2k}(\tau) d\tau \right\|_X \leq CT \|u\|_X^{2k+1}.$$

Putting together the previous estimates we have

$$\|\Phi(u)\|_X \leq 3\|u_0\|_{H_x^s} + C_1 T \|u\|_X^{2k+1}$$

which gives that $\Phi : X \rightarrow X$.

Let $R = 6\|u_0\|_{H_x^s}$ and consider $B_R = \{u \in X; \|u\|_X \leq R\} \subset X$. Then, by choosing T such that $C_1 T R^{2k} \leq 1/2$, we get that, for all $u \in B_R$

$$\|\Phi(u)\|_X \leq R/2 + C_1 T R^{2k+1} \leq R,$$

which gives that Φ sends B_R in B_R .

We now prove that Φ is a contraction.

Let $u, v \in B_R$, and consider

$$\begin{aligned} \|\Phi(u) - \Phi(v)\|_X &= \left\| \int_0^t W_\alpha(t, \tau) (u|u|^{2k}(\tau) - v|v|^{2k}(\tau)) d\tau \right\|_X \\ &= \|t^{\alpha/2} D_x^{s+1/2} \int_0^t W_\alpha(t, \tau) (u|u|^{2k}(\tau) - v|v|^{2k}(\tau)) d\tau\|_{L_t^\infty L_x^2([0, T])} \\ &\quad + \left\| \int_0^t W_\alpha(t, \tau) (u|u|^{2k}(\tau) - v|v|^{2k}(\tau)) d\tau \right\|_{L_t^\infty([0, T]) \dot{H}_x^s} \\ &\quad + \left\| \int_0^t W_\alpha(t, \tau) (u|u|^{2k}(\tau) - v|v|^{2k}(\tau)) d\tau \right\|_{L_t^\infty([0, T]) L_x^2} \\ &= I + II + III. \end{aligned}$$

By using the same estimates used before we have that

$$\begin{aligned} I &\leq \|D^s(u|u|^{2k} - v|v|^{2k})\|_{L_t^1([0, T]) L_x^2} \\ &\leq T \|u|u|^{2k} - v|v|^{2k}\|_{L_t^\infty([0, T]) H_x^s} \\ &\stackrel{\text{by (27)}}{\leq} CT (\|u\|_{L_t^\infty([0, T]) H_x^s}^{2k} + \|v\|_{L_t^\infty([0, T]) H_x^s}^{2k}) \|u - v\|_{L_t^\infty([0, T]) H_x^s} \\ &\leq CT (\|u\|_X^{2k} + \|v\|_X^{2k}) \|u - v\|_X \\ &\leq CTR^{2k} \|u - v\|_X. \end{aligned}$$

For II we estimate the $L_t^\infty([0, T]) \dot{H}_x^s$ - norm as before and get, once more,

$$\begin{aligned} II &\leq \|D^s(u|u|^{2k} - v|v|^{2k})\|_{L_t^1([0, T]) L_x^2} \\ &\leq CTR^{2k} \|u - v\|_X, \end{aligned}$$

while, once again,

$$\begin{aligned} III &\leq \|u|u|^{2k} - v|v|^{2k}\|_{L_t^1([0, T]) L_x^2} \\ &\leq CTR^{2k} \|u - v\|_X. \end{aligned}$$

We then have

$$\|\Phi(u) - \Phi(v)\|_X \leq C_2 T R^{2k} \|u - v\|_X,$$

where T was chosen in such a way that $C_1 T R^{2k} \leq 1/2$. We then choose the time T such that $T = \min\{\frac{1}{C_1 R^{2k}}, \frac{1}{C_2 R^{2k}}\}$, and conclude that Φ is a contraction. The result then follows by fixed point arguments.

We now assume that $n > 1$. We proceed like above and we consider the space

$$X = \{u : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{C}; \|t^{\alpha/2} D_x^{s+1/2} u\|_T < \infty, \|u\|_{L_{[0,T]}^\infty H_x^s} < \infty\},$$

where

$$\|\cdot\|_T = \sup_{\beta \in \mathbb{Z}^n} \|\cdot\|_{L_x^2(Q_\beta) L_t^2([0, T])},$$

and

$$d_X(u, v) = \|t^{\alpha/2} D_x^{s+1/2}(u - v)\|_T + \|u - v\|_{L_t^\infty([0, T]) \dot{H}_x^s} + \|u - v\|_{L_t^\infty([0, T]) L_x^2}.$$

Once again we consider

$$\Phi : X \rightarrow X, \quad \Phi(u) = W_\alpha(t) u_0 + \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau,$$

and prove that Φ is a contraction.

We have that

$$\|\Phi(u)\|_X \leq \|W_\alpha(t) u_0\|_X + \left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_X,$$

and we estimate the two terms on the RHS separately.

For the homogeneous term we have

$$\begin{aligned} \|W_\alpha(t) u_0\|_X &= \|t^{\alpha/2} D_x^{s+1/2} W_\alpha(t) u_0\|_T + \|W_\alpha(t) u_0\|_{L_t^\infty([0, T]) \dot{H}_x^s} + \|W_\alpha(t) u_0\|_{L_t^\infty([0, T]) L_x^2} \\ &\leq \|t^{\alpha/2} D_x^{1/2} W_\alpha(t) D_x^s u_0\|_T + 2 \|W_\alpha(t) u_0\|_{L_t^\infty([0, T]) H_x^s} \\ &= \sup_{\beta \in \mathbb{Z}^n} \|t^{\alpha/2} D_x^{1/2} W_\alpha(t) D_x^s u_0\|_{L^2(Q_\beta) L_t^2([0, T])} + 2 \|W_\alpha(t) u_0\|_{L_t^\infty([0, T]) H_x^s} \\ &\stackrel{(8)}{\leq} \|D_x^s u_0\|_{L^2(\mathbb{R}^n)} + 2 \|u_0\|_{H_x^s} \\ &\leq 3 \|u_0\|_{H_x^s}. \end{aligned}$$

For the inhomogeneous term we have

$$\begin{aligned} (28) \quad \left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_X &\leq \underbrace{\left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_T}_{(28.1)} \\ &\quad + \underbrace{2 \left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_{L_t^\infty([0, T]) H_x^s}}_{(28.2)}, \end{aligned}$$

where

$$\begin{aligned}
(28.1) &= \sup_{\beta \in \mathbb{Z}^n} \|t^{\alpha/2} D_x^{1/2} \int_0^t W_\alpha(t, \tau) D_x^s u |u|^{2k}(\tau) d\tau\|_{L^2(Q_\beta) L_t^2([0, T])} \\
&\stackrel{(8)}{\leq} \|D_x^s u |u|^{2k}\|_{L_t^1([0, T]) L^2(\mathbb{R}^n)} \leq T \|u |u|^{2k}\|_{L_t^\infty([0, T]) H^s(\mathbb{R}^n)} \\
&\leq T \|u\|_{L_t^\infty([0, T]) H^s(\mathbb{R}^n)}^{2k+1} \leq CT \|u\|_X^{2k+1},
\end{aligned}$$

and

$$\begin{aligned}
(28.2) &= \sup_{t \in [0, T]} \left\| \int_0^t W_\alpha(t, \tau) u |u|^{2k}(\tau) d\tau \right\|_{H^s(\mathbb{R}^n)} \\
&= \sup_{t \in [0, T]} \left\| \int_0^t W_\alpha(t, \tau) (1 + D_x^{2s})^{1/2} u |u|^{2k}(\tau) d\tau \right\|_{L_x^2} \\
&\leq \sup_{t \in [0, T]} \left(\int_{\mathbb{R}^n} \left| \int_0^t |W_\alpha(t, \tau) (1 + D_x^{2s})^{1/2} u |u|^{2k}(\tau)| d\tau \right|^2 dx \right)^{1/2} \\
&\leq \sup_{\text{Minkowski } t \in [0, T]} \left(\int_0^t \|(1 + D_x^{2s})^{1/2} u |u|^{2k}(\tau)\|_{L^2(\mathbb{R}^n)}^2 d\tau \right) \\
&= \int_0^T \|u |u|^{2k}(\tau)\|_{H^s(\mathbb{R}^n)}^2 d\tau \\
&= \|u |u|^{2k}\|_{L_t^1([0, T]) H^s(\mathbb{R}^n)} \leq T \|u |u|^{2k}\|_{L_t^\infty([0, T]) H^s(\mathbb{R}^n)} \\
&\leq T \|u\|_{L_t^\infty([0, T]) H^s(\mathbb{R}^n)}^{2k+1} \leq CT \|u\|_X^{2k+1}.
\end{aligned}$$

Therefore, putting together the previous estimates, we get

$$\|\Phi(u)\|_X \leq 3\|u_0\|_{H^s} + C_1 T \|u\|_X^{2k+1},$$

hence $\Phi : X \rightarrow X$.

Let now $R = 6\|u_0\|_{H^s(\mathbb{R}^n)}$ and T such that $C_1 T R^{2k} \leq 1/2$, then $\|\Phi(u)\|_X \leq R$ for all $u \in B_R$, that is, Φ sends B_R into B_R .

We now end the proof by showing that Φ is a contraction.

Let $u, v \in B_R$, then

$$\begin{aligned}
\|\Phi(u) - \Phi(v)\|_X &= \left\| \int_0^t W_\alpha(t, \tau) (u |u|^{2k}(\tau) - v |v|^{2k}(\tau)) d\tau \right\|_X \\
&\leq \|t^{\alpha/2} D_x^{s+1/2} \int_0^t W_\alpha(t, \tau) (u |u|^{2k}(\tau) - v |v|^{2k}(\tau)) d\tau\|_T \\
&+ \left\| \int_0^t W_\alpha(t, \tau) (u |u|^{2k}(\tau) - v |v|^{2k}(\tau)) d\tau \right\|_{L_t^\infty([0, T]) H_x^s(\mathbb{R}^n)} \\
&= I + II,
\end{aligned}$$

where, by (27) and the procedure used before,

$$I, II \leq \|u |u|^{2k} - v |v|^{2k}\|_{L_t^1([0, T]) H_x^s(\mathbb{R}^n)}$$

$$\leq CT(\|u\|_{L_t^\infty([0,T])H_x^s(\mathbb{R}^n)}^{2k} + \|v\|_{L_t^\infty([0,T])H_x^s(\mathbb{R}^n)}^{2k})\|u - v\|_{L_t^\infty([0,T])H_x^s(\mathbb{R}^n)}^2.$$

Now, since $u, v \in B_R$, we get

$$\|\Phi(u) - \Phi(v)\|_X \leq C_2TR^{2k}\|u - v\|_{L_t^\infty([0,T])H_x^s(\mathbb{R}^n)}^2.$$

Finally, by suitably choosing the time T we get that the operator Φ is a contraction and the result follows.

5. THE CASE $b \neq 0$: SMOOTHING ESTIMATES

We now consider the more general case (1) where $b \neq 0$. For simplicity of notation let us rename

$$(29) \quad \mathcal{L}_\alpha = i\partial_t + t^\alpha \Delta_x + b(t, x) \cdot \nabla_x,$$

where $\nabla_x = (\partial_{x_1}, \dots, \partial_{x_n})$, $\Delta_x = \sum_{j=1}^n D_j^2$, $b(t, x) = (b_1(t, x), \dots, b_n(t, x))$, $b_j \in C([0, T], C_b^\infty(\mathbb{R}^n))$ for all $j = 1, \dots, n$, and $C_b^\infty(\mathbb{R}^n) = \{f \in C^\infty(\mathbb{R}^n) : \partial^\alpha f \in L^\infty(\mathbb{R}^n), \forall \alpha \in \mathbb{Z}_+^n\}$. Moreover we assume that there exist $c_0 > 0$ and $\sigma > 1$ such that, for all $j = 1, \dots, n$,

$$|\operatorname{Im} \partial_x^\gamma b_j(t, x)|, |\operatorname{Re} \partial_x^\gamma b_j(t, x)| \lesssim t^\alpha \langle x \rangle^{-\sigma-|\gamma|}, \quad \text{for } |\gamma| = 0, 1.$$

Our goal here is to prove some weighted smoothing estimates similar to the previous ones for the operator above.

Remark 5.1. *The smoothing estimates we are going to prove in this section are better than the ones proved in the case $b \equiv 0$, in the sense that for the non homogenous term we will be able to obtain some smoothing estimates with a gain of one derivative with respect to the regularity of initial data in the IVP. The smoothing estimates below can, of course, be applied to the case $b \equiv 0$ as well. Moreover, one can prove these estimates in a more direct way in the case $b \equiv 0$. However, since we are interested in the case $b \neq 0$ in which a direct proof is not applicable, we shall give the proof of the result for the general case directly.*

As we shall see, a key point in the proof of the smoothing properties for (29) is the use of Doi's lemma (Lemma A.0.1 in the Appendix).

We use Lemma A.0.1 on the symbol $a^w := a = a_2 + ia_1 + a_0$ such that $a_2(x, \xi) = |\xi|^2$ and $a_1 = a_0 = 0$. In this case conditions (B1) and (B2) of Lemma A.0.1 are trivially satisfied, while (A6) holds with $q(x, \xi) = x \cdot \xi \langle \xi \rangle^{-1}$. Therefore, by Lemma A.0.1 with $\lambda'(|x|) = C' \langle x \rangle^{-\sigma}$ (see Remark A.1), with C' is to be chosen later, we get that there exists $p \in S^0$ and $C > 0$ such that (54) holds.

We then consider the pseudo-differential operator K with symbol $K(x, \xi) = e^{p(x, \xi)} \Lambda^s$, where $\Lambda^s := \langle \xi \rangle^s$ and $p(x, \xi)$ is given by Doi's lemma, and define the norm N on $H^s(\mathbb{R}^n)$, equivalent to the standard one (see [8] for the proof of the equivalence), given by

$$(30) \quad N(u)^2 = \|Ku\|_0^2 + \|u\|_{s-1}^2,$$

where $\|\cdot\|_s$ stands for the standard norm in the Sobolev space $H^s(\mathbb{R}^n)$.

Finally, following the technique used in [8], we make use of the norm N to prove smoothing properties of the solutions of the IVP (9) that we stated in Theorem 1.4.

The proof of this theorem is essentially reduced to the proof of the following lemma.

Lemma 5.0.1. *Let $s \in \mathbb{R}$, $\lambda(|x|) := \langle x \rangle^{-\sigma}$ and $\sigma > 1$ such that (10) holds. Then there exists $C_1, C_2 > 0$ such that, for all $u \in C([0, T]; H^{s+2}(\mathbb{R}^n)) \cap C^1([0, T]; H^s(\mathbb{R}^n))$, we have*

(31)

$$\sup_{0 \leq t \leq T} \|u(t)\|_s \leq C_1 e^{C_2(\frac{T^{\alpha+1}}{\alpha+1} + T)} \left(\|u_0\|_s + \int_0^T \|(\partial_t - it^\alpha \Delta_x - ib(t, x) \cdot \nabla_x)u(t, \cdot)\|_s dt \right);$$

(32)

$$\sup_{0 \leq t \leq T} \|u(t)\|_s \leq C_1 e^{C_2(\frac{T^{\alpha+1}}{\alpha+1} + T)} \left(\|u(\cdot, T)\|_s + \int_0^T \|(\partial_t - it^\alpha \Delta_x - ib(t, x) \cdot \nabla_x)^* u(t, \cdot)\|_s dt \right);$$

(33)

$$\begin{aligned} & \sup_{0 \leq t \leq T} \|u(t)\|_s^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \left| \Lambda^{s+1/2} u \right|^2 \lambda(|x|) dx dt \\ & \leq C_1 e^{C_2(\frac{T^{\alpha+1}}{\alpha+1} + T)} \left(\|u_0\|_s^2 + \int_0^T \|(\partial_t - it^\alpha \Delta_x - ib(t, x) \cdot \nabla_x)u(t, \cdot)\|_s^2 dt \right); \end{aligned}$$

(34)

$$\begin{aligned} & \sup_{0 \leq t \leq T} \|u(t)\|_s^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \left| \Lambda^{s+1/2} u \right|^2 \lambda(|x|) dx dt \\ & \leq C_1 e^{C_2 \frac{T^{\alpha+1}}{\alpha+1}} \left(\|u_0\|_s^2 + \int_0^T \int_{\mathbb{R}^n} t^{-\alpha} \lambda(|x|)^{-1} \left| \Lambda^{s-1/2} (\partial_t - it^\alpha \Delta_x - ib(t, x) \cdot \nabla_x) u(t, \cdot) \right|^2 dx dt \right). \end{aligned}$$

Proof. Recall that we defined the norm $N(u)$ on $H^s(\mathbb{R}^n)$ to be the one given in (30). We then estimate the quantity

$$\partial_t N(u)^2 = \partial_t \|Ku\|_0^2 + \partial_t \|u\|_{s-1}^2 = I + II.$$

For the term II we have

$$\begin{aligned} II &= \partial_t \|u\|_{s-1}^2 = \partial_t \langle \Lambda^{s-1} u, \Lambda^{s-1} u \rangle = 2\operatorname{Re} \langle \Lambda^{s-1} \partial_t u, \Lambda^{s-1} u \rangle \\ &= 2\operatorname{Re} \langle \Lambda^{s-1} (it^\alpha \Delta_x + ib(t, x) \cdot \nabla_x) u, \Lambda^{s-1} u \rangle \\ &= 2 \underbrace{\operatorname{Re} \langle it^\alpha \Lambda^{s-1} \Delta_x u, \Lambda^{s-1} u \rangle}_{=0} - 2\operatorname{Re} \langle \Lambda^{s-1} b(t, x) \cdot D_x u, \Lambda^{s-1} u \rangle + 2\operatorname{Re} \langle \Lambda^{s-1} f, \Lambda^{s-1} u \rangle \\ &\leq Ct^\alpha \|u\|_s^2 + 2\operatorname{Re} \langle \Lambda^{s-1} f, \Lambda^{s-1} u \rangle, \end{aligned}$$

where (recall) $D_x = (D_{x_1}, \dots, D_{x_n}) = (-i\partial_{x_1}, \dots, -i\partial_{x_n})$ and $b(t, x) \cdot D_x = \sum_{j=1}^n b_j(t, x) D_{x_j}$, with $b_j \in C([0, T]; C_b^\infty(\mathbb{R}^n))$ such that (10) holds.

Observe that the following estimates hold:

(35)

$$2\operatorname{Re} \langle \Lambda^{s-1} f, \Lambda^{s-1} u \rangle \leq 2\|f\|_{s-1} \|u\|_{s-1} \leq CN(f)N(u),$$

and

$$\begin{aligned} & 2\operatorname{Re} \langle \Lambda^{s-1} f, \Lambda^{s-1} u \rangle = 2\operatorname{Re} \langle t^{-\alpha/2} \lambda(|x|)^{-1/2} \Lambda^{s-1/2} f, t^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s-3/2} u \rangle \\ & \leq \|t^{-\alpha/2} \lambda(|x|)^{-1/2} \Lambda^{s-1/2} f\|_0^2 + \|t^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s-3/2} u\|_0^2 \\ & \leq \langle t^{-\alpha} \lambda(|x|)^{-1} \Lambda^{s-1/2} f, \Lambda^{s-1/2} f \rangle + t^\alpha N(u)^2. \end{aligned}$$

(36)

Therefore, by using (35) and (36), we get that II can be estimated by

$$(37) \quad II \leq Ct^\alpha N(u)^2 + C' \min\{N(f)N(u); \langle t^{-\alpha} \lambda(|x|)^{-1} \Lambda^{s-1/2} f, \Lambda^{s-1/2} f \rangle\},$$

with C and C' new suitable constants.

We now consider the term I

$$(38) \quad \begin{aligned} \partial_t \|Ku\|_0^2 &= 2\operatorname{Re}\langle \partial_t Ku, Ku \rangle = 2\operatorname{Re}\langle K\partial_t u, Ku \rangle \\ &= 2\operatorname{Re}\langle K(it^\alpha \Delta_x + ib(t, x) \cdot \nabla_x)u, Ku \rangle + 2\operatorname{Re}\langle Kf, Ku \rangle \\ &= 2\operatorname{Re}\langle it^\alpha [K, \Delta_x]u, Ku \rangle + \underbrace{2\operatorname{Re}\langle it^\alpha \Delta_x Ku, Ku \rangle}_{=0} \\ &\quad - 2\operatorname{Re}\langle K b(t, x) \cdot D_x u, Ku \rangle + 2\operatorname{Re}\langle Kf, Ku \rangle \\ &= 2\operatorname{Re}\langle it^\alpha [K, \Delta_x]u, Ku \rangle - 2\operatorname{Re}\langle [K, b(t, x) \cdot D_x]u, Ku \rangle \\ &\quad - 2\operatorname{Re}\langle b(t, x) \cdot D_x Ku, Ku \rangle + 2\operatorname{Re}\langle Kf, Ku \rangle, \end{aligned}$$

and estimate the term $2\operatorname{Re}\langle [K, b(t, x) \cdot D_x]u, Ku \rangle$ in the the fifth line of (38). Recall that, given two symbols $p_1 \in S^{m_1}, p_2 \in S^{m_2}$ associated with two operators P_1 and P_2 , then we have that the symbol of the commutator $[P_1, P_2](x, D)$ is given by $-i\{p_1, p_2\}(x, \xi) + p_3(x, \xi)$, where $p_3 \in S^{m_1+m_2-2}$. Therefore, since $K(x, \xi) = e^{p(x, \xi)}(1 + |\xi|^2)^{s/2}$ and the symbol of $b \cdot D_x = b(t, x) \cdot D_x$ is $\sum_{j=1}^n b_j(t, x)\xi_j$, we have that the operator $[K, b(t, x) \cdot D_x]$ is of order s and has symbol

$$(39) \quad \begin{aligned} [K, b(t, x) \cdot D_x](t, x, \xi) &= -i\{K(x, \xi), \sum_{j=1}^n b_j(t, x)\xi_j\} + r_{s-1}(t, x, \xi) \\ &= -i \sum_{k=1}^n \left[e^{p(x, \xi)} (\Lambda^s(\xi) \partial_{\xi_k} p(x, \xi) + s\Lambda^{s-1}(\xi)\xi_k) \sum_{j=1}^n (\partial_{x_k} b_j(t, x))\xi_j \right. \\ &\quad \left. - e^{p(x, \xi)} (\partial_{x_k} p(x, \xi)) \Lambda^s(\xi) b_k(t, x) \right] + r_{s-1}(t, x, \xi). \end{aligned}$$

Therefore, by the properties of $b(t, x)$ (recall that $b \in C_b^\infty$ and is bounded, together with its derivatives in space, by $t^\alpha \lambda(|x|)$) we get

$$-2\operatorname{Re}\langle [K, b(t, x)D_x]u, Ku \rangle \leq Ct^\alpha \|u\|_s^2,$$

where we used $\|r_{s-1}(t, x, D)u\|_0 \leq Ct^\alpha \|u\|_{s-1}$ (this estimate is deduced by using the properties of b following [10] Theorem 1.1.20 pag.14).

Note also that, once more by using the pseudo-differential calculus, we get $[K, \Delta_x](x, D) = [p, \Delta_x]K(x, D) + r_s(x, D)$, where r_s is of order s and $p = p(x, D)$ is the operator of order 0 appearing in the definition of the norm $N(\cdot)$.

Now we can estimate (38) in the following way

$$\begin{aligned}
(38) &\leq Ct^\alpha \|u\|_s^2 + 2\operatorname{Re}\langle (it^\alpha[p, \Delta_x](x, D) - b(t, x) \cdot D_x)Ku, Ku \rangle + |2\operatorname{Re}\langle it^\alpha r_s(x, D)u, Ku \rangle| \\
(40) &\leq Ct^\alpha \|u\|_s^2 + 2\operatorname{Re}\langle (it^\alpha[p, \Delta_x](x, D) - b(t, x)D_x)Ku, Ku \rangle,
\end{aligned}$$

where C is a new suitable positive constant.

We denote $Q(x; D) := it^\alpha[p, \Delta_x](x, D) - b(t, x) \cdot D_x$ whose symbol is such that

$$\begin{aligned}
\operatorname{Re}Q(x, \xi) &= \operatorname{Re}(it^\alpha(-i)\{p, -|\xi|^2\}(x, \xi) - b(t, x) \cdot \xi) + r_0 \\
&\leq -t^\alpha\{p, |\xi|^2\}(x, \xi) + |\operatorname{Re}b(t, x) \cdot \xi| + r_0 \\
&\leq -t^\alpha\{p, |\xi|^2\}(x, \xi) + |\operatorname{Re}b(t, x)||\xi| + C_4 \\
&\stackrel{\text{by (54)}}{\leq} -C't^\alpha\lambda(|x|)|\xi| + C_2t^\alpha + C_0t^\alpha\lambda(|x|)|\xi| + C \\
&\leq t^\alpha(C_0 - C')\lambda(|x|)|\xi| + C_2t^\alpha + C_4 \\
&\leq -Ct^\alpha\lambda(|x|)|\xi| + C_2t^\alpha + C \\
&\leq -Ct^\alpha\lambda(|x|)(1 + |\xi|) + Ct^\alpha\lambda(|x|) + C_2t^\alpha + C_4 \\
&\leq -Ct^\alpha\lambda(|x|)(1 + |\xi|^2)^{1/2} + C_3t^\alpha + C_4 \\
&= t^\alpha(-C\lambda(|x|)(1 + |\xi|^2)^{1/2} + C_3) + C_4
\end{aligned}$$

where we chose C' (which is possible by Doi's lemma, see Remark A.1) in order to have $C_0 - C' < 0$.

Due to the property of the symbol of Q we can apply the Gårding inequality and get

$$\begin{aligned}
2\operatorname{Re}\langle Q(x, D)Ku, Ku \rangle &\leq -Ct^\alpha\langle \lambda(|x|)\Lambda^1 Ku, Ku \rangle + C_3t^\alpha\|Ku\|_0^2 + C_4\|Ku\|_0^2 \\
&\leq -Ct^\alpha\langle \lambda(|x|)\Lambda^1 Ku, Ku \rangle + C_3t^\alpha\|u\|_s^2 + C_4\|u\|_s^2
\end{aligned}$$

Since $\lambda \in C_b^\infty$, by using the symbolic calculus we get that

$$\lambda(|x|)\Lambda^1(x, D) = (\lambda(|x|)^{1/2}\Lambda^{1/2})^2(x, D) + r_0(x, D),$$

where $r_0(x, D)$ has order 0.

Then, by the latter property, we get

$$(41) \quad 2\operatorname{Re}\langle Q(x, D)Ku, Ku \rangle \leq -Ct^\alpha\|\lambda(|x|)^{1/2}\Lambda^{1/2}Ku\|_0^2 + C_3t^\alpha\|u\|_s^2 + C_4\|u\|_s^2,$$

where $C > 0$ is a new suitable constant.

By plugging (41) in (40) we get

$$(42) \quad \partial_t\|Ku\|_0 \leq Ct^\alpha N(u)^2 + C'N(u)^2 - C''t^\alpha\|\lambda(|x|)^{1/2}\Lambda^{1/2}Ku\|_0^2 + C'''N(f)N(u),$$

where in the latter we used $\operatorname{Re}\langle Kf, Ku \rangle \leq C'''N(f)N(u)$.

Finally, by using (37) and the equivalence between the norms $\|\cdot\|_s$ and $N(\cdot)$ (see [8] pag.390), we obtain

$$\begin{aligned}
(43) \quad \partial_t N(u)^2 &= \partial_t\|Ku\|^2 + \partial_t\|u\|_{s-1}^2 \\
&\leq Ct^\alpha N(u)^2 + C'N(u)^2 - C''t^\alpha\|\lambda(|x|)^{1/2}\Lambda^{1/2}Ku\|_0^2 + C'''N(f)N(u) + \\
&\quad + C_3 \min\{N(f)N(u); \langle t^{-\alpha}\lambda(|x|)^{-1}\Lambda^{s-1/2}f, \Lambda^{s-1/2}f \rangle\},
\end{aligned}$$

where the constants are (eventually) new suitable constants.

From (43) we will get (31), (32) and (33) as we shall prove below.

Proof of (31). As regards (31) we observe that, from (43) we have

$$\partial_t N(u)^2 \leq C_1(t^\alpha + 1)N(u)^2 + C_2N(u)N(f)$$

(again C_1 and C_2 new constants) which gives,

$$2\partial_t N(u) \leq C_1(t^\alpha + 1)N(u) + C_2N(f)$$

and

$$\partial_t \left(2e^{-\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} N(u) \right) \leq C_2 e^{-\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} N(f).$$

Hence, by integrating in time from 0 to t we get

$$\begin{aligned} N(u(t)) &\leq C e^{\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} \left[N(u(0)) + C_2 \int_0^t e^{-\frac{1}{2}C_1(s^{\alpha+1}/(\alpha+1)+s)} N(f) ds \right] \\ &\leq C' e^{\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} \left[N(u(0)) + \int_0^t N(f) ds \right], \end{aligned}$$

which finally gives (31) by the equivalence of the norms. \square

Proof of (32). The proof of (32) follows from (31) applied to the adjoint operator and with $u(t, \cdot)$ replaced by $u(T - t, \cdot)$. \square

proof of (33). To obtain (33) we first observe that there exists a pseudo-differential operator \tilde{K} such that

$$I = \tilde{K}K + \Psi_{r_{-1}},$$

where $\Psi_{r_{-1}}$ is an operator with symbol r_{-1} of order -1 (see [8] pag.390 for the proof). By using this property we get

$$\begin{aligned} (44) \quad \|\lambda(|x|)^{1/2} \Lambda^{s+1/2} u\|_0 &\leq \|(\lambda(|x|)^{1/2} \Lambda^{1/2})(\Lambda^s \tilde{K})(K \Lambda^{1/2})u\|_0 + O(N(u)) \\ &\leq \|(\Lambda^s \tilde{K})(\lambda(|x|)^{1/2} \Lambda^{1/2})(K \Lambda^{1/2})u\|_0 + cN(u) \\ &\leq c \left(\|(\lambda(|x|)^{1/2} \Lambda^{1/2})(K \Lambda^{1/2})u\|_0 + N(u) \right), \end{aligned}$$

where, in the second line, we used the fact that $[\Lambda^s \tilde{K}, \lambda(|x|)^{1/2} \Lambda^{1/2}]K \Lambda^{1/2}$ is a pseudo-differential operator of order s together with the equivalence of the norms $\|\cdot\|_s$ and $N(\cdot)$. Therefore, again from (43), we have

$$\begin{aligned} \partial_t N(u)^2 &\leq C_1(t^\alpha + 1)N(u)^2 - C''t^\alpha \|\lambda(|x|)^{1/2} \Lambda^{1/2} K u\|_0^2 + C'''N(f)N(u) \\ &\leq C_1(t^\alpha + 1)N(u)^2 - C_2 t^\alpha \langle \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle + C_3 N(u)^2 + C_4 N(f)^2 \\ &\leq C_1(t^\alpha + 1)N(u)^2 - C_2 t^\alpha \langle \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle + C_4 N(f)^2, \end{aligned}$$

where the constants are new suitable constant. Hence

$$\partial_t N(u)^2 + C_2 \langle t^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, t^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle \leq C_1(t^\alpha + 1)N(u)^2 + C_4 N(f)^2,$$

so that by integrating in time from 0 to t ,

$$\begin{aligned}
& N(u(t))^2 + C_2 e^{\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} \times \\
(45) \quad & \times \int_0^t e^{-\frac{1}{2}C_1(s^{\alpha+1}/(\alpha+1)+s)} \langle s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle ds \\
& \lesssim e^{\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} \left[N(u(0))^2 + \int_0^t e^{-\frac{1}{2}C_1(s^{\alpha+1}/(\alpha+1)+s)} N(f)^2 ds \right] \\
& \lesssim e^{\frac{1}{2}C_1(T^{\alpha+1}/(\alpha+1)+T)} \left[N(u(0))^2 + \int_0^t N(f)^2 ds \right].
\end{aligned}$$

From the previous estimate we get

$$(46) \quad \sup_{t \in [0, T]} N(u(t))^2 \lesssim e^{\frac{1}{2}C_1(T^{\alpha+1}/(\alpha+1)+T)} \left[N(u(0))^2 + \int_0^T N(f)^2 ds \right].$$

Moreover, the second term on the LHS of (45) satisfies

$$\begin{aligned}
& e^{\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} \int_0^t e^{-\frac{1}{2}C_1(s^{\alpha+1}/(\alpha+1)+s)} \langle s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle ds \\
& \geq e^{\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} \left(\inf_{s \in [0, T]} e^{-\frac{1}{2}C_1(s^{\alpha+1}/(\alpha+1)+s)} \right) \times \\
& \quad \times \int_0^t \langle s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle ds \\
& \geq \int_0^t \langle s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle ds.
\end{aligned}$$

Therefore, using the previous inequality and (45),

$$\begin{aligned}
(47) \quad & C_2 \int_0^T \langle s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle ds \\
& = C_2 \sup_{t \in [0, T]} \int_0^t \langle s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle ds \\
& \leq C_2 \sup_{t \in [0, T]} e^{\frac{1}{2}C_1(t^{\alpha+1}/(\alpha+1)+t)} \int_0^t e^{-\frac{1}{2}C_1(s^{\alpha+1}/(\alpha+1)+s)} \langle s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u, s^{\alpha/2} \lambda(|x|)^{1/2} \Lambda^{s+1/2} u \rangle ds \\
& \stackrel{(45)}{\leq} e^{\frac{1}{2}C_1(T^{\alpha+1}/(\alpha+1)+T)} \left[N(u(0))^2 + \int_0^T N(f)^2 ds \right].
\end{aligned}$$

Finally, by summing up (46) and (47) we get (33). \square

Proof of (34). To prove (34), denoting by $\lambda := \lambda(|x|)$, we write

$$\begin{aligned}
(48) \quad 2\operatorname{Re}\langle Kf, Ku \rangle &= 2\operatorname{Re}\langle t^{\alpha/2}\lambda^{1/2}\Lambda^{1/2}Kf, t^{-\alpha/2}\lambda^{-1/2}\Lambda^{-1/2}Ku \rangle \\
&\leq \varepsilon \|t^{\alpha/2}\lambda^{1/2}\Lambda^{1/2}Ku\|_0^2 + \frac{1}{\varepsilon} \|t^{-\alpha/2}\lambda^{-1/2}\Lambda^{-1/2}Kf\|_0^2 \\
&= \varepsilon \|t^{\alpha/2}\lambda^{1/2}\Lambda^{1/2}K\Lambda^{-s-1/2}\Lambda^{s+1/2}u\|_0^2 \\
&\quad + \frac{1}{\varepsilon} \|t^{-\alpha/2}\lambda^{-1/2}\Lambda^{-1/2}K\Lambda^{-s+1/2}\Lambda^{s-1/2}f\|_0^2,
\end{aligned}$$

Since $\Lambda^{1/2}K\Lambda^{-s-1/2}$ and $\Lambda^{-1/2}K\Lambda^{-s+1/2}$ are both pseudo-differential operators of order 0 in x , we have

$$t^{\alpha/2}\lambda^{1/2}\Lambda^{1/2}K\Lambda^{-s-1/2} = \Lambda^{1/2}K\Lambda^{-s-1/2}t^{\alpha/2}\lambda^{1/2} + t^{\alpha/2}\Psi_{r-1},$$

where Ψ_{r-1} denotes an operator of order -1 in the space variable. Of course the same property holds for the operator $t^{-\alpha/2}\lambda^{-1/2}\Lambda^{-1/2}K\Lambda^{-s+1/2}$.

We use these properties in (48) to get

$$\begin{aligned}
(49) \quad 2\operatorname{Re}\langle Kf, Ku \rangle &= 2\operatorname{Re}\langle t^{\alpha/2}\lambda^{1/2}\Lambda^{1/2}Kf, t^{-\alpha/2}\lambda^{-1/2}\Lambda^{-1/2}Ku \rangle \\
&\leq c_1\varepsilon \|t^{\alpha/2}\lambda^{1/2}\Lambda^{s+1/2}u\|_0^2 + c_2\frac{1}{\varepsilon} \|t^{-\alpha/2}\lambda^{-1/2}\Lambda^{s-1/2}f\|_0^2 + c_3t^\alpha \|u\|_s^2.
\end{aligned}$$

By using (44) and (49) in (43), and the equivalence between the norms $N(\cdot)$ and $\|\cdot\|_s$, we obtain

$$\partial_t N(u)^2 + (c_0 - c_1\varepsilon) \|t^{\alpha/2}\lambda^{1/2}\Lambda^{s+1/2}u\|_0^2 \leq c_3t^\alpha N(u)^2 + c_2\frac{1}{\varepsilon} \|t^{-\alpha/2}\lambda^{-1/2}\Lambda^{s-1/2}f\|_0^2,$$

where c_j , $j = 0, 1, 2, 3$ are new suitable constants, and we choose $\varepsilon > 0$ such that $c_0 - c_1\varepsilon \geq c > 0$.

Since ε is now fixed, we have

$$\partial_t N(u)^2 + c \|t^{\alpha/2}\lambda^{1/2}\Lambda^{s+1/2}u\|_0^2 \leq c_3t^\alpha N(u)^2 + c_2\frac{1}{\varepsilon} \|t^{-\alpha/2}\lambda^{-1/2}\Lambda^{s-1/2}f\|_0^2,$$

which gives, by integrating in time from 0 to t , and by using the same argument as in the proof of (33), the proof of (34). \square

The proof is then complete. \square

Lemma 5.0.1 allows to prove the well-posedness and smoothing result in Theorem 1.4.

Proof of Theorem 1.4. From (31) of Lemma 5.0.1 we immediately get the uniqueness of the solution. In fact, let u be a solution of the homogeneous IVP for (29), i.e., with $f = 0$, and initial data $u_0 = 0$. Then, by (31) of Lemma 5.0.1, we get $u = 0$ and thus the uniqueness (even in the general case $f \neq 0$ and $u_0 \neq 0$).

About the existence, we get the results by using density arguments as we will prove below.

Case 1: $f \in \mathcal{S}(\mathbb{R}^{n+1})$ and $u_0 \in \mathcal{S}(\mathbb{R}^n)$.

We consider the subspace

$$E = \{P^*\varphi; \varphi \in C_0^\infty(\mathbb{R}^n \times [0, T])\} = (\partial_t - it^\alpha \Delta_x + b(t, x) \cdot D_x)^*(C_0^\infty(\mathbb{R}^{n+1})) \subset L^1([0, T]; H^{-s}(\mathbb{R}^n))$$

and the linear functional

$$\ell^* : E \rightarrow \mathbb{C}, \quad \ell^*(P^*\varphi) = \int_0^T \langle f, \varphi \rangle_{L^2 \times L^2} dt + \langle u_0, \varphi(\cdot, 0) \rangle_{L^2 \times L^2}.$$

Then, by (32) of Lemma 5.0.1 (applied on φ) with s replaced by $-s$, for $\eta = P^*\varphi$, with $\varphi \in C_0^\infty(\mathbb{R}^n \times [0, T])$ we get

$$\begin{aligned} |\ell^*(\eta)| &\leq \|f\|_{(L^1[0, T]; H_x^s)} \sup_{t \in [0, T]} \|\varphi\|_{H_x^{-s}} + \|u_0\|_{H_x^s} \|\varphi(0)\|_{H_x^{-s}} \\ &\leq e^{C(T^{\alpha+1}/(\alpha+1)+T)} \left(\|f\|_{L_t^1([0, T]; H_x^s)} + \|u_0\|_{H_x^s} \right) \|\eta\|_{L_t^1([0, T]; H_x^{-s})}, \end{aligned}$$

which gives the continuity of ℓ^* on E (the last inequality follows from (32) of Lemma 5.0.1 applied both on the term $\sup_{t \in [0, T]} \|\varphi\|_{H_x^{-s}}$ and $\|\varphi(0)\|_{H_x^{-s}}$ together with the compactness of the support of φ). By the Hahn-Banach theorem we can extend ℓ^* on $L^1([0, T]; H^{-s}(\mathbb{R}^n))$ and finally get the existence of $u \in L^1([0, T]; H^{-s}(\mathbb{R}^n))^* = L^\infty([0, T]; H^s(\mathbb{R}^n))$ such that

$$\ell^*(P^*\varphi) = \langle u, P^*\varphi \rangle_{L^2 \times L^2} = \int_0^T \langle f, \varphi \rangle_{L^2 \times L^2} dt + \langle u_0, \varphi(\cdot, 0) \rangle_{L^2 \times L^2}, \quad \forall \varphi \in C_0^\infty(\mathbb{R}^n \times [0, T])$$

and thus $Pu = f$ in the sense of distributions for $0 < t < T$.

Notice that $Pu \stackrel{\mathcal{D}'}{=} f$ means that $(\partial_t - it^\alpha \Delta_x + b(t, x) \cdot D_x)u \stackrel{\mathcal{D}'}{=} f$ (as distributions on $C_0^\infty([0, T] \times \mathbb{R}^n)$), therefore, since $f \in \mathcal{S}(\mathbb{R}^{n+1})$, we have that $\partial_t u \in (L^\infty[0, T]; H^{s-2}(\mathbb{R}^n))$, which gives $u \in (C([0, T]; H^{s-2}(\mathbb{R}^n)))$. We then use the equation once more, that is $\partial_t u = it^\alpha \Delta_x + b(t, x) \cdot D_x u + f$, and get, by doing the same consideration, that $u \in (C^1[0, T]; H^{s-4}(\mathbb{R}^n))$ and $u(x, 0) = u_0(x)$. Finally, since $u_0 \in H^s(\mathbb{R}^n)$, repeating the previous argument with $s+4$ in place of s we conclude that there exists a solution u of the IVP associated to (29) to which parts (i)-(iv) of Lemma 5.0.1 apply.

Case 2: $f \in L^1([0, T]; H^s(\mathbb{R}^n))$ and $u_0 \in H^s(\mathbb{R}^n)$.

In this case we take two sequences $f_j \in \mathcal{S}(\mathbb{R}^{n+1})$, $v_j \in \mathcal{S}(\mathbb{R}^n)$ such that $f_j \rightarrow f$ in $(L^1([0, T]); H^s(\mathbb{R}^n))$ and $v_j \rightarrow u_0$ in $H^s(\mathbb{R}^n)$.

By the arguments of case 1 we find a solution u_j of the IVP associated with (29) with f_j and v_j in place of f and u_0 respectively. Since u_j satisfies (31) of Lemma 5.0.1, we have that u_j is a Cauchy sequence, therefore, passing to the limit, we get that $u = \lim_{j \rightarrow \infty} u_j$ is a solution of the IVP with initial data f and in initial data u_0 satisfying (32) of Lemma 5.0.1, which proves (ii) of the theorem.

Case 3: $f \in L^2([0, T]; H^s(\mathbb{R}^n))$ and $u_0 \in H^s(\mathbb{R}^n)$.

Here we proceed as in case 2 where, instead, $f_j \in \mathcal{S}(\mathbb{R}^{n+1})$ is such that $f_j \rightarrow f$ in $(L^2([0, T]; H^s(\mathbb{R}^n)))$. Under this hypothesis we get point (ii) of the theorem, that is, it exists a solution $u \in (C[0, T]; H^s(\mathbb{R}^n))$ satisfying (33) of Lemma 5.0.1.

Case 4: $\Lambda^{s-1/2}f \in (L^2(\mathbb{R}^n \times [0, T]) : t^{-\alpha}\lambda(|x|)^{-1}dxdt)$ and $u_0 \in H^s(\mathbb{R}^n)$.

In this case it is possible to prove that there exists $g_j \in \mathcal{S}(\mathbb{R}^{n+1})$ such that $g_j \rightarrow \Lambda^{s-1/2}f$ in $(L^2(\mathbb{R}^n) \times [0, T] : t^{-\alpha}\lambda(|x|)^{-1}dxdt)$. Applying once again the procedure used in case 1 with f_j replaced by $\Lambda^{-s+1/2}g_j$ in (34) of Lemma 5.0.1, and passing to the limit, we finally get point (iii) of Theorem 1.4. \square

6. THE CASE $b \neq 0$: LOCAL WELL-POSEDNESS OF THE NONLINEAR CAUCHY PROBLEM

We now analyze the local well-posedness of the IVP

$$(50) \quad \begin{cases} \partial_t u = it^\alpha \Delta_x u + ib(t, x) \cdot \nabla_x u + P(u, \bar{u}, \nabla u, \nabla \bar{u}) \\ u(0, x) = u_0(x). \end{cases}$$

under the previous hypotheses on the term b in (50). Note that, with some abuse of notation, the quantity ∇u in the nonlinear term is $\nabla u := \operatorname{div}(u) := \sum_{j=1}^n \partial_{x_j} u$, and, similarly, the quantity $\nabla \bar{u}$.

First we shall consider the case $P(u, \bar{u}, \nabla u, \nabla \bar{u}) = P(u) = \pm u|u|^{2k}$, $k \geq 1$, which is treated in Theorem 1.5 and, afterwards, the case $P(u, \bar{u}, \nabla u, \nabla \bar{u}) = \pm t^\beta \nabla_x u \cdot u^{2k}$ in Theorem 1.6. The proof is based on the contraction argument, which, once again, is obtained through the use of the smoothing estimates proved in the previous section.

Remark 6.1. *Observe that in the proof below we will assume that the solution of the homogeneous problem is again of the form $W_\alpha(t)u_0$. Since we know from Theorem 1.4 that the solution of the linear problem exists, we assume that there exists a two-parameter family of operators, denoted $W_\alpha(t, \tau)$, giving the solution at time t of the homogeneous problem with initial condition at time τ (recall, $W_\alpha(t, 0) := W_\alpha(t)$). Under this assumption one can prove that Duhamel's formula still applies, therefore it makes sense to consider the operator Φ in the form given above. This is important to keep in mind since we will apply the same strategy in the subsequent case as well, that is, in the case $P(u, \bar{u}, \nabla u, \nabla \bar{u}) = \pm t^\beta \nabla_x u \cdot u^{2k}$.*

Proof of Theorem 1.5. We shall make use of the result in Theorem 1.4 concerning the linear problem to prove the result in the nonlinear case. Once again, we give the proof in the defocusing case since the proof in the focusing case applies with no modifications. According to Theorem 1.4 we have the local well-posedness in H^s , $s > n/2$, for the linear IVP (9). We now write the solution of (50) as

$$(51) \quad u(t, x) = W_\alpha(t)u_0 + \int_0^t W_\alpha(t, \tau)P(u, \bar{u}, \nabla u, \nabla \bar{u})d\tau,$$

where $W_\alpha(t, \tau)$ is a new suitable two-parameter family of unitary operators.

Because of the previous assumption, solving the IVP (50) with $P(u, \bar{u}, \nabla u, \nabla \bar{u}) = u|u|^{2k}$ is equivalent to find the solution of the integral equation

$$u(t, x) = W_\alpha(t)u_0 + \int_0^t W_\alpha(t, \tau)u|u|^{2k}d\tau,$$

therefore, as in the proof of Theorem 1.4, we look for the solution given by the fixed point of the map

$$\Phi(u) := W_\alpha(t)u_0 + \int_0^t W_\alpha(t, \tau)u|u|^{2k} d\tau,$$

defined on

$$X_T^s := \{u : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{C}; \|u\|_{L_t^\infty H_x^s} < \infty, \left(\int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} u|^2 dx dt \right)^{1/2} < \infty\},$$

where, recall, $\lambda(|x|) := \langle x \rangle^\sigma$, with $\sigma > 1$ and such that (10) holds.

Proving the existence of a fixed point for Φ is, once more, equivalent to prove that the map Φ is a contraction, and, in order to do that, we will first show that Φ sends X_T^s into itself. The key point here will be to reduce ourself to a linear case to which the linear smoothing estimates apply.

Observe that, denoting by $v := \Phi(u)$, we have that v solves the linear problem

$$\begin{cases} \partial_t v - it^\alpha \Delta_x v - ib(t, x) \cdot \nabla_x v = u|u|^{2k} \\ v(0, x) = u_0(x), \end{cases}$$

and we can estimate $\|\Phi(u)\|_{X_T^s} = \|v\|_{X_T^s}$ through the smoothing estimates given by point (i) and (ii) of Theorem 1.4.

In particular we define

$$\begin{aligned} \|v\|_{X_T^s} &:= \|v\|_{L_t^\infty H_x^s} + \left(\int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} v|^2 dx dt \right)^{1/2} \\ &= I + II, \end{aligned}$$

where, by (i) of Theorem 1.4,

$$I = \|v\|_{L_t^\infty H_x^s} \leq C e^{C' \frac{T^{\alpha+1}}{\alpha+1} + T} \left(\|u_0\|_s + \int_0^T \|u|u|^{2k}\|_s dt \right)$$

and, by (ii) of Theorem 1.4,

$$\begin{aligned} II &= \left(\int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} v|^2 dx dt \right)^{1/2} \\ &\leq C e^{\frac{C'}{2} \left(\frac{T^{\alpha+1}}{\alpha+1} + T \right)} \left(\|u_0\|_s^2 + \int_0^T \|u|u|^{2k}\|_s^2 dt \right)^{1/2}. \\ &\leq C e^{C' \left(\frac{T^{\alpha+1}}{\alpha+1} + T \right)} \left(\|u_0\|_s^2 + \left(\int_0^T \|u|u|^{2k}\|_s^2 dt \right)^{1/2} \right). \end{aligned}$$

Following the same computations of Theorem 1.6 we have

$$\int_0^T \|u|u|^{2k}\|_{H_x^s} dt = \|u|u|^{2k}\|_{L_t^1 H_x^s} \leq CT \|u\|_{L_t^\infty H_x^s}^{2k+1}$$

and

$$\int_0^T \| |u|^{2k} \|_{H_x^s}^2 dt \leq \| |u|^{2k} \|_{L_t^\infty H_x^s} \| |u|^{2k} \|_{L_t^1 H_x^s} \leq T^2 \| |u|^{4k+2} \|_{L_t^\infty H_x^s},$$

so, fixing an upper bound for T , $T \leq 1$ for instance (but not necessarily),

$$\begin{aligned} \|v\|_{X_T^s} &\leq C e^{C' \left(\frac{T^{\alpha+1}}{\alpha+1} + T \right)} \left(\|u_0\|_s + \int_0^T \| |u|^{2k} \|_s dt + \left(\int_0^T \| |u|^{2k} \|_{H_x^s}^2 dt \right)^{1/2} \right) \\ &\leq C e^{C' \left(\frac{1}{\alpha+1} + 1 \right)} \left(\|u_0\|_{H_x^s} + T \| |u|^{2k+1} \|_{L_t^\infty H_x^s} \right) \\ &\leq C \|u_0\|_{H_x^s} + CT \| |u|^{2k+1} \|_{L_t^\infty H_x^s} \\ &\leq C \|u_0\|_{H_x^s} + CT \| |u|^{2k+1} \|_{X_T^s}. \end{aligned}$$

where, with some abuse of notations, C is a new suitable constant.

From the previous estimate we get that Φ sends X_T^s into itself. Moreover, let R be $R = \frac{C}{2} \|u_0\|_{H_x^s}$, then, once again from the previous estimate, for all $u \in B_R \subset X_T^s$ (where B_R denotes the ball of radius R in X_T^s) we have

$$\|\Phi(u)\|_{X_T^s} = \|v\|_{X_T^s}^2 \leq R/2 + CTR^{2k+1},$$

which gives, by choosing $T < 1$ sufficiently small so that $CTR^{2k+1} < R/2$, that Φ sends B_R into B_R .

What is left now is to prove that Φ is a contraction. We then consider $v := \Phi(u)$ and $w := \Phi(u')$, and have

$$\|v - w\|_{X_T^s} = \left\| \int_0^t W_\alpha(t, \tau) (|u|^{2k} - |u'|^{2k}) d\tau \right\|_{X_T^s}.$$

By the previous argument applied to $v - w$, which, in particular, is the solution of the linear problem with $f = |u|^{2k} - |u'|^{2k}$ and initial datum 0, we have (for $T < 1$)

$$\|v - w\|_{X_T^s} \leq C \left(\int_0^T \| |u|^{2k} - |u'|^{2k} \|_s dt + \left(\int_0^T \| |u|^{2k} - |u'|^{2k} \|_s^2 dt \right)^{1/2} \right).$$

By using the estimates used in Theorem 1.5 and Theorem 1.6, we have

$$\begin{aligned} \|v - w\|_{X_T^s} &\leq CT (\| |u|^{2k} \|_{L_t^\infty H_x^s} + \| |u'|^{2k} \|_{L_t^\infty H_x^s}) \| |u - u'| \|_{L_t^\infty H_x^s} \\ &\leq CT (\| |u|^{2k} \|_{X_T^s} + \| |u'|^{2k} \|_{X_T^s}) \| |u - u'| \|_{X_T^s}. \end{aligned}$$

Recalling that $v = \Phi(u)$ and $w = \Phi(u')$, we obtain, for any $u, u' \in B_R$,

$$\|\Phi(u) - \Phi(u')\|_{X_T^s} \leq CTR^{2k} \| |u - u'| \|_{X_T^s}.$$

Finally, eventually by taking T smaller in such a way that $CTR^{2k} < 1$, we conclude that Φ is a contraction, which gives, after application of the standard fixed point argument, the desired result. \square

We now consider the case in which the nonlinearity $P(u, \bar{u}, \nabla u, \nabla \bar{u}) = \pm t^\beta \nabla_x u \cdot u^{2k}$, that is, $P(u, \bar{u}, \nabla u, \nabla \bar{u}) = \pm t^\beta \sum_{j=1}^n \partial_{x_j} u \cdot u^{2k}$. To deal with this case we will need some lemmas that we will borrow from [6] and that we recall in the Appendix (see Lemma A.0.2 and Lemma A.0.3). Additionally, we will make use of the following lemma.

Lemma 6.0.1. *Let $f, g \in H_x^s(\mathbb{R}^n)$, $s > n/2$, such that $\langle x \rangle^{2N} f, \langle x \rangle^{2N} g \in H^{n/2+\varepsilon}$ for some $\varepsilon > 0$ and $N \in \mathbb{N}$, then*

$$\|\langle x \rangle^{2N} f g\|_s \lesssim \|\langle x \rangle^{2N} f\|_{n/2+\varepsilon}^2 \|g\|_s^2 + \|\langle x \rangle^{2N} g\|_{n/2+\varepsilon}^2 \|f\|_s^2.$$

Proof. Since for $|\xi - \eta| \leq |\eta|$ we have $\langle \xi \rangle = (1 + |\xi - \eta + \eta|^2)^{1/2} \lesssim \langle \eta \rangle$, then

$$\begin{aligned} \|\langle x \rangle^{2N} f g\|_s^2 &:= \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} |(I - \Delta_\xi)^N \widehat{f g}(\xi)|^2 d\xi \\ &= \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} \left| (I - \Delta_\xi)^N \left(\int_{\mathbb{R}^n} \widehat{f}(\xi - \eta) \widehat{g}(\eta) d\eta \right) \right|^2 d\xi \\ &= \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} \left| (I - \Delta_\xi)^N \left(\int_{|\xi - \eta| > |\eta|} \widehat{f}(\xi - \eta) \widehat{g}(\eta) d\eta + \int_{|\xi - \eta| \leq |\eta|} \widehat{f}(\xi - \eta) \widehat{g}(\eta) d\eta \right) \right|^2 d\xi \\ &\leq \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} \left| (I - \Delta_\xi)^N \left(\int_{|\xi - \eta| > |\eta|} \widehat{f}(\xi - \eta) \widehat{g}(\eta) d\eta \right) \right|^2 d\xi \\ &\quad + \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} \left| (I - \Delta_\xi)^N \left(\int_{|\xi - \eta| \leq |\eta|} \widehat{f}(\xi - \eta) \widehat{g}(\eta) d\eta \right) \right|^2 d\xi \\ &\lesssim \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} \left| (I - \Delta_\xi)^N \left(\int_{|\gamma| > |\xi - \gamma|} \widehat{f}(\gamma) \widehat{g}(\xi - \gamma) d\eta \right) \right|^2 d\xi \\ &\quad + \int_{\mathbb{R}^n} \left(\int_{|\xi - \eta| \leq |\eta|} \langle \eta \rangle^s |(I - \Delta_\xi)^N \widehat{f}(\xi - \eta)| |\widehat{g}(\eta)| d\eta \right)^2 d\xi \\ &\lesssim \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} \langle \gamma \rangle^s |\widehat{f}(\gamma)| |(I - \Delta_\xi)^N \widehat{g}(\xi - \gamma)| d\eta \right)^2 d\xi \\ &\quad + \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} \langle \eta \rangle^s |(I - \Delta_\xi)^N \widehat{f}(\xi - \eta)| |\widehat{g}(\eta)| d\eta \right)^2 d\xi \\ &\lesssim \| |(I - \Delta_\xi^N) \widehat{g}| * \langle \xi \rangle^s |\widehat{f}| \|_{L^2}^2 + \| |(I - \Delta_\xi)^N \widehat{f}| * \langle \xi \rangle^s |\widehat{g}| \|_{L^2}^2 \\ &\stackrel{\text{Joung's ineq.}}{\lesssim} \| \langle \xi \rangle^s |\widehat{f}| \|_{L^2}^2 \| (I - \Delta_\xi)^N \widehat{g} \|_{L^1}^2 + \| (I - \Delta_\xi)^N \widehat{f} \|_{L^1}^2 \| \langle \xi \rangle^s |\widehat{g}| \|_{L^2}^2 \\ &\lesssim \| f \|_s^2 \int_{\mathbb{R}^n} \frac{1}{\langle \xi \rangle^{n/2+\varepsilon}} \langle \xi \rangle^{n/2+\varepsilon} |(I - \Delta_\xi)^N \widehat{g}| d\xi + \| g \|_s^2 \int_{\mathbb{R}^n} \frac{1}{\langle \xi \rangle^{n/2+\varepsilon}} \langle \xi \rangle^{n/2+\varepsilon} |(I - \Delta_\xi)^N \widehat{f}| d\xi \\ &\lesssim \| f \|_s^2 \| \langle \xi \rangle^{n/2+\varepsilon} (I - \Delta_\xi)^N \widehat{g} \|_{L^2}^2 + \| g \|_s^2 \| \langle \xi \rangle^{n/2+\varepsilon} (I - \Delta_\xi)^N \widehat{f} \|_{L^2}^2 \end{aligned}$$

$$\lesssim \|f\|_s^2 \|\langle x \rangle^{2N} g\|_{n/2+\varepsilon} + \|g\|_s^2 \|\langle x \rangle^{2N} f\|_{n/2+\varepsilon},$$

which concludes the proof. \square

We are now ready to finish the proof of Theorem 1.6.

Proof of Theorem 1.6. First we assume that $\beta > \alpha$. Once more we consider the focusing case and write the solution of the IVP under consideration as

$$u(t, x) = W_\alpha(t)u_0 + \int_0^t W_\alpha(t, \tau) \tau^\beta \nabla_x u \cdot u^{2k} d\tau.$$

We look for the solution given by the fixed point of the map

$$\Phi(u) := W_\alpha(t)u_0 + \int_0^t W_\alpha(t, \tau) \tau^\beta \nabla_x u \cdot u^{2k} d\tau,$$

now defined on

$$X_T^s := \{u : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{C}; \|u\|_{L_t^\infty H_x^s} < \infty, \left(\int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} u|^2 dx dt \right)^{1/2} < \infty, \\ \|\lambda(|x|)^{-1} u\|_{L_t^\infty H_x^{s-2N-3/2}} < \infty\},$$

where

$$\|u\|_{X_T^s}^2 = \|u\|_{L_t^\infty H_x^s}^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} u|^2 dx dt + \|\lambda(|x|)^{-1} u\|_{L_t^\infty H_x^{s-2N-3/2}}^2.$$

We then call $v := \Phi(u)$ the solution of the linear problem

$$\begin{cases} \partial_t v - it^\alpha \Delta_x v - ib(t, x) \cdot \nabla_x v = t^\beta \nabla_x u \cdot u^{2k} \\ v(0, x) = u_0(x), \end{cases}$$

and, as before, we make use of the linear smoothing estimates to prove that Φ is a contraction. In the sequel, for shortness, we will often use the notations $\lambda := \lambda(|x|) = \langle x \rangle^{-2N}$, with $N \geq 1$ (i.e. $\sigma = 2N$), and $\nabla := \nabla_x := \sum_{j=1}^n \partial_{x_j}$. We have

$$\|v\|_{X_T^s}^2 := \|v\|_{L_t^\infty H_x^s}^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} v|^2 dx dt + \|\lambda(|x|)^{-1} v\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\ = I + II + III,$$

(recall $s > n + 4N + 3$), and we estimate the three terms separately. By application of (iii) of Theorem 1.4 (we assume $T \leq 1$ and estimate the exponentials with exponent depending on time directly with a suitable uniform constant) we have

$$I + II = \|v\|_{L_t^\infty H_x^s}^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} v|^2 dx dt \\ \lesssim \|u_0\|_{L_t^\infty H_x^s}^2 + \int_0^T \int_{\mathbb{R}^n} t^{-\alpha} \lambda(|x|)^{-1} |\Lambda^{s-1/2} (t^\beta \nabla_x u \cdot u^{2k})|^2 dx dt$$

$$\begin{aligned}
&= \|u_0\|_{L_t^\infty H_x^s}^2 + \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda(|x|)^{-1} |\Lambda^{s-1/2}(\nabla u \cdot u^{2k})|^2 dx dt \\
&= \|u_0\|_{L_t^\infty H_x^s}^2 + II'.
\end{aligned}$$

Since $s - 1/2 \in 2\mathbb{N}$, then $\Lambda^{s-1/2}$ is a differential operator and, by Leibnitz rule, we have

$$\begin{aligned}
\Lambda^{s-1/2}(\nabla u \cdot u^{2k}) &= (\Lambda^{s-1/2} \nabla u) u^{2k} + \sum_{\substack{|\gamma_1|+|\gamma_2| \leq s-1/2 \\ s/2-1/4 \leq |\gamma_1| < s-1/2, |\gamma_2| \leq s/2-1/4}} C_{\gamma_1, \gamma_2, s} (D^{\gamma_1} \nabla_x u) (D^{\gamma_2} u^{2k}) \\
&+ \sum_{\substack{|\gamma_1|+|\gamma_2| \leq s-1/2 \\ |\gamma_1| < s/2-1/4, |\gamma_2| > s/2-1/4}} C_{\gamma_1, \gamma_2, s} (D^{\gamma_1} \nabla_x u) (D^{\gamma_2} u^{2k}),
\end{aligned}$$

and

$$\begin{aligned}
II' &\leq \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda(|x|)^{-1} |(\Lambda^{s-1/2} \nabla u) u^{2k}|^2 dx dt \\
&+ \sum_{\substack{|\gamma_1|+|\gamma_2| \leq s-1/2 \\ s/2-1/4 \leq |\gamma_1| < s-1/2, |\gamma_2| \leq s/2-1/4}} C_{\gamma_1, \gamma_2, s} \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda(|x|)^{-1} |(D^{\gamma_1} \nabla_x u) (D^{\gamma_2} u^{2k})|^2 dx dt \\
&+ \sum_{\substack{|\gamma_1|+|\gamma_2| \leq s-1/2 \\ |\gamma_1| < s/2-1/4, |\gamma_2| > s/2-1/4}} C_{\gamma_1, \gamma_2, s} \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda(|x|)^{-1} |(D^{\gamma_1} \nabla_x u) (D^{\gamma_2} u^{2k})|^2 dx dt \\
&= II'_a + II'_b + II'_c.
\end{aligned}$$

For II'_a we have

$$\begin{aligned}
II'_a &\leq T^{2\beta-2\alpha} \int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s-1/2} \nabla u|^2 \cdot |\lambda(|x|)^{-1} u^{2k}|^2 dx dt \\
&\leq T^{2\beta-2\alpha} \left(\int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2} u|^2 dx dt \right) \cdot \|\lambda^{-1} u^{2k}\|_{L_t^\infty L_x^\infty}^2 \\
&\leq T^{2\beta-2\alpha} \|u\|_{X_T^s}^2 \|\lambda^{-1} u\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \|u\|_{L_t^\infty H_x^s}^{4k-2} \\
&\leq T^{2\beta-2\alpha} \|u\|_{X_T^s}^{4k+2}.
\end{aligned}$$

For II'_b we have

$$II'_b = \sum_{\substack{|\gamma_1|+|\gamma_2| \leq s-1/2 \\ s/2-1/4 \leq |\gamma_1| < s-1/2, |\gamma_2| \leq s/2-1/4}} C_{\gamma_1, \gamma_2, s} \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda(|x|)^{-1} |(D^{\gamma_1} \nabla_x u) (D^{\gamma_2} u^{2k})|^2 dx dt$$

$$\begin{aligned}
&= \sum_{\substack{|\gamma_1|+|\gamma_2|\leq s-1/2 \\ |\gamma_1|<s-1/2, |\gamma_2|\leq s/2-1/4}} C_{\gamma_1, \gamma_2, s} \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} |(D^{\gamma_1} \nabla_x u)|^2 |\lambda(|x|)^{-1/2} D^{\gamma_2} u^{2k}|^2 dx dt \\
&\leq \sum_{\substack{|\gamma_1|+|\gamma_2|\leq s-1/2 \\ |\gamma_1|<s-1/2, |\gamma_2|\leq s/2-1/4}} C_{\gamma_1, \gamma_2, s} T^{2\beta-\alpha} \left(\int_0^T \int_{\mathbb{R}^n} |(D^{\gamma_1} \nabla_x u)|^2 dx dt \right) \cdot \|\lambda(|x|)^{-1} D^{\gamma_2} u^{2k}\|_{L_t^\infty L_x^\infty}^2 \\
&\leq T^{2\beta-\alpha} \sum_{\substack{|\gamma_1|+|\gamma_2|\leq s-1/2 \\ |\gamma_1|<s-1/2, |\gamma_2|\leq s/2-1/4}} C_{\gamma_1, \gamma_2, s} \|u\|_{L_t^\infty H^{\gamma_1+1}}^2 \cdot \|\lambda(|x|)^{-1} D^{\gamma_2} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2.
\end{aligned}$$

Note that, denoting by Ψ^k a pseudo-differential operator (with constant coefficients) of order k , by using Lemma A.0.2 we have

$$\begin{aligned}
\|\lambda(|x|)^{-1} D^{\gamma_2} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2 &\lesssim \|D^{\gamma_2} \lambda(|x|)^{-1} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2 + \sum_{j=1}^n \|\Psi^{|\gamma_2|-1} x_j \langle x \rangle^{2N-2} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2 \\
&\quad + \sum_{|\alpha+\beta|\leq 2N, |\alpha|>2, |\beta|\leq 2N-2} \|\Psi^{|\gamma_2|-|\alpha|} x^\beta u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2 \\
&\lesssim \|\lambda(|x|)^{-1} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|}}^2 + \sum_{j=1}^n \|x_j \langle x \rangle^{2N-2} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|-1}}^2 \\
&\quad + \sum_{|\alpha+\beta|\leq 2N, |\alpha|>2, |\beta|\leq 2N-2} \|x^\beta u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|-|\alpha|}}^2 \\
&\lesssim \|\lambda(|x|)^{-1} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|}}^2 + \sum_{j=1}^n \left\| \frac{x_j}{\langle x \rangle^2} \langle x \rangle^{2N} u^{2k} \right\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|-1}}^2 \\
&\quad + \sum_{|\alpha+\beta|\leq 2N, |\alpha|>2, |\beta|\leq 2N-2} \left\| \frac{x^\beta}{\langle x \rangle^{2N}} \langle x \rangle^{2N} u^{2k} \right\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|-|\alpha|}}^2 \\
&\lesssim \|\lambda(|x|)^{-1} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|}}^2 \\
&\lesssim \|\lambda(|x|)^{-1} u\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|}}^2 \|u\|_{L_t^\infty H_x^{n/2+\varepsilon+|\gamma_2|}}^{4k-2},
\end{aligned}$$

where we used the H^s -boundedness of $\frac{x_j}{\langle x \rangle^2}$ and $\frac{x^\beta}{\langle x \rangle^{2N}}$ as pseudo-differential operators of order 0, together with Sobolev inequalities.

By using the previous estimate in II'_b and using $\|\cdot\|_{H^{n/2+\varepsilon+|\gamma_2|}} \leq \|\cdot\|_{H^{s-2N-3/2}}$ for ε sufficiently small such that $s/2 + n/2 + \varepsilon - 1/4 \leq s - 2N - 3/2$ (recall $s \geq n + 4N + 3$), we get

$$II'_b \lesssim T^{2\beta-\alpha} \|u\|_{X_T^s}^{4k+2}.$$

For II'_c , repeating the steps in the estimate of II'_b , we have

$$\begin{aligned}
II'_c &= \sum_{\substack{|\gamma_1|+|\gamma_2|\leq s-1/2 \\ |\gamma_1|<s/2-1/4, |\gamma_2|>s/2-1/4}} C_{\gamma_1, \gamma_2, s} \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda(|x|)^{-1} |(D^{\gamma_1} \nabla_x u)(D^{\gamma_2} u^{2k})|^2 dx dt \\
&\leq \sum_{\substack{|\gamma_1|+|\gamma_2|\leq s-1/2 \\ |\gamma_1|<s/2-1/4, |\gamma_2|>s/2-1/4}} C_{\gamma_1, \gamma_2, s} T^{2\beta-\alpha} \left(\int_0^T \int_{\mathbb{R}^n} |(D^{\gamma_2} u^{2k})|^2 dx dt \right) \cdot \|\lambda(|x|)^{-1} D^{\gamma_1} \nabla u\|_{L_t^\infty L_x^\infty}^2 \\
&\leq T^{2\beta-\alpha} \sum_{\substack{|\gamma_1|+|\gamma_2|\leq s-1/2 \\ |\gamma_1|<s/2-1/4, |\gamma_2|>s/2-1/4}} C_{\gamma_1, \gamma_2, s} \|u\|_{L_t^\infty H_x^{2k}}^{4k} \|\lambda(|x|)^{-1} D^{\gamma_1+1} u\|_{L_t^\infty H^{n/2+\varepsilon}}^2 \\
&\lesssim T^{2\beta-\alpha} \|u\|_{X_T^s}^{4k+2}, \\
&\quad \text{by Lemma A.0.2}
\end{aligned}$$

and finally, for $T \leq 1$,

$$I + II \leq CT^{2\beta-2\alpha} \|u\|_{X_T^s}^{4k+2}.$$

To estimate III we use Lemma A.0.3, so we have

$$\begin{aligned}
III &= \|\lambda(|x|)^{-1} v\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\
&\lesssim \|\lambda(|x|)^{-1} W_\alpha(t) u_0\|_{L_t^\infty H_x^{s-2N-3/2}}^2 + \left\| \int_0^t \lambda(|x|)^{-1} W_\alpha(t, \tau) \tau^\beta \nabla u \cdot u^{2k} d\tau \right\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\
&\lesssim_{\text{by Lemma A.0.3 and Minkowski}} C(1 + T^{2N})^2 \|\lambda^{-1} u_0\|_{H_x^{s-3/2}}^2 + \\
&\quad \left(\int_0^T \|\lambda(|x|)^{-1} W_\alpha(t, \tau) \tau^\beta \nabla u \cdot u^{2k}\|_{L_\tau^\infty H_x^{s-2N-3/2}} dt \right)^2 \\
&\lesssim C(1 + T^{2N})^2 \|\lambda^{-1} u_0\|_{H_x^{s-3/2}}^2 + \left(T \sup_{0 \leq \tau \leq t \leq T} \|\lambda(|x|)^{-1} W_\alpha(t, \tau) \tau^\beta \nabla u \cdot u^{2k}\|_{H_x^{s-2N-3/2}} \right)^2 \\
&\stackrel{\text{by Lemma A.0.3}}{\lesssim} C(1 + T^{2N})^2 \|\lambda^{-1} u_0\|_{H_x^{s-3/2}}^2 + \left(CT(1 + T^{2N}) \sup_{0 \leq t \leq T} \|\lambda^{-1} t^\beta \nabla u \cdot u^{2k}\|_{H_x^{s-3/2}} \right)^2 \\
&\lesssim C(1 + T^{2N})^2 \|\lambda^{-1} u_0\|_{H_x^{s-3/2}}^2 + CT^{2(\beta+1)} (1 + T^{2N})^2 \|\lambda^{-1} \nabla u \cdot u^{2k}\|_{L_t^\infty H_x^{s-3/2}}^2 \\
&\stackrel{\text{by Lemma 6.0.1}}{\lesssim} C(1 + T^{2N})^2 \|\lambda^{-1} u_0\|_{H_x^{s-3/2}}^2 + CT^{2(\beta+1)} (1 + T^{2N})^2 (\|\lambda^{-1} \nabla u\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2 \|u^{2k}\|_{L_t^\infty H_x^{s-3/2}}^2 \\
&\quad + \|\lambda^{-1} u^{2k}\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2 \|\nabla u\|_{L_t^\infty H_x^{s-3/2}}^2).
\end{aligned}$$

By Lemma A.0.2 we have (recall $\nabla := \sum_{j=1}^n \partial_{x_j}$)

$$\lambda^{-1} \nabla(u) = \nabla(\lambda^{-1} u) + \sum_{j=1}^n D_{x_j} (x_j \langle x \rangle^{2N-2} u),$$

therefore,

$$\begin{aligned} \|\lambda^{-1}\nabla u\|_{L_t^\infty H_x^{n/2+\varepsilon}} &\lesssim \|\nabla\lambda^{-1}u\|_{L_t^\infty H_x^{n/2+\varepsilon}} + \left\| \sum_{j=1}^n D_{x_j} x_j \langle x \rangle^{2N-2} u \right\|_{L_t^\infty H_x^{n/2+\varepsilon}} \\ &= \|\nabla\lambda^{-1}u\|_{L_t^\infty H_x^{n/2+\varepsilon}} + \left\| \sum_{j=1}^n D_{x_j} \frac{x_j}{\langle x \rangle^2} \langle x \rangle^{2N} u \right\|_{L_t^\infty H_x^{n/2+\varepsilon}} \\ &\lesssim \|\lambda^{-1}u\|_{L_t^\infty H_x^{n/2+\varepsilon+1}} \lesssim \|\lambda^{-1}u\|_{L_t^\infty H_x^{n/2+\varepsilon+1}}, \end{aligned}$$

since $\frac{x_j}{\langle x \rangle^2} \in S^0$ and, for ε sufficiently small, $n/2 + \varepsilon + 1 \leq s - 2N - 3/2$. We then have

$$\begin{aligned} III &\lesssim C(1 + T^{2N})^2 \|\lambda^{-1}u_0\|_{H_x^{s-3/2}}^2 + CT^{2(\beta+1)}(1 + T^{2N})^2 \times \\ &\times \left(\|\lambda^{-1}u\|_{L_t^\infty H_x^{n/2+\varepsilon+1}}^2 \|u\|_{L_t^\infty H_x^{4k}}^{4k} + \|\lambda^{-1}u\|_{L_t^\infty H_x^{n/2+\varepsilon}}^2 \|u\|_{L_t^\infty H_x^{4k-2}}^{4k-2} \|u\|_{L_t^\infty H_x^s}^2 \right) \\ &\leq C(1 + T^{2N})^2 \|\lambda^{-1}u_0\|_{H_x^{s-3/2}}^2 + CT^{2(\beta+1)}(1 + T^{2N})^2 \|u\|_{X_T^s}^{4k+2} \end{aligned}$$

From the previous estimates we get

$$\begin{aligned} \|v\|_{X_T^s}^2 &= I + II + III \\ &\lesssim \|u_0\|_{L_t^\infty H_x^s}^2 + CT^{2\beta-2\alpha} \|u\|_{X_T^s}^{4k+2} + C(1 + T^{2N})^2 \|\lambda^{-1}u_0\|_{H_x^{s-3/2}}^2 + CT^{2(\beta+1)}(1 + T^{2N})^2 \|u\|_{X_T^s}^{4k+2}. \\ &\leq_{T \leq 1} C(\|u_0\|_{L_t^\infty H_x^s}^2 + \|\lambda^{-1}u_0\|_{H_x^{s-3/2}}^2) + C(T^{2\beta-2\alpha} + T^{2(\beta+1)}(1 + T^{2N})^2) \|u\|_{X_T^s}^{4k+2}, \end{aligned}$$

where, recall, $\beta > \alpha$. Finally we have, with new suitable constants that we keep denoting simply C ,

$$\|\Phi(u)\|_{X_T^s} = \|v\|_{X_T^s} \leq C(\|u_0\|_{L_t^\infty H_x^s} + \|\lambda^{-1}u_0\|_{H_x^{s-3/2}}) + C(T^{2\beta-2\alpha} + T^{2(\beta+1)}(1 + T^{2N})^2)^{1/2} \|u\|_{X_T^s}^{2k+1},$$

hence, by choosing $R = C/2(\|u_0\|_{L_t^\infty H_x^s} + \|\lambda^{-1}u_0\|_{H_x^{s-3/2}})$ and T sufficiently small such that $C(T^{2\beta-2\alpha} + T^{2(\beta+1)}(1 + T^{2N})^2)^{1/2} R^{2k} < 1/2$, we get that Φ sends the ball $B_R \subset X_T^s$ into itself.

What is left to prove to conclude the proof is that Φ is a contraction. We then consider, as in the proof of Theorem 1.5, $v := \Phi(u)$ and $w := \Phi(u')$ as the solutions of the linear IVP with the same initial datum u_0 . By application of the linear smoothing estimates on $v - w$ we have

$$\begin{aligned} \|v-w\|_{X_T^s}^2 &\lesssim \|v-w\|_{L_t^\infty H_x^s}^2 + \int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda(|x|) |\Lambda^{s+1/2}(v-w)|^2 dx dt + \|\lambda(|x|)^{-1}(v-w)\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\ &\lesssim \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda^{-1}(|x|) |\Lambda^{s-1/2}(\nabla u \cdot u^{2k} - \nabla u' \cdot u'^{2k})|^2 dx dt + \|\lambda(|x|)^{-1}(v-w)\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\ &\lesssim \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda^{-1}(|x|) |\Lambda^{s-1/2}(\nabla(u^{2k+1} - u'^{2k+1}))|^2 dx dt + \|\lambda(|x|)^{-1}(v-w)\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\ &= IV + V. \end{aligned}$$

For the term IV we proceed like in the estimate of II'_a , II'_b and II'_c (recall $s - 1/2 \in 2\mathbb{N}$, so $\Lambda^{s-1/2}$ is a differential operator on which Leibnitz rule applies) and have

$$\begin{aligned}
IV &= \int_0^T \int_{\mathbb{R}^n} t^{2\beta-\alpha} \lambda^{-1}(|x|) |\Lambda^{s-1/2} \nabla (u^{2k+1} - u'^{2k+1})|^2 dx dt \\
&\leq T^{2\beta-2\alpha} \int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda^{-1}(|x|) |\nabla \Lambda^{s-1/2} (u^{2k+1} - u'^{2k+1})|^2 dx dt \\
&\leq T^{2\beta-2\alpha} \int_0^T \int_{\mathbb{R}^n} t^\alpha \lambda^{-1}(|x|) \left| \nabla \Lambda^{s-1/2} \left((u - u') \sum_{j=0}^{2k} u^{2k-j} u'^j \right) \right|^2 dx dt \\
&\lesssim T^{2\beta-2\alpha} \|u - u'\|_{X_T^s}^2 \|\lambda^{-1} \sum_{j=0}^{2k} u^{2k-j} u'^j\|_{L_t^\infty L_x^\infty}^2 \\
&\quad + T^{2\beta-\alpha} \|u - u'\|_{X_T^s}^2 \|\lambda^{-1} \sum_{j=0}^{2k} u^{2k-j} u'^j\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\
&\quad + T^{2\beta-\alpha} \|\lambda^{-1} (u - u')\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \left\| \sum_{j=0}^{2k} u^{2k-j} u'^j \right\|_{L_t^\infty H_x^s}^2 \\
&\lesssim_{T \leq 1} T^{2\beta-2\alpha} \|u - u'\|_{X_T^s}^2 \sum_{j=0}^{2k} \|\lambda^{-1} u^{2k-j} u'^j\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\
&\quad + T^{2\beta-\alpha} \|u - u'\|_{X_T^s}^2 \sum_{j=0}^{2k} \|u^{2k-j}\|_{L_t^\infty H_x^s}^2 \|u'^j\|_{L_t^\infty H_x^s}^2 \\
&\lesssim T^{2\beta-2\alpha} \|u - u'\|_{X_T^s}^2 \sum_{j=0}^{2k} \|\lambda^{-1} u^{2k-j}\|_{L_t^\infty H_x^{s-2N-3/2}} \|\lambda^{-1} u'^j\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \\
&\quad + T^{2\beta-\alpha} \|u - u'\|_{X_T^s}^2 (\|u\|_{X_T^s} + \|u'\|_{X_T^s})^{4k} \\
&\lesssim T^{2\beta-2\alpha} \|u - u'\|_{X_T^s}^2 \left(\|\lambda^{-1} u^{2k}\|_{L_t^\infty H_x^{s-2N-3/2}}^2 + \sum_{j=1}^{2k} \|u^{2k-j} (\lambda^{-1} u'^j)\|_{L_t^\infty H_x^{s-2N-3/2}}^2 \right) \\
&\quad + T^{2\beta-\alpha} \|u - u'\|_{X_T^s}^2 (\|u\|_{X_T^s} + \|u'\|_{X_T^s})^{4k} \\
&\lesssim T^{2\beta-2\alpha} \|u - u'\|_{X_T^s}^2 \left(\|u\|_{X_T^s}^{4k} + \sum_{j=1}^{2k} \|u\|_{X_T^s}^{4k-2j} \|u'\|_{X_T^s}^{2j} \right) + T^{2\beta-\alpha} \|u - u'\|_{X_T^s}^2 (\|u\|_{X_T^s} + \|u'\|_{X_T^s})^{4k} \\
&\lesssim T^{2\beta-2\alpha} \|u - u'\|_{X_T^s}^2 (\|u\|_{X_T^s} + \|u'\|_{X_T^s})^{4k}.
\end{aligned}$$

where we estimated the sum $\sum_{j=1}^{2k} \|\lambda^{-1} u^{2k-j} u'^j\|_{L_t^\infty H_x^{s-2N-3/2}}^2$ by decoupling each term in the form $u^{2k-j} (\lambda^{-1} u') u'^{j-1}$ and then by using Sobolev inequalities.

For the term V we use the procedure used in the estimate of III above (once again, we make use of Lemma A.0.2 and Lemma A.0.3) and some strategies used for IV , and finally we obtain

$$\|v - w\|_{X_T^s} \leq C(T^{2\beta-2\alpha} + T^{2(\beta+1)}(1 + T^{2N})^2)^{1/2} (\|u\|_{X_T^s} + \|u'\|_{X_T^s})^{2k} \|u - u'\|_{X_T^s}.$$

By taking $u, u' \in B_R$, and eventually by taking the time T smaller, we can conclude that

$$\|\Phi(u) - \Phi(u')\|_{X_T^s} = \|v - w\|_{X_T^s} \leq C \|u - u'\|_{X_T^s},$$

with $C < 1$, so Φ is a contraction. After application of the standard fixed point argument the result follows.

We now assume that $\beta = \alpha$. In this case we are not able to use the time factor $T^{\beta-\alpha}$ in order to obtain a contraction unless we assume that the initial data is small. This is a similar issue to the one faces in [7] and [8], particularly in the part involving the norm λ_3 in page 479. It was resolved by using a version of a mean value theorem in the time variable. More precisely we can write

$$t^\alpha |u|^{2k} \nabla_x u = t^\alpha (|u|^{2k} - |u_0|^{2k}) \nabla_x u + t^\alpha |u_0|^{2k} \nabla_x u = \nabla_x u t^\alpha \int_0^t (|u|^{2k})'(s) ds + t^\alpha |u_0|^{2k} \nabla_x u$$

Then in the course of the proof $\int_0^t (|u|^{2k})'(s) ds$ will bring down t which together with t^α will give t^β , $\beta = \alpha + 1$, while $t^\alpha |u_0|^{2k} \nabla_x u$ will be incorporated in the $b(x, t)$ term. We do not write down all the details since there are essentially contained in the references mentioned earlier. □

7. FINAL REMARKS

We conclude this paper with some remarks about the general case $b \not\equiv 0$. However we recall, once again, that the results proved in what we call *general case* are still true for the particular case $b \equiv 0$.

1. We assumed, in the case $b \not\equiv 0$, that for all $j = 1, \dots, n$, b_j is such that there exist $\sigma > 1$ for which

$$|\operatorname{Im} \partial_x^\gamma b_j(t, x)|, |\operatorname{Re} \partial_x^\gamma b_j(t, x)| \lesssim t^\alpha \langle x \rangle^{-\sigma-|\gamma|}.$$

The first condition on the real part of b_j is natural (see also [1, 12]) since it is needed to be able to apply the Sharp Gårding inequality which is the key point even in the proof of the local well-posedness of the linear problem (since it gives the control on the first order term $b \cdot \nabla_x$). Instead, the second condition, imposed both on the real and on the imaginary part of b_j , is needed in order to have the control $\|r_{s-1}(t, x, D)u\|_0 \leq C t^\alpha \|u\|_{s-1}$ for the error term r_{s-1} in (39) and get the smoothing estimates needed to deal with the nonlinear

problem. Finally the condition on σ , that is $\sigma > 1$, which is imposed in the nondegenerate case as well, is required in order to avoid a loss of derivatives of the solution (see [3]).

2. The nonlinear term $\nabla u \cdot u^{2k}$ is chosen for convenience but it can be generalized. For instance, like in [8], one can consider nonlinearities given by polynomials in u and ∇u and their complex conjugates.

3. Possibly by using the techniques in [8, 12, 3], one can obtain the same smoothing and local well-posedness results for the equation

$$i\partial_t u + t^\alpha \Delta_x u + b(t, x) \cdot \nabla_x u + c(t, x) \cdot \nabla_x \bar{u}$$

assuming on the term $c(t, x)$ suitable conditions (possibly similar to that assumed for $b(t, x)$). In this case the equation should be reduced to a systems, that, after diagonalization, satisfies the desired smoothing properties from which the local well-posedness follows.

4. The results proved in this paper are likely still valid for equations of the form

$$i\partial_t u + g(t)\Delta_x + b(t, x) \cdot \nabla_x,$$

provided that g satisfies suitable properties, as, for example, g having constant sign, vanishing at $t = 0$ and such that $|\partial_x^\gamma b(t, x)| \lesssim |g(t)|\lambda(|x|)$.

APPENDIX A.

In this section we shall recall the statement of some key lemmas we used throughout the paper.

Before giving the statement of the first lemma, which is also the crucial one, that is Doi's lemma (Lemma 2.3 in [3]), we state below the conditions needed to apply this result.

According with the notation used by Doi in [3], we shall denote by (B1), (B2) and (A6) the following conditions:

Let $a^w(t, x, \xi)$ be the Weyl symbol of a pseudo-differential operator $A = A(t, x, D_x)$ (see [5]). We shall say that $a^w := a$ satisfies (B1), (B2) and (A6) if

(B1) $a(t, x, \xi) = ia_2(x, \xi) + a_1(t, x, \xi) + a_0(t, x, \xi)$, where $a_2 \in S_{1,0}^2$ is real-valued and $a_j \in S_{1,0}^j$, for $j = 0, 1$;

(B2) $|a_2(x, \xi)| \geq \delta|\xi|^2$ with $x \in \mathbb{R}^n$, $|\xi|^2 \geq C$, and $\delta, C > 0$;

(A6) *There exists a real-valued function $q \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n)$ such that, with $C_{\alpha\beta}, C_1, C_2 > 0$,*

$$|\partial_\xi^\alpha \partial_x^\beta q(x, \xi)| \leq C_{\alpha\beta} \langle x \rangle \langle \xi \rangle^{-|\alpha|}, \quad x, \xi \in \mathbb{R}^n,$$

$$H_{a_2} q(x, \xi) = \{a_2, q\}(x, \xi) \geq C_1 |\xi| - C_2, \quad x, \xi \in \mathbb{R}^n,$$

where we denoted by $S_{1,0}^j = S_{\rho=1,\delta=0}^j =: S^j$ the standard class of pseudo-differential symbols of order j , and by $\{\cdot, \cdot\}$ the Poisson bracket.

Lemma A.0.1 (Doi [3], Lemma 2.3). *Assume (B1), (B2) and (A6). Let $\lambda(s)$ be a positive non increasing function in $C([0, \infty))$. Then*

- (1) *If $\lambda \in L^1([0, \infty))$ there exists a real-valued symbol $p \in S^0$ and $C > 0$ such that*
- $$(52) \quad H_{a_2} p \geq \lambda(|x|)|\xi| - C, \quad x, \xi \in \mathbb{R}^n;$$
- (2) *If $\int_0^t \lambda(\tau) d\tau \leq C \log(t+1) + C'$, $t \geq 0$, $C, C' > 0$, then there exists a real-valued symbol $p \in S_1^0(\log\langle \xi \rangle)$ such that*
- $$(53) \quad H_{a_2} p \geq \lambda(|x|)|\xi| - C_1 \log\langle \xi \rangle - C_2, \quad x, \xi \in \mathbb{R}^n.$$

Remark A.1. *We remark that, by taking $\lambda'(|x|) = C'\lambda(|x|)$ in Doi's lemma, where C' is any positive constant and λ is as in Lemma A.0.1, then we get that there exists a real-valued symbol $p \in S^0$ and a constant $C > 0$ such that*

$$(54) \quad H_{a_2} p \geq C'\lambda(|x|)|\xi| - C, \quad x, \xi \in \mathbb{R}^n.$$

We conclude the section by giving other two useful lemmas taken from [8].

Lemma A.0.2 (Lemma 6.1.2 of [8]). *Let $p \in S_{0,1}^m$, $N \in \mathbb{N}$. Then*

$$(1 + |x|^2)^N \Psi_p f = \Psi_p [(1 + |x|^2)^N f] + 2N \sum_j \Psi_{i\partial_{\xi_j} p} [x_j (1 + |x|^2)^{N-1} f] \\ + \sum_{|\alpha+\beta| \leq N, |\alpha| \geq 2, |\beta| \leq 2N-2} c_{\alpha\beta} \Psi_{\partial_{\xi}^{\alpha} p} [x^{\beta} f],$$

where Ψ_a stands for the pseudo-differential operator with symbol a .

Lemma A.0.3 (Lemma 6.1.3 of [8]). *Let $N \in \mathbb{N}$ and $s \in \mathbb{R}$. Suppose $\langle x \rangle^{2N} u_0 \in H^{s+2N}$. Then*

$$\sup_{0 \leq t \leq T} \|\langle x \rangle^{2N} W_1(t) u_0\|_{H^s}^2 \leq \sum_{j=0}^{2N} c_j T^j \|\langle x \rangle^{2N-j} u_0\|_{H^{s+j}}^2$$

and

$$\sup_{0 \leq t \leq T} \|\langle x \rangle^{2N} W_1(t) u_0\|_{H^s}^2 \leq c(1 + T^{2N}) \|\langle x \rangle^{2N} u_0\|_{H^{s+2N}}^2,$$

where W_1 denotes the solution operator of (6.1) in [8] with $f = 0$.

We remark that Lemma A.0.3 still works in our case where the operator W_1 will be the solution operator of the homogeneous IVP associated with \mathcal{L}_{α} and that we denoted by $W_{\alpha}(t) := W_{\alpha}(t, 0)$ (for details see the proof in [8] pag.474).

ACKNOWLEDGEMENT

We wish to thank Alberto Parmeggiani for useful discussions and suggestions which helped to improve the present paper.

REFERENCES

- [1] M. CICOGNANI AND M. REISSIG, *Well-Posedness for degenerate Schrödinger equations*, Evolution Equations and Control Theory Volume 3, n. 1, March 2014, 15–33.
- [2] P. CONSTANTIN, J.C. SAUT, *Local smoothing properties of dispersive equations*, J. Amer. Math. Soc. 1(1989) 413–446.
- [3] S. DOI, *Remarks on the Cauchy problem for Schrödinger-type equations*, Comm. Partial Differential Equations 21 (1996), 163–178.
- [4] S. DOI, *On the Cauchy problem for Schrödinger type equations and the regularity of solutions*, J. Math. Kyoto Univ. 34 (1994) 319–328.
- [5] L. HÖRMANDER, *The Analysis of Linear Partial Differential Operators. III. Pseudodifferential Operators*, Grundlehren der Mathematischen Wissenschaften 274. Springer-Verlag, Berlin, 1985. viii+525 pp.
- [6] C. KENIG, G. PONCE AND L. VEGA, *Small solutions to nonlinear Schrödinger equations*, Annales de L'I. H. P. Section C, tome 10, n. 3 (1993), p. 255–288.
- [7] C. KENIG, G. PONCE AND L. VEGA, *The Cauchy problem for quasi-linear Schrödinger equations* Invent. math. 158, (2004), 343–388.
- [8] C. KENIG, G. PONCE, C. ROLVUNG AND L. VEGA, *Variable coefficients Schrödinger flows and ultrahyperbolic operators*, Advances in Mathematics 196 (2005), 373–486.
- [9] J. L. MARZUOLA, J. METCALFE AND D. TATARU *Quasilinear Schrödinger equations III: Large Data and Short Time*. Preprint, arXiv:2001.01014.
- [10] N. LERNER, *Metrics on the phase space and non-selfadjoint pseudo-differential operators*, Pseudo-Differential Operators. Theory and Applications, 3. Birkhäuser Verlag, Basel, 2010. xii+397 pp.
- [11] J. L. MARZUOLA, J. METCALFE AND D. TATARU *Quasilinear Schrödinger equations I: Small data and quadratic interactions*, Advances in Math 231 (2012), 1151–1172.
- [12] S. MIZOHATA, *On the Cauchy Problem*, Notes and Reports in Mathematics in Science and Engineering, Vol. 3, Science Press, Academic Press, New York, 1985.
- [13] C. ROLVUNG, *Non-isotropic Schrödinger equations*, Ph.D. dissertation, University of Chicago, 1998.

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS, GHENT UNIVERSITY, KRIJGSLAAN 281, BUILDING S8, B 9000 GHENT, BELGIUM

E-mail address, Serena Federico: serena.federico@ugent.be

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA, 02139, USA

E-mail address, Gigliola Staffilani: gigliola@math.mit.edu