

FUNCTIONAL GROUP	SPECIES	PERC 5%	PERC 25%	PERC 50%	PERC 75%	PERC 95%
Crustose	<i>Acolium inquinans</i>	-2.80	0.00	1.00	2.00	2.40
Crustose	<i>Acolium karelicum</i>	-4.00	-3.00	-1.50	-0.25	2.20
Crustose	<i>Arthonia mediella</i>	-4.30	-0.75	1.50	5.00	5.65
Crustose	<i>Bacidia globulosa</i>	-2.70	-1.50	0.00	0.00	0.00
Crustose	<i>Biatora chrysantha</i>	-1.50	0.25	1.00	1.00	1.75
Crustose	<i>Biatora efflorescens</i>	0.00	0.00	0.50	1.50	2.70
Crustose	<i>Buellia griseovirens</i>	-4.70	-1.25	1.00	2.00	6.85
Crustose	<i>Buellia schaeferi</i>	-6.00	-2.25	-1.00	3.00	10.20
Crustose	<i>Calicium glaucellum</i>	-4.30	-1.75	-0.50	1.75	8.00
Crustose	<i>Calicium pinicola</i>	-2.70	-1.50	0.00	0.00	0.00
Crustose	<i>Calicium tigillare</i>	-2.70	-1.50	0.00	0.00	0.00
Crustose	<i>Calicium trabinellum</i>	0.00	0.00	0.00	0.50	0.90
Crustose	<i>Calicium viride</i>	-11.20	-5.00	-3.00	1.00	7.20
Crustose	<i>Catillaria nigroclavata</i>	0.00	0.00	0.00	2.00	3.60
Crustose	<i>Chaenotheca chrysocephala</i>	-7.70	-3.00	-2.00	0.50	5.05
Crustose	<i>Chaenotheca ferruginea</i>	-6.80	-3.50	0.00	0.50	1.70
Crustose	<i>Chaenotheca hispidula</i>	0.00	0.00	0.50	1.25	1.85
Crustose	<i>Chaenotheca laevigata</i>	-7.10	-3.75	-2.50	-1.25	0.00
Crustose	<i>Chaenotheca phaeocephala</i>	0.00	0.00	0.00	1.00	1.80
Crustose	<i>Chaenotheca stemonea</i>	0.00	0.00	0.00	2.50	4.50
Crustose	<i>Chaenotheca subroscida</i>	-10.55	-4.25	-0.50	1.75	3.55
Crustose	<i>Chaenotheca trichialis</i>	-8.80	-3.00	0.00	2.00	3.95
Crustose	<i>Chaenothecopsis pusilla</i>	0.40	2.00	2.00	3.00	4.60
Crustose	<i>Chrysothrix candelaris</i>	-3.00	-1.50	0.00	1.50	7.50

FUNCTIONAL GROUP	SPECIES	PERC 5%	PERC 25%	PERC 50%	PERC 75%	PERC 95%
Crustose	<i>Coenogonium pineti</i>	0.00	0.00	0.00	1.00	1.80
Crustose	<i>Frutidella furfuracea</i>	-4.50	-2.50	0.00	0.00	0.00
Crustose	<i>Lecanora argentata</i>	-0.50	1.50	2.00	3.50	8.50
Crustose	<i>Lecanora cadubriae</i>	-3.75	-1.00	2.00	2.25	6.50
Crustose	<i>Lecanora circumborealis</i>	-7.45	-3.75	-2.50	1.00	4.00
Crustose	<i>Lecanora hypoptoides</i>	-4.50	-2.50	0.00	0.50	0.90
Crustose	<i>Lecanora pulicaris</i>	-6.70	-4.00	-2.00	0.00	0.70
Crustose	<i>Lecanora symmicta</i>	-0.90	-0.50	0.00	0.50	0.90
Crustose	<i>Lecanora varia</i>	-14.10	-8.50	-3.00	-2.00	-0.60
Crustose	<i>Lecidea leprarioides</i>	-2.50	-0.75	0.00	0.75	4.00
Crustose	<i>Lepra amara</i>	-2.60	-2.00	0.00	1.00	2.20
Crustose	<i>Lepraria elobata</i>	-4.00	-3.00	0.00	2.50	6.40
Crustose	<i>Lepraria incana</i>	-1.80	-1.00	0.00	0.00	0.00
Crustose	<i>Lepraria sp</i>	0.80	1.00	2.00	4.00	5.40
Crustose	<i>Lopadium disciforme</i>	0.00	0.00	1.00	2.00	2.00
Crustose	<i>Micarea melaena</i>	-3.60	-2.00	0.00	1.00	1.80
Crustose	<i>Micarea prasina agg</i>	0.00	0.00	1.00	3.00	3.00
Crustose	<i>Microcalicium disseminatum</i>	0.00	0.75	2.00	2.25	7.55
Crustose	<i>Mycoblastus affinis</i>	-2.45	-0.75	2.00	3.75	4.30
Crustose	<i>Mycoblastus sanguinarius</i>	-9.75	0.00	0.50	1.00	1.00
Crustose	<i>Ochrolechia alboflavescens</i>	-4.80	-2.00	1.00	2.00	2.20
Crustose	<i>Ochrolechia androgyna</i>	-4.75	-3.25	-2.00	0.00	2.25
Crustose	<i>Ochrolechia arborea</i>	-2.00	-1.25	-0.50	1.25	2.65
Crustose	<i>Ochrolechia microstictoides</i>	-7.20	1.00	3.00	5.00	10.00

FUNCTIONAL GROUP	SPECIES	PERC 5%	PERC 25%	PERC 50%	PERC 75%	PERC 95%
Crustose	<i>Protoparmelia ochrococca</i>	-1.80	-1.00	0.00	4.00	7.20
Crustose	<i>Protoparmelia oleagina</i>	-5.40	-3.00	0.00	0.00	0.00
Crustose	<i>Rinodina capensis</i>	-1.85	-1.25	-0.50	0.00	0.00
Crustose	<i>Schismatomma pericleum</i>	-2.80	-1.00	1.00	3.00	4.40
Crustose	<i>Strangospora deplanata</i>	-1.80	-1.00	0.00	0.00	0.00
Crustose	<i>Toniniopsis separabilis</i>	-6.30	-3.50	0.00	0.00	0.00
Crustose	<i>Violella fucata</i>	-2.40	0.00	0.00	2.00	2.80
Foliose broad-lobed	<i>Cetraria pinastri</i>	-5.00	-1.25	1.00	1.00	3.00
Foliose broad-lobed	<i>Cetraria sepincola</i>	-4.50	-2.50	0.00	0.00	0.00
Foliose broad-lobed	<i>Melanohalea exasperatula</i>	0.00	0.00	0.50	1.00	1.00
Foliose broad-lobed	<i>Nephromopsis chlorophylla</i>	-4.20	-1.00	-1.00	1.00	3.60
Foliose broad-lobed	<i>Nephromopsis laureri</i>	-1.00	-1.00	-1.00	1.00	1.60
Foliose broad-lobed	<i>Parmelia omphalodes</i>	-2.70	-1.50	0.00	0.00	0.00
Foliose broad-lobed	<i>Parmelia saxatilis</i>	-3.10	-2.00	1.00	1.00	3.05
Foliose broad-lobed	<i>Parmelia sulcata</i>	-2.45	-2.00	-1.00	1.00	1.90
Foliose broad-lobed	<i>Platismatia glauca</i>	-8.30	-4.00	-2.00	-0.50	3.00
Foliose narrow-lobed	<i>Hypogymnia austerodes</i>	-1.85	-1.25	-1.00	-0.75	-0.15
Foliose narrow-lobed	<i>Hypogymnia bitteri</i>	-2.50	-1.25	1.00	2.00	2.00
Foliose narrow-lobed	<i>Hypogymnia farinacea</i>	-2.00	-1.00	1.00	1.50	2.50
Foliose narrow-lobed	<i>Hypogymnia physodes</i>	-8.00	-3.00	-1.00	1.00	2.00
Foliose narrow-lobed	<i>Hypogymnia tubulosa</i>	-5.00	-1.00	-1.00	1.00	1.40
Foliose narrow-lobed	<i>Imshaugia aleurites</i>	-6.15	-2.00	-1.00	2.75	4.45
Foliose narrow-lobed	<i>Parmeliopsis ambigua</i>	-3.40	1.00	2.00	3.00	6.00
Foliose narrow-lobed	<i>Parmeliopsis hyperopta</i>	-1.00	1.00	2.00	4.00	6.00

FUNCTIONAL GROUP	SPECIES	PERC 5%	PERC 25%	PERC 50%	PERC 75%	PERC 95%
Fruticose (alectorioid)	<i>Alectoria sarmentosa</i>	-1.70	-0.50	0.00	0.25	0.85
Fruticose (alectorioid)	<i>Bryoria capillaris</i>	-6.00	-4.00	-2.00	-1.00	1.00
Fruticose (alectorioid)	<i>Bryoria fuscescens</i>	-5.00	-3.00	-2.00	-1.00	1.00
Fruticose (alectorioid)	<i>Bryoria implexa</i>	-6.30	-4.25	-3.50	-2.50	-0.35
Fruticose (alectorioid)	<i>Evernia divaricara</i>	-7.50	-2.00	-1.00	-1.00	1.00
Fruticose (alectorioid)	<i>Ramalina thrausta</i>	-0.90	-0.50	0.00	0.00	0.00
Fruticose (alectorioid)	<i>Usnea cavernosa</i>	-6.30	-3.50	0.00	0.00	0.00
Fruticose (alectorioid)	<i>Usnea filipendula</i>	-3.70	-2.50	-2.00	-1.50	-0.30
Fruticose (alectorioid)	<i>Usnea hirta</i>	-9.85	-4.25	-2.00	1.25	3.85
Fruticose (alectorioid)	<i>Usnea sp</i>	-8.00	-5.00	-2.00	-1.00	3.20
Fruticose (non alectorioid)	<i>Cladonia coniocraea</i>	0.00	0.25	1.50	2.75	4.50
Fruticose (non alectorioid)	<i>Cladonia digitata</i>	0.50	2.00	3.00	4.00	7.50
Fruticose (non alectorioid)	<i>Evernia mesomorpha</i>	-2.70	-1.50	0.00	0.00	0.00
Fruticose (non alectorioid)	<i>Letharia vulpina</i>	-1.05	-1.00	1.00	2.00	6.05
Fruticose (non alectorioid)	<i>Pseudevernia furfuracea</i>	-5.00	-3.00	-1.00	1.00	3.40
Fruticose (non alectorioid)	<i>Pseudevernia furfuracea v ceratea</i>	-8.55	-5.00	-2.00	1.00	2.00
Fruticose (non alectorioid)	<i>Ramalina farinacea</i>	-0.85	-0.25	0.00	0.25	0.85
Fruticose (non alectorioid)	<i>Ramalina obtusata</i>	-2.70	-1.50	0.00	0.00	0.00

FUNCTIONAL GROUP	SPECIES	PERC 5%	PERC 25%	PERC 50%	PERC 75%	PERC 95%
Fruticose (non alectorioid)	<i>Ramalina pollinaria</i>	0.00	0.00	0.50	1.00	1.00
Squamulose	<i>Cladonia sp</i>	0.00	0.00	0.50	1.00	1.00
Squamulose	<i>Hypocenomyce scalaris</i>	-1.40	0.50	1.00	3.00	9.20