

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Multidisciplinary approaches for studying rhizobium-legume symbioses

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Dicenzo, G.C., Zamani, M., Checcucci, A., Fondi, M., Griffiths, J.S., Finan, T.M., et al. (2019).
Multidisciplinary approaches for studying rhizobium-legume symbioses. CANADIAN JOURNAL OF
MICROBIOLOGY, 65(1), 1-33 [10.1139/cjm-2018-0377].

Availability:

This version is available at: <https://hdl.handle.net/11585/709547> since: 2019-12-19

Published:

DOI: <http://doi.org/10.1139/cjm-2018-0377>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **TITLE PAGE**

2 **Title:** Multi-disciplinary approaches for studying rhizobium – legume symbioses

3

4 **Running title:** Techniques for studying rhizobia

5

6 **Authors:** George C diCenzo^{1*}, Maryam Zamani², Alice Checcucci¹, Marco Fondi¹, Joel S
7 Griffitts³, Turlough M Finan², Alessio Mengoni¹

8

9 **Affiliations:** ¹ Department of Biology, University of Florence, Sesto Fiorentino, FI, 50019, Italy.

10 ² Department of Biology, McMaster University, Hamilton, ON, L8S 4K1, Canada.

11 ³ Department of Microbiology and Molecular Biology, Brigham Young University, Provo, UT,
12 84602, USA.

13

14 *** Corresponding author:** George diCenzo
15 Department of Biology
16 University of Florence
17 Via Madonna del Piano, 6
18 Sesto Fiorentino, FI, 50019
19 Italy
20 Email: georgecolin.dicenzo@unifi.it

21

ABSTRACT

The rhizobium-legume symbiosis is a major source of fixed nitrogen (ammonia) in the biosphere. The potential for this process to increase agricultural yield while reducing the reliance on nitrogen-based fertilizers has generated interest in understanding and manipulating this process. For decades, rhizobium research has benefited from the use of leading techniques from a very broad set of fields, including population genetics, molecular genetics, genomics, and systems biology. In this review, we summarize many of the research strategies that have been employed in the study of rhizobia and the unique knowledge gained from these diverse tools, with a focus on genome and systems-level approaches. We then describe ongoing synthetic biology approaches aimed at improving existing symbioses or engineering completely new symbiotic interactions. The review concludes with our perspective of the future directions and challenges of the field, with an emphasis on how the application of a multi-disciplinary approach and the development of new methods will be necessary to ensure successful biotechnological manipulation of the symbiosis.

Keywords: Rhizobia, nitrogen fixation, symbiosis, systems biology, multi-omics

40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE OF CONTENTS

TITLE PAGE.....1

ABSTRACT2

TABLE OF CONTENTS3

SYMBIOTIC NITROGEN FIXATION: CHALLENGES AND PROSPECTS4

RHIZOBIAL LIFE-HISTORY.....6

GENETIC MANIPULATIONS AND FUNCTIONAL GENOMICS8

LESSONS FROM COMPARATIVE GENOMICS11

LARGE-SCALE GENOME MANIPULATIONS13

TRANSCRIPTOME STUDIES OF RHIZOBIA16

PROTEOME STUDIES OF RHIZOBIA19

METABOLITE ANALYSES20

IN SILICO METABOLIC MODELLING.....21

SYNTHETIC BIOLOGY APPROACHES TO ENGINEERING THE SYMBIOSIS.....23

CONCLUSIONS AND PERSPECTIVE.....26

ACKNOWLEDGEMENTS.....27

SUPPLEMENTARY FILE DESCRIPTIONS27

REFERENCES28

FIGURE LEGENDS.....91

SYMBIOTIC NITROGEN FIXATION: CHALLENGES AND PROSPECTS

Biological nitrogen fixation (BNF) is an agriculturally and ecologically crucial process that is performed by prokaryotes in two ecological groups: i) BNF performed by free-living cells often in close association with plants, or ii) by rhizobia that fix nitrogen during an endosymbiotic relationship with legumes. The fixation of N₂-gas into ammonia by root-nodule bacteria (rhizobia) is referred to as symbiotic nitrogen fixation (SNF), and it is a more efficient process in terms of supplying nitrogen to the plant. Phylogenetically, most rhizobia are α -proteobacteria, but some rhizobia are β -proteobacteria (Figure 1). The various genetic, biochemical, and evolutionary aspects of the symbiotic interaction have been reviewed over the years (Long 1996, Gage 2004, MacLean et al. 2007, Jones et al. 2007, Gibson et al. 2008, Oldroyd and Downie 2008, Masson-Boivin et al. 2009, Downie 2010, Oldroyd et al. 2011, Udvardi and Poole 2013, Haag et al. 2013, Remigi et al. 2016, Mus et al. 2016, Poole et al. 2018). In brief, the symbiosis is initiated following an exchange of signals between the roots of the plant and free-living rhizobia in root-proximal soil (the rhizosphere). As root nodule tissue develops (Figure 2), the rhizobia enter this specialized tissue through an extracellular infection thread; as these inwardly growing threads reach differentiated nodule cells, the bacteria become surrounded by a plant-derived membrane and taken up into the plant cytosol, where they are known as bacteroids. The nascent bacteroids then undergo major morphological and transcriptional changes, leading to active nitrogen fixation.

It has been estimated that BNF contributes several teragrams of nitrogen (the equivalent of billions of USD of nitrogen-based fertilizer) annually to global agricultural systems (de Vries et al., 2011; Herridge et al., 2008; Ladha et al., 2016). Rhizobial inoculants are inexpensive alternatives to industrial nitrogen fertilizers that can improve crop yields (Figure 2), resulting in greater profits and potentially significant impacts on the livelihood of the community, especially

for the world's poorest farmers (Bloem et al. 2009, Mutuma et al. 2014). BNF also has a substantial positive impact on the environment through reducing the application of nitrogen fertilizers. These fertilizers contribute to elevated atmospheric levels of the potent greenhouse gas nitrous oxide (Park et al. 2012), to toxic nitrate-laden water that contributes to algae blooms and eutrophication (Conley et al., 2009; Randall and Mulla, 2001; Ward, 2009), and to the depletion of fossil fuel as industrial ammonia synthesis by the Haber-Bosch process accounts for 1-2% of total human energy consumption (Erisman et al. 2008). These negative consequences could be ameliorated by relying more on biological solutions for supplying ammonia to agricultural systems.

The benefits of SNF can be maximized in agricultural systems by two general strategies. The first is optimizing the amount of nitrogen fixed by rhizobial bio-inoculants in extant legume symbioses. This will involve not only increasing the rate of nitrogen fixation in the nodule, but also increasing the competitiveness of inoculant strains in the soil and rhizosphere. The second strategy is to engineer completely new nitrogen-fixing symbioses with non-leguminous plants. Many of the world's staple crops, such as the cereals, do not enter into a N_2 -fixing symbiosis with rhizobia (Charpentier and Oldroyd 2010, Oldroyd and Dixon 2014). Engineering synthetic symbioses with these crops therefore presents a monumental and challenging opportunity for exploiting SNF.

Successful implementation of either strategy requires an intricate understanding of both symbiotic partners, in terms of abiotic interactions, responses to complex microbial communities, symbiotic communication, developmental processes, and the metabolism that underpins the nitrogen fixation process. Below, we will explore leading-edge questions in rhizobial biology, and describe how a variety of sound experimental approaches have propelled our understanding of symbiosis biology. The current synthetic biology approaches and future directions and challenges

for engineering BNF for agricultural gains will be described, highlighting the need for multi-level studies of rhizobial biology to provide the necessary data to guide rational improvement of SNF through systems and synthetic biology approaches.

RHIZOBIAL LIFE-HISTORY

Nitrogen-fixing rhizobia have complex life-cycles (Figure 3) (Poole et al. 2018). Rhizobia are found as free-living bacteria in general soil environments and in the rhizosphere. These two environments provide unique nutritional and stress conditions (Hinsinger et al. 2009) that the rhizobia must effectively manage. Rhizobia must also compete against other microbial community members to establish a stable population (Smalla et al. 2001, Li et al. 2016). From the rhizosphere, the rhizobia can infect a compatible legume host, wherein growth conditions vary according to the stage of symbiotic development, from the acidic and oxidative conditions in the pocket of a curled root hair (Santos et al. 2001, Hawkins et al. 2017), to linear growth along the oxidative penetrating infection threads (Santos et al. 2001), to symbiotic differentiation after release into host cells. Each stage of free-living and symbiotic growth requires a unique set of genes and metabolic capabilities. Development of next-generation commercial inoculant strains must account for these and other biological properties, including industrial-scale growth and survival during desiccation (O'Callaghan 2016).

The rhizobial life-cycle does not occur in isolation. It is also necessary to account for social interactions between rhizobia and host plants, and between rhizobia and competing microbes (Checcucci et al. 2017). The rhizobia-host interaction involves a high degree of cooperation between the partners, involving multiple signal exchange events as well as a massive exchange of metabolic resources (Oldroyd and Downie 2008, Oldroyd et al. 2011, Udvardi and Poole 2013).

Yet, hints of conflict and aspects of an evolutionary arms race can also be detected (Sachs et al. 2018). For example, plant immunity plays a key role in the infection process, which must be overcome by the rhizobia to establish an effective symbiosis (Gourion et al. 2015, Berrabah et al. 2018). Additionally, plant-produced nodule-specific cysteine-rich (NCR) peptides (Mergaert et al. 2003, Van de Velde et al. 2010) and corresponding bacterial NCR peptidases (Price et al. 2015) represent a late-stage dialogue in many legumes in which bacteroids improve their fitness at the cost of the host plant by resisting symbiotic enslavement (for reviews, refer to: Kondorosi et al. 2013, Haag et al. 2013, Alunni and Gourion 2016, Pan and Wang 2017). As multiple rhizobial strains may inhabit the same or different nodules on a single plant (Hagen and Hamrick 1996, Checcucci et al. 2016), cheating behaviours have emerged in the rhizobia (Singleton and Tavares 1986, Sachs et al. 2010, Checcucci et al. 2016, Regus et al. 2017), while plants have been observed to limit cheating through sanctions and partner discrimination (Kiers et al. 2003, Kiers and Denison 2008, Heath and Tiffin 2009, Sachs et al. 2010, Quides et al. 2017, Westhoek et al. 2017, Daubech et al. 2017).

Rhizobia also interact with a diverse soil microbial community, where they must effectively compete by antagonism and scavenging of nutrients. This can involve, for example, the production of bacteriocins that have anti-bacterial activity against closely related taxa (Hirsch 1979, Wilson et al. 1998, Oresnik et al. 1999, Venter et al. 2001). The advantage of bacteriocin production is highlighted by the example of trifolitoxin, whose production by *Rhizobium* strains improves rhizosphere colonization and increases competitiveness for nodule occupancy (Triplett and Barta 1987, Robleto et al. 1997, Robleto et al. 1998). The native microbiome may also promote symbiotic development by a rhizobial inoculant; it was observed that soil-isolated *Rhizobium fabae* can promote nodulation of *R. etli* in a quorum sensing-dependent mechanism (Miao et al. 2018).

Rhizobia also encounter numerous bacteriophages in nature that may influence the effectiveness of a rhizobial inoculant. Phages generally are virulent to only some strains of a species, and they can be used to reduce the population density of a poor symbiont to allow the better symbiont to flourish (Evans et al. 1979, Hashem and Angle 1990). However, phages can also reduce the effectiveness of rhizobial inoculants (Mendum et al. 2001).

Nodulation efficiency and efficiency of nitrogen fixation are distinct phenotypes (Bourion et al. 2017). It is clear that understanding the diverse mechanisms underlying all stages of the rhizobial life-history, including competitive fitness in the rhizosphere and each stage of symbiotic development, will be imperative in designing and implementing improved rhizobia-based agricultural strategies. To this end, moving beyond artificial, sterile symbiotic assays in lab conditions will be necessary to generalize results to field conditions.

GENETIC MANIPULATIONS AND FUNCTIONAL GENOMICS

Most of our knowledge about core symbiotic processes comes from classical molecular genetic analyses involving forward- and reverse-genetic techniques, wherein single mutations are associated with defective phenotypes. Tn5-based transposon mutagenesis, coupled with plasmid-based complementation and DNA sequencing, played a foundational role in the discovery of the rhizobial *nod* and *nif/fix* genes (Beringer et al. 1978, Rolfe et al. 1980, Meade et al. 1982, Jagadish and Szalay 1984, Chua et al. 1985, Swanson et al. 1987, Long et al. 1988, Becker et al. 1993a, 1993b, 1993c, Glucksmann et al. 1993). The availability of suitable suicide plasmids and broad host-range plasmids have permitted studies that rely on random transposon mutagenesis, targeted gene disruption, and complementation. Such plasmid-based gene manipulation systems continue to play a prominent role in genetic studies in rhizobia (see for example: VanYperen et al. 2015,

177 Hood et al. 2017, Zamani et al. 2017, Ledermann et al. 2018, Wongdee et al. 2018), and targeted
 178 genetic studies remain essential to complement and validate observations from systems- and
 179 genome-level analyses.

180 High-throughput functional genomic studies have become more common for cataloguing
 181 rhizobial gene functions (see Table 1 for a summary). *Sinorhizobium meliloti* ORFeome
 182 (Schroeder et al. 2005) and fusion (Cowie et al. 2006) libraries have been developed to facilitate
 183 functional and expression analyses in free-living and symbiotic states. *S. meliloti* libraries for use
 184 with *in vivo* expression technology (IVET) have been produced and used to identify genes
 185 functioning in early (Zhang and Cheng 2006) and late (Oke and Long 1999) symbiotic processes.
 186 Similarly, a *Rhizobium leguminosarum* library for use with IVET was prepared and used to identify
 187 genes expressed specifically in the legume rhizosphere (Barr et al. 2008).

188 Several groups have employed signature-tagged mutagenesis (STM) for genome-wide
 189 identification of genes involved in symbiosis or competition for nodule occupancy. STM involves
 190 generating insertion mutants using a library of transposons, with each transposon containing a
 191 distinct sequence tag. Mutants containing different transposons are pooled and passed through a
 192 selective regimen. The relative abundance of each mutant can be measured using microarrays that
 193 detect each sequence tag, and these values can be compared for the library before and after the
 194 selective regimen (Becker and Pobigaylo 2013). A STM library of *S. meliloti* was constructed
 195 (Pobigaylo et al. 2006), and nearly 9,600 insertions were mapped to ~ 3,700 genes throughout the
 196 genome (Serrania et al. 2017). This STM library was employed for the identification of 38 new
 197 genes influencing symbiosis (Pobigaylo et al. 2008). In further work, this library was used to show
 198 that ~ 2% of *S. meliloti* genes contribute to rhizosphere colonization (Salas et al. 2017). Screening
 199 of a *Mesorhizobium loti* STM library on host plants identified 13 Nod⁻ or Fix⁻ mutants as well as

33 mutants impaired in symbiotic competitiveness (Borjigin et al. 2011). Recently, a *Sinorhizobium fredii* STM library consisting of 25,500 mutants was reported, and an initial screen of 10% of these mutants identified four mutants impaired in symbiosis (Wang et al. 2016). STM has also been employed in *Rhizobium leguminosarum* bv. *viciae* (Garcia-Fraile et al. 2015), and a screen for genes relevant to rhizosphere colonization revealed the importance of arabinose and protocatechuate catabolism specifically during rhizosphere growth (Garcia-Fraile et al. 2015).

Although STM facilitates much higher throughput than classical analyses, the more recently developed massively parallel transposon insertion sequencing (known as Tn-seq, or INseq), allows for a further increase in throughput and quantitative assessment of gene functions. In this technique, a saturation insertion-mutagenesis is performed to produce a pool of colonies each containing a single insertion, and next-generation sequencing is employed to identify the position of hundreds-of-thousands of insertions throughout the genome. In this way, mutations contributing to fitness in a given environment can be mapped across an entire genome with a resolution of just several nucleotides (Chao et al. 2016). Example data are shown in Figure 4, illustrating the sensitivity of this method in identifying essential or condition-specific essential genes. Multiple research groups have recently adapted Tn-seq for use in rhizobia. A mariner transposon was used to examine the core genome of *R. leguminosarum* bv. *viciae* (Perry and Yost 2014, Perry et al. 2016), as well as to investigate the effects of carbon source and O₂ levels, a key signal during symbiosis, on the fitness of gene disruptions (Wheatley et al. 2017). Similarly, a Tn5 derived transposon was used to interrogate the core genome of *S. meliloti*, and to inspect how chromosomal gene phenotypes differ when the extra-chromosomal replicons are removed (diCenzo et al. 2018a). Additionally, a mariner transposon was used to investigate the genes contributing to sensitivity of *S. meliloti* to a NCR peptide, which plays a key role during rhizobial

differentiation in the nodule (Arnold et al. 2017). Future use of this technique might prove valuable in the genome-scale elucidation of genes involved at various stages of the rhizobial life-style.

LESSONS FROM COMPARATIVE GENOMICS

Genetic variations present in natural rhizobium strains can be powerfully leveraged to understand SNF through the comparison of global genetic features, a process known as comparative genomics. This approach has become increasingly feasible due to the ever-growing number of rhizobial species and strains with whole-genome sequences available through public repositories. These genome sequences have facilitated many high-quality comparative genomic analyses of rhizobia that have contributed to our understanding of symbiosis and plant growth promotion.

Multiple comparative genomic studies have led to the perhaps surprising observation that there is significant variation between rhizobia in terms of their genomic adaptations to symbiosis (González et al. 2003, Masson-Boivin et al. 2009, Black et al. 2012). Indeed, while there are many genes that appear specific to symbiotic nitrogen-fixing bacteria (Young et al. 2006, Pini et al. 2011), there appears to be no gene unique to all bacteria capable of performing SNF (Amadou et al. 2008). Even the common *nodABC* genes required for Nod factor synthesis are not 100% conserved, as they are absent in certain *Bradyrhizobium* species capable of forming nodules in a Nod factor-independent fashion (Giraud et al. 2007). Other examples of symbiotic genes showing variable presence/absence are accessory *nod* genes involved in determining host specificity, the *fixNOPQ* genes encoding a cytochrome cbb3 oxidase complex, as well as genes encoding polysaccharide biosynthesis and secretion systems (Black et al. 2012, Sugawara et al. 2013, De Meyer et al. 2016). In addition, the operon structure and genomic organization of symbiotic genes

within rhizobial genomes differs among species (González et al. 2003, MacLean et al. 2007, Black et al. 2012, De Meyer et al. 2016). Overall, while many of the core symbiotic genes are highly conserved, the observed genomic variation reflects multiple mechanisms and adaptations for establishing an effective symbiosis, as discussed by Masson-Boivin et al. (2009). Such diversity is bound to complicate our understanding of the minimal set of genes necessary for establishing a symbiosis, and it suggests that strategies to optimize symbiotic interactions may be strain specific.

Despite the differences in the symbiotic gene clusters of distinct rhizobia, comparative genomics can help identify new symbiotically-relevant genes. A comparative analysis of 163 rhizobial genomes identified 184 protein families putatively involved in optimizing the symbiotic process (Seshadri et al. 2015). In another study using a similar comparative genomic approach, 139 genes specific to symbiotic rhizobia were identified (Queiroux et al. 2012). Further characterization of 13 of these genes in *S. meliloti* showed that eight were expressed more highly in the nodule than in free-living cells (Queiroux et al. 2012). Additionally, mutation of the one gene chosen for follow-up study was shown to increase competitiveness for nodule occupancy (Queiroux et al. 2012). These results illustrate how comparative genomics coupled with experimental validation is a powerful approach to identify new candidate genes for improving rhizobial symbiotic efficiency.

Analysis of genetic variation present within a population (population genetics) has uncovered high genetic variability in rhizobia, which is reflected in the large, open pangenomes of many rhizobial species (González et al. 2010, Galardini et al. 2011, Tian et al. 2012, Galardini et al. 2013, Sugawara et al. 2013, Porter et al. 2017). This is demonstrated in Figure 5, which displays the distribution pattern of 17,494 genes found in at least one of the 20 included strains of *S. meliloti*. A clearer understanding of the eco-evolutionary drivers of this genomic variation constitutes an

important research frontier in this field. The influence of soil properties, plant host, and other biotic factors including microbial communities, bacteriophages, plasmids, and other invasive DNA elements all surely play a role in the generation of rhizobial diversity (Bromfield et al. 1987, Harrison et al. 1989, Carelli et al. 2000, Silva et al. 2007, Talebi et al. 2008, Toro et al. 2018). Understanding these relationships could potentially allow for the exploitation of the large genetic diversity of rhizobial species for developing elite rhizobial bioinoculants in precision agriculture (Checcucci et al. 2017).

LARGE-SCALE GENOME MANIPULATIONS

Several *Rhizobium*, *Sinorhizobium*, and *Mesorhizobium* derivatives have been produced that are ‘cured’ of one or more of their large extra-chromosomal replicons (Table 2), revealing the influence of these replicons on numerous free-living phenotypes. These include the ability to catabolize a wide-range of nutrient sources, growth and survival in soil environments (including the rhizosphere), biosynthesis of essential nutrients, production of polysaccharides, and motility (Barbour and Elkan 1989, Hynes et al. 1989, Hynes and McGregor 1990, Baldani et al. 1992, Brom et al. 1992, Moënné-Loccoz and Weaver 1995a, 1995b, Hu et al. 2007, García-de-los-Santos et al. 2008, Stasiak et al. 2014). Similarly, genome reduction studies have highlighted the importance of megaplasms and chromids to the symbiotic process, revealing that key symbiotic genes can be located on megaplasms, that optimal symbiotic performance often requires more than one megaplasmid, and that megaplasms can carry loci required for effective competition for nodule occupancy (Barbour and Elkan 1989, Hynes and McGregor 1990, Baldani et al. 1992, Brom et al. 1992, Hu et al. 2007, Barreto et al. 2012, Stasiak et al. 2014). In contrast, some rhizobial plasmids impair symbiotic development. Loss of a large *Mesorhizobium* plasmid can improve both

nitrogen fixation and nodulation competitiveness (Pankhurst et al. 1986, Hu et al. 2007), while several large *S. meliloti* accessory plasmids may result in legume-host incompatibility (Crook et al. 2012, Price et al. 2015).

A more targeted approach is the production of a library of large-scale deletion mutations that cumulatively remove all of the extra-chromosomal replicon(s) (Table 2). We are aware of this approach being employed with *S. meliloti* and *R. etli* (Landeta et al. 2011, Yurgel et al. 2013a, Milunovic et al. 2014). These libraries have been used to localize and identify novel symbiotic genes, as well as to screen for phenotypes such as carbon metabolic abilities, cytochrome c oxidase activity, soil growth, and osmotolerance (Dominguez-Ferreras et al. 2006, Landeta et al. 2011, Yurgel et al. 2013a, diCenzo et al. 2016a, Zamani et al. 2017).

In addition to genome reduction approaches, genome expansion studies have been used to evaluate the genetic basis of symbiosis. The large symbiotic plasmids of several *Rhizobium* spp. and *S. meliloti* have been transferred to related, non-symbiotic organisms such as *Agrobacterium* spp. and *Ensifer adhaerens* (Hooykaas et al. 1982, Wong et al. 1983, Truchet et al. 1984, Hirsch et al. 1984, Hynes et al. 1986, Finan et al. 1986, Martínez et al. 1987, Novikova and Safronova 1992, Abe et al. 1998, Rogel et al. 2001, Nakatsukasa et al. 2008). The primary conclusion of these studies is that poor, if any, symbiotic abilities are transferred to the recipient strain. This observation is perhaps surprising considering that natural lateral transfer of symbiotic plasmids and islands between related strains results in a gain of symbiotic abilities (Mozo et al. 1988, Laguerre 1992, Sullivan et al. 1995, Sullivan and Ronson 1998, Pérez Carrascal et al. 2016, Nandasena et al. 2007). However, non-fixing outcomes after inter-species heterologous gene transfer may be due to an incomplete complement of symbiotic functions in the transconjugant strains, or due to incompatible alleles of a common gene (diCenzo et al. 2017c). To overcome

315 these limitations, a *S. meliloti* strain lacking the pSymA and pSymB symbiotic replicons was
316 constructed, which represents a 45% reduction of the genome (diCenzo et al. 2014). As expected,
317 this strain is Nod⁻ Fix⁻, and re-introduction of all pSymA and pSymB genes fully restored
318 symbiosis (diCenzo et al. 2016b). The chromosome-only strain, known to be permissive to SNF,
319 is being used for identification of the necessary and sufficient set of symbiotic genes on these
320 replicons.

321 In a particularly notable case, inter-species transfer of a symbiotic plasmid was coupled
322 with laboratory evolution in an ongoing attempt to yield a symbiosis-competent strain. The
323 researchers established *Ralstonia solanacearum* (a plant pathogen) carrying the symbiotic plasmid
324 of the rhizobium species *Cupriavidus taiwanensis* as the test system (Marchetti et al. 2010, Guan
325 et al. 2013, Marchetti et al. 2014, Remigi et al. 2014, Marchetti et al. 2017, Capela et al. 2017).
326 Although initially unable to form nodules, adaptive mutations, such as in the type III secretion
327 system, were identified that allowed for nodulation to occur (Marchetti et al. 2010). Sequential
328 passage of these nodulating strains led to the recovery of derivatives with improved nodulation
329 capabilities, although not yet nitrogen fixation (Marchetti et al. 2017). Nevertheless, this long-term
330 study lends insight into the genomic changes required for the evolution of a new rhizobial species,
331 such as a need to rewire transcriptional networks and to limit induction of the legume immunity
332 response (Marchetti et al. 2014, Capela et al. 2017).

333 In contrast to the inter-species plasmid transfers described above, intra-species transfer of
334 megaplasms in *S. meliloti* yielded strains displaying plant host cultivar-specific improvements
335 in symbiotic effectiveness (Checcucci et al. 2018), suggesting that plasmid transfer is a viable
336 method for ‘breeding’ improved inoculants. Similarly, intra-species transfer of symbiotic plasmids
337 in *R. leguminosarum* resulted in increased nitrogen fixation efficiency (Brewin et al. 1980, DeJong

et al. 1981) and competitiveness for nodule occupancy (Brewin et al. 1983). Remarkably, *Mesorhizobium* strains with the same symbiotic island have widely varying symbiotic phenotypes, from non-effective to highly effective, and varying competitiveness for nodule occupancy (Nandasena et al. 2007, Haskett et al. 2016). Transferring the symbiotic islands from three of these strains to a common *Mesorhizobium* strain confirmed that the differences in symbiotic phenotypes were dependent on differences in the genetic backgrounds of the strains (Haskett et al. 2016). Collectively, these studies underscore the importance of functional interactions between symbiotic replicons/islands and other genomic regions in bringing about efficient symbiotic nitrogen fixation.

TRANSCRIPTOME STUDIES OF RHIZOBIA

Transcriptomic studies, in the form of both microarrays and RNA deep-sequencing (RNA-seq) have been used to investigate transcriptional responses to a broad range of genetic and environmental conditions, to elucidate the regulons of key transcriptional regulators, and to study RNA-based regulation (see Table 3 for a detailed list and description of rhizobium transcriptomic studies). Notable studies of symbiotically-relevant regulators include the FixL/FixJ/FixK system involved in activation of symbiotic genes in response to low oxygen (Bobik et al. 2006, Mesa et al. 2008), and NodD involved in induction of nodulation genes (Capela et al. 2005).

Transcriptomic studies have also probed early and late aspects of symbiotic development (Table 3). This has involved characterizing processes involved in rhizosphere colonization, such as growth in the presence of root exudates (Liu et al. 2017b, Klonowska et al. 2018), or during adaptation to various plant rhizosphere environments (Ramachandran et al. 2011). Several studies have examined the transcriptional consequences of flavonoid detection (Ampe et al. 2003, Barnett

et al. 2004, Capela et al. 2005, Pérez-Montaña et al. 2016a, 2016b), a key initial step in the establishment of the symbiosis (reviewed by (Jiménez-Guerrero et al. 2017). In some species, flavonoid perception results in the induction of many genes (Pérez-Montaña et al. 2016b) and can influence a range of phenotypes including growth and biofilm formation (Spini et al. 2015). Differentiation within the nodule involves changes in the cell cycle, and studies have examined the transcriptional control of the cell cycle of *S. meliloti* and cell cycle-dependent gene expression (De Nisco et al. 2014, Pini et al. 2015). Others have examined the transcriptional effects of plant-produced NCR peptide application to free-living rhizobia, with the results revealing changes in cell cycle behaviour (Penterman et al. 2014) and changes associated with membrane depolarization (Tiricz et al. 2013). Moreover, the transcriptome of *S. meliloti* has been compared between cells growing in oxic and microoxic conditions (Ampe et al. 2003, Becker et al. 2004, Bobik et al. 2006, Pessi et al. 2007), as oxygen limitation is a major trigger for symbiotic gene expression. These studies have confirmed that FixJ is the major regulator of genes differentially expressed under these conditions (Bobik et al. 2006), and that not all genes induced by oxygen limitation in free-living bacteria are expressed in the nodule (Becker et al. 2004).

Transcriptomic experiments have been carried out to evaluate gene regulatory events occurring in root nodules (Table 3). Some of these studies have monitored gene expression of both symbiotic partners (for example, (Barnett et al. 2004, Heath et al. 2012, Roux et al. 2014, Mitsch et al. 2018), or have focused on comparing free-living versus symbiotic bacterial cells (Ampe et al. 2003, Becker et al. 2004, Barnett et al. 2004, Pessi et al. 2007, Karunakaran et al. 2009). A general consensus from these studies is that the bacteria undergo dramatic transcriptional changes in the symbiotic state, with possibly a thousand genes differentially expressed. The majority of differentially-expressed genes are down-regulated in the nodule, consistent with a general

metabolic repression during SNF. Transcriptomics has also been used to decipher the role of specific genes, regulators, or metabolic processes in symbiosis (Table 3), such as the genes *fixJ* (Barnett et al. 2004) and *bacA* (Karunakaran et al. 2010), and carbon import (Mitsch et al. 2018). Others have examined the transcriptome differences of bacteroids displaying distinct morphologies (Lamouche et al. 2018). Additionally, researchers have examined how genetic variations in the plant or rhizobium partner influence the nodule transcriptome (Heath et al. 2012, Burghardt et al. 2017), as well as how the bacteroid transcriptome of the same strain differs depending on the plant hosts (Ampe et al. 2003, Karunakaran et al. 2009, Li et al. 2013). These studies highlight how variation in symbiotic compatibility can result in significant transcriptional alterations.

Researchers have developed imaginative approaches to tease apart the various stages of symbiotic development by either isolating RNA from nodules of different ages (Ampe et al. 2003, Capela et al. 2006) or by using plant or bacterial mutants that are blocked at various stages of the developmental process (Ampe et al. 2003, Tian et al. 2006, Starker et al. 2006, Capela et al. 2006, Maunoury et al. 2010). Others have used laser-capture microdissection to separately capture each zone of the nodule to determine the spatial-expression patterns within nodules (Roux et al. 2014). Such studies have clearly demonstrated a cascade of transcriptional events occurring throughout symbiotic development. This is demonstrated in Figure 6, which displays the unique gene expression patterns of ten genes across four developmental stages. These results highlight the complexity of the differentiation process and the need for detailed studies of each developmental stage in addition to whole-nodule analyses.

PROTEOME STUDIES OF RHIZOBIA

Numerous proteomic studies of rhizobia have been prepared, and a detailed list of these studies are provided in Table 3. Large-scale proteomic studies have been used for purposes such as improving genome annotation, identifying the phosphorylated proteome, characterizing the functions of specific proteins and non-coding RNAs, and evaluating the cellular response to varying environmental conditions (Table 3). Additionally, comparative proteomic studies have been used in the identification of putative symbiotic genes (Gomes et al. 2012b). Several processes of more direct relevance to SNF have also been studied using proteomics (Table 3), such as the proteomic response to flavonoids (Guerreiro et al. 1997, Chen et al. 2000b, Hempel et al. 2009, da Silva Batista and Hungria 2012, Tolín et al. 2013, Arrigoni et al. 2013, Gao et al. 2015, Meneses et al. 2017), and to micro-aerobic or anaerobic conditions (Dainese-Hatt et al. 1999). The secretome of rhizobia, with a particular focus on the proteins exported by type III secretion systems, has also been investigated (Saad et al. 2005, Rodrigues et al. 2007, Hempel et al. 2009, Okazaki et al. 2009).

Nodule-associated rhizobia have also been the subject of proteomic studies (Table 3). Comparison of the proteomes of nitrogen-fixing rhizobia to that of their free-living counterparts have helped identify biochemical processes active during SNF (Djordjevic 2004, Sarma and Emerich 2005, 2006, Djordjevic et al. 2007, Delmotte et al. 2010, Resendis-Antonio et al. 2011, Tatsukami et al. 2013). Similarly, comparing the proteomes of bacteroids isolated from naturally effective and ineffective symbioses implicates pathways contributing to symbiotic efficiency (Cooper et al. 2018). The proteome of the peribacteroid space and peribacteroid membrane (Saalbach et al. 2002) as well as the bacteroid periplasm (Strodtman et al. 2017) have been determined, and the results suggest strong partitioning of proteins among these functionally

distinct compartments (Strodtman et al. 2017). Furthermore, nodule proteomics has been used to examine the proteome during field growth conditions (Delmotte et al. 2010), to compare root-nodules and stem-nodules (Delmotte et al. 2014), and to identify NifA-regulated proteins (Salazar et al. 2010). Studies have compared the proteomes of nodules of different maturities/ages (Delmotte et al. 2014, Nambu et al. 2015, Marx et al. 2016), and recently, proteomes of three zones from the same nodules were independently isolated and evaluated (Ogden et al. 2017). These studies have revealed how the proteome changes during symbiotic development, and they complement and extend conclusions generated using transcriptomics. A more detailed review on the use of proteomics to characterize the symbiosis was recently published (Larrainzar and Wienkoop 2017).

METABOLITE ANALYSES

Metabolomics, the large-scale study of the metabolites of the cell or its environment, has been used to characterize several rhizobial processes. Several studies have examined how the intracellular metabolome of free-living rhizobia differs depending on nutritional or stress conditions, or as a result of genetic manipulations (see Table 3 for a detailed list of relevant studies). Additionally, lipidomics has been used to characterize the membrane-lipid composition and the effects of phosphate or acid stress, as well as to identify proteins involved in membrane-lipid metabolism (Table 3). Studies have also used a fluxomic approach (the study of the rates of reaction fluxes) to characterize central carbon metabolism during growth with either glucose (Fuhrer et al. 2005) or succinate (Terpolilli et al. 2016) as the sole source of carbon.

Metabolomics has been employed to characterize the metabolic shifts occurring during SNF (Table 3). Studies have compared the metabolomes of bacteroids and free-living rhizobia to

identify biochemical processes that differ between these physiologically distinct cell populations (Resendis-Antonio et al. 2012, Vauclare et al. 2013), such as the role of lipogenesis in nitrogen-fixing bacteroids (Terpolilli et al. 2016). Others have compared the metabolite profile of effective and non-effective nodules to identify metabolic processes specifically active during SNF (Ye et al. 2013, Gemperline et al. 2015). As with transcriptomics and proteomics, metabolomics of nodules of different age/maturity has been performed to examine developmental progression (Lardi et al. 2016), and zone-specific metabolite analyses have been performed (Ogden et al. 2017). Metabolomics has further been employed to compare the metabolome of nodules from different plants containing the same rhizobial strain (Lardi et al. 2016). Furthermore, metabolomics has been used to characterize metabolism in root hairs during the early stages of symbiosis (Brechenmacher et al. 2010). Overall, such studies provide insights into the metabolic processes occurring during SNF, and how these processes differ depending on the partners involved.

***IN SILICO* METABOLIC MODELLING**

Genome-scale metabolic network reconstruction refers to the process of building an *in silico* representation of the complete metabolism of the cell (Thiele and Palsson 2010). Reconstructions link metabolism to genetics by listing all reactions expected to be present in the cell together with the corresponding genes encoding the enzymes that catalyze the reaction; they can therefore serve as a valuable knowledge base of the metabolic properties of an organism. The reconstructions can then be used to simulate the metabolism of cells under defined environmental conditions, using mathematical approaches such as flux balance analysis (FBA) (Schellenberger et al. 2011). This approach can predict the metabolic fluxes through the reactions in a given nutritional condition, can simulate how environmental perturbations change the flux distributions,

and can predict how deletion of one or more genes influences growth rate and flux distribution. As an example, Figures 7 and S1 display the predicted distribution of flux through the central carbon metabolic pathways of *S. meliloti* when grown with glucose or succinate as a carbon source.

Metabolic modelling is a relatively new method in the study of rhizobia. The first rhizobial reconstruction was reported in 2007 for *R. etli* (Resendis-Antonio et al. 2007) followed by several refinements (Resendis-Antonio et al. 2011, 2012). In 2012, the first reconstruction for *S. meliloti* was reported (Zhao et al. 2012). These reconstructions were relatively small and were designed solely for the purpose of representing bacteroid metabolism. Simulations of flux distribution and analyses of genes essentiality during SNF generally displayed good correlation with experimental data (Resendis-Antonio et al. 2007, Zhao et al. 2012). Recently, genome-scale metabolic network reconstructions capable of representing both free-living and symbiotic metabolism were reported for *S. meliloti* (diCenzo et al. 2016a) and *Bradyrhizobium diazoefficiens* (Yang et al. 2017c). Simulations with both of these models were consistent with transcriptomic data that indicate a major metabolic shift and general down-regulation, particularly in biosynthesis of cellular components, during symbiosis compared to free-living bacteria (diCenzo et al. 2016a, Yang et al. 2017c). A metabolic reconstruction for the legume *Medicago truncatula* has also been developed (Pfau et al. 2016).

We expect that metabolic reconstruction and constraint-based modelling will serve as a useful tool in the field of SNF, and it will provide an *in silico* method for examining the progression of symbiotic development, which can be experimentally difficult to dissect. Results of the simulations may establish novel hypotheses amenable to experimental testing, and they may contribute to the identification of putative targets for biotechnological manipulation and experimental validation. Several methods have been developed for the integration of

transcriptomics, metabolomics, proteomics, and functional genomics (Tn-seq) data into modelling procedures (Fondi and Liò 2015, diCenzo et al. 2017a), and rhizobial models have been combined with each of these data sets (Resendis-Antonio et al. 2011, 2012, Yang et al. 2017c, diCenzo et al. 2018a). This process results in the creation of high-quality, context-specific representations of cellular metabolism. However, current incomplete knowledge of the metabolic exchange between the plant and bacterium may still limit attempts to accurately represent SNF through a metabolic modelling approach.

SYNTHETIC BIOLOGY APPROACHES TO ENGINEERING THE SYMBIOSIS

Developing improved rhizobium bio-inoculants has been the topic of a few recent reviews and book chapters (Lupwayi et al. 2006, Archana 2010, Dwivedi et al. 2015, Checcucci et al. 2017), and has been of interest to the scientific community for decades (Paau 1991). Yet, despite the many genetic and systems-level studies that shed light on this symbiosis, genetically-engineered rhizobial inoculants are not available. In order for a rhizobial strain to serve as an effective agricultural inoculant, it must be able to i) fix large amounts of nitrogen, and ii) effectively outcompete the indigenous rhizobial population for nodule occupancy. Unfortunately, the native rhizobial community that is often the most competitive for nodule occupancy tends to exhibit low nitrogen fixation rates (see for example: Cardoso et al. 2017a, Chibeba et al. 2017). Indeed, for decades the failure of rhizobia to improve crop yields has often been attributed to the poor competitive abilities of the inoculant and not due to poor nitrogen fixation abilities (Triplett and Sadowsky 1992, Streeter 1994, Ndungu et al. 2018). In strong support of this, it has been observed that the natural transfer of symbiotic islands from inoculant *Mesorhizobium* strains to the native *Mesorhizobium* population often, and rapidly, results in highly competitive strains with poor

nitrogen fixation capabilities (Sullivan et al. 1995, Nandasena et al. 2007). A genomic-based approach comparing the genomes of nodule-isolated rhizobia with the soil rhizobium population, coupled with Tn-seq studies of highly competitive strains in non-sterile environments, may assist in the identification of genes contributing to high competitiveness for nodule occupancy. At the same time, studies such as those described in this review provide a framework for targeted attempts at improving the rate of nitrogen fixation and exchange of nitrogen during the symbiosis. Reports over the last three decades support the feasibility of such an approach (Paa 1991, Ramírez et al. 1999, van Dillewijn et al. 2001, Orikasa et al. 2010).

Many studies have highlighted strong host-strain compatibility properties in SNF (see references below). Consequently, a “personalized” (i.e. cultivar-specific, environment-specific) selection of elite inoculant rhizobia should be considered. However, additional studies are necessary to identify the G (plant genotype) x E (environment) x M (root and soil microbiota) x R (rhizobium) interactions, and to sort out the relevant rhizobial functions of this multi-component interaction (Busby et al. 2017, Finkel et al. 2017, Pahua et al. 2018). For example, *M. truncatula* cultivar-specific NCR peptides can block symbiosis in a bacterial strain-specific manner (Wang et al. 2017, Yang et al. 2017b); many other examples of plant genes restricting symbiosis in a rhizobial strain-dependent fashion exist in the literature (for example: Fan et al. 2017, Yamaya-Ito et al. 2018). Others have shown that relatively small differences between rhizobial genomes can result in significant variations in symbiotic ability (Jozefkiewicz et al. 2017). Additionally, studies have highlighted rhizobial lineage-specific adaptations to symbiosis (Tian et al. 2012, Liu et al. 2017a), and have identified rhizobial genes blocking symbiosis in a plant cultivar-specific manner (Crook et al. 2012, Price et al. 2015, Nguyen et al. 2017). Moreover, studies have demonstrated

that both the environment (Ji et al. 2017) and microbiome (Jozefkiewicz et al. 2017) influence the symbiotic properties of rhizobia.

A major factor limiting our ability to make the most of SNF in agriculture is the inability of many staple crops, such as the cereals, of entering into a SNF relationship. As such, there is significant interest in engineering synthetic symbioses between these plants and rhizobia, and there have been a few recent reviews on this topic (Beatty and Good 2011, Rogers and Oldroyd 2014, Mus et al. 2016, Burén and Rubio 2018). As one can imagine, engineering new inter-kingdom interactions is daunting (Mus et al. 2016), and will require engineering a complex developmental pathway into the plant genome. Fortunately, legume-rhizobium symbioses make use of the common symbiosis signalling pathway that is conserved in plants (including cereals) that participate in mycorrhizal interactions (Oldroyd 2013). In the short term, however, a simpler option would be to improve the beneficial interaction between free-living “associative” nitrogen-fixing bacteria and these crop plants (Geddes et al. 2015), even if these looser symbiotic arrangements transfer nitrogen to the plant less efficiently (Beatty and Good 2011).

An alternative to achieving effective nitrogen fixation in plants is to bypass the bacterial component altogether, and directly engineer plants to perform their own nitrogen fixation. Several reviews and perspective articles discuss this idea and its associated challenges (Beatty and Good 2011, Oldroyd and Dixon 2014, Rogers and Oldroyd 2014, Curatti and Rubio 2014). The main objective is to engineer plants that express a functional nitrogenase inside the mitochondrion or plastid, as these organelles may have the necessary conditions to support this oxygen-sensitive enzyme. Several recent advances have been made in this area that moves the community closer to reaching this goal. Using *Saccharomyces cerevisiae* as a model eukaryote, methods have been developed to express functional NifB protein (Burén et al. 2017a), NifDK tetramers (Burén et al.

2017b), and a functional version of the oxygen-labile NifH protein in mitochondria (López-Torrejón et al. 2016). Other groups were able to individually express each of the *Klebsiella pneumoniae* Nif proteins within the mitochondria of a plant species (Allen et al. 2017), as well as to express active NifH in a plant plastid (Ivleva et al. 2016). Others have re-wired the transcriptional regulatory network of the *nif* genes (Wang et al. 2013), and have identified a minimal nitrogenase gene cluster (Yang et al. 2014). Additionally, it has been shown that the functioning of nitrogenase in free-living bacterial diazotrophs can be supported by the electron-transport chain proteins of plant organelles (Yang et al. 2017a). Each of these studies demonstrates the feasibility of this ambitious research avenue.

CONCLUSIONS AND PERSPECTIVE

Leading-edge techniques from a wide range of disciplines have been applied to the study of rhizobia, facilitating the development of a strong multi-disciplinary understanding of their interactions with leguminous plant hosts (Table 4). Going forward, it will be valuable to integratively employ these methods in orchestrated multi-disciplinary studies. A major challenge of the future will be the development of tools to study the spatio-temporal function of genes in the nodule and during earlier stages of symbiotic development. While transcriptomic, proteomic, and metabolic approaches can help provide insights into the unique developmental stages, genetic and cell biology tools are required to validate conclusions reached from these systems-level analyses. The inducible Cre/*lox* system by (Harrison et al. 2011) is a step in this direction, as are the *lux* bioreporters for the *in vivo* analysis of the spatial and temporal presence of plant metabolites (Pini et al. 2017). Similarly, the *S. meliloti* multiple gene-expression reporter strain for examining the role of plant genes in nodule development helps address this problem (Lang et al. 2018). Moreover,

tools and experimental settings for the fast analysis of host preference in a GxExMxR interaction framework are necessary, such as the recently described select and re-sequence approach (Burghardt et al. 2018). The development of additional tools and their application to studying SNF are a necessity to develop the required knowledge to be able to effectively manipulate this process.

Our understanding of the intricacies of SNF has rapidly advanced over the last few decades, and the emergence of synthetic biology in recent years presents an unprecedented opportunity to engineer improved rhizobia and SNF relationships. However, despite the progress reviewed here, there is still a long road ahead to achieve the ambitious goal of completely replacing nitrogen fertilizer with BNF, and we should continue to balance discovery-oriented and applied research efforts.

ACKNOWLEDGEMENTS

GCD is funded through a post-doctoral fellowship from the Natural Sciences and Engineering Research Council of Canada (NSERC). AC is supported by Fondazione Buzzati-Traverso Fellowship for 2018. JSG is supported by the USDA/NIFA Agricultural and Food Research Initiative (AFRI) award no. 2015-67013-22915. Research in the lab of TMF is supported by NSERC. AM is supported by the University of Florence, “PROGETTI STRATEGICI DI ATENEIO ANNO 2014”.

SUPPLEMENTARY FILE DESCRIPTIONS

File S1. Contains Figure S1, and the methods for generating the data underlying Figures 1, 5, 7, and S1.

Figure S1. *In silico* predicted flux distributions and reaction essentialities. Example data obtained through metabolic modelling. Results from *in silico* metabolic modelling of *S. meliloti* metabolism using the iGD726 metabolic reconstruction are shown (diCenzo et al. 2018a). Growth was simulated using (A) glucose or (B) succinate as the sole carbon source. The pathways represent central carbon metabolism, with the exception of the pentose phosphate pathway (for simplicity). The width of the line represents the amount of flux through the reaction; a doubling in the width corresponds to a ten-fold flux increase. Arrows indicate the direction of flux. Colours indicate if the reaction is essential (red) or non-essential (blue). Genes associated with reactions are based on the information present in iGD726, but they are not necessarily comprehensive. See the Supplementary Materials for details on the modelling procedures used. A version without gene names is provided as Figure 7 in the manuscript.

File S2a. An enlarged, high quality version of the phylogeny provided in Figure 1, including taxonomic names.

File S2b. The phylogeny of Figure 1 in Newick format, including bootstrap values and branch lengths.

File S2c. The annotation file for annotating the phylogeny of File S2b.

REFERENCES

Abe, M., Kawamura, R., Higashi, S., Mori, S., Shibata, M., and Uchiumi, T. 1998. Transfer of the symbiotic plasmid from *Rhizobium leguminosarum* biovar *trifolii* to *Agrobacterium tumefaciens*. J. Gen. Appl. Microbiol. **44**(1): 65–74. doi:10.2323/jgam.44.65.

- Alexandre, A., Laranjo, M., and Oliveira, S. 2014. Global transcriptional response to heat shock of the legume symbiont *Mesorhizobium loti* MAFF303099 comprises extensive gene downregulation. *DNA Res.* **21**(2): 195–206. doi:10.1093/dnares/dst050.
- Allen, R.S., Tilbrook, K., Warden, A.C., Campbell, P.C., Rolland, V., Singh, S.P., and Wood, C.C. 2017. Expression of 16 nitrogenase proteins within the plant mitochondrial matrix. *Front. Plant. Sci* **8**: 287. doi:10.3389/fpls.2017.00287.
- Alunni, B., and Gourion, B. 2016. Terminal bacteroid differentiation in the legume-rhizobium symbiosis: nodule-specific cysteine-rich peptides and beyond. *New Phytol.* **211**(2): 411–417. doi:10.1111/nph.14025.
- Amadou, C., Pascal, G., Mangenot, S., Glew, M., Bontemps, C., Capela, D., Carrère, S., Cruveiller, S., Dossat, C., Lajus, A., Marchetti, M., Poinso, V., Rouy, Z., Servin, B., Saad, M., Schenowitz, C., Barbe, V., Batut, J., Médigue, C., and Masson-Boivin, C. 2008. Genome sequence of the β -rhizobium *Cupriavidus taiwanensis* and comparative genomics of rhizobia. *Genome Res.* **18**(9): 1472–1483. doi:10.1101/gr.076448.108.
- Ampe, F., Kiss, E., Sabourdy, F., and Batut, J. 2003. Transcriptome analysis of *Sinorhizobium meliloti* during symbiosis. *Genome Biol.* **4**(2): R15. doi:10.1186/gb-2003-4-2-r15.
- Arango Pinedo, C., and Gage, D.J. 2009. Plasmids that insert into the rhamnose utilization locus, *rha*: a versatile tool for genetic studies in *Sinorhizobium meliloti*. *J. Mol. Microbiol. Biotechnol.* **17**(4): 201–210. doi:10.1159/000242446.
- Archana, G. 2010. Engineering Nodulation Competitiveness of Rhizobial Bioinoculants in Soils. *In* *Microbes for Legume Improvement*. Springer Vienna, Vienna. pp. 157–194. doi:10.1007/978-3-211-99753-6_8.

- 654 Arnold, M.F.F., Shabab, M., Penterman, J., Boehme, K.L., Griffiths, J.S., and Walker, G.C. 2017.
 655 Genome-wide sensitivity analysis of the microsymbiont *Sinorhizobium meliloti* to
 656 symbiotically important, defensin-like host peptides. MBio **8**(4): e01060–17.
 657 doi:10.1128/mBio.01060-17.
- 658 Arrigoni, G., Tolin, S., Moscatiello, R., Masi, A., Navazio, L., and Squartini, A. 2013. Calcium-
 659 dependent regulation of genes for plant nodulation in *Rhizobium leguminosarum* detected by
 660 iTRAQ quantitative proteomic analysis. J. Proteome Res. **12**(11): 5323–5330.
 661 doi:10.1021/pr400656g.
- 662 Baldani, J.I., Weaver, R.W., Hynes, M.F., and Eardly, B.D. 1992. Utilization of carbon substrates,
 663 electrophoretic enzyme patterns, and symbiotic performance of plasmid-cured clover rhizobia.
 664 Appl. Environ. Microbiol. **58**(7): 2308–2314.
- 665 Barbour, W.M., and Elkan, G.H. 1989. Relationship of the presence and copy number of plasmids
 666 to exopolysaccharide production and symbiotic effectiveness in *Rhizobium fredii* USDA 206.
 667 Appl. Environ. Microbiol. **55**(4): 813–818.
- 668 Barnett, M.J., and Long, S.R. 2015. The *Sinorhizobium meliloti* SyrM regulon: effects on global
 669 gene expression are mediated by *syrA* and *nodD3*. J. Bacteriol. **197**(10): 1792–1806.
 670 doi:10.1128/JB.02626-14.
- 671 Barnett, M.J., Toman, C.J., Fisher, R.F., and Long, S.R. 2004. A dual-genome Symbiosis Chip for
 672 coordinate study of signal exchange and development in a prokaryote–host interaction. Proc.
 673 Natl. Acad. Sci. U.S.A. **101**(47): 16636–16641. doi:10.1073/pnas.0407269101.
- 674 Barr, M., East, A.K., Leonard, M., Mauchline, T.H., and Poole, P.S. 2008. *In vivo* expression
 675 technology (IVET) selection of genes of *Rhizobium leguminosarum* biovar *viciae* A34

- expressed in the rhizosphere. FEMS Microbiol. Lett. **282**(2): 219–227. doi:10.1111/j.1574-6968.2008.01131.x.
- Barra-Bily, L., Fontenelle, C., Jan, G., Flechard, M., Trautwetter, A., Pandey, S.P., Walker, G.C., and Blanco, C. 2010. Proteomic alterations explain phenotypic changes in *Sinorhizobium meliloti* lacking the RNA chaperone Hfq. J. Bacteriol. **192**(6): 1719–1729. doi:10.1128/JB.01429-09.
- Barreto, E.F., Stralio, R., and Baldani, J.I. 2012. Curing of a non-symbiotic plasmid of the *Rhizobium tropici* strain CIAT 899 affected nodule occupancy and competitiveness of the bacteria in symbiosis with common beans. Eur. J. Soil Biol. **50**(C): 91–96. doi:10.1016/j.ejsobi.2011.12.003.
- Barsch, A., Patschkowski, T., and Niehaus, K. 2004. Comprehensive metabolite profiling of *Sinorhizobium meliloti* using gas chromatography-mass spectrometry. Funct. Integr. Genomic **4**(4): 219–230. doi:10.1007/s10142-004-0117-y.
- Basconcillo, L.S., Zaheer, R., Finan, T.M., and McCarry, B.E. 2009a. Cyclopropane fatty acyl synthase in *Sinorhizobium meliloti*. Microbiology **155**(Pt 2): 373–385. doi:10.1099/mic.0.022608-0.
- Basconcillo, L.S., Zaheer, R., Finan, T.M., and McCarry, B.E. 2009b. A shotgun lipidomics approach in *Sinorhizobium meliloti* as a tool in functional genomics. J. Lipid Res. **50**(6): 1120–1132. doi:10.1194/jlr.M800443-JLR200.
- Basconcillo, L.S., Zaheer, R., Finan, T.M., and McCarry, B.E. 2009c. A shotgun lipidomics study of a putative lysophosphatidic acid acyl transferase (PlsC) in *Sinorhizobium meliloti*. J. Chromatogr. B. **877**(26): 2873–2882. doi:10.1016/j.jchromb.2009.05.014.

- 698 Batista, J.S.D.S., Torres, A.R., and Hungria, M. 2010. Towards a two-dimensional proteomic
699 reference map of *Bradyrhizobium japonicum* CPAC 15: spotlighting "hypothetical proteins".
700 Proteomics **10**(17): 3176–3189. doi:10.1002/pmic.201000092.
- 701 Beatty, P.H., and Good, A.G. 2011. Future prospects for cereals that fix nitrogen. Science
702 **333**(6041): 416–417. doi:10.1126/science.1209467.
- 703 Becker, A., Kleickmann, A., Arnold, W., and Pühler, A. 1993. Analysis of the *Rhizobium meliloti*
704 *exoH/exoK/exoL* fragment: ExoK shows homology to excreted endo- β -1,3-1,4-glucanases and
705 ExoH resembles membrane proteins. Mol. Gen. Genet. **238**(1-2): 145–154.
- 706 Becker, A., Kleickmann, A., Keller, M., Arnold, W., and Pühler, A. 1993. Identification and
707 analysis of the *Rhizobium meliloti* *exoAMONP* genes involved in exopolysaccharide
708 biosynthesis and mapping of promoters located on the *exoHKLAMONP* fragment. Mol. Gen.
709 Genet. **241**(3-4): 367–379.
- 710 Becker, A., Kleickmann, A., Küster, H., Keller, M., Arnold, W., and Pühler, A. 1993. Analysis of
711 the *Rhizobium meliloti* genes *exoU*, *exoV*, *exoW*, *exoT*, and *exoI* involved in exopolysaccharide
712 biosynthesis and nodule invasion: *exoU* and *exoW* probably encode glucosyltransferases. Mol.
713 Plant Microbe Interact. **6**(6): 735–744. doi:10.1094/MPMI-6-735.
- 714 Becker, A., and Pobigaylo, N. 2013. Construction of signature-tagged mutant libraries and its
715 application to plant-symbiotic bacteria. In Molecular Microbial Ecology of the Rhizosphere.
716 Edited by F.J. de Bruijn. John Wiley & Sons, Inc., Hoboken, NJ, USA. pp. 921–928.
717 doi:10.1002/9781118297674.ch86.
- 718 Becker, A., Bergès, H., Krol, E., Bruand, C., Rüberg, S., Capela, D., Lauber, E., Meilhoc, E.,
719 Ampe, F., de Bruijn, F.J., Fourment, J., Francez-Charlot, A., Kahn, D., Küster, H., Liebe, C.,
720 Pühler, A., Weidner, S., and Batut, J. 2004. Global changes in gene expression in

- 721 *Sinorhizobium meliloti* 1021 under microoxic and symbiotic conditions. Mol. Plant Microbe
722 Interact. **17**(3): 292–303. doi:10.1094/MPMI.2004.17.3.292.
- 723 Beringer, J.E., Beynon, J.L., Buchanan-Wollaston, A.V., and Johnston, A.W.B. 1978. Transfer of
724 the drug-resistance transposon Tn5 to *Rhizobium*. Nature **276**: 633–634.
- 725 Berrabah, F., Balliau, T., Aït-Salem, E.H., George, J., Zivy, M., Ratet, P., and Gourion, B. 2018.
726 Control of the ethylene signaling pathway prevents plant defenses during intracellular
727 accommodation of the rhizobia. New Phytol. **219**(1): 310–323. doi:10.1111/nph.15142.
- 728 Bhat, S.V., Booth, S.C., McGrath, S.G.K., and Dahms, T.E.S. 2014. *Rhizobium leguminosarum*
729 bv. *viciae* 3841 adapts to 2,4-dichlorophenoxyacetic acid with “auxin-like” morphological
730 changes, cell envelope remodeling and upregulation of central metabolic pathways. PLOS One
731 **10**(4): e0123813. doi:10.1371/journal.pone.0123813.
- 732 Black, M., Moolhuijzen, P., Chapman, B., Barrero, R., Howieson, J., Hungria, M., and Bellgard,
733 M. 2012. The genetics of symbiotic nitrogen fixation: comparative genomics of 14 rhizobia
734 strains by resolution of protein clusters. Genes **3**(1): 138–166. doi:10.3390/genes3010138.
- 735 Bloem, J.F., Trytsman, G., and Smith, H.J. 2009. Biological nitrogen fixation in resource-poor
736 agriculture in South Africa. Symbiosis **48**(1-3): 18–24. doi:10.1007/BF03179981.
- 737 Bobik, C., Meilhoc, E., and Batut, J. 2006. FixJ: a major regulator of the oxygen limitation
738 response and late symbiotic functions of *Sinorhizobium meliloti*. J. Bacteriol. **188**(13): 4890–
739 4902. doi:10.1128/JB.00251-06.
- 740 Borjigin, N., Furukawa, K., Shimoda, Y., Tabata, S., Sato, S., Eda, S., Minamisawa, K., and
741 Mitsui, H. 2011. Identification of *Mesorhizobium loti* genes relevant to symbiosis by using
742 signature-tagged mutants. Microbes Environ. **26**(2): 165–171. doi:10.1264/jsme2.ME10213.

- 743 Bourdès, A., Rudder, S., East, A.K., and Poole, P.S. 2012. Mining the *Sinorhizobium meliloti*
 744 transportome to develop FRET biosensors for sugars, dicarboxylates and cyclic polyols. PLOS
 745 One **7**(9): e43578. doi:10.1371/journal.pone.0043578.
- 746 Bourion, V., Heulin-Gotty, K., Aubert, V., Tisseyre, P., Chabert-Martinello, M., Pervent, M.,
 747 Delaitre, C., Vile, D., Siol, M., Duc, G., Brunel, B., Burstin, J., and Lepetit, M. 2017. Co-
 748 inoculation of a pea core-collection with diverse rhizobial strains shows competitiveness for
 749 nodulation and efficiency of nitrogen fixation are distinct traits in the interaction. Front. Plant
 750 Sci **8**: 2249. doi:10.3389/fpls.2017.02249.
- 751 Brechenmacher, L., Lei, Z., Libault, M., Findley, S., Sugawara, M., Sadowsky, M.J., Sumner,
 752 L.W., and Stacey, G. 2010. Soybean metabolites regulated in root hairs in response to the
 753 symbiotic bacterium *Bradyrhizobium japonicum*. Plant Physiol. **153**(4): 1808–1822.
 754 doi:10.1104/pp.110.157800.
- 755 Brewin, N.J., DeJong, T.M., Philips, D.A., and Johnston, A.W.B. 1980. Co-transfer of
 756 determinants for hydrogenase activity and nodulation ability in *Rhizobium leguminosarum*.
 757 Nature **288**: 77-79
- 758 Brewin, N.J., Wood, E.A., and Young, J.P.W. 1983. Contribution of the symbiotic plasmid to the
 759 competitiveness of *Rhizobium leguminosarum*. J. Gen. Microbiol. **129**: 2973-2977.
 760 doi:10.1099/00221287-129-10-2973.
- 761 Bringhurst, R.M., Cardon, Z.G., and Gage, D.J. 2001. Galactosides in the rhizosphere: Utilization
 762 by *Sinorhizobium meliloti* and development of a biosensor. Proc. Natl. Acad. Sci. U.S.A.
 763 **98**(8): 4540–4548. doi:10.1073/pnas.071375898.
- 764 Brom, S., García-de los Santos, A., Cervantes, L., Palacios, R., and Romero, D. 2000. In
 765 *Rhizobium etli* symbiotic plasmid transfer, nodulation competitiveness and cellular growth

- 766 require interaction among different replicons. *Plasmid* **44**(1): 34–43.
 767 doi:10.1006/plas.2000.1469.
- 768 Brom, S., García-de los Santos, A., Stepkowsky, T., Flores, M., Davila, G., Romero, D., and
 769 Palacios, R. 1992. Different plasmids of *Rhizobium leguminosarum* bv. *phaseoli* are required
 770 for optimal symbiotic performance. *J. Bacteriol.* **174**(16): 5183–5189.
 771 doi:10.1128/jb.174.16.5183-5189.1992.
- 772 Bromfield, E.S.P., Thurman, N.P., Whitwill, S.T., and Barran, L.R. 1987. Plasmids and symbiotic
 773 effectiveness of representative phage types from two indigenous populations of *Rhizobium*
 774 *meliloti*. *Microbiology* **133**(12): 3457–3466. doi:10.1099/00221287-133-12-3457.
- 775 Burén, S., Jiang, X., López-Torrejón, G., Echavarri-Erasun, C., and Rubio, L.M. 2017a.
 776 Purification and *in vitro* activity of mitochondria targeted nitrogenase cofactor maturase NifB.
 777 *Front. Plant Sci.* **8**: 1567. doi:10.3389/fpls.2017.01567.
- 778 Burén, S., Young, E.M., Sweeny, E.A., López-Torrejón, G., Veldhuizen, M., Voigt, C.A., and
 779 Rubio, L.M. 2017b. Formation of nitrogenase NifDK tetramers in the mitochondria of
 780 *Saccharomyces cerevisiae*. *ACS Synth. Biol.* **6**(6): 1043–1055.
 781 doi:10.1021/acssynbio.6b00371.
- 782 Burghardt, L.T., Epstein, B., Guhlin, J., Nelson, M.S., Taylor, M.R., Young, N.D., Sadowsky,
 783 M.J., and Tiffin, P. 2018. Select and resequence reveals relative fitness of bacteria in symbiotic
 784 and free-living environments. *Proc. Natl. Acad. Sci. U.S.A.* **115**(10): 2425–2430.
 785 doi:10.1073/pnas.1714246115.
- 786 Burghardt, L.T., Guhlin, J., Chun, C.L., Liu, J., Sadowsky, M.J., Stupar, R.M., Young, N.D., and
 787 Tiffin, P. 2017. Transcriptomic basis of genome by genome variation in a legume-rhizobia
 788 mutualism. *Mol. Ecol.* **25**(2): 4946. doi:10.1111/mec.14285.

789 Busby, P.E., Soman, C., Wagner, M.R., Friesen, M.L., Kremer, J., Bennett, A., Morsy, M., Eisen,
 790 J.A., Leach, J.E., and Dangl, J.L. 2017. Research priorities for harnessing plant microbiomes
 791 in sustainable agriculture. *PLOS Biol.* **15**(3): e2001793. doi:10.1371/journal.pbio.2001793.
 792 Calatrava-Morales, N., Nogales, J., Amezttoy, K., van Steenberg, B., and Soto, M.J. 2017. The
 793 NtrY/NtrX system of *Sinorhizobium meliloti* GR4 regulates motility, EPS I production, and
 794 nitrogen metabolism but Is dispensable for symbiotic nitrogen fixation. *Mol. Plant Microbe*
 795 *Interact* **30**(7): 566–577. *Mol. Plant Microbe Interact.* doi:10.1094/MPMI-01-17-0021-R.
 796 Cantero, L., Palacios, J.M., Ruiz-Argüeso, T., and Imperial, J. 2006. Proteomic analysis of quorum
 797 sensing in *Rhizobium leguminosarum* biovar *viciae* UPM791. *Proteomics* **6**(S1): S97–106.
 798 doi:10.1002/pmic.200500312.
 799 Capela, D., Carrère, S., and Batut, J. 2005. Transcriptome-based identification of the
 800 *Sinorhizobium meliloti* NodD1 regulon. *Appl. Environ. Microbiol.* **71**(8): 4910–4913.
 801 doi:10.1128/AEM.71.8.4910-4913.2005.
 802 Capela, D., Filipe, C., Bobik, C., Batut, J., and Bruand, C. 2006. *Sinorhizobium meliloti*
 803 differentiation during symbiosis with alfalfa: a transcriptomic dissection. *Mol. Plant Microbe*
 804 *Interact.* **19**(4): 363–372. doi:10.1094/MPMI-19-0363.
 805 Capela, D., Marchetti, M., Clerissi, C., Perrier, A., Guetta, D., Gris, C., Valls, M., Jauneau, A.,
 806 Cruveiller, S., Rocha, E.P.C., and Masson-Boivin, C. 2017. Recruitment of a lineage-specific
 807 virulence regulatory pathway promotes intracellular infection by a plant pathogen
 808 experimentally evolved into a legume symbiont. *Mol. Biol. Evol.* **34**(10): 2503–2521.
 809 doi:10.1093/molbev/msx165.

- 810 Cardoso, A.A., Andraus, M. de P., Borba, T.C. de O., Martin-Didonet, C.C.G., and Ferreira, E.P.
 811 de B. 2017a. Characterization of rhizobia isolates obtained from nodules of wild genotypes of
 812 common bean. *Braz. J. Microbiol.* **48**(1): 43–50. doi:10.1016/j.bjm.2016.09.002.
- 813 Cardoso, P., Santos, M., Freitas, R., Rocha, S.M., and Figueira, E. 2017b. Response of *Rhizobium*
 814 to Cd exposure: A volatile perspective. *Environ. Pollut.* **231**(Pt 1): 802–811.
 815 doi:10.1016/j.envpol.2017.08.067.
- 816 Carelli, M., Gnocchi, S., Fancelli, S., Mengoni, A., Paffetti, D., Scotti, C., and Bazzicalupo, M.
 817 2000. Genetic diversity and dynamics of *Sinorhizobium meliloti* populations nodulating
 818 different alfalfa cultivars in Italian soils. *Appl. Environ. Microbiol.* **66**(11): 4785–4789.
 819 doi:10.1128/AEM.66.11.4785-4789.2000.
- 820 Cervantes, L., Bustos, P., Girard, L., Santamaría, R.I., Dávila, G., Vinuesa, P., Romero, D., and
 821 Brom, S. 2011. The conjugative plasmid of a bean-nodulating *Sinorhizobium fredii* strain is
 822 assembled from sequences of two *Rhizobium* plasmids and the chromosome of a
 823 *Sinorhizobium* strain. *BMC Microbiol.* **11**(1): 149. doi:10.1186/1471-2180-11-149.
- 824 Chain, P.S., Hernandez-Lucas, I., Golding, B., Finan, T.M. 2000. *oriT*-directed cloning of defined
 825 large regions from bacterial genomes: identification of the *Sinorhizobium meliloti* pExo
 826 megaplasmid replicator region. *J. Bacteriol.* **182**(19): 5486-5494.
 827 doi:10.1128/JB.182.19.5486-5494.2000.
- 828 Charles, T.C., and Finan, T.M. 1991. Analysis of a 1600-kilobase *Rhizobium meliloti* megaplasmid
 829 using defined deletions generated *in vivo*. *Genetics* **127**(1): 5–20.
- 830 Chao, M.C., Abel, S., Davis, B.M., and Waldor, M.K. 2016. The design and analysis of transposon
 831 insertion sequencing experiments. *Nat. Rev. Microbiol.* **14**(2): 119–128.
 832 doi:10.1038/nrmicro.2015.7.

- 833 Chao, T.-C., Buhrmester, J., Hansmeier, N., Pühler, A., and Weidner, S. 2005. Role of the
834 regulatory gene *rirA* in the transcriptional response of *Sinorhizobium meliloti* to iron
835 limitation. Appl. Environ. Microbiol. **71**(10): 5969–5982. doi:10.1128/AEM.71.10.5969-
836 5982.2005.
- 837 Charpentier, M., and Oldroyd, G. 2010. How close are we to nitrogen-fixing cereals? Curr. Opin.
838 Plant Biol. **13**(5): 556–564. doi:10.1016/j.pbi.2010.08.003.
- 839 Chaudri, A.M., Lawlor, K., Preston, S., Paton, G.I., Killham, K., and McGrath, S.P. 2000. Response
840 of a *Rhizobium*-based luminescence biosensor to Zn and Cu in soil solutions from sewage
841 sludge treated soils. Soil Biol Biochem **32**(3): 383–388. doi:10.1016/S0038-0717(99)00166-
842 2.
- 843 Checcucci, A., Azzarello, E., Bazzicalupo, M., Galardini, M., Lagomarsino, A., Mancuso, S.,
844 Marti, L., Marzano, M.C., Mocali, S., Squartini, A., Zanardo, M., and Mengoni, A. 2016.
845 Mixed nodule infection in *Sinorhizobium meliloti*-*Medicago sativa* symbiosis suggest the
846 presence of cheating behavior. Front. Plant Sci. **7**: 835. doi:10.3389/fpls.2016.00835.
- 847 Checcucci, A., diCenzo, G.C., Bazzicalupo, M., and Mengoni, A. 2017. Trade, diplomacy, and
848 warfare: the quest for elite rhizobia inoculant strains. Front. Microbiol. **8**: 2207.
849 doi:10.3389/fmicb.2017.02207.
- 850 Checcucci, A., diCenzo, G.C., Ghini, V., Bazzicalupo, M., Beker, A., Decorosi, F., Döhlemann,
851 J., Fagorzi, C., Finan, T.M., Fondi, M., Luchinat, C., Turano, P., Vignolini, T., Viti, C., and
852 Mengoni, A. 2018. Creation and multi-omics characterization of a genomically hybrid strain
853 in the nitrogen-fixing symbiotic bacterium *Sinorhizobium meliloti*. bioRxiv **296483**.
854 doi:10.1101/296483.

- 855 Chen, H., Gao, K., Kondorosi, E., Kondorosi, A., and Rolfe, B.G. 2005. Functional genomic
856 analysis of global regulator NolR in *Sinorhizobium meliloti*. *Mol. Plant Microbe Interact.*
857 **18**(12): 1340–1352. doi:10.1094/MPMI-18-1340.
- 858 Chen, H., Higgins, J., Kondorosi, E., Kondorosi, A., Djordjevic, M.A., Weinman, J.J., and Rolfe,
859 B.G. 2000a. Identification of *nolR*-regulated proteins in *Sinorhizobium meliloti* using
860 proteome analysis. *Electrophoresis* **21**(17): 3823–3832. doi:10.1002/1522-
861 2683(200011)21:17<3823::AID-ELPS3823>3.0.CO;2-K.
- 862 Chen, H., Higgins, J., Oresnik, I.J., Hynes, M.F., Natera, S., Djordjevic, M.A., Weinman, J.J., and
863 Rolfe, B.G. 2000b. Proteome analysis demonstrates complex replicon and luteolin interactions
864 in pSyma-cured derivatives of *Sinorhizobium meliloti* strain 2011. *Electrophoresis* **21**(17):
865 3833–3842. doi:10.1002/1522-2683(200011)21:17<3833::AID-ELPS3833>3.0.CO;2-I.
- 866 Chen, H., Teplitski, M., Robinson, J.B., Rolfe, B.G., and Bauer, W.D. 2003. Proteomic analysis
867 of wild-type *Sinorhizobium meliloti* responses to N-acyl homoserine lactone quorum-sensing
868 signals and the transition to stationary phase. *J. Bacteriol.* **185**(17): 5029–5036.
869 doi:10.1128/JB.185.17.5029-5036.2003.
- 870 Chibeba, A.M., Kyei-Boahen, S., Guimarães, M. de F., Nogueira, M.A., and Hungria, M. 2017.
871 Isolation, characterization and selection of indigenous *Bradyrhizobium* strains with
872 outstanding symbiotic performance to increase soybean yields in Mozambique. *Agric.*
873 *Ecosyst. Environ.* **246**: 291–305. doi:10.1016/j.agee.2017.06.017.
- 874 Chua, K.Y., Pankhurst, C.E., Macdonald, P.E., Hopcroft, D.H., Jarvis, B.D., and Scott, D.B. 1985.
875 Isolation and characterization of transposon Tn5-induced symbiotic mutants of *Rhizobium loti*.
876 *J. Bacteriol.* **162**(1): 335–343.

- 877 Cooper, B., Campbell, K.B., Beard, H.S., Garrett, W.M., Mowery, J., Bauchan, G.R., and Elia, P.
 878 2018. A proteomic network for symbiotic nitrogen fixation efficiency in *Bradyrhizobium*
 879 *elkanii*. Mol. Plant Microbe Interact. **31**(3): 334–343. doi:10.1094/MPMI-10-17-0243-R.
- 880 Cowie, A., Cheng, J., Sibley, C.D., Fong, Y., Zaheer, R., Patten, C.L., Morton, R.M., Golding,
 881 G.B., and Finan, T.M. 2006. An integrated approach to functional genomics: construction of
 882 a novel reporter gene fusion library for *Sinorhizobium meliloti*. Appl. Environ. Microbiol.
 883 **72**(11): 7156–7167. doi:10.1128/AEM.01397-06.
- 884 Crook, M.B., Lindsay, D.P., Biggs, M.B., Bentley, J.S., Price, J.C., Clement, S.C., Clement, M.J.,
 885 Long, S.R., and Griffiths, J.S. 2012. Rhizobial plasmids that cause impaired symbiotic nitrogen
 886 fixation and enhanced host invasion. Mol. Plant Microbe Interact. **25**(8): 1026–1033.
 887 doi:10.1094/MPMI-02-12-0052-R.
- 888 Curatti, L., and Rubio, L.M. 2014. Challenges to develop nitrogen-fixing cereals by direct *nif*-gene
 889 transfer. Plant Sci. **225**: 130–137. doi:10.1016/j.plantsci.2014.06.003.
- 890 Cytryn, E.J., Sangurdekar, D.P., Streeter, J.G., Franck, W.L., Chang, W.-S., Stacey, G., Emerich,
 891 D.W., Joshi, T., Xu, D., and Sadowsky, M.J. 2007. Transcriptional and physiological
 892 responses of *Bradyrhizobium japonicum* to desiccation-induced stress. J. Bacteriol. **189**(19):
 893 6751–6762. doi:10.1128/JB.00533-07.
- 894 D'Alessio, M., Nordeste, R., Doxey, A.C., and Charles, T.C. 2017. Transcriptome analysis of
 895 polyhydroxybutyrate cycle mutants reveals discrete loci connecting nitrogen utilization and
 896 carbon storage in *Sinorhizobium meliloti*. mSystems **2**(5): e00035–17.
 897 doi:10.1128/mSystems.00035-17.

- 898 da Silva Batista, J.S., and Hungria, M. 2012. Proteomics reveals differential expression of proteins
 899 related to a variety of metabolic pathways by genistein-induced *Bradyrhizobium japonicum*
 900 strains. J. Proteom. **75**(4): 1211–1219. doi:10.1016/j.jprot.2011.10.032.
- 901 Dainese-Hatt, P., Fischer, H.M., Hennecke, H., and James, P. 1999. Classifying symbiotic proteins
 902 from *Bradyrhizobium japonicum* into functional groups by proteome analysis of altered gene
 903 expression levels. Electrophoresis **20**(18): 3514–3520. doi:10.1002/(SICI)1522-
 904 2683(19991201)20:18<3514::AID-ELPS3514>3.0.CO;2-T.
- 905 Daubech, B., Remigi, P., Doin de Moura, G., Marchetti, M., Pouzet, C., Auriac, M.-C., Gokhale,
 906 C.S., Masson-Boivin, C., and Capela, D. 2017. Spatio-temporal control of mutualism in
 907 legumes helps spread symbiotic nitrogen fixation. Elife **6**: 57. doi:10.7554/eLife.28683.
- 908 DeJong, T.M., Brewin, N.J., and Phillips, D.A. 1981. Effects of plasmid content in *Rhizobium*
 909 *leguminosarum* on pea nodule activity and plant growth. J. Gen. Microbiol. **124**: 1-7.
 910 doi:10.1099/00221287-124-1-1.
- 911 De Meyer, S.E., Briscoe, L., Martínez-Hidalgo, P., Agapakis, C.M., de-Los Santos, P.E., Seshadri,
 912 R., Reeve, W., Weinstock, G., O'Hara, G., Howieson, J.G., and Hirsch, A.M. 2016. Symbiotic
 913 *Burkholderia* species show diverse arrangements of *nif*/*fix* and *nod* genes and lack typical
 914 high-affinity cytochrome *cbb3* oxidase genes. Mol. Plant Microbe Interact. **29**(8): 609–619.
 915 doi:10.1094/MPMI-05-16-0091-R.
- 916 De Nisco, N.J., Abo, R.P., Wu, C.M., Penterman, J., and Walker, G.C. 2014. Global analysis of
 917 cell cycle gene expression of the legume symbiont *Sinorhizobium meliloti*. Proc. Natl. Acad.
 918 Sci. U.S.A. **111**(9): 3217–3224. doi:10.1073/pnas.1400421111.

- 919 de Rudder, K.E., López-Lara, I.M., and Geiger, O. 2000. Inactivation of the gene for phospholipid
 920 N-methyltransferase in *Sinorhizobium meliloti*: phosphatidylcholine is required for normal
 921 growth. *Mol. Microbiol.* **37**(4): 763–772. doi:10.1046/j.1365-2958.2000.02032.x.
- 922 de Rudder, K.E.E., Sohlenkamp, C., and Geiger, O. 1999. Plant-exuded choline is used for
 923 rhizobial membrane lipid biosynthesis by phosphatidylcholine synthase. *J. Biol. Chem.*
 924 **274**(28): 20011–20016. doi:10.1074/jbc.274.28.20011.
- 925 Delmotte, N., Ahrens, C.H., Knief, C., Qeli, E., Koch, M., Fischer, H.-M., Vorholt, J.A.,
 926 Hennecke, H., and Pessi, G. 2010. An integrated proteomics and transcriptomics reference
 927 data set provides new insights into the *Bradyrhizobium japonicum* bacteroid metabolism in
 928 soybean root nodules. *Proteomics* **10**(7): 1391–1400. doi:10.1002/pmic.200900710.
- 929 Delmotte, N., Mondy, S., Alunni, B., Fardoux, J., Chaintreuil, C., Vorholt, J., Giraud, E., and
 930 Gourion, B. 2014. A proteomic approach of *Bradyrhizobium/Aeschynomene* root and stem
 931 symbioses reveals the importance of the *fixA* locus for symbiosis. *Int. J. Mol. Sci.* **15**(3): 3660–
 932 3670. doi:10.3390/ijms15033660.
- 933 diCenzo, G., Milunovic, B., Cheng, J., and Finan, T.M. 2013. The tRNA^{arg} gene and *engA* are
 934 essential genes on the 1.7-Mb pSymB megaplasmid of *Sinorhizobium meliloti* and were
 935 translocated together from the chromosome in an ancestral strain. *J. Bacteriol.* **195**(2): 202–
 936 212. doi:10.1128/JB.01758-12.
- 937 diCenzo, G.C., Benedict, A.B., Fondi, M., Walker, G.C., Finan, T.M., Mengoni, A., and Griffiths,
 938 J.S. 2018a. Robustness encoded across essential and accessory replicons in an ecologically
 939 versatile bacterium. *PLOS Genet.* **14**(4): e1007357. doi:10.1101/209916.
- 940 diCenzo, G.C., Checcucci, A., Bazzicalupo, M., Mengoni, A., Viti, C., Dziewit, L., Finan, T.M.,
 941 Galardini, M., and Fondi, M. 2016a. Metabolic modelling reveals the specialization of

- 942 secondary replicons for niche adaptation in *Sinorhizobium meliloti*. Nat. Commun. **7**: 12219.
 943 doi:10.1038/ncomms12219.
- 944 diCenzo, G.C., Debiec, K., Krzysztoforski, J., Uhrynowski, W., Mengoni, A., Fagorzi, C.,
 945 Gorecki, A., Dziewit, L., Bajda, T., Rzepa, G., and Drewniak, L. 2018b. Genomic and
 946 biotechnological characterization of the heavy-metal resistant, arsenic-oxidizing bacterium
 947 *Ensifer* sp. M14. Genes **9**(8): 379. doi:10.3390/genes9080379.
- 948 diCenzo, G.C., and Finan, T.M. 2018. Techniques for Large-Scale Bacterial Genome Manipulation
 949 and Characterization of the Mutants with Respect to *In Silico* Metabolic Reconstructions.
 950 Methods Mol. Biol. **1716**(Chapter 13): 291–314. Springer New York, New York, NY.
 951 doi:10.1007/978-1-4939-7528-0_13.
- 952 diCenzo, G.C., MacLean, A.M., Milunovic, B., Golding, G.B., and Finan, T.M. 2014. Examination
 953 of prokaryotic multipartite genome evolution through experimental genome reduction. PLOS
 954 Genet. **10**(10): e1004742. doi:10.1371/journal.pgen.1004742.
- 955 diCenzo, G.C., Mengoni, A., and Fondi, M. 2017a. Tn-Core: context-specific reconstruction of
 956 core metabolic models using Tn-seq data. bioRxiv. **221325**: doi:10.1101/221325.
- 957 diCenzo, G.C., Muhammed, Z., Østerås, M., O'Brien, S.A.P., and Finan, T.M. 2017b. A key
 958 regulator of the glycolytic and gluconeogenic central metabolic pathways in *Sinorhizobium*
 959 *meliloti*. Genetics **207**(3): 961–974. doi:10.1534/genetics.117.300212.
- 960 diCenzo, G.C., Zamani, M., Ludwig, H.N., and Finan, T.M. 2017c. Heterologous
 961 complementation reveals a specialized activity for BacA in the *Medicago-Sinorhizobium*
 962 *meliloti* symbiosis. Mol. Plant Microbe Interact. **30**(4): 312–324. doi:10.1094/MPMI-02-17-
 963 0030-R.

- 964 diCenzo, G.C., Zamani, M., Milunovic, B., and Finan, T.M. 2016b. Genomic resources for
 965 identification of the minimal N₂-fixing symbiotic genome. *Environ. Microbiol.* **18**(8): 2534–
 966 2547. doi:10.1111/1462-2920.13221.
- 967 Djordjevic, M.A. 2004. *Sinorhizobium meliloti* metabolism in the root nodule: A proteomic
 968 perspective. *Proteomics* **4**(7): 1859–1872. doi:10.1002/pmic.200300802.
- 969 Djordjevic, M.A., Chen, H.C., Natera, S., Van Noorden, G., Menzel, C., Taylor, S., Renard, C.,
 970 Geiger, O., and Weiller, G.F. 2007. A global analysis of protein expression profiles in
 971 *Sinorhizobium meliloti*: discovery of new genes for nodule occupancy and stress adaptation.
 972 *Mol. Plant Microbe Interact.* **16**(6): 508–524. . doi:10.1094/MPMI.2003.16.6.508.
- 973 Döhlemann, J., Brennecke, M., and Becker, A. 2016. Cloning-free genome engineering in
 974 *Sinorhizobium meliloti* advances applications of Cre/*loxP* site-specific recombination. *J.*
 975 *Biotechnol.* **233**: 160–170. doi:10.1016/j.jbiotec.2016.06.033.
- 976 Dominguez-Ferreras, A., Perez-Arnedo, R., Becker, A., Olivares, J., Soto, M.J., and Sanjuán, J.
 977 2006. Transcriptome profiling reveals the importance of plasmid pSymB for osmoadaptation
 978 of *Sinorhizobium meliloti*. *J. Bacteriol.* **188**(21): 7617–7625. doi:10.1128/JB.00719-06.
- 979 Donati, A.J., Jeon, J.-M., Sangurdekar, D., So, J.-S., and Chang, W.-S. 2011. Genome-wide
 980 transcriptional and physiological responses of *Bradyrhizobium japonicum* to paraquat-
 981 mediated oxidative stress. *Appl. Environ. Microbiol.* **77**(11): 3633–3643.
 982 doi:10.1128/AEM.00047-11.
- 983 Dong, R., Zhang, J., Huan, H., Bai, C., Chen, Z., and Liu, G. 2017. High salt tolerance of a
 984 *Bradyrhizobium* strain and its promotion of the growth of *Stylosanthes guianensis*. *Int. J. Mol.*
 985 *Sci.* **18**(8): 1625. doi:10.3390/ijms18081625.

- 986 Downie, J.A. 2010. The roles of extracellular proteins, polysaccharides and signals in the
 987 interactions of rhizobia with legume roots. *FEMS Microbiol. Rev.* **34**(2): 150–170.
 988 doi:10.1111/j.1574-6976.2009.00205.x.
- 989 Draghi, W.O., Del Papa, M.F., Hellweg, C., Watt, S.A., Watt, T.F., Barsch, A., Lozano, M.J.,
 990 Salas, M.E., López, J.L., Albicoro, F.J., Nilsson, J.F., Torres Tejerizo, G.A., Luna, M.F.,
 991 Pistorio, M., Boiardi, J.L., Pühler, A., Weidner, S., Niehaus, K., and Lagares, A. 2016. A
 992 consolidated analysis of the physiologic and molecular responses induced under acid stress in
 993 the legume-symbiont model-soil bacterium *Sinorhizobium meliloti*. *Sci. Rep.* **6**(1): 29278.
 994 doi:10.1038/srep29278.
- 995 Dwivedi, S.L., Sahrawat, K.L., Upadhyaya, H.D., Mengoni, A., Galardini, M., Bazzicalupo, M.,
 996 Biondi, E.G., Hungria, M., Kaschuk, G., Blair, M.W., and Ortiz, R. 2015. Advances in host
 997 plant and rhizobium genomics to enhance symbiotic nitrogen fixation in grain legumes. *Adv.*
 998 *Agron.* 129:1–116. doi:10.1016/bs.agron.2014.09.001.
- 999 Encarnación, S., del Carmen Vargas, M., Dunn, M.F., Dávalos, A., Mendoza, G., Mora, Y., and
 1000 Mora, J. 2002. AniA regulates reserve polymer accumulation and global protein expression in
 1001 *Rhizobium etli*. *J. Bacteriol.* **184**(8): 2287–2295. doi:10.1128/JB.184.8.2287-2295.2002.
- 1002 Encarnación, S., Guzmán, Y., Dunn, M.F., Hernández, M., del Carmen Vargas, M., and Mora, J.
 1003 2003. Proteome analysis of aerobic and fermentative metabolism in *Rhizobium etli* CE3.
 1004 *Proteomics* **3**(6): 1077–1085. doi:10.1002/pmic.200300427.
- 1005 Erisman, J.W., Sutton, M.A., Galloway, J., Klimont, Z., and Winiwarter, W. 2008. How a century
 1006 of ammonia synthesis changed the world. *Nat. Geosci.* **1**(10): 636–639. doi:10.1038/ngeo325.

- 1007 Evans, J., Barnet, Y.M., and Vincent, J.M. 1979. Effect of a bacteriophage on colonisation and
1008 nodulation of clover roots by paired strains of *Rhizobium trifolii*. Can. J. Microbiol. **25**(9):
1009 974–978.
- 1010 Fan, Y., Liu, J., Lyu, S., Wang, Q., Yang, S., and Zhu, H. 2017. The soybean *RfgI* gene restricts
1011 nodulation by *Sinorhizobium fredii* USDA193. Front. Plant Sci. **8**: 1548.
1012 doi:10.3389/fpls.2017.01548.
- 1013 Fei, F., diCenzo, G.C., Bowdish, D.M.E., McCarry, B.E., and Finan, T.M. 2016. Effects of
1014 synthetic large-scale genome reduction on metabolism and metabolic preferences in a
1015 nutritionally complex environment. Metabolomics **12**(2): 23. doi:10.1007/s11306-015-0928-
1016 y.
- 1017 Finan, T.M., Kunkel, B., De Vos, G.F., and Signer, E.R. 1986. Second symbiotic megaplasmid in
1018 *Rhizobium meliloti* carrying exopolysaccharide and thiamine synthesis genes. J. Bacteriol.
1019 **167**(1): 66–72. doi:10.1128/jb.167.1.66-72.1986.
- 1020 Finkel, O.M., Castrillo, G., Herrera Paredes, S., Salas González, I., and Dangl, J.L. 2017.
1021 Understanding and exploiting plant beneficial microbes. Curr. Opin. Plant Biol. **38**: 155–163.
1022 doi:10.1016/j.pbi.2017.04.018.
- 1023 Fondi, M., and Liò, P. 2015. Multi -omics and metabolic modelling pipelines: challenges and tools
1024 for systems microbiology. Microbiol Res **171**: 52–64. doi:10.1016/j.micres.2015.01.003.
- 1025 Frederix, M., Edwards, A., Swiderska, A., Stanger, A., Karunakaran, R., Williams, A., Abbruscato,
1026 P., Sanchez-Contreras, M., Poole, P.S., and Downie, J.A. 2014. Mutation of *praR* in
1027 *Rhizobium leguminosarum* enhances root biofilms, improving nodulation competitiveness by
1028 increased expression of attachment proteins. Mol. Microbiol. **93**(3): 464–478.
1029 doi:10.1111/mmi.12670.

- 1030 Fuhrer, T., Fischer, E., and Sauer, U. 2005. Experimental identification and quantification of
 1031 glucose metabolism in seven bacterial species. *J. Bacteriol.* **187**(5): 1581–1590.
 1032 doi:10.1128/JB.187.5.1581-1590.2005.
- 1033 Gage, D.J. 2004. Infection and invasion of roots by symbiotic, nitrogen-fixing rhizobia during
 1034 nodulation of temperate legumes. *Microbiol. Mol. Biol. Rev.* **68**(2): 280–300.
 1035 doi:10.1128/MMBR.68.2.280-300.2004.
- 1036 Galardini, M., Mengoni, A., Brilli, M., Pini, F., Fioravanti, A., Lucas, S., Lapidus, A., Cheng, J.-
 1037 F., Goodwin, L., Pitluck, S., Land, M., Hauser, L., Woyke, T., Mikhailova, N., Ivanova, N.,
 1038 Daligault, H., Bruce, D., Detter, C., Tapia, R., Han, C., Teshima, H., Mocali, S., Bazzicalupo,
 1039 M., and Biondi, E.G. 2011. Exploring the symbiotic pangenome of the nitrogen-fixing
 1040 bacterium *Sinorhizobium meliloti*. *BMC Genomics* **12**(1): 235. doi:10.1186/1471-2164-12-
 1041 235.
- 1042 Galardini, M., Pini, F., Bazzicalupo, M., Biondi, E.G., and Mengoni, A. 2013. Replicon-dependent
 1043 bacterial genome evolution: the case of *Sinorhizobium meliloti*. *Genome Biol. Evol.* **5**(3): 542–
 1044 558. doi:10.1093/gbe/evt027.
- 1045 Gao, J.-L., Weissenmayer, B., Taylor, A.M., Thomas-Oates, J., López-Lara, I.M., and Geiger, O.
 1046 2004. Identification of a gene required for the formation of lyso-ornithine lipid, an
 1047 intermediate in the biosynthesis of ornithine-containing lipids. *Mol. Microbiol.* **53**(6): 1757–
 1048 1770. doi:10.1111/j.1365-2958.2004.04240.x.
- 1049 Gao, M., Barnett, M.J., Long, S.R., and Teplitski, M. 2010. Role of the *Sinorhizobium meliloti*
 1050 global regulator Hfq in gene regulation and symbiosis. *Mol. Plant Microbe Interact.* **23**(4):
 1051 355–365. doi:10.1094/MPMI-23-4-0355.

- 1052 Gao, M., Chen, H., Eberhard, A., Gronquist, M.R., Robinson, J.B., Connolly, M., Teplitski, M.,
 1053 Rolfe, B.G., and Bauer, W.D. 2007. Effects of AiiA-mediated quorum quenching in
 1054 *Sinorhizobium meliloti* on quorum-sensing signals, proteome patterns, and symbiotic
 1055 interactions. *Mol. Plant Microbe Interact.* **20**(7): 843–856. doi:10.1094/MPMI-20-7-0843.
- 1056 Gao, M., Chen, H., Eberhard, A., Gronquist, M.R., Robinson, J.B., Rolfe, B.G., and Bauer, W.D.
 1057 2005. *sinI*- and *expR*-dependent quorum sensing in *Sinorhizobium meliloti*. *J. Bacteriol.*
 1058 **187**(23): 7931–7944. doi:10.1128/JB.187.23.7931-7944.2005.
- 1059 Gao, T.G., Xu, Y.Y., Jiang, F., Li, B.Z., Yang, J.S., Wang, E.T., and Yuan, H.L. 2015. Nodulation
 1060 characterization and proteomic profiling of *Bradyrhizobium liaoningense* CCBAU05525 in
 1061 response to water-soluble humic materials. *Sci. Rep.* **5**(1): 10836. doi:10.1038/srep10836.
- 1062 Garcia-Fraile, P., Seaman, J.C., Karunakaran, R., Edwards, A., Poole, P.S., and Downie, J.A. 2015.
 1063 Arabinose and protocatechuate catabolism genes are important for growth of *Rhizobium*
 1064 *leguminosarum* biovar *viciae* in the pea rhizosphere. *Plant Soil* **390**(1-2): 251–264.
 1065 doi:10.1007/s11104-015-2389-5.
- 1066 García-de-los-Santos, A., López, E., Cubillas, C.A., Noel, K.D., Brom, S., and Romero, D. 2008.
 1067 Requirement of a plasmid-encoded catalase for survival of *Rhizobium etli* CFN42 in a
 1068 polyphenol-rich environment. *Appl. Environ. Microbiol.* **74**(8): 2398–2403.
 1069 doi:10.1128/AEM.02457-07.
- 1070 Geddes, B.A., Ryu, M.-H., Mus, F., Garcia Costas, A., Peters, J.W., Voigt, C.A., and Poole, P.
 1071 2015. Use of plant colonizing bacteria as chassis for transfer of N₂-fixation to cereals. *Curr.*
 1072 *Opin. Biotech.* **32**: 216–222. doi:10.1016/j.copbio.2015.01.004.

- 1073 Gemperline, E., Jayaraman, D., Maeda, J., Ané, J.-M., and Li, L. 2015. Multifaceted investigation
1074 of metabolites during nitrogen fixation in *Medicago* via high resolution MALDI-MS imaging
1075 and ESI-MS. *J. Am. Soc. Mass Spectrom.* **26**(1): 149–158. doi:10.1007/s13361-014-1010-0.
- 1076 Ghobakhlou, A., Laberge, S., Antoun, H., Wishart, D.S., Xia, J., Krishnamurthy, R., and Mandal,
1077 R. 2013. Metabolomic analysis of cold acclimation of Arctic *Mesorhizobium* sp. strain N33.
1078 *PLOS One* **8**(12): e84801. doi:10.1371/journal.pone.0084801.
- 1079 Ghobakhlou, A.-F., Johnston, A., Harris, L., Antoun, H., and Laberge, S. 2015. Microarray
1080 transcriptional profiling of Arctic *Mesorhizobium* strain N33 at low temperature provides
1081 insights into cold adaption strategies. *BMC Genomics* **16**(1): 383. doi:10.1186/s12864-015-
1082 1611-4.
- 1083 Gibson, K.E., Barnett, M.J., Toman, C.J., Long, S.R., and Walker, G.C. 2007. The symbiosis
1084 regulator CbrA modulates a complex regulatory network affecting the flagellar apparatus and
1085 cell envelope proteins. *J. Bacteriol.* **189**(9): 3591–3602. doi:10.1128/JB.01834-06.
- 1086 Gibson, K.E., Kobayashi, H., and Walker, G.C. 2008. Molecular determinants of a symbiotic
1087 chronic infection. *Annu. Rev. Genet.* **42**(1): 413–441.
1088 doi:10.1146/annurev.genet.42.110807.091427.
- 1089 Giraud, E., Moulin, L., Vallenet, D., Barbe, V., Cytryn, E., Avarre, J.-C., Jaubert, M., Simon, D.,
1090 Cartieaux, F., Prin, Y., Bena, G., Hannibal, L., Fardoux, J., Kojadinovic, M., Vuillet, L., Lajus,
1091 A., Cruveiller, S., Rouy, Z., Mangenot, S., Segurens, B., Dossat, C., Franck, W.L., Chang, W.-
1092 S., Saunders, E., Bruce, D., Richardson, P., Normand, P., Dreyfus, B., Pignol, D., Stacey, G.,
1093 Emerich, D., Verméglio, A., Médigue, C., and Sadowsky, M. 2007. Legumes symbioses:
1094 absence of *nod* genes in photosynthetic bradyrhizobia. *Science* **316**(5829): 1307–1312.
1095 doi:10.1126/science.1139548.

- 1096 Glucksmann, M.A., Reuber, T.L., and Walker, G.C. 1993. Genes needed for the modification,
 1097 polymerization, export, and processing of succinoglycan by *Rhizobium meliloti*: a model for
 1098 succinoglycan biosynthesis. *J. Bacteriol.* **175**(21): 7045–7055. doi:10.1128/jb.175.21.7045-
 1099 7055.1993.
- 1100 Gomes, D.F., Batista, J.S.D.S., Schiavon, A.L., Andrade, D.S., and Hungria, M. 2012a. Proteomic
 1101 profiling of *Rhizobium tropici* PRF 81: identification of conserved and specific responses to
 1102 heat stress. *BMC Microbiol.* **12**(1): 84. doi:10.1186/1471-2180-12-84.
- 1103 Gomes, D.F., Batista, J.S.D.S., Torres, A.R., de Souza Andrade, D., Galli-Terasawa, L.V., and
 1104 Hungria, M. 2012b. Two-dimensional proteome reference map of *Rhizobium tropici* PRF 81
 1105 reveals several symbiotic determinants and strong resemblance with agrobacteria. *Proteomics*
 1106 **12**(6): 859–863. doi:10.1002/pmic.201100406.
- 1107 Gomes, D.F., da Silva Batista, J.S., Rolla, A.A.P., da Silva, L.P., Bloch, C., Galli-Terasawa, L.V.,
 1108 and Hungria, M. 2014. Proteomic analysis of free-living *Bradyrhizobium diazoefficiens*:
 1109 highlighting potential determinants of a successful symbiosis. *BMC Genomics* **15**(1): 643.
 1110 doi:10.1186/1471-2164-15-643.
- 1111 González, V., Acosta, J.L., Santamaría, R.I., Bustos, P., Fernández, J.L., Hernández González,
 1112 I.L., Díaz, R., Flores, M., Palacios, R., Mora, J., and Dávila, G. 2010. Conserved symbiotic
 1113 plasmid DNA sequences in the multireplicon pangenomic structure of *Rhizobium etli*. *Appl.*
 1114 *Environ. Microbiol.* **76**(5): 1604–1614. doi:10.1128/AEM.02039-09.
- 1115 González, V., Bustos, P., Ramírez-Romero, M.A., Medrano-Soto, A., Salgado, H., Hernández-
 1116 González, I., Hernández-Celis, J.C., Quintero, V., Moreno-Hagelsieb, G., Girard, L.,
 1117 Rodríguez, O., Flores, M., Cevallos, M.A., Collado-Vides, J., Romero, D., and Dávila, G.
 1118 2003. The mosaic structure of the symbiotic plasmid of *Rhizobium etli* CFN42 and its relation

- 1119 to other symbiotic genome compartments. *Genome Biol.* **4**(6): R36. doi:10.1186/gb-2003-4-
1120 6-r36.
- 1121 Gourion, B., Berrabeh, F., Ratet, P., and Stacey, G. 2015. Rhizobium-legume symbioses: the
1122 crucial role of plant immunity. *Trends Plant Sci.* **20**(3): 186-194.
1123 doi:10.1016/j.tplants.2014.11.008.
- 1124 Gourion, B., Sulser, S., Frunzke, J., Francez-Charlot, A., Stiefel, P., Pessi, G., Vorholt, J.A.,
1125 Fischer, H.M. 2009. The PhyR-sigma(EcfG) signalling cascade is involved in stress response
1126 and symbiotic efficiency in *Bradyrhizobium japonicum*. *Mol. Microbiol.* **73**(2): 291-305.
1127 doi:10.1111/j.1365-2958.2009.06769.x.
- 1128 Guan, S.H., Gris, C., Cruveiller, S., Pouzet, C., Tasse, L., Leru, A., Maillard, A., Médigue, C.,
1129 Batut, J., Masson-Boivin, C., and Capela, D. 2013. Experimental evolution of nodule
1130 intracellular infection in legume symbionts. *ISME J.* **7**(7): 1367–1377.
1131 doi:10.1038/ismej.2013.24.
- 1132 Guerreiro, N., Djordjevic, M.A., and Rolfe, B.G. 1999. Proteome analysis of the model
1133 microsymbiont *Sinorhizobium meliloti*: isolation and characterisation of novel proteins.
1134 Electrophoresis **20**(4-5): 818–825. doi:10.1002/(SICI)1522-
1135 2683(19990101)20:4/5<818::AID-ELPS818>3.0.CO;2-6.
- 1136 Guerreiro, N., Redmond, J.W., Rolfe, B.G., and Djordjevic, M.A. 1997. New *Rhizobium*
1137 *leguminosarum* flavonoid-induced proteins revealed by proteome analysis of differentially
1138 displayed proteins. *Mol. Plant Microbe Interact.* **10**(4): 506–516.
1139 doi:10.1094/MPMI.1997.10.4.506.
- 1140 Guerreiro, N., Stepkowski, T., Rolfe, B.G., and Djordjevic, M.A. 1998. Determination of plasmid-
1141 encoded functions in *Rhizobium leguminosarum* biovar *trifolii* using proteome analysis of

- 1142 plasmid-cured derivatives. Electrophoresis **19**(11): 1972–1979.
 1143 doi:10.1002/elps.1150191115.
- 1144 Haag, A.F., Arnold, M.F.F., Myka, K.K., Kerscher, B., Dall'Angelo, S., Zanda, M., Mergaert, P.,
 1145 and Ferguson, G.P. 2013. Molecular insights into bacteroid development during *Rhizobium*-
 1146 legume symbiosis. FEMS Microbiol. Rev. **37**(3): 364–383. doi:10.1111/1574-6976.12003.
- 1147 Hagen, M.J., and Hamrick, J.L. 1996. Population level processes in *Rhizobium leguminosarum* bv.
 1148 *trifolii*: the role of founder effects. Mol. Ecol. **5**(6): 707–714. doi:10.1111/j.1365-
 1149 294X.1996.tb00367.x.
- 1150 Harrison, C.L., Crook, M.B., Peco, G., Long, S.R., and Griffitts, J.S. 2011. Employing site-specific
 1151 recombination for conditional genetic analysis in *Sinorhizobium meliloti*. Appl. Environ.
 1152 Microbiol. **77**(12): 3916–3922. doi:10.1128/AEM.00544-11.
- 1153 Harrison, S.P., Jones, D.G., and Young, J.P.W. 1989. *Rhizobium* population genetics: genetic
 1154 variation within and between populations from diverse locations. Microbiology **135**(5): 1061–
 1155 1069. doi:10.1099/00221287-135-5-1061.
- 1156 Hashem, F.M., and Angle, J.S. 1990. Rhizobiophage effects on nodulation, nitrogen fixation, and
 1157 yield of field-grown soybeans (*Glycine max* L. Merr.). Biol. Fert. Soils **9**(4): 330–334.
 1158 doi:10.1007/BF00634110.
- 1159 Haskett, T.L., Terpolilli, J.J., Bekuma, A., O'Hara, G.W., Sullivan, J.T., Wang, P., Ronson, C.W.,
 1160 and Ramsay, J.P. 2016. Assembly and transfer of tripartite integrative and conjugative genetic
 1161 elements. Proc. Natl. Acad. Sci. U.S.A. **113**(43): 12268–12273.
 1162 doi:10.1073/pnas.1613358113.
- 1163 Haskett, T.L., Terpolilli, J.J., Ramachandran, V.K., Verdonk, C.J., Poole, P.S., O'Hara, G.W., and
 1164 Ramsay, J.P. 2018. Sequential induction of three recombination directionality factors directs

- assembly of tripartite integrative and conjugative elements. PLOS Genet. **14**(3): e1007292.
doi:10.1371/journal.pgen.1007292.
- Hawkins, J.P., Geddes, B.A., and Oresnik, I.J. 2017. Succinoglycan production contributes to
acidic pH tolerance in *Sinorhizobium meliloti* Rm1021. Mol. Plant Microbe Interact. **30**(12):
1009–1019. doi:10.1094/MPMI-07-17-0176-R.
- Hawkins, J.P., Ordonez, P.A., and Oresnik, I.J. 2018. Characterization of mutants that affect the
non-oxidative pentose phosphate pathway in *Sinorhizobium meliloti*. J. Bacteriol. **200**(2):
e00436–17. doi:10.1128/JB.00436-17.
- Heath, K.D., and Tiffin, P. 2009. Stabilizing mechanisms in a legume-rhizobium mutualism.
Evolution **63**(3): 652–662. doi:10.1111/j.1558-5646.2008.00582.x.
- Heath, K.D., Burke, P.V., and Stinchcombe, J.R. 2012. Coevolutionary genetic variation in the
legume-rhizobium transcriptome. Mol. Ecol. **21**(19): 4735–4747. doi:10.1111/j.1365-
294X.2012.05629.x.
- Heil, J.R., Cheng, J., and Charles, T.C. 2012. Site-specific bacterial chromosome engineering:
ΦC31 integrase mediated cassette exchange (IMCE). J. Vis. Exp. **61**: e3698.
doi:10.3791/3698.
- Heinz, E.B., and Streit, W.R. 2003. Biotin limitation in *Sinorhizobium meliloti* strain 1021 alters
transcription and translation. Appl. Environ. Microbiol. **69**(2): 1206–1213.
doi:10.1128/AEM.69.2.1206-1213.2003.
- Hellweg, C., Pühler, A., and Weidner, S. 2009. The time course of the transcriptomic response of
Sinorhizobium meliloti 1021 following a shift to acidic pH. BMC Microbiol. **9**(1): 37.
doi:10.1186/1471-2180-9-37.

- 1187 Hempel, J., Zehner, S., Göttfert, M., and Patschkowski, T. 2009. Analysis of the secretome of the
1188 soybean symbiont *Bradyrhizobium japonicum*. J. Biotechnol. **140**(1-2): 51–58.
1189 doi:10.1016/j.jbiotec.2008.11.002.
- 1190 Hinsinger, P., Bengough, A.G., Vetterlein, D., and Young, I.M. 2009. Rhizosphere: biophysics,
1191 biogeochemistry and ecological relevance. Plant Soil **321**(1-2): 117–152. doi:10.1007/s11104-
1192 008-9885-9.
- 1193 Hirsch, A.M., Wilson, K.J., Jones, J.D., Bang, M., Walker, V.V., and Ausubel, F.M. 1984.
1194 *Rhizobium meliloti* nodulation genes allow *Agrobacterium tumefaciens* and *Escherichia coli*
1195 to form pseudonodules on alfalfa. J. Bacteriol. **158**(3): 1133–1143.
- 1196 Hirsch, P.R. 1979. Plasmid-determined bacteriocin production by *Rhizobium leguminosarum*.
1197 Microbiology **113**(2): 219–228. doi:10.1099/00221287-113-2-219.
- 1198 Hood, G., Ramachandran, V., East, A.K., Downie, J.A., and Poole, P.S. 2017. Manganese transport
1199 is essential for N₂-fixation by *Rhizobium leguminosarum* in bacteroids from galeoid but not
1200 phaseoloid nodules. Environ. Microbiol. **19**(7): 2715–2726. doi:10.1111/1462-2920.13773.
- 1201 Hooykaas, P.J., Snijdwint, F.G., and Schilperoort, R.A. 1982. Identification of the Sym plasmid
1202 of *Rhizobium leguminosarum* strain 1001 and its transfer to and expression in other rhizobia
1203 and *Agrobacterium tumefaciens*. Plasmid **8**(1): 73–82. doi:10.1016/0147-619X(82)90042-7.
- 1204 House, B.L., Mortimer, M.W., and Kahn, M.L. 2004. New recombination methods for
1205 *Sinorhizobium meliloti* genetics. Appl. Environ. Microbiol. **70**(5): 2806–2815.
1206 doi:10.1128/AEM.70.5.2806-2815.2004.
- 1207 Hu, G., Li, Y., and Zhou, J. 2007. Biological characteristics of plasmids of *Mesorhizobium huakuii*
1208 HN3015 from *Astragalus sinicus*. World J. Microbiol. Biotechnol. **23**(6): 845–851.
1209 doi:10.1007/s11274-006-9308-0.

- 1210 Hu, G., Li, Y., and Zhou, J. 2008. Incompatibility behavior of a megaplasmid pMhHN3015c in
 1211 *Mesorhizobium huakuii* HN3015. World J. Microbiol. Biotechnol. **24**(8): 1281–1287.
 1212 doi:10.1007/s11274-007-9611-4.
- 1213 Hynes, M.F., and McGregor, N.F. 1990. Two plasmids other than the nodulation plasmid are
 1214 necessary for formation of nitrogen-fixing nodules by *Rhizobium leguminosarum*. Mol.
 1215 Microbiol. **4**(4): 567–574. doi:10.1111/j.1365-2958.1990.tb00625.x.
- 1216 Hynes, M.F., Quandt, J., O'Connell, M.P., and Pühler, A. 1989. Direct selection for curing and
 1217 deletion of *Rhizobium* plasmids using transposons carrying the *Bacillus subtilis sacB* gene.
 1218 Gene **78**(1): 111–120. doi:10.1016/0378-1119(89)90319-3.
- 1219 Hynes, M.F., Simon, R., Müller, P., Niehaus, K., Labes, M., and Pühler, A. 1986. The two
 1220 megaplasms of *Rhizobium meliloti* are involved in the effective nodulation of alfalfa. Mol.
 1221 Gen. Genet. **202**(3): 356–362. doi:10.1007/BF00333262.
- 1222 Irar, S., González, E.M., Arrese-Igor, C., and Marino, D. 2014. A proteomic approach reveals new
 1223 actors of nodule response to drought in split-root grown pea plants. Physiol. Plant. **152**(4):
 1224 634–645. doi:10.1111/ppl.12214.
- 1225 Ivleva, N.B., Groat, J., Staub, J.M., and Stephens, M. 2016. Expression of active subunit of
 1226 nitrogenase via integration into plant organelle genome. PLOS One **11**(8): e0160951.
 1227 doi:10.1371/journal.pone.0160951.
- 1228 Jagadish, M.N., and Szalay, A.A. 1984. Directed transposon Tn5 mutagenesis and
 1229 complementation in slow-growing, broad host range cowpea *Rhizobium*. Mol. Gen. Genet.
 1230 **196**(2): 290–300.
- 1231 Jebbar, M., Sohn-Bosser, L., Bremer, E., Bernard, T., and Blanco, C. 2005. Ectoine-induced
 1232 proteins in *Sinorhizobium meliloti* include an ectoine ABC-type transporter involved in

- 1233 osmoprotection and ectoine catabolism. J. Bacteriol. **187**(4): 1293–1304.
 1234 doi:10.1128/JB.187.4.1293-1304.2005.
- 1235 Jeon, J.-M., Lee, H.-I., Donati, A.J., So, J.-S., Emerich, D.W., and Chang, W.-S. 2011. Whole-
 1236 genome expression profiling of *Bradyrhizobium japonicum* in response to hydrogen peroxide.
 1237 Mol. Plant Microbe Interact. **24**(12): 1472–1481. doi:10.1094/MPMI-03-11-0072.
- 1238 Ji, Z.J., Yan, H., Cui, Q.G., Wang, E.T., Chen, W.F., and Chen, W.X. 2017. Competition between
 1239 rhizobia under different environmental conditions affects the nodulation of a legume. Syst.
 1240 Appl. Microbiol. **40**(2): 114–119. doi:10.1016/j.syapm.2016.12.003.
- 1241 Jiménez-Guerrero, I., Acosta Jurado, S., Del Cerro, P., Navarro-Gómez, P., López-Baena, F.J.,
 1242 Ollero, F.J., Vinardell, J.M., and Pérez-Montaña, F. 2017. Transcriptomic studies of the effect
 1243 of *nod* gene-inducing molecules in rhizobia: different weapons, one purpose. Genes **9**(1): 1.
 1244 doi:10.3390/genes9010001.
- 1245 Jones, K.M., Kobayashi, H., Davies, B.W., Taga, M.E., and Walker, G.C. 2007. How rhizobial
 1246 symbionts invade plants: the *Sinorhizobium–Medicago* model. Nat. Rev. Microbiol. **5**(8):
 1247 619–633. doi:10.1038/nrmicro1705.
- 1248 Jozefkiewicz, C., Brambilla, S., Frare, R., Stritzler, M., Puente, M., Piccinetti, C., Soto, G., and
 1249 Ayub, N. 2017. Microevolution rather than large genome divergence determines the
 1250 effectiveness of legume-rhizobia symbiotic interaction under field conditions. J. Mol. Evol.
 1251 **85**(3-4): 79–83. doi:10.1007/s00239-017-9808-6.
- 1252 Kajiwar, H., Kaneko, T., Ishizaka, M., Tajima, S., and Kouchi, H. 2003. Protein profile of
 1253 symbiotic bacteria *Mesorhizobium loti* MAFF303099 in mid-growth phase. Biosci.
 1254 Biotechnol. Biochem. **67**(12): 2668–2673. doi:10.1271/bbb.67.2668.

- 1255 Karunakaran, R., Haag, A.F., East, A.K., Ramachandran, V.K., Prell, J., James, E.K., Scocchi, M.,
 1256 Ferguson, G.P., and Poole, P.S. 2010. BacA is essential for bacteroid development in nodules
 1257 of galeoid, but not phaseoloid, legumes. *J. Bacteriol.* **192**(11): 2920–2928.
 1258 doi:10.1128/JB.00020-10.
- 1259 Karunakaran, R., Ramachandran, V.K., Seaman, J.C., East, A.K., Mouhsine, B., Mauchline, T.H.,
 1260 Prell, J., Skeffington, A., and Poole, P.S. 2009. Transcriptomic analysis of *Rhizobium*
 1261 *leguminosarum* biovar *viciae* in symbiosis with host plants *Pisum sativum* and *Vicia cracca*.
 1262 *J. Bacteriol.* **191**(12): 4002–4014. doi:10.1128/JB.00165-09.
- 1263 Keum, Y.S., Seo, J.S., Li, Q.X., and Kim, J.H. 2008. Comparative metabolomic analysis of
 1264 *Sinorhizobium* sp. C4 during the degradation of phenanthrene. *Appl. Microbiol. Biotechnol.*
 1265 **80**(5): 863–872. doi:10.1007/s00253-008-1581-4.
- 1266 Kiers, E.T., and Denison, R.F. 2008. Sanctions, cooperation, and the stability of plant-rhizosphere
 1267 mutualisms. *Annu. Rev. Ecol. Evol. Syst.* **39**(1): 215–236.
 1268 doi:10.1146/annurev.ecolsys.39.110707.173423.
- 1269 Kiers, E.T., Rousseau, R.A., West, S.A., and Denison, R.F. 2003. Host sanctions and the legume-
 1270 rhizobium mutualism. *Nature* **425**(6953): 78–81. doi:10.1038/nature01931.
- 1271 Klonowska, A., Melkonian, R., Miché, L., Tisseyre, P., and Moulin, L. 2018. Transcriptomic
 1272 profiling of *Burkholderia phymatum* STM815, *Cupriavidus taiwanensis* LMG19424 and
 1273 *Rhizobium mesoamericanum* STM3625 in response to *Mimosa pudica* root exudates
 1274 illuminates the molecular basis of their nodulation competitiveness and symbiotic
 1275 evolutionary history. *BMC Genomics* **19**(1): 105. doi:10.1186/s12864-018-4487-2.

- 1276 Kondorosi, E., Mergaert, P., and Kereszt, A. 2013. A paradigm for endosymbiotic life: cell
 1277 differentiation of *Rhizobium* bacteria provoked by host plant factors. *Annu. Rev. Microbiol.*
 1278 **67**(1): 611–628. doi:10.1146/annurev-micro-092412-155630.
- 1279 Krol, E., and Becker, A. 2004. Global transcriptional analysis of the phosphate starvation response
 1280 in *Sinorhizobium meliloti* strains 1021 and 2011. *Mol. Genet. Genomics* **272**(1).
 1281 doi:10.1007/s00438-004-1030-8.
- 1282 Krol, E., and Becker, A. 2011. ppGpp in *Sinorhizobium meliloti*: biosynthesis in response to
 1283 sudden nutritional downshifts and modulation of the transcriptome. *Mol. Microbiol.* **81**(5):
 1284 1233–1254. doi:10.1111/j.1365-2958.2011.07752.x.
- 1285 Krol, E., Klaner, C., Gnau, P., Kaeffer, V., Essen, L.-O., and Becker, A. 2016. Cyclic
 1286 mononucleotide- and Clr-dependent gene regulation in *Sinorhizobium meliloti*. *Microbiology*
 1287 **162**(10): 1840–1856. doi:10.1099/mic.0.000356.
- 1288 Krysciak, D., Grote, J., Rodriguez Orbegoso, M., Utpatel, C., Förstner, K.U., Li, L., Schmeisser,
 1289 C., Krishnan, H.B., and Streit, W.R. 2014. RNA sequencing analysis of the broad-host-range
 1290 strain *Sinorhizobium fredii* NGR234 identifies a large set of genes linked to quorum sensing-
 1291 dependent regulation in the background of a *trai* and *ngrI* deletion mutant. *Appl. Environ.*
 1292 *Microbiol.* **80**(18): 5655–5671. doi:10.1128/AEM.01835-14.
- 1293 Kumar, D., Yadav, A.K., Kadimi, P.K., Nagaraj, S.H., Grimmond, S.M., and Dash, D. 2013.
 1294 Proteogenomic analysis of *Bradyrhizobium japonicum* USDA110 using GenoSuite, an
 1295 automated multi-algorithmic pipeline. *Mol. Cell Proteomics* **12**(11): 3388–3397.
 1296 doi:10.1074/mcp.M112.027169.

- 1297 Lagares, A., Borella, G.C., Linne, U., Becker, A., and Valverde, C. 2017. Regulation of
1298 polyhydroxybutyrate accumulation in *Sinorhizobium meliloti* by the trans-encoded small RNA
1299 MmgR. J. Bacteriol. **199**(8): e00776–16. doi:10.1128/JB.00776-16.
- 1300 Laguerre, G. 1992. Plasmid profiles and restriction fragment length polymorphism of *Rhizobium*
1301 *leguminosarum* bv. *viciae* in field populations. FEMS Microbio. Lett. **101**(1): 17–26.
- 1302 Lamouche, F., Gully, D., Chaumeret, A., Nouwen, N., Verly, C., Pierre, O., Sciallano, C., Fardoux,
1303 J., Jeudy, C., Szűcs, A., Mondy, S., Salon, C., Nagy, I., Kereszt, A., Dessaux, Y., Giraud, E.,
1304 Mergaert, P., and Alunni, B. 2018. Transcriptomic dissection of *Bradyrhizobium* sp. strain
1305 ORS285 in symbiosis with *Aeschynomene* spp. inducing different bacteroid morphotypes with
1306 contrasted symbiotic efficiency. Environ. Microbiol. doi:10.1111/1462-2920.14292.
- 1307 Landeta, C., Dávalos, A., Cevallos, M.Á., Geiger, O., Brom, S., and Romero, D. 2011. Plasmids
1308 with a chromosome-like role in rhizobia. J. Bacteriol. **193**(6): 1317–1326.
1309 doi:10.1128/JB.01184-10.
- 1310 Lang, C., Smith, L.S., and Long, S.R. 2018. Characterization of novel plant symbiosis mutants
1311 using a new multiple gene-expression reporter *Sinorhizobium meliloti* strain. Front. Plant Sci.
1312 **9**: 76. doi:10.3389/fpls.2018.00076.
- 1313 Laranjo, M., Alexandre, A., and Oliveira, S. 2014. Genes commonly involved in acid tolerance
1314 are not overexpressed in the plant microsymbiont *Mesorhizobium loti* MAFF303099 upon
1315 acidic shock. Appl. Microbiol. Biotechnol. **98**(16): 7137–7147. doi:10.1007/s00253-014-
1316 5875-4.
- 1317 Laranjo, M., Alexandre, A., and Oliveira, S. 2017. Global transcriptional response to salt shock of
1318 the plant microsymbiont *Mesorhizobium loti* MAFF303099. Res. Microbiol. **168**(1): 55–63.
1319 doi:10.1016/j.resmic.2016.07.006.

- 1320 Lardi, M., Murset, V., Fischer, H.-M., Mesa, S., Ahrens, C.H., Zamboni, N., and Pessi, G. 2016.
 1321 Metabolomic profiling of *Bradyrhizobium diazoefficiens*-induced root nodules reveals both
 1322 host plant-specific and developmental signatures. *Int. J. Mol. Sci.* **17**(6): 815.
 1323 doi:10.3390/ijms17060815.
- 1324 Larrainzar, E., and Wienkoop, S. 2017. A proteomic view on the role of legume symbiotic
 1325 interactions. *Front. Plant Sci.* **8**: 1267. doi:10.3389/fpls.2017.01267.
- 1326 Ledermann, R., Bartsch, I., Müller, B., Wülser, J., and Fischer, H.-M. 2018. A functional general
 1327 stress response of *Bradyrhizobium diazoefficiens* is required for early stages of host plant
 1328 infection. *Mol. Plant Microbe Interact.* **31**(5): 537-547. doi:10.1094/MPMI-11-17-0284-R.
- 1329 Ledermann, R., Strebel, S., Kampik, C., and Fischer, H.-M. 2016. Versatile vectors for efficient
 1330 mutagenesis of *Bradyrhizobium diazoefficiens* and other Alphaproteobacteria. *Appl. Environ.*
 1331 *Microbiol.* **82**(9): 2791–2799. doi:10.1128/AEM.04085-15.
- 1332 Lehman, A.P., and Long, S.R. 2018. OxyR-dependent transcription response of *Sinorhizobium*
 1333 *meliloti* to oxidative stress. *J. Bacteriol.* **200**(7): e00622–17. doi:10.1128/JB.00622-17.
- 1334 Li, Y., Tian, C.F., Chen, W.F., Wang, L., Sui, X.H., and Chen, W.X. 2013. High-resolution
 1335 transcriptomic analyses of *Sinorhizobium* sp. NGR234 bacteroids in determinate nodules of
 1336 *Vigna unguiculata* and indeterminate nodules of *Leucaena leucocephala*. *PLOS One* **8**(8):
 1337 e70531. doi:10.1371/journal.pone.0070531.
- 1338 Li, Z., Zu, C., Wang, C., Yang, J., Yu, H., and Wu, H. 2016. Different responses of rhizosphere
 1339 and non-rhizosphere soil microbial communities to consecutive *Piper nigrum* L. monoculture.
 1340 *Sci. Rep.* **6**(1): 35825. doi:10.1038/srep35825.
- 1341 Liu, L.X., Li, Q.Q., Zhang, Y.Z., Hu, Y., Jiao, J., Guo, H.J., Zhang, X.X., Zhang, B., Chen, W.X.,
 1342 and Tian, C.F. 2017a. The nitrate-reduction gene cluster components exert lineage-dependent

- 1343 contributions to optimization of *Sinorhizobium* symbiosis with soybeans. Environ. Microbiol.
 1344 **19**(12): 4926–4938. doi:10.1111/1462-2920.13948.
- 1345 Liu, T., Tian, C.F., and Chen, W.X. 2015. Site-specific Ser/Thr/Tyr phosphoproteome of
 1346 *Sinorhizobium meliloti* at stationary phase. PLOS One **10**(9): e0139143.
 1347 doi:10.1371/journal.pone.0139143.
- 1348 Liu, X., Luo, Y., Mohamed, O.A., Liu, D., and Wei, G. 2014. Global transcriptome analysis of
 1349 *Mesorhizobium alhagi* CCNWXJ12-2 under salt stress. BMC Microbiol. **14**(1): 1.
 1350 doi:10.1186/s12866-014-0319-y.
- 1351 Liu, Y., Jiang, X., Guan, D., Zhou, W., Ma, M., Zhao, B., Cao, F., Li, L., and Li, J. 2017b.
 1352 Transcriptional analysis of genes involved in competitive nodulation in *Bradyrhizobium*
 1353 *diazoefficiens* at the presence of soybean root exudates. Sci. Rep. **7**(1): 10946.
 1354 doi:10.1038/s41598-017-11372-0.
- 1355 Long, S., Reed, J.W., Himawan, J., and Walker, G.C. 1988. Genetic analysis of a cluster of genes
 1356 required for synthesis of the calcofluor-binding exopolysaccharide of *Rhizobium meliloti*. J.
 1357 Bacteriol. **170**(9): 4239–4248. doi:10.1128/jb.170.9.4239-4248.1988.
- 1358 Long, S.R. 1996. Rhizobium symbiosis: nod factors in perspective. Plant Cell **8**(10): 1885–1898.
 1359 doi:10.1105/tpc.8.10.1885.
- 1360 López-Lara, I.M., Gao, J.-L., Soto, M.J., Solares-Pérez, A., Weissenmayer, B., Sohlenkamp, C.,
 1361 Verroios, G.P., Thomas-Oates, J., and Geiger, O. 2005. Phosphorus-free membrane lipids of
 1362 *Sinorhizobium meliloti* are not required for the symbiosis with alfalfa but contribute to
 1363 increased cell yields under phosphorus-limiting conditions of growth. Mol. Plant Microbe
 1364 Interact. **18**(9): 973–982.. doi:10.1094/MPMI-18-0973.

- 1365 López-Torrejón, G., Jiménez-Vicente, E., Buesa, J.M., Hernandez, J.A., Verma, H.K., and Rubio,
 1366 L.M. 2016. Expression of a functional oxygen-labile nitrogenase component in the
 1367 mitochondrial matrix of aerobically grown yeast. *Nat. Commun.* **7**: 11426.
 1368 doi:10.1038/ncomms11426.
- 1369 Lu, M., Jiao, S., Gao, E., Song, X., Li, Z., Hao, X., Rensing, C., and Wei, G. 2017. Transcriptome
 1370 response to heavy metals in *Sinorhizobium meliloti* CCNWSX0020 reveals new metal
 1371 resistance determinants that also promote bioremediation by *Medicago lupulina* in metal
 1372 contaminated soil. *Appl. Environ. Microbiol.* **83**(20): e01244–17. doi:10.1128/AEM.01244-
 1373 17.
- 1374 Lupwayi, N.Z., Clayton, G.W., and Rice, W.A. 2006. Rhizobial inoculants for legume crops. *J.*
 1375 *Crop Im.* **15**(2): 289–321. doi:10.1300/J411v15n02_09.
- 1376 MacLean, A.M., Finan, T.M., and Sadowsky, M.J. 2007. Genomes of the symbiotic nitrogen-
 1377 fixing bacteria of legumes. *Plant Physiol.* **144**(2): 615–622. doi:10.1104/pp.107.101634.
- 1378 Mandal, S., Mandal, M., Pati, B., Das, A., and Ghosh, A. 2009. Proteomics view of a *Rhizobium*
 1379 isolate response to arsenite [As(III)] stress. *Acta Microbiol. Immunol. Hung.* **56**(2): 157–167.
 1380 doi:10.1556/AMicr.56.2009.2.4.
- 1381 Marchetti, M., Capela, D., Glew, M., Cruveiller, S., Chane-Woon-Ming, B., Gris, C., Timmers,
 1382 T., Poinot, V., Gilbert, L.B., Heeb, P., Médigue, C., Batut, J., and Masson-Boivin, C. 2010.
 1383 Experimental evolution of a plant pathogen into a legume symbiont. *PLOS Biol.* **8**(1):
 1384 e1000280–10. doi:10.1371/journal.pbio.1000280.
- 1385 Marchetti, M., Clerissi, C., Yousfi, Y., Gris, C., Bouchez, O., Rocha, E., Cruveiller, S., Jauneau,
 1386 A., Capela, D., and Masson-Boivin, C. 2017. Experimental evolution of rhizobia may lead to

- 1387 either extra- or intracellular symbiotic adaptation depending on the selection regime. Mol.
 1388 Ecol. **26**(7): 1818–1831. doi:10.1111/mec.13895.
- 1389 Marchetti, M., Jauneau, A., Capela, D., Remigi, P., Gris, C., Batut, J., and Masson-Boivin, C.
 1390 2014. Shaping bacterial symbiosis with legumes by experimental evolution. Mol. Plant
 1391 Microbe Interact. **27**(9): 956–964. doi:10.1094/MPMI-03-14-0083-R.
- 1392 Martínez, E., Palacios, R., and Sánchez, F. 1987. Nitrogen-fixing nodules induced by
 1393 *Agrobacterium tumefaciens* harboring *Rhizobium phaseoli* plasmids. J. Bacteriol. **169**(6):
 1394 2828–2834. doi:10.1128/jb.169.6.2828-2834.1987.
- 1395 Martínez-Salazar, J.M., Salazar, E., Encarnación, S., Ramírez-Romero, M.A., and Rivera, J. 2009.
 1396 Role of the extracytoplasmic function sigma factor RpoE4 in oxidative and osmotic stress
 1397 responses in *Rhizobium etli*. J. Bacteriol. **191**(13): 4122–4132. doi:10.1128/JB.01626-08.
- 1398 Marx, H., Minogue, C.E., Jayaraman, D., Richards, A.L., Kwiecien, N.W., Siahipirani, A.F.,
 1399 Rajasekar, S., Maeda, J., Garcia, K., Del Valle-Echevarria, A.R., Volkening, J.D., Westphall,
 1400 M.S., Roy, S., Sussman, M.R., Ané, J.-M., and Coon, J.J. 2016. A proteomic atlas of the
 1401 legume *Medicago truncatula* and its nitrogen-fixing endosymbiont *Sinorhizobium meliloti*.
 1402 Nat. Biotechnol. **34**(11): 1198–1205. doi:10.1038/nbt.3681.
- 1403 Masson-Boivin, C., Giraud, E., Perret, X., and Batut, J. 2009. Establishing nitrogen-fixing
 1404 symbiosis with legumes: how many rhizobium recipes? Trends Microbiol. **17**(10): 458–466.
 1405 doi:10.1016/j.tim.2009.07.004.
- 1406 Mathis, J.N., Barbour, W.M., and Elkan, G.H. 1985. Effect of sym plasmid curing on symbiotic
 1407 effectiveness in *Rhizobium fredii*. Appl. Environ. Microbiol. **49**(6): 1385–1388.
- 1408 Mauchline, T.H., Fowler, J.E., East, A.K., Sartor, A.L., Zaheer, R., Hosie, A.H.F., Poole, P.S., and
 1409 Finan, T.M. 2006. Mapping the *Sinorhizobium meliloti* 1021 solute-binding protein-dependent

- 1410 transportome. Proc. Natl. Acad. Sci. U.S.A. **103**(47): 17933–17938.
 1411 doi:10.1073/pnas.0606673103.
- 1412 Maunoury, N., Redondo-Nieto, M., Bourcy, M., Van de Velde, W., Alunni, B., Laporte, P.,
 1413 Durand, P., Agier, N., Marisa, L., Vaubert, D., Delacroix, H., Duc, G., Ratet, P., Aggerbeck,
 1414 L., Kondorosi, E., and Mergaert, P. 2010. Differentiation of symbiotic cells and
 1415 endosymbionts in *Medicago truncatula* nodulation are coupled to two transcriptome-switches.
 1416 PLOS One **5**(3): e9519. doi:10.1371/journal.pone.0009519.
- 1417 Maynaud, G., Brunel, B., Mornico, D., Durot, M., Severac, D., Dubois, E., Navarro, E., Cleyet-
 1418 Marel, J.-C., and Le Quéré, A. 2013. Genome-wide transcriptional responses of two metal-
 1419 tolerant symbiotic *Mesorhizobium* isolates to zinc and cadmium exposure. BMC Genomics
 1420 **14**(1): 292. doi:10.1186/1471-2164-14-292.
- 1421 Meade, H.M., Long, S.R., Ruvkun, G.B., Brown, S.E., and Ausubel, F.M. 1982. Physical and
 1422 genetic characterization of symbiotic and auxotrophic mutants of *Rhizobium meliloti* induced
 1423 by transposon Tn5 mutagenesis. J. Bacteriol. **149**(1): 114-122.
- 1424 Mendum, T.A., Clark, I.M., and Hirsch, P.R. 2001. Characterization of two novel *Rhizobium*
 1425 *leguminosarum* bacteriophages from a field release site of genetically-modified rhizobia.
 1426 Antonie Van Leeuwenhoek **79**(2): 189–197. doi:10.1023/A:1010238412538.
- 1427 Meneses, N., Mendoza-Hernández, G., and Encarnación, S. 2010. The extracellular proteome of
 1428 *Rhizobium etli* CE3 in exponential and stationary growth phase. Proteome Sci. **8**(1): 51.
 1429 doi:10.1186/1477-5956-8-51.
- 1430 Meneses, N., Taboada, H., Dunn, M.F., Vargas, M.D.C., Buchs, N., Heller, M., and Encarnación,
 1431 S. 2017. The naringenin-induced exoproteome of *Rhizobium etli* CE3. Arch. Microbiol.
 1432 **199**(5): 737–755. doi:10.1007/s00203-017-1351-8.

- 1433 Mergaert, P., Nikovics, K., Kelemen, Z., Maunoury, N., Vaubert, D., Kondorosi, A., and
 1434 Kondorosi, E. 2003. A novel family in *Medicago truncatula* consisting of more than 300
 1435 nodule-specific genes coding for small, secreted polypeptides with conserved cysteine motifs.
 1436 Plant Physiol. **132**(1): 161–173. doi:10.1104/pp.102.018192.
- 1437 Mesa, S., Hauser, F., Friberg, M., Malaguti, E., Fischer, H.-M., and Hennecke, H. 2008.
 1438 Comprehensive assessment of the regulons controlled by the FixLJ-FixK2-FixK1 cascade in
 1439 *Bradyrhizobium japonicum*. J. Bacteriol. **190**(20): 6568–6579. doi:10.1128/JB.00748-08.
- 1440 Miao, J., Zhang, N., Liu, H., Wang, H., Zhong, Z., and Zhu, J. 2018. Soil commensal rhizobia
 1441 promote *Rhizobium etli* nodulation efficiency through CinR-mediated quorum sensing. Arch.
 1442 Microbiol. **27**: 462. doi:10.1007/s00203-018-1478-2.
- 1443 Milunovic, B., diCenzo, G.C., Morton, R.A., and Finan, T.M. 2014. Cell growth inhibition upon
 1444 deletion of four toxin-antitoxin loci from the megaplasmids of *Sinorhizobium meliloti*. J.
 1445 Bacteriol. **196**(4): 811–824. doi:10.1128/JB.01104-13.
- 1446 Mitsch, M.J., diCenzo, G.C., Cowie, A., and Finan, T.M. 2018. Succinate transport is not essential
 1447 for symbiotic nitrogen fixation by *Sinorhizobium meliloti* nor *Rhizobium leguminosarum*.
 1448 Appl. Environ. Microbiol. **84**(1): e01561–17. doi:10.1128/AEM.01561-17.
- 1449 Moënné-Loccoz, Y., Baldani, J.I., and Weaver, R.W. 1995. Sequential heat-curing of
 1450 Tn5-Mob-sac labelled plasmids from *Rhizobium* to obtain derivatives with various
 1451 combinations of plasmids and no plasmid. Lett. Appl. Microbiol. **20**(3): 175–179.
 1452 doi:10.1111/j.1472-765X.1995.tb00420.x.
- 1453 Moënné-Loccoz, Y., and Weaver, R.W. 1995a. Plasmids and saprophytic growth of *Rhizobium*
 1454 *leguminosarum* bv. *trifolii* W14-2 in soil. FEMS Microbiol. Ecol. **18**(2): 139-144.
 1455 doi:10.1111/j.1574-6941.1995.tb00171.x.

- 1456 Moënne-Loccoz, Y., and Weaver, R.W. 1995b. Plasmids influence growth of rhizobia in the
 1457 rhizosphere of clover. *Soil Biol. Biochem.* **27**(8): 1001–1004. doi:10.1016/0038-
 1458 0717(95)00035-D.
- 1459 Morrison, N.A., Hau, C.Y., Trinick, M.J., Shine, J., and Rolfe, B.G. 1983. Heat curing of a sym
 1460 plasmid in a fast-growing *Rhizobium* sp. that is able to nodulate legumes and the nonlegume
 1461 *Parasponia* sp. *J. Bacteriol.* **153**(1): 527–531.
- 1462 Mozo, T., Cabrera, E., and Ruiz-Argüeso, T. 1988. Diversity of plasmid profiles and conservation
 1463 of symbiotic nitrogen fixation genes in newly isolated *Rhizobium* strains nodulating sulla
 1464 (*Hedysarum coronarium* L.). *Appl. Environ. Microbiol.* **54**(5): 1262–1267.
- 1465 Mus, F., Crook, M.B., Garcia, K., Garcia Costas, A., Geddes, B.A., Kouri, E.D., Paramasivan, P.,
 1466 Ryu, M.-H., Oldroyd, G.E.D., Poole, P.S., Udvardi, M.K., Voigt, C.A., Ané, J.-M., and Peters,
 1467 J.W. 2016. Symbiotic nitrogen fixation and the challenges to its extension to nonlegumes.
 1468 *Appl. Environ. Microbiol.* **82**(13): 3698–3710. doi:10.1128/AEM.01055-16.
- 1469 Mutuma, S.P., Okello, J.J., Karanja, N.K., and Woomer, P.I. 2014. Smallholder farmers' use and
 1470 profitability of legume inoculants in western Kenya. *Afr. Crop Sci. J.* **22**(3): 205–214.
 1471 doi:10.4314/acsj.v22i3.
- 1472 Münchbach, M., Dainese, P., Staudenmann, W., Narberhaus, F., and James, P. 1999. Proteome
 1473 analysis of heat shock protein expression in *Bradyrhizobium japonicum*. *Eur. J. Biochem.*
 1474 **264**(1): 39–48. doi:10.1046/j.1432-1327.1999.00567.x.
- 1475 Nakatsukasa, H., Uchiumi, T., Kucho, K.-I., Suzuki, A., Higashi, S., and Abe, M. 2008.
 1476 Transposon mediation allows a symbiotic plasmid of *Rhizobium leguminosarum* bv. *trifolii* to
 1477 become a symbiosis island in *Agrobacterium* and *Rhizobium*. *J. Gen. Appl. Microbiol.* **54**(2):
 1478 107–118. doi:10.2323/jgam.54.107.

- 1479 Nambu, M., Tatsukami, Y., Morisaka, H., Kuroda, K., and Ueda, M. 2015. Quantitative time-
 1480 course proteome analysis of *Mesorhizobium loti* during nodule maturation. J. Proteom. **125**:
 1481 112–120. doi:10.1016/j.jprot.2015.04.034.
- 1482 Nandasena, K.G., O’Hara, G.W., Tiwari, R.P., Sezmiş, E., and Howieson, J.G. 2007. *In situ* lateral
 1483 transfer of symbiosis islands results in rapid evolution of diverse competitive strains of
 1484 mesorhizobia suboptimal in symbiotic nitrogen fixation on the pasture legume *Biserrula*
 1485 *pelecinus* L. Environ. Microbiol. **9**(10): 2496-2511. doi:10.1111/j.1462-2920.2007.01368.x.
- 1486 Natera, S.H., Guerreiro, N., and Djordjevic, M.A. 2000. Proteome analysis of differentially
 1487 displayed proteins as a tool for the investigation of symbiosis. Mol. Plant Microbe Interact.
 1488 **13**(9): 995–1009. doi:10.1094/MPMI.2000.13.9.995.
- 1489 Ndungu, S.M., Messmer, M.M., Ziegler, D., Thuita, M., Vanlauwe, B., Frossard, E., and Thonar,
 1490 C. 2018. Evaluation of MALDI-TOF mass spectrometry for the competitiveness analysis of
 1491 selected indigenous cowpea (*Vigna unguiculata* L. Walp.) *Bradyrhizobium* strains from
 1492 Kenya. Appl. Microbiol. Biotechnol. **7**(8): 988. doi:10.1007/s00253-018-9005-6.
- 1493 Nguyen, H.P., Miwa, H., Kaneko, T., Sato, S., and Okazaki, S. 2017. Identification of
 1494 *Bradyrhizobium elkanii* genes involved in incompatibility with *Vigna radiata*. Genes **8**(12):
 1495 374. doi:10.3390/genes8120374.
- 1496 Nogales, J., Domínguez-Ferreras, A., Amaya-Gomez, C.V., van Dillewijn, P., Cuellar, V.,
 1497 Sanjuán, J., Olivares, J., and Soto, M.J. 2010. Transcriptome profiling of a *Sinorhizobium*
 1498 *meliloti* *fadD* mutant reveals the role of rhizobactin 1021 biosynthesis and regulation genes in
 1499 the control of swarming. BMC Genomics **11**(1): 157. doi:10.1186/1471-2164-11-157.

- 1500 Novikova, N., and Safronova, V. 1992. Transconjugants of *Agrobacterium radiobacter* harbouring
1501 sym genes of *Rhizobium galegae* can form an effective symbiosis with *Medicago sativa*.
1502 FEMS Microbio. Lett. **93**(3): 261–268. doi:10.1111/j.1574-6968.1992.tb05107.x.
- 1503 O'Callaghan, M. 2016. Microbial inoculation of seed for improved crop performance: issues and
1504 opportunities. Appl. Microbiol. Biotechnol. **100**(13): 5729–5746. doi:10.1007/s00253-016-
1505 7590-9.
- 1506 Ogden, A.J., Gargouri, M., Park, J., Gang, D.R., and Kahn, M.L. 2017. Integrated analysis of zone-
1507 specific protein and metabolite profiles within nitrogen-fixing *Medicago truncatula*-
1508 *Sinorhizobium medicae* nodules. PLOS One **12**(7): e0180894.
1509 doi:10.1371/journal.pone.0180894.
- 1510 Oger, E., Marino, D., Guigonis, J.-M., Pauly, N., and Puppo, A. 2012. Sulfenylated proteins in the
1511 *Medicago truncatula*-*Sinorhizobium meliloti* symbiosis. J. Proteom. **75**(13): 4102–4113.
1512 doi:10.1016/j.jprot.2012.05.024.
- 1513 Okazaki, S., Zehner, S., Hempel, J., Lang, K., and Göttfert, M. 2009. Genetic organization and
1514 functional analysis of the type III secretion system of *Bradyrhizobium elkanii*. FEMS
1515 Microbio. Lett. **295**(1): 88–95. doi:10.1111/j.1574-6968.2009.01593.x.
- 1516 Oke, V., and Long, S.R. 1999. Bacterial genes induced within the nodule during the *Rhizobium*-
1517 legume symbiosis. Mol. Microbiol. **32**(4): 837–849. doi:10.1046/j.1365-2958.1999.01402.x.
- 1518 Oldroyd, G.E., and Dixon, R. 2014. Biotechnological solutions to the nitrogen problem. Curr.
1519 Opin. Biotech. **26**: 19–24. doi:10.1016/j.copbio.2013.08.006.
- 1520 Oldroyd, G.E.D. 2013. Speak, friend, and enter: signalling systems that promote beneficial
1521 symbiotic associations in plants. Nat. Rev. Microbiol. **11**(4): 252–263.
1522 doi:10.1038/nrmicro2990.

- 1523 Oldroyd, G.E.D., and Downie, J.A. 2008. Coordinating nodule morphogenesis with rhizobial
 1524 infection in legumes. *Annu. Rev. Plant Biol.* **59**(1): 519–546.
 1525 doi:10.1146/annurev.arplant.59.032607.092839.
- 1526 Oldroyd, G.E.D., Murray, J.D., Poole, P.S., and Downie, J.A. 2011. The rules of engagement in
 1527 the legume-rhizobial symbiosis. *Annu. Rev. Genet.* **45**(1): 119–144. doi:10.1146/annurev-
 1528 genet-110410-132549.
- 1529 Oresnik, I.J., Liu, S.L., Yost, C.K., and Hynes, M.F. 2000. Megaplasmid pRme2011a of
 1530 *Sinorhizobium meliloti* is not required for viability. *J. Bacteriol.* **182**(12): 3582–3586.
 1531 doi:10.1128/JB.182.12.3582-3586.2000.
- 1532 Oresnik, I.J., Twelker, S., and Hynes, M.F. 1999. Cloning and characterization of a *Rhizobium*
 1533 *leguminosarum* gene encoding a bacteriocin with similarities to RTX toxins. *Appl. Environ.*
 1534 *Microbiol.* **65**(7): 2833–2840.
- 1535 Orikasa, Y., Nodasaka, Y., Ohyama, T., Okuyama, H., Ichise, N., Yumoto, I., Morita, N., Wei,
 1536 M., and Ohwada, T. 2010. Enhancement of the nitrogen fixation efficiency of genetically-
 1537 engineered *Rhizobium* with high catalase activity. *J. Biosci. Bioeng.* **110**(4): 397–402.
 1538 doi:10.1016/j.jbiosc.2010.04.007.
- 1539 Paau, A.S. 1991. Improvement of *Rhizobium* inoculants by mutation, genetic engineering and
 1540 formulation. *Biotechnol. Adv.* **9**(2): 173–184.
- 1541 Page, A.J., Cummins, C.A., Hunt, M., Wong, V.K., Reuter, S., Holden, M.T.G., Fookes, M.,
 1542 Falush, D., Keane, J.A., and Parkhill, J. 2015. Roary: rapid large-scale prokaryote pan genome
 1543 analysis. *Bioinformatics* **31**(22): 3691–3693. doi:10.1093/bioinformatics/btv421.

- 1544 Pahua, V.J., Stokes, P.J.N., Hollowell, A.C., Regus, J.U., Gano-Cohen, K.A., Wendlandt, C.E.,
 1545 Quides, K.W., Lyu, J.Y., and Sachs, J.L. 2018. Fitness variation among host species and the
 1546 paradox of ineffective rhizobia. *J. Evol. Biol.* **31**(4): 599–610. doi:10.1111/jeb.13249.
- 1547 Pan, H., and Wang, D. 2017. Nodule cysteine-rich peptides maintain a working balance during
 1548 nitrogen-fixing symbiosis. *Nat. Plants* **3**(5): 17048. doi:10.1038/nplants.2017.48.
- 1549 Pankhurst, C.E., Macdonald, P.E., and Reeves, J.M. 1986. Enhanced nitrogen fixation and
 1550 competitiveness for nodulation of *Lotus pedunculatus* by a plasmid-cured derivative of
 1551 *Rhizobium loti*. *Microbiology* **132**(8): 2321–2328. doi:10.1099/00221287-132-8-2321.
- 1552 Park, S., Croteau, P., and Boering, K.A. 2012. Trends and seasonal cycles in the isotopic
 1553 composition of nitrous oxide since 1940. *Nat. Geosci.* **5**(4): 261–265.
- 1554 Paton, G.I., Palmer, G., Burton, M., Rattray, E.A., McGrath, S.P., Glover, L.A., and Killham, K.
 1555 1997. Development of an acute and chronic ecotoxicity assay using *lux*-marked *Rhizobium*
 1556 *leguminosarum* biovar *trifolii*. *Lett. Appl. Microbiol.* **24**(4): 296–300. doi:10.1046/j.1472-
 1557 765X.1997.00071.x.
- 1558 Pech-Canul, Á., Nogales, J., Miranda-Molina, A., Álvarez, L., Geiger, O., Soto, M.J., and López-
 1559 Lara, I.M. 2011. FadD is required for utilization of endogenous fatty acids released from
 1560 membrane lipids. *J. Bacteriol.* **193**(22): 6295–6304. doi:10.1128/JB.05450-11.
- 1561 Penterman, J., Abo, R.P., De Nisco, N.J., Arnold, M.F.F., Longhi, R., Zanda, M., and Walker,
 1562 G.C. 2014. Host plant peptides elicit a transcriptional response to control the *Sinorhizobium*
 1563 *meliloti* cell cycle during symbiosis. *Proc. Natl. Acad. Sci. U.S.A.* **111**(9): 3561–3566.
 1564 doi:10.1073/pnas.1400450111.
- 1565 Penttinen, P., Räsänen, L.A., Lortet, G., and Lindström, K. 2013. Stable isotope labelling reveals
 1566 that NaCl stress decreases the production of *Ensifer* (*Sinorhizobium*) *arboris*

- lipochitooligosaccharide signalling molecules. FEMS Microbio. Lett. **349**(2): 117–126.
doi:10.1111/1574-6968.12303.
- Peralta, H., Aguilar, A., Díaz, R., Mora, Y., Martínez-Batallar, G., Salazar, E., Vargas-Lagunas, C., Martínez, E., Encarnación, S., Girard, L., and Mora, J. 2016. Genomic studies of nitrogen-fixing rhizobial strains from *Phaseolus vulgaris* seeds and nodules. BMC Genomics **17**(1): 711. doi:10.1186/s12864-016-3053-z.
- Perry, B.J., Akter, M.S., and Yost, C.K. 2016. The use of transposon insertion sequencing to interrogate the core functional genome of the legume symbiont *Rhizobium leguminosarum*. Front. Microbiol. **7**: 1873. doi:10.3389/fmicb.2016.01873.
- Perry, B.J., and Yost, C.K. 2014. Construction of a *mariner*-based transposon vector for use in insertion sequence mutagenesis in selected members of the *Rhizobiaceae*. BMC Microbiol. **14**(1): 298. doi:10.1186/s12866-014-0298-z.
- Pessi, G., Ahrens, C.H., Rehrauer, H., Lindemann, A., Hauser, F., Fischer, H.-M., and Hennecke, H. 2007. Genome-wide transcript analysis of *Bradyrhizobium japonicum* bacteroids in soybean root nodules. Mol. Plant Microbe Interact. **20**(11): 1353–1363. doi:10.1094/MPMI-20-11-1353.
- Pérez Carrascal, O.M., VanInsberghe, D., Juárez, S., Polz, M.F., Vinuesa, P., and González, V. 2016. Population genomics of the symbiotic plasmids of sympatric nitrogen-fixing *Rhizobium* species associated with *Phaseolus vulgaris*. Environ. Microbiol. **18**(8): 2660–2676. doi:10.1111/1462-2920.13415.
- Pérez-Giménez, J., Covelli, J.M., López, M.F., Althabegoiti, M.J., Ferrer-Navarro, M., Mongiardini, E.J., and Lodeiro, A.R. 2012. Soybean seed lectin prevents the accumulation of S-adenosyl methionine synthetase and the S1 30S ribosomal protein in *Bradyrhizobium*

- 1590 *japonicum* under C and N starvation. Curr. Microbiol. **65**(4): 465–474. doi:10.1007/s00284-
1591 012-0180-x.
- 1592 Pérez-Montaña, F., Del Cerro, P., Jiménez-Guerrero, I., López-Baena, F.J., Cubo, M.T., Hungria,
1593 M., Megías, M., and Ollero, F.J. 2016a. RNA-seq analysis of the *Rhizobium tropici* CIAT 899
1594 transcriptome shows similarities in the activation patterns of symbiotic genes in the presence
1595 of apigenin and salt. BMC Genomics **17**(1): 198. doi:10.1186/s12864-016-2543-3.
- 1596 Pérez-Montaña, F., Jiménez-Guerrero, I., Acosta-Jurado, S., Navarro-Gómez, P., Ollero, F.J.,
1597 Ruiz-Sainz, J.E., López-Baena, F.J., and Vinardell, J.M. 2016b. A transcriptomic analysis of
1598 the effect of genistein on *Sinorhizobium fredii* HH103 reveals novel rhizobial genes putatively
1599 involved in symbiosis. Sci. Rep. **6**(1): 31592. doi:10.1038/srep31592.
- 1600 Pfau, T., Christian, N., Masakapalli, S.K., Sweetlove, L.J., Poolman, M.G., and Ebenhoeh, O.
1601 2016. The intertwined metabolism of *Medicago truncatula* and its nitrogen fixing symbiont
1602 *Sinorhizobium meliloti* elucidated by genome-scale metabolic models. bioRxiv: **067348**.
1603 doi:10.1101/067348.
- 1604 Pini, F., De Nisco, N.J., Ferri, L., Penterman, J., Fioravanti, A., Brilli, M., Mengoni, A.,
1605 Bazzicalupo, M., Viollier, P.H., Walker, G.C., and Biondi, E.G. 2015. Cell cycle control by
1606 the master regulator CtrA in *Sinorhizobium meliloti*. PLOS Genet. **11**(5): e1005232.
1607 doi:10.1371/journal.pgen.1005232.
- 1608 Pini, F., East, A.K., Appia-Ayme, C., Tomek, J., Karunakaran, R., Mendoza-Suárez, M., Edwards,
1609 A., Terpolilli, J.J., Roworth, J., Downie, J.A., and Poole, P.S. 2017. Bacterial biosensors for
1610 *in vivo* spatiotemporal mapping of root secretion. Plant Physiol. **174**(3): 1289–1306.
1611 doi:10.1104/pp.16.01302.

- 1612 Pini, F., Galardini, M., Bazzicalupo, M., and Mengoni, A. 2011. Plant-bacteria association and
 1613 symbiosis: are there common genomic traits in *Alphaproteobacteria*? *Genes* **2**(4): 1017–1032.
 1614 doi:10.3390/genes2041017.
- 1615 Pobigaylo, N., Szymczak, S., Nattkemper, T.W., and Becker, A. 2008. Identification of genes
 1616 relevant to symbiosis and competitiveness in *Sinorhizobium meliloti* using signature-tagged
 1617 mutants. *Mol. Plant Microbe Interact.* **21**(2): 219–231. doi:10.1094/MPMI-21-2-0219.
- 1618 Pobigaylo, N., Wetter, D., Szymczak, S., Schiller, U., Kurtz, S., Meyer, F., Nattkemper, T.W., and
 1619 Becker, A. 2006. Construction of a large signature-tagged mini-Tn5 transposon library and its
 1620 application to mutagenesis of *Sinorhizobium meliloti*. *Appl. Environ. Microbiol.* **72**(6): 4329–
 1621 4337. doi:10.1128/AEM.03072-05.
- 1622 Poole, P., Ramachandran, V., and Terpolilli, J. 2018. Rhizobia: from saprophytes to
 1623 endosymbionts. *Nat. Rev. Microbiol.* **16**(5): 291–303. doi:10.1038/nrmicro.2017.171.
- 1624 Porter, S.S., Chang, P.L., Conow, C.A., Dunham, J.P., and Friesen, M.L. 2017. Association
 1625 mapping reveals novel serpentine adaptation gene clusters in a population of symbiotic
 1626 *Mesorhizobium*. *ISME J.* **11**(1): 248–262. doi:10.1038/ismej.2016.88.
- 1627 Price, P.A., Tanner, H.R., Dillon, B.A., Shabab, M., Walker, G.C., and Griffiths, J.S. 2015.
 1628 Rhizobial peptidase HrrP cleaves host-encoded signaling peptides and mediates symbiotic
 1629 compatibility. *Proc. Natl. Acad. Sci. U.S.A.* **112**(49): 15244–15249.
 1630 doi:10.1073/pnas.1417797112.
- 1631 Quandt, J., and Hynes, M.F. 1993. Versatile suicide vectors which allow direct selection for gene
 1632 replacement in Gram-negative bacteria. *Gene* **127**(1): 15–21. doi:10.1016/0378-
 1633 1119(93)90611-6.

- 1634 Queiroux, C., Washburn, B.K., Davis, O.M., Stewart, J., Brewer, T.E., Lyons, M.R., and Jones,
1635 K.M. 2012. A comparative genomics screen identifies a *Sinorhizobium meliloti* 1021 *sodM*-
1636 like gene strongly expressed within host plant nodules. BMC Microbiol. **12**(1): 74.
1637 doi:10.1186/1471-2180-12-74.
- 1638 Quides, K.W., Stomackin, G.M., Lee, H.-H., Chang, J.H., and Sachs, J.L. 2017. *Lotus japonicus*
1639 alters *in planta* fitness of *Mesorhizobium loti* dependent on symbiotic nitrogen fixation. PLOS
1640 One **12**(9): e0185568. doi:10.1371/journal.pone.0185568.
- 1641 Rachwał, K., Matczyńska, E., and Janczarek, M. 2015. Transcriptome profiling of a *Rhizobium*
1642 *leguminosarum* bv. *trifolii* *rosR* mutant reveals the role of the transcriptional regulator RosR
1643 in motility, synthesis of cell-surface components, and other cellular processes. BMC
1644 Genomics **16**(1): 1111. doi:10.1186/s12864-015-2332-4.
- 1645 Ramachandran, V.K., East, A.K., Karunakaran, R., Downie, J.A., and Poole, P.S. 2011. Adaptation
1646 of *Rhizobium leguminosarum* to pea, alfalfa and sugar beet rhizospheres investigated by
1647 comparative transcriptomics. Genome Biol. **12**(10): R106. doi:10.1186/gb-2011-12-10-r106.
- 1648 Ramírez, M., Valderrama, B., Arredondo-Peter, R., Soberón, M., Mora, J., and Hernández, G.
1649 1999. *Rhizobium etli* genetically engineered for the heterologous expression of *Vitreoscilla* sp.
1650 hemoglobin: effects on free-living and symbiosis. Mol. Plant Microbe Interact. **12**(11): 1008–
1651 1015. doi:10.1094/MPMI.1999.12.11.1008.
- 1652 Ramsay, J.P., Sullivan, J.T., Stuart, G.S., Lamont, I.L., and Ronson, C.W. 2006. Excision and
1653 transfer of the *Mesorhizobium loti* R7A symbiosis island requires an integrase IntS, a novel
1654 recombination directionality factor RdfS, and a putative relaxase RlxS. Mol. Microbiol. **62**(3):
1655 723–734. doi:10.1111/j.1365-2958.2006.05396.x.

- 1656 Reeve, W.G., Tiwari, R.P., Guerreiro, N., Stubbs, J., Dilworth, M.J., Glenn, A.R., Rolfe, B.G.,
 1657 Djordjevic, M.A., and Howieson, J.G. 2004. Probing for pH-regulated proteins in
 1658 *Sinorhizobium medicae* using proteomic analysis. J. Mol. Microbiol. Biotechnol. **7**(3): 140–
 1659 147. doi:10.1159/000078657.
- 1660 Remigi, P., Capela, D., Clerissi, C., Tasse, L., Torchet, R., Bouchez, O., Batut, J., Cruveiller, S.,
 1661 Rocha, E.P.C., and Masson-Boivin, C. 2014. Transient hypermutagenesis accelerates the
 1662 evolution of legume endosymbionts following horizontal gene transfer. PLOS Biol. **12**(9):
 1663 e1001942. doi:10.1371/journal.pbio.1001942.
- 1664 Remigi, P., Zhu, J., Young, J.P.W., and Masson-Boivin, C. 2016. Symbiosis within symbiosis:
 1665 Evolving nitrogen-fixing legume symbionts. Trends Microbiol. **24**(1): 63–75.
 1666 doi:10.1016/j.tim.2015.10.007.
- 1667 Resendis-Antonio, O., Hernández, M., Mora, Y., and Encarnación, S. 2012. Functional modules,
 1668 structural topology, and optimal activity in metabolic networks. PLOS Comput. Biol. **8**(10):
 1669 e1002720. doi:10.1371/journal.pcbi.1002720.
- 1670 Resendis-Antonio, O., Hernández, M., Salazar, E., Contreras, S., Batallar, G.M., Mora, Y., and
 1671 Encarnación, S. 2011. Systems biology of bacterial nitrogen fixation: high-throughput
 1672 technology and its integrative description with constraint-based modeling. BMC Syst. Biol.
 1673 **5**(1): 120. doi:10.1186/1752-0509-5-120.
- 1674 Resendis-Antonio, O., Reed, J.L., Encarnación, S., Collado-Vides, J., and Palsson, B.Ø. 2007.
 1675 Metabolic reconstruction and modeling of nitrogen fixation in *Rhizobium etli*. PLOS Comput.
 1676 Biol. **3**(10): 1887–1895. doi:10.1371/journal.pcbi.0030192.

- 1677 Reyes-Pérez, A., Vargas, M.D.C., Hernández, M., Aguirre-von-Wobeser, E., Pérez-Rueda, E., and
 1678 Encarnación, S. 2016. Transcriptomic analysis of the process of biofilm formation in
 1679 *Rhizobium etli* CFN42. Arch. Microbiol. **198**(9): 847–860. doi:10.1007/s00203-016-1241-5.
- 1680 Robleto, E.A., Scupham, A.J., and Triplett, E.W. 1997. Trifolitoxin production in *Rhizobium etli*
 1681 strain CE3 increases competitiveness for rhizosphere colonization and root nodulation of
 1682 *Phaseolus vulgaris* in soil. Mol. Plant Microbe Interact. **10**(2): 228-233.
 1683 doi:10.1094/MPMI.1997.10.2.228.
- 1684 Robleto, E.A., Kmiecik, K. Oplinger, E.S., Neinhuis, J., and Triplett, E.W. 1998. Trifolitoxin
 1685 production increases nodulation competitiveness of *Rhizobium etli* CE3 under agricultural
 1686 conditions. Appl. Environ. Microbiol. **64**(7): 2630-2633.
- 1687 Rodrigues, J.A., López-Baena, F.J., Ollero, F.J., Vinardell, J.M., Espuny, M.D.R., Bellogín, R.A.,
 1688 Ruiz-Sainz, J.E., Thomas, J.R., Sumpton, D., Ault, J., and Thomas-Oates, J. 2007. NopM and
 1689 NopD are rhizobial nodulation outer proteins: identification using LC-MALDI and LC-ESI
 1690 with a monolithic capillary column. J. Proteome Res. **6**(3): 1029–1037.
 1691 doi:10.1021/pr060519f.
- 1692 Rogel, M.A., Hernández-Lucas, I., Kuykendall, L.D., Balkwill, D.L., and Martinez-Romero, E.
 1693 2001. Nitrogen-fixing nodules with *Ensifer adhaerens* harboring *Rhizobium tropici* symbiotic
 1694 plasmids. Appl. Environ. Microbiol. **67**(7): 3264–3268. doi:10.1128/AEM.67.7.3264-
 1695 3268.2001.
- 1696 Rogers, C., and Oldroyd, G.E.D. 2014. Synthetic biology approaches to engineering the nitrogen
 1697 symbiosis in cereals. J. Exp. Bot. **65**(8): 1939–1946. doi:10.1093/jxb/eru098.

- 1698 Rolfe, G., Gresshoff, P.M., and Shine, J. 1980. Rapid screening for symbiotic mutants of
1699 *Rhizobium* and white clover. *Plant Sci. Lett.* **19**(3): 277-284. doi:10.1016/0304-
1700 4211(80)90082-6.
- 1701 Romero, D., Brom, S., Martínez-Salazar, J., Girard, M.L., Palacios, R., and Davila, G. 1991.
1702 Amplification and deletion of a *nod-nif* region in the symbiotic plasmid of *Rhizobium phaseoli*.
1703 *J. Bacteriol.* **173**(8): 2435–2441. doi:10.1128/jb.173.8.2435-2441.1991.
- 1704 Roux, B., Rodde, N., Jardinaud, M.-F., Timmers, T., Sauviac, L., Cottret, L., Carrère, S., Sallet,
1705 E., Courcelle, E., Moreau, S., Debellé, F., Capela, D., de Carvalho-Niebel, F., Gouzy, J.,
1706 Bruand, C., and Gamas, P. 2014. An integrated analysis of plant and bacterial gene expression
1707 in symbiotic root nodules using laser-capture microdissection coupled to RNA sequencing.
1708 *Plant J.* **77**(6): 817–837. doi:10.1111/tpj.12442.
- 1709 Saad, M.M., Kobayashi, H., Marie, C., Brown, I.R., Mansfield, J.W., Broughton, W.J., and
1710 Deakin, W.J. 2005. NopB, a type III secreted protein of *Rhizobium* sp. strain NGR234, is
1711 associated with pilus-like surface appendages. *J. Bacteriol.* **187**(3): 1173–1181.
1712 doi:10.1128/JB.187.3.1173-1181.2005.
- 1713 Saalbach, G., Erik, P., and Wienkoop, S. 2002. Characterisation by proteomics of peribacteroid
1714 space and peribacteroid membrane preparations from pea (*Pisum sativum*) symbiosomes.
1715 *Proteomics* **2**(3): 325–337.
- 1716 Sachs, J.L., Ehinger, M.O., and Simms, E.L. 2010. Origins of cheating and loss of symbiosis in
1717 wild *Bradyrhizobium*. *J. Evol. Biol.* **23**(5): 1075–1089. doi:10.1111/j.1420-
1718 9101.2010.01980.x.
- 1719 Sachs, J.L., Quides, K.W., and Wendlandt, C.E. 2018. Legumes versus rhizobia: a model for
1720 ongoing conflict in symbiosis. *New Phytol.* **29**: 925. doi:10.1111/nph.15222.

- 1721 Salas, M.E., Lozano, M.J., López, J.L., Draghi, W.O., Serrania, J., Torres Tejerizo, G.A., Albicoro,
 1722 F.J., Nilsson, J.F., Pistorio, M., Del Papa, M.F., Parisi, G., Becker, A., and Lagares, A. 2017.
 1723 Specificity traits consistent with legume-rhizobia coevolution displayed by *Ensifer meliloti*
 1724 rhizosphere colonization. *Environ. Microbiol.* **282**: 219. doi:10.1111/1462-2920.13820.
- 1725 Salazar, E., Díaz-Mejía, J.J., Moreno-Hagelsieb, G., Martínez-Batallar, G., Mora, Y., Mora, J., and
 1726 Encarnación, S. 2010. Characterization of the NifA-RpoN regulon in *Rhizobium etli* in free
 1727 life and in symbiosis with *Phaseolus vulgaris*. *Appl. Environ. Microbiol.* **76**(13): 4510–4520.
 1728 doi:10.1128/AEM.02007-09.
- 1729 Santos, R., Hérouart, D., Sigaud, S., Touati, D., and Puppo, A. 2001. Oxidative burst in alfalfa-
 1730 *Sinorhizobium meliloti* symbiotic interaction. *Mol. Plant Microbe Interact.* **14**(1): 86-89.
 1731 doi:10.1094/MPMI.2001.14.1.86.
- 1732 Saramago, M., Peregrina, A., Robledo, M., Matos, R.G., Hilker, R., Serrania, J., Becker, A.,
 1733 Arraiano, C.M., and Jiménez-Zurdo, J.I. 2017. *Sinorhizobium meliloti* YbeY is an
 1734 endoribonuclease with unprecedented catalytic features, acting as silencing enzyme in
 1735 riboregulation. *Nucleic Acids Res.* **45**(3): 1371–1391. doi:10.1093/nar/gkw1234.
- 1736 Sarma, A.D., and Emerich, D.W. 2005. Global protein expression pattern of *Bradyrhizobium*
 1737 *japonicum* bacteroids: a prelude to functional proteomics. *Proteomics* **5**(16): 4170–4184.
 1738 doi:10.1002/pmic.200401296.
- 1739 Sarma, A.D., and Emerich, D.W. 2006. A comparative proteomic evaluation of culture grown vs
 1740 nodule isolated *Bradyrhizobium japonicum*. *Proteomics* **6**(10): 3008–3028.
 1741 doi:10.1002/pmic.200500783.
- 1742 Schellenberger, J., Que, R., Fleming, R.M.T., Thiele, I., Orth, J.D., Feist, A.M., Zielinski, D.C.,
 1743 Bordbar, A., Lewis, N.E., Rahmanian, S., Kang, J., Hyduke, D.R., and Palsson, B.Ø. 2011.

- 1744 Quantitative prediction of cellular metabolism with constraint-based models: the COBRA
 1745 Toolbox v2.0. Nat. Protoc. **6**(9): 1290–1307. doi:10.1038/nprot.2011.308.
- 1746 Schlüter, J.-P., Reinkensmeier, J., Daschkey, S., Evguenieva-Hackenberg, E., Janssen, S., Jänicke,
 1747 S., Becker, J.D., Giegerich, R., and Becker, A. 2010. A genome-wide survey of sRNAs in the
 1748 symbiotic nitrogen-fixing alpha-proteobacterium *Sinorhizobium meliloti*. BMC Genomics
 1749 **11**(1): 245. doi:10.1186/1471-2164-11-245.
- 1750 Schroeder, B.K., House, B.L., Mortimer, M.W., Yurgel, S.N., Maloney, S.C., Ward, K.L., and
 1751 Kahn, M.L. 2005. Development of a functional genomics platform for *Sinorhizobium meliloti*:
 1752 construction of an ORFeome. Appl. Environ. Microbiol. **71**(10): 5858–5864.
 1753 doi:10.1128/AEM.71.10.5858-5864.2005.
- 1754 Serrania, J., Johnner, T., Rupp, O., Goesmann, A., and Becker, A. 2017. Massive parallel insertion
 1755 site sequencing of an arrayed *Sinorhizobium meliloti* signature-tagged mini-Tn5 transposon
 1756 mutant library. J. Biotechnol. doi:10.1016/j.jbiotec.2017.02.019.
- 1757 Seshadri, R., Reeve, W.G., Ardley, J.K., Tennessen, K., Woyke, T., Kyrpides, N.C., and Ivanova,
 1758 N.N. 2015. Discovery of novel plant interaction determinants from the genomes of 163 root
 1759 nodule bacteria. Sci. Rep. **5**(1): 16825. doi:10.1038/srep16825.
- 1760 Shamseldin, A., Nyalwidhe, J., and Werner, D. 2006. A proteomic approach towards the analysis
 1761 of salt tolerance in *Rhizobium etli* and *Sinorhizobium meliloti* strains. Curr. Microbiol. **52**(5):
 1762 333–339. doi:10.1007/s00284-005-6472-7.
- 1763 Sharma, P.K., and Laxminarayana, K. 1989. Effect of high temperature on plasmid curing of
 1764 *Rhizobium* spp. in relation to nodulation of pigeonpea [*Cajanus cajan* (L.) Millsp.]. Biol Fert
 1765 Soils **8**(1): 75–79. doi:10.1007/BF00260520.

- 1766 Silva, C., Kan, F.L., and Martínez-Romero, E. 2007. Population genetic structure of *Sinorhizobium*
 1767 *meliloti* and *S. medicae* isolated from nodules of *Medicago* spp. in Mexico. FEMS Microbiol.
 1768 Ecol. **60**(3): 477–489. doi:10.1111/j.1574-6941.2007.00301.x.
- 1769 Smalla, K., Wieland, G., Buchner, A., Zock, A., Parzy, J., Kaiser, S., Roskot, N., Heuer, H., and
 1770 Berg, G. 2001. Bulk and rhizosphere soil bacterial communities studied by denaturing gradient
 1771 gel electrophoresis: plant-dependent enrichment and seasonal shifts revealed. Appl. Environ.
 1772 Microbiol. **67**(10): 4742–4751. doi:10.1128/AEM.67.10.4742-4751.2001.
- 1773 Sobrero, P., Schlüter, J.-P., Lanner, U., Schlosser, A., Becker, A., and Valverde, C. 2012.
 1774 Quantitative proteomic analysis of the Hfq-regulon in *Sinorhizobium meliloti* 2011. PLOS One
 1775 **7**(10): e48494. doi:10.1371/journal.pone.0048494.
- 1776 Sohlenkamp, C., de Rudder, K.E., Rohrs, V., López-Lara, I.M., and Geiger, O. 2000. Cloning and
 1777 characterization of the gene for phosphatidylcholine synthase. J. Biol. Chem. **275**(25): 18919–
 1778 18925. doi:10.1074/jbc.M000844200.
- 1779 Sohlenkamp, C., de Rudder, K.E.E., and Geiger, O. 2004. Phosphatidylethanolamine is not
 1780 essential for growth of *Sinorhizobium meliloti* on complex culture media. J. Bacteriol. **186**(6):
 1781 1667–1677. doi:10.1128/JB.186.6.1667-1677.2004.
- 1782 Spini, G., Decorosi, F., Cerboneschi, M., Tegli, S., Mengoni, A., Viti, C., and Giovannetti, L.
 1783 2015. Effect of the plant flavonoid luteolin on *Ensifer meliloti* 3001 phenotypic responses.
 1784 Plant Soil **399**(1): 159–178. doi:10.1007/s11104-015-2659-2.
- 1785 Starker, C.G., Parra-Colmenares, A.L., Smith, L., Mitra, R.M., and Long, S.R. 2006. Nitrogen
 1786 fixation mutants of *Medicago truncatula* fail to support plant and bacterial symbiotic gene
 1787 expression. Plant Physiol. **140**(2): 671–680. doi:10.1104/pp.105.072132.

- 1788 Stasiak, G., Mazur, A., Wielbo, J., Marczak, M., Żebracki, K., Koper, P., and Skorupska, A. 2014.
 1789 Functional relationships between plasmids and their significance for metabolism and
 1790 symbiotic performance of *Rhizobium leguminosarum* bv. *trifolii*. J. Appl. Genet. **55**(4): 515–
 1791 527. doi:10.1007/s13353-014-0220-2.
- 1792 Stockwell, S.B., Reutimann, L., and Guerinot, M.L. 2012. A role for *Bradyrhizobium japonicum*
 1793 ECF16 sigma factor EcfS in the formation of a functional symbiosis with soybean. Mol. Plant
 1794 Microbe Interact. **25**(1): 119–128. doi:10.1094/MPMI-07-11-0188.
- 1795 Streeter, J.G. 1994. Failure of inoculant rhizobia to overcome the dominance of indigenous strains
 1796 for nodule formation. Can. J. Microbiol. **40**(7): 513–522. doi:10.1139/m94-084.
- 1797 Strodtman, K.N., Stevenson, S.E., Waters, J.K., Mawhinney, T.P., Thelen, J.J., Polacco, J.C., and
 1798 Emerich, D.W. 2017. The bacteroid periplasm in soybean nodules is an interkingdom
 1799 symbiotic space. Mol. Plant Microbe Interact. **30**(12): 997–1008. doi:10.1094/MPMI-12-16-
 1800 0264-R.
- 1801 Sugawara, M., Epstein, B., Badgley, B.D., Unno, T., Xu, L., Reese, J., Gyaneshwar, P., Denny,
 1802 R., Mudge, J., Bharti, A.K., Farmer, A.D., May, G.D., Woodward, J.E., Médigue, C., Vallenet,
 1803 D., Lajus, A., Rouy, Z., Martinez-Vaz, B., Tiffin, P., Young, N.D., and Sadowsky, M.J. 2013.
 1804 Comparative genomics of the core and accessory genomes of 48 *Sinorhizobium* strains
 1805 comprising five genospecies. Genome Biol. **14**(2): R17. doi:10.1186/gb-2013-14-2-r17.
- 1806 Sullivan, J.T., and Ronson, C.W. 1998. Evolution of rhizobia by acquisition of a 500-kb symbiosis
 1807 island that integrates into a phe-tRNA gene. Proc. Natl. Acad. Sci. U.S.A. **95**(9): 5145–5149.
 1808 doi:10.1073/pnas.95.9.5145.

- 1809 Sullivan, J.T., Patrick, H.N., Lowther, W.L., Scott, D.B., and Ronson, C.W. 1995. Nodulating
1810 strains of *Rhizobium loti* arise through chromosomal symbiotic gene transfer in the
1811 environment. *Proc. Natl. Acad. Sci. U.S.A.* **92**(19): 8985–8989. doi:10.1073/pnas.92.19.8985.
- 1812 Swanson, J.A., Tu, J.K., Ogawa, J., Sanga, R., Fisher, R.F., and Long, S.R. 1987. Extended region
1813 of nodulation genes in *Rhizobium meliloti* 1021. I. Phenotypes of Tn5 insertion mutants.
1814 *Genetics* **117**(2): 181–189.
- 1815 Talebi, M.B., Bahar, M., Saeidi, G., Mengoni, A., and Bazzicalupo, M. 2008. Diversity of
1816 *Sinorhizobium* strains nodulating *Medicago sativa* from different Iranian regions. *FEMS*
1817 *Microbio. Lett.* **288**(1): 40–46. doi:10.1111/j.1574-6968.2008.01329.x.
- 1818 Tanthanuch, W., Tittabutr, P., Mohammed, S., Matthiesen, R., Yamabhai, M., Manassila, M.,
1819 Jensen, O.N., Boonkerd, N., and Teaumroong, N. 2010. Identification of salt-tolerant
1820 *Sinorhizobium* sp. strain BL3 membrane proteins based on proteomics. *Microbes Environ.*
1821 **25**(4): 275–280. doi:10.1264/jsme2.ME09185.
- 1822 Tatsukami, Y., Nambu, M., Morisaka, H., Kuroda, K., and Ueda, M. 2013. Disclosure of the
1823 differences of *Mesorhizobium loti* under the free-living and symbiotic conditions by
1824 comparative proteome analysis without bacteroid isolation. *BMC Microbiol.* **13**(1): 180.
1825 BioMed Central. doi:10.1186/1471-2180-13-180.
- 1826 Terpolilli, J.J., Masakapalli, S.K., Karunakaran, R., Webb, I.U.C., Green, R., Watmough, N.J.,
1827 Kruger, N.J., Ratcliffe, R.G., and Poole, P.S. 2016. Lipogenesis and redox balance in nitrogen-
1828 fixing pea bacteroids. *J. Bacteriol.* **198**(20): 2864–2875. doi:10.1128/JB.00451-16.
- 1829 Thiele, I., and Palsson, B.Ø. 2010. A protocol for generating a high-quality genome-scale
1830 metabolic reconstruction. *Nat. Protoc.* **5**(1): 93–121. doi:10.1038/nprot.2009.203.

- 1831 Tian, C.F., Zhou, Y.J., Zhang, Y.M., Li, Q.Q., Zhang, Y.Z., Li, D.F., Wang, S., Wang, J., Gilbert,
1832 L.B., Li, Y.R., and Chen, W.X. 2012. Comparative genomics of rhizobia nodulating soybean
1833 suggests extensive recruitment of lineage-specific genes in adaptations. *Proc. Natl. Acad. Sci.*
1834 U.S.A. **109**(22): 8629–8634. doi:10.1073/pnas.1120436109.
- 1835 Tian, Z., Zou, H., Li, J., Zhang, Y., Liu, Y., Yu, G., Zhu, J., Rüberg, S., Becker, A., and Wang, Y.
1836 2006. Transcriptome analysis of *Sinorhizobium meliloti* nodule bacteria in *nifA* mutant
1837 background. *Chinese Sci. Bull.* **51**(17): 2079–2086. doi:10.1007/s11434-006-2092-2.
- 1838 Tiricz, H., Szűcs, A., Farkas, A., Pap, B., Lima, R.M., Maróti, G., Kondorosi, E., and Kereszt, A.
1839 2013. Antimicrobial nodule-specific cysteine-rich peptides induce membrane depolarization-
1840 associated changes in the transcriptome of *Sinorhizobium meliloti*. *Appl. Environ. Microbiol.*
1841 **79**(21): 6737–6746. doi:10.1128/AEM.01791-13.
- 1842 Todd, J.D., Sawers, G., and Johnston, A.W.B. 2005. Proteomic analysis reveals the wide-ranging
1843 effects of the novel, iron-responsive regulator RirA in *Rhizobium leguminosarum* bv. *viciae*.
1844 *Mol. Genet. Genomics* **273**(2): 197–206. doi:10.1007/s00438-005-1127-8.
- 1845 Tolin, S., Arrigoni, G., Moscatiello, R., Masi, A., Navazio, L., Sablok, G., and Squartini, A. 2013.
1846 Quantitative analysis of the naringenin-inducible proteome in *Rhizobium leguminosarum* by
1847 isobaric tagging and mass spectrometry. *Proteomics* **13**(12-13): 1961–1972.
1848 doi:10.1002/pmic.201200472.
- 1849 Toro, N., Martínez-Abarca, F., Molina-Sánchez, M.D., García-Rodríguez, F.M., and Nisa-
1850 Martínez, R. 2018. Contribution of mobile group II introns to *Sinorhizobium meliloti* genome
1851 evolution. *Front. Microbiol.* **9**: 627. doi:10.3389/fmicb.2018.00627.
- 1852 Torres-Quesada, O., Millán, V., Nisa-Martínez, R., Bardou, F., Crespi, M., Toro, N., and Jiménez-
1853 Zurdo, J.I. 2013. Independent activity of the homologous small regulatory RNAs AbcR1 and

- 1854 AbcR2 in the legume symbiont *Sinorhizobium meliloti*. PLOS One **8**(7): e68147.
 1855 doi:10.1371/journal.pone.0068147.
- 1856 Torres-Quesada, O., Oruezabal, R.I., Peregrina, A., Jofré, E., Lloret, J., Rivilla, R., Toro, N., and
 1857 Jiménez-Zurdo, J.I. 2010. The *Sinorhizobium meliloti* RNA chaperone Hfq influences central
 1858 carbon metabolism and the symbiotic interaction with alfalfa. BMC Microbiol. **10**(1): 71.
 1859 doi:10.1186/1471-2180-10-71.
- 1860 Torres-Quesada, O., Reinkensmeier, J., Schlüter, J.-P., Robledo, M., Peregrina, A., Giegerich, R.,
 1861 Toro, N., Becker, A., and Jiménez-Zurdo, J.I. 2014. Genome-wide profiling of Hfq-binding
 1862 RNAs uncovers extensive post-transcriptional rewiring of major stress response and symbiotic
 1863 regulons in *Sinorhizobium meliloti*. RNA Biol. **11**(5): 563–579. doi:10.4161/rna.28239.
- 1864 Triplett, E.W., and Barta, T.M. 1987. Trifolitoxin production and nodulation are necessary for the
 1865 expression of superior nodulation competitiveness by *Rhizobium leguminosarum* bv. *trifolii*
 1866 strain T24 on clover. Plant Physiol. **85**(2): 335-342. doi:10.1104/pp.85.2.335.
- 1867 Triplett, E.W., and Sadowsky, M.J. 1992. Genetics of competition for nodulation of legumes.
 1868 Annu. Rev. Microbiol. **46**(1): 399–422. doi:10.1146/annurev.mi.46.100192.002151.
- 1869 Truchet, G., Rosenberg, C., Vasse, J., Julliot, J.S., Camut, S., and Dénarié, J. 1984. Transfer of
 1870 *Rhizobium meliloti* pSym genes into *Agrobacterium tumefaciens*: host-specific nodulation by
 1871 atypical infection. J. Bacteriol. **157**(1): 134–142.
- 1872 Udvardi, M., and Poole, P.S. 2013. Transport and metabolism in legume-rhizobia symbioses.
 1873 Annu. Rev. Plant Biol. **64**(1): 781–805. doi:10.1146/annurev-arplant-050312-120235.
- 1874 Ulvé, V.M., Sevin, E.W., Cheron, A., and Barloy-Hubler, F. 2007. Identification of chromosomal
 1875 alpha-proteobacterial small RNAs by comparative genome analysis and detection in
 1876 *Sinorhizobium meliloti* strain 1021. BMC Genomics **8**(1): 467. doi:10.1186/1471-2164-8-467.

- 1877 Van de Velde, W., Zehirov, G., Szatmari, A., Debreczeny, M., Ishihara, H., Kevei, Z., Farkas, A.,
 1878 Mikulass, K., Nagy, A., Tiricz, H., Satiat-Jeunemaître, B., Alunni, B., Bourge, M., Kucho, K.-
 1879 I., Abe, M., Kereszt, A., Maróti, G., Uchiumi, T., Kondorosi, E., and Mergaert, P. 2010. Plant
 1880 peptides govern terminal differentiation of bacteria in symbiosis. *Science* **327**(5969): 1122–
 1881 1126. doi:10.1126/science.1184057.
- 1882 van Dillewijn, P., Soto, M.J., Villadas, P.J., and Toro, N. 2001. Construction and environmental
 1883 release of a *Sinorhizobium meliloti* strain genetically modified to be more competitive for
 1884 alfalfa nodulation. *Appl. Environ. Microbiol.* **67**(9): 3860–3865.
 1885 doi:10.1128/AEM.67.9.3860-3865.2001.
- 1886 VanYperen, R.D., Orton, T.S., and Griffiths, J.S. 2015. Genetic analysis of signal integration by
 1887 the *Sinorhizobium meliloti* sensor kinase FeuQ. *Microbiology* **161**(Pt 2): 244–253.
 1888 doi:10.1099/mic.0.000002.
- 1889 Vauclare, P., Bligny, R., Gout, E., and Widmer, F. 2013. An overview of the metabolic differences
 1890 between *Bradyrhizobium japonicum* 110 bacteria and differentiated bacteroids from soybean
 1891 (*Glycine max*) root nodules: an *in vitro* ¹³C- and ³¹P-nuclear magnetic resonance
 1892 spectroscopy study. *FEMS Microbio. Lett.* **343**(1): 49–56. doi:10.1111/1574-6968.12124.
- 1893 Vences-Guzmán, M.A., Geiger, O., and Sohlenkamp, C. 2008. *Sinorhizobium meliloti* mutants
 1894 deficient in phosphatidylserine decarboxylase accumulate phosphatidylserine and are strongly
 1895 affected during symbiosis with alfalfa. *J. Bacteriol.* **190**(20): 6846–6856.
 1896 doi:10.1128/JB.00610-08.
- 1897 Venter, A.P., Twelker, S., Oresnik, I.J., and Hynes, M.F. 2001. Analysis of the genetic region
 1898 encoding a novel rhizobiocin from *Rhizobium leguminosarum* bv. *viciae* strain 306. *Can. J.*
 1899 *Microbiol.* **47**(6): 495-502. doi:10.1139/w01-043.

- 1900 Vercruysse, M., Fauvart, M., Jans, A., Beullens, S., Braeken, K., Cloots, L., Engelen, K., Marchal,
1901 K., and Michiels, J. 2011. Stress response regulators identified through genome-wide
1902 transcriptome analysis of the (p)ppGpp-dependent response in *Rhizobium etli*. *Genome Biol.*
1903 **12**(2): R17. doi:10.1186/gb-2011-12-2-r17.
- 1904 Wang, C., Sheng, X., Equi, R.C., Trainer, M.A., Charles, T.C., and Sobral, B.W.S. 2007. Influence
1905 of the poly-3-hydroxybutyrate (PHB) granule-associated proteins (PhaP1 and PhaP2) on PHB
1906 accumulation and symbiotic nitrogen fixation in *Sinorhizobium meliloti* Rm1021. *J. Bacteriol.*
1907 **189**(24): 9050–9056. doi:10.1128/JB.01190-07.
- 1908 Wang, D., Wang, Y.C., Wu, L.J., Liu, J.X., Zhang, P., Jiao, J., Yan, H., Liu, T., Tian, C.F., and
1909 Chen, W.X. 2016. Construction and pilot screening of a signature-tagged mutant library of
1910 *Sinorhizobium fredii*. *Arch. Microbiol.* **198**(2): 91–99. doi:10.1007/s00203-015-1161-9.
- 1911 Wang, Q., Yang, S., Liu, J., Terecskei, K., Ábrahám, E., Gombár, A., Domonkos, Á., Szűcs, A.,
1912 Körmöczi, P., Wang, T., Fodor, L., Mao, L., Fei, Z., Kondorosi, E., Kaló, P., Kereszt, A., and
1913 Zhu, H. 2017. Host-secreted antimicrobial peptide enforces symbiotic selectivity in *Medicago*
1914 *truncatula*. *Proc. Natl. Acad. Sci. U.S.A.* **114**(26): 6854–6859. doi:10.1073/pnas.1700715114.
- 1915 Wang, X., Yang, J.-G., Chen, L., Wang, J.-L., Cheng, Q., Dixon, R., and Wang, Y.-P. 2013. Using
1916 synthetic biology to distinguish and overcome regulatory and functional barriers related to
1917 nitrogen fixation. *PLOS One* **8**(7): e68677. doi:10.1371/journal.pone.0068677.
- 1918 Weissenmayer, B., Gao, J.-L., López-Lara, I.M., and Geiger, O. 2002. Identification of a gene
1919 required for the biosynthesis of ornithine-derived lipids. *Mol. Microbiol.* **45**(3): 721–733.
1920 doi:10.1046/j.1365-2958.2002.03043.x.
- 1921 Weissenmayer, B., Geiger, O., and Benning, C. 2000. Disruption of a gene essential for
1922 sulfoquinovosyldiacylglycerol biosynthesis in *Sinorhizobium meliloti* has no detectable effect

- 1923 on root nodule symbiosis. *Mol. Plant Microbe Interact.* **13**(6): 666–672.
- 1924 doi:10.1094/MPMI.2000.13.6.666.
- 1925 Westhoek, A., Field, E., Rehling, F., Mulley, G., Webb, I., Poole, P.S., and Turnbull, L.A. 2017.
- 1926 Policing the legume-Rhizobium symbiosis: a critical test of partner choice. *Sci. Rep.* **7**(1):
- 1927 1419. doi:10.1038/s41598-017-01634-2.
- 1928 Wheatley, R.M., Ramachandran, V.K., Geddes, B.A., Perry, B.J., Yost, C.K., and Poole, P.S. 2017.
- 1929 The role of O₂ in the growth of *Rhizobium leguminosarum* bv. *viciae* 3841 on glucose and
- 1930 succinate. *J. Bacteriol.* **199**(1): e00572–16–16. doi:10.1128/JB.00572-16.
- 1931 Wilson, R.A., Handley, B.A., and Beringer, J.E. 1998. Bacteriocin production and resistance in a
- 1932 field population of *Rhizobium leguminosarum* biovar *viciae*. *Soil Biol. Biochem.* **30**(3): 413–
- 1933 417. doi:10.1016/S0038-0717(97)00123-5.
- 1934 Wong, C.H., Pankhurst, C.E., Kondorosi, A., and Broughton, W.J. 1983. Morphology of root
- 1935 nodules and nodule-like structures formed by *Rhizobium* and *Agrobacterium* strains
- 1936 containing a *Rhizobium meliloti* megaplasmid. *J. Cell Biol.* **97**(3): 787–794.
- 1937 doi:10.1083/jcb.97.3.787.
- 1938 Wongdee, J., Boonkerd, N., Teaumroong, N., Tittabutr, P., and Giraud, E. 2018. Regulation of
- 1939 nitrogen fixation in *Bradyrhizobium* sp. strain DOA9 involves two distinct NifA regulatory
- 1940 proteins that are functionally redundant during symbiosis but not during free-living growth.
- 1941 *Front. Microbiol.* **9**: 1644. doi:10.3389/fmicb.2018.01644.
- 1942 Yamaya-Ito, H., Shimoda, Y., Hakoyama, T., Sato, S., Kaneko, T., Hossain, M.S., Shibata, S.,
- 1943 Kawaguchi, M., Hayashi, M., KOUCHI, H., and Umehara, Y. 2018. Loss-of-function of
- 1944 ASPARTIC PEPTIDASE NODULE-INDUCED 1 (APN1) in *Lotus japonicus* restricts

- 1945 efficient nitrogen-fixing symbiosis with specific *Mesorhizobium loti* strains. *Plant J.* **93**(1): 5–
 1946 16. doi:10.1111/tpj.13759.
- 1947 Yang, J., Xie, X., Wang, X., Dixon, R., and Wang, Y.-P. 2014. Reconstruction and minimal gene
 1948 requirements for the alternative iron-only nitrogenase in *Escherichia coli*. *Proc. Natl. Acad.*
 1949 *Sci. U.S.A.* **111**(35): E3718–25. doi:10.1073/pnas.1411185111.
- 1950 Yang, J., Xie, X., Yang, M., Dixon, R., and Wang, Y.-P. 2017a. Modular electron-transport chains
 1951 from eukaryotic organelles function to support nitrogenase activity. *Proc. Natl. Acad. Sci.*
 1952 *U.S.A.* **114**(12): E2460–E2465. doi:10.1073/pnas.1620058114.
- 1953 Yang, S., Wang, Q., Fedorova, E., Liu, J., Qin, Q., Zheng, Q., Price, P.A., Pan, H., Wang, D.,
 1954 Griffiths, J.S., Bisseling, T., and Zhu, H. 2017b. Microsymbiont discrimination mediated by a
 1955 host-secreted peptide in *Medicago truncatula*. *Proc. Natl. Acad. Sci. U.S.A.* **114**(26): 6848–
 1956 6853. doi:10.1073/pnas.1700460114.
- 1957 Yang, Y., Hu, X.-P., and Ma, B.-G. 2017c. Construction and simulation of the *Bradyrhizobium*
 1958 *diazoefficiens* USDA110 metabolic network: a comparison between free-living and symbiotic
 1959 states. *Mol. Biosyst.* **13**(3): 607–620. doi:10.1039/c6mb00553e.
- 1960 Ye, H., Gemperline, E., Venkateshwaran, M., Chen, R., Delaux, P.-M., Howes-Podoll, M., Ané,
 1961 J.-M., and Li, L. 2013. MALDI mass spectrometry-assisted molecular imaging of metabolites
 1962 during nitrogen fixation in the *Medicago truncatula*-*Sinorhizobium meliloti* symbiosis. *Plant*
 1963 *J.* **75**(1): 130–145. doi:10.1111/tpj.12191.
- 1964 Young, J.P.W., Crossman, L.C., Johnston, A.W., Thomson, N.R., Ghazoui, Z.F., Hull, K.H.,
 1965 Wexler, M., Curson, A.R., Todd, J.D., Poole, P.S., Mauchline, T.H., East, A.K., Quail, M.A.,
 1966 Churcher, C., Arrowsmith, C., Cherevach, I., Chillingworth, T., Clarke, K., Cronin, A., Davis,
 1967 P., Fraser, A., Hance, Z., Hauser, H., Jagels, K., Moule, S., Mungall, K., Norbertczak, H.,

- 1968 Rabbinoiwitsch, E., Sanders, M., Simmonds, M., Whitehead, S., and Parkhill, J. 2006. The
 1969 genome of *Rhizobium leguminosarum* has recognizable core and accessory components.
 1970 Genome Biol. **7**(4): R34. doi:10.1186/gb-2006-7-4-r34.
- 1971 Yuan, Z.-C., Zaheer, R., Morton, R., and Finan, T.M. 2006. Genome prediction of PhoB regulated
 1972 promoters in *Sinorhizobium meliloti* and twelve proteobacteria. Nucleic Acids Res. **34**(9):
 1973 2686–2697. doi:10.1093/nar/gkl365.
- 1974 Yurgel, S.N., Mortimer, M.W., Rice, J.T., Humann, J.L., and Kahn, M.L. 2013a. Directed
 1975 construction and analysis of a *Sinorhizobium meliloti* pSymA deletion mutant library. Appl.
 1976 Environ. Microbiol. **79**(6): 2081–2087. doi:10.1128/AEM.02974-12.
- 1977 Yurgel, S.N., Rice, J., and Kahn, M.L. 2013b. Transcriptome analysis of the role of GlnD/GlnBK
 1978 in nitrogen stress adaptation by *Sinorhizobium meliloti* Rm1021. PLOS One **8**(3): e58028.
 1979 doi:10.1371/journal.pone.0058028.
- 1980 Zamani, M., diCenzo, G.C., Milunovic, B., and Finan, T.M. 2017. A putative 3-hydroxyisobutyryl-
 1981 CoA hydrolase is required for efficient symbiotic nitrogen fixation in *Sinorhizobium meliloti*
 1982 and *Sinorhizobium fredii* NGR234. Environ. Microbiol. **19**(1): 218–236. doi:10.1111/1462-
 1983 2920.13570.
- 1984 Zavaleta-Pastor, M., Sohlenkamp, C., Gao, J.L., Guan, Z., Zaheer, R., Finan, T.M., Raetz, C.R.H.,
 1985 López-Lara, I.M., and Geiger, O. 2010. *Sinorhizobium meliloti* phospholipase C required for
 1986 lipid remodeling during phosphorus limitation. Proc. Natl. Acad. Sci. U.S.A. **107**(1): 302–307.
 1987 doi:10.1073/pnas.0912930107.
- 1988 Zhang, X.-S., and Cheng, H.-P. 2006. Identification of *Sinorhizobium meliloti* early symbiotic
 1989 genes by use of a positive functional screen. Appl. Environ. Microbiol. **72**(4): 2738–2748.
 1990 doi:10.1128/AEM.72.4.2738-2748.2006.

- 1991 Zhang, Y., Smallbone, L.A., diCenzo, G.C., Morton, R., and Finan, T.M. 2016. Loss of malic
 1992 enzymes leads to metabolic imbalance and altered levels of trehalose and putrescine in the
 1993 bacterium *Sinorhizobium meliloti*. **16**(1): 491. doi:10.1186/s12866-016-0780-x.
- 1994 Zhao, H., Li, M., Fang, K., Chen, W., and Wang, J. 2012. *In silico* insights into the symbiotic
 1995 nitrogen fixation in *Sinorhizobium meliloti* via metabolic reconstruction. PLOS One **7**(2):
 1996 e31287. doi:10.1371/journal.pone.0031287.
- 1997 Zou, L., Gastebois, A., Mathieu-Demazière, C., Sorroche, F., Masson-Boivin, C., Batut, J., and
 1998 Garnerone, A.-M. 2017. Transcriptomic insight in the control of legume root secondary
 1999 infection by the *Sinorhizobium meliloti* transcriptional regulator Clr. Front. Microbiol. **8**:
 2000 1236. doi:10.3389/fmicb.2017.01236.

FIGURE LEGENDS

Figure 1. Phylogenetic distribution of biological nitrogen fixation. A maximum likelihood phylogeny of the α - and β -proteobacteria based on 23 highly conserved proteins (Frr, NusA, RplB, RplC, RplD, RplK, RplL, RplM, RplN, RplP, RplS, RplT, RpmA, RpoB, RpsB, RpsC, RpsE, RpsI, RpsJ, RpsK, RpsM, RpsS, Tsf). Species with at least one strain carrying the *nifHDK* nitrogen fixation genes are coloured red (sample genera are indicated), while species with at least one strain carrying the *nifHDK* nitrogen fixation genes and the *nodABC* nodulation genes are coloured blue (the genera and families are indicated). No strains have the nodulation genes but lack the nitrogen fixation genes. Branch lengths for the taxa that include the genera *Tremblaya*, *Hodgkinia*, and *Liberibacter* (light grey) were shortened for presentation. The enlarged phylogeny, including taxa names and bootstrap values, is provided as File S2. See the Methods in File S1 for details on phylogeny construction and gene identification (diCenzo et al. 2017c, diCenzo et al. 2018b).

Figure 2. Visualizing the rhizobium – legume symbiosis. (A) A picture of *Medicago alba* (white sweet clover) plants grown in nitrogen free conditions. Plants in columns 1-3 and 5 are inoculated with nitrogen-fixing *S. meliloti*; plants in columns 4 and 6 are inoculated with a *S. meliloti* mutant unable to fix nitrogen; and plants in column 7 are not inoculated with *S. meliloti*. (B) *Medicago sativa* (alfalfa) root nodules, induced by *S. meliloti*. On the left are ineffective, white nodules containing a *S. meliloti* mutant unable to fix nitrogen; on the right are effective, pink nodules containing *S. meliloti* cells actively fixing nitrogen. (C) Effective, pink nodules of *Vigna unguiculata* (cowpea) containing *S. fredii* NGR234 cells actively fixing nitrogen. (D, E) Confocal micrographs of alfalfa root nodules filled with *S. meliloti* cells. The nodule sections were stained with a fluorescent nucleic acid binding dye (Syto9); the green colour is indicative of the presence of *S. meliloti*. Scale bar represents 250 μ m. (F, G) Transmission electron micrographs of alfalfa

root nodules, showing plant cells filled with nitrogen-fixing *S. meliloti* bacteroids. Scale bar represents 25 μm .

Figure 3. Schematic representation of the life-cycle of rhizobia. Inoculant strains are typically grown at fermenter scale and used to coat seeds of compatible legumes. In the soil, the rhizobia must compete with the indigenous microbes and initiate symbiosis with an exchange of specific nodulation signals. Curling of a host root hair traps a rhizobium cell, allowing colonization to proceed in a growing infection thread that progresses inward to the cortical cells. Here, the rhizobia are released into the cytoplasm of specialized nodule cells, where they are enclosed in a plant-derived membrane (peribacteroid membrane). The rhizobia and plant undergo a differentiation process that involves massive transcriptional and metabolic shifts, resulting in the formation of a nitrogen-fixing nodule. Five distinct developmental zones of an indeterminate-type nodule are shown (not drawn to scale): zone I – apical meristem; zone II – infection and differentiation zone; interzone II-III – a small region between zone II and zone III; zone III – the nitrogen-fixing zone; zone IV – the senescence zone found in mature nodules. We note that this figure shows an indeterminate nodule, that many of the features shown in this image are not universal, and that many types of nodulation exist; see Masson-Boivin et al. (2009) for a description of the differences.

Figure 4. Example output of a Tn-seq study. A *S. meliloti* gene region illustrating the ability of Tn-seq to identify essential genes and environment-specific essential genes is shown. Published Tn-seq data (diCenzo et al. 2018a) for *S. meliloti* grown in a nutritionally complex medium (rich medium) and a nutritionally defined medium (defined medium) were mapped to the *S. meliloti* chromosome. A five-gene region is shown. The location of the transposon insertions in the cell populations recovered from both conditions are indicated by the red bars, with the height of the

2047 bar representing the number of reads (log scale). Genes are colour coded based on their fitness
 2048 classifications.

2049 **Figure 5. The *S. meliloti* pangenome.** The pangenome of 20 *S. meliloti* strains was determined
 2050 using Roary (Page et al. 2015). Strains possess a median of 6,556 genes per genome, with a total
 2051 of 17,494 sets of orthologous proteins, of which 4,000 were present in all strains. **(A)** A heatmap
 2052 showing gene presence (dark blue) or absence (light blue) in each of the 20 strains. A phylogeny
 2053 built based on the 4,000 core genes is shown on the left, and the strain names are indicated on the
 2054 right. **(B)** A histogram displaying the distribution of how many genomes each gene is found within.
 2055 See the Methods in File S1 for details on pangenome construction.

2056 **Figure 6. Expression patterns of select *S. meliloti* genes during symbiosis with *M. truncatula*.**
 2057 The spatial expression patterns across a *M. truncatula* nodule of 10 *S. meliloti* genes, each
 2058 displaying a unique pattern, is shown. Data were taken from the study of Roux *et al.* (Roux et al.
 2059 2014). The nodule zone is indicated along the bottom, and the gene names are given on the right.
 2060 The plotted data is normalized across each zone; the darker the blue, the higher the expression of
 2061 the gene in the indicated zone relative to other zones. For example, *aglE* is expressed strongly in
 2062 the distal zone II but lowly in zone III, whereas *fixU* is primarily expressed in zone III. Colours
 2063 provide no information on the expression of a gene compared to the other genes shown in the
 2064 figure. Locations of each nodule zone are shown in Figure 3, with zone II distal and zone II
 2065 proximal referring to the sections of zone II that are distal and proximal to the root, respectively.

2066 **Figure 7. *In silico* predicted flux distributions and reaction essentialities.** Example data
 2067 obtained through metabolic modelling. Results from *in silico* metabolic modelling of *S. meliloti*
 2068 metabolism using the iGD726 metabolic reconstruction are shown (diCenzo et al. 2018a). Growth
 2069 was simulated using **(A)** glucose or **(B)** succinate as the sole carbon source. The pathways represent

2070 central carbon metabolism, with the exception of the pentose phosphate pathway (for simplicity).
2071 The width of the line represents the amount of flux through the reaction; a doubling in the width
2072 corresponds to a ten-fold flux increase. Arrows indicate the direction of flux. Colours indicate if
2073 the reaction is essential (red) or non-essential (blue). See the Methods in File S1 for details on the
2074 modelling procedures used. A version with gene names is provided as Figure S1.
2075

Table 1. Tools for functional genomics and genome manipulation of rhizobia, and rhizobium biosensor and reporter strains.

Functional genomic tools and resources		
Tool	Description	References
Marker-less gene deletion	Sample systems for the construction of marker-less gene deletions in diverse rhizobia	(Quandt and Hynes 1993, Ledermann et al. 2016)
Site-specific DNA integration	Site-specific gene integration in a neutral location	(Arango Pinedo and Gage 2009)
Cloning-free genome engineering	Cloning-free method for DNA insertion and deletion	(Döhlemann et al. 2016)
<i>S. meliloti</i> ORFeome	Plasmid-based ORF library representing all <i>S. meliloti</i> Rm1021 open reading frames	(Schroeder et al. 2005)
<i>S. meliloti</i> fusion libraries	Libraries of <i>S. meliloti</i> strains each with an integrated or replicative transcriptional reporter plasmid	(Cowie et al. 2006, Mauchline et al. 2006)
IVET* libraries	Libraries to identify expressed promoters in specific conditions (nodule, rhizosphere), in <i>S. meliloti</i> , and <i>R. leguminosarum</i>	(Oke and Long 1999, Zhang and Cheng 2006)
STM† libraries	Signature-tagged mutageneses libraries of <i>S. meliloti</i> , <i>S. fredii</i> , <i>R. leguminosarum</i> , and <i>M. loti</i>	(Pobigaylo et al. 2006, Borjigin et al. 2011, Garcia-Fraile et al. 2015, Wang et al. 2016)
Tn-seq / INseq transposons	Mariner and Tn5 transposons for Tn-seq / INseq analyses	(Perry and Yost 2014, Arnold et al. 2017, diCenzo et al. 2018a)
Site-specific recombinases	Systems for using site specific recombinases (Flp/ <i>FRT</i> , Cre/ <i>lox</i> , ΦC31 integrase/ <i>attB/attP</i> , lambda/ <i>attB/attP</i>) for large-scale genome deletions, <i>in vivo</i> cloning, and gene insertion	(Chain et al. 2000, House et al. 2004, Landeta et al. 2011, Harrison et al. 2011, Heil et al. 2012, diCenzo et al. 2013, Yurgel et al. 2013a, Milunovic et al. 2014, diCenzo et al. 2016b, Döhlemann et al. 2016, diCenzo and Finan 2018)
Biosensors and reporter strains		
Description		Reference
Biosensors for measuring ecotoxicity and heavy metal presence based on constitutive <i>lux</i> expression and cell viability		(Paton et al. 1997, Chaudri 2000)
Biosensor for galactosides, based on promoter fusion to <i>gfp</i>		(Bringinghurst et al. 2001)
Biosensors for sugars, dicarboxylates, and polyols based on the use of solute binding proteins of transporters and FRET‡		(Bourdès et al. 2012)
Biosensors for the <i>in vivo</i> mapping of the distribution of root exudate metabolites based on promoter fusions to <i>lux</i>		(Pini et al. 2017)
<i>S. meliloti</i> reporter strain to identify the stage of symbiosis in which plant mutants are blocked		(Lang et al. 2018)

* *In vivo* expression technology
† Signature-tagged mutagenesis
‡ Förster resonance energy transfer

Table 2. Studies reporting synthetic, large-scale reductions of the genomes of rhizobia

Organism	Description	References
<i>S. meliloti</i> Rm1021 and RmP110	Libraries of large-scale deletion mutants of the pSymA megaplasmid and pSymB chromid	(Charles and Finan 1991, Yurgel et al. 2013a, Milunovic et al. 2014)
<i>S. meliloti</i> Rm2011	Complete removal of one or both of the pSymA megaplasmid and pSymB chromid	(Oresnik et al. 2000, diCenzo et al. 2014)
<i>S. meliloti</i> Rm2011	Deletions of the pSymA megaplasmid	(Hynes et al. 1989)
<i>S. meliloti</i> MV11	Complete removal of four cryptic plasmids	(Hynes et al. 1989)
<i>S. meliloti</i> various	Complete removal large host-range determining plasmids	(Crook et al. 2012)
<i>S. fredii</i> NGR234	Complete removal of the symbiotic megaplasmid (pNGR234a)	(Morrison et al. 1983)
<i>S. fredii</i> USDA 206	Complete removal of one to three non-symbiotic plasmids	(Mathis et al. 1985, Barbour and Elkan 1989)
<i>S. fredii</i> GR64	Complete removal of one or both of the (mega)-plasmids, including the symbiotic plasmid	(Cervantes et al. 2011)
<i>R. leguminosarum</i> various	Complete removal or large deletions of many (mega)-plasmids, including symbiotic plasmids	(Hynes et al. 1989)
<i>R. leguminosarum</i> VF39	Complete removal of each of the six (mega)-plasmids	(Hynes and McGregor 1990)
<i>R. leguminosarum</i> bv. <i>trifolii</i> various	Complete removal of two to four of the (mega)-plasmids	(Baldani et al. 1992)
<i>R. leguminosarum</i> bv. <i>trifolii</i> W14-2	Complete removal of each of the four (mega)-plasmids and (mega)-plasmid combinations, including preparing a strain lacking plasmids	(Moënne-Loccoz et al. 1995)
<i>R. leguminosarum</i> bv. <i>trifolii</i> TA1	Complete removal of two megaplasmids, and deletions of the symbiotic megaplasmid	(Stasiak et al. 2014)
<i>R. etli</i> CFN42	Complete removal of five of the six (mega)-plasmids, deletions in the sixth, and combinations with up to five (mega)-plasmids removed in one strain	(Brom et al. 1992, 2000)
<i>R. etli</i> CFN42	Library of large-scale deletion mutants of the p42e chromid	(Landeta et al. 2011)
<i>R. etli</i> CFN42	Deletion of the symbiotic region of the symbiotic megaplasmid	(Romero et al. 1991)
<i>R. tropici</i> CIAT 899	Complete removal of three (mega)-plasmids including the symbiotic megaplasmid, with one to three plasmids removed per strain	(Barreto et al. 2012)
<i>R. sp.</i> P1	Complete removal of the symbiosis megaplasmid	(Sharma and Laxminarayana 1989)
<i>M. loti</i> NZP2213, NZP2037	Complete removal of the large non-symbiotic plasmid	(Chua et al. 1985)
<i>M. loti</i> R7A	Deletion of the large chromosomal symbiotic island	(Ramsay et al. 2006)
<i>M. hawaii</i> HN308SR, 7653R	Complete removal of plasmids	(Hu et al. 2008)

Table 3. Studies of rhizobial species employing -omics technologies.

Topic of the study	Organisms	Approaches	References
Characterization of proteins			
SyrM (Nod factor synthesis)	<i>S. meliloti</i>	Transcriptomics	(Barnett and Long 2015)
NodD (Nod factor synthesis)	<i>S. meliloti</i>	Transcriptomics	(Capela et al. 2005)
NolR (Nod factor synthesis)	<i>S. meliloti</i>	Proteomics	(Chen et al. 2000a, 2005)
FixL/FixJ/FixK (nitrogen fixation)	<i>S. meliloti</i> ; <i>B. diazoefficiens</i>	Transcriptomics	(Bobik et al. 2006, Mesa et al. 2008)
CbrA (cell cycle and symbiosis)	<i>S. meliloti</i>	Transcriptomics	(Gibson et al. 2007)
CtrA (cell cycle)	<i>S. meliloti</i>	Transcriptomics	(Pini et al. 2015)
RosR (exopolysaccharide synthesis)	<i>R. leguminosarum</i> bv. <i>trifolii</i>	Transcriptomics	(Rachwał et al. 2015)
FeuP/FeuQ (osmoadaptation and symbiosis)	<i>S. meliloti</i>	Transcriptomics	(Griffitts et al. 2008)
ECF sigma factors (stress response and symbiosis)	<i>R. etli</i> ; <i>B. diazoefficiens</i>	Transcriptomics	(Gourion et al. 2009, Martínez-Salazar et al. 2009, Stockwell et al. 2012)
Clr (cAMP dependent regulator, nodulation)	<i>S. meliloti</i>	Transcriptomics	(Krol et al. 2016, Zou et al. 2017)
PraR (quorum sensing and biofilms)	<i>R. leguminosarum</i> bv. <i>viciae</i>	Transcriptomics	(Frederix et al. 2014)
PckR (central carbon metabolism)	<i>S. meliloti</i>	Transcriptomics	(diCenzo et al. 2017b)
PhoB (phosphate limitation response)	<i>S. meliloti</i>	Transcriptomics	(Krol and Becker 2004, Yuan et al. 2006)
GlnD/GlnBK (nitrogen limitation response)	<i>S. meliloti</i>	Transcriptomics	(Yurgel et al. 2013b)
NtrY/NtrX (nitrogen metabolism)	<i>S. meliloti</i>	Transcriptomics	(Calatrava-Morales et al. 2017)
Tkt2/Tal (central carbon metabolism)	<i>S. meliloti</i>	Proteomics; metabolomics	(Hawkins et al. 2018)
OxyR (oxidative stress)	<i>S. meliloti</i>	Transcriptomics	(Lehman and Long 2018)
RelA/Rsh (stringent response)	<i>S. meliloti</i> ; <i>R. etli</i>	Transcriptomics	(Vercruysse et al. 2011, Krol and Becker 2011)
RirA (iron limitation)	<i>S. meliloti</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i>	Transcriptomics; proteomics	(Todd et al. 2005, Chao et al. 2005)
LeuB (leucine biosynthesis)	<i>S. meliloti</i>	Metabolomics	(Barsch et al. 2004)
AniA (PHB accumulation)	<i>R. etli</i>	Proteomics	(Encarnación et al. 2002)
FadD (Fatty acid metabolism)	<i>S. meliloti</i>	Lipid analysis	(Nogales et al. 2010)
PHB cycle mutations	<i>S. meliloti</i>	Transcriptomics; proteomics	(Wang et al. 2007, D'Alessio et al. 2017)
Dme/Tme (Malic enzymes)	<i>S. meliloti</i>	Transcriptomics; metabolomics	(Zhang et al. 2016)
Membrane lipid biosynthetic proteins (Cfa, PlsC, PlcP, Pcs, Pmt, OlsA, OlsB, PssA, Psd, SqdB, BtaA)	<i>S. meliloti</i>	Lipidomics and lipid analysis	(de Rudder et al. 1999, Weissenmayer et al. 2000, Sohlenkamp et al. 2000, de Rudder et al. 2000, Weissenmayer et al. 2002, Sohlenkamp et al. 2004, Gao et al. 2004, López-Lara et al. 2005, Vences-Guzmán et al. 2008, Basconcillo et al. 2009a, 2009c, Zavaleta-Pastor et al. 2010, Pech-Canul et al. 2011)
Riboregulation			
Small RNAs (sRNA)	<i>S. meliloti</i>	Transcriptomics	(Ulvé et al. 2007, Schlüter et al. 2010)
Hfq (RNA binding protein)	<i>S. meliloti</i>	Transcriptomics; proteomics	(Barra-Bily et al. 2010, Torres-Quesada et al. 2010, Gao et al. 2010, Sobrero et al. 2012)
Hfq binding RNAs	<i>S. meliloti</i>	Transcriptomics	(Torres-Quesada et al. 2014)
YbeY (endoribonuclease)	<i>S. meliloti</i>	Transcriptomics	(Saramago et al. 2017)
AbrC1/AbrC2 (sRNAs, nutrient uptake)	<i>S. meliloti</i>	Proteomics	(Torres-Quesada et al. 2013)
MmgR (sRNA, PHB accumulation)	<i>S. meliloti</i>	Proteomics	(Lagares et al. 2017)
Environmental adaptation and stresses			
Growth in root hairs	<i>B. diazoefficiens</i>	Metabolomics	(Brechenmacher et al. 2010)
Oxic versus microoxic growth	<i>S. meliloti</i> ; <i>B. diazoefficiens</i>	Transcriptomics	(Ampe et al. 2003, Becker et al. 2004, Bobik et al. 2006, Pessi et al. 2007)

Aerobic, micro-aerobic, anaerobic growth	<i>B. diazoefficiens</i>	Proteomics	(Dainese-Hatt et al. 1999)
Rhizosphere adaptation	<i>R. leguminosarum</i> bv. <i>viciae</i>	Transcriptomics	(Ramachandran et al. 2011)
Growth in presence of root exudates	<i>B. diazoefficiens</i> ; <i>B. phymatum</i> ; <i>C. taiwanensis</i> ; <i>R. mesoamericanum</i>	Transcriptomics	(Liu et al. 2017b, Klonowska et al. 2018)
Seed endophytic growth	<i>R. phaseoli</i> ; <i>S. americanum</i>	Proteomics	(Peralta et al. 2016)
Stationary phase adaptation	<i>S. meliloti</i> ; <i>R. etli</i>	Proteomics	(Guerreiro et al. 1999, Meneses et al. 2010)
Phosphate limitation	<i>S. meliloti</i>	Lipidomics and lipid analysis	(Basconcillo et al. 2009a, 2009b, 2009c, Zavaleta-Pastor et al. 2010)
Biotin limitation	<i>S. meliloti</i>	Proteomics	(Heinz and Streit 2003)
Carbon limitation	<i>S. meliloti</i>	Proteomics	(Djordjevic et al. 2007)
Nitrogen limitation	<i>S. meliloti</i>	Proteomics	(Djordjevic et al. 2007)
Osmotic/salt stress	<i>S. meliloti</i> ; <i>M. loti</i> ; <i>M. alhagi</i> ; <i>R. etli</i> ; <i>S. sp.</i> BL3; <i>B. spp.</i>	Transcriptomics; proteomics	(Jebbar et al. 2005, Shamseldin et al. 2006, Dominguez-Ferreras et al. 2006, Tanthanuch et al. 2010, Liu et al. 2014, Laranjo et al. 2017, Dong et al. 2017)
Oxidative stress	<i>B. diazoefficiens</i>	Transcriptomics	(Donati et al. 2011, Jeon et al. 2011)
Acid stress	<i>S. meliloti</i> ; <i>M. loti</i> ; <i>S. medicae</i>	Transcriptomics; proteomics; metabolomics; lipidomics	(Reeve et al. 2004, Basconcillo et al. 2009a, Hellweg et al. 2009, Laranjo et al. 2014, Draghi et al. 2016)
Heat stress	<i>M. loti</i> ; <i>R. tropici</i> ; <i>B. diazoefficiens</i>	Transcriptomics; proteomics	(Münchbach et al. 1999, Gomes et al. 2012a, Alexandre et al. 2014)
Cold stress	<i>M. sp.</i> N33	Transcriptomics; metabolomics	(Ghobakhlou et al. 2013, 2015)
Heavy metal stress	<i>S. meliloti</i> ; <i>M. metallidurans</i> ; <i>M. sp.</i> STM 4661; <i>R. sp.</i> VMA301; <i>R. sp.</i> E20-8	Transcriptomics; proteomics; metabolomics	(Mandal et al. 2009, Maynaud et al. 2013, Lu et al. 2017, Cardoso et al. 2017b)
Dessication stress	<i>B. diazoefficiens</i>	Transcriptomics	(Cytryn et al. 2007)
Herbicide exposure	<i>R. leguminosarum</i> bv. <i>viciae</i>	Metabolomics	(Bhat et al. 2014)
Cellular processes			
Flavonoid perception	<i>S. meliloti</i> ; <i>S. fredii</i> ; <i>R. tropici</i> ; <i>R. leguminosarum</i> bv. <i>trifolii</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i> ; <i>R. etli</i> ; <i>B. liaoningense</i> ; <i>B. japonicum</i> ; <i>B. diazoefficiens</i>	Transcriptomics; proteomics	(Guerreiro et al. 1997, Chen et al. 2000b, Ampe et al. 2003, Barnett et al. 2004, Capela et al. 2005, Hempel et al. 2009, da Silva Batista and Hungria 2012, Tolin et al. 2013, Arrigoni et al. 2013, Gao et al. 2015, Pérez-Montaño et al. 2016a, 2016b, Meneses et al. 2017)
Nod factor production	<i>S. arboris</i>	Metabolite analysis	(Penttinen et al. 2013)
Cell cycle	<i>S. meliloti</i>	Transcriptomics	(De Nisco et al. 2014, Pini et al. 2015)
Response to NCR peptides	<i>S. meliloti</i>	Transcriptomics	(Tiricz et al. 2013, Penterman et al. 2014)
Secretome / Type III secretion	<i>B. diazoefficiens</i> ; <i>S. fredii</i> ; <i>B. elkanii</i>	Proteomics	(Saad et al. 2005, Rodrigues et al. 2007, Hempel et al. 2009, Okazaki et al. 2009)
Biofilm formation	<i>R. etli</i>	Transcriptomics	(Reyes-Pérez et al. 2016)
Quorum sensing	<i>S. fredii</i> ; <i>S. meliloti</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i>	Transcriptomics; proteomics	(Chen et al. 2003, Gao et al. 2005, Cantero et al. 2006, Gao et al. 2007, Krysiak et al. 2014)
Motility	<i>S. meliloti</i>	Transcriptomics	(Nogales et al. 2010)
Aerobic vs fermentative growth	<i>R. etli</i>	Proteomics	(Encarnación et al. 2003)
Lectin perception	<i>B. diazoefficiens</i>	Proteomics	(Pérez-Giménez et al. 2012)
Catabolism of phenanthrene	<i>S. sp.</i> C4	Metabolomics	(Keum et al. 2008)
Symbiotic island transfer	<i>M. ciceri</i>	Transcriptomics	(Haskett et al. 2018)
Nodule analyses			
Integrated plant + bacterium analyses	<i>S. meliloti</i> - <i>M. truncatula</i> ; <i>S. meliloti</i> - <i>M. alba</i> ; <i>S. medicae</i> - <i>M. truncatula</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i> - <i>P. sativum</i>	Transcriptomics; proteomics	(Natera et al. 2000, Barnett et al. 2004, Heath et al. 2012, Roux et al. 2014, Marx et al. 2016, Ogden et al. 2017, Mitsch et al. 2018)
Bacteroids vs free-living	<i>S. meliloti</i> - <i>M. truncatula</i> ; <i>S. meliloti</i> - <i>M. sativa</i> ; <i>S. meliloti</i> - <i>M. alba</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i> - <i>P. sativum</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i> - <i>V. cracca</i> ;	Transcriptomics; proteomics; metabolomics	(Ampe et al. 2003, Becker et al. 2004, Djordjevic 2004, Barnett et al. 2004, Sarma and Emerich 2006, Djordjevic et al. 2007, Pessi et al. 2007, Karunakaran et al. 2009, Resendis-Antonio et al. 2012, Vauclare et al. 2013, Tatsukami et al. 2013)

Nodules of varying maturity	<i>B. diazoefficiens</i> - <i>G. max</i> ; <i>M. loti</i> - <i>L. japonicus</i> ; <i>R. etli</i> - <i>P. vulgaris</i> <i>S. meliloti</i> - <i>M. truncatula</i> ; <i>S. meliloti</i> - <i>M. sativa</i> ; <i>B. sp.</i> ORS 278 - <i>A. indica</i> ; <i>M. loti</i> - <i>L. japonicus</i>	Transcriptomics; proteomics	(Ampe et al. 2003, Capela et al. 2006, Delmotte et al. 2014, Nambu et al. 2015, Lardi et al. 2016, Marx et al. 2016)
Plant/bacterium mutants	<i>S. meliloti</i> - <i>M. truncatula</i> ; <i>S. meliloti</i> - <i>M. sativa</i> ; <i>B. diazoefficiens</i> - <i>G. max</i>	Transcriptomics; metabolomics	(Tian et al. 2006, Starker et al. 2006, Capela et al. 2006, Maunoury et al. 2010, Ye et al. 2013, Gemperline et al. 2015, Lardi et al. 2016)
Nodule zones	<i>S. meliloti</i> - <i>M. truncatula</i> ; <i>S. medicae</i> - <i>M. truncatula</i>	Transcriptomics; proteomics; metabolomics	(Roux et al. 2014, Ogden et al. 2017)
Reference nodule proteome/transcriptome	<i>B. diazoefficiens</i> - <i>G. max</i> ; <i>R. etli</i> - <i>P. vulgaris</i>	Transcriptomics; proteomics	(Sarma and Emerich 2005, Delmotte et al. 2010, Resendis-Antonio et al. 2011)
Sulfenylated nodule proteome	<i>S. meliloti</i> - <i>M. truncatula</i>	Proteomics	(Oger et al. 2012)
Peribacteroid space/membrane proteome	<i>R. leguminosarum</i> bv. <i>viciae</i> - <i>P. sativum</i>	Proteomics	(Saalbach et al. 2002)
Bacteroid periplasm	<i>B. diazoefficiens</i> - <i>G. max</i>	Proteomics	(Strodtman et al. 2017)
FixJ in nodules	<i>S. meliloti</i> - <i>M. truncatula</i>	Transcriptomics	(Barnett et al. 2004)
BacA in nodules	<i>R. leguminosarum</i> bv. <i>viciae</i>	Transcriptomics	(Karunakaran et al. 2010)
NifA in nodules	<i>R. etli</i> - <i>P. vulgaris</i>	Proteomics	(Salazar et al. 2010)
Control of plant immunity (plant <i>symCRK</i> mutant)	<i>M. truncatula</i> - <i>S. medicae</i>	Proteomics	(Berrabah et al. 2018)
C4-dicarboxylate transport	<i>R. leguminosarum</i> bv. <i>viciae</i> - <i>P. sativum</i>	Transcriptomics	(Mitsch et al. 2018)
Lipogenesis	<i>R. leguminosarum</i> bv. <i>viciae</i> - <i>P. sativum</i>	Metabolomics	(Terpolilli et al. 2016)
Effect of genetic variation	<i>S. meliloti</i> - <i>M. truncatula</i> ; <i>S. medicae</i> - <i>M. truncatula</i>	Transcriptomics	(Heath et al. 2012, Burghardt et al. 2017)
Effect of plant host	<i>S. meliloti</i> - <i>M. truncatula</i> ; <i>S. meliloti</i> - <i>M. sativa</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i> - <i>P. sativum</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i> - <i>V. cracca</i> ; <i>S. fredii</i> - <i>V. unguiculata</i> ; <i>S. fredii</i> - <i>L. leucocephala</i> ; <i>B. diazoefficiens</i> - <i>G. max</i> ; <i>B. diazoefficiens</i> - <i>V. radiata</i> ; <i>B. diazoefficiens</i> - <i>V. unguiculata</i> ; <i>B. diazoefficiens</i> - <i>M. atropurpureum</i>	Transcriptomics; metabolomics	(Ampe et al. 2003, Karunakaran et al. 2009, Li et al. 2013, Lardi et al. 2016)
Bacteroid morphology	<i>B. sp.</i> ORS285 - <i>Aeschynomene</i> spp.	Transcriptomics	(Lamouche et al. 2018)
Effect of field growth	<i>B. diazoefficiens</i> - <i>G. max</i>	Proteomics	(Delmotte et al. 2010)
Root versus stem nodules	<i>B. sp.</i> ORS278 - <i>A. indica</i>	Proteomics	(Delmotte et al. 2014)
Drought stress	<i>R. leguminosarum</i> - <i>P. sativum</i>	Proteomics	(Irar et al. 2014)
Other			
Reference free-living proteome	<i>S. meliloti</i> ; <i>M. loti</i> ; <i>B. diazoefficiens</i> ; <i>B. japonicum</i> ; <i>R. tropici</i>	Proteomics	(Kajiwarra et al. 2003, Djordjevic et al. 2007, da Silva Batista et al. 2010, Gomes et al. 2012b, Kumar et al. 2013, Gomes et al. 2014)
Phosphorylated free-living proteome	<i>S. meliloti</i>	Proteomics	(Liu et al. 2015)
Effects of genome reduction	<i>S. meliloti</i> ; <i>R. leguminosarum</i> bv. <i>trifolii</i>	Transcriptomics; proteomics; metabolomics	(Guerreiro et al. 1998, Chen et al. 2000b, Fei et al. 2016)
Growth with different carbon sources	<i>S. meliloti</i> ; <i>R. leguminosarum</i> bv. <i>viciae</i>	Metabolomics; fluxomics	(Barsch et al. 2004, Fuhrer et al. 2005, Terpolilli et al. 2016)
Membrane lipid composition	<i>S. meliloti</i>	Lipidomics	(Basconcello et al. 2009b)

Table 4. Summary of the fields covered in this review.

Field	Goal	Key lessons
Socio-microbiology	Study the interactions between rhizobia and other organisms that influence environmental growth and the symbiotic process.	Rhizobia interact with many organisms (eukaryotes, prokaryotes, viruses), influencing symbiotic efficiency and competition for nodule occupancy.
Genetics and molecular biology	Characterize the precise function of individual genes and their role in complex biological processes including symbiosis.	Laid the groundwork for modern studies with rhizobia, establishing the fundamentals of the symbiotic interactions.
High-throughput functional genomics	Perform genome-scale analyses to identify genomic elements that contribute to a particular process, growth in a specific environment, or symbiosis.	Elucidated countless loci of relevance to symbiosis or other phenotypes (e.g., rhizosphere growth), highlighting the complexity of these processes.
Large-scale genome manipulation	Generate massive addition or deletions in the genomic content of the cell, and characterize the phenotypic effects and symbiotic effects.	Genes related to symbiosis are spread throughout the genome, and gain of classical symbiotic genes is not necessary sufficient for symbiosis.
Genomics	Compare genome content between strains or species to identify loci likely contributing to a particular phenotype, such as symbiosis.	Rhizobia show massive genome variations that could be related to symbiotic efficiency, and it appears that no gene is common and specific to rhizobia.
Transcriptomics	Identify changes in gene expression across environments or during symbiosis to gain insights into important functions and regulatory mechanisms.	Major transcriptional changes occur during symbiosis, and there is a constant transcriptional rewiring throughout the developmental process.
Proteomics	Identify changes in protein levels across environments or during symbiosis to gain insights into important functions and regulatory mechanisms.	Have provided information into the biochemical pathways that are active during symbiosis, and the roles of proteins and ncRNAs in free-living cells.
Metabolite analysis	Identify changes in metabolite concentrations, or reaction fluxes, across environments or during symbiosis to gain insights into metabolism.	Has been used to identify metabolic pathways active in symbiosis, and has been instrumental in studying the genetics of membrane lipid biosynthesis.
Metabolic modelling	Use computer simulations to predict effect of genomic or environmental alterations on metabolism, and to integrate multi-level omics data.	Used to predict changes in metabolism during shifts in environments and during symbiosis, but application to rhizobia has so far been limited.
Synthetic biology	Engineer cells for the purpose of studying a biological process, or for engineering improved abilities, such as greater symbiotic efficiency.	Has demonstrated the feasibility of at least the initial steps of engineering improved rhizobial inoculants and of transferring BNF directly to plants.




Figure 1. Phylogenetic distribution of biological nitrogen fixation. A maximum likelihood phylogeny of the α - and β -proteobacteria based on 23 highly conserved proteins (Frr, NusA, RplB, RplC, RplD, RplK, RplL, RplM, RplN, RplP, RplS, RplT, RpmA, RpoB, RpsB, RpsC, RpsE, RpsI, RpsJ, RpsK, RpsM, RpsS, Tsf). Species with at least one strain carrying the *nifHDK* nitrogen fixation genes are coloured red (sample genera are indicated), while species with at least one strain carrying the *nifHDK* nitrogen fixation genes and the *nodABC* nodulation genes are coloured blue (the genera and families are indicated). No strains have the nodulation genes but lack the nitrogen fixation genes. Branch lengths for the taxa that include the genera *Tremblaya*, *Hodgkinia*, and *Liberibacter* (light grey) were shortened for presentation. The complete phylogeny, including bootstrap values, is provided in File S2. See the Methods in File S1 for details on phylogeny construction and gene identification (diCenzo et al. 2017c, diCenzo et al. 2018b).

83x62mm (300 x 300 DPI)




Figure 2. Visualizing the rhizobium – legume symbiosis. (A) A picture of *Medicago alba* (white sweet clover) plants grown in nitrogen free conditions. Plants in columns 1-3 and 5 are inoculated with nitrogen-fixing *S. meliloti*; plants in columns 4 and 6 are inoculated with a *S. meliloti* mutant unable to fix nitrogen; and plants in column 7 are not inoculated with *S. meliloti*. (B) *Medicago sativa* (alfalfa) root nodules, induced by *S. meliloti*. On the left are ineffective, white nodules containing a *S. meliloti* mutant unable to fix nitrogen; on the right are effective, pink nodules containing *S. meliloti* cells actively fixing nitrogen. (C) Effective, pink nodules of *Vigna unguiculata* (cowpea) containing *S. fredii* NGR234 cells actively fixing nitrogen. (D, E) Confocal micrographs of alfalfa root nodules filled with *S. meliloti* cells. The nodule sections were stained with a fluorescent nucleic acid binding dye (Syto9); the green colour is indicative of the presence of *S. meliloti*. Scale bar represents 250 μm . (F, G) Transmission electron micrographs of alfalfa root nodules, showing plant cells filled with nitrogen-fixing *S. meliloti* bacteroids. Scale bar represents 25 μm .

83x153mm (300 x 300 DPI)




Figure 3. Schematic representation of the life-cycle of rhizobia. For inoculants, the life-cycle begins in an industrial environment, where the strain is grown in a fermenter. Inoculants are transported to the agricultural field, possibly as a seed coating. In the soil, the rhizobia must exploit nutrients, cope with the stresses, and compete with the indigenous microbiomes. The symbiotic interaction begins following an exchange of signals by the rhizobia and legume. Curling of a legume root hair traps a rhizobium cell, which then penetrates the cell wall of the root hair. The rhizobium proceeds to proliferates in a growing infection thread that progresses to the cortical cells. Here, the rhizobia are released into the cytoplasm of specialized cortical cells, resulting in them being enclosed in a plant derived membrane (peribacteroid membrane). The rhizobia and plant undergo a differentiation process that involves massive transcriptional and metabolic shifts, resulting in the formation of a nitrogen-fixing nodule. Five distinct developmental zones of the nodule are shown, but they are not drawn to scale: zone I – apical meristem; zone II – infection and differentiation zone; interzone II-III – a small region between zone II and zone III; zone III – the nitrogen-fixing zone; zone IV – the senescence zone found in mature nodules. This figure displays an indeterminate root nodule, formed through a Nod factor dependent mechanism and with infection proceeding through an infection thread. However, we note that none of these features are universal and that many types of nodulation exist; see Masson-Boivin et al. (2009) for a description of the differences.

168x119mm (300 x 300 DPI)




Figure 4. Example output of a Tn-seq study. A *S. meliloti* gene region illustrating the ability of Tn-seq to identify essential genes and environment-specific essential genes is shown. Published Tn-seq data (diCenzo et al. 2018a) for *S. meliloti* grown in a nutritionally complex medium (rich medium) and a nutritionally defined medium (defined medium) were mapped to the *S. meliloti* chromosome. A five-gene region is shown. The location of the transposon insertions in the cell populations recovered from both conditions are indicated by the red bars, with the height of the bar representing the number of reads (log scale). Genes are colour coded based on their fitness classifications.

167x44mm (300 x 300 DPI)




Figure 5. The *S. meliloti* pangenome. The pangenome of 20 *S. meliloti* strains with complete genomes was determined using Roary (Page et al. 2015). There was a median of 6,556 genes (standard deviation 362) per genome, and a total of 17,494 sets of orthologous proteins were detected of which 4,000 were present in all strains. (A) A heatmap showing gene presence (dark blue) or absence (light blue) in each of the 20 strains. A phylogeny built based on the 4,000 core genes is shown on the left, and the strain names are indicated on the right. (B) A histogram displaying the distribution of how many genomes each gene is found within. See the Methods in File S1 for details on pangenome construction.

83x116mm (300 x 300 DPI)




Figure 6. Expression patterns of select *S. meliloti* genes during symbiosis with *M. truncatula*. The spatial expression patterns across a *M. truncatula* nodule of 10 *S. meliloti* genes, each displaying a unique pattern, is shown. Data were taken from the study of Roux et al. (Roux et al. 2014). The nodule zone is indicated along the bottom, and the gene names are given on the right. The plotted data is normalized across each zone; the darker the blue, the higher the expression of the gene in the indicated zone relative to other zones. For example, *aglE* is expressed strongly in the distal zone II but lowly in zone III, whereas *fixU* is primarily expressed in zone III. Colours provide no information on the expression of a gene compared to the other genes shown in the figure. Locations of each nodule zone are shown in Figure 3, with zone II distal and zone II proximal referring to the sections of zone II that are distal and proximal to the root, respectively.

83x39mm (300 x 300 DPI)




Figure 7. In silico predicted flux distributions and reaction essentialities. Example data obtained through metabolic modelling. Results from in silico metabolic modelling of *S. meliloti* metabolism using the iGD726 metabolic reconstruction are shown (diCenzo et al. 2018a). Growth was simulated using (A) glucose or (B) succinate as the sole carbon source. The pathways represent central carbon metabolism, with the exception of the pentose phosphate pathway (for simplicity). The width of the line represents the amount of flux through the reaction; a doubling in the width corresponds to a ten-fold flux increase. Arrows indicate the direction of flux. Colours indicate if the reaction is essential (red) or non-essential (blue). See the Methods in File S1 for details on the modelling procedures used. A version with gene names is provided as Figure S1.

83x60mm (300 x 300 DPI)