



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Timing and mechanisms of sediment accumulation and pedogenesis: Insights from the Po Plain (northern Italy)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Luigi Bruno, Bruno Campo, Irka Hajdas, Wan Hong, Alessandro Amorosi (2022). Timing and mechanisms of sediment accumulation and pedogenesis: Insights from the Po Plain (northern Italy). PALAEOGEOGRAPHY PALAEOCLIMATOLOGY PALAEOECOLOGY, 591, 110881-110881 [10.1016/j.palaeo.2022.110881].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/892863> since: 2024-05-15

*Published:*

DOI: <http://doi.org/10.1016/j.palaeo.2022.110881>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Luigi Bruno; Bruno Campo; Irka Hajdas; Wan Hong; Alessandro Amorosi: Timing and mechanisms of sediment accumulation and pedogenesis: *Insights from the Po Plain (northern Italy)*

PALAEOGEOGRAPHY PALAEOCLIMATOLOGY PALAEOECOLOGY 591. ISSN 0031-0182

DOI: 10.1016/j.palaeo.2022.110881

The final published version is available online at:

<https://dx.doi.org/10.1016/j.palaeo.2022.110881>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

26       **Abstract**

27

28       Paleosol stratigraphy in the Po Plain, reconstructed through sedimentological analysis and  
29 correlation of ca. 170 core data chronologically constrained by 376 radiocarbon dates, revealed  
30 strict relationships between pedogenetic and fluvial processes in alluvial and paralic environments.  
31 Vertically stacked, weakly developed paleosols within Upper Pleistocene and Holocene mud-prone  
32 strata testify to intermittent pedogenesis, periodically interrupted by overbank sedimentation.  
33 Individual paleosols are laterally traceable for tens of km and exhibit A-Bk-Bw, A-Bk or A-Bw profiles.  
34 Stratigraphically ordered <sup>14</sup>C calibrated ages from A horizons testify to slow aggradation during 4-6  
35 thousand years-long exposure periods. Burial ages, with an error of few centuries, are provided by  
36 plant debris at the top of A horizons.

37       Millennial-scale climate oscillations and glacio-eustasy are the main drivers of the pedo-  
38 sedimentary evolution of the area during the last 50 kyr. Pleistocene (P) paleosols developed in well-  
39 drained floodplain environments, dissected by fluvial incision, during relatively warm periods.  
40 Overbank sedimentation and paleosol burial occurred during colder phases. High-sediment-supply  
41 during the Last Glacial Maximum hindered pedogenesis and led to the accumulation of 3-10 m-thick  
42 overbank strata. Widespread soil development (paleosol PH) occurred at the end of Last Glacial  
43 Maximum, following glaciers retreat and afforestation of drainage basins. At distal locations,  
44 paleosol PH was progressively buried under estuarine sediments during the Holocene phases of  
45 post-glacial sea-level rise. Beyond the area of marine influence, burial ages of paleosol PH change  
46 from a place to another without specific spatial trends and reflect upstream fluvial sedimentation  
47 dominated by avulsions and deposition of spatially restricted alluvial units. Holocene (H) paleosols  
48 show a poor correlation potential and laterally variable degree of maturity that reflect avulsive

49 sedimentation patterns and crevassing. Deforestation and land reclamation affected pedo-  
50 sedimentary processes during the last 6 kyr.

51

52

53 **Keywords**

54 Paleosol stratigraphy, pedo-sedimentary evolution, radiocarbon dates, climate change, eustasy,

55 Late Quaternary

56

57

58 **1. INTRO**

59

60 In alluvial settings, the development and burial of soil horizons are intimately related to fluvial  
61 activity (Kraus and Bown, 1993) and governed by a variety of factors. High sediment supply and  
62 fluvial aggradation inhibit pedogenesis (Demko et al., 2004), whereas low sediment supply and  
63 fluvial incision promote soil development in interfluvial areas (Blum and Törnqvist, 2000). Climate  
64 influences the amount of sediment delivered to the basins, modulating the magnitude and type of  
65 precipitations, the vegetation cover of drainage areas and the river's transport capacity  
66 (Vandenbergh, 2003). Alluvial sedimentation can be locally hindered or enhanced by tectonic  
67 movements (Holbrook and Schumm, 1999; Viseras et al., 2003; Benvenuti et al., 2008; Pati et al.,  
68 2019), differential subsidence (Cohen et al., 2003) or by avulsions and other processes related to  
69 the intrinsic dynamics of the fluvial apparatus (Kraus, 1999; Cleveland et al., 2007; Hajek and  
70 Wolinsky, 2012). In coastal settings, fluvial dynamics are strongly influenced by sea-level oscillations  
71 (Atchley et al., 2004; Blum et al., 2013). Incised valleys (and associated interfluvial paleosols) related

72 to glacio-eustasy are a widely recognized trait in the sedimentary record (Shanley and McCabe,  
73 1994; Aitken and Flint, 1996; McCarthy and Plint, 1998; Plint et al., 2001; Gibling et al., 2011).

74 The causal relationships between paleosol formation and alluvial sedimentation have been  
75 reconstructed in ancient alluvial successions, where the geometric relationships between paleosols  
76 and associated fluvial sediment bodies are superbly exposed (Gibling and Bird, 1994; McCarthy and  
77 Plint, 2003; Dubiel and Hasiotis, 2011; Soares et al., 2020). However, the role of distinct controlling  
78 factors on paleosol development and burial can hardly be assessed (Wright and Marriott, 1993;  
79 Atchley et al., 2004), because of the temporal resolution of pre-Quaternary dating methods,  
80 insufficient to describe pedo-sedimentary processes which typically act at millennial time scales  
81 (Rasbury et al., 1997; Hofmann et al., 2017). Furthermore, diagenesis may have substantially  
82 modified soil properties after burial, hindering comparison with modern soils and the reconstruction  
83 of the soil-forming environment (Trendell et al., 2012).

84 Improved techniques of subsurface investigation have recently prompted the study of buried  
85 upper Quaternary alluvial successions (Choi, 2005; Tsatskin et al., 2015; Amorosi et al., 2017a,b). In  
86 this relatively narrow time window, drainage areas, ecosystems and soil-forming environment did  
87 not change significantly. Moreover, the high temporal resolution of Quaternary dating methods  
88 permits to link unequivocally pedogenetic processes to a well-known climate and eustatic history.

89 Radiocarbon is probably the most reliable dating method for the last 50 kyr (Reimer et al., 2020).  
90 However, numerous uncertainties remain in the interpretation of <sup>14</sup>C dates from paleosols. Organic  
91 carbon is continuously incorporated in soils, and subjected to complex, and not totally understood,  
92 biological and physical transformations (Larionova et al., 2017; Meingast et al., 2020). These  
93 processes can act at different velocity and magnitude, depending on climate, morphology of the  
94 exposed surface, parent material, and vegetation (Alexandrovskiy and Chichagova, 1998; Kleber,  
95 2010).

96 This study is based on refined pedostratigraphic reconstructions and on a vast <sup>14</sup>C dataset which  
97 permitted to chronologically constrain the major architectural and facies changes. The primary aim  
98 of this research is to understand the causes and mechanisms of soil development and burial in an  
99 alluvial-deltaic setting and its relation to fluvial and coastal dynamics. To this purpose, we focused  
100 on Upper Pleistocene-Holocene paleosols and on their stratigraphic relationship with older, coeval  
101 and younger pedo-sedimentary units. Secondary objectives are to discuss: (i) the relative role of  
102 global and local controlling factors on soil development and burial; (ii) the potential use (and major  
103 limitations) of radiocarbon to assess the exposure age of paleosols.

104

105

## 106 **2. GEOLOGICAL SETTING**

107

### 108 **2.1. Structural setting and basin-scale stratigraphy**

109

110 The Po sedimentary basin is enclosed in a natural amphitheatre bordered by the Alps, to west  
111 and to the north, and by the Apennines, to the south. The eastward prolongation of the Po Plain is  
112 the Adriatic epicontinental sea (Fig. 1). The most external thrusts of the Southern Alps and of  
113 Apennines, showing opposite vergence, locally face each other beneath the Po Plain (Fantoni et al.,  
114 2004). South-Alpine buried thrusts are arranged in a single wide arc running from Milan to the Garda  
115 Lake (Vannoli et al., 2015). The Apennine front is composed of three tectonically active arcuate  
116 thrust systems: from west to east, the Monferrato, Emilia and Ferrara-Romagna Arcs (Ghielmi et al.,  
117 2013). Thrusts of the Emilia and Ferrara Arcs are sealed beneath a pile of sediments with extremely  
118 variable thicknesses (from a few hundred metres to ca. 8 km; Amadori et al., 2019). These lateral  
119 variations in thickness are controlled by lateral changes in subsidence rates (Campo et al., 2020),

120 ranging between zero and 2 mm/y (Bruno et al., 2020a). The northern margin of the Apennines,  
121 which corresponds to an out-of-sequence system of N-NE verging thrusts (Maestrelli et al., 2018),  
122 has been involved in a generalized uplift since the Early Pleistocene (Argnani et al., 2003).

123 The Po Basin fill shows an overall regressive trend from deep-marine to continental and coastal  
124 units (Ghielmi et al., 2013). Widespread alluvial sedimentation started in the Po Plain around 870  
125 kyr B.P. (Muttoni et al., 2003, 2011; Gunderson et al., 2014). The Po River, flowing in W-E direction,  
126 divides the alluvial plain in two sectors (Fig. 1A). In the northern sector, Alpine tributaries flow  
127 entrenched in Pleistocene sandy units (Ravazzi et al., 2012; Fontana et al., 2014; Bruno et al., 2018).  
128 To the south, paleosol-bearing Pleistocene muds are sealed by a thick (up to 30 m) Holocene alluvial  
129 and paralic succession (Amorosi et al., 2017a, b). Early Holocene estuarine deposits typically onlap  
130 onto Upper Pleistocene alluvial strata, testifying to the progressive drowning of the alluvial plain  
131 after 11.5 cal kyr B.P. (Bruno et al., 2017). Following stabilization of sea-level, the Middle-Late  
132 Holocene records the progressive filling of the Po estuary and the successive delta progradation  
133 (Amorosi et al., 2017b, 2019).

134

## 135 **2.2. Paleosols of the Po Basin**

136

137 Highly weathered rubified soils have been reported from the Po Basin margin (Cremaschi, 1987;  
138 Garzanti et al., 2011) and from isolated hills in the alluvial plain, generated by South-Alpine (Zerboni  
139 et al., 2015) and Apennine blind thrusts (Zuffetti et al., 2018). In depocentral areas, mature paleosols  
140 are replaced by sequences of aggradationally stacked, weakly developed paleosols laterally  
141 traceable for tens of km (Amorosi et al., 2017a). A series of closely spaced paleosols, marking the  
142 MIS 3/MIS 2 transition, has been described as the equivalent of the Interfluvial Sequence Boundary  
143 (McCarty and Plint, 2003) in a high accommodation setting, due to its peculiar geometric

144 relationships with genetically related fluvial-channel bodies (Amorosi et al., 2017a). The most  
145 widespread paleosol reported in the Po Plain marks the Pleistocene-Holocene boundary and  
146 responds to several local names: 'Caranto' in the Venetian Plain (Mozzi et al., 2003), 'Ringhiera  
147 paleosol' in the Romagna area (Ravazzi et al., 2006), 'YD paleosol' in the Po Delta Plain (Amorosi et  
148 al., 2017b; Morelli et al., 2017), 'PH paleosol' in the Bologna urban area (Amorosi et al., 2014; Bruno  
149 et al., 2020b; this name will be used in this paper). Although a wide literature reported the existence  
150 of this pedogenized horizon, the mechanisms and causes of its formation and burial are not yet fully  
151 understood. More discontinuous paleosols have been described from the Holocene succession  
152 (Bruno et al., 2015, 2013).

153

154

### 155 **3. METHODS**

156

157 This work is based on the sedimentological and stratigraphic analysis of core material. Sixty-nine  
158 cores, 30-50 m deep (red circles in Fig. 2), were analysed and sampled for radiocarbon dating. For  
159 each core lithology, grainsize, colour and accessory materials (vegetal and shell macrofossils,  
160 carbonate concretions, redoximorphic features) were reported. The reaction to HCl was tested as a  
161 qualitative indicator of the presence of calcium carbonate. Resistance was measured on fine-grained  
162 sediments with a Pocket Penetrometer (PP). One hundred and seventy-two wood, peat, vegetal-  
163 remains and bulk-soil samples were dated at the KIGAM AMS laboratory (Daejeon, Republic of  
164 Korea) and at the Ion Beam Physics laboratory of the ETH (Zurich, Switzerland, Synal et al., 2007).  
165 Samples were collected fresh from the innermost part of the core and desiccated in a 40° oven. The  
166 humin fraction was separated through acid-alkali-acid (AAA) pre-treatment and AMS dated. Humic



167 acids, extracted from 9 bulk soil samples were also dated. Conventional <sup>14</sup>C ages were calibrated  
168 using OxCal 4.4, with the IntCal 20 curve (Reimer et al., 2020).

169 More than 100 core descriptions from previous works and from the database of the Geological,  
170 Seismic and Soil survey of Regione Emilia-Romagna (yellow circles in Fig. 2) were reinterpreted  
171 following calibration with new cores. These descriptions provide information on lithology, colour,  
172 consistency, reaction to 10% HCl, pocket penetrometer measurement and radiocarbon dates.  
173 Additional 204 radiocarbon dates derive from previously published works (Geological Map of Italy  
174 to scale 1:50.000; Ravazzi et. al., 2006; Cacciari et al., 2017; Marabini and Vai, 2020).

175

176

## 177 **4. RESULTS**

178

### 179 **4.1. Sedimentary facies**

180

181 The sedimentological analysis of the studied cores led to the identification of several facies  
182 associations which belong to alluvial, deltaic, estuarine and shallow-marine depositional  
183 environments. To the purposes of this paper, we will describe in detail four main facies associations,  
184 which are part of the alluvial, upper delta-plain and inner estuary depositional systems. For other  
185 facies associations, the reader is referred to Amorosi et al. (2017b, 2019) and Bruno et al. (2017).

186

#### 187 **4.1.1. Fluvial channel-related facies association (FCR)**

188

189 This facies association is composed of cross-stratified or laminated coarse-to-fine sand with local  
190 mud intercalations (Fig. 3). The occasional presence of floated woods or plant debris is recorded.

191 Marine or lagoonal macrofossils are absent, whereas shell fragments of freshwater gasteropods  
192 are seldom encountered.

193 Based on lithology, sedimentary structures and on the lack of marine and lagoonal macrofossil,  
194 these coarse-grained deposits are interpreted as related to fluvial activity (in-channel deposition,  
195 bar migration, crevassing, proximal overbank; Miall, 1985; Toonen et al., 2012; Aslan, 2013).

196

#### 197 4.1.2. Freshwater swamp and marsh (SM)

198

199 This facies association is dominated by fine-grained material, with abundant wood, vegetal  
200 remains and peat (Fig. 3). Peat horizons, 5-30 cm thick, are composed of poorly decomposed woody  
201 material and are separated by grey (2.5 YR 7.1, 7.5 YR 8/1), peat-free clays. Freshwater fossils were  
202 locally encountered. Consistency is very weak and PP values are  $< 1 \text{ kg/cm}^2$ . Carbonate concretions  
203 and redoximorphic features are absent. This facies association shows poor lateral extent in the  
204 Upper Pleistocene succession. On the contrary, in Holocene distal deposits it is volumetrically  
205 dominant and only locally interrupted by channel-related sands.

206 The grain size of this facies group denotes a low-energy depositional environment. The  
207 abundance of poorly decomposed organic material and the lack of oxidation suggest the setting of  
208 reducing conditions in submerged environments. The latter were likely isolated ponds during the  
209 Late Pleistocene, and swamp-marshes part of wider delta-plain and bayhead delta systems during  
210 the Holocene (Amorosi et al., 2017b; Bruno et al., 2017; Cacciari et al., 2020). Peat horizons  
211 separated by peat-free clays testify to intermittent sedimentation, with peat accumulation during  
212 sedimentation breaks (Hijma and Cohen, 2011; Ishii et al., 2016; Bruno et al., 2019).

213

#### 214 4.1.3. Poorly drained floodplain (PDFP)

215

216 This facies association is composed of grey (7.5 YR 6/1) silty clays with faint lamination given by  
217 millimetre-scale silt intercalations. Soft and isolated carbonate concretions (CC in Fig. 4) and sparse  
218 plant debris (UOM in Fig. 4) were seldom encountered. Redoximorphic features are lacking.  
219 Freshwater gasteropods were seldom encountered. Consistency is weak to firm. Pocket  
220 penetrometer values are in the range of 1-2 kg/cm<sup>2</sup>.

221 Based on grainsize and fossil content, this facies association is interpreted as deposited in a distal  
222 fluvial environment, occasionally subjected to river flooding. Lack of redoximorphic features  
223 suggests a groundwater table fluctuating close to the topographic surface (poorly drained  
224 floodplain). The amount, distribution and type of secondary calcite are consistent with this  
225 interpretation.

226

#### 227 4.1.4. Well-drained floodplain (WDFP)

228

229 This facies association is composed of firm-to-hard clayey silts and silty clays. Sedimentary  
230 structures are absent. By contrast, polygonal structures, desiccation cracks and rootlets are  
231 commonly observed. A well-expressed horizonation is given by colour and CaCO<sub>3</sub> distribution (Fig.  
232 3). Dark horizons, showing weak or no reaction to HCl alternate with lighter horizons showing strong  
233 reaction to HCl. Carbonate concretions, in the form of coalescent nodules, coatings, filaments and  
234 sub-horizontal bands are observed in lighter horizons. Redoximorphic features, in the form of  
235 orange mottles or bands (RF in Fig. 6), are commonly observed at the mesoscale. Plant debris and  
236 freshwater gasteropods are seldom encountered. PP values are > 2 kg/cm<sup>2</sup>.

237 As for facies PDFP, grainsize and fossil content point to a distal fluvial environment, occasionally  
238 inundated by river waters (Collinson, 1996). The alternation of dark and light horizons is interpreted

239 as the result of multiple phases of soil development interrupted by overbank deposition and  
240 paleosol burial. Dark colour reflects the accumulation and decomposition of organic matter in the  
241 topsoil (Bruno et al., 2020b). Carbonate distribution, with carbonate-free horizons overlying  $\text{CaCO}_3$   
242 enriched horizons, is interpreted as the result of carbonate leaching from the topsoil (A horizon)  
243 and illuviation to the underlying Bk horizon (Boul et al., 2011). Carbonate mobilization requires the  
244 percolation of meteoric waters in a vadose zone. We, thus, argue that the groundwater was located  
245 a few metres below the topographic surface during subaerial exposure (well-drained floodplain).

246

247

#### 248 **4.2. Paleosol stratigraphy**

249

250 Paleosols were identified at three stratigraphic intervals in the Upper Pleistocene-Holocene  
251 succession of the Po Plain.

252 The lower interval is composed of a series of closely-spaced paleosols (paleosols P in Figs. 4, 5  
253 and 6), whose cumulative thickness is in general  $< 5$  m. Calibrated ages span from the  $^{14}\text{C}$  detection  
254 limit ( $\sim 50$  kyr B.P.) to  $\sim 23$  kyr B.P. (Figs. 5 and 6). These paleosols show A-Bk-Bw or A-Bk profiles. A  
255 horizons, 30-50 cm thick, are decalcified (Bruno et al., 2020b; Amorosi et al., this volume) and  
256 present diffuse organic matter (DOM in Fig. 4) and slight bands of Fe oxides and/or hydroxides (Fig.  
257 3). Undecomposed organic matter was not observed and carbonate concretions have locally been  
258 reported from the base and the top of these horizons. Bk horizons are typified by high  $\text{CaCO}_3$  content  
259 (30-40% of the total volume, Bruno et al., 2020b). Carbonate concretions are abundant as  
260 coalescent nodules or whitish sub-horizontal bands (evolutionary stage II of Gile et al., 1981;  
261 Machette, 1985). Transition to overlying and underlying horizons is in general gradational. Bw  
262 horizons show moderate  $\text{CaCO}_3$  content (Bruno et al., 2020b; Amorosi et al., this volume) and faint

263 mottles of Fe oxides and-or hydroxides. Evidences of clays mobilization (Bt horizons) were not  
264 observed. At distal locations, these paleosols may show less clear horizonation (A-Bw profile) or may  
265 be replaced by coeval peat-bearing swamp deposits (Fig. 5).

266 In the second interval, a single paleosol (paleosol PH, Figs. 4, 5 and 6) is encountered at the top  
267 of a 3-10 m thick sedimentary unit composed of overbank muds and subordinated channel-related  
268 sands (Last Glacial Maximum Sedimentary Unit, LGM SU in Figs. 4, 5 and 6). Textural characters of  
269 paleosol PH vary from proximal to distal locations. At inland locations (core EM15, Fig. 3), paleosol  
270 PH shares many characteristics with older P paleosols: (i) a carbonate-free, brown (10YR 4/4, 3/3) A  
271 horizon, up to 100 cm thick, with DOM and RF; (ii) a lighter Bk horizon with abundant nodules or  
272 bands of secondary calcite (evolutionary stage II); (iii) a Bw horizon showing evidence of incipient  
273 alteration. At more distal locations (cores EM1 and EM3, Fig. 3), paleosol PH exhibit a grey (5Y 4/1,  
274 2.5Y 6/1) decalcified A horizon, 30-50 cm thick, with DOM. Plant debris, absent in the lower part,  
275 become increasingly abundant upsection and, at places, grade into the overlying peats (see core B3,  
276 Fig. 4). The underlying Bk horizon, when present, is relatively thin (a few dm), with secondary calcite  
277 in the form of nodules, coatings or filaments (evolutionary stages I and II). Redoximorphic features  
278 were not observed at the mesoscale along the whole soil profile.

279 In the third interval, dated to the Holocene, paleosols (H1 and H2 in Figs. 4, 5 and 6) have been  
280 observed only at inland locations. At distal locations, they are replaced by coeval estuarine and  
281 deltaic deposits. H paleosols have poor correlation potential and their textural features vary  
282 dramatically from a place to another, without specific spatial trends. In few localities, they show  
283 characteristics similar to those of their Pleistocene counterparts. At places, they are barely  
284 identifiable due to a poorly developed horizonation. Archeological artifacts are increasingly  
285 abundant in younger paleosols.

286 Paleosols P and PH bound nearly tabular sediment packages and are locally replaced by coeval  
287 incisions filled with FCR or SM deposits (see cores CNT, RN3 and EM2, Fig. 5). On the contrary,  
288 Holocene paleosols overlie or are overlain by CR facies (see cores MW and 15, Figs. 5 and 6). Alluvial  
289 units delimited by these soil horizons are poorly extended and typically wedge out laterally or  
290 downdip (Figs. 5 and 6).

291

292

### 293 **4.3. Radiocarbon dating of paleosols**

294

295 Upper Quaternary successions provide the great opportunity to link pedo-sedimentary events to  
296 the detailed climatic and eustatic history described in a vast literature (eg. Clark and Huybers, 2009;  
297 Lambeck et al., 2011; Rasmussen et al., 2014). Assessing the timing of soil development and burial  
298 is crucial to disentangle the relative influence of global and local factors. This chapter summarizes  
299 observations deriving from the analysis of a chronological database composed of 376 radiocarbon  
300 dates (160 from paleosols).

301 Along single cores, multiple dating of A horizons yielded stratigraphically ordered ages, spanning  
302 3-5 kyr (see cores B3, RW28 and 13, Figs. 4, 5C and 6). Only in core SPR (Fig. 5C) two samples from  
303 the base and the top of the same A horizon provided similar ages. In distal cores, plant debris from  
304 topmost A horizons yielded ages that are considerably younger (up to 5 kyr) than those from bulk-  
305 soil samples and just a few centuries older than those from overlying SM peats (see cores EM1, EM3  
306 and B3, Figs. 3, 4 and 5). Dates from A horizons indicate aggradation at extremely low rates (0,05-  
307 0,23 mm yr<sup>-1</sup>) during soil development. Accumulation rates above or between paleosols are at least  
308 one order of magnitude greater (> 1 mm/yr). Distinct ages were also obtained from different

309 fractions extracted from the same sample. Particularly, humic acids yielded, in 9 cases out of 10,  
310 ages older than those obtained from the humin fraction (Table 1).

311 Dates from bulk-soil samples change also laterally, along the same paleosol, spanning in a range  
312 of 4-6 kyr (Fig. 3). No specific trend was observed from proximal to distal locations. Ages also change  
313 over short distances, as observed in the Bologna area. In the Two Tower Medieval district, where  
314 distance between cores is just 10-20 m, six samples from paleosol P3 yielded ages in the range of 3-  
315 4 kyr (Fig. 6).

316 The frequency distribution of radiocarbon dates from paleosols (grey rectangles in Fig. 7) shows  
317 a major peak between 14 and 10 kyr B.P., related to the PH paleosol. Samples from the P paleosols  
318 yielded ages clustered around three minor peaks (P1, P2 and P3, Fig. 7), at 46-41, 38-32 and 29-25  
319 cal kyr B.P., respectively. The overall decay with time in the number of  $^{14}\text{C}$  dates across the  
320 Pleistocene simply reflects the fact that older sediments were penetrated by a progressively lower  
321 number of cores. A decline in the number of  $^{14}\text{C}$  dates from soil samples is also recorded towards  
322 younger ages, together with a sharp increase in dates from SM and PDFP deposits. Ages from  
323 Holocene soil samples are clustered around two peaks. The older peak reflects, in part, the youngest  
324 ages from paleosol PH and in part dates from a distinct paleosol (H1 in Figs 4, 5 and 6) observed at  
325 several inland locations. In the Bubano Quarry (BQ in Fig. 5C), 5 m-thick overbank strata separate  
326 H1 from PH. In the Bologna area (core 16, Figs. 4 and 6), paleosol H1 bears Neolithic and Eneolithic  
327 artefacts (Bruno et al., 2013). The second peak (H2, Fig. 7) corresponds to another archeological  
328 level, rich in Late Bronze Age, Iron-Age and Roman rests (IR paleosol of Bruno et al., 2013), which  
329 was buried in large part of the Po Plain during the Middle Ages (see core CRV, Fig. 5A). Locally, this  
330 paleosol is still exposed (Fig. 5). Interestingly, the maximum number of  $^{14}\text{C}$  dates from facies FCR is  
331 recorded between H2 and H1.

332

333

## 334 5. DISCUSSION

335

### 336 5.1 Use and limitations of radiocarbon for the estimation of the exposure age of paleosols

337

338 The use and reliability of radiocarbon dates from paleosols have been widely debated because,  
339 unlike plant macrofossils, which preserve the particular  $^{14}\text{C}$  content of the moment of grow, soils  
340 are open systems where organic carbon is continuously incorporated, humified and-or oxidized to  
341  $\text{CO}_2$  before burial (Orlova and Panychev, 1993; Martin and Johnson, 1995; Gaudinski et al., 2000).  
342 Therefore,  $^{14}\text{C}$  ages from paleosols reflect the mean residence time of the organic carbon plus the  
343 time of burial (Matthews, 1980; Geyh et al., 1983). The amount of organic carbon added to a soil  
344 and the velocity of its transformation processes change from place to place as a function of climate,  
345 vegetation and parent material (Reimer et al., 2020). Moreover, turnover and breakdown of organic  
346 matter may lead to the redistribution of  $^{14}\text{C}$  ratios in different soil fractions. As a result, apparently  
347 conflicting ages can be obtained from the same paleosol (Figs. 5 and 6) and from different soil  
348 fractions (Table 1). Because of this inherent variability, many authors concluded that radiocarbon  
349 dates from soils are unreliable and the concept of “age” cannot be strictly applied to paleosols (Gilet-  
350 Blein et al., 1980; Becker-Heidmann et al., 2002). However, if not subjected to post-burial  
351 contamination (e.g. rootlets), paleosols should yield ages scattered within its formation period. The  
352 estimation of the length of this period, which can attain several millennia, is an intriguing goal for  
353 Quaternary paleosol stratigraphers.

354 The lack of evidence of clay mobilization, the evolutionary stage of secondary calcite and the  
355 limited rubefaction suggest that Po Plain paleosols evolved over relatively short periods.  
356 Radiocarbon ages from deposits below and above paleosols constrain their exposure period within



357 10-12 thousand years (Figs. 5 and 6). Dates from paleosols are within the narrower range of 4-6  
358 millennia and provide a robust constrain for stratigraphic correlations over long distances (Amorosi  
359 et al., 2017a). More complex is the precise assessment of the ages that mark the onset and end of  
360 pedogenesis for each paleosol, even if radiocarbon dates are available from the base and top of A  
361 horizons. Previous work has demonstrated that, although some organic carbon can remain in soils  
362 for millennia (Feng, 2009), the oldest organic fraction accumulated at the onset of pedogenesis may  
363 have been completely replaced by younger organic carbon (Scharpenseel, 1971; Alexandrovskiy and  
364 Chichagova, 1998) introduced by roots in the deepest soil horizons. Thereby, radiocarbon dates  
365 from organic-sediment samples collected from the base of the A horizon may not be representative  
366 of the onset of soil formation (Gilet-Blein et al., 1980; Alexandrovskiy and Chichagova, 1998). Several  
367 factors influence carbon turnover times in soils (Kleber, 2010; Chen et al., 2013; Pandey et al., 2014).  
368 Particularly, carbon renewal is slower in colder climates and in deep soil horizons (Alexandrovskiy  
369 and Chichagova, 1998; Gaudinski et al., 2000; Fontaine et al., 2007). Therefore, for soils that  
370 developed in cold climatic conditions, such as paleosols P3, which formed at the onset of Last Glacial  
371 Maximum, dates from the base of the A horizon should better approximate the onset of  
372 pedogenesis. This assumption is corroborated by dates from core B3 (Figs. 3, 4 and 5), where the  
373 oldest age from paleosols P3 partially overlaps the age from underlying paleosol P2. For paleosols  
374 developed in warmer periods (PH and H paleosols), ages from the base of A horizons may post-date  
375 the onset of pedogenesis.

376 Sediment samples collected from the top of A horizons may be older than burial time, because a  
377 certain amount of time is required for litter humification and mineralization (Chichagova and  
378 Cherkinsky, 1993). At inland locations of our study area, where O horizons are generally not  
379 preserved, burial ages cannot be assessed precisely. Available dates permit to assert, for example,  
380 that paleosol PH was buried after 10.3 kyr B.P. in core 13 (Figs. 4 and 6) and between 14.3 and 8.6

381 kyr B.P. in core 201S4 (Fig. 5). Conversely, at distal locations, plant macrofossils preserved in the  
382 topsoil provides ages that approximate the paleosol burial age with an error of a few centuries. For  
383 example, in EM1, where estuarine sedimentation was already established around 8.1 kyr B.P., burial  
384 of paleosol PH occurred around 8.5 kyr B.P. (Figs. 3, 4 and 5).

385 Age/depth gradients, like those observed in the Po Plain paleosols, have been reported from  
386 several soil profiles (Matthews, 1981; Matthews and Dresser, 1983; Alexandrovskiy and Chichagova,  
387 1998; Becker-Heidmann et al., 2002) and can possibly be explained by the fact that the amount of  
388 carbon added to the topsoil through litter accumulation is in general order of magnitudes greater  
389 than the one introduced in deeper soil horizons via roots exudation (Fontaine et al., 2007). The  
390 measured accumulation rates, incompatible with alluvial sedimentation, suggest that litter  
391 accumulation accounted for soil aggradation during exposure, with a possible contribution of wind-  
392 transported dust. Therefore, paleosols from the Po Plain represent condensation horizons, rather  
393 than depositional hiatuses. On the contrary, sedimentation between pedogenetic phases seems a  
394 quasi-instantaneous process (see ages from P2 and P3 in core B3, Figs 4 and 5). Paleosols  
395 aggradation is also testified by the presence of carbonate concretions in lower A horizons, likely  
396 related to the upward migration of the eluvial-illuvial (A-B) boundary, and by vertically stacked sub-  
397 horizontal bands of secondary calcite in Bk horizons (Fig. 3). Carbonate concretions in upper A  
398 horizons, on the other hand, reflect post-burial illuviation from overlying paleosols. For this reason,  
399 we avoided radiocarbon dating of carbonates and removed secondary calcite in acid steps of the  
400 AAA sample preparation, before AMS analysis. Humic acids are also considered as a mobile  
401 compound, which may yield anomalously young ages due to contamination from overlying strata  
402 (Matthews, 1980). In our case, the oldest ages are provided by humic acids (Table 1). We thus  
403 excluded contamination from overlying strata during each phase of pedogenesis.

404

405

## 406 **5.2. Pedo-sedimentary evolution of the Po Plain and controlling factors**

407

408 With the chronological constraint of 376 radiocarbon dates, the alluvial succession of the  
409 southern Po Plain provides an unprecedented record of pedo-sedimentary events across the last 50  
410 kyr.

411 Aggradationally stacked paleosols 'P' developed during the Late Pleistocene, when the Adriatic  
412 coastline migrated in response to the stepwise sea-level fall toward its lowstand position, ~300 km  
413 away from the modern one (Pellegrini et al., 2018). Among P paleosols, only P3 developed during a  
414 phase of sea-level drop (~30–50m between 30 and 24 cal ka BP, Peltier and Fairbanks, 2006; Fig.  
415 7B). Conversely, P1 and P2 developed in a period characterized by low-amplitude sea-level  
416 oscillations (Chappell, 2002; Fig. 7B). Far from the influence of sea-level change, pedo-alluvial  
417 processes are controlled by the balance between sediment supply and water discharge (Blum and  
418 Törnqvist, 2000). Low sediment supply-discharge ratios result in channel incision and soil  
419 development in the adjacent interfluves, whereas high sediment supply-discharge ratios promote  
420 fluvial aggradation and paleosol burial (Demko et al., 2004). Bruno et al. (2020b) documented that  
421 soil development and burial during the Late Pleistocene was mainly controlled by changes in  
422 vegetation driven by climatic oscillations at the scale of Bond cycles (Bond et al., 1993). Soils started  
423 to form at the onset of Bond cycles, when warm climate conditions favoured the spread of dense  
424 forests (Magri, 1999; Fletcher et al., 2010), which protected the Apennine drainage basins from  
425 erosion (low sediment supply). Soil development was systematically interrupted during the  
426 subsequent cold phases, at the culmination of Bond cycles, corresponding to North Atlantic Heinrich  
427 events (light blue vertical lines in Fig. 7B). These periods record generalized forest retreats (Fletcher  
428 et al., 2010; Helmens, 2014), which enhanced erosion in the drainage basins and sediment transfer  
429 to the alluvial plains. Increased sediment supply promoted the filling of fluvial incisions and the

430 burial of formerly exposed interfluves by overbank sedimentation. Thin sediment packages were  
431 deposited between two successive phases of subaerial exposure (Figs. 3 to 6). These deposits (and,  
432 partly, the underlying paleosols) were modified during the successive pedogenetic phase. A similar  
433 response to millennial-scale climate oscillations has been observed in loess paleosol-bearing  
434 successions (Rousseau et al., 2011, 2007; Moine et al., 2017; Zeeden et al., 2018).

435 The Last Glacial Maximum was characterized by deposition of thick and widespread fluvial-  
436 channel gravel-sand bodies in the Po Plain (Ravazzi et al., 2012; Campo et al., 2016) and in other  
437 alluvial systems worldwide (Lewis et al., 2001; Briant et al., 2005). Alpine glaciers, which extended  
438 down to the Po Plain margin, supplied huge amounts of sediments to the related fluvio-glacial  
439 systems (Fontana et al. 2014). In the Apennine domain, high sediment supply, favoured by a  
440 predominantly herbaceous vegetation cover of drainage basins (Vescovi et al., 2010), resulted in the  
441 deposition of the LGM SU, which bears poor evidence of pedogenesis (Figs. 4, 5 and 6).

442 Above the LGM SU, PH is the most widespread paleosol in the Po Plain. This surface yielded  
443 radiocarbon dates spanning between ~17 and 6 kyr B.P. (Fig. 8), but mostly clustered around 14 and  
444 10 kyr B.P. (Fig.7). The precise assessment of the age of PH formation is hampered by paucity of  
445 dateable material from the underlying LGM SU. However, available dates indicate that the onset of  
446 post-LGM pedogenesis was diachronous across the Po Plain, without a specific spatial trend (Fig. 8).  
447 Whereas in the Bologna area PH started to develop at least 16.8 kyr B.P. (see core RW28, Figs 6 and  
448 8), in the Romagna plain (Bubano Quarry - BQ, in Figs 5C and 8) pedogenesis started after 14.4 kyr  
449 B.P. (Ravazzi et al., 2006). We argue that post-LGM pedogenesis was favoured by decreased  
450 sediment supply related to the progressive afforestation of the area (Pini et al., 2010; Vescovi et al.,  
451 2010; Campo et al., 2020), with diachroneity possibly related to particular local conditions (e.g.  
452 distance from active channels). North of the Po River, in the Alpine tributary system, fluvial incision

453 occurred soon after glacial retreat (~17 kyr B.P., Monegato et al., 2017), due to sediment trapping  
454 in lakes which formed in place of abandoned glacial amphitheatres (Fontana et al., 2014).

455 Interestingly most of the  $^{14}\text{C}$  dates from paleosol PH post-date the Oldest Dryas (OD in Fig. 7).  
456 The clustering of dates around the Younger Dryas in the Po delta plain (Figs. 7A and 8), suggested a  
457 possible control of this cold event on paleosol formation (Amorosi et al., 2017b, Morelli et al., 2017).  
458 The wider dataset shown in this work highlights that pedogenesis started in many sites of the Po  
459 Plain at least during the Bølling and Allerød warm phases (BA in Fig. 7A). Unlike cold phases at the  
460 culmination of pre-LGM Bond cycles, the Younger Dryas cold spell did not lead to paleosol burial.  
461 This event was probably too short-lived to promote the filling of fluvial incisions and overbank  
462 sedimentation. Paleosol development recorded after LGM in other alluvial (Mol, 1997; Choi, 2005;  
463 van Balen et al., 2010; Janssens et al., 2012) and eolian (Kaiser et al., 2009; Hošek et al., 2017)  
464 successions, indicates a likely climate control at a supra-regional spatial scale. Progressively warmer  
465 conditions lead to widespread afforestation and reduced sediment supply (Fletcher et al., 2010;  
466 Helmens, 2014).

467 Whereas Alpine rivers still flow entrenched in Pleistocene deposits, Apennine incisions were filled  
468 during the Early Holocene (see cores RN3 and EM2, Fig. 3C) and paleosol PH was buried soon after.  
469 In the coastal plain, where paleosol PH is encountered at greater depths beneath estuarine deposits,  
470 burial ages show an overall inverse power law with elevation (Fig. 9) that reflects the progressive  
471 drowning of the area during the early Holocene sea-level rise (Bruno et al., 2017). Slight dispersion  
472 of data may reflect post-burial deformation of paleosol PH by differential subsidence and  
473 neotectonics (Amorosi et al., 2021). In the area under marine influence (shades of blue in Fig. 10),  
474 burial ages decrease landwards from ~11.5 to 5.4 kyr B.P.. This overall spatial trend is not recorded  
475 in two isolated areas, where paleosol PH and older paleosols are upbended (see also Fig. 5C). In  
476 cross-sections, the progressive drowning of paleosol PH is testified by onlap terminations of

477 Holocene flooding surfaces (Fig. 5A). These horizons, associated to prominent facies backstepping,  
478 are coeval to the main Holocene melt water pulses (MWP-1b, 1c, 1d; Fig. 7B), which forced the  
479 landward migration of the Po estuary across the study area. The effects of MWP-1A are not recorded  
480 in the study area, which was too far from the coastline at that time (Storms et al., 2008).

481 Pedo-sedimentary features of paleosol PH reflect this evolutionary history. Carbonate  
482 redistribution along the soil profile and relatively low amounts of organic carbon in the A horizons  
483 (Amorosi et al., this volume) are indicative of a low groundwater table (Kalisz et al., 2010), enhanced  
484 by fluvial incision during the early phases of soil development (Tebbens et al., 1999). Preserved plant  
485 debris in the topsoil, radiocarbon dated to the early Holocene (Figs. 4 and 5), reflects the progressive  
486 rise in groundwater table (Bohacs and Suter, 1997) up to the setting of waterlogged conditions and  
487 to the accumulation of peat beds (Diessel, 1992; Richardson and Vepraskas, 2001; Stolt and  
488 Rabenhorst, 2011). The rise of the groundwater table, forced by sea-level rise (Kosters and Suter,  
489 1993; Törnqvist, 1993; Wadsworth et al., 2003), led to the progressive filling of Apennine fluvial  
490 incisions by fine-grained and organic-matter-rich estuarine strata (facies SM, cores RN3 and EM2 in  
491 Fig. 5C). The lack of fluvial sands (facies FCR) at the base of fluvial incisions, suggests that valley  
492 filling was promoted by eustasy rather than by an upstream increase in sediment supply (Kvale and  
493 Archer, 2007).

494 Beyond the area of marine influence, at elevations  $> -3$  m, the burial age of paleosol PH is  
495 substantially unpredictable (Figs. 9 and 10). The dispersion of burial dates in this upstream sector  
496 reflects the dynamics of Holocene sedimentation, dominated by avulsion processes and by the  
497 deposition of poorly extended alluvial units (Figs. 5 and 6). This aspect is particularly evident in the  
498 Bologna area (Fig. 6): in core 16, for example, paleosol PH was already buried around 8.2 kyr B.P.,  
499 when paleosol H1 was still developing (see also Fig. 4); in the nearby core 15, at a distance of just  
500 224 m, the burial of paleosol PH occurred only around 7 kyr B.P.. The transition to an avulsive

501 sedimentation pattern at the onset of the Holocene, recorded in several worldwide fluvial systems  
502 (Phillips, 2011; Stouthamer et al., 2011), is an indirect consequence of the landward migration of  
503 the coastline, which dramatically reduced the area available for sediment storing and lowered rivers  
504 longitudinal gradients. Decreased stream power, and capability to erode banks and create space for  
505 sediment storage, led to river aggradation, crevassing and avulsions (Blum et al., 2013).

506 The discontinuity of Holocene paleosols (Figs. 5 and 6) reflects the limited areal extent of the  
507 sedimentary units on which they developed. Avulsive sedimentation patterns and crevassing also  
508 determined laterally variable maturity of Holocene paleosols. Indeed, soil development was  
509 interrupted only in restricted areas affected by alluvial deposition. In sediment-starved areas, soils  
510 continued to evolve. Lateral changes in soil maturity are observable in modern soils, which started  
511 to develop during the Late Bronze age (areas where H2 is still exposed – Fig. 5), and after the Middle  
512 Ages (areas where H2 is buried - Fig. 5).

513 The peak in radiocarbon dates from fluvial channel-related deposits, centred around 5-6 kyr B.P.  
514 (Fig. 7), corresponds to a generalized re-organization of the river network at the onset of  
515 Neoglaciation (Piovan et al., 2012; Bruno et al., 2015; Amorosi et al., 2017b), which led to the burial  
516 of paleosol H1. Slope instability and enhanced sediment production, recorded in the Northern  
517 Apennines since 5.5 kyr B.P. (Soldati et al., 2006; Picotti and Pazzaglia, 2008), likely reflect  
518 widespread deforestation of the Apennine drainage basins during the early Eneolithic (Cremaschi  
519 and Nicosia, 2012). The progressive return to stable floodplain conditions was paralleled by  
520 improved techniques of land reclamation and management from the late Bronze age to late  
521 Antiquity (Bruno et al., 2013).

522

523

524 **6. CONCLUSIONS**

525

526       Based on the sedimentological analysis and correlation of more than 170 core data, with the  
527 chronological constraint of 376 radiocarbon dates, this study provided new data on: (i) the complex  
528 relationships between pedogenetic, fluvial and costal processes in alluvial and paralic environments;  
529 (ii) the relative influence of global (i.e. climate and eustasy) and local (e.g. river avulsions) factors on  
530 soil development and burial; (ii) the use and limitations of radiocarbon to assess the exposure age  
531 of paleosols.

532       In the Upper Pleistocene-Holocene mud-prone succession of the Po Basin, vertically stacked,  
533 weakly developed paleosols testify to intermittent pedogenesis, periodically interrupted by  
534 overbank sedimentation. Radiocarbon analyses permitted to define the duration of the main phases  
535 of pedogenesis and to compare them to climate and eustatic curves.

536       Radiocarbon dates from paleosols (bulk-soil and plant-debris samples) and from underlying and  
537 overlying sedimentary units indicate exposure periods encompassing 4-6 millennia. This estimate is  
538 consistent with pedological features (e.g. type of secondary calcite, lack of illuvial clay, limited  
539 rubefaction), which denote incipient pedogenesis. Stratigraphically ordered <sup>14</sup>C calibrated ages  
540 within A horizons revealed that Po Plain paleosols are condensation horizons, which experienced  
541 slow aggradation during exposure (0,05-0,23 mm/yr) due to accumulation of plant litter, with a likely  
542 contribution of wind-transported dust.

543       Millennial-scale climate oscillations and glacio-eustasy concurred in driving the pedo-  
544 sedimentary evolution of the Po Plain during the Late Pleistocene. In particular, during the MIS 3,  
545 pedogenetic events were mainly controlled by changes in vegetation driven by climatic oscillations  
546 at the scale of Bond cycles. Paleosols developed in well-drained floodplain environments, dissected  
547 by fluvial incision, during relatively warm periods. Overbank sedimentation and paleosol burial



548 occurred during colder phases. During the Last Glacial Maximum, high-sediment-supply hindered  
549 pedogenesis and led to the accumulation of 3-10 m-thick overbank strata.

550 Widespread soil development (paleosol PH) occurred at the end of the Last Glacial Maximum,  
551 following glaciers retreat and afforestation of drainage basins. Burial of paleosol PH was  
552 diachronous as indicated by radiocarbon dates from vegetal remains preserved at the topsoil. At  
553 distal locations, paleosol PH was progressively drowned and buried under estuarine sediments  
554 during the Holocene phases of post-glacial sea-level rise. Beyond the area of marine influence, burial  
555 ages of paleosol PH change from a place to another without specific spatial trends, reflecting  
556 upstream fluvial sedimentation dominated by avulsions.

557 Holocene (H) paleosols, observed at inland locations, grade seaward into coeval estuarine and  
558 deltaic deposits. These pedogenized horizons show a poor correlation potential and laterally  
559 variable degree of maturity that reflect avulsive sedimentation patterns and widespread crevassing.  
560 Human activities (e.g. deforestation and land reclamation) became an increasingly important factor  
561 affecting pedo-sedimentary processes during the last 6 kyr.

562 This study highlights the potential for paleoenvironmental reconstructions of upper Quaternary  
563 sedimentary successions. The pedo-stratigraphic approach adopted here, relying upon a robust  
564 chronological framework, permitted to assess the temporal and spatial scale of pedogenetic events  
565 and to discern the relative contribution of distinct controlling factors that typically act  
566 contemporaneously.

567

568

## 569 **Acknowledgments**

570 We thank Mantana Maurer and Lukas Wacker and the AMS team of LIP ETH Zurich for their  
571 support with sample preparation and AMS analysis.

572

573

574 **REFERENCES**

575

576 Aitken, J.F., Flint, S.S., 1996. Variable expressions of interfluvial sequence boundaries in the

577 Breathitt Group (Pennsylvanian), eastern Kentucky, USA. *Geol. Soc. London, Spec. Publ.* 104,

578 193–206.

579 Alexandrovskiy, A.L., Chichagova, O.A., 1998. Age of humic substances in paleosols. *Radiocarbon*

580 40, 991–997.

581 Amadori, C., Toscani, G., Di Giulio, A., Maesano, F.E., D’Ambrogi, C., Ghielmi, M., Fantoni, R., 2019.

582 From cylindrical to non-cylindrical foreland basin: Pliocene–Pleistocene evolution of the Po

583 Plain–Northern Adriatic basin (Italy). *Basin Res.* 991–1015. <https://doi.org/10.1111/bre.12369>

584 Amorosi, A., Bruno, L., Rossi, V., Severi, P., Hajdas, I., 2014. Paleosol architecture of a late

585 Quaternary basin – margin sequence and its implications for high-resolution , non-marine

586 sequence stratigraphy. *Glob. Planet. Change* 112, 12–25.

587 <https://doi.org/10.1016/j.gloplacha.2013.10.007>

588 Amorosi, A., Bruno, L., Cleveland, D.M., Morelli, A., Hong, W., 2017a. Paleosols and associated

589 channel-belt sand bodies from a continuously subsiding late quaternary system (Po basin,

590 Italy): New insights into continental sequence stratigraphy. *Bull. Geol. Soc. Am.* 129, 449–463.

591 <https://doi.org/10.1130/B31575.1>

592 Amorosi, A., Bruno, L., Campo, B., Morelli, A., Rossi, V., Scarponi, D., Hong, W., Bohacs, K.M.,

593 Drexler, T.M., 2017b. Global sea-level control on local parasequence architecture from the

594 Holocene record of the Po Plain, Italy. *Mar. Pet. Geol.* 87, 99–111.

595 <https://doi.org/10.1016/j.marpetgeo.2017.01.020>

596 Amorosi, A., Barbieri, G., Bruno, L., Campo, B., Drexler, T.M., Hong, W., Rossi, V., Sammartino, I.,  
597 Scarponi, D., Vaiani, S.C., 2019. Three-fold nature of coastal progradation during the  
598 Holocene eustatic highstand, Po Plain, Italy—close correspondence of stratal character with  
599 distribution patterns. *Sedimentology* 66, 3029–3052.

600 Amorosi, A., Bruno, L., Campo, B., Costagli, B., Hong, W., Picotti, V., Vaiani, S.C., 2021. Deformation  
601 patterns of upper Quaternary strata and their relation to active tectonics, Po Basin, Italy.  
602 *Sedimentology* 68, 402–424. <https://doi.org/10.1111/sed.12784>

603 Amorosi, A., Bruno, L., Campo, B., Di Martino, A., I. Sammartino. Patterns of geochemical variability  
604 across weakly developed paleosol profiles and their role as regional stratigraphic markers (Late  
605 Pleistocene, Po Plain). Submitted to this volume.

606 Argnani, A., Barbacini, G., Bernini, M., Camurri, F., Ghielmi, M., Papani, G., Rizzini, F., Rogledi, S.,  
607 Torelli, L., 2003. Gravity tectonics driven by Quaternary uplift in the Northern Apennines:  
608 Insights from the La Spezia-Reggio Emilia geo-transect. *Quat. Int.* 101–102, 13–26.  
609 [https://doi.org/10.1016/S1040-6182\(02\)00088-5](https://doi.org/10.1016/S1040-6182(02)00088-5)

610 Aslan, A., 2013. Sediments. *Encycl. Quat. Sci. Second Ed.* 663–675. [https://doi.org/10.1016/B978-](https://doi.org/10.1016/B978-0-444-53643-3.00111-4)  
611 [0-444-53643-3.00111-4](https://doi.org/10.1016/B978-0-444-53643-3.00111-4)

612 Atchley, S.C., Nordt, L.C., Dworkin, S.I., 2004. Eustatic control on alluvial sequence stratigraphy: A  
613 possible example from the Cretaceous-Tertiary transition of the Tornillo Basin, Big Bend  
614 National Park, West Texas, U.S.A. *J. Sediment. Res.* 74, 391–404.  
615 <https://doi.org/10.1306/102203740391>

616 Becker-Heidmann, P., Andresen, O., Kalmar, D., Scharpenseel, H.W., Yaalon, D.H., 2002. Carbon  
617 dynamics in Vertisols as revealed by high-resolution sampling. *Radiocarbon* 44, 63–73.  
618 <https://doi.org/10.1017/S0033822200064687>

619 Benvenuti, M., Bonini, M., Moratti, G., Ricci, M., Tanini, C., 2008. Tectonic and climatic controls on

620 historical landscape modifications: The avulsion of the lower Cecina River (Tuscany, central  
621 Italy). *Geomorphology* 100, 269–284. <https://doi.org/10.1016/j.geomorph.2007.12.009>

622 Blum, M.D., Törnqvist, T.E., 2000. Fluvial responses to climate and sea-level change: A review and  
623 look forward. *Sedimentology* 47, 2–48. <https://doi.org/10.1046/j.1365-3091.2000.00008.x>

624 Blum, M., Martin, J., Milliken, K., Garvin, M., 2013. Paleovalley systems: Insights from Quaternary  
625 analogs and experiments. *Earth-Science Rev.* 116, 128–169.  
626 <https://doi.org/10.1016/j.earscirev.2012.09.003>

627 Bohacs, K., Suter, J., 1997. Sequence stratigraphic distribution of coaly rocks: Fundamental  
628 controls and paralic examples. *Am. Assoc. Pet. Geol. Bull.* 81, 1612–1639.  
629 <https://doi.org/10.1306/7834e192-1721-11d7-8645000102c1865d>

630 Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., McManus, J., 1992. andrews J. Huon S., Jantschik  
631 R., clasen S., Simet ch., Tedesco K., Klas M., Bonani G. Ivy S 245–249.

632 Bond, G., Broecker, W., Johnsen, S., McManus, J., Labeyrie, L., Jouzel, J., Bonani, G., 1993. Bond  
633 1993. *Nature* 365, 143–147.

634 Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., DeMenocal, P., Priore, P., Cullen, H.,  
635 Hajdas, I., Bonani, G., 1997. A pervasive millennial-scale cycle in North Atlantic Holocene and  
636 glacial climates. *Science* (80-. ). 278, 1257–1266.

637 Boul, S.W., Southard, R.J., Graham, R.C., McDaniel, P.A., 2011. *Soil genesis and classification:*  
638 *Hoboken*, 6th ed. Wiley-Blackwell, Chichester. <https://doi.org/10.1002/9780470960622>.

639 Briant, R.M., Bateman, M.D., Coope, G.R., Gibbard, P.L., 2005. Climatic control on quaternary  
640 fluvial sedimentology of a Fenland Basin river, England. *Sedimentology* 52, 1397–1423.  
641 <https://doi.org/10.1111/j.1365-3091.2005.00747.x>

642 Bruno, L., Amorosi, A., Curina, R., Severi, P., Bitelli, R., 2013. Human-landscape interactions in the  
643 Bologna area (northern Italy) during the mid-late Holocene, with focus on the Roman period.

644 Holocene 23, 1560–1571. <https://doi.org/10.1177/0959683613499054>

645 Bruno, L., Amorosi, A., Severi, P., Bartolomei, P., 2015. High-frequency depositional cycles within  
646 the late Quaternary alluvial succession of Reno River (Northern Italy). *Ital. J. Geosci.* 134, 339–  
647 354. <https://doi.org/10.3301/IJG.2014.49>

648 Bruno, L., Bohacs, K.M., Campo, B., Drexler, T.M., Rossi, V., Sammartino, I., Scarponi, D., Hong, W.,  
649 Amorosi, A., 2017. Early Holocene transgressive palaeogeography in the Po coastal plain  
650 (northern Italy). *Sedimentology* 64, 1792–1816. <https://doi.org/10.1111/sed.12374>

651 Bruno, L., Piccin, A., Sammartino, I., Amorosi, A., 2018. Decoupled geomorphic and sedimentary  
652 response of Po River and its Alpine tributaries during the last glacial/post-glacial episode.  
653 *Geomorphology* 317, 184–198. <https://doi.org/10.1016/j.geomorph.2018.05.027>

654 Bruno, L., Campo, B., Di Martino, A., Hong, W., Amorosi, A., 2019. Peat layer accumulation and  
655 post-burial deformation during the mid-late Holocene in the Po coastal plain (Northern Italy).  
656 *Basin Res.* 31, 621–639. <https://doi.org/10.1111/bre.12339>

657 Bruno, L., Campo, B., Costagli, B., Stouthamer, E., Teatini, P., Zoccarato, C., Amorosi, A., 2020a.  
658 Factors controlling natural subsidence in the Po Plain. *Proc. Int. Assoc. Hydrol. Sci.* 382, 285–  
659 290. <https://doi.org/10.5194/piahs-382-285-2020>

660 Bruno, L., Marchi, M., Bertolini, I., Gottardi, G., Amorosi, A., 2020b. Climate control on stacked  
661 paleosols in the Pleistocene of the Po Basin (northern Italy). *J. Quat. Sci.* 35, 559–571.  
662 <https://doi.org/10.1002/jqs.3199>

663 Burrato, P., Ciucci, F., Valensise, G., 2003. An inventory of river anomalies in the Po Plain, Northern  
664 Italy: Evidence for active blind thrust faulting. *Ann. Geophys.* 46, 865–882.  
665 <https://doi.org/10.4401/ag-3459>

666 Cacciari, M., Cremonini, S., Marchesini, M., Vianello, G., Antisari, L.V., 2017. When a pedomarker is  
667 lacking: palynological and chemical multianalysis of a lateglacial-holocene soils suite

668 (Bologna). *EQA-International J. Environ. Qual.* 24, 47–73.

669 Cacciari, M., Amorosi, A., Marchesini, M., Kaniewski, D., Bruno, L., Campo, B., Rossi, V., 2020.

670 Linking Holocene vegetation dynamics , palaeoclimate variability and depositional patterns in

671 coastal successions : Insights from the Po Delta plain of northern Italy. *Palaeogeogr.*

672 *Palaeoclimatol. Palaeoecol.* 538, 109468. <https://doi.org/10.1016/j.palaeo.2019.109468>

673 Campo, B., Amorosi, A., Bruno, L., 2016. Contrasting alluvial architecture of Late Pleistocene and

674 Holocene deposits along a 120-km transect from the central Po Plain (northern Italy).

675 *Sediment. Geol.* 341, 265–275.

676 Campo, B., Bruno, L., Amorosi, A., 2020. Basin-scale stratigraphic correlation of late Pleistocene-

677 Holocene (MIS 5e-MIS 1) strata across the rapidly subsiding Po Basin (northern Italy). *Quat.*

678 *Sci. Rev.* 237, 106300. <https://doi.org/10.1016/j.quascirev.2020.106300>

679 Chappell, J., 2002. Sea level changes forced ice breakouts in the Last Glacial cycle: New results

680 from coral terraces. *Quat. Sci. Rev.* 21, 1229–1240. [https://doi.org/10.1016/S0277-](https://doi.org/10.1016/S0277-3791(01)00141-X)

681 [3791\(01\)00141-X](https://doi.org/10.1016/S0277-3791(01)00141-X)

682 Chen, S., Huang, Y., Zou, J., Shi, Y., 2013. Mean residence time of global topsoil organic carbon

683 depends on temperature, precipitation and soil nitrogen. *Glob. Planet. Change* 100, 99–108.

684 <https://doi.org/10.1016/j.gloplacha.2012.10.006>

685 Chichagova, O.A., Cherkinsky, A.E., 1993. Problems in radiocarbon dating of soils. *Radiocarbon* 35,

686 351–362. <https://doi.org/10.1017/S0033822200060355>

687 Choi, K., 2005. Pedogenesis of late Quaternary deposits, northern Kyonggi Bay, Korea: Implications

688 for relative sea-level change and regional stratigraphic correlation. *Palaeogeogr.*

689 *Palaeoclimatol. Palaeoecol.* 220, 387–404. <https://doi.org/10.1016/j.palaeo.2005.02.006>

690 Clark, P.U., Huybers, P., 2009. Global change: Interglacial and future sea level. *Nature* 462, 856–

691 857. <https://doi.org/10.1038/462856a>

692 Cleveland, D.M., Atchley, S.C., Nordt, L.C., 2007. Continental sequence stratigraphy of the Upper  
693 Triassic (Norian-Rhaetian) chinle strata, northern New Mexico, U.S.A.: Allocyclic and  
694 autocyclic origins of paleosol-bearing alluvial successions. *J. Sediment. Res.* 77, 909–924.  
695 <https://doi.org/10.2110/jsr.2007.082>

696 Cohen, K.M., Gouw, M.J.P., Holten, J.P., 2003. Fluvio-deltaic floodbasin deposits recording  
697 differential subsidence within a coastal prism (central Rhine-Meuse delta, the Netherlands).  
698 *Ned. Geogr. Stud.* 40–68. <https://doi.org/10.1002/9781444304350.ch17>

699 Cremaschi, M., 1987. Paleosols and Vetusols in the Central Po Plain (Northern Italy): a study in  
700 Quaternary geology and soil development, Unicopli. ed. *Paleosols and Vetusols in the Central*  
701 *Po Plain (Northern Italy): A Study in Quaternary Geology and Soil Development.*, Milano.

702 Cremaschi, M., Nicosia, C., 2012. Sub-Boreal aggradation along the Apennine margin of the Central  
703 Po Plain: geomorphological and geoarchaeological aspects. *Géomorphologie Reli. Process.*  
704 *Environ.* 18, 155–174.

705 Demko, T.M., Currie, B.S., Nicoll, K.A., 2004. Regional paleoclimatic and stratigraphic implications  
706 of paleosols and fluvial/overbank architecture in the Morrison Formation (Upper Jurassic),  
707 Western Interior, USA. *Sediment. Geol.* 167, 115–135.  
708 <https://doi.org/10.1016/j.sedgeo.2004.01.003>

709 Diessel, C.F.K., 1992. Coal facies and depositional environment, in: *Coal-Bearing Depositional*  
710 *Systems.* Springer, pp. 161–264.

711 Dubiel, R.F., Hasiotis, S.T., 2011. Deposystems, paleosols, and climatic variability in a continental  
712 system: the Upper Triassic Chinle Formation, Colorado Plateau, USA. *From River to Rock Rec.*  
713 *Preserv. Fluv. Sediments their Subseq. Interpret. SEPM (Society Sediment. Geol. Spec. Publ.*  
714 *97, 393–421.*

715 Fantoni, R., Bersezio, R., Forcella, F., 2004. Alpine structure and deformation chronology at the

716 Southern Alps-Po Plain border in Lombardy. *Boll. della Soc. Geol. Ital.* 123, 463–476.

717 Feng, Y., 2009. Fundamental considerations of soil organic carbon dynamics: a new theoretical  
718 framework. *Soil Sci.* 174, 467–481.

719 Fletcher, W.J., Sánchez Goñi, M.F., Allen, J.R.M., Cheddadi, R., Combourieu-Nebout, N., Huntley,  
720 B., Lawson, I., Londeix, L., Magri, D., Margari, V., Müller, U.C., Naughton, F., Novenko, E.,  
721 Roucoux, K., Tzedakis, P.C., 2010. Millennial-scale variability during the last glacial in  
722 vegetation records from Europe. *Quat. Sci. Rev.* 29, 2839–2864.  
723 <https://doi.org/10.1016/j.quascirev.2009.11.015>

724 Fontaine, S., Barot, S., Barré, P., Bdioui, N., Mary, B., Rumpel, C., 2007. Stability of organic carbon  
725 in deep soil layers controlled by fresh carbon supply. *Nature* 450, 277–280.  
726 <https://doi.org/10.1038/nature06275>

727 Fontana, A., Mozzi, P., Marchetti, M., 2014. Alluvial fans and megafans along the southern side of  
728 the Alps. *Sediment. Geol.* 301, 150–171. <https://doi.org/10.1016/j.sedgeo.2013.09.003>

729 Garzanti, E., Vezzoli, G., Andò, S., 2011. Paleogeographic and paleodrainage changes during  
730 Pleistocene glaciations (Po Plain, Northern Italy). *Earth-Science Rev.* 105, 25–48.  
731 <https://doi.org/10.1016/j.earscirev.2010.11.004>

732 Gaudinski, J., Trumbore, S., Davidson, E., 2000. Soil carbon cycling in a temperate forest:  
733 radiocarbon-based estimates of residence times, .... *Biogeochemistry* 51, 33–69.

734 Geyh, M.A., Roeschmann, G., Wijmstra, T.A., Middeldorp, A.A., 1983. The Unreliability of 14 C  
735 Dates Obtained from Buried Sandy Podzols . *Radiocarbon* 25, 409–416.  
736 <https://doi.org/10.1017/s0033822200005695>

737 Ghielmi, M., Minervini, M., Nini, C., Rogledi, S., Rossi, M., 2013. Late Miocene-Middle Pleistocene  
738 sequences in the Po Plain - Northern Adriatic Sea (Italy): The stratigraphic record of  
739 modification phases affecting a complex foreland basin. *Mar. Pet. Geol.* 42, 50–81.



740 <https://doi.org/10.1016/j.marpetgeo.2012.11.007>

741 Gibling, M.R., Bird, D.J., 1994. Late Carboniferous cyclothems and alluvial paleovalleys in the  
742 Sydney Basin, Nova Scotia. *Geol. Soc. Am. Bull.* 106, 105–117. [https://doi.org/10.1130/0016-](https://doi.org/10.1130/0016-7606(1994)106<0105:LCCAAP>2.3.CO;2)  
743 [7606\(1994\)106<0105:LCCAAP>2.3.CO;2](https://doi.org/10.1130/0016-7606(1994)106<0105:LCCAAP>2.3.CO;2)

744 Gibling, M.R., Fielding, C.R., Sinha, R., Davidson, S.K., Leleu, S., North, C.P., 2011. Alluvial valleys  
745 and alluvial sequences: towards a geomorphic assessment. *From River to Rock Rec. Preserv.*  
746 *Fluv. Sediments Their Subseq. Interpret.* SEPM, Spec. Publ. 97, 423–447.

747 Gile, L.H., Hawley, J.W., Grossman, R.B., 1981. Soils and geomorphology in the Basin and Range  
748 area of southern New Mexico: Guidebook to the Desert Project, New Mexico, Bureau of  
749 Mines and Mineral Resources Memoir.

750 Gilet-Blein, N., Marien, G., Evin, J., 1980. of and of of and. *Radiocarbon* 22, 919–929.

751 Gunderson, K.L., Pazzaglia, F.J., Picotti, V., Anastasio, D.A., Kodama, K.P., Rittenour, T., Frankel,  
752 K.F., Ponza, A., Berti, C., Negri, A., Sabbatini, A., 2014. Unraveling tectonic and climatic  
753 controls on synorogenic growth strata (Northern Apennines, Italy). *Bull. Geol. Soc. Am.* 126,  
754 532–552. <https://doi.org/10.1130/B30902.1>

755 Hajek, E.A., Wolinsky, M.A., 2012. Simplified process modeling of river avulsion and alluvial  
756 architecture: Connecting models and field data. *Sediment. Geol.* 257–260, 1–30.  
757 <https://doi.org/10.1016/j.sedgeo.2011.09.005>

758 Helmens, K.F., 2014. The last interglacial-glacial cycle (MIS 5-2) re-examined based on long proxy  
759 records from central and northern europe. *Quat. Sci. Rev.* 86, 115–143.  
760 <https://doi.org/10.1016/j.quascirev.2013.12.012>

761 Hijma, M.P., Cohen, K.M., 2011. Holocene transgression of the Rhine river mouth area, The  
762 Netherlands/Southern North Sea: Palaeogeography and sequence stratigraphy.  
763 *Sedimentology* 58, 1453–1485. <https://doi.org/10.1111/j.1365-3091.2010.01222.x>

- 764 Hofmann, F., Reichenbacher, B., Farley, K.A., 2017. Evidence for >5 Ma paleo-exposure of an  
765 Eocene–Miocene paleosol of the Bohnerz Formation, Switzerland. *Earth Planet. Sci. Lett.* 465,  
766 168–175. <https://doi.org/10.1016/j.epsl.2017.02.042>
- 767 Holbrook, J., Schumm, S.A., 1999. Geomorphic and sedimentary response of rivers to tectonic  
768 deformation: A brief review and critique of a tool for recognizing subtle epeirogenic  
769 deformation in modern and ancient settings. *Tectonophysics* 305, 287–306.  
770 [https://doi.org/10.1016/S0040-1951\(99\)00011-6](https://doi.org/10.1016/S0040-1951(99)00011-6)
- 771 Hošek, J., Lisá, L., Hambach, U., Petr, L., Vejrostová, L., Bajer, A., Grygar, T.M., Moska, P., Gottvald,  
772 Z., Horsák, M., 2017. Middle Pleniglacial pedogenesis on the northwestern edge of the  
773 Carpathian basin: A multidisciplinary investigation of the Bíňa pedo-sedimentary section, SW  
774 Slovakia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 487, 321–339.  
775 <https://doi.org/10.1016/j.palaeo.2017.09.017>
- 776 Ishii, Y., Hori, K., Momohara, A., Nakanishi, T., Hong, W., 2016. Middle to late-Holocene decreased  
777 fluvial aggradation and widespread peat initiation in the Ishikari lowland (northern Japan).  
778 *The Holocene* 26, 1924–1938.
- 779 Janssens, M.M., Kasse, C., Bohncke, S.J.P., Greaves, H., Cohen, K.M., Wallinga, J., Hoek, W.Z., 2012.  
780 Climate-driven fluvial development and valley abandonment at the last glacial-interglacial  
781 transition (Oude IJssel-Rhine, Germany). *Geol. en Mijnbouw/Netherlands J. Geosci.* 91, 37–  
782 62. <https://doi.org/10.1017/s001677460000055x>
- 783 Kaiser, K., Hilgers, A., Schlaak, N., Jankowski, M., Kühn, P., Bussemer, S., Przegietka, K., 2009.  
784 Palaeopedological marker horizons in northern central Europe: Characteristics of Lateglacial  
785 Usselo and Finow soils. *Boreas* 38, 591–609. [https://doi.org/10.1111/j.1502-](https://doi.org/10.1111/j.1502-3885.2008.00076.x)  
786 [3885.2008.00076.x](https://doi.org/10.1111/j.1502-3885.2008.00076.x)
- 787 Kalisz, B., Lachacz, A., Glazewski, R., 2010. Transformation of some organic matter components in

788 organic soils exposed to drainage. *Turkish J. Agric. For.* 34, 245–256.  
789 <https://doi.org/10.3906/tar-0905-33>

790 Kleber, M., 2010. What is recalcitrant soil organic matter? *Environ. Chem.* 7, 320–332.  
791 <https://doi.org/10.1071/EN10006>

792 Kosters, E.C., Suter, J.R., 1993. Facies relationships and systems tracts in the late Holocene  
793 Mississippi Delta Plain. *J. Sediment. Petrol.* 63, 727–733. [https://doi.org/10.1306/D4267BCC-](https://doi.org/10.1306/D4267BCC-2B26-11D7-8648000102C1865D)  
794 [2B26-11D7-8648000102C1865D](https://doi.org/10.1306/D4267BCC-2B26-11D7-8648000102C1865D)

795 Kraus, M.J., 1999. Paleosols in clastic sedimentary rocks: Their geologic applications. *Earth Sci. Rev.*  
796 47, 41–70. [https://doi.org/10.1016/S0012-8252\(99\)00026-4](https://doi.org/10.1016/S0012-8252(99)00026-4)

797 Kraus, M.J., Bown, T.M., 1993. Palaeosols and sandbody prediction in alluvial sequences. *Geol.*  
798 *Soc. London, Spec. Publ.* 73, 23–31.

799 Kvale, E.P., Archer, A.W., 2007. Paleovalley fills: Trunk vs. tributary. *Am. Assoc. Pet. Geol. Bull.* 91,  
800 809–821. <https://doi.org/10.1306/11060606022>

801 Lambeck, K., Antonioli, F., Anzidei, M., Ferranti, L., Leoni, G., Scicchitano, G., Silenzi, S., 2011. Sea  
802 level change along the Italian coast during the Holocene and projections for the future. *Quat.*  
803 *Int.* 232, 250–257. <https://doi.org/10.1016/j.quaint.2010.04.026>

804 Larionova, A.A., Maltseva, A.N., Lopes de Gerenyu, V.O., Kvitkina, A.K., Bykhovets, S.S., Zolotareva,  
805 B.N., Kudeyarov, V.N., 2017. Effect of temperature and moisture on the mineralization and  
806 humification of leaf litter in a model incubation experiment. *Eurasian Soil Sci.* 50, 422–431.  
807 <https://doi.org/10.1134/S1064229317020089>

808 Lewis, S.G., Maddy, D., Scaife, R.G., 2001. The fluvial system response to abrupt climate change  
809 during the last cold stage: The Upper Pleistocene River Thames fluvial succession at Ashton  
810 Keynes, UK. *Glob. Planet. Change* 28, 341–359. [https://doi.org/10.1016/S0921-](https://doi.org/10.1016/S0921-8181(00)00083-7)  
811 [8181\(00\)00083-7](https://doi.org/10.1016/S0921-8181(00)00083-7)

812 Machette, M.N., 1985. Calcific soils of the southwestern United States. *Soils Quat. Geol.*  
813 *Southwest. United States Geol. Soc. Am. Spec. Pap.* 203, 1–21.

814 Maestrelli, D., Benvenuti, M., Bonini, M., Carnicelli, S., Piccardi, L., Sani, F., 2018. The structural  
815 hinge of a chain-foreland basin: Quaternary activity of the Pede-Apennine Thrust front  
816 (Northern Italy). *Tectonophysics* 723, 117–135. <https://doi.org/10.1016/j.tecto.2017.12.006>

817 Magri, D., 1999. Late Quaternary vegetation history at Lagaccione near Lago di Bolsena (central  
818 Italy). *Rev. Palaeobot. Palynol.* 106, 171–208. [https://doi.org/10.1016/S0034-6667\(99\)00006-](https://doi.org/10.1016/S0034-6667(99)00006-8)  
819 [8](https://doi.org/10.1016/S0034-6667(99)00006-8)

820 Marabini, S., Vai G.B., 2020. *Carta Geologica della Pianura tra Imola e Ravenna*, Thèodolite Editore,  
821 Imola, 81pp. (in Italian).

822 Martin, C.W., Johnson, W.C., 1995. Variation in radiocarbon ages of soil organic matter fractions  
823 from late Quaternary buried soils. *Quat. Res.* 43, 232–237.

824 Matthews, J.A., 1980. Some Problems and Implications of 14C Dates from a Podzol Buried beneath  
825 an End Moraine at Haugabreen , Southern Norway Author ( s ): John A . Matthews Published  
826 by : Taylor & Francis , Ltd . on behalf of the Swedish Society for Anthropology and Geography  
827 62, 185–208.

828 Matthews, J.A., 1981. Natural 14 C age/depth gradient in a buried soil. *Naturwissenschaften* 68,  
829 472–474.

830 Matthews, J.A., Dresser, P.Q., 1983. Intensive 14C dating of a buried palaeosol horizon. *Geol.*  
831 *Föreningen i Stock. Förhandlingar* 105, 59–63.

832 McCarthy, P.J., Plint, A.G., 1998. Recognition of interfluvial sequence boundaries: integrating  
833 paleopedology and sequence stratigraphy. *Geology* 26, 387–390.  
834 [https://doi.org/10.1130/0091-7613\(1998\)026<0387:ROISBI>2.3.CO;2](https://doi.org/10.1130/0091-7613(1998)026<0387:ROISBI>2.3.CO;2)

835 McCarthy, P.J., Plint, A.G., 2003. Spatial variability of palaeosols across Cretaceous interfluvial in

836 the Dunvegan Formation, NE British Columbia, Canada: Palaeohydrological,  
837 palaeogeomorphological and stratigraphic implications. *Sedimentology* 50, 1187–1220.  
838 <https://doi.org/10.1111/j.1365-3091.2003.00600.x>

839 Meingast, K.M., Grunert, B.K., Green, S.A., Kane, E.S., Khademimoshgenani, N., 2020. Insights on  
840 dissolved organic matter production revealed by removal of charge-transfer interactions in  
841 senescent leaf leachates. *Water (Switzerland)* 12. <https://doi.org/10.3390/W12092356>

842 Miall, A.D., 1985. Architectural-element analysis: a new method of facies analysis applied to fluvial  
843 deposits. *Earth-Science Rev.* 22, 261–308.

844 Moine, O., Antoine, P., Hatté, C., Landais, A., Mathieu, J., Prud'homme, C., Rousseau, D.D., 2017.  
845 The impact of Last Glacial climate variability in west-European loess revealed by radiocarbon  
846 dating of fossil earthworm granules. *Proc. Natl. Acad. Sci. U. S. A.* 114, 6209–6214.  
847 <https://doi.org/10.1073/pnas.1614751114>

848 Mol, J., 1997. Fluvial response to Weichselian climate changes in the Niederlausitz (Germany). *J.*  
849 *Quat. Sci.* 12, 43–60. [https://doi.org/10.1002/\(SICI\)1099-1417\(199701/02\)12:1<43::AID-](https://doi.org/10.1002/(SICI)1099-1417(199701/02)12:1<43::AID-JQS291>3.0.CO;2-0)  
850 [JQS291>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1099-1417(199701/02)12:1<43::AID-JQS291>3.0.CO;2-0)

851 Monegato, G., Scardia, G., Hajdas, I., Rizzini, F., Piccin, A., 2017. The Alpine LGM in the boreal ice-  
852 sheets game. *Sci. Rep.* 7, 1–8. <https://doi.org/10.1038/s41598-017-02148-7>

853 Morelli, A., Bruno, L., Cleveland, D.M., Drexler, T.M., Amorosi, A., 2017. Reconstructing Last Glacial  
854 Maximum and Younger Dryas paleolandscapes through subsurface paleosol stratigraphy: An  
855 example from the Po coastal plain, Italy. *Geomorphology* 295, 790–800.  
856 <https://doi.org/10.1016/j.geomorph.2017.08.013>

857 Mozzi, P., Bini, C., Zilocchi, L., Becattini, R., Lippi, M.M., 2003. Stratigraphy, palaeopedology and  
858 palynology of late Pleistocene and Holocene deposits in the landward sector of the Lagoon of  
859 Venice (Italy), in relation to the Caranto level. *Alp. Mediterr. Quat.* 16, 193–210.

860 Muttoni, G., Carcano, C., Garzanti, E., Ghielmi, M., Piccin, A., Pini, R., Rogledi, S., Sciunnach, D.,  
861 2003. Onset of major Pleistocene glaciations in the Alps. *Geology* 31, 989–992.  
862 <https://doi.org/10.1130/G19445.1>

863 Muttoni, G., Scardia, G., Kent, D. V., Morsiani, E., Tremolada, F., Cremaschi, M., Peretto, C., 2011.  
864 First dated human occupation of Italy at ~0.85Ma during the late Early Pleistocene climate  
865 transition. *Earth Planet. Sci. Lett.* 307, 241–252. <https://doi.org/10.1016/j.epsl.2011.05.025>

866 Orlova, L.A., Panychev, V.A., 1993. The reliability of radiocarbon dating buried soils. *Radiocarbon*  
867 35, 369–377. <https://doi.org/10.1017/S0033822200060379>

868 Pandey, D., Agrawal, M., Singh Bohra, J., Adhya, T.K., Bhattacharyya, P., 2014. Recalcitrant and  
869 labile carbon pools in a sub-humid tropical soil under different tillage combinations: A case  
870 study of rice-wheat system. *Soil Tillage Res.* 143, 116–122.  
871 <https://doi.org/10.1016/j.still.2014.06.001>

872 Pati, P., Verma, A.K., Dash, C., Patel, N.K., Gupta, A., Sharma, V., Jakhmola, R.P., Parkash, B.,  
873 Awasthi, A.K., Saraf, A.K., 2019. Influence of neotectonism on geomorphology and  
874 depositional architecture of the Gandak megafan, middle Ganga plain, India. *Geomorphology*  
875 327, 489–503. <https://doi.org/10.1016/j.geomorph.2018.11.029>

876 Pellegrini, C., Asioli, A., Bohacs, K.M., Drexler, T.M., Feldman, H.R., Sweet, M.L., Maselli, V., Rovere,  
877 M., Gamberi, F., Valle, G.D., Trincardi, F., 2018. The late Pleistocene Po River lowstand wedge  
878 in the Adriatic Sea: Controls on architecture variability and sediment partitioning. *Mar. Pet.*  
879 *Geol.* 96, 16–50. <https://doi.org/10.1016/j.marpetgeo.2018.03.002>

880 Peltier, W.R., Fairbanks, R.G., 2006. Global glacial ice volume and Last Glacial Maximum duration  
881 from an extended Barbados sea level record. *Quat. Sci. Rev.* 25, 3322–3337.  
882 <https://doi.org/10.1016/j.quascirev.2006.04.010>

883 Phillips, J.D., 2011. Universal and local controls of avulsions in southeast Texas Rivers.

884 Geomorphology 130, 17–28. <https://doi.org/10.1016/j.geomorph.2010.10.001>

885 Picotti, V., Pazzaglia, F.J., 2008. A new active tectonic model for the construction of the Northern  
886 Apennines mountain front near Bologna (Italy). *J. Geophys. Res. Solid Earth* 113, 1–24.  
887 <https://doi.org/10.1029/2007JB005307>

888 Pini, R., Ravazzi, C., Reimer, P.J., 2010. The vegetation and climate history of the last glacial cycle in  
889 a new pollen record from Lake Fimon (southern Alpine foreland, N-Italy). *Quat. Sci. Rev.* 29,  
890 3115–3137. <https://doi.org/10.1016/j.quascirev.2010.06.040>

891 Piovan, S., Mozzi, P., Zecchin, M., 2012. The interplay between adjacent Adige and Po alluvial  
892 systems and deltas in the late Holocene (Northern Italy). *Géomorphologie Reli. Process.*  
893 *Environ.* 18, 427–440.

894 Plint, A.G., Moore, J., Plint, A., Sanders, J., Ulic, D., 2001. stratigraphy : Updip expression Formation  
895 , Alberta foreland. October 11, 1967–2001.

896 Rasbury, E.T., Hanson, G.N., Meyers, W.J., Saller, A.H., 1997. Dating of the time of sedimentation  
897 using U-Pb ages for paleosol calcite. *Geochim. Cosmochim. Acta* 61, 1525–1529.  
898 [https://doi.org/10.1016/S0016-7037\(97\)00043-4](https://doi.org/10.1016/S0016-7037(97)00043-4)

899 Rasmussen, S.O., Bigler, M., Blockley, S.P., Blunier, T., Buchardt, S.L., Clausen, H.B., Cvijanovic, I.,  
900 Dahl-Jensen, D., Johnsen, S.J., Fischer, H., Gkinis, V., Guillevic, M., Hoek, W.Z., Lowe, J.J.,  
901 Pedro, J.B., Popp, T., Seierstad, I.K., Steffensen, J.P., Svensson, A.M., Vallelonga, P., Vinther,  
902 B.M., Walker, M.J.C., Wheatley, J.J., Winstrup, M., 2014. A stratigraphic framework for abrupt  
903 climatic changes during the Last Glacial period based on three synchronized Greenland ice-  
904 core records: Refining and extending the INTIMATE event stratigraphy. *Quat. Sci. Rev.* 106,  
905 14–28. <https://doi.org/10.1016/j.quascirev.2014.09.007>

906 Ravazzi, C., Donegana, M., Vescovi, E., Arpentini, E., Caccianiga, M., Kaltenrieder, P., Londeix, L.,  
907 Marabini, S., Mariani, S., Pini, R., Vai, G.B., Wick, L., 2006. A new Late-glacial site with *Picea*

908 abies in the northern Apennine foothills: An exception to the model of glacial refugia of trees.  
909 Veg. Hist. Archaeobot. 15, 357–371. <https://doi.org/10.1007/s00334-006-0055-9>

910 Ravazzi, C., Deaddis, M., De Amicis, M., Marchetti, M., Vezzoli, G., Zanchi, A., 2012. Évolution de la  
911 plaine centrale du Pô entre les rivières Adda et Serio au cours des 40 derniers millénaires.  
912 Geomorphol. Reli. Process. Environ. 131–154. <https://doi.org/10.4000/geomorphologie.9794>

913 Regione Emilia-Romagna & Eni-Agip, 1998. Riserve idriche sotterranee della Regione Emilia-  
914 Romagna. S.EL.CA, Firenze (*in Italian*).

915 Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M.,  
916 Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J.,  
917 Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C.,  
918 Van Der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M.,  
919 Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich,  
920 R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo,  
921 S., 2020. The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0-55 cal  
922 kBP). Radiocarbon 62, 725–757. <https://doi.org/10.1017/RDC.2020.41>

923 Richardson, J.L., Vepraskas, M.J., 2001. Wetland soilsgenesis, hydrology, landscapes, and  
924 classification.

925 Rousseau, D.D., Sima, A., Antoine, P., Hatté, C., Lang, A., Zöller, L., 2007. Link between European  
926 and North Atlantic abrupt climate changes over the last glaciation. Geophys. Res. Lett. 34, 2–  
927 7. <https://doi.org/10.1029/2007GL031716>

928 Rousseau, D.D., Antoine, P., Gerasimenko, N., Sima, A., Fuchs, M., Hatté, C., Moine, O., Zoeller, L.,  
929 2011. North Atlantic abrupt climatic events of the last glacial period recorded in Ukrainian  
930 loess deposits. Clim. Past 7, 221–234. <https://doi.org/10.5194/cp-7-221-2011>

931 Scharpenseel, H.W., 1971. Radiocarbon dating of soils--problems, troubles, hopes. Paleopedol.



932 Orig. Nat. Dating Paleosols. Pap. 77–88.

933 Shanley, K.W., McCabe, P.J., 1994. Perspectives on the sequence stratigraphy of continental strata.  
934 Am. Assoc. Pet. Geol. Bull. 78, 544–568. [https://doi.org/10.1306/bdff9258-1718-11d7-](https://doi.org/10.1306/bdff9258-1718-11d7-8645000102c1865d)  
935 [8645000102c1865d](https://doi.org/10.1306/bdff9258-1718-11d7-8645000102c1865d)

936 Synal, H.A., Stocker, M., and Suter, M., 2007. MICADAS: A new compact radiocarbon AMS system.  
937 Nuclear Instruments & Methods in Physics Research Section B-Beam Interactions with  
938 Materials and Atoms 259, 7-13. <https://doi.org/10.1016/j.nimb.2007.01.138>

939 Soares, M.V.T., Basilici, G., Lorenzoni, P., Colombera, L., Mountney, N.P., Martinelli, A.G., Ferreira  
940 Mesquita, Á., da Silva Marinho, T., Vázquez García, R.G., Marconato, A., 2020. Landscape  
941 and depositional controls on palaeosols of a distributive fluvial system (Upper Cretaceous,  
942 Brazil). Sediment. Geol. 410, 105774. <https://doi.org/10.1016/j.sedgeo.2020.105774>

943 Soldati, M., Borgatti, L., Cavallin, A., De Amicis, M., Frigerio, S., Giardino, M., Mortara, G.,  
944 Pellegrini, G.B., Ravazzi, C., Surian, N., Tellini, C., Zanchi, A., Alberto, W., Albanese, D., Chelli,  
945 A., Corsini, A., Marchetti, M., Palomba, M., Panizza, M., 2006. Geomorphological evolution of  
946 slopes and climate changes in Northern Italy during the late quaternary: Spatial and temporal  
947 distribution of landslides and landscape sensitivity implications. Geogr. Fis. e Din. Quat. 29,  
948 164–183.

949 Stolt, M.H., Rabenhorst, M.C., 2011. Introduction and historical development of subaqueous soil  
950 concepts. 36: 1-14.

951 Storms, J.E.A., Weltje, G.J., Terra, G.J., Cattaneo, A., Trincardi, F., 2008. Coastal dynamics under  
952 conditions of rapid sea-level rise: Late Pleistocene to Early Holocene evolution of barrier-  
953 lagoon systems on the northern Adriatic shelf (Italy). Quat. Sci. Rev. 27, 1107–1123.  
954 <https://doi.org/10.1016/j.quascirev.2008.02.009>

955 Stouthamer, E., Cohen, K.M., Gouw, M.J.P., 2011. Avulsion and its implications for fluvial-deltaic

956 architecture: insights from the Holocene Rhine-Meuse delta. *SEPM Spec. Publ.* 97, 215–232.

957 Synal HA, Stocker M, and Suter M. 2007. MICADAS: A new compact radiocarbon AMS system.  
958 *Nuclear Instruments & Methods in Physics Research Section B-Beam Interactions with*  
959 *Materials and Atoms* 259: 7-13.

960 Tebbens, L.A., Veldkamp, A., Westerhoff, W., Kroonenberg, S.B., 1999. Fluvial incision and channel  
961 downcutting as a response to Late-glacial and Early Holocene climate change: The lower  
962 reach of the river Meuse (Maas), The Netherlands. *J. Quat. Sci.* 14, 59–75.  
963 [https://doi.org/10.1002/\(SICI\)1099-1417\(199902\)14:1<59::AID-JQS408>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1099-1417(199902)14:1<59::AID-JQS408>3.0.CO;2-Z)

964 Toonen, W.H.J., Kleinhans, M.G., Cohen, K.M., 2012. Sedimentary architecture of abandoned  
965 channel fills. *Earth Surf. Process. Landforms* 37, 459–472. <https://doi.org/10.1002/esp.3189>

966 Törnqvist, T.E., 1993. HOLOCENE ALTERNATION OF MEANDERING AND ANASTOMOSING  
967 Formation Kreftenheye Formation m below Dutch Ordnance Datum ( O . D . ) \_ Twent ~ e F ~  
968 rmation ~ Betuwe and Formation Broek Formation / T ~ wente Formation. *J. Sediment.*  
969 *Petrol.* 63, 683–693.

970 Trendell, A.M., Atchley, S.C., Nordt, L.C., 2012. Depositional and diagenetic controls on reservoir  
971 attributes within a fluvial outcrop analog: Upper Triassic Sonsela member of the Chinle  
972 Formation, Petrified Forest National Park, Arizona. *Am. Assoc. Pet. Geol. Bull.* 96, 679–707.  
973 <https://doi.org/10.1306/08101111025>

974 Tsatskin, A., Sandler, A., Avnaim-Katav, S., 2015. Quaternary subsurface paleosols in Haifa Bay,  
975 Israel: A new perspective on stratigraphic correlations in coastal settings. *Palaeogeogr.*  
976 *Palaeoclimatol. Palaeoecol.* 426, 285–296. <https://doi.org/10.1016/j.palaeo.2015.03.018>

977 van Balen, R.T., Busschers, F.S., Tucker, G.E., 2010. Modeling the response of the Rhine-Meuse  
978 fluvial system to Late Pleistocene climate change. *Geomorphology* 114, 440–452.  
979 <https://doi.org/10.1016/j.geomorph.2009.08.007>

980 Vandenberghe, J., 2003. Climate forcing of fluvial system development: An evolution of ideas.  
981 Quat. Sci. Rev. 22, 2053–2060. [https://doi.org/10.1016/S0277-3791\(03\)00213-0](https://doi.org/10.1016/S0277-3791(03)00213-0)

982 Vannoli, P., Burrato, P., Valensise, G., 2015. The Seismotectonics of the Po Plain (Northern Italy):  
983 Tectonic Diversity in a Blind Faulting Domain. Pure Appl. Geophys. 172, 1105–1142.  
984 <https://doi.org/10.1007/s00024-014-0873-0>

985 Vescovi, E., Ammann, B., Ravazzi, C., Tinner, W., 2010. A new Late-glacial and Holocene record of  
986 vegetation and fire history from Lago del Greppo, northern Apennines, Italy. Veg. Hist.  
987 Archaeobot. 19, 219–233. <https://doi.org/10.1007/s00334-010-0243-5>

988 Viseras, C., Calvache, M.L., Soria, J.M., Fernández, J., 2003. Differential features of alluvial fans  
989 controlled by tectonics or eustatic accommodation space. Examples from the Betic Cordillera,  
990 Spain. Geomorphology 50, 181–202. [https://doi.org/10.1016/S0169-555X\(02\)00214-3](https://doi.org/10.1016/S0169-555X(02)00214-3)

991 Wadsworth, J., Boyd, R., Diessel, C., Leckie, D., 2003. Stratigraphic style of coal and non-marine  
992 strata in a high accommodation setting: Falher Member and Gates Formation (Lower  
993 Cretaceous), western Canada. Bull. Can. Pet. Geol. 51, 275–303.  
994 <https://doi.org/10.2113/51.3.275>

995 Wright, V.P., Marriott, S.B., 1993. The sequence stratigraphy of fluvial depositional systems: the  
996 role of floodplain sediment storage. Sediment. Geol. 86, 203–210.

997 Zeeden, C., Hambach, U., Veres, D., Fitzsimmons, K., Obreht, I., Böskén, J., Lehmkuhl, F., 2018.  
998 Millennial scale climate oscillations recorded in the Lower Danube loess over the last glacial  
999 period. Palaeogeogr. Palaeoclimatol. Palaeoecol. 509, 164–181.  
1000 <https://doi.org/10.1016/j.palaeo.2016.12.029>

1001 Zerboni, A., Trombino, L., Frigerio, C., Livio, F., Berlusconi, A., Michetti, A.M., Rodnight, H., Spötl,  
1002 C., 2015. The loess-paleosol sequence at Monte Netto: a record of climate change in the  
1003 Upper Pleistocene of the central Po Plain, northern Italy. J. Soils Sediments 15, 1329–1350.

1004 <https://doi.org/10.1007/s11368-014-0932-2>


1005 Zuffetti, C., Trombino, L., Zembo, I., Bersezio, R., 2018. Soil evolution and origin of landscape in a  
1006 late Quaternary tectonically mobile setting: The Po Plain-Northern Apennines border in  
1007 Lombardy (Italy). *Catena* 171, 376–397. <https://doi.org/10.1016/j.catena.2018.07.026>

1008

1009

1010 **Figures and captions**

1011




1012

1013 Fig. 1 – A. Geological map of the Po Plain, with location of the study area and of the seismic profile  
1014 of Fig. 1B (modified after Bruno et al., 2019). The extent of Pleistocene alluvial deposits is from  
1015 Fontana et al. (2014). Buried structures are modified after Burrato et al., (2003). B. Interpreted deep  
1016 seismic profile (from Regione Emilia-Romagna & Eni-Agip, 1998, location in Figure 1A), with  
1017 identification of the major blind thrusts (red lines) and main stratigraphic unconformities (lines from  
1018 A–F) within the Po Basin fill (coloured area).

1019







1025

1026 Fig. 3 – Representative photographs of paleosols from cores EM1, EM3 and B3 (location in Fig. 2).

1027 WDFP: well-drained floodplain; PDFP: poorly-drained floodplain; SW: swamp; FCR: fluvial-channel-

1028 related deposits. For pedological horizons A, Bk and Bw, see text. The white dates (cal kyr B.P.) are

1029 from bulk-sediment samples, whereas the yellow ones are from plant macrofossils.

1030

depth (m)		lithology	colour	consistency	accessory material	reaction to HCl	facies association		
6 14720±320 16760±100	RW28	PH	clayey silt	brown (10 YR 4/4)	hard	DOM, RF	weak	WDFP (A horizon)	
				wheat (10 YR 8/4)		CC (3), RF	strong	WDFP (Bk horizon)	
10 10340±120 11880±120 13540±80	13	PH	clayey silt	brown (10 YR 3/3)	hard	DOM, RF	weak	WDFP (A horizon)	
				beige (10 YR 8/2)		CC (1, 2)	strong	WDFP (Bk horizon)	
8 5050±260 8210±160 10 11870±270 12 14 25850±360	16	H1	silty clay	brown (7.5YR 5/1)	firm	DOM, RF, Eneolithic artifacts	weak	WDFP (A horizon)	
			clayey silt	grey (7.5YR 6/1)					WDFP (Bw horizon)
			clayey and sandy silt	grey (7.5YR 6/1) with orange mottles		RF	moderate	FCR	
			silty clay with silt intercalations	grey (7.5YR 6/1)	weak			PDFP	
			PH		dark grey (10YR 3/1)		DOM, UOM on top	weak	WDFP (A horizon)
				clayey silt	grey (5Y 7/1)	firm	CC (2), RF	strong	WDFP (Bk horizon)
14 25850±360	P3			dark grey (5Y 4/1)	hard	DOM, RF	weak	WDFP (A horizon)	
				olive grey (5Y 6/1)		CC (2), RF	strong	WDFP (Bk horizon)	
17 8870±90 19 11660±160	EM20	PH		grey (2.5YR 7/1)	weak	Abundant UOM	weak	SM	
				light grey (2.5YR 8/1)		CC (1) sparse UOM	moderate	PDFP	
				grey (2.5Y 6/1)		DOM, sparse UOM on top	weak	WDFP (A horizon)	
				light grey (2.5YR 8/1)	firm	CC (2), RF	strong	WDFP (Bk horizon)	
17 7870±50 8125±105 8510±50 19 11430±195	EM1	PH		light grey (7.5YR 8/1)	very weak	Abundant UOM, peat, locally DOM	moderate, weak on peat	SM	
				grey (2.5Y 6/1)	firm	DOM, sparse UOM on top	weak	WDFP (A horizon)	
				light grey (2.5YR 8/1)	hard	CC (2)	strong	WDFP (Bk horizon)	
8 7210±30 7380±40 7630±20 7960±40 10 10000±120 12 27820±130 14 32790±440 32630±370	B3	PH		light grey (clay, 5YR 8/1)	very weak	UOM, peat at the base	moderate, weak on peat	SM	
					brown (peat, 2.5Y 3/2)	weak	sparse UOM		SM (O horizon)
					dark grey (5Y 4/1)	firm	DOM	weak	PDFP
									WDFP (A horizon)
				silty clay- sandy silt	grey (10R 7/1) with orange (10 YR 8/6) mottles		RF	moderate	WDFP (Bw horizon)
				silty clay, clayey silt		firm			FCR
									WDFP (Bw horizon)
			P3		dark grey (7.5Y 5/1)	hard	DOM, RF	weak	WDFP (A horizon)
14 32790±440 32630±370	P2			beige-brown (10YR 6/3)		CC (2,3), RF	strong	WDFP (Bk horizon)	
				beige (7.5Y 8/2)	firm	RF		WDFP (Bw horizon)	
				dark grey (10YR 5/1)		DOM, RF	weak	WDFP (A horizon)	

1031

1032

1033


1034

1035

Fig. 4 – Sedimentological characters and radiocarbon ages of paleosols P2, P3, PH and H1 from cores RW28, 16, 17, EM1, EM20 and B3 (for location see Fig. 2). DOM: dissolved organic matter; UOM: undecomposed organic matter; RF: redoximorphic features; CC carbonate concretions (the number indicates the evolutionary stage, from Gile et al., 1981; Machette et al., 1985). WDFP: well-drained




1036 floodplain; PDFP: poorly-drained floodplain; SW: swamp; FCR: fluvial-channel-related deposits. The  
 1037 black dates (cal kyr B.P.) are from bulk-sediment samples, whereas the red ones are from plant  
 1038 macrofossils.  
 1039



1040  
 1041 Fig. 5 – Upper Pleistocene and Holocene stratigraphy of the Po Plain depicted in cross-sections with  
 1042 W-E (A) and S-N (B and C) orientations (location in Fig. 2A). LGM SU: Last Glacial Maximum  
 1043 Stratigraphic Unit.

1044  
 1045




1046

1047 Fig. 6 – Paleosol-based correlations in the Bologna Area. LGM SU: Last Glacial Maximum  
 1048 Stratigraphic Unit. Location of cores in Fig 2.

1049

1050

1051



1052

1053

1054

1055

1056

1057


1058

1059

1060

1061


Fig. 7 – A. Distribution of radiocarbon dates from bulk-soil and plant-macrofossil samples. A major peak in dates from bulk-soil samples corresponds to paleosol PH; five minor peaks are related to paleosols P1, P2, P3, H1 and H2. Notable is the lack of radiocarbon dated soils around 20 cal ky B.P. (LGM). The sharp increase in radiocarbon dates from plant-macrofossils after 10 ky B.P., paralleled by a decline in the number of dates from bulk-soil samples, reflects the onset of estuarine-deltaic sedimentation in the distal Po Plain and the burial of paleosol PH. Red areas correspond to interstadials, according to Bond et al., (1997) and Rasmussen et al. (2014). OD: Oldest Dryas; YD: Younger Dryas. Light blue vertical bars indicate Henrich events (Bond et al., 1992). B. Sea-level oscillations during the last 50 ky.



1062

1063 Fig. 8 – Radiocarbon dates (black numbers, cal ky B.P.) from paleosol PH (bulk soil samples) in the  
 1064 Po Plain (A) and in the Bologna urban area (B). Blue numbers are the youngest ages from the  
 1065 underlying LGM stratigraphic unit. The present-day elevation (metres above modern sea-level) of  
 1066 paleosol PH is depicted through 5 m-spaced isolines. BQ: Bubano Quarry.

1067




1068

1069 Fig. 9 – Burial ages of paleosol PH plotted against its present-day elevation (metres above modern  
 1070 sea-level). Black rectangles are radiocarbon dates from bulk soil samples. Red rectangles are  
 1071 radiocarbon dates from plant macrofossils. Burial ages from samples at elevation < - 3 m (dashed  
 1072 line) show an overall inverse proportionality with elevation. On the contrary, no specific trends are  
 1073 observable in burial-age distribution at elevation > - 3 m.

1074





1075

1076 Fig. 10 – Map showing burial ages (cal kyr B.P.) of paleosol PH in the Po Plain (A) and in the Bologna  
 1077 urban area (B). The shades of blue depict progressively younger burial ages upland. In the green  
 1078 area burial ages do not show specific spatial trends. Red numbers represent burial ages or intervals  
 1079 measured along single cores (black circles). The map also depicts the present-day elevation (metres  
 1080 above modern sea-level) of paleosol PH through 5 m-spaces isolines. Black dots are the countered  
 1081 data points. The dashed line represents the maximum landward extent of Holocene estuarine and  
 1082 delta plain deposits.

1083

1084

Figure 1







Figure 2






Figure 3




Figure 4

depth (m)		lithology	colour	consistency	accessory material	reaction to HCl	facies association
6	<b>RW28</b>						
14720 <sub>±</sub> 220			brown (10 YR 4/4)		DOM, RF	weak	WDFP (A horizon)
16760 <sub>±</sub> 100		PH	clayey silt	hard	CC (1), RF	strong	WDFP (Bk horizon)
8			wheat (10 YR 8/4)				
10	<b>13</b>						
10340 <sub>±</sub> 120			brown (10 YR 3/3)		DOM, RF	weak	WDFP (A horizon)
11880 <sub>±</sub> 120		PH	clayey silt	hard	CC (1, 2)	strong	WDFP (Bk horizon)
12			beige (10 YR 8/2)				
13540 <sub>±</sub> 80							
8	<b>16</b>						
5050 <sub>±</sub> 260		H1	silty clay	firm	DOM, RF, Eneolithic artifacts	weak	WDFP (A horizon)
8210 <sub>±</sub> 160			clayey silt		RF		WDFP (Bw horizon)
			clayey and sandy silt			moderate	FCR
			silty clay with silt intercalations	weak			PDFP
10			grey (7.5YR 6/1) with orange mottles				
11870 <sub>±</sub> 270		PH	dark grey (10YR 3/1)		DOM, UOM on top	weak	WDFP (A horizon)
12			grey (5Y 7/1)	firm	CC (2), RF	strong	WDFP (Bk horizon)
			clayey silt				
14		P3	dark grey (5Y 4/1)	hard	DOM, RF	weak	WDFP (A horizon)
25850 <sub>±</sub> 200			olive grey (5Y 6/1)		CC (2), RF	strong	WDFP (Bk horizon)
17	<b>EM20</b>		grey (2.5YR 7/1)	weak	Abundant UOM	weak	SM
8870 <sub>±</sub> 90			light grey (2.5YR 8/1)		CC (1)	moderate	PDFP
			grey (2.5Y 6/1)		sparse UOM		
10		PH	light grey (2.5YR 8/1)	firm	DOM, sparse UOM on top	weak	WDFP (A horizon)
11660 <sub>±</sub> 160			light grey (2.5YR 8/1)		CC (2), RF	strong	WDFP (Bk horizon)
17	<b>EM1</b>		light grey (7.5YR 8/1)	very weak	Abundant UOM, peat, locally DOM	moderate, weak on peat	SM
7870 <sub>±</sub> 10			grey (2.5Y 6/1)	weak			
8125 <sub>±</sub> 105			light grey (2.5YR 8/1)	firm	DOM, sparse UOM on top	weak	WDFP (A horizon)
8510 <sub>±</sub> 50			light grey (2.5YR 8/1)	hard	CC (2)	strong	WDFP (Bk horizon)
10		PH	dark grey (5Y 4/1)	firm	DOM	weak	WDFP (A horizon)
11430 <sub>±</sub> 195			grey (10R 7/1) with orange (10 YR 8/6) mottles		RF	moderate	FCR
8	<b>B3</b>		light grey (clay, 5YR 8/1)	very weak	UOM, peat at the base	moderate, weak on peat	SM
7210 <sub>±</sub> 60			brown (peat, 2.5Y 3/2)				SM (O horizon)
7380 <sub>±</sub> 60			dark grey (5Y 4/1)	weak	sparse UOM		PDFP
7630 <sub>±</sub> 20				firm	DOM	weak	WDFP (A horizon)
7960 <sub>±</sub> 60							WDFP (Bw horizon)
10		PH	grey (10R 7/1) with orange (10 YR 8/6) mottles	firm	RF	moderate	FCR
10000 <sub>±</sub> 120			clayey silt				WDFP (Bw horizon)
12			clayey silt				
27820 <sub>±</sub> 120			dark grey (7.5Y 5/1)	hard	DOM, RF	weak	WDFP (A horizon)
14		P3	beige-brown (10YR 6/3)		CC (2,3), RF	strong	WDFP (Bk horizon)
32790 <sub>±</sub> 60			beige (7.5Y 8/2)	firm	RF		WDFP (Bw horizon)
32630 <sub>±</sub> 170		P2	dark grey (10YR 5/1)		DOM, RF	weak	WDFP (A horizon)

Figure 5






Figure 6




Figure 7




Figure 8






Figure 9





Figure 10





**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: