

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Bezout-like polynomial equations associated with dual univariate interpolating subdivision schemes

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version: Bezout-like polynomial equations associated with dual univariate interpolating subdivision schemes / Gemignani, Luca; Romani, Lucia; Viscardi, Alberto. - In: ADVANCES IN COMPUTATIONAL MATHEMATICS. -ISSN 1019-7168. - STAMPA. - 48:1(2022), pp. 4.1-4.27. [10.1007/s10444-021-09912-4]

Availability: This version is available at: https://hdl.handle.net/11585/844091 since: 2022-01-13

Published:

DOI: http://doi.org/10.1007/s10444-021-09912-4

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Gemignani, L., Romani, L. & Viscardi, A. Bezout-like polynomial equations associated with dual univariate interpolating subdivision schemes. *Adv Comput Math* 48, 4 (2022)

The final published version is available online at <u>https://doi.org/10.1007/s10444-</u> 021-09912-4

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

Noname manuscript No. (will be inserted by the editor)

# Bezout-like polynomial equations associated with dual univariate interpolating subdivision schemes

- <sup>3</sup> Luca Gemignani Lucia Romani Alberto
- 4 Viscardi

6 the date of receipt and acceptance should be inserted later

7 Abstract The algebraic characterization of dual univariate interpolating subdivi-

 $_{\ensuremath{\scriptscriptstyle 8}}$  sion schemes is investigated. Specifically, we provide a constructive approach for

<sup>9</sup> finding dual univariate interpolating subdivision schemes based on the solutions

<sup>10</sup> of certain associated polynomial equations. The proposed approach also makes it

<sup>11</sup> possible to identify conditions for the existence of the sought schemes.

Keywords Bezout equation; Univariate dual subdivision; Higher arity; Interpo lation

<sup>14</sup> Mathematics Subject Classification (2000) 65F05 68W30 65D05 65D17

#### 15 1 Introduction

Subdivision schemes are useful tools for the fast generation of graphs of functions, mooth curves and surfaces by the application of iterative refinements to an ini-

tial set of discrete data. The major fields of application of subdivision schemes

<sup>19</sup> are Computer Graphics and Animation, Computer-Aided Geometric Design and

<sup>20</sup> Signal/Image Processing, but a further motivation for their study is also their

21 close relation to multiresolution analysis and wavelets. The last connection was

22 especially investigated in the case of interpolating subdivision schemes and it was

<sup>23</sup> pointed out that the interpolatory subdivision schemes of Dubuc-Deslauriers [11]

 $_{\rm 24}$  are connected to orthonormal wavelets of Daubechies [6,23]. Interpolating subdi-

 $_{\rm 25}$   $\,$  vision schemes were also deeply studied, because they are considered to be very

<sup>26</sup> efficient in representing smooth curves and surfaces passing through a given set

 $_{\rm 27}~$  of points. In fact, after five or six subdivision iterations only, they are capable of

Luca Gemignani Dipartimento di Informatica, Università di Pisa, Largo Bruno Pontecorvo, 3 - 56127 Pisa, Italy E-mail: luca.gemignani@unipi.it

Lucia Romani, Alberto Viscardi Dipartimento di Matematica and  $AM^2$ , Alma Mater Studiorum Università di Bologna, Piazza di Porta San Donato, 5 - 40126 Bologna, Italy E-mail: lucia.romani@unibo.it, E-mail: alberto.viscardi@unibo.it

providing the refined set of points needed to represent on the screen the desired 28 smooth limit shape interpolating the given data. The main properties of inter-29 polating subdivision schemes were investigated over the past 20 years by several 30 researchers (see, e.g., [12, 15, 18]) and many approaches were proposed to design 31 their refinement rules. However, as far as we are aware, before the papers [25, 26], 32 no one ever tried to construct interpolating subdivision schemes that do not sat-33 isfy the stepwise interpolation property and are thus not defined via refinement 34 rules that at each stage of the iteration leave the previous set of points unchanged. 35 Stepwise interpolating subdivision schemes - also known as primal interpolating 36 subdivision schemes [11, 17] - are defined by finite subdivision masks of odd width 37 that contain as a special submask the sequence  $\boldsymbol{\delta} = \{\delta_{0,j}, j \in \mathbb{Z}\}$ . Differently, mem-38 bers of the most recently introduced class of non-stepwise interpolating subdivision 39 schemes -- also known as dual interpolating subdivision schemes-- are characterized 40 by finite masks with an even number of entries that do not necessarily contain as a 41 special submask the  $\delta$  sequence. A first algorithm to construct dual interpolating 42 quaternary schemes was proposed in [25] and successively extended to arbitrary 43 44 arity greater than two in [26]. Precisely, in the latter it was shown that, under some suitable auxiliary assumptions, the coefficients of the subdivision mask of 45 dual interpolating scheme can be (possibly) determined by the solution of an 46 associated rectangular linear system. This system can be clearly inconsistent for 47 some choices of input data and/or size (length) of the mask. For a given input 48 data set the approach taken in [26] consists of an exhaustive analysis of the as-49 sociated linear systems of increasing sizes in order to identify possible consistent 50 configurations. 51

In this paper we pursue a different method for constructing dual interpolating 52 subdivision schemes based on the reduction of the matrix formulation into a func-53 tional setting to solving a certain Bezout-like polynomial equation. The method 54 makes it possible to address the consistency issues by detecting suitable condi-55 tions on the input data which guarantee the existence of a dual interpolating 56 scheme. Additionally, it yields a full characterization of the set of solutions which 57 can be exploited to fulfil additional demands and properties of the solution mask. 58 From the point of view of applications, such a computational approach allows 59 the user to meet specific requests in terms of polynomial reproduction, support 60 size and regularity. Even though a general result concerning convergence and/or 61 smoothness of a dual interpolating subdivision scheme is not yet available, in all 62 the considered examples the regularity analysis is done via joint spectral radius 63 techniques (see [4, 20, 22]), rather than by means of the restricted spectral radius 64 approach (see, e.g., [3]), and the best Hölder exponent for each scheme is computed 65

 $_{\rm 66}$   $\,$  up to the 15th decimal digit.

#### 67 2 Background and notation

- In this section we briefly recall some needed background on subdivision schemes of arbitrary arity  $m \in \mathbb{N}$ ,  $m \ge 2$ .
- Any linear, stationary subdivision scheme is identified by a *refinement mask*  $a := \{a_i \in \mathbb{R}, i \in \mathbb{Z}\}$  that is usually assumed to have finite support, *i.e.* to satisfy  $a_i = 0$  for  $i \notin [-L, L]$  for suitable L > 0.

The subdivision scheme identified by the mask a consists of the subsequent application of the subdivision operator

$$S_{\boldsymbol{a}} : \ell(\mathbb{Z}) \to \ell(\mathbb{Z}) , \qquad (S_{\boldsymbol{a}} \boldsymbol{p})_i := \sum_{j \in \mathbb{Z}} a_{i-mj} p_j, \quad i \in \mathbb{Z} ,$$

which provides the linear rules determining the successive refinements of the ini-73

tial sequence of discrete data  $\boldsymbol{p} := (p_i \in \mathbb{R}, i \in \mathbb{Z}) \in \ell(\mathbb{Z})$ . Introducing the notation 74  $p^{(0)} := p$ , we can thus describe the subdivision scheme as an iterative method that 75 at the k-th step generates the refined scalar sequence 76

$$\boldsymbol{p}^{(k+1)} := S_{\boldsymbol{a}} \boldsymbol{p}^{(k)}, \qquad k \ge 0.$$
(1)

Attaching the data  $p_i^{(k)}$  generated at the k-th step to the parameter values  $t_i^{(k)}$ with

$$t_i^{(k)} < t_{i+1}^{(k)}, \quad \text{and} \quad t_{i+1}^{(k)} - t_i^{(k)} = m^{-k}, \quad k \ge 0$$

(these are usually set as  $t_i^{(k)} := m^{-k}i$ ) we see that the subdivision process generates denser and denser sequences of data so that a notion of convergence can be estable 77

78

lished by taking into account the piecewise linear function  $P^{(k)}$  that interpolates 79 the data, namely 80

$$P^{(k)}(t_i^{(k)}) = p_i^{(k)}, \qquad P^{(k)}|_{[t_i^{(k)}, t_{i+1}^{(k)}]} \in \Pi_1, \qquad i \in \mathbb{Z}, \quad k \ge 0,$$

where  $\Pi_1$  is the space of linear polynomials. If the sequence of the continuous 81 functions  $\{P^{(k)}, k \ge 0\}$  converges uniformly, then we denote its limit by 82

$$f \boldsymbol{p} := \lim_{k \to \infty} P^{(k)}$$

and say that  $f_{p}$  is the *limit function* of the subdivision scheme based on the rule (1) for the data p [2]. When  $\mathbf{p} = \boldsymbol{\delta}$ ,  $f_{\boldsymbol{\delta}}$  is called *basic limit function*.

The analysis of convergence of a subdivision scheme can be accomplished by studying the properties of the so-called *symbol* of the subdivision mask [14]. The symbol of a finitely supported sequence a is defined as the Laurent polynomial

$$a(z) := \sum_{i \in \mathbb{Z}} a_i z^i, \qquad z \in \mathbb{C} \setminus \{0\}$$

Besides convergence and smoothness, many other properties of a subdivision scheme, 83

like polynomial generation and reproduction, can be checked by investigating al-84 gebraic conditions on the subdivision symbol [7]. While the term polynomial gen-85

eration refers to the capability of the subdivision scheme of providing polynomials 86

as limit functions, with *polynomial reproduction* we mean the capability of a subdi-87 vision scheme of reproducing in the limit exactly the same polynomial from which 88

the data are sampled. The property of polynomial reproduction is very impor-89

tant since strictly connected to the approximation order of the subdivision scheme 90

and to its regularity [5, 16]. With respect to the capability of reproducing poly-91

nomials up to a certain degree, the standard parametrization (corresponding to 92  $_{1}(k)$  $-k_i$  i  $( \mathbb{Z})$  is stimol ahoid ot olr +h Indeed the

<sup>93</sup> the choice 
$$t_i^{(k)} := m^{-k}(i + \sigma/(m-1))$$
 with  $\sigma = a^{(1)}(1)/m$ , turns out to be the recommended

spectral 
$$v_i$$
 is the control of  $v_i$  is the control of  $\sigma \in \mathbb{Z}$  are termed primal, whereas  
selection [8]. The subdivision schemes for which  $\sigma \in \mathbb{Z}$  are termed primal, whereas

the ones for which  $\sigma \in (2\mathbb{Z}+1)/2$  are called *dual*. The target of this work are dual 96 schemes. While dual approximating schemes were investigated extensively (see, 97 e.g., [8,13] and references therein), to the best of our knowledge dual interpolating 98 schemes were only considered in the recent papers [25, 26]. However, as already 99 acknowledged in [25], the open problem treated in these papers was suggested by 100 Malcolm Sabin, who has the merit of being the first who foresaw the existence 101 of dual *m*-ary schemes (with m > 2) that are capable of interpolating the initial 102 data. 103

#### <sup>104</sup> **3** The proposed approach

The aim of this section is to investigate the algebraic characterization of univariate dual interpolating subdivision schemes of arity m. According to the results shown in [26], the construction of such schemes requires as input the desired degree of polynomial reproduction (denoted in the following by d - 1,  $d \in \mathbb{N}$ ) and some samples of the resulting basic limit function  $f_{\delta}$ , i.e.,

$$f_{\delta}\left(\frac{1}{2}+\ell\right) = \varphi\left(\frac{1}{2}+\ell\right), \qquad \forall \ell \in \mathbb{Z},$$
(2)

for a given  $\varphi: (2\mathbb{Z} + 1)/2 \to \mathbb{R}$ . A similar procedure was investigated in [9, 10], where the samples of the basic limit function at the integers were required: here instead the samples at the integers are fixed to be the  $\delta$  sequence and information about the samples at the half-integers are required.

More specifically, in [26] it is seen that taking Fourier transforms on both sides of 114 the refinement equation for the basic limit function  $f_{\delta}$  allows one to describe the 115 mask of dual interpolatory schemes in a matrix setting in terms of the solution 116 of certain bi-infinite Toeplitz-like linear systems in banded form. In this paper we 117 exploit the interplay between the functional and the matrix settings into more 118 details. In particular, from the matrix setting we come back to the functional 119 one by relying upon the connection of Toeplitz-like systems with corresponding 120 Bezout-like polynomial equations. This connection yields a constructive approach 121 to determine the associated symbols. Moreover, the proposed approach also makes 122 it possible to identify conditions for the existence of the sought dual interpolatory 123 schemes. In the following, to simplify the presentation, we distinguish between the 124 odd and even arity cases. 125

<sup>126</sup> 3.1 The odd arity case

Now let us consider the solution of the linear system (35) in [26] for the case where m is an odd integer. The system is defined as follows:

$$M\boldsymbol{a} = \boldsymbol{c}, \qquad M = (\mu_{i,j})_{i,j\in\mathbb{Z}}, \qquad \boldsymbol{c} = (c_i)_{i\in\mathbb{Z}},$$

4

$$\mu_{i,j} = \begin{cases} \varphi\left(\frac{i+1}{2} - j\right), & \text{if } i \in 2m\mathbb{Z}, \\ 1, & \text{if } i \in m(2\mathbb{Z}+1), \ j = \frac{i+1}{2}, \\ 0, & \text{otherwise}, \end{cases}$$
$$c_i = \begin{cases} 1, & \text{if } i = 0, \\ \varphi\left(\frac{i}{2m}\right), & \text{if } i \in m(2\mathbb{Z}+1), \\ 0, & \text{otherwise}. \end{cases}$$

- 130
- By suppressing the zero rows in both M and c we obtain the equivalent linear system

$$\widehat{M}\boldsymbol{a} = \widehat{\boldsymbol{c}}, \qquad \widehat{M} = (\widehat{\mu}_{i,j})_{i,j\in\mathbb{Z}}, \qquad \widehat{\boldsymbol{c}} = (\widehat{c}_i)_{i\in\mathbb{Z}}, \qquad (3)$$

133 where

134

$$\hat{\mu}_{i,j} = \begin{cases} \varphi \left( \frac{im+1}{2} - j \right), & \text{if } \mod(i,2) = 0, \\ 1, & \text{if } \mod(i,2) = 1, \ j = \frac{im+1}{2}, \\ 1, & \text{if } i = 0, \\ \hat{c}_i = \begin{cases} 1, & \text{if } i = 0, \\ \varphi \left( \frac{i}{2} \right), & \text{if } \mod(i,2) = 1, \\ 0, & \text{otherwise.} \end{cases}$$

The interplay between computations with polynomials and Toeplitz-like matrices can be exploited to recast the solution of the linear system (3) in terms of solving an associated Bezout-like polynomial equation. Indeed from the proof of Theorem

 $_{138}$  4.1 in [26] one deduces that the entries of the unknown vector  $\boldsymbol{a}$  satisfy

$$\begin{cases}
\sum_{\alpha \in m(2\mathbb{Z}+1)} \varphi\left(\frac{\alpha}{2m}\right) z^{\alpha} = \sum_{\alpha \in m(2\mathbb{Z}+1)} a_{\frac{\alpha+1}{2}} z^{\alpha}, \\
1 = \sum_{\alpha \in 2m\mathbb{Z}} \sum_{\beta \in \mathbb{Z}} a_{\beta} \varphi\left(\frac{\alpha+1}{2} - \beta\right) z^{\alpha},
\end{cases}$$
(4)

139 which implies

$$\begin{cases} a_{mi+\frac{m+1}{2}} = \varphi\left(\frac{2i+1}{2}\right), & i \in \mathbb{Z}, \\ 1 - \sum_{\alpha \in 2m\mathbb{Z}} \sum_{\beta \in m\mathbb{Z} + \frac{m+1}{2}} a_{\beta}\varphi\left(\frac{\alpha+1}{2} - \beta\right) z^{\alpha} = \\ \sum_{\alpha \in 2m\mathbb{Z}} \sum_{\substack{\beta \in \mathbb{Z} \\ \text{mod } (m,\beta) \neq \frac{m+1}{2}}} a_{\beta}\varphi\left(\frac{\alpha+1}{2} - \beta\right) z^{\alpha}. \end{cases}$$
(5)

The system (5) can be rewritten into a more compact form by using the decom-140 position of  $a(z) = \sum_{i \in \mathbb{Z}} a_i z^i$  that involves the sub-symbols of the scheme given 141

by 142

$$a(z) = \sum_{i=0}^{m-1} a_i(z^m) z^i, \qquad a_\ell(z) = \sum_{i \in \mathbb{Z}} a_{mi+\ell} z^i, \qquad 0 \le \ell \le m-1.$$
(6)

Let us introduce the corresponding decomposition of the Laurent polynomial 143  $\phi(z) = \sum_{\ell \in \mathbb{Z}} \varphi\left(\frac{1}{2} + \ell\right) z^{\ell}$  defined by 144

$$\phi(z) = \sum_{i=0}^{m-1} \phi_i(z^m) z^{-i}, \qquad \phi_\ell(z) = \sum_{i \in \mathbb{Z}} \varphi\left(\frac{2mi+1}{2} - \ell\right) z^i, \qquad 0 \le \ell \le m-1.$$
(7)

The first equation of (5) determines 145

$$a_{\frac{m+1}{2}}(z) = \phi(z). \tag{8}$$

Then the second equation can be read as follows 146

$$1 - a_{\frac{m+1}{2}}(z^m)\phi_{\frac{m+1}{2}}(z^m) = \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} a_i(z^m)\phi_i(z^m)$$

or, equivalently, 147

$$1 - \phi(z)\phi_{\frac{m+1}{2}}(z) = \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} a_i(z)\phi_i(z).$$
(9)

Our computational task is therefore reduced to computing a Laurent polynomial 148

a(z) defined as in (6) satisfying the Bezout-like polynomial equation (9). It is quite 149

natural for convergence and reproducibility issues to impose some other constraints 150 of the form 151

$$a_{i}(1) = 1, \quad 0 \leq i \leq m - 1,$$

$$a(z) = \left(\frac{1 + z + \dots + z^{m-1}}{m}\right)^{d} b(z),$$
(10)

152

for some  $b(z) \in \mathbb{R}[z, z^{-1}]$  with  $b(\xi_k) \neq 0, 1 \leq k \leq m-1$ , where  $\xi_k = e^{2\pi i k/m}$ . Our proposed construction of such a polynomial a(z) works under some additional assumptions on the input data  $\{\varphi((2k+1)/2)\}_{k=-\kappa}^{\kappa-1}$  encoded in the function  $\phi(z)$ . 153 154

More specifically: 155

**ASSUMPTION 1** : We suppose that 156

$$1 - z\phi(z^2) = (z-1)^d \gamma(z), \tag{11}$$

for a certain  $\gamma(z) \in \mathbb{R}[z, z^{-1}]$  the ring of Laurent polynomials in  $z, z^{-1}$  over  $\mathbb{R}$ . 157

**ASSUMPTION 2** : We suppose that 158

$$g(z) := gcd\left\{\phi_0(z), \dots, \phi_{\frac{m-1}{2}}(z), \phi_{\frac{m+3}{2}}(z), \dots, \phi_{m-1}(z)\right\},\$$

divides  $1 - \phi(z)\phi_{\frac{m+1}{2}}(z)$  and is such that  $g(1) \neq 0$ . 159

6

Assumption 1 is necessary in order to achieve polynomial reproduction of order

<sup>161</sup> *d.* Indeed, by definition of polynomial reproduction, we should have, for every <sup>162</sup> polynomial p of degree d - 1,

$$p(x) = \sum_{k \in \mathbb{Z}} p(k) f_{\delta}(x-k), \quad \forall x \in \mathbb{R}.$$

In particular, taking x = 1/2 + i,  $i \in \mathbb{Z}$ , the (compactly supported) vector  $[f_{\delta}(1/2 + \ell)]_{\ell \in \mathbb{Z}}$  defines column-wise a (bandlimited) Toeplitz matrix **T** such

that, for every polynomial p of degree d-1,

$$\mathbf{T} \left[ p\left( k \right) \right]_{k \in \mathbb{Z}} = \left[ p\left( \frac{1}{2} + i \right) \right]_{i \in \mathbb{Z}}.$$

<sup>166</sup> Thus, one can naively define a binary primal interpolating refinement mask as

$$\mathbf{r} \ = \ \{ \ r_i \in \mathbb{R}, \ i \in \mathbb{Z} \ \}, \quad \text{ with } \quad r_i \ = \ \begin{cases} \varphi \left( \frac{1}{2} + \ell \right), \ \text{ if } i = 2\ell + 1 \\ \\ \delta_{0,\ell}, \quad \text{ if } i = 2\ell, \end{cases}$$

which is not guaranteed to be associated with a convergent subdivision scheme,
 but it always satisfies

$$\sum_{i \in \mathbb{Z}} r_i z^i = (z+1)^d \tilde{\gamma}(z), \qquad (12)$$

for some Laurent polynomial  $\tilde{\gamma}(z)$ . Now it is easy to check that, replacing z with -z in (12) and using (7), one indeed obtains Assumption 1, i.e.,  $1 - z\phi(z^2) = (z-1)^d \gamma(z)$ , with  $\gamma(z) = (-1)^d \tilde{\gamma}(-z)$ .

Remark 1 The previous observation is also the reason why a suitable way to construct the starting sequence  $\{\varphi((2k+1)/2)\}_{k\in\mathbb{Z}}$  is using the mask of a binary primal interpolating scheme with the desired reproduction properties. In the following Example 1 and 2 ((26) and (42) respectively) we choose the mask of the binary 6-point Dubuc-Deslauriers interpolating scheme [11] since it forms the shortest symmetric sequence that guarantees polynomial reproduction of order 6.

As for Assumption 2, requiring g(z) to divide  $1 - \phi(z)\phi_{\frac{m+1}{2}}(z)$  is also necessary due to equation (9), while asking  $g(1) \neq 0$  is only a sufficient condition as it will be clear in the following. When 1 is a root of g(z), the construction we propose is still viable but a price has to be paid in terms of polynomial reproduction (see Remark 2).

<sup>183</sup> Under Assumption 1 and Assumption 2 our composite approach for computing <sup>184</sup> a(z) proceeds by the following steps. The first step consists of determining the <sup>185</sup> values  $a_i^{(s)}(1), 0 \le i \le m-1, s = 0, ..., d-1$ . From (10) one gets immediately <sup>186</sup>  $a_i^{(0)}(1) = a_i(1) = 1, 0 \le i \le m-1$ . Due to (8) and Assumption 1, we have that

$$1 - z\phi(z^2) = 1 - za_{\frac{m+1}{2}}(z^2) = (z-1)^d \gamma(z),$$

from which we can compute the values of  $a_{\frac{m+1}{2}}(z)$  and its derivatives at z = 1.

**Theorem 1** Under Assumption 1, it holds

$$\begin{cases} a_{\frac{m+1}{2}}(1) = \phi(1) = 1, \\ \\ a_{\frac{m+1}{2}}^{(k)}(1) = \phi^{(k)}(1) = (-1)^k \frac{(2k-1)!!}{2^k}, \quad 1 \le k \le d-1. \end{cases}$$

189 Proof Substituting  $z = \sqrt{w}$  in (11), we get

$$\phi(w) - w^{-1/2} = \frac{(1 - \sqrt{w})^d (-1)^{d+1} \gamma(\sqrt{w})}{\sqrt{w}}.$$

<sup>190</sup> The proof easily follows by differentiating this relation at w = z = 1.

<sup>191</sup> The remaining unknowns  $a_i^{(s)}(1), 0 \leq i \leq m-1, i \neq (m+1)/2, s = 1, \dots, d-1,$ 

<sup>192</sup> are computed by solving the linear system obtained by differentiation of (10). <sup>193</sup> Specifically, by differentiating s times the expression of a(z) in (6) with respect to <sup>194</sup> the variable z we find that

$$a^{(s)}(z) = \sum_{i=0}^{m-1} \sum_{p=0}^{s} \frac{a_i^{(p)}(z^m)}{p!} \left( \sum_{j=max\{s-i,p\}}^{s} \binom{s}{j} A_{j,p}(z) \frac{i!}{(i-(s-j))!} z^{i-(s-j)} \right), \quad (13)$$

where  $A_{j,p}(z)$  are polynomials defined by Hoppe's formula (see, e.g., [21]) for the differentiation of composite function according to

$$A_{j,p}(z) = \sum_{\ell=0}^{j} {\binom{p}{\ell}} (-f(z))^{p-\ell} \frac{d^{j}}{dz^{j}} (f(z))^{\ell}, \qquad f(z) = z^{m}$$

If  $\xi_k = e^{2\pi i k/m}$ ,  $1 \leq k \leq m-1$ , are the *m*-th roots of unity, then from (10) it follows that  $a^{(s)}(\xi_k) = 0$ ,  $s = 0, \ldots, d-1$ ,  $1 \leq k \leq m-1$ . In the view of (13) this implies that the values  $a_i^{(s)}(1)$ ,  $0 \leq i \leq m-1$ ,  $i \neq (m+1)/2$ ,  $s = 1, \ldots, d-1$ , can be computed recursively by solving

$$\sum_{i=0}^{m-1} \sum_{p=0}^{s} \frac{a_i^{(p)}(1)}{p!} \left( \sum_{j=max\{s-i,p\}}^{s} {s \choose j} A_{j,p}(\xi_k) \frac{i!}{(i-(s-j))!} \xi_k^{i-(s-j)} \right) = 0,$$

with  $1 \leq k \leq m-1$ . The system can be expressed in matrix form as

$$m^{s} \mathcal{M}\left[a_{0}^{(s)}(1), \dots, a_{\frac{m-1}{2}}^{(s)}(1), a_{\frac{m+3}{2}}^{(s)}(1), \dots, a_{m-1}^{(s)}(1)\right]^{T} = \boldsymbol{b}_{s},$$
(14)

202 with

$$\mathcal{M} := \mathcal{D}\left(\xi_1^{(m-1)s}, \dots, \xi_{m-1}^{(m-1)s}\right) \mathcal{V}(\xi_1, \dots, \xi_{m-1}).$$

Here  $\mathcal{D}(\boldsymbol{v}), \, \boldsymbol{v} = [v_1, \dots, v_{m-1}]^T$ , is the diagonal matrix with diagonal entries  $v_k, \, 1 \leq k \leq m-1, \, \mathcal{V}(\xi_1, \dots, \xi_{m-1})$  is the Vandermonde matrix with nodes  $\xi_k$ ,  $1 \leq k \leq m-1$ , and

$$(\boldsymbol{b}_{s})_{k} = -\sum_{i=0}^{m-1} \sum_{p=0}^{s-1} \frac{a_{i}^{(p)}(1)}{p!} \left( \sum_{j=max\{s-i,p\}}^{s} \binom{s}{j} A_{j,p}(\xi_{k}) \frac{i!}{(i-(s-j))!} \xi_{k}^{i-(s-j)} \right) - \frac{a_{m+1}^{(s)}(1)}{s!} A_{s,s}(\xi_{k}) \xi_{k}^{\frac{m+1}{2}}, \qquad 1 \le k \le m-1.$$

Since  $\xi_k$ ,  $1 \leq k \leq m-1$ , are distinct and non-zero, the coefficient matrix is nonsingular and  $a_i^{(s)}(1)$ ,  $0 \leq i \leq m-1$ ,  $i \neq (m+1)/2$ , are uniquely determined.

Once the quantities  $a_i^{(s)}(1), 0 \le i \le m-1, s = 0, \dots, d-1$ , are calculated, then the sub-symbols  $a_i(z), 0 \le i \le m-1, i \ne (m+1)/2$ , can be represented as follows

$$a_i(z) = 1 + \sum_{j=1}^{d-1} \frac{a_i^{(j)}(1)}{j!} (z-1)^j + (z-1)^d \hat{a}_i(z) = \check{a}_i(z) + (z-1)^d \hat{a}_i(z), \quad (15)$$

for suitable  $\hat{a}_i(z) \in \mathbb{R}[z, z^{-1}]$ . This representation is exploited in the second step to find a solution of (9). Combining (9) with (15) we obtain

$$1 - \phi(z)\phi_{\frac{m+1}{2}}(z) - \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} \check{a}_i(z)\phi_i(z) = (z-1)^d \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} \hat{a}_i(z)\phi_i(z).$$
(16)

<sup>212</sup> Thus, setting

$$\theta(z) := 1 - \phi(z)\phi_{\frac{m+1}{2}}(z) - \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1}\check{a}_i(z)\phi_i(z), \tag{17}$$

213 the condition

$$\theta^{(s)}(1) = 0, \qquad 0 \le s \le d-1,$$

<sup>214</sup> is needed, but it is always guaranteed by the following result.

Theorem 2 The function  $\theta(z)$  in (17) satisfies  $\theta^{(s)}(1) = 0$  for  $s = 0, \dots, d-1$ .

<sup>216</sup> Proof Let us introduce the truncated representation  $\check{a}(z)$  of the symbol a(z), that <sup>217</sup> is,

$$\check{a}(z) = \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} \check{a}_i(z^m) z^i + \phi(z^m) z^{\frac{m+1}{2}},$$

and consider the auxiliary function  $q(z) = z^{-\frac{m+1}{2}}\check{a}(z^2)z\phi(z^2)$ . From (11) it follows that  $q(z) = z^{-\frac{m+1}{2}}\check{a}(z^2) - z^{-\frac{m+1}{2}}\check{a}(z^2)(-1)^d(1-z)^d\gamma(z)$ . By construction  $\check{a}(z)$ satisfies relations (10). By using the representation of  $\check{a}(z)$  provided by (10) this gives

$$q(z) = z^{-\frac{m+1}{2}} \check{a}(z^2) + \frac{(1-z^m)^d (1+z^m)^d}{(1+z)^d} \hat{\rho}(z)$$

with  $\hat{\rho}(z) \in \mathbb{R}[z, z^{-1}]$ . Observe that

$$z^{-\frac{m+1}{2}}\check{a}(z^{2}) = z^{\frac{m+1}{2}}\phi(z^{2m}) + \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1}\check{a}_{i}(z^{2m})z^{2i-\frac{m+1}{2}},$$

223 and, hence,

$$q(z) = z^{\frac{m+1}{2}}\phi(z^{2m}) + \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1}\check{a}_i(z^{2m})z^{2i-\frac{m+1}{2}} + \frac{(1-z^m)^d(1+z^m)^d}{(1+z)^d}\hat{\rho}(z).$$
(18)

Moreover it can be easily seen that the two sets  $[0, m-1] \cap \mathbb{N}$  and  $\{n \in \mathbb{N} : n = 2i - (m+1)/2 \pmod{m}, 0 \leq i \leq m-1\}$  coincide. Besides this, by direct multiplication of  $a(z^2)$  and  $\phi(z^2)$ , we can write

$$q(z) = z^{\frac{1-m}{2}} \left( \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} \check{a}_i(z^{2m})\phi_i(z^{2m}) + \phi(z^{2m})\phi_{\frac{m+1}{2}}(z^{2m}) \right) + z^{\frac{1-m}{2}} \sum_{\substack{0 \le i,j \le m-1\\i\neq j}} z^{2(i-j)}\eta_{i,j}(z^{2m}),$$
(19)

for suitable Laurent polynomials  $\eta_{i,j}(z) \in \mathbb{R}[z, z^{-1}]$ . Since  $(1-m)/2 \equiv (m+1)/2$ (mod m) the class of integers congruent to (1-m)/2 modulo m is  $\{n \in \mathbb{Z} : n = (1-m)/2 + \ell m, \ell \in \mathbb{Z}\}$ . It follows that  $n = (1-m)/2 + 2(i-j), i \neq j, 0 \leq i, j \leq m-1$ , is such that  $n \not\equiv (1-m)/2$  (mod m). Hence, by comparison of classes mod m in (18) and (19), we obtain that

$$z^{m}\phi(z^{2m}) = \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} \check{a}_{i}(z^{2m})\phi_{i}(z^{2m}) + \phi(z^{2m})\phi_{\frac{m+1}{2}}(z^{2m}) + (1-z^{m})^{d}\tilde{\rho}(z),$$

for some  $\tilde{\rho}(z) \in \mathbb{R}[z, z^{-1}]$ . From (11) this implies that

$$\sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} \check{a}_i(z^2)\phi_i(z^2) + \phi(z^2)\phi_{\frac{m+1}{2}}(z^2) = 1 + (1-z)^d\rho(z), \qquad \rho(z) \in \mathbb{R}[z, z^{-1}],$$

 $_{233}$  which concludes the proof.

Theorem 2, along with Assumption 2, guarantees the existence of  $\hat{\theta}(z) \in \mathbb{R}[z, z^{-1}]$ such that

$$\theta(z) = (z-1)^d g(z) \hat{\theta}(z).$$
(20)

Thus, due to (16), the polynomial corrections  $\hat{a}_i(z)$ ,  $0 \le i \le m-1$ ,  $i \ne (m+1)/2$ , must satisfy the Bezout equation

$$\hat{\theta}(z) = \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1} \hat{a}_i(z) \frac{\phi_i(z)}{g(z)}.$$
(21)

Under Assumption 2 the Laurent polynomials  $\phi_i(z)/g(z)$ ,  $i \neq (m+1)/2$ , are relatively prime and thus equation (21) is solvable. In particular, following [19] every solution of (21) can be written as

$$\hat{a}_{i}(z) = \tilde{a}_{i}(z) + \sum_{\substack{j=i+1\\ j \neq \frac{m+1}{2}}}^{m-1} H_{i,j}(z) \frac{\phi_{j}(z)}{g(z)} - \sum_{\substack{j=0\\ j \neq \frac{m+1}{2}}}^{i-1} H_{j,i}(z) \frac{\phi_{j}(z)}{g(z)},$$
(22)

where  $\{\tilde{a}_i(z), i \neq (m+1)/2\}$  is a particular solution of (21) and  $H_{i,j}(z)$  is any element of  $\mathbb{R}[z, z^{-1}]$ . Upper bounds for the minimal length of the coefficient vectors associated to the solution of (21) are known a priori [19]. Using these bounds the computation of a particular solution  $\tilde{a}_i(z), 0 \leq i \leq m-1, i \neq \frac{m+1}{2}$ , reduces to solving a square linear system.

Remark 2 If  $g(z) = (z-1)^q \hat{g}(z)$  for some  $q \in \mathbb{N}, q < d$ , and  $\hat{g}(z) \in \mathbb{R}[z, z^{-1}]$ , with 246  $\hat{g}(1) \neq 0$ , then the result of Theorem 2 is unchanged but, differently from (20), we 247 can only factorize  $\theta(z)$  as 248

$$\theta(z) = (z-1)^{d-q} g(z) \hat{\theta}(z)$$

Thus, in this case, one should consider 249

$$a_i(z) = 1 + \sum_{j=1}^{d-q-1} \frac{a_i^{(j)}(1)}{j!} (z-1)^j + (z-1)^{d-q} \hat{a}_i(z) = \check{a}_i(z) + (z-1)^{d-q} \hat{a}_i(z), \quad (23)$$

instead of (15), and the illustrated procedure will lead to a symbol a(z) of the 250 form 251

$$a(z) = \left(\frac{1+z+\ldots+z^{m-1}}{m}\right)^{d-q} b(z)$$
 (24)

- rather than (10). This means that the scheme associated to a(z) would reproduce 252 only polynomials up to degree d - q - 1. 253
- Remark 3 Combining (6), (15) and (22), we get 254

$$\begin{aligned} a(z) &= \sum_{i=0}^{m-1} \check{a}_i(z^m) z^i + (z^m - 1)^d \sum_{i=0}^{m-1} \widetilde{a}_i(z^m) z^i + \\ &+ (z^m - 1)^d \sum_{i=0}^{m-1} \left( \sum_{\substack{j=i+1\\ j \neq \frac{m+1}{2}}}^{m-1} H_{i,j}(z^m) \frac{\phi_j(z^m)}{g(z^m)} - \sum_{\substack{j=0\\ j \neq \frac{m+1}{2}}}^{i-1} H_{j,i}(z^m) \frac{\phi_j(z^m)}{g(z^m)} \right) z^i, \end{aligned}$$

where on the right-hand-side the unique unknowns are the coefficients of the Lau-255 rent polynomials  $H_{i,j}(z)$ , i, j = 0, ..., m-1. Knowing the first and the last non-zero

256 coefficients of 257

$$\sum_{i=0}^{m-1} \check{a}_i(z^m) z^i + (z^m - 1)^d \sum_{i=0}^{m-1} \tilde{a}_i(z^m) z^i,$$
(25)

it is possible to establish the indices of the first and the last non-zero coefficients 258 of each  $H_{i,j}(z)$ , so that the range of the powers in (25) and in 259

$$(z^m - 1)^d \sum_{i=0}^{m-1} \left( \sum_{\substack{j=i+1\\j\neq\frac{m+1}{2}}}^{m-1} H_{i,j}(z^m) \frac{\phi_j(z^m)}{g(z^m)} - \sum_{\substack{j=0\\j\neq\frac{m+1}{2}}}^{i-1} H_{j,i}(z^m) \frac{\phi_j(z^m)}{g(z^m)} \right) z^i,$$

are the same. After that one can start imposing the first (or the last) coefficient 260 of a(z) to be 0, which is a linear condition with respect to the coefficients of all 261  $H_{i,j}(z)$ . It is possible then to add linear constraints in the same unknowns in 262 order to annihilate additional coefficients of a(z) as long as the new added linear 263 condition is compatible with the previous ones. Since (22) encodes all possible 264 solutions of (21), when there are no more compatible conditions to be added, the 265 mask with minimal support has been reached. A naive implementation of this 266 strategy has been used in our experiments to produce the interpolatory mask of 267 minimal support. A more general algorithmic description of this strategy should 268 incorporate some preprocessing algebraic computations such as the reduction of 269 the polynomials in reduced form as described in [1]. 270

271 Remark 4 Of great importance for applications is the case of symbols a(z) that 272 satisfy the symmetry condition  $a(z) = za(z^{-1})$ . The existence of such a sym-273 metric symbol can be proven under the auxiliary assumption that  $\varphi(1/2 + \ell) =$ 274  $\varphi(-1/2 - \ell), \ell \in \mathbb{N} \cup \{0\}$ . Under this assumption, we obtain that the coefficients 275 of a(z) satisfy (4) if and only if the coefficients of  $za(z^{-1})$  also satisfy (4). By 276 linearity this implies that the coefficients of  $(a(z) + za(z^{-1}))/2$  satisfy (4) too, with 277  $(a(z) + za(z^{-1}))/2$  fulfilling the symmetry condition.

The presented procedure for the odd arity case can be summarized as in Algorithm 1, at the end of which Remark 3 and Remark 4 can be exploited to reduce the support of the resulting mask and/or to obtain a symmetric mask. The following example is used to illustrate our composite approach for the odd arity case. Here we construct the dual ternary interpolating scheme, reproducing quintic polynomials, sharing with the primal binary Dubuc-Deslauriers 6-point scheme the same samples at the half integers, and having symmetric mask with shortest support.

Example 1 We choose m = 3, d = 6 and (see Remark 1)

$$\varphi\left(\frac{1}{2}+\ell\right) = \begin{cases} \frac{3}{256}, & \text{if } \ell \in \{-3,2\}, \\ -\frac{25}{256}, & \text{if } \ell \in \{-2,1\}, \\ \frac{75}{128}, & \text{if } \ell \in \{-1,0\}, \\ 0, & \text{otherwise.} \end{cases}$$
(26)

These values are taken from the mask of the primal binary 6-point interpolating scheme which reproduces quintic polynomials and it has a basic limit function supported in [-5, 5] with best Hölder exponent 2.830074998557687. Its primal ternary counterpart (see, e.g., [24]) reproduces quintic polynomials as well, but it has a basic limit function supported in [-4, 4] with best Hölder exponent 2.319856140753624. According to (7), we have

$$\begin{split} \phi(z) &= \frac{3}{256\,z^3} - \frac{25}{256\,z^2} + \frac{75}{128\,z} + \frac{75}{128} - \frac{25\,z}{256} + \frac{3\,z^2}{256} \\ &= \phi_0(z^3) + \phi_1(z^3)z^{-1} + \phi_2(z^3)z^{-2}, \end{split}$$

292 with

$$\phi_0(z) = \frac{3}{256 z} + \frac{75}{128}, \qquad \phi_1(z) = \frac{75}{128} + \frac{3 z}{256}, \qquad \phi_2(z) = -\frac{25}{256} - \frac{25 z}{256}.$$

<sup>293</sup> In particular, we observe that

$$1 - z\phi(z^2) = -(z-1)^6 \frac{3z^4 + 18z^3 + 38z^2 + 18z + 3}{256z^5}$$
(27)

294 and

$$g(z) = gcd\{\phi_0(z), \phi_1(z)\} = 1$$

Thus, Assumption 1 and Assumption 2 are satisfied. After solving the linear system (14), we have from (15)

$$a_0(z) = \check{a}_0(z) + (z-1)^6 \hat{a}_0(z), \qquad a_1(z) = \check{a}_1(z) + (z-1)^6 \hat{a}_1(z)$$

#### Algorithm 1 [odd arity case]

**Input:**  $m \in 2\mathbb{N} + 1$  and a compactly supported sequence  $\left\{\varphi\left(\frac{2k+1}{2}\right) \in \mathbb{R}\right\}_{k \in \mathbb{Z}}$  such that the Laurent polynomials

$$\begin{split} \phi_{\ell}(z) &= \sum_{i \in \mathbb{Z}} \varphi\left(\frac{2mi+1}{2} - \ell\right) z^{i}, \qquad \ell \in \{0, \dots, m-1\}, \\ g(z) &= gcd\left\{\phi_{0}(z), \dots, \phi_{\frac{m-1}{2}}(z), \phi_{\frac{m+3}{2}}(z), \dots, \phi_{m-1}(z)\right\}, \\ \phi(z) &= \sum_{i=0}^{m-1} \phi_{i}(z^{m}) z^{-i}, \end{split}$$

satisfy

- (a)  $1 z\phi(z^2) = (z-1)^d \gamma(z)$  for some  $d \in \mathbb{N}, \gamma \in \mathbb{R}[z, z^{-1}];$
- (b) g(z) divides  $1 \phi(z)\phi_{\frac{m+1}{2}}(z);$
- (c)  $g(z) = (z-1)^q \hat{g}(z)$  for some  $0 \leq q < d$ , and  $\hat{g}(z) \in \mathbb{R}[z, z^{-1}]$  with  $\hat{g}(1) \neq 0$ .

#### **Procedure:**

- (i) set  $a_{\frac{m+1}{2}}(z) = \phi(z);$
- (ii) for  $s \in \{1, \dots, d-q-1\}$ , solve linear system (14) for  $\{a_i^{(s)}(1)\}_{i \in \{0, \dots, m-1\} \setminus \{\frac{m+1}{2}\}};$
- (iii) for  $i \in \{0, \ldots, m-1\} \setminus \left\{\frac{m+1}{2}\right\}$ , define

$$\check{a}_i(z) \ = \ 1 \ + \ \sum_{s=1}^{d-q-1} \frac{a_i^{(s)}(1)}{s!} (z-1)^s;$$

(iv) compute

$$\hat{\theta}(z) = \frac{(z-1)^{q-d}}{g(z)} \left( 1 - \phi(z)\phi_{\frac{m+1}{2}}(z) - \sum_{\substack{i=0\\i\neq\frac{m+1}{2}}}^{m-1}\check{a}_i(z)\phi_i(z) \right);$$

(v) follow the strategy in [19] to compute Laurent polynomials  $\{\hat{a}_i(z)\}_{i\in\{0,...,m-1\}\setminus\left\{\frac{m+1}{2}\right\}}$  such that

$$\hat{\theta}(z) = \sum_{\substack{i=0\\i \neq \frac{m+1}{2}}}^{m-1} \hat{a}_i(z) \frac{\phi_i(z)}{g(z)};$$

Output: the symbol

$$a(z) \ = \ \sum_{i=0}^{m-1} a_i(z^m) z^i \quad \text{with} \quad a_i(z) \ = \ \check{a}_i(z) \ + \ \hat{a}_i(z)(z-1)^{d-q}, \ i \neq \frac{m+1}{2},$$

of an *m*-ary dual interpolating subdivision scheme reproducing polynomials up to degree d-q-1 and having basic limit function with the given samples at  $\mathbb{Z}/2$ .

297 with

304

$$\begin{split} \breve{a}_0(z) &= 1 + \frac{(z-1)}{6} - \frac{5(z-1)^2}{72} + \frac{55(z-1)^3}{1296} - \frac{935(z-1)^4}{31104} + \frac{4301(z-1)^5}{186624}, \\ \breve{a}_1(z) &= 1 - \frac{(z-1)}{6} + \frac{7(z-1)^2}{72} - \frac{91(z-1)^3}{1296} + \frac{1729(z-1)^4}{31104} - \frac{8645(z-1)^5}{186624}, \end{split}$$

299 and

$$a_2(z) = \phi(z).$$

300 To search for compatible  $\hat{a}_0(z)$  and  $\hat{a}_1(z)$ , we first compute

$$\widehat{\theta}(z) \;=\; \frac{8645\,z^3 + 215471\,z^2 - 24300\,z + 18225}{15925248\,z^3}$$

<sup>301</sup> in such a way that (20) holds, i.e.,

$$(z-1)^{6}\widehat{\theta}(z) = 1 - a_{2}(z) \phi_{2}(z) - \sum_{i=0}^{1} \check{a}_{i}(z) \phi_{i}(z).$$

Then we look for particular solutions  $\tilde{a}_0(z)$  and  $\tilde{a}_1(z)$  such that

$$\widehat{\theta}(z) = \widetilde{a}_0(z) \phi_0(z) + \widetilde{a}_1(z) \phi_1(z)$$

303 A possible choice is

$$\widetilde{a}_0(z) = -\frac{9903400 \, z - 45544275}{466373376 \, z^2}, \\ \widetilde{a}_1(z) = \frac{21603855 \, z - 46560721}{466373376 \, z^2}.$$

To obtain a shorter mask, according to Remark 3, we search for a suitable  $H_{0,1}(z)$ so that replacing

$$\hat{a}_{0}(z) = \tilde{a}_{0}(z) + H_{0,1}(z) \phi_{1}(z),$$
$$\hat{a}_{1}(z) = \tilde{a}_{1}(z) - H_{0,1}(z) \phi_{0}(z),$$

in the previous expressions of  $a_0(z)$  and  $a_1(z)$ , leads to a symbol

$$a(z) = a_0(z^3) + a_1(z^3) z + a_2(z^3) z^2$$

with a shorter associated mask. The choice of  $H_{0,1}(z)$  that leads to the shortest mask is

$$H_{0,1}(z) = -\frac{844799}{5465313 z^2}$$

and, after symmetrization (see Remark 4), the resulting symmetric mask a is such that  $a_i = 0$  for  $i \notin [-14, 15]$ , with the first half of its entries being

$$\left\{\frac{16567}{466373376}, 0, -\frac{414175}{233186688}, \frac{224821}{66624768}, \frac{3}{256}, \frac{589847}{33312384}, -\frac{83995}{2776032}, -\frac{25}{256}, -\frac{2042857}{22208256}, \frac{1290971}{8328096}, \frac{75}{128}, \frac{63152905}{66624768}\right\}.$$

$$(28)$$

The basic limit function  $f_{\delta}$  related to the mask in (28) is shown in Figure 1, and two examples of interpolating curves can be found in Figure 2. We have that  $\operatorname{supp}(f_{\delta}) = [-23/4, 23/4]$  and  $f_{\delta} \in C^{\omega}(\mathbb{R})$  with the best Hölder exponent  $\omega$ being 3.006664260760692. By construction the corresponding subdivision scheme reproduces polynomials of degree 5.



Fig. 1: The graph of the basic limit function  $f_{\pmb{\delta}}$  related to the mask in (28).



Fig. 2: Two examples of interpolating curves given by the subdivision scheme associated to the mask in (28). On the left, the first level of subdivision starting with the dotted control polygons; on the right, the corresponding interpolating limit curves.

### 318 3.2 The even arity case

Let us now consider the solution of the linear system (35) in [26] for the case where m is an even integer. The system is defined as follows:

$$M\boldsymbol{a} = \boldsymbol{c}, \qquad M = (\mu_{i,j})_{i,j\in\mathbb{Z}}, \qquad \boldsymbol{c} = (c_i)_{i\in\mathbb{Z}}$$
 (29)

321 where

$$\mu_{i,j} = \begin{cases} \varphi\left(\frac{i+1}{2} - j\right), & \text{if } i \in m\mathbb{Z}, \\ 0, & \text{otherwise,} \end{cases} \qquad c_i = \begin{cases} 1, & \text{if } i = 0, \\ \varphi\left(\frac{i}{2m}\right), & \text{if } i \in m(2\mathbb{Z}+1), \\ 0, & \text{otherwise.} \end{cases}$$

By suppressing the zero rows in both M and c we obtain the equivalent linear system

$$\widehat{M}\boldsymbol{a} = \widehat{\boldsymbol{c}}, \qquad \widehat{M} = (\widehat{\mu}_{i,j})_{i,j\in\mathbb{Z}}, \qquad \widehat{\boldsymbol{c}} = (\widehat{c}_i)_{i\in\mathbb{Z}}$$
(30)

324 where

$$\hat{\mu}_{i,j} = \varphi\left(\frac{im+1}{2} - j\right), \quad i,j \in \mathbb{Z}, \qquad \hat{c}_i = \begin{cases} 1, & \text{if } i = 0, \\ \varphi\left(\frac{i}{2}\right), & \text{if } \mod(i,2) = 1, \\ 0, & \text{otherwise.} \end{cases}$$

According to [26], (29) and (30) can be expressed in functional form as

$$\sum_{\alpha \in m\mathbb{Z}} \sum_{\beta \in \mathbb{Z}} a_{\beta} \varphi \left( \frac{\alpha + 1}{2} - \beta \right) z^{\alpha} = 1 + \sum_{\alpha \in m(2\mathbb{Z} + 1)} \varphi \left( \frac{\alpha}{2m} \right) z^{\alpha}$$

326 which can be rewritten as

$$\sum_{\ell \in \mathbb{Z}} \sum_{\beta \in \mathbb{Z}} a_{\beta} \varphi \left( \frac{m\ell + 1}{2} - \beta \right) z^{\ell} = 1 + \sum_{\ell \in \mathbb{Z}} \varphi \left( \frac{2\ell + 1}{2} \right) z^{2\ell + 1}$$

$$= 1 + \sum_{\ell \in \mathbb{Z}} \varphi \left( \ell + \frac{1}{2} \right) z^{2\ell + 1}.$$
(31)

327 Supposing that, as in the odd arity case, Assumption 1 holds for

$$\phi(z) = \sum_{\ell \in \mathbb{Z}} \varphi\left(\frac{1}{2} + \ell\right) z^{\ell},$$

the right-hand side of (31) satisfies

$$\begin{aligned} 1 + \sum_{\ell \in \mathbb{Z}} \varphi\left(\ell + \frac{1}{2}\right) z^{2\ell+1} &= 1 + z\phi(z^2) \\ &= (z+1)^d (-1)^d \gamma(-z) \\ &= (z+1)^d \tilde{\gamma}(z), \qquad \tilde{\gamma}(z) \in \mathbb{R}[z, z^{-1}]. \end{aligned}$$

Concerning the representation of the left-hand side of (31), let us introduce the modified subsymbols defined by

$$\hat{\phi}_{\ell}(z) = \sum_{i \in \mathbb{Z}} \varphi\left(\frac{mi+1}{2} - \ell\right) z^{i}, \qquad 0 \leq \ell \leq m-1.$$
(32)

Notice that if  $\phi_{\ell}(z)$ ,  $0 \leq \ell \leq m/2 - 1$ , denote the subsymbols of the mask of arity m/2, then we have

$$\hat{\phi}_{\ell}(z) = \phi_{\ell}(z), \qquad \hat{\phi}_{\ell+m/2}(z) = z\hat{\phi}_{\ell}(z), \qquad 0 \le \ell \le m/2 - 1.$$
(33)

333 In particular this implies that

$$\hat{\phi}_{\ell+m/2}(-1) = -\hat{\phi}_{\ell}(-1), \qquad \hat{\phi}_{\ell+m/2}(1) = \hat{\phi}_{\ell}(1), \qquad 0 \le \ell \le m/2 - 1.$$

Moreover, from  $1 + z\phi(z^2) = (z+1)^d \tilde{\gamma}(z)$  and  $1 - z\phi(z^2) = (z-1)^d \gamma(z)$  one deduces that

$$(z+1)^{d}\tilde{\gamma}(z) = 2 - (z-1)^{d}\gamma(z).$$
(34)

 $_{336}$  Then for the left-hand side of (31) it holds

$$\sum_{\ell \in \mathbb{Z}} \sum_{\beta \in \mathbb{Z}} a_{\beta} \varphi \left( \frac{m\ell+1}{2} - \beta \right) z^{\ell} = a_0(z^2) \widehat{\phi}_0(z) + \ldots + a_{m-1}(z^2) \widehat{\phi}_{m-1}(z) .$$

Hence, it follows that relation (31) can be reformulated as the Bezout-like poly nomial equation

$$a_0(z^2)\hat{\phi}_0(z) + \ldots + a_{m-1}(z^2)\hat{\phi}_{m-1}(z) = (z+1)^d\tilde{\gamma}(z).$$
(35)

From (33) it follows that equation (35) can be equivalently rewritten as

$$\left(a_0(z^2) + za_{m/2}(z^2)\right)\phi_0(z) + \ldots + \left(a_{m/2-1}(z^2) + za_{m-1}(z^2)\right)\phi_{m/2-1}(z) = (z+1)^d\tilde{\gamma}(z).$$
(36)

To proceed we consider the following assumption that plays the same role as Assumption 2 in the odd arity case.

342 ASSUMPTION 3 : We suppose that

$$g(z) := gcd \left\{ \phi_0(z), \dots, \phi_{m/2-1}(z) \right\},$$

<sup>343</sup> divides  $(z+1)^d \tilde{\gamma}(z)$  and is such that  $g(\pm 1) \neq 0$ .

Requiring g(z) to divide  $(z + 1)^d \tilde{\gamma}(z)$  and satisfy  $g(1) \neq 0$  is clearly a necessary condition because of (36) and Assumption 1. Condition  $g(-1) \neq 0$  however is only sufficient to construct a(z) as in (10) and, when it is not satisfied, a price has to be paid in terms of polynomial reproduction (see Remark 5).

Under Assumption 3 the solution to equation (35) can be found similarly to the odd arity case. Specifically, at the first step the unknowns  $a_i^{(s)}(1), 0 \le i \le m-1$ ,  $s = 1, \ldots, d-1$ , are computed by solving a Vandermonde linear system. The system is formed as follows. The first m-1 equations are obtained by differentiation of (10) complemented with relation (35). The last equation is found by imposing the

 $\leq m$ .

property (34) on the left hand-side of (35). If  $\xi_k = e^{2\pi i k/m}$ ,  $1 \leq k \leq m$ , denote the *m*-th roots of unity, then the system is of the form

$$m^{s} \mathcal{D}\left(\xi_{1}^{(m-1)s}, \dots, \xi_{m-1}^{(m-1)s}, (2/m)^{s}\right) \mathcal{V}(\xi_{1}, \dots, \xi_{m-1}, \phi) \left[a_{0}^{(s)}(1), \dots, a_{m-1}^{(s)}(1)\right]^{T} = \boldsymbol{b}_{s}, \quad (37)$$

where  $\mathcal{D}(\boldsymbol{v})$  is the diagonal matrix with diagonal entries  $v_k$  and  $\mathcal{V}(\xi_1, \ldots, \xi_{m-1}, \boldsymbol{\phi})$ is a Vandermonde-like matrix with nodes  $\xi_k$  of the form

$$\mathcal{V}(\xi_1, \dots, \xi_{m-1}, \phi) = \begin{bmatrix} \xi_1^0 & \dots & \xi_1^{m-1} \\ \vdots & \vdots & \vdots & \vdots \\ \xi_{m-1}^0 & \dots & \xi_{m-1}^{m-1} \\ (\phi)_1 & \dots & (\phi)_m \end{bmatrix} \text{ with } (\phi)_j = \hat{\phi}_{j-1}(1), \ 1 \le j$$

<sup>357</sup> The solvability of the systems (37) follows from the next lemma.

**Lemma 1** For any  $v = [v_1, \ldots, v_{m/2}] \in \mathbb{R}^{m/2}$  and  $w = [v, v] \in \mathbb{R}^m$ , it holds

$$\det \mathcal{V}(\xi_1,\ldots,\xi_{m-1},\boldsymbol{w}) = \frac{2}{m} \prod_{1 \leq i,j \leq m} (\xi_i - \xi_j) \sum_{i=1}^{m/2} v_i$$

Proof By Laplace's rule we find that det  $\mathcal{V}(\xi_1, \ldots, \xi_{m-1}, \boldsymbol{w})$  is linear in  $v_1, \ldots, v_{m/2}$ . If  $\sum_{i=1}^{m/2} v_i = 0$ , then det  $\mathcal{V}(\xi_1, \ldots, \xi_{m-1}, \boldsymbol{w}) = 0$  since the all-ones vector **1** belongs to the kernel of the matrix. This implies that det  $\mathcal{V}(\xi_1, \ldots, \xi_{m-1}, \boldsymbol{w}) = \gamma \sum_{i=1}^{m/2} v_i$ for a suitable  $\gamma$ . The value of  $\gamma$  can be determined by setting  $\boldsymbol{w} = \mathbf{1}$  which amounts to consider the customary Vandermonde matrix.  $\square$ 

As a consequence of Assumption 1 with z = 1, we have that

$$\sum_{i=1}^{m/2} \hat{\phi}_{i-1}(1) = \sum_{i=1}^{m/2} \phi_{i-1}(1) = 1.$$

Therefore, by Lemma 1, the coefficient matrix in (37) is non-singular and  $a_i^{(s)}(1)$ ,  $0 \leq i \leq m-1$ , are uniquely determined. Once these quantities are computed, the sub-symbols can be represented as follows

$$a_{i}(z) = \check{a}_{i}(z) + (z-1)^{d} \hat{a}_{i}(z), \qquad 0 \leq i \leq m-1,$$
  
$$\check{a}_{i}(z) = 1 + \sum_{j=1}^{d-1} \frac{a_{i}^{(j)}(1)}{j!} (z-1)^{j},$$
(38)

for suitable  $\hat{a}_i(z) \in \mathbb{R}[z, z^{-1}]$ . This representation is exploited in the second step to find a solution of (35). If we set

$$\theta(z) := (z+1)^{d} \tilde{\gamma}(z) - \sum_{j=0}^{m-1} \check{a}_{j}(z^{2}) \hat{\phi}_{j}(z), \qquad (39)$$

by using similar arguments as in the proof of Theorem 2, together with Assumption 3, it is shown that there exists  $\hat{\theta}(z) \in \mathbb{R}[z, z^{-1}]$  such that

$$\theta(z) = (z^2 - 1)^d g(z) \hat{\theta}(z).$$
(40)

<sup>372</sup> In this way equation (35) can be simplified as follows

$$\hat{a}_0(z^2)\frac{\hat{\phi}_0(z)}{g(z)} + \ldots + \hat{a}_{m-1}(z^2)\frac{\hat{\phi}_{m-1}(z)}{g(z)} = \hat{\theta}(z),$$

<sup>373</sup> which yields to its reduced analogue

$$\left(\hat{a}_0(z^2) + z\hat{a}_{m/2}(z^2)\right)\frac{\phi_0(z)}{g(z)} + \ldots + \left(\hat{a}_{m/2-1}(z^2) + z\hat{a}_{m-1}(z^2)\right)\frac{\phi_{m/2-1}(z)}{g(z)} = \hat{\theta}(z).$$

By setting  $\tilde{a}_i(z) = \hat{a}_i(z^2) + z\hat{a}_{i+m/2}(z^2), 0 \le i \le m/2 - 1$ , thanks to Assumption 3 we deduce that the equation

$$\widetilde{a}_0(z)\frac{\phi_0(z)}{g(z)} + \ldots + \widetilde{a}_{m/2-1}(z)\frac{\phi_{m/2-1}(z)}{g(z)} = \widehat{\theta}(z)$$
(41)

376 is solvable and every solution can be written as

$$\overline{a}_i(z) = \widetilde{a}_i(z) + \sum_{j=i+1}^{m/2-1} H_{i,j}(z) \frac{\phi_j(z)}{g(z)} - \sum_{j=0}^{i-1} H_{j,i}(z) \frac{\phi_j(z)}{g(z)}$$

where  $\tilde{a}_i(z)$  is a particular solution of (41) and  $H_{i,j}(z)$  is any element of  $\mathbb{R}[z, z^{-1}]$ .

Remark 5 If  $g(z) = (z+1)^q \hat{g}(z)$  for some  $q \in \mathbb{N}$ , q < d, and  $\hat{g}(z) \in \mathbb{R}[z, z^{-1}]$ , with  $\hat{g}(-1) \neq 0$ , then one can only factorize  $\theta(z)$  as

$$\theta(z) = (z^2 - 1)^{d-q} (z - 1)^q g(z) \hat{\theta}(z)$$

rather than (40). In this case, similarly to what was done in Remark 2, one should consider (23) instead of (38) and the illustrated procedure will lead to a symbol a(z) of the form (24) instead of (10). This means that the subdivision scheme associated to a(z) would reproduce only polynomials up to degree d - q - 1.

Remark 6 For m = 2 equation (35) becomes

$$(a_0(z^2) + za_1(z^2))\phi(z) = 1 + z\phi(z^2)$$

which implies that the first and the last non-zero elements of a(z) must be equal to 1. It follows that the associated subdivision scheme cannot be convergent [26].

The presented procedure for the even arity case can be summarized as in 387 Algorithm 2, after which similar arguments as in Remark 3 and Remark 4 can 388 be exploited, to reduce the support of the resulting mask and/or to obtain a 389 symmetric one. Next, we conclude with the illustration of our composite approach, 390 in the even arity case, by means of a computational example where we construct the 391 dual quaternary interpolating scheme, reproducing quintic polynomials, sharing 392 with the primal binary Dubuc-Deslauriers 6-point scheme the same samples at the 393 half integers, and having symmetric mask with shortest support. 394

#### Algorithm 2 [even arity case]

**Input:**  $m \in 2\mathbb{N} \setminus \{2\}$  and a compactly supported sequence  $\left\{\varphi\left(\frac{2k+1}{2}\right) \in \mathbb{R}\right\}_{k \in \mathbb{Z}}$  such that the Laurent polynomials

$$\begin{split} \phi_{\ell}(z) &= \sum_{i \in \mathbb{Z}} \varphi\left(\frac{mi+1}{2} - \ell\right) z^{i}, \qquad \ell \in \{0, \dots, m/2 - 1\}, \\ g(z) &= gcd\left\{\phi_{0}(z), \dots, \phi_{m/2 - 1}(z)\right\}, \\ \phi(z) &= \sum_{i=0}^{m/2 - 1} \phi_{i}(z^{m}) z^{-i}, \end{split}$$

satisfy

- (a)  $1 z\phi(z^2) = (z-1)^d \gamma(z)$  for some  $d \in \mathbb{N}, \gamma \in \mathbb{R}[z, z^{-1}];$
- (b) g(z) divides  $1 + z\phi(z^2)$ ;

(c) 
$$g(z) = (z+1)^q \hat{g}(z)$$
 for some  $0 \leq q < d$ , and  $\hat{g}(z) \in \mathbb{R}[z, z^{-1}]$  with  $\hat{g}(-1) \neq 0$ .

#### **Procedure:**

- (i) for  $s \in \{1, ..., d q 1\}$ , solve linear system (37) for  $\{a_i^{(s)}(1)\}_{i=0,...,m-1}$ ; (ii) for  $i \in \{0, ..., m 1\}$ , define

$$\check{a}_i(z) = 1 + \sum_{s=1}^{d-q-1} \frac{a_i^{(s)}(1)}{s!} (z-1)^s;$$

(iii) compute

$$\hat{\theta}(z) = \frac{(z^2 - 1)^{q-d}}{(z - 1)^q g(z)} \left( 1 + z\phi(z^2) - \sum_{i=0}^{m/2-1} \left(\check{a}_i(z^2) + z\check{a}_{m/2+i}(z^2)\right)\phi_i(z) \right);$$

(iv) follow the strategy in [19] to compute Laurent polynomials  $\{\tilde{a}_i(z)\}_{i=0,\dots,m/2-1}$  such that

$$\hat{\theta}(z) = \sum_{i=0}^{m/2-1} \tilde{a}_i(z) \frac{\phi_i(z)}{g(z)};$$

(v) for  $i \in \{0, ..., m/2 - 1\}$ , compute the Laurent polynomials  $\hat{a}_i(z)$  and  $\hat{a}_{m/2+i}(z)$  uniquely defined by the relation

$$\widetilde{a}_i(z) = \widehat{a}_i(z^2) + z \widehat{a}_{m/2+i}(z^2).$$

Output: the symbol

$$a(z) = \sum_{i=0}^{m-1} a_i(z^m) z^i$$
 with  $a_i(z) = \check{a}_i(z) + \hat{a}_i(z)(z-1)^{d-q}$ ,

of an m-ary dual interpolating subdivision scheme reproducing polynomials up to degree d-q-1 and having basic limit function with the given samples at  $\mathbb{Z}/2$ .

#### *Example 2* We choose m = 4, d = 6 and (see Remark 1) 395

$$\varphi\left(\frac{1}{2}+\ell\right) = \begin{cases} \frac{3}{256}, & \text{if } \ell \in \{-3, 2\}, \\ -\frac{25}{256}, & \text{if } \ell \in \{-2, 1\}, \\ \frac{75}{128}, & \text{if } \ell \in \{-1, 0\}, \\ 0, & \text{otherwise.} \end{cases}$$
(42)

These values are again taken from the mask of the primal binary 6-point inter-396 polating scheme which reproduces quintic polynomials and it has a basic limit 397 function supported in [-5,5] with best Hölder exponent 2.830074998557687. Its 398 primal quaternary counterpart (see, e.g., [24]) reproduces quintic polynomials as 399 well, but it has a basic limit function supported in [-11/3, 11/3] with best Hölder 400 exponent 2.099550050039848. In view of (32) and (33), we have 401

$$\hat{\phi}_0(z) = \phi_0(z) = -\frac{25}{256 z} + \frac{75}{128} + \frac{3 z}{256}$$

402

$$\hat{\phi}_1(z) = \phi_1(z) = \frac{3}{256 z} + \frac{75}{128} - \frac{25 z}{256},$$
$$\hat{\phi}_2(z) = z\phi_0(z) = -\frac{25}{256} + \frac{75 z}{128} + \frac{3 z^2}{256},$$

403 404

$$\hat{\phi}_3(z) = z\phi_1(z) = \frac{3}{256} + \frac{75z}{128} - \frac{25z^2}{256}.$$

Assumption 1 is satisfied since  $\phi(z)$  is the same as in Example 1 (27), while As-405

sumption 3 holds because 406

$$\phi_1(z) = \phi_0(1/z) \implies gdc\{\phi_0(z), \phi_1(z)\} = 1.$$

After solving the linear system (37), from (38) we obtain 407

$$a_i(z) = \check{a}_i(z) + (z-1)^d \hat{a}_i(z), \qquad 0 \le i \le 3,$$

with 408

$$\breve{a}_0(z) = 1 + \frac{(z-1)}{8} - \frac{7(z-1)^2}{128} + \frac{35(z-1)^3}{1024} - \frac{805(z-1)^4}{32768} + \frac{4991(z-1)^5}{262144},$$

$$\breve{a}_1(z) = 1 - \frac{(z-1)}{8} + \frac{9(z-1)^2}{128} - \frac{51(z-1)^3}{1024} + \frac{1275(z-1)^4}{32768} - \frac{8415(z-1)^5}{262144},$$

$$\check{a}_1(z) = 1 - \frac{(z-1)}{z} + \frac{9(z)}{z}$$

$$\check{a}_{1}(z) = 1 - \frac{(z-1)}{8} + \frac{9(z-1)^2}{128} - \frac{51(z-1)^3}{1024} + \frac{1275(z-1)^4}{32768} - \frac{8415(z-1)^4}{262144} + \frac{1275(z-1)^4}{32768} - \frac{1275(z-1)^4}{262144} + \frac{1275(z-1)^4}{32768} - \frac{1275(z-1)^4}{3276$$

$$\check{a}_{2}(z) = 1 - \frac{3(z-1)}{8} + \frac{33(z-1)^{2}}{128} - \frac{209(z-1)^{3}}{1024} + \frac{5643(z-1)^{4}}{32768} - \frac{39501(z-1)^{5}}{262144}$$

...

$$\check{a}_3(z) = 1 - \frac{5(z-1)}{8} + \frac{65(z-1)^2}{128} - \frac{455(z-1)^3}{1024} + \frac{13195(z-1)^4}{32768} - \frac{97643(z-1)^5}{262144} + \frac{13195(z-1)^4}{262144} - \frac{1100}{262144} + \frac{1100}{26214} + \frac{1100}{262144} + \frac{1100}{262$$

To search for compatible  $\hat{a}_0(z)$ ,  $\hat{a}_1(z)$ ,  $\hat{a}_2(z)$  and  $\hat{a}_3(z)$ , we first compute 412

$$\widehat{\theta}(z) \ = \ \frac{3}{256\,z^5} \ - \ \frac{7}{256\,z^3} \ + \ \frac{5086563}{16777216\,z} \ - \ \frac{580643}{16777216}$$

 $_{413}$  such that, according to (39) and (40),

$$(z^{2}-1)^{6}\widehat{\theta}(z) = (z+1)^{6} \widetilde{\gamma}(z) - \sum_{i=0}^{3} \widecheck{a}_{i}(z^{2}) \widehat{\phi}_{i}(z),$$

414 with

$$\widetilde{\gamma}(z) = \frac{3}{256 z^5} - \frac{9}{128 z^4} + \frac{19}{128 z^3} - \frac{9}{128 z^2} + \frac{3}{256 z^5}$$

due to (34). Then we search for  $\tilde{a}_0(z)$  and  $\tilde{a}_1(z)$  that solve the reduced Bezout equation in (41), i.e.,

$$\hat{\theta}(z) = \tilde{a}_0(z) \phi_0(z) + \tilde{a}_1(z) \phi_1(z).$$
(43)

417 A possible choice is

$$\begin{split} \widetilde{a}_0(z) &= \frac{2126507351527}{157810688\,z} - \frac{176620228675}{78905344}, \\ \widetilde{a}_1(z) &= \frac{1}{z^4} - \frac{50}{z^3} + \frac{2506}{z^2} - \frac{2118539063675}{157810688\,z} - \frac{21194427441}{78905344}. \end{split}$$

<sup>419</sup> Once we have a solution of (43), we search for

$$\overline{a}_0(z) = \widetilde{a}_0(z) + H_{0,1}(z) \phi_1(z),$$
  
$$\overline{a}_1(z) = \widetilde{a}_1(z) - H_{0,1}(z) \phi_0(z),$$

421 so that  $\{\hat{a}_k(z)\}_{k=0,\dots,3}$  fulfilling

$$\overline{a}_i(z) = \hat{a}_i(z^2) + z \, \hat{a}_{i+2}(z^2), \qquad i \in \{0, 1\},$$

<sup>422</sup> lead to a symbol a(z) satisfying  $a(z) = za(z^{-1})$ . For example, the choice

$$H_{0,1}(z) = -\frac{7064809147}{308224 z} + \frac{281633113}{616448 z^2} - \frac{2817667}{308224 z^3} + \frac{119853}{616448 z^4} + \frac{7302199}{596413440 z^5} - \frac{3127}{1232896 z^6} + \frac{947}{1331280 z^7}$$

423 leads to

$$\overline{a}_{0}(z) = \frac{39501}{262144 z} - \frac{4991}{262144 z^{2}} - \frac{5643}{262144 z^{3}} + \frac{24415849}{4362338304 z^{4}} + \frac{394938757}{40715157504 z^{5}} \\ - \frac{61600783}{43623383040 z^{6}} + \frac{15760091}{40715157504 z^{7}} + \frac{947}{113602560 z^{8}} \\ \overline{a}_{1}(z) = \frac{97643}{262144 z} + \frac{8415}{262144 z^{2}} - \frac{7917}{262144 z^{3}} - \frac{49446367}{7270563840 z^{4}} + \frac{482174039}{40715157504 z^{5}} \\ + \frac{116624327}{43623383040 z^{6}} - \frac{27054815}{40715157504 z^{7}} + \frac{4735}{68161536 z^{8}},$$

425 and so

424

$$\hat{a}_{0}(z) = -\frac{4991}{262144 z} + \frac{24415849}{4362338304 z^{2}} - \frac{61600783}{43623383040 z^{3}} + \frac{947}{113602560 z^{4}},$$

$$\hat{a}_{1}(z) = \frac{8415}{262144 z} - \frac{49446367}{7270563840 z^{2}} + \frac{116624327}{43623383040 z^{3}} + \frac{4735}{68161536 z^{4}},$$

427

428

â (n)	39501	5643	394938757	15760091
$a_2(z) =$	$\overline{262144z}$	$\overline{262144  z^2}$	$+$ $\frac{1}{40715157504 z^3}$	$+ \frac{1}{40715157504 z^4},$
$\hat{z}$ (v)	97643	7917	482174039	27054815
$a_{3}(z) =$	$\overline{262144z}$	$\overline{262144z^2}$	$+$ $\overline{40715157504 z^3}$	$\overline{40715157504 z^4}$ .

Replacing the previous expressions in the above equations of  $a_0(z)$ ,  $a_1(z)$ ,  $a_2(z)$ and  $a_3(z)$  and using

$$a(z) = a_0(z^4) + a_1(z^4) z + a_2(z^4) z^2 + a_3(z^4) z^3,$$

431 the first half of the resulting symmetric mask a is

5	947	4735	15760091	27054815	63782671				
J	113602560	$\overline{68161536}$ ,	40715157504'	$\overline{40715157504}$	$\overline{43623383040}$				
	$\frac{98441927}{43623383040}$	$, \frac{42911173}{581645107}$	$\frac{3}{72}, \frac{92071847}{5816451072}$	$, \frac{154804477}{10905845760},$	$-\frac{79247347}{3635281920},$	(44)			
_	$\frac{143318065}{1938817024}$	$, -\frac{2156430}{1938817}$	$\frac{11}{024}, -\frac{7170639}{9694085}$	$\frac{19}{12}, \frac{4869166267}{43623383040}$	$\frac{2428957997}{5816451072},$	(44)			
$\left. \frac{4331006815}{5816451072}, \ \frac{528433771}{545292288} \right\}.$									

<sup>432</sup> The basic limit function  $f_{\delta}$  related to this mask is shown in Figure 3, and two <sup>433</sup> examples of interpolating curves can be found in Figure 4. We have that  $\operatorname{supp}(f_{\delta}) =$ <sup>434</sup> [-11/2, 11/2] and, via joint spectral radius techniques, one can prove that  $f_{\delta} \in$ <sup>435</sup>  $\mathcal{C}^{\omega}(\mathbb{R})$  with the best Hölder exponent  $\omega$  being 3.050871089158321. By construction

<sup>436</sup> the corresponding subdivision scheme reproduces polynomials of degree 5.

437 Acknowledgements The authors would like to thank the anonymous referees for their con438 structive comments and suggested improvements. All the authors are members of INdAM 439 GNCS, which partially supported this work. The first author is also partially supported by
440 the project PRA\_2020\_61 of the University of Pisa. The research of the last two authors has
441 been done within the Italian Network on Approximation (RITA) and the thematic group on

442 "Approximation Theory and Applications" of the Italian Mathematical Union.

443

444 Competing interests The authors have no competing interests to declare that are relevant 445 to the content of this article.

#### 446 References

- 447 1. Michael Alekhnovich. Linear Diophantine equations over polynomials and soft decoding
   448 of Reed-Solomon codes. *IEEE Trans. Inform. Theory*, 51(7):2257–2265, 2005.
- Alfred S. Cavaretta, Wolfgang Dahmen, and Charles A. Micchelli. Stationary subdivision.
   Mem. Amer. Math. Soc., 93(453), 1991.
- 451 3. Maria Charina, Costanza Conti, and Tomas Sauer. Regularity of multivariate vector 452 subdivision schemes. *Numer. Algorithms*, 39(1-3):97–113, 2005.
- 4. Maria Charina and Thomas Mejstrik. Multiple multivariate subdivision schemes: matrix and operator approaches. J. Comput. Appl. Math., 349:279–291, 2019.



Fig. 3: The graph of the basic limit function  $f_{\delta}$  related to the mask in (44).



Fig. 4: Two examples of interpolating curves given by the subdivision scheme associated to the mask in (44). On the left, the first level of subdivision starting with the dotted control polygons; on the right, the corresponding interpolating limit curves.

- Sung Woo Choi, Byung-Gook Lee, Yeon Ju Lee, and Jungho Yoon. Stationary subdivision
   schemes reproducing polynomials. *Comput. Aided Geom. Design*, 23(4):351 360, 2006.
- Charles Chui and Johan M. de Villiers. Wavelet Subdivision Methods: GEMS for Rendering Curves and Surfaces. CRC Press Boca Raton, FL, 2010.
- 459 7. Costanza Conti and Kai Hormann. Polynomial reproduction for univariate subdivision
   460 schemes of any arity. J. Approx. Theory, 163(4):413-437, 2011.
- 461 8. Costanza Conti and Lucia Romani. Dual univariate m-ary subdivision schemes of de
   462 Rham-type. J. Math. Anal. Appl., 407(2):443-456, 2013.
- 463 9. Johan de Villiers and Mpfareleni Rejoyce Gavhi. Local interpolation with optimal poly 464 nomial exactness in refinement spaces. *Math. Comp.*, 85(298):759–782, 2016.
- Io. Johan M. de Villiers, Charles A. Micchelli, and Tomas Sauer. Building refinable functions
   from their values at integers. *Calcolo*, 37(3):139–158, 2000.
- 467 11. Gilles Deslauriers and Serge Dubuc. Symmetric iterative interpolation processes. Constr.
   468 Approx., 5(1):49-68, 1989.
- 469 12. Serge Dubuc. Interpolation through an iterative scheme. J. Math. Anal. Appl., 114(1):185
   470 204, 1986.
- 471 13. Serge Dubuc. de Rham transforms for subdivision schemes. J. Approx. Theory, 163(8):966
   472 987, 2011.
- 473 14. Nira Dyn. Subdivision schemes in computer-aided geometric design. In Advances in Numerical Analysis, Vol. II (Lancaster, 1990), Oxford Sci. Publ., pages 36–104. Oxford Univ. Press, New York, 1992.
- <sup>476</sup> 15. Nira Dyn. Interpolatory subdivision schemes. In Floater M.S. Iske A., Quak E., edi<sup>477</sup> tor, *Tutorials on Multiresolution in Geometric Modelling*, pages 25–50. Springer, Berlin,
  <sup>478</sup> Heidelberg, 2002.
- 479 16. Nira Dyn, Kai Hormann, Malcolm A. Sabin, and Zuowei Shen. Polynomial reproduction
  480 by symmetric subdivision schemes. J. Approx. Theory, 155(1):28–42, 2008.
- 17. Nira Dyn and David Levin. Subdivision schemes in geometric modelling. Acta Numer.,
   11:73-144, 2002.
- 18. Mpfareleni Rejoyce Gavhi-Molefe and Johan de Villiers. On interpolatory subdivision
  symbol formulation and parameter convergence intervals. J. Comput. Appl. Math., 349:354
   365, 2019.
- 486 19. Graziano Gentili and Daniele C. Struppa. Minimal degree solutions of polynomial equa 487 tions. *Kybernetika*, 23(1):44–53, 1987.
- 20. Nicola Guglielmi and Vladimir Protasov. Exact computation of joint spectral character istics of linear operators. Found. Comput. Math., 13(1):37–97, 2013.
- 490 21. Warren P. Johnson. The curious history of Faà di Bruno's formula. Amer. Math. Monthly,
   491 109(3):217-234, 2002.
- 492 22. Thomas Mejstrik. Algorithm 1011: improved invariant polytope algorithm and applica 493 tions. ACM Trans. Math. Software, 46(3):Art. 29, 26, 2020.
- 494 23. Charles A. Micchelli. Interpolatory subdivision schemes and wavelets. J. Approx. Theory, 86:41-71, 1996.
- 496 24. Georg Muntingh. Symbols and exact regularity of symmetric pseudo-splines of any arity.
   497 BIT, 57(3):867–900, 2017.
- 498 25. Lucia Romani. Interpolating m-refinable functions with compact support: The second
   499 generation class. Appl. Math. Comput., 361:735-746, 2019.
- Lucia Romani and Alberto Viscardi. Dual univariate interpolatory subdivision of every arity: Algebraic characterization and construction. J. Math. Anal. Appl., 484(1):123713, 2020.