

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

The positive impact of conservation action

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Langhammer, P.F., Bull, J.W., Bicknell, J.E., Oakley, J.L., Brown, M.H., Bruford, M.W., et al. (2024). The positive impact of conservation action. SCIENCE, 384(6694), 453-458 [10.1126/science.adj6598].

Availability:

This version is available at: <https://hdl.handle.net/11585/996574> since: 2024-11-15

Published:

DOI: <http://doi.org/10.1126/science.adj6598>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Title: The Positive Impact of Conservation Action

Authors: Penny F. Langhammer^{1,2*}, Joseph W. Bull³⁻⁵, Jake E. Bicknell⁴, Joseph L. Oakley⁶, Mary H. Brown¹, Michael W. Bruford^{7,8†}, Stuart H. M. Butchart^{9,10}, Jamie A. Carr¹¹⁻¹³, Don Church¹, Rosie Cooney^{14,15}, Simone Cutajar¹⁶, Wendy Foden^{13,17-19}, Matthew N. Foster¹, Claude Gascon²⁰, Jonas Geldmann^{10,21}, Piero Genovesi²²⁻²⁴, Michael Hoffmann^{25,26}, Jo Howard-McCombe^{27,28}, Tiffany Lewis²⁹, Nicholas B.W. Macfarlane³⁰, Zoe E. Melvin^{27,31}, Rossana Stoltz Merizalde³², Meredith G. Morehouse³³, Shyama Pagad³⁴, Beth Polidoro^{25,35}, Wes Sechrest¹, Gernot Segelbacher^{8,36}, Kevin G. Smith³⁷, Janna Steadman⁴, Kyle Strongin³⁸, Jake Williams³⁹, Stephen Woodley⁴⁰, Thomas M. Brooks⁴¹⁻⁴³

Affiliations:

¹Re:wild; PO Box 129, Austin, TX 78767, USA.

²Arizona State University, School of Life Sciences; PO Box 874501, Tempe, AZ 85287, USA.

³Department of Biology, University of Oxford; 11a Mansfield Rd, Oxford OX1 3SZ.

⁴Durrell Institute of Conservation and Ecology (DICE), School of Anthropology and Conservation, University of Kent; Canterbury, CT2 7NR, UK.

⁵Wild Business Ltd; London, UK.

⁶44 Hillary Road, Maidstone ME14 2JT, UK.

⁷School of Biosciences and Sustainable Places Research Institute; Cathays Park, Cardiff CF10 3AX, UK.

⁸IUCN SSC Conservation Genetics Specialist Group; 28 rue Mauverney, Gland 1196, Switzerland.

⁹BirdLife International; David Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, UK.

¹⁰Department of Zoology, University of Cambridge; Downing St., Cambridge CB2 3EJ, UK.

¹¹Leverhulme Centre for Anthropocene Biodiversity, University of York, York YO10 15DD, UK.

¹²Department of Environment and Geography, University of York; York YO10 5DD, UK.

¹³IUCN SSC Climate Change Specialist Group; 28 rue Mauverney, Gland 1196, Switzerland.

¹⁴CEESP/SSC IUCN Sustainable Use and Livelihoods Specialist Group; 28 rue Mauverney, Gland 1196, Switzerland.

¹⁵Fenner School of Environment and Society, Australian National University; 2601, ACT, Australia.

¹⁶University of Malta; Msida MSD 2080, Malta.

¹⁷South African National Parks, Cape Research Centre, Tokai; Cape Town, 7966, South Africa.

¹⁸FitzPatrick Institute of African Ornithology, Rondebosch; Cape Town, 7701, South Africa.

¹⁹Global Change Biology Group, Department of Botany and Zoology, Stellenbosch University; Stellenbosch, South Africa.

²⁰The Global Environment Facility; 1818 H Street NW, Washington, DC 20433, USA.

²¹Center for Macroecology, Evolution and Climate, GLOBE Institute, University of Copenhagen; Universitetsparken 15, 2100 Copenhagen E, Denmark.

²²Institute for Environmental Protection and Research; Via Vitaliano Brancati 48, 00144 Rome, Italy.

²³IUCN SSC Invasive Species Specialist Group; 00144 Rome, Italy.

²⁴Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University; Stellenbosch, South Africa.

²⁵IUCN Species Survival Commission; 28 rue Mauverney, Gland 1196, Switzerland.

²⁶Zoological Society of London; Regent's Park, London NW1 4RY, UK.

²⁷School of Biosciences, Cardiff University; The Sir Martin Evans Building, Museum Avenue, Cardiff, CF10 3AX, UK

²⁸RZSS WildGenes, Conservation Department, Royal Zoological Society of Scotland; Edinburgh EH12 6TS, UK.

²⁹Arizona State University; 427 E. Tyler Mall, Tempe, AZ 85281, USA.

³⁰IUCN; 1630 Connecticut Ave. NW, Washington DC 20009, USA.

³¹Bangor University, School of Natural Sciences; Deiniol Road, Bangor, Gwynedd, LL57 2UW, Wales, UK.

³²Conservation International; 2011 Crystal Drive #600, Arlington, VA 22202, USA.

³³LLaves: Keys to Bilingual Conservation, LLC; 346 Mayberry Hill Road, Casco, Maine 04015, USA.

³⁴University of Auckland; Auckland, 1072 New Zealand.

³⁵Arizona State University; 4701 W. Thunderbird Rd, Glendale, AZ 85382, USA.

³⁶University Freiburg; Tennenbacher Str. 4, 79106 Freiburg, Germany.

³⁷IUCN; The David Attenborough Building, Pembroke St, Cambridge, CB2 3QZ, UK.

³⁸Arizona State University; 800 S. Cady Mall, Tempe, AZ 85281, USA.

³⁹Imperial College London; Silwood Park, Ascot SL5 7PY, UK.

⁴⁰IUCN World Commission on Protected Areas; 64 Juniper Road, Chelsea, Quebec J9B 1T3, Canada.

⁴¹IUCN; 28 rue Mauverney, Gland 1196, Switzerland.

⁴²World Agroforestry Center, University of The Philippines Los Baños; Laguna, Philippines.

⁴³Institute for Marine & Antarctic Studies, University of Tasmania; Hobart, Australia.

*Corresponding author. Email: planghammer@rewild.org

†Deceased

‡Present address: ACT Government Office of Nature Conservation; 480 Northbourne Ave, 2602, ACT, Australia.

One-Sentence Summary: Meta-analysis reveals significant positive impacts of conservation actions on biodiversity, compared with outcomes anticipated without intervention

Abstract: Governments recently adopted new global targets to halt and reverse the loss of biodiversity. It is therefore crucial to understand the outcomes of conservation actions. We conducted a global meta-analysis of 186 studies (including 665 trials) that measured biodiversity over time and compared outcomes under conservation action with a suitable counterfactual of no action. We find that in two-thirds of cases, conservation either improved the state of biodiversity, or at least slowed declines. Specifically, we find that interventions targeted at species and ecosystems, such as invasive species control, habitat loss reduction and restoration, protected areas, and sustainable management, are highly effective, with large effect sizes. This provides the strongest evidence to date that conservation actions are successful, but require transformational scaling-up to meet global targets.

Main Text: Ongoing and dramatic declines in global biodiversity, and the associated negative consequences for human well-being, are among the most pressing contemporary risks to society (1). Governments have thus adopted goals to tackle biodiversity loss and its drivers: 20 Aichi Biodiversity Targets in the Strategic Plan for Biodiversity 2010-2020 through the Convention on Biological Diversity, and now 4 goals for 2050 and 23 targets for 2030 in the 2022 Kunming-Montreal Global Biodiversity Framework (GBF) (2). Similar targets are echoed in the 17 Sustainable Development Goals (3). More than \$121 billion is invested annually into biodiversity conservation worldwide (4). Yet despite this, none of the Aichi Targets were fully met (1). It could be concluded that responses to the ongoing biodiversity crisis are either insufficient, ineffective, or both (5, 6) and that the targets established in the new GBF will also likely not be achieved. However, such conclusions are premature: conservation interventions could represent progress, and be at least partly effective, even if global policy targets have not yet been achieved in full. There is consequently a need for a robust evaluation of policy targets, and an assessment of whether conservation interventions are working – that is, having positive impacts and providing better outcomes than the absence of interventions – as governments start committing resources and implementing the GBF.

Robust impact assessment using a counterfactual approach (7) reveals that conservation action has prevented extinctions (8) and reduced extinction risk (9) for species across taxonomic groups compared with an absence of conservation action. There has been an increase over the last decade in studies evaluating the impact of specific conservation actions, from global to local scales, using counterfactual comparisons, including for protected areas (10), payments for environmental services (11), invasive alien species (IAS) eradications (12), and sustainable management of ecosystems (13). Other studies have undertaken meta-analyses or systematic reviews, but only for individual conservation actions (14–16). Similarly, the Conservation Evidence website (17) provides a compendium of evidence for the effectiveness of a wide range of individual conservation interventions. However, since conservation actions started over a century ago, there has been no comprehensive meta-analysis of the impact of conservation across the full suite of conservation actions and intervention types, multiple levels and metrics of biodiversity, and over time. Such an assessment is critically needed to inform implementation of the GBF.

Here we conduct a meta-analysis of the impact of a wide range of conservation interventions globally. Meta-analysis provides a powerful and informative method to summarize results from multiple studies and accounts for unequal precision among studies in the calculation of effect sizes. We evaluate the impact of conservation actions that address direct pressures on biodiversity, promote restoration and recovery of populations and habitat, and aim to safeguard the environment across different levels of biological organization, compared with outcomes expected without intervention. Specifically, we consider seven intervention types aiming to tackle the direct drivers of environmental degradation: establishment and management of protected areas; other measures to reduce habitat loss and degradation (including policy and restoration); sustainable use of species; sustainable management of ecosystems; control of pollution; eradication and control of invasive alien (and problematic native) species; and climate change adaptation. These classes of intervention were identified based on the strategies of the intergovernmental environmental agreements, especially the nine Aichi Targets in Strategic Goals B and C of the Strategic Plan for Biodiversity 2011-2020 (Table S1), aligned in turn to the targets of the GBF and Sustainable Development Goals 14 (Life Below Water) & 15 (Life on Land). We consider impacts on biodiversity at ecosystem, species, and genetic levels.

We conducted a Rapid Evidence Assessment (see Methods) and meta-analysis of studies published in English that present a counterfactual-based analysis of the impact of conservation actions over time (18). Our literature search yielded 1,445 studies (published papers) spanning spatial scales from local to continental, and more than a century (1890-2019) (Table S2; Fig. S1). Of these, we retained only studies that contained temporal data, where we could express outcomes as effect sizes generated from the ‘rate of change under the intervention’ compared with ‘rate of change under a counterfactual scenario’ (19), yielding 186 studies that were included in our meta-analysis (Table S3). Using a rate of change to calculate the effect size allowed us to assess conservation actions over different time scales, avoiding premature conclusions based on study duration. Where studies measured change in biodiversity using more than one metric (e.g. different species), each was treated as a distinct trial nested within that study, totaling 665 trials across the dataset.

Our comprehensive dataset shows the variable outcomes of conservation outcomes. As such, interventions that generate gains in the state of biodiversity compared with a counterfactual in which biodiversity declines, stays the same, or improves to a lesser degree than the intervention, reveal absolute positive impacts of conservation action (Fig. 1a). Relative positive impacts of conservation action result when biodiversity declines, but the intervention slows the decline compared with the counterfactual (Fig. 1b). Conversely, relative negative impacts of conservation action occur when biodiversity improves, but the counterfactual reveals greater improvements than the intervention (Fig. 1c). Absolute negative impacts of conservation action result when biodiversity declines following the intervention while it improves, stays the same, or declines to a lesser degree in the counterfactual (Fig. 1d). These four categories are mutually exclusive.

Results

We find that in the majority of cases, biodiversity conservation works. Our meta-analysis shows that the ‘Overall’ impact of conservation is positive and significant (mean Hedges’ g [\pm 95% CI] = 3.24 [2.95 – 3.52], $p < 0.001$), indicating that, conservation interventions yield beneficial outcomes for biodiversity compared with the outcome in the absence of an intervention (Fig. 2a). We also show that conservation actions can yield positive impacts in both an absolute and a relative sense (Fig. 1). In two-thirds of trials, conservation either improved the state of biodiversity (“absolute positive impacts”, 45.4%), or at least slowed declines (“relative positive impacts”, 20.6%). However, in one-fifth of trials, biodiversity under the intervention declined more than no action (“absolute negative impacts”, 20.6%), while in a smaller number of cases biodiversity improved in both the intervention and counterfactual, but the counterfactual revealed greater improvements (“relative negative impacts”, 11.6%). There was no difference between intervention and counterfactual for 1.8% of trials. Moreover, we find that the effect sizes of some individual interventions are high in magnitude and positive, indicating a substantial positive impact of those actions on the whole.

All types of intervention assessed that had more than five trials showed a significant positive effect compared with a counterfactual (Fig. 2a; Fig. S2) – eradication and control of invasive alien and problematic native species (7.07 [6.1–8.04], $p < 0.001$), sustainable management of ecosystems (5.70 [4.66–6.74], $p < 0.001$), habitat loss reduction & restoration (5.58 [4.5–6.7], $p < 0.001$), and establishment and management of protected areas (1.41 [1.03–1.78], $p < 0.001$).

The impact of efforts towards sustainable use of species is inconclusive (2.07 [-0.71–4.84], $p=0.15$), with large confidence intervals possibly related to the small number of studies ($N_{\text{studies}}=5$, $N_{\text{trials}}=7$). There were too few studies assessing the impact of interventions of pollution control or climate change adaptation to analyze separately (each had five or fewer trials) but these were included in the calculation of the Overall effect size. Likewise ‘Other’ studies that did not fit into one of the seven key intervention categories (e.g. food supplementation ($N_{\text{studies}}=1$), culling of diseased individuals ($N_{\text{studies}}=1$), and captive breeding and release ($N_{\text{studies}}=5$)) were only included in the calculation of the Overall effect size (categorized as “Other” in the Figures). Studies focused on the impact of controlling problematic native species were combined with studies on IAS given the similarity of the interventions involved.

The impact of actions targeting different levels of ecological organization are positive and significant, with the greatest impact shown for actions targeting species (3.56 [3.14–3.97], $p<0.001$), followed by ecosystems (2.88 [2.46–3.29], $p<0.001$), and then genetic diversity (3.84 [1.35–6.32], $p=0.002$), which had wide confidence intervals because there were few studies (Fig. 2b). In terms of geographic breakdown, effect sizes were positive and significant on all continents (Table S4). Our dataset also shows that effect sizes are positive and significant across each of the differing approaches to measuring the impact of an intervention (Table S5).

The timespan of datasets analyzed in the different studies was highly variable, with the shortest being 1 month and the longest 110 years (median=4.7, mean=7.4) (Fig. 3). These date back to 1890, and show a trend that studies mainly focused on protected areas until the 1990s, but after this, diversified to a wider range of interventions. Studies with a longer duration were not significantly more likely to show more beneficial or detrimental impacts of conservation actions than shorter studies (Fig. S3). Meta-regression of effect sizes against year of publication (Fig. 4) indicated that more recent studies were more likely to show a positive effect of conservation action, although the low R^2 shows that other factors influence effect sizes, as would be expected (because many factors in addition to the intervention determine effect sizes).

Half of the studies that met our meta-analysis inclusion criteria (95/186) were conducted in Western Europe, North America, Australia, and New Zealand (Fig. 3). Of the seven intervention types examined, the largest proportion focused on terrestrial and marine protected areas (38%) and the eradication and control of invasive alien and problematic native species (25%). Fewer studies evaluated other conservation actions for ecosystems (e.g. restoration, habitat conservation policy, sustainable management), for species (e.g. sustainable use, reintroductions), and for genetic diversity (e.g. supplementation, supportive breeding; Table S3).

Our results remain largely unchanged in sensitivity analyses designed to test the impact of different methodological considerations (Fig. S4), including: imputing the rate of change when it was zero in either the intervention or counterfactual; nesting trials within studies; and undertaking a supplemental literature search. Cumulative meta-analysis revealed that effect sizes stabilize after the addition of studies published from approximately 2011 onwards (Fig S5), and in assessing publication bias, the symmetrical nature of our funnel plots (Fig. S6), combined with a Fail-safe N of 1,280, led us to conclude that any publication bias in our dataset is minimal.

Discussion

We have shown that across a full suite of conservation actions and intervention types, multiple levels and metrics of biodiversity, and over a century of action, conservation has improved the state of biodiversity – or at least slowed its decline – compared with no conservation action. Our calculated effect sizes are often large and positive, meaning that the outcomes from conservation actions are substantially better than no action at all.

Among the different conservation actions evaluated in our study, the eradication, control and management of IAS showed the largest impact of conservation action (as highlighted by the largest effect size in our intervention groupings), followed by actions to reduce habitat loss and degradation, sustainable management of ecosystems, and protected areas (Fig. 2a). IAS eradication and control has generated some of the most striking conservation successes reported to date, particularly on islands (20, 21). There were numerous studies evaluating the impact of IAS control and eradication that were excluded from our meta-analysis because they report data for only one time point, but these tend to also show a positive impact.

Although few in number, studies assessing actions to reduce habitat loss and degradation, including restoration, did show positive results (Fig. 2a) consistent with global analyses (22, 23). Efforts to ensure sustainable management of ecosystems, particularly on land, also generally increased native species abundance and habitat cover (Fig. 2a). These results are consistent with global analyses of sustainable management of ecosystems under agriculture and forestry (24, 25).

Protected areas have been shown to be effective in reducing conversion of natural land cover (10), terrestrial habitat loss (15), coral loss (26), tropical forest fires (27), species extinction risk (28), and in increasing biomass and density of marine organisms (14, 29). Protected area effectiveness varies geographically, in terms of their effectiveness in preventing deforestation (30) and reducing anthropogenic pressure (31). Poor performance of protected areas often results from shortfalls in human and financial capacity (32, 33), while protected area downgrading, downsizing and degazettement presents another major challenge in some regions (34). However, our results concur that while their effectiveness is not universal (31), protected areas are an important tool for achieving conservation outcomes (Fig. 2a).

The impacts of efforts to address unsustainable use of species in the ocean and on land were mixed (Fig. 2a), but the number of studies meeting the criteria for inclusion in our meta-analysis was small ($N_{\text{studies}}=5$).

Across our dataset, 137 trials (21% of cases) provide examples in which conservation interventions were not only associated with a negative rate of change, but performed more poorly than counterfactuals for biodiversity state. Eradicating and controlling invasive alien and problematic native species can negatively impact non-target species through incidental damage or mortality, or meso-predator release, and this explains the negative impacts observed for this group of papers. For example, in the United States, application of herbicides to invasive alien plants harmed native forbs (35), and in India, physical removal of invasive alien algae caused further spread and establishment elsewhere (36) (Fig. 1). Protected areas can show negative impacts if there is poor enforcement and insufficient resourcing, leading to higher rates of resource extraction, poaching, or agricultural expansion compared with counterfactual areas (37), or if protection increases the abundance of both predators and the target species in a no-take marine protected area (38). These unintended outcomes emphasize the importance of evidence-

based interventions alongside effective monitoring, so that negative responses can be quickly detected and conservation measures adjusted accordingly.

More and better counterfactual studies are needed for a wider range of conservation interventions and geographic regions. Particular gaps include assessments of pollution control, climate change adaptation, sustainable use of species, habitat loss reduction (beyond protected areas), actions targeting species and genetic diversity, and conservation actions in the Global South.

The finding that contemporary studies were more likely to show a positive effect of conservation actions (Fig. 4) may be due to increases in funding and project-level resources, more targeted interventions, or indeed, that conservation practice is improving over time as lessons are learned from previous failures, and methods are improved. These same factors may help explain the larger effect sizes of more rigorous experimental/quasi-experimental study designs (Table S5), since their application (particularly quasi-experimental) tends to be more recent. That the relationship between effect size vs. study duration (Fig. S3) showed no general trend in direction indicates that valuable insights can be gained even for studies that are short in duration, provided the study design is appropriate. This further underscores the necessity of counterfactual evaluation and meta-analysis to assess conservation impact.

Overall, global conservation efforts have helped to slow declines in biodiversity, and could eventually bend the curve of absolute biodiversity loss (39). Quantifying and evidencing the relative biodiversity outcomes (typically gains) of conservation is crucial, to contextualize and explain declining state indicator and increasing response indicator trends. While the state of biodiversity is declining across the globe in absolute terms, conservation actions work most of the time – the challenge is now to expand these to the scale necessary to reverse the global biodiversity crisis. That is, conservation interventions are working, but there are simply not enough conservation actions implemented, or in the right places. Realizing the highly ambitious vision of the GBF not simply to slow declines of biodiversity by the end of the decade, but to reverse them (2), will require ongoing assessment of the impact of specific conservation interventions, to inform adaptive management with evidence. Even more importantly, however, it will require substantially scaled-up funding and commitment for implementation of demonstrably effective conservation interventions – a real transformational change – which in turn depends on increased political will and investment.

Such an increase in conservation action and associated outcomes will require expanded implementation and significant additional investment across many sectors of society, particularly beyond the traditional conservation sector. Meeting global biodiversity conservation targets to reduce the extinction risk of all species and safeguard sites of international biodiversity importance was estimated to cost around \$80 billion annually over a decade ago (40). A comprehensive global conservation program would require an investment of \$178-524 billion annually (4), much of it focused in highly biodiverse countries. Although high, these costs are dwarfed by the value that biodiversity provides to society through the delivery of ecosystem services (41). Thus, conservation actions are investments rather than payments – and, as our study demonstrates, they are typically investments that yield genuine, high-magnitude positive impacts.

References and Notes:

1. D. Díaz *et al.*, Pervasive human-driven decline of life on Earth points to the need for profound change. *Science* **366**, eaax3100 (2019).
2. Convention on Biological Diversity. Kunming-Montreal Global Biodiversity Framework. CBD/COP/15/L.25. CBD, Montreal, Canada (2022).
<https://www.cbd.int/doc/c/e6d3/cd1d/daf663719a03902a9b116c34/cop-15-l-25-en.pdf>
3. United Nations. Transforming our world: The 2030 Agenda for Sustainable Development. United Nations, New York, USA. <https://sdgs.un.org/publications/transforming-our-world-2030-agenda-sustainable-development-17981> (2015).
4. A. Seidl, K. Mulungu, M. Arlaud, O. van den Heuvel, M. Riva, Finance for nature: A global estimate of public biodiversity investments. *Ecosystem Services* **46**, 101216 (2020).
5. J. Terborgh, *Requiem for Nature* (Island Press, Washington, D.C., 1999).
6. M. Marvier, P. Kareiva, R. Lalasz, Conservation in the Anthropocene: Beyond solitude and fragility. *Breakthrough Journal* <https://thebreakthrough.org/journal/issue-2/conservation-in-the-anthropocene> (2012).
7. P.J. Ferraro, S.K. Pattanayak, Money for nothing? A call for empirical evaluation of biodiversity conservation investments. *PLoS Biol.* **4**, e105 (2006).
8. F.C. Bolam, *et al.*, How many bird and mammal extinctions has recent conservation action prevented? *Conserv. Lett.* e12762 (2020).
9. M. Hoffmann *et al.*, The impact of conservation on the status of the world's vertebrates. *Science* **330**, 1503-1509 (2010).
10. L.N. Joppa, A. Pfaff, Global protected area impacts. *P. Roy. Soc. B-Biol. Sci.* **278**, 1633-1638 (2011).
11. E. Wiik, *et al.*, Experimental evaluation of the impact of a payment for environmental services program on deforestation. *Conserv. Sci. Pract.* **1**, e8 (2019).
12. D.P. Armstrong, *et al.*, Strategic rat control for restoring populations of native species in forest fragments. *Conserv. Biol.* **28**, 713-723 (2014).
13. J.M. Holland, B.M. Smith, J. Storkey, P.J. Lutman, N.J. Aebischer, Managing habitats on English farmland for insect pollinator conservation. *Conserv. Biol.* **182**, 215-222 (2015).
14. S.E. Lester, *et al.*, Biological effects within no-take marine reserves: a global synthesis. *Mar. Ecol. Prog. Ser.* **384**, 33-46 (2009).
15. J. Geldmann, *et al.*, Effectiveness of terrestrial protected areas in reducing habitat loss and population declines. *Biol. Conserv.* **161**, 230-238 (2013).
16. J.E. Bicknell, M.J. Struebig, D.P. Edwards, Z.G. Davies, Improved timber harvest techniques maintain biodiversity in tropical forests. *Curr. Biol.* **24**, R1119-R1120 (2014).
17. Conservation Evidence; www.conservationevidence.com [accessed 08 July 2023].
18. H.S. Wauchope, *et al.*, Evaluating impact using time-series data. *Trends Ecol. Evol.* **36**, 196-205 (2021).
19. I.M. Côté, J.A. Gill, T.A. Gardner, A.R. Watkinson, Measuring coral reef decline through meta-analyses. *Philos. Trans. R. Soc. B-Biol. Sci.* **360**, 385-395 (2005).
20. H.P. Jones, *et al.*, Invasive mammal eradication on islands results in substantial conservation gains. *Proc. Natl Acad. Sci. USA* **113**, 4033-4038 (2016).
21. D.R. Spatz, *et al.*, The global contribution of invasive vertebrate eradication as a key island restoration tool. *Scientific Reports* **12**, 13391 (2022).
22. J.M. Rey Benayas, A.C. Newton, A. Diaz, J.M. Bullock, Enhancement of biodiversity and ecosystem services by ecological restoration: A meta-analysis. *Science* **325**, 1121-1124 (2009).

23. R. Crouzeilles, *et al.*, A global meta-analysis on the ecological drivers of forest restoration success. *Nat. Commun.* **7**, 11666 (2016).
24. K. Fedrowitz, *et al.*, Can retention forestry help conserve biodiversity? A meta-analysis. *J. Appl. Ecol.* **51**, 1669-1679 (2014).
- 5 25. J. Bengtsson, J. Ahnström, A.C. Weibull, The effects of organic agriculture on biodiversity and abundance: a meta-analysis. *J. Appl. Ecol.* **42**, 261-269 (2005).
26. E.R. Selig, J.F. Bruno, A global analysis of the effectiveness of marine protected areas in preventing coral loss. *PLOS ONE* **5**, e9278 (2010).
- 10 27. A. Nelson, K.M. Chomiz, Effectiveness of strict vs. multiple use protected areas in reducing tropical forest fires: A global analysis using matching methods. *PLOS ONE* **6**, e22722 (2011).
28. S.H.M. Butchart, *et al.*, Protecting important sites for biodiversity contributes to meeting global conservation targets. *PLOS ONE* **7**, e32529 (2012).
- 15 29. M. Sciberras, *et al.*, Evaluating the relative conservation value of fully and partially protected marine areas. *Fish Fish.* **16**, 58-77 (2015).
30. M. Heino, *et al.*, Forest loss in protected areas and intact forest landscapes: A global analysis. *PLOS ONE* **10**, e0138918 (2015).
- 20 31. J. Geldmann, A. Manica, N.D. Burgess, L. Coad, A. Balmford, A global-level assessment of the effectiveness of protected areas at resisting anthropogenic pressures. *Proc. Natl Acad. Sci. USA* **16**, 23209-23215 (2019).
32. D.A. Gill, *et al.*, Capacity shortfalls hinder the performance of marine protected areas globally. *Nature* **543**, 665-669 (2017).
33. J. Geldmann, *et al.*, A global analysis of management capacity and ecological outcomes in terrestrial protected areas. *Conserv. Lett.* **11**, e12434 (2018).
- 25 34. R.E. Golden Kroner, *et al.*, The uncertain future of protected lands and waters. *Science* **364**, 881-886 (2019).
35. E.E. Crone, M. Marler, D.E. Pearson, Non-target effects of broadleaf herbicide on a native perennial forb: A demographic framework for assessing and minimizing impacts. *J. Appl. Ecol.* **46**, 673-682 (2009).
- 30 36. B. Kamalakannan, J.J.J. Jeevamani, N.A. Nagendran, D. Pandiaraja, S. Chandrasekaran, Impact of removal of invasive species *Kappaphycus alvarezii* from coral reef ecosystem in Gulf of Mannar, India. *Curr. Sci.* **106**, 1401-1408 (2014).
37. A. Blackman, A. Pfaff, J. Robalino, Paper park performance: Mexico's natural protected areas in the 1990s. *Glob. Environ. Change* **31**, 50-61 (2015).
- 35 38. D. Harasti, K. Martin-Smith, W. Gladstone, Does a no-take marine protected area benefit seahorses? *PLOS ONE* **9**, e105462 (2014).
39. G.M. Mace, *et al.*, Aiming higher to bend the curve of biodiversity loss. *Nature Sustainability* **1**, 448-451 (2018).
- 40 40. D.P. McCarthy, *et al.*, Financial costs of meeting global biodiversity conservation targets: Current spending and unmet needs. *Science* **338**, 946-949 (2012).
41. A. Balmford, *et al.*, Economic reasons for conserving wild nature. *Science* **297**, 950-953 (2003).
42. M.O. O'Brien, J.D. Wilson, Population changes of breeding waders on farmland in relation to agri-environment management. *Bird Study* **58**, 399-408 (2011).
- 45 43. D. Nepstad, *et al.*, Inhibition of Amazon deforestation and fire by parks and indigenous lands. *Conserv. Biol.* **20**, 65-73 (2006).
44. <https://doi.org/10.22024/UniKent/01.01.146>

45. E. Burton, G. Butler, J. Hodgkinson, S. Marshal. in *Community Safety: Innovation and Evaluation* (eds. E. Hogard, R. Ellis, J. Warren) pp. 50–62 (Chester Academic Press, Chester, 2007).
46. A.S. Pullin, G.B. Stewart, Guidelines for systematic review in conservation and environmental management. *Conserv. Biol.* **20**, 1647-1656 (2006).
47. S. Nakagawa, D.W.A. Noble, A.M. Senior, M. Lagisz, Meta-evaluation of meta-analysis: ten appraisal questions for biologists. *BMC Biol.* **15**, **18** (2017).
48. J.W. Bull, M. Maron, How humans drive speciation as well as extinction. *P. Roy. Soc. B-Biol. Sci.* **283**, 20160600 (2016).
49. B.W.T. Coetzee, K.J. Gaston, An appeal for more rigorous use of counterfactual thinking in biological conservation. *Conserv. Sci. Pract.* **3**, e409 (2021).
50. A. Rohatgi, WebPlotDigitizer Version 4.2. <https://automeris.io/WebPlotDigitizer/> (2019).
51. P.J. Ferraro, *et al.*, More strictly protected areas are not necessarily more protective: evidence from Bolivia, Costa Rica, Indonesia, and Thailand. *Environ. Res. Lett.* **8**, 25011 (2013).
52. J. Koricheva, J. Gurevitch, K. Mengersen, *Handbook of Meta-analysis in Ecology and Evolution* (Princeton University Press, Princeton, 2013).
53. M. Borenstein, L.V. Hedges, J.P.T. Higgins, H.R. Rothstein, *Introduction to Meta-analysis* (Wiley & Sons Ltd., Chichester, 2009).
54. C.J. Weir, *et al.*, Dealing with missing standard deviation and mean values in meta-analysis of continuous outcomes: a systematic review. *BMC Med. Res. Methodol.* **18**, 25 (2018).
55. J. Koricheva, J. Gurevitch, Uses and misuses of meta-analysis in plant ecology. *J. Ecol.* **102**, 828–844 (2014).
56. M. Borenstein, L.V. Hedges, J.P.T. Higgins, H.R. Rothstein, *Comprehensive Meta-analysis*, Version 4 (Biostat, Englewood, 2023).
57. P.J.K. McGowan, The need to redress the geographical imbalance in the publication of conservation science. *Oryx* **44**, 328-329 (2010).
58. P.R. Adams, D.B. Orr, C. Arellano, Y.J. Cardoza. Soil and foliar arthropod abundance and diversity in five cropping systems in the coastal plains of North Carolina. *Environ. Entomol.* **46**, 771-783 (2017).
59. C.L. Alados, C.L. *et al.*, Variations in landscape patterns and vegetation cover between 1957 and 1994 in a semiarid Mediterranean ecosystem. *Landscape Ecol.* **19**, 543-559 (2004).
60. D. Alemany, O.O. Iribarne, E.M. Acha, Effects of a large-scale and offshore marine protected area on the demersal fish assemblage in the Southwest Atlantic. *ICES J. Mar. Sci.* **70**, 123-134 (2013).
61. D.C. Allen, H.S. Galbraith, C.C. Vaughn, D.E.A. Spooner, Tale of two rivers: implications of water management practices for mussel biodiversity outcomes during droughts. *Ambio* **42**, 881-891 (2013).
62. C.A. Alo, R.G. Pontius, Jr., Identifying systematic land-cover transitions using remote sensing and GIS: the fate of forests inside and outside protected areas of Southwestern Ghana. *Environ. Plan. B Plan. Des.* **35**, 280-295 (2008).
63. A.F.A. Al-Zankana, T. Matheson, D.M. Harper, Secondary production of macroinvertebrates as indicators of success in stream rehabilitation. *River Res. and Appl.* **37**, 408-422 (2021).
64. J.M. Ament, G.S. Cumming, Scale dependency in effectiveness, isolation, and social-ecological spillover of protected areas. *Conserv. Biol.* **30**, 846-855 (2016).
65. K.S. Andam, P.J. Ferraro, M.M. Hanauer, The effects of protected area systems on ecosystem restoration: a quasi-experimental design to estimate the impact of Costa Rica's protected area system on forest regrowth. *Conserv. Lett.* **6**, 317-323 (2013).

66. R.A. Arriagada, C.M. Echeverria, D.E. Moya, Creating protected areas on public lands: Is there room for additional conservation? *PLOS ONE* **11**, e0148094 (2016).
67. G.A. Balme, R. Slotow, L.T.B. Hunter, Impact of conservation interventions on the dynamics and persistence of a persecuted leopard (*Panthera pardus*) population. *Biol. Conserv.* **142**, 2681-2690 (2009).
68. T.A. Belligan, *et al.*, Rapid recovery of macroinvertebrates in a South African stream treated with rotenone. *Hydrobiologica* **834**, 1-11 (2019).
69. L.D. Bennion, J.A. Ferguson, L.F. New, C.B. Schultz, Community-level effects of herbicide-based restoration treatments: structural benefits but at what cost? *Restor. Ecol.* **28**, 553-563 (2020).
70. B.A. Berejikian, D.M. Van Doornik, Increased natural reproduction and genetic diversity one generation after cessation of a steelhead trout (*Oncorhynchus mykiss*) conservation hatchery program. *PLOS ONE* **13**, e0190799 (2018).
71. A.E. Beresford, *et al.*, Protection reduces loss of natural land-cover at sites of conservation importance across Africa. *PLOS ONE* **8**, e65370 (2013).
72. D. Bhaskar, P.S. Easa, K.A. Sreejith, J. Skejo, A. Hochkirch, Large scale burning for a threatened ungulate in a biodiversity hotspot is detrimental for grasshoppers (Orthoptera: Caelifera). *Biodivers. Conserv.* **28**, 3221-3237 (2019).
73. T.O. Bickel, G.P. Closs, Impact of partial removal of the invasive macrophyte *Lagarosiphon major* (Hydrocharitaceae) on invertebrates and fish. *River Res. Appl.* **25**, 734-744 (2009).
74. T.L.F. Bird, A. Bouskila, E. Groner, P. Bar Kutiel, Can vegetation removal successfully restore coastal dune biodiversity? *Applied Sciences-Basel* **10**, 2310 (2020).
75. M. Biro, J. Boloni, Z. Molnar, Use of long-term data to evaluate loss and endangerment status of Natura 2000 habitats and effects of protected areas. *Conserv. Biol.* **32**, 660-671 (2018).
76. R.E. Blyth-Skyrme, M.J. Kaiser, J.G. Hiddink, G. Edwards-Jones, P.J.B. Hart, Conservation benefits of temperate marine protected areas: variation among fish species. *Conserv. Biol.* **20**, 811-820 (2006).
77. R.U. Bobiles, V.S. Soliman, Y. Nakamura, Partially protected marine area renders non-fishery benefits amidst high fishing pressure: A case study from eastern Philippines. *Reg. Stud. Mar. Sci.* **3**, 225-233 (2016).
78. E. Bonnaud, *et al.*, Top-predator control on islands boosts endemic prey but not mesopredator. *Anim. Conserv.* **13**, 556-567 (2010).
79. A.B. Bos, *et al.*, Global data and tools for local forest cover loss and REDD plus performance assessment: Accuracy, uncertainty, complementarity and impact. *Int. J. Appl. Earth Obs. Geoinf.* **80**, 295-311 (2019).
80. A.D. Bosch-Serra, R. Padro, R.R. Boixadera-Bosch, J. Orobítg, M.R. Yague, Tillage and slurry over-fertilization affect oribatid mite communities in a semiarid Mediterranean environment. *Appl. Soil Ecol.* **84**, 124-139 (2014).
81. P.P. Bosu, M.M. Apetorgbor, E.E. Nkrumah, K.P. Bando, The impact of *Broussonetia papyrifera* (L.) vent. on community characteristics in the forest and forest-savannah transition ecosystems of Ghana. *Afr. J. Ecol.* **51**, 528-535 (2013).
82. N.A. Bourg, W.J. McShea, V. Herrmann, C.M. Stewart, Interactive effects of deer exclusion and exotic plant removal on deciduous forest understory communities. *AoB Plants* **9**, plx046 (2017).
83. J.S. Brandt, V. Butsic, B. Schwab, T. Kuemmerle, V.C. Radeloff, The relative effectiveness of protected areas, a logging ban, and sacred areas for old-growth forest protection in southwest China. *Biol. Conserv.* **181**, 1-8 (2015).

84. C.L.M. Brenes, K.W. Jones, P. Schlesinger, J. Robalino, L. Vierling, The impact of protected area governance and management capacity on ecosystem function in Central America. *PLOS ONE* **13**, e0205964 (2018).
85. T.M. Brereton, M.S. Warren, D.B. Roy, K. Stewart, The changing status of the Chalkhill blue butterfly *Polyommatus coridon* in the UK: the impacts of conservation policies and environmental factors. *J. Insect Conserv.* **12**, 629-638 (2008).
86. J. Bried, V.C. Neves, Habitat restoration on Praia Islet, Azores Archipelago, proved successful for seabirds, but new threats have emerged. *Airo* **23**, 25-35 (2015).
87. A.B. Brink, *et al.*, Indicators for assessing habitat values and pressures for protected areas-an integrated habitat and land cover change approach for the Udzungwa Mountains National Park in Tanzania. *Remote Sens.* **8**, 862 (2016).
88. E. Bro, P. Mayot, F. Reitz, Effectiveness of habitat management for improving grey partridge populations: a BACI experimental assessment. *Anim. Biodivers. Conserv.* **35**, 405-413 (2012).
89. J.M. Brooke, *et al.*, Effects of fertilization and crown release on white oak (*Quercus alba*) masting and acorn quality. *For. Ecol. Manag.* **433**, 305-312 (2019).
90. L.P. Brower, *et al.*, Quantitative changes in forest quality in a principal overwintering area of the monarch butterfly in Mexico, 1971–1999. *Conserv. Biol.* **16**, 346-359 (2002).
91. D. Bruggeman, P. Meyfroidt, E.F. Lambin, Impact of land-use zoning for forest protection and production on forest cover changes in Bhutan. *Appl. Geogr.* **96**, 153-165 (2018).
92. P.G. Cardoso, D. Raffaelli, A.I. Lillebo, T. Verdelhos, M.A. Pardal, The impact of extreme flooding events and anthropogenic stressors on the macrobenthic communities' dynamics. *Estuar. Coast. Shelf Sci.* **76**, 553-565 (2008).
93. T. Carranza, A. Balmford, V. Kapos, A. Manica, Protected area effectiveness in reducing conversion in a rapidly vanishing ecosystem: the Brazilian Cerrado. *Conserv. Lett.* **7**, 216-223 (2014).
94. B. Cavallo, J. Merz, J. Setka, Effects of predator and flow manipulation on Chinook salmon (*Oncorhynchus tshawytscha*) survival in an imperiled estuary. *Environ. Biol. Fishes* **96**, 393-403 (2013).
95. R.S. Ceia, *et al.*, Throwing the baby out with the bathwater: does laurel forest restoration remove a critical winter food supply for the critically endangered Azores bullfinch? *Biol. Invasions* **13**, 93-104 (2011).
96. C. Chatelain, A. Bakayoko, P. Martin, L. Gautier, Monitoring tropical forest fragmentation in the Zagne-Tai area (west of Tai National Park, Cote d'Ivoire). *Biodivers. Conserv.* **19**, 2405-2420 (2010).
97. T.K. Christensen, J.P. Hounisen, Managing hunted populations through sex-specific season lengths: a case of the common eider in the Baltic-Wadden Sea flyway population. *Eur. J. Wildl. Res.* **60**, 717-726 (2014).
98. N.E. Clark, E.H. Boakes, P.J.K. McGowan, G.M. Mace, R.A. Fuller, Protected areas in South Asia have not prevented habitat loss: a study using historical models of land-use change. *PLOS ONE* **8**, e65298 (2013).
99. J. Claudet, D. Pelletier, J.Y. Jouvenel, F. Bachet, R. Galzin, Assessing the effects of marine protected area (MPA) on a reef fish assemblage in a northwestern Mediterranean marine reserve: identifying community-based indicators. *Biol. Conserv.* **130**, 349-369 (2006).
100. R. Crates, *et al.*, Sustained and delayed noisy miner suppression at an avian hotspot. *Austral Ecol.* **45**, 636-643 (2020).
101. M.A. Coleman, A. Palmer-Brodie, B.P. Kelaher, Conservation benefits of a network of marine reserves and partially protected areas. *Biol. Conserv.* **167**, 257-264 (2013).

102. P. Cuenca, R. Arriagada, C. Echeverria, How much deforestation do protected areas avoid in tropical Andean landscapes? *Environ. Sci. Policy* **56**, 56-66 (2016).
103. L.M. Curran, S.N. Trigg, A.K. McDonald, D. Astiani, Y.M. Hardiono, Lowland forest loss in protected areas of Indonesian Borneo. *Science* **303**, 1000 (2004).
- 5 104. S.A. Cushman, D.O. Wallin, Rates and patterns of landscape change in the Central Sikhote-alin Mountains, Russian Far East. *Landsc. Ecol.* **15**, 643-659 (2000).
105. C.Z. Ding, *et al.*, Fish assemblage responses to a low-head dam removal in the Lancang River. *Chin. Geogr. Sci.* **29**, 26-36 (2019).
106. M.J. Dornbusch, R. Limb, K.K. Sedivec, Alternative grazing management strategies combat invasive grass dominance. *Nat. Areas J.* **40**, 86-95 (2020).
- 10 107. P. Eguiguren, R. Fischer, S. Gunter, Degradation of ecosystem services and deforestation in landscapes with and without incentive-based forest conservation in the Ecuadorian Amazon. *Forests* **10**, 442 (2019).
108. R.M. Engeman, *et al.*, Dramatic and immediate improvements in insular nesting success for threatened sea turtles and shorebirds following predator management. *J. Exp. Mar. Biol. Ecol.* **395**, 147-152 (2010).
- 15 109. K. Engst, *et al.*, Functional community ecology meets restoration ecology: Assessing the restoration success of alluvial floodplain meadows with functional traits. *J. Appl. Ecol.* **53**, 751-764 (2016).
- 20 110. G. Epstein, A. Foggo, D.A. Smale, Inconspicuous impacts: Widespread marine invader causes subtle but significant changes in native macroalgal assemblages. *Ecosphere* **10**, e02814 (2019).
111. P.J. Ferraro, C. McIntosh, M. Ospina, The effectiveness of the US endangered species act: an econometric analysis using matching methods. *J. Environ. Econ. Manage.* **54**, 245-261 (2007).
- 25 112. A. Ferreira, A.S. Alves, J.C. Marques, S. Seixas, Ecosystem response to different management options in Marine Protected Areas (MPA): A case study of intertidal rocky shore communities. *Ecol. Indic.* **81**, 471-480 (2017).
113. R.H. Field, S. Benke, K. Badonyi, R.B. Bradbury, Influence of conservation tillage on winter bird use of arable fields in Hungary. *Agric. Ecosyst. Environ.* **120**, 399-404 (2007).
- 30 114. A.D. Flesch, A. Esquer, Impacts of riparian restoration on vegetation and avifauna on private and communal lands in northwest Mexico and implications for future efforts. *Air Soil Water Res.* **13** (2020).
115. L. Flores, *et al.*, Effects of wood addition on stream benthic invertebrates differed among seasons at both habitat and reach scales. *Ecol. Eng.* **106**, 116-123 (2017).
- 35 116. Y.S. Foo, S. Numata, Deforestation and forest fragmentation in and around Endau-Rompin National Park, Peninsular Malaysia. *Tropics* **28**, 23-37 (2019).
117. J.L. Forrest, *et al.*, Patterns of land cover change in and around Madidi National Park, Bolivia. *Biotropica* **40**, 285-294 (2008).
- 40 118. S.V. Fowler, Biological control of an exotic scale, *Orthezia insignis* Browne (Homoptera: Ortheziidae), saves the endemic gumwood tree, *Commidendrum robustum* (Roxb.) DC. (Asteraceae) on the island of St. Helena. *Biol. Control* **29**, 367-374 (2004).
119. S. Fox, J.M. Potts, D. Pemberton, D. Crosswell, Roadkill mitigation: trialing virtual fence devices on the west coast of Tasmania. *Aust. Mammal.* **41**, 205-211 (2019).
- 45 120. M.L. Fujitani, E.P. Fenichel, J. Torre, L.R. Gerber, Synthesizing ecological and human use information to understand and manage coastal change. *Ocean Coast. Manag.* **162**, 100-109 (2018).

121. D.L.A. Gaveau, *et al.*, Evaluating whether protected areas reduce tropical deforestation in Sumatra. *J. Biogeogr.* **36**, 2165-2175 (2009).
122. D.L.A. Gaveau, H. Wandono, F. Setiabudi, Three decades of deforestation in southwest Sumatra: have protected areas halted forest loss and logging, and promoted re-growth? *Biol. Conserv.* **134**, 495-504 (2007).
123. D.L.A. Gaveau, *et al.*, Reconciling forest conservation and logging in Indonesian Borneo. *PLOS ONE* **8**, e69887 (2013).
124. R. Giudice, J. Borner, S. Wunder, E. Cisneros, Selection biases and spillovers from collective conservation incentives in the Peruvian Amazon. *Environ. Res. Lett.* **14**, 45004 (2019).
125. L. Gonsalves, B. Law, R. Blakey, Experimental evaluation of the initial effects of large-scale thinning on structure and biodiversity of river red gum (*Eucalyptus camaldulensis*) forests. *Wildl. Res.* **45**, 397-410 (2018).
126. D. Gorman, A. Turra, The role of mangrove revegetation as a means of restoring macrofaunal communities along degraded coasts. *Sci. Total Environ.* **566**, 223-229 (2017).
127. J.M.H. Green, *et al.*, Deforestation in an African biodiversity hotspot: extent, variation and the effectiveness of protected areas. *Biol. Conserv.* **164**, 62-72 (2013).
128. R. Hagglund, *et al.*, Restoration measures emulating natural disturbances alter beetle assemblages in boreal forest. *For. Ecol. Manag.* **462**, 117934 (2020).
129. J.K. Hanford, C.E. Webb, D.F. Hochuli, Management of urban wetlands for conservation can reduce aquatic biodiversity and increase mosquito risk. *J. Appl. Ecol.* **57**, 794-805 (2020).
130. L. Hannan, D.S. Le Roux, R.N.C. Milner, P. Gibbons, Erecting dead trees and utility poles to offset the loss of mature trees. *Biol. Conserv.* **236**, 340-346 (2019).
131. E. Hardt, *et al.*, Does certification improve biodiversity conservation in Brazilian coffee farms? *For. Ecol. Manag.* **357**, 181-194 (2015).
132. K.M. Hartman, B.C. McCarthy, Restoration of a forest understory after the removal of an invasive shrub, Amur honeysuckle (*Lonicera maackii*). *Restor. Ecol.* **12**, 154-165 (2004).
133. J.A. Henden, D. Ehrich, E.M. Soininen, R.A. Ims, Accounting for food web dynamics when assessing the impact of mesopredator control on declining prey populations. *J. Appl. Ecol.* **58**, 104-113 (2021).
134. D. Hervieux, M. Hebblewhite, D. Stepnisky, M. Bacon, S. Boutin, Managing wolves (*Canis lupus*) to recover threatened woodland caribou (*Rangifer tarandus caribou*) in Alberta. *Can. J. Zool.* **92**, 1029-1037 (2014).
135. M.A. Hess, *et al.*, Supportive breeding boosts natural population abundance with minimal negative impacts on fitness of a wild population of Chinook salmon. *Mol. Ecol.* **21**, 5236-5250 (2012).
136. R. Hilborn, *et al.*, Effective enforcement in a conservation area. *Science* **314**, 1266 (2006).
137. K.M. Hinkson, S.C. Richter, Temporal trends in genetic data and effective population size support efficacy of management practices in critically endangered dusky gopher frogs (*Lithobates sevosus*). *Ecol. Evol.* **6**, 2667-2678 (2016).
138. W.W. Hochstedler, B.S. Slaughter, D.L. Gorchov, L.P. Saunders, M.H.H. Stevens, Forest floor plant community response to experimental control of the invasive biennial, *Alliaria petiolata* (garlic mustard). *J. Torrey Bot. Soc.* **134**, 155-165 (2007).
139. J.A. Hostetler, D.P. Onorato, D. Jansen, M.K. Oli, A cat's tale: the impact of genetic restoration on Florida panther population dynamics and persistence. *J. Anim. Ecol.* **82**, 608-620 (2013).

140. L. Huang, Q.Q. Shao, J.Y. Liu, Assessing the conservation effects of nature reserve networks under climate variability over the northeastern Tibetan plateau. *Ecol. Indic.* **96**, 163-173 (2019).
141. X.F. Huang, *et al.*, Larva fish assemblage structure in three-dimensional floating wetlands and non-floating wetlands in the Changjiang River estuary. *Journal of Oceanology and Limnology* **39**, 721-731 (2021).
142. J.M. Igual, M.G. Forero, T. Gomez, J.F. Orueta, D. Oro, Rat control and breeding performance in Cory's shearwater (*Calonectris diomedea*): effects of poisoning effort and habitat features. *Anim. Conserv.* **9**, 59-65 (2006).
143. T. Ito, *et al.*, Responses of soil nematode community structure to soil carbon changes due to different tillage and cover crop management practices over a nine-year period in Kanto, Japan. *Appl. Soil Ecol.* **89**, 50-58 (2015).
144. J. Jimenez, *et al.*, Restoring apex predators can reduce mesopredator abundances. *Biol. Conserv.* **238**, 108234 (2019).
145. G.R. Johnston, Drought increases the impact of introduced European foxes on breeding Australian pelicans. *Wildl. Res.* **43**, 507-514 (2016).
146. L.J. Jones, S.M. Ostojka, M.L. Brooks, M. Hutten, Short-term response of *Holcus lanatus* L. (common velvetgrass) to chemical and manual control at Yosemite National Park, USA. *Invasive Plant Sci. Manag.* **8**, 262-268 (2015).
147. M. Kayal, *et al.*, Marine reserve benefits and recreational fishing yields: The winners and the losers. *PLOS ONE* **15**, e0237685 (2020).
148. T.A. Kennedy, J.C. Finlay, S.E. Hobbie, Eradication of invasive *Tamarix ramosissima* along a desert stream increases native fish density. *Ecol. Appl.* **15**, 2072-2083 (2005).
149. B.K. Kerns, M.A. Day, Prescribed fire regimes subtly alter ponderosa pine forest plant community structure. *Ecosphere* **9**, e02529 (2018).
150. A.H. Khalyani, A.L. Mayer, C.R. Webster, M.J. Falkowski, Ecological indicators for protection impact assessment at two scales in the Bozin and Marakhil protected area, Iran. *Ecol. Indic.* **25**, 99-107 (2013).
151. M.F. Kinnaird, E.W. Sanderson, T.G. O'Brien, H.T. Wibisono, G. Woolmer, Deforestation trends in a tropical landscape and implications for endangered large mammals. *Conserv. Biol.* **17**, 245-257 (2003).
152. M.T. Koenen, R.B. Utych, D.M. Leslie, Jr., Methods used to improve least tern and snowy plover nesting success on alkaline flats. *J. Field Ornithol.* **67**, 281-291 (1996).
153. S. Lachish, H. McCallum, D. Mann, C.E. Pukk, M.E. Jones, Evaluation of selective culling of infected individuals to control Tasmanian devil facial tumor disease. *Conserv. Biol.* **24**, 841-851 (2010).
154. D.C. Laughlin, *et al.*, The hierarchy of predictability in ecological restoration: are vegetation structure and functional diversity more predictable than community composition? *J. Appl. Ecol.* **54**, 1058-1069 (2017).
155. R. Laughton, P.J. Cosgrove, L.C. Hastie, I. Sime, Effects of aquatic weed removal on freshwater pearl mussels and juvenile salmonids in the River Spey, Scotland. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **18**, 44-54 (2008).
156. M.G. Linke, R.S. Godoy, A.S. Rolon, L. Maltchik, Can organic rice crops help conserve aquatic plants in southern Brazil wetlands? *Appl. Veg. Sci.* **17**, 346-355 (2014).
157. J. Liu, *et al.*, Ecological degradation in protected areas: the case of Wolong Nature Reserve for giant pandas. *Science* **292**, 98-101 (2001).
158. A. Lundberg, J. Kapfer, I.E. Maren, Reintroduced mowing can counteract biodiversity loss in abandoned meadows. *Erdkunde* **71**, 127-142 (2017).

159. H. Marchante, H. Freitas, J.H. Hoffmann, Post-clearing recovery of coastal dunes invaded by *Acacia longifolia*: is duration of invasion relevant for management success? *J. Appl. Ecol.* **48**, 1295-1304 (2011).
160. J.C. Marks, G.A. Haden, M. O'Neill, C. Pace, Effects of flow restoration and exotic species removal on recovery of native fish: lessons from a dam decommissioning. *Restor. Ecol.* **18**, 934-943 (2010).
161. C. Marquez, J.P. Gibbs, V. Carrion, S. Naranjo, A. Llerena, Population response of giant galapagos tortoises to feral goat removal. *Restor. Ecol.* **21**, 181-185 (2013).
162. T. Martelloni, *et al.*, Artificial soft sediment resuspension and high density opportunistic macroalgal mat fragmentation as method for increasing sediment zoobenthic assemblage diversity in a eutrophic lagoon. *Mar. Pollut. Bull.* **110**, 212-220 (2016).
163. N. Mashavakure, *et al.*, Soil dwelling beetle community response to tillage, fertilizer and weeding intensity in a sub-humid environment in Zimbabwe. *Appl. Soil Ecol.* **135**, 120-128 (2019).
164. D. Mateos-Molina, M.T. Scharer-Umpierre, R.S. Appeldoorn, J.A. Garcia-Charton, Measuring the effectiveness of a Caribbean oceanic island no-take zone with an asymmetrical BACI approach. *Fish. Res.* **150**, 1-10 (2014).
165. A. Martinez-Abraín, *et al.*, Assessing the effectiveness of a hunting moratorium on target and non-target species. *Biol. Conserv.* **165**, 171-178 (2013).
166. T.M. Mau-Crimmins, Effects of removing *Cynodon dactylon* from a recently abandoned agricultural field. *Weed Res.* **47**, 212-221 (2007).
167. K.G. McAlpine, S.L. Lamoureaux, S.M. Timmins, D.M. Wotton, Native woody plant recruitment in lowland forests invaded by non-native ground cover weeds and mammals. *N. Z. J. Ecol.* **41**, 65-73 (2017).
168. E. Mendoza, R. Dirzo, Deforestation in Lacandonia (southeast Mexico): evidence for the declaration of the northernmost tropical hot-spot. *Biodivers. Conserv.* **8**, 1621-1641 (1999).
169. M. Merkohasanaj, *et al.*, Assessing the environmental effectiveness of the Spanish marine reserve network using remote sensing. *Ecol. Ind.* **107**, 105583 (2019).
170. J.P. Messina, S.J. Walsh, C.F. Mena, P.L. Delamater, Land tenure and deforestation patterns in the Ecuadorian Amazon: conflicts in land conservation in frontier settings. *Appl. Geogr.* **26**, 113-128 (2006).
171. J.J. Miranda, L. Corral, A. Blackman, G. Asner, E. Lima, Effects of protected areas on forest cover change and local communities: evidence from the Peruvian Amazon. *World Dev.* **78**, 288-307 (2015).
172. F. Monti, *et al.*, The price of success: integrative long-term study reveals ecotourism impacts on a flagship species at a UNESCO site. *Anim. Conserv.* **21**, 448-458 (2018).
173. J.H. Moos, S. Schrader, H.M. Paulsen, G. Rahmann, Occasional reduced tillage in organic farming can promote earthworm performance and resource efficiency. *Appl Soil Ecol.* **103**, 22-30 (2016).
174. R. Moreno-Opo, *et al.*, Is it necessary managing carnivores to reverse the decline of endangered prey species? Insights from a removal experiment of mesocarnivores to benefit demographic parameters of the Pyrenean capercaillie. *PLOS ONE* **10**, e0139837 (2015).
175. J. Morsing, S. Kepfer-Rojas, L. Bastrup-Spohr, A.L. Rodriguez, K. Raulund-Rasmussen, Litter legacy after spruce plantation removal hampers initial vegetation establishment. *Basic Appl. Ecol.* **42**, 4-14 (2020).
176. P.J. Mumby, A.R. Harborne, Marine reserves enhance the recovery of corals on Caribbean reefs. *PLOS ONE* **5**, e8657 (2010).

177. M.A.K. Mwangi, *et al.*, Tracking trends in key sites for biodiversity: a case study using Important Bird Areas in Kenya. *Bird Conserv. Int.* **20**, 215-230 (2010).
178. H. Nagendra, S. Pareeth, B. Sharma, C.M. Schweik, K.R. Adhikari, Forest fragmentation and regrowth in an institutional mosaic of community, government and private ownership in Nepal. *Landsc. Ecol.* **23**, 41-54 (2008).
179. M. Narvarte, R. Gonzalez, M. Fernandez, Comparison of Tehuelche octopus (*Octopus tehueltchus*) abundance between an open-access fishing ground and a marine protected area: evidence from a direct development species. *Fish. Res.* **79**, 112-119 (2006).
180. C. Nolte, A. Agrawal, K.M. Silvius, B.S. Soares-Filho, Governance regime and location influence avoided deforestation success of protected areas in the Brazilian Amazon. *Proc. Natl Acad. Sci. USA* **110**, 4956-4961 (2013).
181. S. Nowak, R.W. Myslajek, Response of the wolf (*Canis lupus* Linnaeus, 1758) population to various management regimes at the edge of its distribution range in western Poland, 1951-2012. *Appl. Ecol. Environ. Res.* **15**, 187-203 (2017).
182. P. Nummi, *et al.*, Alien predation in wetlands - the raccoon dog and waterbird breeding success. *Baltic Forestry* **25**, 228-237 (2019).
183. M.J. Osborne, E.W. Carson, T.F. Turner, Genetic monitoring and complex population dynamics: insights from a 12-year study of the Rio Grande silvery minnow. *Evol. Appl.* **5**, 553-574 (2012).
184. W.K. Ottichilo, J. De Leeuw, A.K. Skidmore, H.H.T. Prins, M.Y. Said, Population trends of large non-migratory wild herbivores and livestock in the Masai Mara ecosystem, Kenya, between 1977 and 1997. *Afr. J. Ecol.* **38**, 202-216 (2000).
185. R.L. Paice, J.M. Chambers, B.J. Robson, Outcomes of submerged macrophyte restoration in a shallow impounded, eutrophic river. *Hydrobiologia* **778**, 179-192 (2016).
186. L. Painter, *et al.*, Reconciliation of cattle ranching with biodiversity and social inclusion objectives in large private properties in Paraguay and collective indigenous lands in Bolivia. *Agric. Syst.* **184**, 102861 (2020).
187. L. Pereda-Briones, F. Tomas, J. Terrados, Field transplantation of seagrass (*Posidonia oceanica*) seedlings: Effects of invasive algae and nutrients. *Mar. Pollut. Bull.* **134**, 160-165 (2018).
188. P.G. Peterson, M.F. Merrett, S.V. Fowler, D.P. Barrett, Q. Paynter, Comparing biocontrol and herbicide for managing an invasive non-native plant species: Efficacy, non-target effects and secondary invasion. *J. Appl. Ecol.* **57**, 1876-1884 (2020).
189. A. Pfaff, F. Santiago-Avila, L. Joppa, Evolving protected-area impacts in Mexico: Political shifts as suggested by impact evaluations. *Forests* **8**, 17 (2017).
190. L. Pham, M.G. Jarvis, D. West, G.P. Closs, Rotenone treatment has a short-term effect on New Zealand stream macroinvertebrate communities. *N. Z. J. Mar. Freshw. Res.* **52**, 42-54 (2018).
191. D. Ramler, H. Keckeis, Effects of large-river restoration measures on ecological fish guilds and focal species of conservation in a large European river (Danube, Austria). *Sci. Total Environ.* **686**, 1076-1089 (2019).
192. J.L. Reidy, F.R. Thompson, C. Schwoppe, S. Rowin, J.M. Mueller, Effects of prescribed fire on fuels, vegetation, and Golden-cheeked Warbler (*Setophaga chrysoparia*) demographics in Texas juniper-oak woodlands. *For. Ecol. Manag.* **376**, 96-106 (2016).
193. G. Ren, *et al.*, Effectiveness of China's national forest protection program and nature reserves. *Conserv. Biol.* **29**, 1368-1377 (2015).

194. A. Robley, A.M. Gormley, D.M. Forsyth, B. Triggs, Long-term and large-scale control of the introduced red fox increases native mammal occupancy in Australian forests. *Biol. Conserv.* **180**, 262-269 (2014).
195. C.B. Rohal, C. Cranney, E.L.G. Hazelton, K.M. Kettenring, Invasive *Phragmites australis* management outcomes and native plant recovery are context dependent. *Ecol. Evol.* **9**, 13835-13849 (2019).
196. A. Roopsind, B. Sohngen, J. Brandt, Evidence that a national REDD plus program reduces tree cover loss and carbon emissions in a high forest cover, low deforestation country. *Proc. Natl Acad. Sci. USA* **116**, 24492-24499 (2019).
197. J. Rudolphi, M.T. Jonsson, L. Gustafsson, Biological legacies buffer local species extinction after logging. *J. Appl. Ecol.* **51**, 53-62 (2014).
198. A. Rumm, F. Foeckler, O. Deichner, M. Scholz, M. Gerisch, Dyke-slotting initiated rapid recovery of habitat specialists in floodplain mollusc assemblages of the Elbe River, Germany. *Hydrobiologia* **771**, 151-163 (2016).
199. G.R. Russ, K.I. Miller, J.R. Rizzari, A.C. Alcala, Long-term no-take marine reserve and benthic habitat effects on coral reef fishes. *Mar. Ecol. Prog. Ser.* **529**, 233-248 (2015).
200. S.A. Sader, D.J. Hayes, J.A. Hepinstall, M. Coan, C. Soza, Forest change monitoring of a remote biosphere reserve. *Int. J. Remote Sens.* **22**, 1937-1950 (2001).
201. U.J. Sanchez-Reyes, S. Nino-Maldonado, L. Barrientos-Lozano, J. Trevino-Carreón, Assessment of land use-cover changes and successional stages of vegetation in the natural protected area Altas Cumbres, Northeastern Mexico, using Landsat satellite imagery. *Remote Sens.* **9**, 712 (2017).
202. F.J. Sanderson, *et al.*, Assessing the performance of EU nature legislation in protecting target bird species in an era of climate change. *Conserv. Lett.* **9**, 172-180 (2015).
203. C.B. Schultz, J.A. Ferguson, Demographic costs and benefits of herbicide-based restoration to enhance habitat for an endangered butterfly and a threatened plant. *Restor. Ecol.* **28**, 564-572 (2020).
204. R. Serrouya, B.N. McLellan, H. van Oort, G. Mowat, S. Boutin, Experimental moose reduction lowers wolf density and stops decline of endangered caribou. *Peer J* **5**, e3736 (2017).
205. C. Seytre, P. Francour, A long-term survey of *Posidonia oceanica* fish assemblages in a Mediterranean marine protected area: emphasis on stability and no-take area effectiveness. *Mar. Freshw. Res.* **65**, 244-254 (2014).
206. P. Shearman, J.A. Bryan, Bioregional analysis of the distribution of rainforest cover, deforestation and degradation in Papua New Guinea. *Austral Ecol.* **36**, 9-24 (2011).
207. E.V. Sheehan, T.F. Stevens, S.C. Gall, S.L. Cousens, M.J. Attrill, Recovery of a temperate reef assemblage in a marine protected area following the exclusion of towed demersal fishing. *PLOS ONE* **8**, e83883 (2013).
208. J. Shimeta, L. Saint, E.R. Verspaandonk, D. Nuggeoda, S. Howe, Long-term ecological consequences of herbicide treatment to control the invasive grass, *Spartina anglica*, in an Australian saltmarsh. *Estuar. Coast. Shelf Sci.* **176**, 58-66 (2016).
209. T. Shumba, *et al.*, Effectiveness of private land conservation areas in maintaining natural land cover and biodiversity intactness. *Global Ecology and Conservation* **22**, e00935 (2020).
210. Z. Siraw, W. Bewket, M.A. Degefu, Effects of community-based watershed development on landscape Greenness and Vegetation Cover in the Northwestern Highlands of Ethiopia. *Earth Syst. Environ.* **4**, 245-256 (2020).
211. M.P. Small, K. Currens, T.H. Johnson, A.E. Frye, J.F. Von Bargen, Impacts of supplementation: genetic diversity in supplemented and unsupplemented populations of

summer chum salmon (*Oncorhynchus keta*) in Puget Sound (Washington, USA). *Can. J. Fish. Aquat. Sci.* **66**, 1216-1229 (2009).

212. J. Smart, *et al.*, Managing uplands for biodiversity: do agri-environment schemes deliver benefits for breeding lapwing *Vanellus vanellus*? *J. Appl. Ecol.* **50**, 794-804 (2013).
213. M. Songer, M. Aung, B. Senior, R. DeFries, P. Leimgruber, Spatial and temporal deforestation dynamics in protected and unprotected dry forests: a case study from Myanmar (Burma). *Biodivers. Conserv.* **18**, 1001-1018 (2009).
214. J. Southworth, H. Nagendra, L.A. Carlson, C. Tucker, Assessing the impact of Celaque National Park on forest fragmentation in western Honduras. *Appl. Geogr.* **24**, 303-322 (2004).
215. B. Stoner-Osborne, The effects of marine protected areas on populations of commercial reef fishes in Moorea, French Polynesia. *Marine Policy* **121**, 104177 (2020).
216. K. Tabor, N.D. Burgess, B.P. Mbilinyi, J.J. Kashaigili, M.K. Steininger, Forest and woodland cover and change in coastal Tanzania and Kenya, 1990 to 2000. *J. East Afr. Nat. Hist.* **99**, 19-45 (2010).
217. M.E. Taylor, M.D. Morecroft, Effects of agri-environment schemes in a long-term ecological time series. *Agric. Ecosyst. Environ.* **130**, 9-15 (2009).
218. J. Tesitel, J. Mladek, K. Fajmon, P. Blazek, O. Mudrak, Reversing expansion of *Calamagrostis epigejos* in a grassland biodiversity hotspot: Hemiparasitic *Rhinanthus major* does a better job than increased mowing intensity. *Appl. Veg. Sci.* **21**, 104-112 (2018).
219. Z. Tonkin, *et al.*, Does localized control of invasive eastern gambusia (Poeciliidae: *Gambusia holbrooki*) increase population growth of generalist wetland fishes? *Austral Ecol.* **39**, 355-366 (2014).
220. C.P. Trentini, *et al.*, Thinning of loblolly pine plantations in subtropical Argentina: Impact on microclimate and understory vegetation. *For. Ecol. Manag.* **384**, 236-247 (2017).
221. I. Tritsch, *et al.*, Do forest-management plans and FSC certification help avoid deforestation in the Congo Basin? *Ecol. Econ.* **175**, 106660 (2020).
222. T.T.A. Truong, *et al.*, Impact of a native invasive weed (*Microstegium ciliatum*) on regeneration of a tropical forest. *Plant Ecol.* **222**, 173-191 (2021).
223. J. Van Den Hoek, M. Ozdogan, A. Burnicki, A.X. Zhu, Evaluating forest policy implementation effectiveness with a cross-scale remote sensing analysis in a priority conservation area of Southwest China. *Appl. Geogr.* **47**, 177-189 (2014).
224. P.A. Vesk, *et al.*, Demographic effects of habitat restoration for the grey-crowned babbler *Pomatostomus temporalis*, in Victoria, Australia. *PLOS ONE* **10**, e0130153 (2015).
225. L. Wan, Y. Zhang, X. Zhang, S. Qi, X. Na, Comparison of land use/land cover change and landscape patterns in Honghe National Nature Reserve and the surrounding Jiansanjiang Region, China. *Ecol. Indic.* **51**, 205-214 (2015).
226. W. Wang, *et al.*, Effectiveness of nature reserve system for conserving tropical forests: a statistical evaluation of Hainan Island, China. *PLOS ONE* **8**, e57561 (2013).
227. C. Watts, *et al.*, Invertebrate community turnover following control of an invasive weed. *Arthropod-Plant Interact.* **9**, 585-597 (2015).
228. K.M. Webb, R.E. Schultz, E.D. Dibble, The influence of invasive aquatic plant removal on diets of bluegill in Minnesota lakes. *J. Aquat. Plant Manag.* **54**, 37-45 (2016).
229. M.J. Weisse, L.C. Naughton-Treves, Conservation beyond park boundaries: The impact of buffer zones on deforestation and mining concessions in the Peruvian Amazon. *Environ. Manage.* **58**, 297-311 (2016).

230. K.J. Wendland, M. Baumann, D.J. Lewis, A. Sieber, V.C. Radeloff, Protected area effectiveness in European Russia: a postmatching panel data analysis. *Land Econ.* **91**, 149-168 (2015).
231. D. Western, S. Russell, I. Cuthill, The status of wildlife in protected areas compared to non-protected areas of Kenya. *PLOS ONE* **4**, e6140 (2009).
232. M.E. Wittmann, *et al.*, Harvesting an invasive bivalve in a large natural lake: species recovery and impacts on native benthic macroinvertebrate community structure in Lake Tahoe, USA. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **22**, 588-597 (2012).
233. X.X. Yao, *et al.*, Effects of long term fencing on biomass, coverage, density, biodiversity and nutritional values of vegetation community in an alpine meadow of the Qinghai-Tibet Plateau. *Ecol. Eng.* **130**, 80-93 (2019).
234. M. Yasue, A. Nellas, A.C.J. Vincent, Seahorses helped drive creation of marine protected areas, so what did these protected areas do for the seahorses? *Environ. Conserv.* **39**, 183-193 (2012).
235. V. Zanzarini, D. Zanchetta, A. Fidelis, Do we need intervention after pine tree removal? The use of different management techniques to enhance Cerrado natural regeneration. *Perspect. Ecol. Conserv.* **17**, 146-150 (2019).
236. I. Zhuravleva, *et al.*, Satellite-based primary forest degradation assessment in the Democratic Republic of the Congo, 2000–2010. *Environ. Res. Lett.* **8**, 24034 (2013).

Acknowledgments: We are grateful to I. Andersen, M. Dalibor, M. Davis, S. Delahaye, A. Hillers, N. Ishii, A. Monard, J.-Y. Pirot, C. Sendashonga, S.N. Stuart, and the late G. Fonseca, for their support in facilitating this project; to J. Koricheva for advice on the meta-analysis; to J. Avaniidhar, A. Lamb, and C. Stengel for assistance with the figures; and to E.J. Milner-Gulland, Bernard W.T. Coetzee, and several anonymous reviewers for their insightful and helpful comments.

Funding: This work was supported by a Global Environment Facility grant to IUCN on “The Global Environmental Commons: Solutions for a Crowded Planet”.

Author contributions:

Conceptualization: PFL, TMB

Methodology: PFL, JWB, JEB, TMB

Investigation: PFL, JWB, JEB, JLO, MHB, MWB, SHMB, JAC, RC, SC, WF, MNF, JG, PG, MH, JH-M, TL, NM, ZM, RM, MGM, SP, BP, GS, KS, JES, KS, JW, SW

Formal analysis: JEB, JWB, PFL

Visualization: JEB, PFL, JWB

Funding acquisition: TMB, CG, NBWM, RGM, DC, WS

Writing – original draft: PFL, JWB, JEB, TMB

Writing – review & editing: PFL, JWB, JEB, JLO, MHB, MWB, SHMB, JAC, DC, RC, SC, WF, MNF, CG, JG, PG, MH, JH-M, TL, NBWM, ZEM, RGM, MGM, SP, BP, WS, GS, KGS, JS, KS, JW, SW, TMB

Competing interests: PFL, JWB, MHB, SHMB, DC, RC, WF, MNF, CG, PG, MH, JH-M, NBWM, RSM, WS, KGS, TMB are employed by organizations engaged in conservation management activities (see author affiliations for details). RC serves on the

External Scientific Advisory Committee for the Louis Vuitton x People for Wildlife partnership.

Data and materials availability: The dataset and code generated and analyzed during the current study is available at (44).

5 **Supplementary Materials**

Materials and Methods

Figs. S1 to S6

Tables S1 to S5

References (45–236)

10




Fig. 1. Schematic representation of different broad categories of conservation impact, with illustrative case studies drawn from our dataset (example reference numbers in superscript). **(A)** ‘Absolute positive impact’ = intervention outperforms counterfactual and there is an increasing biodiversity trend under the intervention. **(B)** ‘Relative positive impact’ = intervention outperforms counterfactual but there is a declining biodiversity trend under both intervention and counterfactual. **(C)** ‘Relative negative impact’ = counterfactual outperforms intervention but there is an increasing biodiversity trend under both intervention and counterfactual. **(D)** ‘Absolute negative impact’ = counterfactual outperforms intervention and there is a declining biodiversity trend under intervention.




Fig. 2. Effect sizes of conservation interventions **(A)** Overall, and by class of intervention, and **(B)** for different levels of ecological organization. The number of intervention vs counterfactual trials of data (n_{trials}), and unique studies (n_{studies}) are reported in parentheses. Mean standardized effect size (Hedges' g) is indicated by the vertical line, and 95% confidence intervals are represented by the bar width. Where the confidence intervals do not overlap zero, the effect size is significant. Vertical dashed lines show zero effect, whilst effect sizes to the right indicate that the intervention is more successful than the counterfactual. Interventions with five or fewer trials (pollution control, climate change adaptation and those classified as 'other'), are not shown, but do contribute to the calculation of the Overall effect size. The icons (Copyright BIP/SCBD) in (A) show the primary Aichi target that the intervention classes align with, however, as shown in Table S1, these interventions also align with a suite of other goals and targets from intergovernmental environmental agreements.




Fig. 3. Characteristics of the studies (n=186) included in the meta-analysis. Each bar represents a single study included within the meta-analysis, and shows (against the x-axis) the timescale (start to end years) covered by the dataset in that study (note break in the scale). The color of each bar denotes the intervention type explored by that study. Points show the year in which the associated study was published in the literature. Inset: world map showing the number of studies carried out in each country represented within the meta-analysis (number of countries=42).




Fig 4. Publication year vs. mean standardized effect sizes for each study, colored by intervention. Blue line and dark grey shaded area show linear line of best fit with 95% confidence intervals. Linear regression is significant ($p < 0.001$) but R^2 is small ($R^2 = 0.03$), as would be expected.