



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

DC3N observations towards high-mass star-forming regions

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

DC3N observations towards high-mass star-forming regions / Rivilla, VM; Colzi, L; Fontani, F; Melosso, M; Caselli, P; Bizzocchi, L; Tamassia, F; Dore, L. - In: MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY. - ISSN 0035-8711. - STAMPA. - 496:2(2020), pp. 1990-1999. [10.1093/mnras/staa1616]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/778791> since: 2022-02-28

*Published:*

DOI: <http://doi.org/10.1093/mnras/staa1616>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

**Rivilla, V.M., Colzi, L., Fontani, F., Melosso, M., Caselli, P., Bizzocchi, L., Tamassia, F., Dore, L., 2020. DC3N observations towards high-mass star-forming regions. Monthly Notices of the Royal Astronomical Society 496, 1990–1999.**

The final published version is available online at:  
<https://doi.org/10.1093/mnras/staa1616>

#### Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

# DC<sub>3</sub>N observations towards high-mass star-forming regions

V. M. Rivilla<sup>1</sup>, L. Colzi<sup>1</sup>, F. Fontani<sup>1</sup>, M. Melosso<sup>2</sup>, P. Caselli<sup>3</sup>, L. Bizzocchi<sup>3</sup>,  
F. Tamassia<sup>4</sup>, and L. Dore<sup>2</sup>

<sup>1</sup> *INAF/Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, I-50125, Florence, Italy*

<sup>2</sup> *Dipartimento di Chimica “Giacomo Ciamician”, Università di Bologna, Via F. Selmi 2, 40126 Bologna, Italy*

<sup>3</sup> *Center for Astrochemical Studies, Max Planck Institut für extraterrestrische Physik Gießenbachstraße 1, 85748 Garching bei München, Germany*

<sup>4</sup> *Dipartimento di Chimica Industriale “Toso Montanari”, Università di Bologna, Viale del Risorgimento 4, 40136 Bologna, Italy*

28 May 2020

## ABSTRACT

We present the study of deuteration of cyanoacetylene (HC<sub>3</sub>N) towards a sample of 28 high-mass star-forming cores divided into different evolutionary stages, from starless to evolved protostellar cores. We report for the first time the detection of DC<sub>3</sub>N towards 15 high-mass cores. The abundance ratios of DC<sub>3</sub>N with respect HC<sub>3</sub>N range in the interval 0.003–0.022, lower than those found in low-mass protostars and dark clouds. No significant trend with the evolutionary stage, or with the kinetic temperature of the region, has been found. We compare the level of deuteration of HC<sub>3</sub>N with those of other molecules towards the same sample, finding weak correlation with species formed only or predominantly in gas-phase (N<sub>2</sub>H<sup>+</sup> and HNC, respectively), and no correlation with species formed only or predominantly on dust grains (CH<sub>3</sub>OH and NH<sub>3</sub>, respectively). We also present a single-dish map of DC<sub>3</sub>N towards the protocluster IRAS 05358+3543, which shows that DC<sub>3</sub>N traces an extended envelope (~0.37 pc) and peaks towards two cold condensations separated from the positions of the protostars and the dust continuum. The observations presented in this work suggest that deuteration of HC<sub>3</sub>N is produced in the gas of the cold outer parts of massive star-forming clumps, giving us an estimate of the deuteration factor prior to the formation of denser gas.

**Key words:** ISM: abundances – ISM: molecules – astrochemistry – stars: formation – stars: massive

## 1 INTRODUCTION

The deuteration level of interstellar molecules is a powerful tool to study the history of star-forming regions. The formation of deuterated species is significantly enhanced during the pre-stellar phases of star-forming regions due to the combination of high gas densities ( $>10^4$  cm<sup>-3</sup>) and low temperatures ( $T \leq 20$  K), which favour the depletion of CO onto interstellar dust surfaces (Caselli et al. 1999; Walmsley et al. 2004). This enhances the gas-phase abundances of several ions (e.g. H<sub>3</sub><sup>+</sup>), in absence of its main destroyer, CO. Then, the reaction of H<sub>3</sub><sup>+</sup> with HD, the main reservoir of deuterium in molecular clouds, produces H<sub>2</sub>D<sup>+</sup> through the reaction  $H_3^+ + HD \rightarrow H_2D^+ + H_2 + 232$  K, when the species involved are in the para-H<sub>2</sub> form. The exothermicity<sup>1</sup> of this reaction prevents the backwards reaction at low temperatures, thus increasing the

gas-phase abundance of H<sub>2</sub>D<sup>+</sup>. This species transfers deuterium to other molecules efficiently, increasing the deuterium fractionation in molecules by several orders of magnitude with respect to the primordial value of D/H  $\sim (1-2) \times 10^{-5}$  in the solar neighborhood (Oliveira et al. 2003; Linsky et al. 2006). Therefore, the deuterated fraction,  $D_{\text{frac}}$ , defined as the column density ratio of one species containing deuterium to its counterpart containing hydrogen, can be used as a useful probe of the evolution of star-forming cores.

In low-mass star-forming cores, Caselli (2002) found that  $D_{\text{frac}}$  significantly increases during the pre-stellar phase, when the core density profile becomes more centrally peaked due to the collapse. As a consequence, the CO freeze-out increases, along with the abundance of deuterated molecules in the gas-phase. Multiple observations have shown that the value of  $D_{\text{frac}}$  reaches its maximum in pre-stellar cores close to gravitational collapse (e.g. Crapsi et al. 2005), and decreases during the later protostellar phases (Emprechtinger et al. 2009).

In high-mass star-forming regions, how  $D_{\text{frac}}$  changes with the evolutionary stage is less clear. In previous works, we have studied the deuteration of different species towards a sample of

<sup>1</sup> The exothermicity of this reaction depends on the orto-para ratio of the species involved, and that can vary between 84 and 256 K ( Millar et al. 1989; Pagani et al. 1992).

high-mass star-forming cores at different evolutionary stages, from pre-stellar to protostellar phases. Fontani et al. (2011) found that enhanced values of  $N_2D^+$  are found towards high-mass pre-stellar cores, similarly to low-mass counterparts. However, other molecular species, such as HNC,  $NH_3$  and  $CH_3OH$ , do not show any clear trend with the evolutionary stage using the same sample of sources (Fontani et al. 2014, 2015; Colzi et al. 2018b).

In this work, we focus on the carbon-chain molecule cyanoacetylene ( $HC_3N$ ). Little is known about its deuteration, since only a few observations have been able to detect it in space.  $DC_3N$  was first detected by Langer et al. (1980) towards the low-mass dark cloud TMC-1, and by Howe et al. (1994) towards several other dark clouds, finding values of  $D_{\text{frac}}$  in the range 0.03–0.13. In the protostellar phase,  $DC_3N$  has been detected only towards a handful of sources: IRAS 16293-2422 (Al-Edhari et al. 2017), L1527 (Sakai et al. 2009 and Araki et al. 2016), Chamaeleon MMS1 (Cordiner et al. 2012), and SVS13-A (Bianchi et al. 2019), with values of  $D_{\text{frac}} \sim 0.03$ –0.4.

There is still no unambiguous detection of  $DC_3N$  towards high-mass star-forming regions. Only two tentative detections have been reported towards Orion KL and the Sgr B2 N2 hot molecular cores (Esplugues et al. 2013; Belloche et al. 2016, respectively). These observations, along with the non detection reported towards NGC 2264 CMM3 (Cordiner et al. 2012), derived upper limits of  $D_{\text{frac}} < 0.015$ , which are lower than those reported in low-mass protostars.

There are two main questions that need to be answered about deuterium fractionation of  $HC_3N$ . First, is  $HC_3N$  deuterated through gas-phase reactions, like simple species such as  $N_2H^+$  (e.g. Fontani et al. 2011), or alternatively on the surface of interstellar dust grains via H-D exchange reactions, as suggested for other carbon-chain molecule ( $c\text{-}C_3H_2$ , Chantzos et al. 2018)? And second, how  $D_{\text{frac}}$  changes with the evolutionary stage of the star-forming core? To properly answer these questions, detections of  $DC_3N$  towards massive cores at different evolutionary stages are needed. With this aim, we present the first study of deuteration of  $HC_3N$  towards a sample of massive star-forming regions spanning a wide range of evolutionary stages, from initial starless cores to more evolved phases. We report the detection of the  $DC_3N$   $J=11-10$  transition towards 15 regions, and present the first maps of  $DC_3N$  in one of the sources of the sample, the high-mass protocluster IRAS 05358+3543.

## 2 OBSERVATIONS

We conducted astronomical observations using the 30m-diameter telescope of the Institut de Radioastronomie Millimétrique (IRAM), located at Pico Veleta (Granada, Spain), as part of the projects 129–12 and 040–19. In the project 129–12, we performed single-pointing observations towards 26 high-mass star-forming cores (see details about the sample in Sect. 3). We used the broad-band Eight MIXer Receiver (EMIR), covering  $\sim 8$  GHz from 89.11 to 96.89 GHz, and the fast Fourier Transform Spectrometer (FTS, Klein et al. 2012) in a mode that provides a channel width of 195 kHz ( $\sim 0.63$  km s $^{-1}$ ). The observed setup includes the  $J=11-10$  transition of  $DC_3N$  at 92.872373 GHz (spectroscopy from the recent laboratory measurements of Melosso et al. *in prep.*), and the  $J=10-9$  transitions of two of the three  $^{13}C$ -isotopologues of  $HC_3N$ ,  $HC^{13}CCN$  and  $HCC^{13}CN$ , at 90.593059 and 90.601777 GHz, respectively (Creswell et al. 1977; Thorwirth et al. 2001). The spatial resolution of the observations, given by the half-power beam width of the antenna (HPBW), can be calculated as  $HPBW('')=2460/\nu(\text{GHz})$ , which gives  $26.5''$  at the frequency of the  $DC_3N$   $J=11-10$  transition.

**Table 1.** List of the observed sources. Cols. 2 and 3 give the equatorial coordinates of the sources. Cols. 4 and 5 give the source distance to the Sun, and the Galactocentric distance, respectively. For the references on distances from the Sun see Fontani et al. (2011). In Col. 6 the  $^{12}C/^{13}C$  ratios, derived using the galactocentric trend by Milam et al. (2005), and the associated uncertainties, are given. In the last column the kinetic temperatures of the clumps derived by Fontani et al. (2015) are listed. For the sources without a direct derivation of  $T_{\text{kin}}$  (indicated with  $^a$ ), the average value for that evolutionary stage was taken.


Source	$\alpha$ (J2000) (h m s)	$\delta$ (J2000) ( $^{\circ}$ $'$ $''$ )	$d$ (kpc)	$D_{GC}$ (kpc)	$^{12}C/^{13}C$	$T_{\text{kin}}$ (K)
HMSC (cold)						
I00117-mm2	00:14:26.3	+64:28:28.0	1.8	9.5	69 $\pm$ 21	14
G034-G2(mm2)	18:56:50.0	+01:23:08.0	2.9	6.3	50 $\pm$ 17	16 $^a$
G034-F1(mm8)	18:53:19.1	+01:26:53.0	3.7	5.8	47 $\pm$ 16	16 $^a$
G034-F2(mm7)	18:53:16.5	+01:26:10.0	3.7	5.8	47 $\pm$ 16	16 $^a$
G028-C1(mm9)	18:42:46.9	-04:04:08.0	5.0	4.7	40 $\pm$ 15	17
I20293-WC	20:31:10.7	+40:03:28.0	2.0	8.3	62 $\pm$ 19	17
I22134-B	22:15:05.8	+58:48:59.0	2.6	9.5	69 $\pm$ 21	17
05358-D1	05:39:11.5	35:45:46.2	1.8	10.3	74 $\pm$ 22	26
05358-D2	05:39:11.5	35:45:23.9	1.8	10.3	74 $\pm$ 22	20
HMSC (warm)						
AFGL5142-EC	05:30:48.7	+33:47:53.0	1.8	10.3	74 $\pm$ 22	25
05358-mm3	05:39:12.5	+35:45:55.0	1.8	10.3	74 $\pm$ 22	30
I22134-G	22:15:10.5	+58:48:59.0	2.6	9.5	69 $\pm$ 21	25
HMPO						
I00117-mm1	00:14:26.1	+64:28:44.0	1.8	9.5	69 $\pm$ 21	20
AFGL5142-mm	05:30:48.0	+33:47:54.0	1.8	10.3	74 $\pm$ 22	34
05358-mm1	05:39:13.1	+35:45:51.0	1.8	10.3	74 $\pm$ 22	39
18089-1732	18:11:51.4	-17:31:28.0	3.6	5.0	43 $\pm$ 15	38
18517+0437	18:54:14.2	+04:41:41.0	2.9	6.4	51 $\pm$ 17	40 $^a$
G75-core	20:21:44.0	+37:26:38.0	3.8	8.4	63 $\pm$ 19	96
I20293-mm1	20:31:12.8	+40:03:23.0	2.0	8.3	62 $\pm$ 19	36
I21307	21:32:30.6	+51:02:16.0	3.2	9.3	68 $\pm$ 20	21
I23385	23:40:54.5	+61:10:28.0	4.9	11.4	81 $\pm$ 23	37
UC HII						
G5.89-0.39	18:00:30.5	-24:04:01.0	1.3	7.2	55 $\pm$ 18	32 $^a$
I19035-VLA1	19:06:01.5	+06:46:35.0	2.2	7.0	54 $\pm$ 18	39
19410+2336	19:43:11.4	+23:44:06.0	2.1	7.7	58 $\pm$ 18	21
ON1	20:10:09.1	+31:31:36.0	2.5	8.0	60 $\pm$ 19	26
I22134-VLA1	22:15:09.2	+58:49:08.0	2.6	9.5	69 $\pm$ 21	47
23033+5951	23:05:24.6	+60:08:09.0	3.5	10.2	74 $\pm$ 21	25
NGC7538-IRS9	23:14:01.8	+61:27:20.0	2.8	9.9	72 $\pm$ 21	32 $^a$

The spectra were calibrated using the GILDAS<sup>2</sup> software package. The antenna temperature units,  $T_A^*$ , were converted to main beam brightness temperature,  $T_{\text{mb}}$ , via the relation  $T_A^* = T_{\text{mb}} \times (B_{\text{eff}}/F_{\text{eff}})$ , using the corresponding telescope efficiencies<sup>3</sup>. The flux density calibration uncertainty of the observations is  $\sim 20\%$ . More details of the observations (e.g. weather conditions, background subtraction, focus and pointing) are presented in Fontani et al. (2015).

In the 040–19 project, we mapped with multiple telescope pointings the molecular emission of one of the star-forming regions of the sample, the high-mass protocluster IRAS 05358+3543 (hereafter 05358). We also observed the  $J=11-10$  transition of  $DC_3N$ , and the  $J=10-9$  transitions of  $HC^{13}CCN$  and  $HCC^{13}CN$ , during

<sup>2</sup> The GILDAS software is available at <http://www.iram.fr/IRAMFR/GILDAS>

<sup>3</sup> <http://www.iram.es/IRAMES/mainWiki/Iram30mEfficiencies>




**Figure 1.** DC<sub>3</sub>N  $J=11-10$  detections towards the sample of high-mass star-forming regions studied in this work. The observed IRAM 30m telescope spectra are shown with a black histogram, while the best LTE fits are shown with red curves. Each panel is centered at the systemic velocity of each source. The name of each source is indicated above each panel, coloured by evolutionary groups: dark green for *cold* HMSCs, black for *warm* HMSCs, red for HMPOs and blue for UC HII regions.

observations performed in the periods 31st July to 5th August 2019 and 18th to 19th September 2019. As for the single-pointing observations, we used the 3mm receiver EMIR, and the FTS spectrometer with a frequency resolution of 195 kHz ( $\sim 0.63$  km s<sup>-1</sup>). The observations were made in position-switching mode. Pointing was checked every 1.5 h, and focus was corrected at the beginning of the observations, at dawn and every 4-6 h. The molecular datacubes were produced from On-The-Fly (OTF) mapping, covering an area of  $120'' \times 120''$  (corresponding to  $\sim 1$  pc  $\times$  1 pc, at the source distance of 1.8 kpc; Heyer et al. 1996), with a central position of RA(J2000) = 05h:39m:13s.0, DEC(J2000) = 35°:45':51''. The integration time for each OTF map (vertical + horizontal + calibration times) was of  $\sim 20$  minutes. The antenna temperature  $T_A^*$  was converted to main beam temperature  $T_{mb}$  by using the same expression given above. Flux calibration uncertainties of 20% are considered in the analysis. The GILDAS packages CLASS and GREG were used to reduce and post-process the data. Baselines were all fitted by constant functions or polynomials of order 1. We built the molecular data cube convolving the OTF map with a Gaussian kernel, using a regularly spaced grid with pixel size of  $9''$ .

### 3 SOURCE SAMPLE

The source sample, already used in several works (Fontani et al. 2011, 2014, 2015; Colzi et al. 2018b,a; Mininni et al. 2018), includes high-mass star-forming cores spanning a wide range of evolutionary stages:

- High-Mass Starless Cores (HMSCs), which are not directly associated with indicators of on-going star formation, such as embedded infrared sources, outflows, or masers. As in previous works, we have divided this group into two subgroups: *cold* and *warm*. The latter includes three regions (AFGL5141-EC, 05358-mm3 and I22134-G) that have temperatures  $T_{kin} \geq 25$  K, and that can be externally heated by nearby protostellar objects detected by high-angular resolution observations (Zhang et al. 2002; Busquet et al. 2011; Sánchez-Monge et al. 2011; Colzi et al. 2019). We note that our observations of these three *warm* HMSCs are likely contaminated by the emission of nearby protostellar sources, which fall in the IRAM 30m beam of  $26.5''$ . This will be discussed in Section 5. We have added to the *cold* HMSC group two new cores that have been identified in the DC<sub>3</sub>N maps of the 05358 region presented in this work (Sect. 4.2): 05358-D1 and D2.



**Figure 2.** Differences between the velocities (upper panel) and FWHM (lower panel) of DC<sub>3</sub>N and the <sup>13</sup>C-isotopologues. Filled(empty) circles correspond to HCC<sup>13</sup>CN(HC<sup>13</sup>CCN). The different colors indicate the different evolutionary groups, as indicated. The shaded area denotes the region  $\pm 1 \text{ km s}^{-1}$ .

- High-Mass Protostellar Objects (HMPOs), which are associated with molecular outflows, and/or faint radio continuum emission ( $S_\nu$  at  $3.6 \text{ cm} < 1 \text{ mJy}$ ) likely tracing a radio-jet, and/or infrared sources.
- Ultracompact (UC) HII regions; associated with a strong radio-continuum emission ( $S_\nu$  at  $3.6 \text{ cm} \geq 1 \text{ mJy}$ ), which likely traces photoionised gas by nascent massive stars.

All the selected sources are located at distances  $d \leq 5 \text{ kpc}$ . The full list of the observed sources is shown in Table 1. More details about how the sources were selected and classified in the different evolutionary stages are given in Fontani et al. (2011). In summary, we observed 28 high-mass star-forming cores divided into: 9 *cold* HMSCs, 3 *warm* HMSCs, 9 HMPOs, and 7 UC HII regions.

## 4 DATA ANALYSIS AND RESULTS

### 4.1 Single-pointing observations

The spectra were imported into the MADCUBA package<sup>4</sup> (Martín et al. 2019). The identification of the molecular lines was performed using the SLIM (Spectral Line Identification and Modeling) tool within MADCUBA. SLIM generates synthetic spectra of molecular species under the assumption of Local Thermodynamic Equilibrium (LTE) conditions. We have implemented into SLIM the new DC<sub>3</sub>N spectroscopic data recently obtained in the laboratory by Melosso et al. (*in prep.*).

We detected the DC<sub>3</sub>N  $J=11-10$  transition towards 15 regions. The transitions observed, shown in Figure 1, include 2 *cold* HMSCs, 3 *warm* HMSCs, 4 HMPOs and 6 UC HII regions. To obtain

the best fit of the observed transitions we used the MADCUBA-AUTOFIT tool that compares the observed spectra with LTE synthetic spectra, and provides the best non-linear least-squares fit using the Levenberg–Marquardt (LM) algorithm (see details in Martín et al. 2019). The free parameters of the fit are: column density of the molecule ( $N$ ), excitation temperature ( $T_{\text{ex}}$ ), velocity ( $v_{\text{LSR}}$ ) and full width half maximum ( $FWHM$ ) of the Gaussian profiles of the lines. We assumed that the molecular emission fills the beam of the telescope, and thus we did not apply any beam dilution correction. This assumption is reasonable, since the low energies of the transitions studied ( $E_{\text{up}}=24-27 \text{ K}$ ), which suggest that these molecules trace relatively extended gas. In the particular case of the 05358 region, the emission maps we present in Sect. 4.2 confirm that this assumption is fulfilled. Since we only have one rotational transition, we fixed the value of  $T_{\text{ex}}$  to the kinetic temperature  $T_{\text{kin}}$  of the cores (Table 1). We note that the column density of DC<sub>3</sub>N can vary by factors  $< 1.3(1.7)$  depending of the  $T_{\text{ex}}$  assumed in the range 15–50 K (15–100 K). The other parameters ( $N$ ,  $v_{\text{LSR}}$  and  $FWHM$ ) were left free whenever possible. Only if the LM algorithm did not converge we fixed  $v_{\text{LSR}}$  and/or  $FWHM$  to values that visually match the observed lines. The resulting fits are shown in red overplotted to the observed spectra in Figure 1, and the resulting parameters are presented in Table 2. In the case of non detections, upper limits for the column density of DC<sub>3</sub>N were derived with SLIM, which uses the formula  $3\sigma \times \Delta v / \sqrt{N_{\text{chan}}}$ , where  $\sigma$  is the *rms* noise of the spectra, and  $N_{\text{chan}}$  is the number of channels covered by the linewidth  $\Delta v$ . The derived upper limits for the column density of DC<sub>3</sub>N are shown in Table 2.


To obtain the value of  $D_{\text{frac}}$ , we derived the HC<sub>3</sub>N column densities towards each source with the same procedure used for DC<sub>3</sub>N. Since HC<sub>3</sub>N is an abundant molecule in high-mass star-forming regions, its emission is expected to be optically thick, which prevents a reliable accurate derivation of its column density. For this reason, we opted to analyze the two <sup>13</sup>C-isotopologues of HC<sub>3</sub>N available in the observed frequency band (HC<sup>13</sup>CCN and HCC<sup>13</sup>CN), which are expected to be optically thin. We have used the spectroscopic entries of the Cologne Database for Molecular Spectroscopy (CDMS, Müller et al. 2001, 2005; Endres et al. 2016), based on the laboratory works by Creswell et al. (1977) and Thorwirth et al. (2001). We calculated the average column density value of the <sup>13</sup>C-isotopologues, and assumed the <sup>12</sup>C/<sup>13</sup>C ratio as a function of the Galactocentric distance found by Milam et al. (2005) (Table 1) to convert to HC<sub>3</sub>N column density. The derived values of  $D_{\text{frac}}$  are presented in the last column of Table 2. We note that, unlike the values of the column densities, the values of  $D_{\text{frac}}$  are nearly independent of the assumed  $T_{\text{ex}}$ , whenever DC<sub>3</sub>N and the <sup>13</sup>C-isotopologues of HC<sub>3</sub>N have a similar  $T_{\text{ex}}$ . We found values of  $D_{\text{frac}}$  in the range 0.003–0.022. For the sources with no DC<sub>3</sub>N detection but HC<sup>13</sup>CCN and HCC<sup>13</sup>CN detections, we give the upper limits of  $D_{\text{frac}}$ . Only in one case, the UC HII region I19035-VLA1, DC<sub>3</sub>N was detected but not the <sup>13</sup>C-isotopologues. For this source we show in Table 2 the derived lower limit of  $D_{\text{frac}}$ ,  $\geq 0.022$ .

To make sure that the <sup>13</sup>C-isotopologues arise from similar gas than DC<sub>3</sub>N, we have checked that their velocities and linewidths are similar. We show in Figure 2 that the differences of the velocities ( $\Delta v_{\text{LSR}}$ ) and linewidths ( $\Delta FWHM$ ) between DC<sub>3</sub>N and the <sup>13</sup>C-isotopologues are always within a narrow range of  $\pm 1 \text{ km s}^{-1}$ , considering the uncertainties derived by the fits. This similarity in the kinematics supports our assumption that DC<sub>3</sub>N and the <sup>13</sup>C-isotopologues are likely tracing similar gas.

To compare the deuteration of HC<sub>3</sub>N with that of other molecules already studied in previous works towards the same sam-

<sup>4</sup> Madrid Data Cube Analysis on ImageJ is a software developed at the Center of Astrobiology (CAB) in Madrid; <http://cab.inta-csic.es/madcuba/Portada.html>.





**Figure 3.** Values of  $D_{\text{frac}}$  derived for  $\text{HC}_3\text{N}$  in the sample of high-mass star-forming regions studied in this work (*warm* HMSC, HMPO and UC HII regions), compared with those derived for other species in previous works. The different colours of the circles correspond to different evolutionary groups, as indicated in each panel. Upper limits are denoted by triangles. The outputs of the Kendall  $\tau$  test ( $\tau$  and double-side  $p$  value) are indicated in the upper right corner of each panel.


ple, we show in Figure 3 the  $D_{\text{frac}}(\text{HC}_3\text{N})$  versus those derived for  $\text{N}_2\text{H}^+$  (Fontani et al. 2011), HNC (Colzi et al. 2018b),  $\text{NH}_3$  (Fontani et al. 2015) and  $\text{CH}_3\text{OH}$  (Fontani et al. 2015). We performed Kendall’s  $\tau$  tests (Kendall 1938) to search for possible correlations between the set of values of  $D_{\text{frac}}$  of the different species. We considered only the sources in which the deuterated species have been detected (no upper limits). The initial guess (null hypothesis) is that the two datasets are not correlated. The Kendall’s  $\tau$  correlation parameter can adopt values between -1 and 1: is equal to 1 if a perfect correlation is found, 0 if there is no correlation (initial guess fulfilled), and -1 if there is perfect anticorrelation. As an output of the test we also give the 2-sided  $p$  value, which gives the level of significance of the test between 0 and 1: 0 when it is statistically significant and 1 when it is totally not significant. The results of the tests are shown in the upper right corner of each panel in Figure 3. The deuteration of  $\text{HC}_3\text{N}$  shows better correlations with  $\text{N}_2\text{H}^+$  and HNC rather than  $\text{NH}_3$  and  $\text{CH}_3\text{OH}$ . There are (weak) correlations with  $\text{N}_2\text{H}^+$  and HNC ( $\tau=0.47$  and  $0.37$ , respectively), which are statistically significant (6% and 13% probability that the initial guess is correct, i.e., no correlation). For  $\text{NH}_3$  and  $\text{CH}_3\text{OH}$ , there is no correlation ( $\tau=0.1$  and  $0.0$ , respectively), with 75% and

100% probability of no correlation, respectively. We discuss the implications of these results in Sect. 5.

## 4.2 Maps of the 05358+3543 protocluster

We produced the integrated maps of the  $J=11-10$  transition of  $\text{DC}_3\text{N}$ , and the combined integrated map of the  $J=10-9$  transitions of the two  $^{13}\text{C}$ -isotopologues ( $\text{HC}^{13}\text{CCN}$  and  $\text{HCC}^{13}\text{CN}$ ), which are shown in Figure 4. We also indicate in the figure the positions of the continuum sources previously identified in the region with interferometric observations (Colzi et al. 2019): mm1, mm2, mm3 and mm4. For comparison, we also show in the right panel of Figure 4 the continuum map at  $850\ \mu\text{m}$  obtained with the Submillimetre Common User Bolometer Array (SCUBA) of the 15m-diameter James Clerk Maxwell Telescope (JCMT), from Di Francesco et al. (2008). The SCUBA observations have a HPBW of  $22.9''$ , similar to our IRAM 30m observations.

The main protostellar activity in the 05358 protocluster is located in the mm1/mm2/mm3 region, which matches with the SCUBA continuum peak, and where multiple molecular outflows have been observed (Beuther et al. 2002). The source mm1 cor-



**Figure 4.** *Left:* The color scale shows the integrated emission map of the  $J=11-10$  transition of  $\text{DC}_3\text{N}$ . The red contour correspond to 0.4 times the emission peak ( $178 \text{ mK km s}^{-1}$ ). The gray contours show to the combined integrated emission of the  $J=10-9$  transitions of  $\text{HC}^{13}\text{CCN}$  and  $\text{HCC}^{13}\text{CN}$  (0.9, 0.8, 0.7, 0.6, 0.5 and 0.4 times the emission peak, which is  $265 \text{ mK km s}^{-1}$ ). The green squares indicate the positions of the continuum sources mm1, mm2, mm3 and mm4 (Colzi et al. 2019), and the  $\text{DC}_3\text{N}$  peak positions (D1 and D2), respectively. The gray filled circle in the lower left corner indicate the beam of the  $\text{DC}_3\text{N}$  observation. *Right:* The color scale shows the SCUBA  $850 \mu\text{m}$  continuum observations, from Di Francesco et al. (2008). The beam size of the SCUBA map ( $22.9''$ ) is indicated in the lower left corner with a blue open circle overplotted to the IRAM 30m beam (gray filled circle). The gray contour levels are the same as in the left panel, while the red contour levels are 0.9, 0.8, 0.7, 0.6, 0.5 and 0.4 times the peak of the integrated emission map of  $\text{DC}_3\text{N}$ .

responds to one of the HMPOs studied in the previous section, and exhibits compact molecular emission with hot core chemistry (Leurini et al. 2007; and Colzi, *priv. communication*). The source mm3 is one of the *warm* HMSCs of our sample, and it is considered to be pre-stellar because no compact line emission has been detected in this position. The evolutionary state of mm2 and the recently discovered mm4 (Colzi et al. 2019) are still unknown. As shown in the Figure 4, the positions of mm1 and mm3 are separated by only  $8.3''$ , while the beam of the IRAM 30m observations is  $\sim 26''$ . Therefore, as previously discussed, it is likely that the spectra from the *warm* HMSC in mm3 is contaminated by the nearby protostellar activity.

Figure 4 shows that the integrated emission of  $\text{DC}_3\text{N}$  and the  $^{13}\text{C}$ -isotopologues of  $\text{HC}_3\text{N}$  are extended, larger than the IRAM 30m beam. To measure the area of the emitting regions we have considered the contour in which the emission falls to a 40% of the peak emission. The emission area of  $\text{DC}_3\text{N}$  and the  $^{13}\text{C}$ -isotopologues are  $\sim 1446''^2$  and  $\sim 2042''^2$ , respectively. These emitting areas correspond to those of circular regions with diameters 0.37 pc and 0.44 pc, respectively. In both cases the sizes are larger than that of the IRAM 30m beam ( $\sim 0.26$  pc). Therefore, the assumption of emitting region filling the telescope beam used in the previous section is fulfilled, at least, in the case of the 05358 region.




The integrated map of the  $^{13}\text{C}$ -isotopologues of  $\text{HC}_3\text{N}$  peak in the region located between mm1/mm2 and mm4, slightly shifted with respect to the peak of the SCUBA continuum by  $\leq 10''$ . This shift is likely not significant, since it is less than half of the beams of the molecular and continuum observations, and considering some uncertainty of the pointings of the different telescopes.

Interestingly, the morphology of the emission of  $\text{DC}_3\text{N}$  is not fully coincident with that of the  $^{13}\text{C}$ -isotopologues. Figure 4 shows that the overall  $\text{DC}_3\text{N}$  emission is shifted towards SW, with two main peaks, hereafter D1 and D2, not coincident with the peak of the  $^{13}\text{C}$ -isotopologues nor with the SCUBA continuum peak. Since no signs of active star formation towards D1 and D2 have been re-

ported, we considered them as *cold* HMSCs, and added them to the sample of sources (Table 1). From the datacube, we extracted the spectra of circular regions centered at the D1 and D2 positions with diameters matching the IRAM 30m beam. We performed the analysis of  $\text{DC}_3\text{N}$ ,  $\text{HC}^{13}\text{CCN}$  and  $\text{HCC}^{13}\text{CN}$  using the same procedure described in the previous section. The values of  $T_{\text{kin}}$  were derived from  $\text{NH}_3$  (as for the other cores of the sample), using the VLA interferometric maps from Lu et al. (2014), and integrating in a circular area with a diameter matching the IRAM 30 beam. The values obtained for D1 and D2 are 26 and 20 K, respectively, lower than that considered for mm1 and mm3 (39 K and 30 K, respectively; Table 1), which supports their *cold* starless nature. The derived values of  $D_{\text{frac}}$  are  $0.020 \pm 0.011$  and  $0.019 \pm 0.011$  for D1 and D2, respectively (Table 2). These values are higher than those found towards mm1 and mm3,  $0.005 \pm 0.02$  and  $0.008 \pm 0.03$ , respectively.

## 5 DISCUSSION

It is still not clear how  $\text{DC}_3\text{N}$  is formed in star-forming regions. Langer et al. (1980) suggested several ion-molecule reactions that can form it during the cold pre-stellar phase:



and



Other possible chemical pathway might be directly linked to the formation of  $\text{HC}_3\text{N}$  itself. Many observational works, from the



**Table 2.** Total DC<sub>3</sub>N column density and D-fractionation of DC<sub>3</sub>N. Values without errors are fixed in the fit procedure.


Source	$N_{\text{tot}}$ ( $\times 10^{11}$ cm <sup>-2</sup> )	$v_{\text{LSR}}$ (km s <sup>-1</sup> )	$\Delta v$ (km s <sup>-1</sup> )	$D_{\text{frac}}(\text{HC}_3\text{N})$
HMSC (cold)				
I00117-mm2	$\leq 1.6$	–	–	–
G034-G2(mm2)	$\leq 1.4$	–	–	–
G034-F1(mm8)	$\leq 1.3$	–	–	–
G034-F2(mm7)	$\leq 1.3$	–	–	–
G028-C1(mm9)	$\leq 1.7$	–	–	–
I20293-WC	$\leq 1.4$	–	–	$\leq 0.010$
I22134-B	$\leq 1.3$	–	–	–
05358-D1	$5.4 \pm 1.4$	$-16.6 \pm 0.4$	$4.5 \pm 0.9$	$0.020 \pm 0.011$
05358-D2	$4.6 \pm 1.2$	$-16.3 \pm 0.4$	$4.1 \pm 0.8$	$0.019 \pm 0.011$
HMSC (warm)				
AFGL5142-EC	$3.7 \pm 0.9$	$-2.6 \pm 0.2$	$2.7 \pm 0.5$	$0.015 \pm 0.004$
05358-mm3	$3.1 \pm 0.9$	$-16.2 \pm 0.2$	$1.9 \pm 0.5$	$0.008 \pm 0.003$
I22134-G	$2.4 \pm 0.8$	$-18.6 \pm 0.3$	$1.9 \pm 0.6$	$0.010 \pm 0.005$
HMPO				
I00117-mm1	$\leq 1.7$	–	–	–
AFGL5142-mm	$4.5 \pm 1.3$	$-3.2 \pm 0.4$	$3.5 \pm 0.9$	$0.012 \pm 0.005$
05358-mm1	$2.3 \pm 0.8$	$-16.4 \pm 0.2$	$0.9 \pm 0.3$	$0.005 \pm 0.002$
18089-1732	$\leq 1.9$	–	–	$\leq 0.003$
18517+0437	$5.2 \pm 1.4$	$42.9 \pm 0.4$	$4.4 \pm 0.8$	$0.022 \pm 0.006$
G75-core	$9 \pm 3$	$-0.7 \pm 0.3$	$4.3 \pm 0.7$	$0.009 \pm 0.002$
I20293-mm1	$\leq 1.4$	–	–	$\leq 0.003$
I21307	$\leq 1.6$	–	–	–
I23385	$\leq 1.1$	–	–	$\leq 0.012$
UC HII				
G5.89-0.39	$16 \pm 4$	$8.3 \pm 0.2$	$3.4 \pm 0.4$	$0.0030 \pm 0.0005$
I19035-VLA1	$2.1 \pm 0.9$	$30.0 \pm 0.5$	$4 \pm 1$	$\geq 0.022$
19410+2336	$2.4 \pm 0.6$	$21.9 \pm 0.1$	$1.6 \pm 0.2$	$0.009 \pm 0.002$
ON1	$4 \pm 1$	$11.2 \pm 0.5$	5	$0.010 \pm 0.003$
I22134-VLA1	$2.9 \pm 1.1$	$-18.5 \pm 0.6$	$3.9 \pm 1.4$	$0.018 \pm 0.009$
23033+5951	$\leq 1.1$	–	–	$\leq 0.006$
NGC 7538-IRS9	$1.6 \pm 0.5$	$-57.1 \pm 0.2$	1	$0.009 \pm 0.002$

study of its <sup>13</sup>C isotopologues, have suggested that HC<sub>3</sub>N is formed through the gas-phase neutral-neutral reaction between acetylene (C<sub>2</sub>H<sub>2</sub>) and CN (e.g. Takano et al. 1998; Taniguchi et al. 2016; Li et al. 2016; Araki et al. 2016; Taniguchi et al. 2017). Therefore, this pathway might also contribute to form DC<sub>3</sub>N via:



An alternative route would imply surface chemistry through D-H exchange reactions after the freeze-out of HC<sub>3</sub>N, as proposed by Chantzos et al. (2018) for other carbon-chain species like cyclopropenylidene (c-C<sub>3</sub>H<sub>2</sub>). A fourth possibility was suggested by Esplugues et al. (2013), based on gas-phase hot chemistry during the protostellar phase.

We discuss in the following how our detections of DC<sub>3</sub>N towards a sample of massive cores can help us to understand how and when HC<sub>3</sub>N is deuterated during high-mass star-formation. The comparison of  $D_{\text{frac}}$  of different species in our sample (Figure 3) shows that HC<sub>3</sub>N is better correlated with N<sub>2</sub>H<sup>+</sup> and DNC than with NH<sub>3</sub> and CH<sub>3</sub>OH. Interestingly, these species are formed only (the former) and predominantly (the latter) in gas-phase. The opposite case is represented by CH<sub>3</sub>OH and NH<sub>3</sub>, which are produced efficiently on grain surfaces through hydrogenation of N and CO, respectively. The (weak) correlation of HC<sub>3</sub>N with the species


**Figure 5.**  $D_{\text{frac}}$  of HC<sub>3</sub>N as a function of the kinetic temperature  $T_{\text{kin}}$  towards the sample of high-mass star-forming regions studied in this work. The different colours of the circles correspond to different evolutionary groups, as indicated in the upper right corner. Upper limits of  $D_{\text{frac}}$  are denoted by triangles pointing downwards, while the only lower limit is denoted by the triangle pointing upwards.

formed in gas phase, and the null correlation with species formed on grains, supports that the deuteration of HC<sub>3</sub>N is likely produced through gas-phase chemistry rather than surface chemistry.

We have not found a clear correlation between the presence of DC<sub>3</sub>N and the evolutionary stage, since we detected it in 2 *cold* HMSCs, 3 *warm* HMSCs, 4 HMPOs and 6 UC HII regions. Supporting this lack of dependence with the evolutionary phase, we do not find a correlation between  $D_{\text{frac}}$  and the kinetic temperature  $T_{\text{kin}}$  of the sources (Figure 5). We note that although the non-detections of DC<sub>3</sub>N in the sample correspond mainly to *cold* HMSCs (Table 2), this does not imply necessarily that deuteration is lower in this early evolutionary stage. These non-detections are likely due to sensitivity limits of the observations, since in most of the cases, the <sup>13</sup>C-isotopologues are also not detected. We have checked that the 6 *cold* HMSCs with no detection of any of the isotopologues of HC<sub>3</sub>N are those with weaker line intensities in other molecular species. As an example, the N<sub>2</sub>H<sup>+</sup> observations by Fontani et al. (2011) showed that these 6 sources have integrated line intensities  $< 3$  K km s<sup>-1</sup>, while the other HMSCs (where DC<sub>3</sub>N is detected) have 7–43 K km s<sup>-1</sup>. Another factor that might limit the detection of DC<sub>3</sub>N in *cold* HMSCs is spectral dilution of the line profiles. These cores exhibit narrow linewidths in other deuterated species, e.g. 0.5–1.65 km s<sup>-1</sup> in the N<sub>2</sub>D<sup>+</sup>(2-1) transition (Fontani et al. 2011). These linewidths are similar or slightly larger than the spectral resolution of the observations ( $\sim 0.65$  km s<sup>-1</sup>), which may produce some dilution of the line intensities.

The presence of similar levels of deuteration of DC<sub>3</sub>N in cores with different evolutionary stages does not favor the hypothesis of formation on the surface of dust grains, or at high temperatures during the protostellar phase. In these two scenarios, the DC<sub>3</sub>N emission should be predominantly concentrated around the protostars, and higher values of  $D_{\text{frac}}$  should be expected in the protostellar phase, due to thermal desorption and hot chemistry, respectively, triggered by the protostellar heating. However, we do not find any preference towards protostellar cores. Moreover, the DC<sub>3</sub>N map towards the 05358 protocluster (Figure 4) shows that the emission is


extended in the region ( $\sim 0.37$  pc), with the main deuteration peaks (D1 and D2) shifted with respect to the dust continuum peak where the protostellar activity is on-going. Furthermore, hot chemistry in the hot core phase seems unlikely, since it is likely too slow to significantly modify the deuteration levels after the desorption of grain mantles (Charnley et al. 1997 and Osamura et al. 2004).

Alternatively, the relatively constant level of  $\text{HC}_3\text{N}$  deuteration during the different evolutionary phases can be naturally explained if  $\text{DC}_3\text{N}$  mainly arises from material in the outer parts of the clumps, which is common to pre-stellar and protostellar cores. In a region with on-going star-formation, only the inner gas is heated by the protostar, while the outer envelope remains cold and nearly unaltered from the pre-stellar phase (e.g. Aikawa et al. 2012). The lower values of  $D_{\text{frac}}$  of  $\text{HC}_3\text{N}$  with respect to other species that mainly trace the inner regions of the cores, such as  $\text{N}_2\text{H}^+$  or  $\text{NH}_3$  (Figure 3), points also towards an origin of  $\text{DC}_3\text{N}$  in the outer envelope, where the gas is less dense, and hence deuteration is less efficient. To further support this hypothesis, we have compared the linewidths of  $\text{DC}_3\text{N}$  with those of  $\text{N}_2\text{D}^+$  and  $o\text{-NH}_3\text{D}$  (from Fontani et al. 2011 and Fontani et al. 2011, respectively). We show in Figure 6 that the linewidths of  $\text{DC}_3\text{N}$  are in most cases larger than those of  $\text{N}_2\text{D}^+$  and  $o\text{-NH}_3\text{D}$ , as expected if the former traces the diffuse outer part of the envelope, and the two latter trace the inner compact regions. We note that the few cases in which  $\text{N}_2\text{D}^+$  or  $o\text{-NH}_3\text{D}$  have larger linewidths than  $\text{DC}_3\text{N}$  might be explained by broadening produced by protostellar activity such as molecular outflows.

In the external part of a star-forming region, the interstellar radiation field (ISRF) is higher, since it is less shielded by the presence of dust. This allows that a higher fraction of carbon is in atomic form, which starts a more rich carbon-chain chemistry. This effect has been detected observationally towards the L1544 pre-stellar core, where many carbon-chain molecules, including  $\text{HC}_3\text{N}$ , arise from the external regions of the core, avoiding the inner dust peak (Spezzano et al. 2017). We have observed a similar behaviour in the 05358 protocluster, where the emission of the  $^{13}\text{C}$ -isotopologues of  $\text{HC}_3\text{N}$ , and mainly  $\text{DC}_3\text{N}$ , are shifted with respect to the dust continuum (Figure 4).

Thus, all the observational evidence presented in this work suggest that  $\text{DC}_3\text{N}$  might be a good tracer of deuteration fraction in star-forming regions prior the formation of the denser gas, which is an important parameter in chemical models to constrain, for instance, the ortho/para- $\text{H}_2$  ratio (Pagani et al. 2011; Kong et al. 2015; Brünken et al. 2014). New studies of deuteration of other carbon-chain species (e.g.  $\text{HC}_5\text{N}$ ,  $\text{C}_2\text{H}$  or  $c\text{-C}_3\text{H}_2$ ) in large samples of regions like the one used in this work will serve to better understand how the deuteration of this family of species proceeds during the star-formation process.

Finally, in Figure 7 we compare the  $D_{\text{frac}}$  values found in our sample of high-mass cores, divided by evolutionary stage, with those reported in the literature towards other interstellar sources (low-mass dark clouds, low-mass protostars, and previous upper limits in high-mass star-forming regions; see references in Figure 7). We found values in the interval 0.003–0.022 in high-mass cores, which are consistent with the previous upper limits found in Orion KL and SgrB2 N2 hot cores: 0.015 and 0.0009, respectively (Espluques et al. 2013; Belloche et al. 2016). These values of  $D_{\text{frac}}$  are lower than those found in protostellar low mass stars (0.025–0.5), and pre-stellar low-mass dark clouds (0.03–0.13), as shown in Figure 7. This might be due to slightly higher temperatures of high-mass clumps, which would make less efficient the ion-molecule gas-phase reactions responsible of the formation of  $\text{DC}_3\text{N}$ . However, we stress that the comparison between low- and high-mass cores should be




**Figure 6.** Comparison of the FWHM values of  $\text{DC}_3\text{N}$  with those of ortho- $\text{NH}_2(1_{1,1}-1_{0,1})$  (open circles, from Fontani et al. 2015) and  $\text{N}_2\text{D}^+(2-1)$  (filled circles, from Fontani et al. 2011). The different colors correspond to different evolutionary groups. The shaded area indicates the parameter space where the FWHM of  $\text{DC}_3\text{N}$  is larger than those of the other deuterated species.

done with caution, since the observations are tracing very different linear sizescales. As previously mentioned, our observations have likely detected the  $\text{DC}_3\text{N}$  arising from the external envelope of the clumps, rather than the inner cores. Therefore, future interferometric observations with high angular resolution are needed to understand if and how  $\text{DC}_3\text{N}$  is formed at smaller core scales ( $\sim 0.01$  pc).

## 6 CONCLUSIONS

We present the first study of  $\text{DC}_3\text{N}$  toward a sample of massive cores in different evolutionary stages, from pre-stellar to protostellar phases. We detected the  $\text{DC}_3\text{N}$   $J=11-10$  transition towards 15 regions, which include 2 *cold* High Mass Starless Cores (HMSCs), 3 *warm* HMSCs, 4 High Mass Protostellar Objects (HMPOs) and 6 Ultra Compact HII regions. We found values of  $D_{\text{frac}}$  (abundance ratio of  $\text{DC}_3\text{N}$  with respect its  $\text{HC}_3\text{N}$ ) of 0.003–0.022, lower than those found in pre-stellar and protostellar low-mass star-forming regions. We do not find any correlation between  $D_{\text{frac}}$  and the evolutionary stage of the cores, or with the kinetic temperature. The comparison with other deuterated species previously studied toward the same sample indicates a weak correlation of  $D_{\text{frac}}$  in those species formed only or predominantly in gas-phase ( $\text{N}_2\text{H}^+$  and HNC, respectively), and no correlation in species formed only or predominantly on dust grains ( $\text{CH}_3\text{OH}$  and  $\text{NH}_3$ , respectively). We also present the first map of  $\text{DC}_3\text{N}$  in a high-mass star-forming region, the protocluster IRAS 05358+3543. The  $\text{DC}_3\text{N}$  emission is extended ( $\sim 0.37$  pc), and it is shifted with respect to the dust continuum peak where the protostellar activity is on-going. Our observational evidence indicates that  $\text{DC}_3\text{N}$  is likely formed by gas-phase ion-molecule reactions in the outer and less dense part of star-forming clumps, where the interstellar radiation field keeps most of the carbon in atomic form, enhancing the formation of carbon-chain molecules such as  $\text{HC}_3\text{N}$ . Thus,  $\text{DC}_3\text{N}$  might be a good tracer of the deuteration level in star-forming regions prior to the formation of denser gas.



**Figure 7.**  $D_{\text{frac}}$  of HC<sub>3</sub>N in different interstellar sources: low-mass dark clouds from Howe et al. (1994) (orange circles); low-mass protostars from Al-Edhari et al. (2017); Sakai et al. (2009); Araki et al. (2016); Bianchi et al. (2019); Cordiner et al. (2012) (magenta circles); high-mass star-forming regions from this work (different colors denote different evolutionary groups, as indicated above the panel; upper limits are denoted by triangles pointing downwards, while the only lower limit is denoted by the triangle pointing upwards); and previous upper limits (gray triangles) from Esplugues et al. (2013); Watanabe et al. (2015); Belloche et al. (2016).

## 7 ACKNOWLEDGEMENTS

We acknowledge the anonymous reviewer for her/his careful reading of the manuscript and her/his useful comments. V.M.R. has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 664931. LC acknowledges support from the Italian Ministero dell’Istruzione, Università e Ricerca through the grant Progetti Premiali 2012 - iALMA (CUP C52I13000140001). This work is based on observations carried out under projects number 129-12 and 040-19 with the IRAM 30m telescope. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain). This publication has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No 730562 [RadioNet]. L.C. and F.F. thank the IRAM staff for the precious help provided during the observations. We warmly thank Xing "Walker" Lu and Qizhou Zhang for sharing with us their VLA map of the 05358+3543 high-mass protocluster.

## REFERENCES

Aikawa Y., Wakelam V., Hersant F., Garrod R. T., Herbst E., 2012, *ApJ*, **760**, 40  
 Al-Edhari A. J., et al., 2017, *Astron. & Astrophys.*, **597**, A40  
 Araki M., Takano S., Sakai N., Yamamoto S., Oyama T., Kuze N., Tsukiyama K., 2016, *ApJ*, **833**, 291  
 Belloche A., Müller H., Garrod R., Menten K., 2016, *Astron. & Astrophys.*, **587**, A91

Beuther H., Schilke P., Gueth F., McCaughrean M., Andersen M., Sridharan T. K., Menten K. M., 2002, *A&A*, **387**, 931  
 Bianchi E., Ceccarelli C., Codella C., Enrique-Romero J., Favre C., Lefloch B., 2019, *ACS Earth and Space Chemistry*, **3**, 2659  
 Brünken S., et al., 2014, *Nature*, **516**, 219  
 Busquet G., Estalella R., Zhang Q., Viti S., Palau A., Ho P. T. P., Sánchez-Monge Á., 2011, *A&A*, **525**, A141  
 Caselli P., 2002, *Planetary and Space Science*, **50**, 1133  
 Caselli P., Walmsley C. M., Tafalla M., Dore L., Myers P. C., 1999, *ApJ*, **523**, L165  
 Chantzos J., Spezzano S., Caselli P., Chacón-Tanarro A., Bizzocchi L., Sipilä O., Giuliano B. M., 2018, *ApJ*, **863**, 126  
 Charnley S. B., Tielens A. G. G. M., Rodgers S. D., 1997, *ApJ*, **482**, L203  
 Colzi L., Fontani F., Rivilla V. M., Sánchez-Monge A., Testi L., Beltrán M. T., Caselli P., 2018a, *MNRAS*, **478**, 3693  
 Colzi L., Fontani F., Caselli P., Ceccarelli C., Hily-Blant P., Bizzocchi L., 2018b, *A&A*, **609**, A129  
 Colzi L., Fontani F., Caselli P., Leurini S., Bizzocchi L., Quaia G., 2019, *MNRAS*, **485**, 5543  
 Cordiner M. A., Charnley S. B., Wirstrom E. S., Smith R. G., 2012, *ApJ*, **744**, 131  
 Crapsi A., Caselli P., Walmsley C. M., Myers P. C., Tafalla M., Lee C. W., Bourke T. L., 2005, *ApJ*, **619**, 379  
 Creswell R., Winnemiss G., Gerry M., 1977, *Journal of Molecular Spectroscopy*, **65**, 420  
 Di Francesco J., Johnstone D., Kirk H., MacKenzie T., Ledwosinska E., 2008, *Astrophys. J. Suppl. S.*, **175**, 277  
 Emprechtinger M., Caselli P., Volgenau N. H., Stutzki J., Wiedner M. C., 2009, *A&A*, **493**, 89  
 Endres C. P., Schlemmer S., Schilke P., Stutzki J., Müller H. S. P., 2016, *Journal of Molecular Spectroscopy*, **327**, 95

- Esplugues G. B., et al., 2013, *Astron. & Astrophys.*, 559, A51
- Fontani F., et al., 2011, *A&A*, 529, L7
- Fontani F., Sakai T., Furuya K., Sakai N., Aikawa Y., Yamamoto S., 2014, *MNRAS*, 440, 448
- Fontani F., Busquet G., Palau A., Caselli P., Sánchez-Monge Á., Tan J. C., Audard M., 2015, *A&A*, 575, A87
- Heyer M. H., Carpenter J. M., Ladd E. F., 1996, *Astrophys. J.*, 463, 630
- Howe D. A., Millar T. J., Schilke P., Walmsley C. M., 1994, *MNRAS*, 267, 59
- Kendall M. G., 1938, *Biometrika*, 30, 81
- Klein B., Hochgürtel S., Krämer I., Bell A., Meyer K., Güsten R., 2012, *Astronomy and Astrophysics*, 542, L3
- Kong S., Caselli P., Tan J. C., Wakelam V., Sipilä O., 2015, *ApJ*, 804, 98
- Langer W., Schloerb F., Snell R., Young J., 1980, *Astrophys. J.*, 239, L125
- Leurini S., Beuther H., Schilke P., Wyrowski F., Zhang Q., Menten K. M., 2007, *A&A*, 475, 925
- Li J., et al., 2016, *ApJ*, 824, 136
- Linsky J. L., et al., 2006, *ApJ*, 647, 1106
- Lu X., Zhang Q., Liu H. B., Wang J., Gu Q., 2014, *ApJ*, 790, 84
- Martín S., Martín-Pintado J., Blanco-Sánchez C., Rivilla V. M., Rodríguez-Franco A., Rico-Villas F., 2019, *A&A*, 631, A159
- Milam S. N., Savage C., Brewster M. A., Ziurys L. M., Wyckoff S., 2005, *ApJ*, 634, 1126
- Millar T. J., Bennett A., Herbst E., 1989, *ApJ*, 340, 906
- Mininni C., Fontani F., Rivilla V. M., Beltrán M. T., Caselli P., Vasyunin A., 2018, *MNRAS*, 476, L39
- Müller H. S. P., Thorwirth S., Roth D. A., Winnewisser G., 2001, *A&A*, 370, L49
- Müller H. S. P., Schlöder F., Stutzki J., Winnewisser G., 2005, *Journal of Molecular Structure*, 742, 215
- Oliveira C. M., Hébrard G., Howk J. C., Kruk J. W., Chayer P., Moos H. W., 2003, *ApJ*, 587, 235
- Osamura Y., Roberts H., Herbst E., 2004, *A&A*, 421, 1101
- Pagani L., Salez M., Wannier P. G., 1992, *A&A*, 258, 479
- Pagani L., Roueff E., Lesaffre P., 2011, *ApJ*, 739, L35
- Sakai N., Sakai T., Hirota T., Yamamoto S., 2009, *ApJ*, 702, 1025
- Sánchez-Monge Á., Pandian J. D., Kurtz S., 2011, *ApJ*, 739, L9
- Spezzano S., Caselli P., Bizzocchi L., Giuliano B. M., Lattanzi V., 2017, *A&A*, 606, A82
- Takano S., et al., 1998, *A&A*, 329, 1156
- Taniguchi K., Saito M., Ozeki H., 2016, *ApJ*, 830, 106
- Taniguchi K., Ozeki H., Saito M., 2017, *ApJ*, 846, 46
- Thorwirth S., Müller H. S. P., Winnewisser G., 2001, *Phys. Chem. Chem. Phys.*, 3, 1236
- Walmsley C. M., Flower D. R., Pineau des Forêts G., 2004, *A&A*, 418, 1035
- Watanabe Y., et al., 2015, *Astrophys. J.*, 809, 162
- Zhang Q., Hunter T. R., Sridharan T. K., Ho P. T. P., 2002, *ApJ*, 566, 982

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.