



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Disability accrual in primary and secondary progressive multiple sclerosis

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Harding-Forrester S, R.I., MSBase investigators (2023). Disability accrual in primary and secondary progressive multiple sclerosis. JOURNAL OF NEUROLOGY, NEUROSURGERY AND PSYCHIATRY, 94(9), 707-717 [10.1136/jnnp-2022-330726].

Availability:

This version is available at: <https://hdl.handle.net/11585/935060> since: 2024-02-10

Published:

DOI: <http://doi.org/10.1136/jnnp-2022-330726>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Harding-Forrester S, Roos I, Nguyen AL, Malpas CB, Diouf I, Moradi N, Sharmin S, Izquierdo G, Eichau S, Patti F, Horakova D, Kubala Havrdova E, Prat A, Girard M, Duquette P, Grand'Maison F, Onofri M, Lugaresi A, Grammond P, Ozakbas S, Amato MP, Gerlach O, Sola P, Ferraro D, Buzzard K, Skibina O, Lechner-Scott J, Alroughani R, Boz C, Van Pesch V, Cartechini E, Terzi M, Maimone D, Ramo-Tello C, Yamout B, Khoury SJ, La Spitaleri D, Sa MJ, Blanco Y, Granella F, Slee M, Butler E, Sidhom Y, Gouider R, Bergamaschi R, Karabudak R, Ampapa R, Sánchez-Menoyo JL, Prevost J, Castillo-Trivino T, McCombe PA, Macdonell R, Laureys G, Van Hijfte L, Oh J, Altintas A, de Gans K, Turkoglu R, van der Walt A, Butzkueven H, Vucic S, Barnett M, Cristiano E, Hodgkinson S, Iuliano G, Kappos L, Kuhle J, Shaygannejad V, Soysal A, Weinstock-Guttman B, Van Wijmeersch B, Kalincik T; MSBase investigators.

Disability accrual in primary and secondary progressive multiple sclerosis.

J Neurol Neurosurg Psychiatry. 2023 Sep;94(9):707-717.

The final published version is available online at: [10.1136/jnnp-2022-330726](https://doi.org/10.1136/jnnp-2022-330726)

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Disability accrual in primary and secondary progressive multiple sclerosis

Sam Harding-Forrester, MD, MPH; Izanne Roos, MBBS, PhD; Ai-Lan Nguyen, MBBS, PhD; Charles B. Malpas, PhD; Ibrahima Diouf, PhD; Nahid Moradi, MD; Sifat Sharmin, PhD; Guillermo Izquierdo, MD, PhD; Sara Eichau, MD, PhD; Francesco Patti, MD; Dana Horáková, MD, PhD; Eva Kubala Havrdová, MD, PhD; Alexandre Prat, MD, PhD; Marc Girard, MD; Pierre Duquette, MD; François Grand'Maison, MD; Marco Onofri, MD; Alessandra Lugaresi, MD, PhD; Pierre Grammond, MD; Serkan Özakbaş, MD; Maria Pia Amato, MD; Oliver Gerlach, MD, PhD; Patrizia Sola, MD, PhD; Diana Ferraro, MD, PhD; Katherine Buzzard, MBBS, PhD; Olga Skibina, MBBS, PhD; Jeannette Lechner-Scott, MD, PhD; Raed Alroughani, MD; Cavit Boz, MD; Vincent van Pesch, MD, PhD; Elisabetta Cartechini, MD; Murat Terzi, MD; Davide Maimone, MD, PhD; Cristina Ramo-Tello, MD; Bassem Yamout, MD; Samia J. Khoury, MD; Daniele L.A. Spitaleri, MD; Maria José Sá, MD, PhD; Yolanda Blanco, MD; Franco Granella, MD; Mark Sleen, MBBS, PhD; Ernest Butler, MBBS, PhD; Youssef Sidhom, MD; Riadh Gouider, MD; Roberto Bergamaschi, MD; Rana Karabudak, MD; Radek Ampapa, MD; José Luis Sánchez-Menoyo, MD; Julie Prévost, MD; Tamara Castillo-Triviño, MD; Pamela McCombe, MBBS, PhD; Richard Macdonell, MD; Guy Laureys, MD, PhD; Liesbeth Van Hijfte; Jiwon Oh, MD, PhD; Ayse Altintas, MD; Koen de Gans, MD; Recai Türkoğlu, MD; Anneke van der Walt, MBBS, PhD; Helmut Butzkueven, MBBS, PhD; Steve Vucic, MBBS, PhD; Michael Barnett, MBBS, PhD; Edgardo Cristiano, MD; Suzanne Hodgkinson, MBBS, PhD; Gerardo Iuliano, MD; Ludwig Kappos, MD; Jens Kuhle, MD, PhD; Vahid Shaygannejad, MD; Aysun Soysal, MD; Bianca Weinstock-Guttman, MD; Bart Van Wijmeersch, MD, PhD; and Tomas Kalincik, MD, PhD, on behalf of the MSBase investigators.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Tomas Kalincik, MD, PhD

Clinical Outcomes Research Unit (CORE)

Royal Melbourne Hospital, 300 Grattan St, Parkville VIC 3050, Australia

E-mail: tomas.kalincik@unimelb.edu.au

Telephone: +61 393 424 404

Word count: Abstract, 256; Main text, 3544

Display items: Tables, 4; Figures, 4

Keywords: multiple sclerosis, chronic progressive; disease progression; prognosis; observational study; MSBase

Glossary:

MS = multiple sclerosis;

RRMS = relapsing-remitting MS;

SPMS = secondary progressive MS;

SPMS-N = SPMS with no superimposed relapse activity;

SPMS-A = SPMS with superimposed relapse activity;

PPMS = primary progressive MS;

PPMS-N = PPMS with no superimposed relapse activity;

PPMS-A = PPMS with superimposed relapse activity;

EDSS = Expanded Disability Status Scale;

DMT = disease-modifying therapy;

IST = immunosuppressant therapy;

ASCT = autologous stem-cell transplant;

CI = confidence interval;

HR = hazard ratio.

Abstract

Background. Some studies comparing primary and secondary progressive multiple sclerosis (PPMS, SPMS) report similar ages at onset of the progressive phase, and similar rates of subsequent disability accrual. Others report later onset and/or faster accrual in SPMS. Comparisons have been complicated by regional cohort effects, phenotypic differences in sex ratio and management, and variable diagnostic criteria for SPMS.

Methods. We compared disability accrual in PPMS and operationally-diagnosed SPMS in the international, clinic-based MSBase cohort. Inclusion required PPMS or SPMS with onset at age ≥ 18 years since 1995. We estimated Andersen-Gill hazard ratios for disability accrual on the Expanded Disability Status Scale (EDSS), adjusted for sex, age, baseline disability, EDSS score frequency, and drug therapies, with center and patient as random effects. We also estimated ages at onset of the progressive phase (Kaplan-Meier) and at EDSS milestones (Turnbull). Analyses were replicated with physician-diagnosed SPMS.

Results. Included patients comprised 1872 with PPMS (47% male; 50% with activity) and 2575 with SPMS (32% male; 40% with activity). Relative to PPMS, SPMS had older age at onset of the progressive phase (median 46.7 years [95% confidence interval 46.2–47.3] versus 43.9 [43.3–44.4]; $P < .001$), greater baseline disability, slower disability accrual (hazard ratio 0.86 [0.78–0.94]; $P < .001$), and similar age at wheelchair dependence.

Conclusions. We demonstrate later onset of the progressive phase and slower disability accrual in SPMS versus PPMS. This may balance greater baseline disability in SPMS, yielding convergent disability trajectories across phenotypes. The different rates of disability accrual should be considered before amalgamating PPMS and SPMS in clinical trials.

Key messages

What is already known on this topic

Some studies comparing PPMS and SPMS report similar ages at onset of the progressive phase and similar rates of disability accrual, while others report later onset and/or faster accrual in SPMS. Comparisons have been complicated by regional cohort effects, phenotypic differences in sex ratio and management, and variable diagnostic criteria for SPMS.

What this study adds

We compared disability accrual in PPMS and SPMS in the international MSBase cohort, using multivariable survival models, with SPMS diagnosed operationally. Relative to PPMS, patients with SPMS have greater baseline disability at onset of the progressive phase; however, we show that patients with SPMS enter their progressive phase at older ages and experience slower disability accrual thereafter. This may yield similar ages at wheelchair dependence across phenotypes.

How this study might affect research, practice, or policy

Our results indicate that disability accrual is slower in SPMS than in PPMS. Caution is warranted about combining the two phenotypes in clinical trials, even as their long-term prognosis may be similar.

Introduction

Over 80% of patients with multiple sclerosis (MS) first present with relapsing-remitting MS (RRMS), in which episodic relapses produce varying degrees of sustained disability. Most convert, after a median 15–20 years, to secondary progressive MS (SPMS),^{1–3} defined by continuous disability accrual (the “progressive phase”) with or without superimposed relapses (“activity”).⁴ In contrast, ~15% of patients with MS have primary progressive MS (PPMS), with the progressive phase apparent from clinical onset.

Several geographically determined cohort studies have compared disability trajectories in PPMS and SPMS, using nonparametric estimates of time to milestones on the Expanded Disability Status Scale (EDSS).⁵ Most concluded that regardless of phenotype, the progressive phase had median onset during the fifth decade of life, and stereotyped disability accrual above EDSS 4.^{6–12} Many authors (not all)¹³ have therefore argued that although SPMS begins with baseline disability established during the relapsing-remitting phase, PPMS and SPMS ultimately converge on a partially age-dependent disability trajectory.^{2,7–12,14,15}

However, whereas several studies found that PPMS and SPMS had similar ages at onset of the progressive phase,^{2,9,10,12} and similar rates of disability accrual thereafter,^{9,12} others found SPMS had later onset^{16,17} or faster disability accrual.^{8,18} The possibility that SPMS displays both greater baseline disability and faster ongoing disability accrual conflicts with a model in which long-term outcomes converge across phenotypes.

Past comparisons of PPMS and SPMS have been complicated by regional cohort effects; phenotypic differences in sex ratio¹⁹ and clinical management; and poorly standardized diagnosis of SPMS.^{20,21} Here, we compare disability accrual among patients with PPMS and SPMS in the international MSBase cohort, using multivariable survival models, and applying operationalized diagnostic criteria for SPMS.²² We also estimate ages at onset of the progressive phase and at EDSS milestones in each phenotype; assess mean disability trajectories; and compare subgroups of each phenotype with and without activity.

Methods

Participants

Patient records were extracted from MSBase on January 7, 2020. MSBase is an international, clinic-based multiple sclerosis registry,²³ approved by the Melbourne Health Human Research Ethics Committee (2006.044) and includes records entered prospectively since July 1, 2004, in addition to retrospectively added records. Participants gave written informed consent as per the MSBase and local regulations. Data quality and generalizability procedures were applied before inclusion screening (**supplementary methods 1.1**).

Patients with PPMS were identified by physician diagnosis, with onset defined as the date of first symptoms. Patients with SPMS were identified operationally (among patients with initial RRMS), using the previously validated Lorscheider criteria.²² These require an EDSS increase of at least 1.5, 1.0, or 0.5 point(s) from baseline(s) 0, 1.0–5.5, and ≥ 6 , respectively; confirmation over ≥ 90 days of the EDSS increase and leading functional system score; and a minimum EDSS score of 4 and pyramidal functional system score of 2 at onset. Only EDSS scores > 30 days after the onset of any preceding relapse (“outside relapse”) are used to identify and confirm EDSS increases.

Study eligibility required PPMS or SPMS with onset at age ≥ 18 years since January 1, 1995, and at least three EDSS scores⁵—including a final “confirmatory score” ≥ 180 days after the second and outside relapse. For patients with SPMS, eligibility further required initial records during RRMS, including an EDSS score ≤ 3 ; this ensured diagnosis at the earliest qualifying date under the Lorscheider criteria. Eligible patients were generally included from the date of their first EDSS score in the progressive phase; if this score occurred during relapse and the first subsequent score outside relapse was lower, the patient was included from the date of the latter. Patients were censored on the date of their final score ≥ 180 days prior to the final confirmatory score.

Patients were considered to have superimposed activity (PPMS-A, SPMS-A) if any relapse was documented during the progressive phase (prospectively or retrospectively, including any diagnosis of “progressive-relapsing MS”); otherwise, patients were considered to have no relapse activity (PPMS-N, SPMS-N). Relapses are defined in MSBase protocols by new or exacerbated symptoms persisting ≥ 24 hours, absent concurrent illness, and beginning > 30 days after onset of any prior relapse.

Outcomes

The primary outcome was disability accrual during the progressive phase. Disability accrual events were defined by an EDSS score increase of at least 1.5, 1.0, or 0.5 point(s) from baseline(s) 0, 1.0–

1
2
3 5.5, and ≥ 6 respectively,²⁴ with the increase confirmed ≥ 180 days later outside relapse, and
4 sustained throughout follow-up. Each event established a new EDSS baseline, defined as the lowest
5 score on or after that date. Periods between baselines were termed “epochs”.

6
7
8 The secondary outcome was age at confirmed EDSS ≥ 7 (wheelchair dependence), defined by the
9 first EDSS score ≥ 7 with no subsequent score < 7 and a confirmatory score ≥ 7 recorded ≥ 180
10 days later outside relapse. For patients observed from a confirmed score ≥ 7 , EDSS ≥ 7 was
11 considered attained in the interval between MS onset (inclusive) and the first available EDSS score
12 (exclusive).
13

14
15
16 Population mean EDSS scores by age, in two-year intervals, were calculated for each phenotype
17 (“mean EDSS trajectories”). For these calculations, if a patient had multiple scores within a two-
18 year interval, the median was taken as the patient’s score for that interval.
19
20

21 *Statistical analysis*

22
23
24 Disability accrual in PPMS and SPMS was visualized using Nelson-Aalen cumulative hazard
25 functions, and compared formally using Andersen-Gill adjusted hazard ratios (HR).
26
27

28
29 All Andersen-Gill models included random-effects terms for treating center and for patient identity
30 (nested within treating center). The initial model, estimating the total effect of phenotype (PPMS
31 or SPMS), included terms for sex and for age at the start of each epoch. The “complete” model
32 (**supplementary methods 1.2–1.3**) added terms for EDSS score at the start of each epoch; the
33 proportion of each epoch receiving disease-modifying therapy (DMT) and immunosuppressant
34 therapy (agents listed in **table 1**); and the annualized frequency of EDSS scores during each epoch
35 (“EDSS score frequency”). Next, a phenotype–activity interaction was added; based on the result,
36 the “complete” model was re-assessed with phenotype comprising PPMS-N, PPMS-A, SPMS-N,
37 and SPMS-A. Finally, to assess whether hazard ratios for DMT differed between phenotypes,
38 models were constructed separately for PPMS and SPMS, with or without an activity–DMT
39 interaction. Ties were handled using the Efron approximation. The proportional hazards
40 assumption was assessed using Schoenfeld residuals (visual and formal evaluation).
41
42
43
44
45

46
47 For each phenotype, age-based survival functions for onset of the progressive phase were estimated
48 using the Kaplan-Meier estimator, and compared using the log-rank test. Age-based survival
49 functions for EDSS ≥ 7 were estimated using the Turnbull estimator (to accommodate interval-
50 censored observations), and compared using a generalized log-rank test.²⁵ Median ages at onset of
51 the progressive phase and at confirmed EDSS ≥ 4 , ≥ 6 , and ≥ 7 were likewise estimated using the
52 Turnbull estimator, for the full dataset (1995–2020; all patients) and for three constituent time
53 periods (1995–2003, 2004–2011, 2012–2020); analyses for each period included only patients
54
55
56
57
58
59
60

1
2
3 diagnosed with PPMS or SPMS within the period, and only EDSS scores recorded before the end
4 of the period.
5

6
7 Five sensitivity analyses were performed. The first required onset of PPMS or SPMS since July 1,
8 2004 (the start of prospective data collection in MSBase). The second examined only the longest
9 period (if any) in each patient's progressive phase during which at least one EDSS score was
10 recorded every 15 months (457 days). The third defined disability accrual and EDSS ≥ 7 to require
11 confirmation over ≥ 365 days, reducing the possibility that "confirmed" EDSS increases might
12 subsequently reverse.^{24,26} The fourth obtained period-specific hazard ratios 0–5, 5–10, and 10–15
13 years from onset of the progressive phase. The fifth assessed disability accrual from EDSS 4–5
14 rather than from onset of the progressive phase (and, accordingly, restricted analysis to EDSS
15 scores ≥ 4 , and to patients with an initial score in the progressive phase ≤ 5).
16
17
18
19

20
21 Finally, all analyses were repeated comparing PPMS with SPMS identified by physician diagnosis
22 (since January 1, 1995), rather than operationalized criteria.
23

24
25 To assess whether the effect of phenotype (PPMS or SPMS) on disability accrual was mediated by
26 DMT exposure and EDSS score frequency, a mediation analysis was performed using a broadly
27 applicable natural effects estimation procedure (**supplementary methods 1.4**).^{27,28} Mediation
28 analysis assumes control for exposure–outcome, mediator–outcome, and exposure–mediator
29 confounding, and the absence of mediator–outcome confounders affected by the exposure.
30
31

32
33 Analyses were performed in R 4.0.0 (packages 'survival' 3.1-12, 'coxme' 2.2-16, 'lme4' 1.1-26,
34 'glrt' 2.0, 'survminer' 0.4.8).
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Main analyses

1872 patients with PPMS (47% male; 50% PPMS-A) and 2575 with operationally diagnosed SPMS (32% male; 40% SPMS-A) were included in the main analyses, drawn from 107 centers in 33 countries (**figure 1**; **supplementary figure 1.1**; **supplementary table 1.1**). Among patients with operationally diagnosed SPMS, 1134 (44%) were physician-diagnosed by the end of follow-up. Patient characteristics are summarized in **table 1** (for relapse characteristics, see **supplementary table 1.2**).

Table 1. Clinical characteristics of included patients

Patient characteristics	PPMS (all)	PPMS-N	PPMS-A	SPMS (all)	SPMS-N	SPMS-A
Patients	1872	935	937	2575	1541	1034
Sex, male (%)	878 (47)	442 (47)	436 (47)	836 (32)	513 (33)	323 (31)
Age, MS onset	43.5 ± 10.5	45.6 ± 9.8	41.5 ± 10.7	32.3 ± 10.2	33.2 ± 10.5	31.0 ± 9.6
Age, progressive phase	43.5 ± 10.5	45.6 ± 9.8	41.5 ± 10.7	47.0 ± 10.1	48.7 ± 10.3	44.5 ± 9.4
Age, inclusion	48.7 ± 10.8	50.7 ± 10.1	46.7 ± 11.2	47.0 ± 10.1	48.7 ± 10.3	44.5 ± 9.4
Time, MS onset to inclusion; years	4.1 [2.0–7.3]	4.1 [2.1–7.3]	4.1 [1.9–7.3]	13.5 [8.2–19.7]	14.3 [8.8–20.8]	12.4 [7.3–18.2]
Time, follow-up; years	4.2 [1.9–7.9]	3.8 [1.7–7.5]	4.4 [2.1–8.2]	3.7 [1.6–6.8]	2.7 [1.2–5.4]	5.3 [2.8–8.6]
EDSS score frequency, annualized	1.53 [1.00–2.28]	1.37 [0.92–2.10]	1.68 [1.12–2.44]	1.79 [1.20–2.56]	1.69 [1.09–2.30]	1.90 [1.38–2.94]
Disability at inclusion; EDSS	4.0 [3.0–6.0]	4.0 [3.0–6.0]	4.0 [3.0–6.0]	4.5 [4.0–6.0]	4.5 [4.0–6.0]	4.5 [4.0–5.5]
Disability at censoring; EDSS	6.0 [4.5–6.5]	6.0 [4.5–6.5]	6.0 [4.0–6.5]	6.0 [4.5–6.5]	5.5 [4.0–6.5]	6.0 [5.0–6.5]
Disability increase, annualized; EDSS	0.15 [0.00–0.38]	0.15 [0.00–0.37]	0.15 [0.00–0.39]	0.00 [0.00–0.22]	0.00 [0.00–0.22]	0.06 [0.00–0.22]
Deaths recorded, from any cause (%)	59 (3)	33 (4)	26 (3)	71 (3)	45 (3)	26 (3)
Relapses during follow-up						
Patients (%)	336 (18)	0 (0)	336 (36)	948 (37)	0 (0)	948 (92)
Annualized relapse rate	0.26 [0.14–0.48]	NA	0.26 [0.14–0.48]	0.34 [0.19–0.60]	NA	0.34 [0.19–0.60]
Cerebrospinal fluid oligoclonal bands; patients (%)						
Present	1149 (61)	546 (58)	603 (64)	1496 (58)	833 (54)	663 (64)
Absent	89 (5)	46 (5)	43 (5)	86 (3)	52 (3)	34 (3)
Not assessed	634 (34)	343 (37)	291 (31)	993 (39)	656 (43)	337 (33)
Disease-modifying therapy, proportion of follow-up receiving treatment; patients (%)						
0%	1104 (59)	652 (70)	452 (48)	541 (21)	395 (26)	146 (14)
> 0–25%	221 (12)	105 (11)	116 (12)	204 (8)	111 (7)	93 (9)
> 25–50%	144 (8)	46 (5)	98 (10)	184 (7)	87 (6)	97 (9)
> 50–75%	106 (6)	36 (4)	70 (7)	185 (7)	80 (5)	105 (10)
> 75%	297 (16)	96 (10)	201 (21)	1461 (57)	868 (56)	593 (57)
Disease-modifying therapy, exposure to specific agents during follow-up; patients (%)						
Interferon beta	318 (17)	85 (9)	233 (25)	1014 (39)	517 (34)	497 (48)
Glatiramer acetate	154 (8)	46 (5)	108 (12)	412 (16)	194 (13)	218 (21)
Fingolimod	118 (6)	39 (4)	79 (8)	510 (20)	225 (15)	285 (28)
Teriflunomide	28 (1)	7 (1)	21 (2)	141 (5)	78 (5)	63 (6)

Dimethyl fumarate	24 (1)	8 (1)	16 (2)	166 (6)	89 (6)	77 (7)
Cladribine	2 (0)	0 (0)	2 (0)	13 (1)	6 (0)	7 (1)
Mitoxantrone	91 (5)	29 (3)	62 (7)	129 (5)	57 (4)	72 (7)
Natalizumab	83 (4)	27 (3)	56 (6)	443 (17)	216 (14)	227 (22)
Alemtuzumab	12 (1)	3 (0)	9 (1)	55 (2)	24 (2)	31 (3)
Daclizumab	1 (0)	0 (0)	1 (0)	3 (0)	2 (0)	1 (0)
Rituximab	62 (3)	16 (2)	46 (5)	52 (2)	28 (2)	24 (2)
Ocrelizumab	169 (9)	90 (10)	79 (8)	78 (3)	51 (3)	27 (3)
Siponimod	0 (0)	0 (0)	0 (0)	2 (0)	2 (0)	0 (0)
ASCT	4 (0)	0 (0)	4 (0)	9 (0)	4 (0)	5 (0)
Immunosuppressant therapy, exposure to specific agents during follow-up; patients (%)						
Azathioprine	233 (12)	61 (7)	172 (18)	231 (9)	94 (6)	137 (13)
Methotrexate	121 (6)	41 (4)	80 (9)	105 (4)	36 (2)	69 (7)
Cyclophosphamide	94 (5)	26 (3)	68 (7)	90 (3)	31 (2)	59 (6)
Mycophenolate mofetil	5 (0)	2 (0)	3 (0)	1 (0)	1 (0)	0 (0)
Pregnancy during follow-up						
Patients (%)	15 (1)	2 (0)	13 (1)	32 (1)	13 (1)	19 (2)

Table 1. Clinical characteristics of included patients (continued)

Values are number (%), mean \pm standard deviation, or median [interquartile range]. Patients classified as having PPMS-A include all eligible patients with a recorded diagnosis of progressive-relapsing MS (n = 540 patients; 58%). Patients classified as having SPMS under the operationalized diagnostic criteria include 1134 patients (44%) with SPMS under physician diagnosis. Annualized relapse rate is calculated for the subset of patients with one or more relapses during follow-up.

Abbreviations: EDSS = Expanded Disability Status Scale; ASCT = autologous stem-cell transplant, assumed effective for 5 years.

The Kaplan-Meier median age at onset of the progressive phase was younger in PPMS (43.9; 95% confidence interval [CI] = 43.3–44.4) versus SPMS (46.7; 95% CI = 46.2–47.3) ($P < .001$), and in the subgroup of each phenotype with activity (PPMS-A, SPMS-A) versus those without ($P < .001$ in both PPMS and SPMS) (table 2).

Table 2. Ages at onset of the progressive phase and at confirmed EDSS ≥ 7

Phenotype	All patients	Female	Male
Age at onset of the progressive phase (median), years (95% CI)			
PPMS	43.9 (43.3–44.4)	44.4 (43.8–45.2)	43.0 (42.3–44.0)
PPMS-N	46.0 (45.2–46.8)	46.5 (45.3–47.7)	45.5 (44.5–46.7)
PPMS-A	41.9 (40.7–42.6)	42.8 (42.0–43.7)	40.0 (38.6–41.8)

SPMS	46.7 (46.2–47.3)	47.3 (46.6–47.8)	45.6 (44.9–46.7)
SPMS-N	48.3 (47.7–49.0)	49.3 (48.3–50.2)	46.7 (45.6–48.0)
SPMS-A	44.1 (43.1–44.8)	44.2 (43.2–45.0)	43.0 (42.1–45.5)
Age at confirmed EDSS \geq 7 (25th percentile), years (95% CI)			
PPMS	60.3 (58.8–62.5)	63.4 (60.5–66.2)	58.1 (56.6–60.1)
PPMS-N	62.5 (60.1–65.8)	66.0 (62.5–68.7)	60.1 (57.3–62.7)
PPMS-A	57.6 (55.4–60.9)	60.9 (56.3–65.3)	55.8 (52.8–58.6)
SPMS	62.2 (60.2–64.0)	63.2 (62.1–66.0)	58.3 (56.5–62.3)
SPMS-N	63.6 (62.1–66.9)	63.6 (62.1–67.1)	64.7 (60.4–70.8)
SPMS-A	58.2 (57.0–62.1)	63.2 (60.7–70.0)	55.4 (50.8–57.8)

Ages at onset of the progressive phase (Kaplan-Meier estimator; log-rank test) were younger in PPMS versus SPMS ($P < .001$), in subgroups with activity (-A) versus those without (-N) ($P < .001$ in both PPMS and SPMS), and in males versus females ($P = .22$ in PPMS, $P = .01$ in SPMS). Ages at confirmed EDSS \geq 7 (Turnbull estimator; generalized log-rank test) were similar in SPMS and PPMS ($P = .06$; among females, $P = .45$; among males, $P = .44$; among patients with activity, $P = .22$; without activity, $P = .38$), but younger in patients with activity than in those without (PPMS-A vs. PPMS-N, $P = .002$; SPMS-A vs. SPMS-N, $P = .007$).

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.

The hazard of disability accrual was lower in SPMS versus PPMS—whether based on unadjusted Nelson-Aalen cumulative hazards (**figure 2A**); the Andersen-Gill model adjusted for sex and age (HR = 0.75; 95% CI = 0.70–0.81; $P < .001$; **supplementary table 1.3**); or the complete Andersen-Gill model further adjusted for baseline EDSS score, DMT and immunosuppressant therapy, and EDSS score frequency (HR = 0.86; 95% CI = 0.78–0.94; $P < .001$; **table 3**). The proportional hazards assumption was violated for EDSS score frequency; correction did not alter findings (**supplementary table 1.4**).

Table 3. Andersen-Gill models for hazard of confirmed disability accrual

Variable	Hazard ratio (95% CI)	<i>P</i>
Model comparing SPMS to PPMS (reference)		
Phenotype, SPMS	0.86 (0.78–0.94)	< .001

Sex, male	1.17 (1.09–1.25)	< .001
Age, at start of epoch	1.00 (0.99–1.00)	.28
EDSS baseline, at start of epoch	0.92 (0.90–0.95)	< .001
DMT, % of epoch on treatment (25% increments)	0.97 (0.94–0.99)	.004
IST, % of epoch on treatment (25% increments)	0.96 (0.92–0.99)	.01
EDSS score frequency, annualized, during epoch	1.14 (1.13–1.16)	< .001
Model comparing PPMS-A, SPMS-N, and SPMS-A to PPMS-N (reference)		
Phenotype		
PPMS-A	0.98 (0.89–1.09)	.75
SPMS-N	0.94 (0.84–1.06)	.34
SPMS-A	0.78 (0.69–0.88)	< .001
Sex, male	1.17 (1.09–1.25)	< .001
Age, at start of epoch	1.00 (0.99–1.00)	.13
EDSS baseline, at start of epoch	0.92 (0.90–0.94)	< .001
DMT, % of epoch on treatment (25% increments)	0.97 (0.95–0.99)	.006
IST, % of epoch on treatment (25% increments)	0.96 (0.92–0.99)	.01
EDSS score frequency, annualized, during epoch	1.15 (1.13–1.16)	< .001

Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale; DMT = disease-modifying therapy; IST = immunosuppressant therapy.

In the Andersen-Gill model adding a phenotype–activity interaction, disability accrual was not associated with phenotype (SPMS versus PPMS; HR = 0.94; 95% CI = 0.84–1.06; $P = .34$) or activity (present versus absent; HR = 0.98; 95% CI = 0.89–1.09; $P = .75$), but an interaction was observed between SPMS and activity (HR = 0.84; 95% CI = 0.73–0.96; $P = .01$). We therefore modified the complete model to estimate hazards of disability accrual in activity subgroups. Relative to PPMS-N, hazard was comparable in PPMS-A and SPMS-N, but lower in SPMS-A (HR

= 0.78; 95% CI = 0.69–0.88; $P < .001$; **table 3**; cf. unadjusted comparison, **figure 2B**). Analyses stratified by phenotype yielded no evidence of interactions between DMT and activity (PPMS, $P = .60$; SPMS, $P = .13$; **supplementary tables 1.5–1.6**).

Turnbull estimates for ages at EDSS ≥ 7 were similar in PPMS (25th percentile = 60.3 years; 95% CI = 58.8–62.5) and SPMS (25th percentile = 62.2 years; 95% CI = 60.2–64.0) ($P = .06$; among females, $P = .45$; males, $P = .44$). However, ages at EDSS ≥ 7 were younger in patients with relapse activity versus those without (PPMS-A versus PPMS-N, $P = .002$; SPMS-A versus SPMS-N, $P = .007$; **figure 3**), consistent with the earlier onset of the progressive phase in those with activity. Turnbull estimates for median ages demonstrated that whereas EDSS ≥ 4 was reached at younger ages in SPMS versus PPMS (and often prior to SPMS onset, as expected), ages at EDSS ≥ 6 and ≥ 7 became more similar across phenotypes (**table 4**). Therefore, the time between EDSS milestones was longer in SPMS (particularly SPMS-A), reflecting slower disability accrual in this phenotype versus PPMS. Across phenotypes, ages at onset of the progressive phase and at each EDSS milestone were older in later time periods.

Table 4. Ages at onset of the progressive phase and at confirmed EDSS scores, by time period

	Median age at progressive multiple sclerosis milestones, years (95% CI)			
	Onset	EDSS ≥ 4	EDSS ≥ 6	EDSS ≥ 7
Complete dataset (1995–2020)				
PPMS ($n = 1872$)	43.9 (43.3–44.4)	50.4 (49.6–51.1)	56.9 (55.8–57.4)	72.8 (70.8–76.4)
PPMS-N ($n = 935$)	46.0 (45.2–46.8)	51.9 (50.8–53.2)	57.5 (56.9–59.0)	75.4 (71.0–NA)
PPMS-A ($n = 937$)	41.9 (40.7–42.6)	47.7 (46.2–49.6)	54.8 (53.6–56.6)	70.8 (68.9–NA)
SPMS ($n = 2575$)	46.7 (46.2–47.3)	46.1 (45.5–46.6)	55.4 (54.6–56.5)	76.0 (74.9–NA)
SPMS-N ($n = 1541$)	48.3 (47.7–49.0)	47.6 (47.1–48.3)	57.8 (56.3–58.9)	76.0 (75.4–NA)
SPMS-A ($n = 1034$)	44.1 (43.1–44.8)	43.5 (42.9–44.3)	53.0 (52.0–54.1)	76.1 (70.0–NA)
1995–2003				
PPMS ($n = 125$)	41.9 (39.3–44.9)	47.0 (45.8–50.9)	51.9 (50.0–55.2)	60.6 (56.0–65.0)
PPMS-N ($n = 56$)	44.5 (41.5–48.7)	50.7 (47.9–53.5)	54.3 (51.0–57.5)	62.2 (56.0–68.5)
PPMS-A ($n = 69$)	41.0 (37.5–43.7)	44.4 (43.3–50.8)	47.6 (46.2–54.5)	58.8 (55.0–65.4)
SPMS ($n = 121$)	43.0 (41.4–46.3)	42.9 (40.5–44.9)	47.4 (45.6–50.4)	57.4 (55.7–63.8)
SPMS-N ($n = 48$)	45.5 (42.6–51.5)	44.5 (42.0–51.5)	48.8 (44.5–52.8)	57.4 (53.1–72.6)
SPMS-A ($n = 73$)	42.1 (39.8–44.9)	41.8 (39.8–44.5)	47.1 (45.1–51.5)	58.0 (56.3–NA)
2004–2011				
PPMS ($n = 266$)	44.4 (43.3–45.9)	49.3 (46.6–50.8)	53.4 (52.2–56.9)	65.1 (62.4–NA)
PPMS-N ($n = 132$)	46.5 (44.5–48.3)	50.5 (48.3–52.5)	54.8 (52.4–57.3)	67.5 (60.0–NA)
PPMS-A ($n = 134$)	42.8 (40.6–44.5)	46.4 (45.1–49.7)	53.0 (50.2–59.0)	65.1 (62.4–NA)

SPMS (<i>n</i> = 687)	45.2 (44.2–46.0)	44.7 (43.8–45.6)	51.6 (49.7–52.5)	65.0 (62.1–68.4)
SPMS-N (<i>n</i> = 329)	46.6 (45.7–48.2)	46.3 (45.3–47.6)	52.3 (50.6–54.6)	67.4 (63.6–71.4)
SPMS-A (<i>n</i> = 358)	43.3 (42.3–44.8)	43.0 (42.0–44.4)	49.8 (48.3–52.2)	61.3 (58.5–68.9)
2012–2020 (ending January 7, 2020)				
PPMS (<i>n</i> = 250)	47.6 (45.9–49.8)	54.2 (52.6–59.1)	63.3 (61.1–70.6)	80.8 (NA–NA)
PPMS-N (<i>n</i> = 129)	48.3 (47.2–51.5)	57.8 (52.6–60.8)	64.8 (60.8–NA)	NA (71.9–NA)
PPMS-A (<i>n</i> = 121)	45.5 (42.8–50.2)	54.2 (50.3–58.7)	62.3 (58.8–NA)	80.8 (NA–NA)
SPMS (<i>n</i> = 1353)	47.9 (47.3–48.5)	47.2 (46.3–47.7)	59.5 (58.2–60.6)	NA (76.0–NA)
SPMS-N (<i>n</i> = 950)	49.0 (48.2–49.9)	48.4 (47.5–49.1)	60.5 (59.5–63.6)	NA (76.0–NA)
SPMS-A (<i>n</i> = 403)	44.7 (43.2–46.7)	44.1 (42.9–45.6)	55.5 (54.4–58.0)	71.8 (71.8–NA)

Table 4. Ages at onset of the progressive phase and at confirmed EDSS scores, by time period (continued)

Median ages at onset of the progressive phase (Kaplan-Meier estimator) and at confirmed EDSS scores \geq 4, 6, and 7 (Turnbull estimator), for the complete dataset and for three constituent time periods. Analyses for each time period include only those patients with onset of PPMS or (operationally diagnosed) SPMS during that period, and only EDSS scores recorded by the end of that period (such that the combined number of patients across the three periods is less than the total number of patients in the complete dataset). Note that in some cases data were insufficient for complete Turnbull estimates of the median ages at EDSS \geq 6 or 7.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.

When mean EDSS trajectories were assessed using only EDSS scores recorded during the progressive phase, patients with SPMS had higher scores at younger ages but flatter slopes over time versus those with PPMS (**figure 4A**). When trajectories were assessed including scores during RRMS, patients with RRMS-SPMS had steeper slopes at younger ages versus those with PPMS (**figure 4B**). These observations are consistent with a model in which the earlier onset of disability accrual in the RRMS-SPMS phenotype is balanced by faster disability accrual in PPMS, such that disability trajectories of the two phenotypes ultimately converge.

Sensitivity and mediation analyses

The sensitivity analyses corroborated the above findings; results are summarized in **supplementary tables 1.7–1.8**.

The analyses with SPMS defined by physician diagnosis (*n* = 4610 SPMS; 32% male; 50% SPMS-A) are presented in **supplementary material section 2**. Results corroborated those from the main analyses with respect to age at onset (Kaplan-Meier median for SPMS = 46.3 years; 95% CI =

1
2
3 45.8–46.7); hazard of disability accrual in SPMS versus PPMS (HR = 0.89; 95% CI = 0.83–0.95;
4 $P < .001$; complete model); ages at EDSS milestones; and mean EDSS trajectories. However, the
5 adjusted hazard of disability accrual was lower in both SPMS-A (HR = 0.85; 95% CI = 0.78–0.93;
6 $P < .001$) and SPMS-N (HR = 0.86; 95% CI = 0.79–0.94; $P = .001$) versus PPMS-N. Additionally,
7 EDSS ≥ 7 was reached at younger ages in SPMS (25th percentile = 54.4 years; 95% CI = 53.7–
8 55.2) versus PPMS ($P < .001$), although still to a lesser extent than EDSS ≥ 4 . Among patients
9 with physician-diagnosed SPMS, 2458 (53%) met the operationalized diagnostic criteria.

10
11
12
13
14 Mediation analysis indicated that the causal effect of phenotype on disability accrual comprises
15 only small indirect effects via DMT exposure (in 25% increments; HR = 1.02; 95% CI = 0.97–
16 1.07) and EDSS score frequency (annualized; HR = 1.01; 95% CI = 1.00–1.02), and a substantial
17 “direct” effect via other pathways (SPMS versus PPMS; HR = 0.78; 95% CI = 0.70–0.86).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

In this observational study comparing 1872 patients with PPMS and 2575 with operationally diagnosed SPMS from the MSBase cohort, SPMS had an older median age at onset by 3 years (46.7 versus 43.9), and a 14% lower hazard of disability accrual (adjusted for sex, age, baseline disability, disease-modifying and immunosuppressant therapy, and EDSS score frequency). However, patients who reached wheelchair dependence did so at similar ages in PPMS and SPMS. These results were robust in sensitivity analyses, in particular when including only patients with regular follow-up; when requiring confirmation of disability accrual events over at least one year; and when restricting analysis to EDSS scores ≥ 4 for both PPMS and SPMS. The older onset and lower hazard of disability accrual in SPMS were also demonstrated when SPMS was identified by physician diagnosis rather than operationalized criteria. Mediation analysis demonstrated that the lower hazard in SPMS versus PPMS was not reducible to differences in DMT exposure or EDSS score frequency between the two phenotypes.

Later onset of SPMS versus PPMS

Whether onset of the progressive phase occurs at older ages in SPMS versus PPMS remains contested.^{2,16} We observe older median age at onset of SPMS versus PPMS by approximately 2–4 years, whether SPMS onset is identified by operationalized or physician diagnosis. This may reflect increasing use of high-efficacy DMT during RRMS, which defers conversion to SPMS.^{29,30} However—consistent with some prior studies,³¹ but not others³²—we find that PPMS and SPMS show similar secular trends toward older ages at onset and at EDSS milestones, suggesting contributing factors beyond DMT uptake.

Slower disability accrual in SPMS versus PPMS; similar long-term trajectories

Modern cohort studies comparing PPMS and SPMS have reported either similar rates of disability accrual in the two phenotypes,^{9,16} or faster accrual in SPMS.^{8,18} These analyses used unadjusted life-table or Kaplan-Meier estimates of intervals between milestones (onset of the progressive phase; EDSS 4, 6, 7, or 8). In contrast, we observe slower disability accrual in SPMS. Our analyses used multivariable proportional hazards models, allowing adjustment for phenotypic differences in baseline disability, sex, age, and clinical management, without restriction to a single EDSS interval. It is possible that disability accrual in our SPMS cohort was tempered by DMT¹⁸—particularly in SPMS-A, where inflammation is treatable.³³ However, our mediation analysis suggests this explanation is insufficient, as do the secular trends noted above.

Our results cohere with the view that disability trajectories in RRMS-SPMS and PPMS ultimately converge as patients age^{2,7–12,14}—perhaps reflecting a shift in pathology from focal inflammation

1
2
3 to diffuse neurodegeneration in the progressive phase,³⁴ accompanied by declining neurologic
4 repair and compensation mechanisms.³⁵ Whereas disability accrual in RRMS-SPMS begins at
5 RRMS onset (mean age 32.3 years in our SPMS cohort), accrual in PPMS typically begins a decade
6 later (mean age 43.5 years). This delayed presentation in PPMS may be counterbalanced by
7 younger onset of the progressive phase and faster disability accrual relative to SPMS, yielding
8 convergence of long-term disability trajectories across phenotypes.
9

10
11
12 Supporting this interpretation, patients with PPMS and operationally diagnosed SPMS reached
13 EDSS ≥ 6 and EDSS ≥ 7 (wheelchair dependence) at similar ages, despite the younger ages at
14 EDSS ≥ 4 in the SPMS cohort. Likewise, mean EDSS trajectories for the two phenotypes
15 converged with age (whether SPMS was diagnosed operationally or by physicians). However,
16 patients with SPMS under physician diagnosis reached EDSS ≥ 7 younger than those with PPMS,
17 as observed in some earlier cohorts.^{7,8}
18
19
20
21

22 *Patients with and without activity*

23
24
25 In both PPMS and SPMS, patients with superimposed activity had younger onset of the progressive
26 phase than those without, consistent with age-related declines in MS activity^{18,36} and prior findings
27 for PPMS.⁸ Using operationalized diagnosis, SPMS-A had slower disability accrual versus PPMS-
28 N, whereas SPMS-N did not. This may reflect a genuine phenotypic difference (e.g., SPMS-A may
29 typically involve milder “progressive” pathology than other phenotypes). However, using
30 physician diagnosis, both SPMS-N and SPMS-A had slower disability accrual versus PPMS-N.
31
32
33

34 *Limitations*

35
36
37 It remains difficult to precisely diagnose the onset of progressive MS.³⁵ Here, we have addressed
38 the particular challenge of identifying the transition from RRMS to SPMS by using operationalized
39 diagnosis, while also demonstrating consistent results under physician diagnosis. However, both
40 methods of diagnosing SPMS are imperfect, as is physician diagnosis of PPMS. Importantly, our
41 finding of slower disability accrual in SPMS is replicated in the estimates of median ages at EDSS
42 milestones, and in the sensitivity analysis modelling accrual from EDSS 4–5 rather than from onset
43 of the progressive phase. These two analyses depend on correct inclusion of patients with PPMS
44 and SPMS, but not on precise identification of dates of onset.
45
46
47
48

49
50 Our main analyses using operationalized diagnosis of conversion from RRMS to SPMS required
51 initial EDSS scores ≤ 3 during RRMS, with follow-up commencing at SPMS onset. This has
52 several implications. First, many patients in our SPMS cohort lack a physician diagnosis,
53 potentially because operationalized diagnosis detects SPMS earlier.²² Second, the early progressive
54 phase is preferentially sampled in SPMS. Third, the requirement for initial records from the
55
56
57
58
59
60

relapsing-remitting phase for patients with SPMS may have upwardly biased our estimated ages at onset and at EDSS milestones in this phenotype. Fourth, operationalized criteria for SPMS are not yet widely applied in clinical trials, which may limit generalizability of our findings. These concerns are addressed by the sensitivity analysis assessing disability accrual from EDSS 4–5, and by the analyses using physician-diagnosed SPMS. (For physician-diagnosed SPMS, ages at onset may be biased upward by diagnostic delay^{20,21}; conversely, ages at onset and/or at EDSS milestones may be biased downward by preferential recognition of SPMS in patients with more aggressive disease.)

Our Kaplan-Meier analyses of ages at onset are right-truncated (restricted to individuals who experienced the outcome), so likely yielded net underestimates. This proved necessary in comparing PPMS and SPMS: whereas more accurate estimates of SPMS onset are obtained by including patients with unconverted RRMS in the risk set,¹⁶ the equivalent is not feasible for PPMS given its covert prodromal pathology.^{2,37}

Identification of patients with activity is complicated by variability in diagnosing relapse; unrecorded relapses; censoring before a patient's first superimposed relapse; and DMT-induced relapse suppression. We therefore assigned "active" status inclusively, if any relapse was recorded during the progressive phase. PPMS-A cohorts may also include cases of misclassified RRMS—although notably, we observe similar hazards of disability accrual in PPMS-A and PPMS-N.

The presented analyses did not differentiate disability accrual events by the magnitude of qualifying EDSS increases. This may yield a detection bias, understating disability accrual in patients who experienced rapid or repeated worsening between clinical visits. However, under both operationalized and physician diagnosis of SPMS, the median annualized frequency of EDSS scores was greater in SPMS versus PPMS (**table 1; supplementary table 2.2**). Moreover, epochs with an EDSS increase at least twice threshold were less common in both operationally diagnosed SPMS (7.5% of epochs in both SPMS-N and SPMS-A) and physician-diagnosed SPMS (11.1% in both SPMS-N and SPMS-A) versus PPMS (13.3% of epochs; PPMS-N, 13.1%; PPMS-A, 13.4%). Therefore, our finding of slower disability accrual in SPMS versus PPMS likely does not result from a preferential failure to capture accrual between visits in SPMS.

The current phenotypic classification⁴ distinguishing RRMS, PPMS, and SPMS itself warrants examination,³⁵ given recent demonstrations of progression independent of activity during RRMS,^{38–40} and evidence that the recognized phenotypes share the same underlying pathologies.³⁴ Nonetheless, our results suggest that the distinction between PPMS and SPMS captures clinically salient differences in disability accrual.

Clinic-based observational registry data are subject to selection biases, confounding, and incomplete records.⁴¹ Here, data limitations precluded analyses of imaging records, and of death

1
2
3 as a competing risk. Additionally, although the EDSS remains widely used in both observational
4 studies and clinical trials, it suffers from poor intra- and inter-rater reliability at low scores; poor
5 responsiveness to upper-limb and cognitive impairments; and non-linearity.²⁴ In future studies,
6 magnetic resonance imaging may help to identify prodromal MS pathology,²¹ refine phenotypic
7 classifications,⁴² and provide paraclinical outcomes.⁴³
8
9

10 11 *Conclusion* 12

13
14 We show that relative to PPMS, the progressive phase in SPMS begins at older ages, but with
15 greater baseline disability due to the preceding relapsing-remitting phase. Patients with SPMS then
16 experience slower disability accrual than those with PPMS, ultimately yielding convergent
17 disability trajectories, with similar ages at more severe disability milestones in the two phenotypes.
18 One should exercise caution about amalgamating PPMS and SPMS in clinical trials, and consider
19 the difference in disability trajectories when comparing disability outcomes. At the same time, the
20 convergence of disability trajectories in PPMS and SPMS coheres with the view that MS pathology
21 interacts with normal central nervous system aging to drive similar long-term outcomes across
22 phenotypes.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article Information

Author affiliations

CORe, Department of Medicine, University of Melbourne, Melbourne, Australia (Harding-Forrester, Roos, Nguyen, Malpas, Diouf, Moradi, Sharmin, Kalincik); Multiple Sclerosis Centre, Department of Neurology, Royal Melbourne Hospital, Melbourne, Australia (Roos, Nguyen, Malpas, Diouf, Moradi, Sharmin, Kalincik); Multiple Sclerosis Unit, Hospital Universitario Virgen Macarena, Sevilla, Spain (Izquierdo, Eichau); Department of Medical and Surgical Sciences and Advanced Technologies 'G. F. Ingrassia', University of Catania, Catania, Italy (Patti); Department of Neurology and Center of Clinical Neuroscience, First Faculty of Medicine and General University Hospital, Charles University, Prague, Czech Republic (Horáková, Havrdová); Centre de Recherche du Centre Hospitalier de l'Université de Montréal (CHUM), Montréal, Canada (Prat, Girard, Duquette); Department of Neuroscience, Faculty of Medicine, Université de Montréal, Montréal, Canada (Prat, Girard, Duquette); Neuro Rive-Sud, Longueuil, Canada (Grand'Maison); Department of Neuroscience, Imaging, and Clinical Sciences, University G. d'Annunzio of Chieti-Pescara, Chieti, Italy (Onofri); IRCCS Istituto delle Scienze Neurologiche di Bologna, Bologna, Italia (Lugaresi); Dipartimento di Scienze Biomediche e Neuromotorie, Università di Bologna, Bologna, Italia (Lugaresi); Centre Intégré de Santé et de Services Sociaux (CISSS) de Chaudière-Appalaches, Lévis, Canada (Grammond); Department of Neurology, Faculty of Medicine, Dokuz Eylül University, İzmir, Turkey (Özakbaş); Department of Neuroscience, Psychology, Pharmacology, and Child Health (NEUROFARBA), University of Florence, Florence, Italy (Amato); IRCCS Fondazione Don Carlo Gnocchi, Florence, Italy (Amato); Department of Neurology, Zuyderland Medical Center, Sittard-Geleen, The Netherlands (Gerlach); Department of Neuroscience, Azienda Ospedaliera-Universitaria di Modena, Modena, Italy (Sola, Ferraro); Department of Biomedical, Metabolic, and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy (Ferraro); Department of Neurology, Box Hill Hospital, Melbourne, Australia (Buzzard, Skibina); School of Medicine and Public Health, University of Newcastle, Newcastle, Australia (Lechner-Scott); Department of Neurology, John Hunter Hospital, Newcastle, Australia (Lechner-Scott); Division of Neurology, Department of Medicine, Al-Amiri Hospital, Kuwait City, Kuwait (Alroughani); Department of Neurology, Faculty of Medicine, Karadeniz Technical University (KTU) Medical Faculty, Farabi Hospital, Trabzon, Turkey (Boz); Cliniques Universitaires Saint-Luc, UCLouvain, Brussels, Belgium (van Pesch); Unità Operative Complesse (UOC) di Neurologia, Azienda Sanitaria Unica Regionale Marche - AV3, Macerata, Italy (Cartechini); Medical Faculty, Ondokuz Mayıs University, Samsun, Turkey (Terzi); Department of Neurology, Garibaldi Hospital, Catania, Italy (Maimone); Department of Neurology, Hospital Germans Trias i Pujol, Badalona, Spain (Ramo-Tello); Nehme and Therese Tohme Multiple Sclerosis Center, Department of Neurology, American University of Beirut Medical Center, Beirut, Lebanon (Yamout, Houry); Department of Neurology, Azienda Ospedaliera di Rilievo Nazionale San Giuseppe Moscati Avellino, Avellino, Italy (Spitaleri); Department of Neurology, Centro Hospitalar Universitário de São João, Porto, Portugal (Sá); Faculty of Health Sciences, University Fernando Pessoa, Porto, Portugal (Sá); Department of Neurology, Hospital Clinic de Barcelona, Barcelona, Spain (Blanco); Department of Medicine and Surgery, University of Parma, Parma, Italy (Granella); Flinders Medical Centre, Flinders University, Adelaide, Australia (Slee); Monash Medical Centre, Monash University, Melbourne, Australia (Butler); Department of Neurology, Razi Hospital, Manouba, Tunisia (Sidhom, Gouider); IRCCS Mondino Foundation, Pavia, Italy (Bergamaschi); Department of Neurology, Hacettepe University, Ankara, Turkey (Karabudak); Department of Neurology, Nemocnice Jihlava, Jihlava, Czech Republic (Ampapa); Department of Neurology, Hospital de Galdakao-Usansolo, Galdakao, Spain (Sánchez-Menoyo); Centre de Santé et de Services Sociaux (CSSS) de Saint-Jérôme, Saint-Jérôme, Canada (Prévost); Instituto de Investigación Sanitaria Biodonostia, Hospital Universitario Donostia, San Sebastián, Spain (Castillo-Triviño); Centre for Clinical Research, Faculty of Medicine, University of Queensland, Brisbane, Australia (McCombe); Department of Neurology, Austin Health, Melbourne, Australia (Macdonell); Department of Neurology, Ghent University Hospital, Ghent, Belgium (Laureys, Van Hijfte); St. Michael's Hospital,

1
2
3 Toronto, Canada (Oh); Department of Neurology, School of Medicine, Koc University, and Koc University Research
4 Center for Translational Medicine (KUTTAM), Istanbul, Turkey (Altintas); Groene Hart Ziekenhuis, Gouda, The
5 Netherlands (de Gans); Haydarpasa Numune Training and Research Hospital, Istanbul, Turkey (Türkoğlu); Multiple
6 Sclerosis and Neuroimmunology Unit, Central Clinical School, Monash University, Melbourne, Australia (van der
7 Walt, Butzkueven); Department of Neurology, The Alfred Hospital, Melbourne, Australia (Butzkueven); Westmead
8 Hospital, Sydney, Australia (Vucic); Brain and Mind Centre, University of Sydney, Sydney, Australia (Barnett);
9 Centro de Esclerosis Múltiple de Buenos Aires (CEMBA), Hospital Italiano de Buenos Aires, Argentina (Cristiano);
10 Liverpool Hospital, Sydney, Australia (Hodgkinson); Ospedali Riuniti di Salerno, Salerno, Italy (Iuliano); Neurologic
11 Clinic and Policlinic and MS Center, Departments of Medicine, Biomedicine, and Clinical Research, University
12 Hospital Basel, University of Basel, Basel, Switzerland (Kappos, Kuhle); Research Center for Clinical
13 Neuroimmunology and Neuroscience, University Hospital Basel, University of Basel, Basel, Switzerland (Kappos,
14 Kuhle); Isfahan University of Medical Sciences, Isfahan, Iran (Shaygannejad); Bakirkoy Education and Research
15 Hospital for Psychiatric and Neurological Diseases, Istanbul, Turkey (Soysal); Department of Neurology, Jacobs
16 School of Medicine and Biomedical Sciences, Buffalo, United States (Weinstock-Guttman); Universitair MS Centrum,
17 Hasselt-Pelt, Belgium (Van Wijmeersch); Rehabilitation & MS Centre, Pelt, Belgium (Van Wijmeersch).

21 *Acknowledgements*

22
23
24 We thank all patients and caregivers who contributed data to the MSBase registry and participated in this study.
25 MSBase Study Group co-investigators: Magd Zakaria, MD (Faculty of Medicine, Ain Shams University, Cairo,
26 Egypt).

29 *Contributors*

30
31 Drs Harding-Forrester and Kalincik had full access to all of the data in the study and take responsibility for the integrity
32 of the data and the accuracy of the data analysis.

33 Concept and design: Harding-Forrester, Kalincik.

34 Acquisition, analysis, or interpretation of data: All authors.

35 Drafting of the manuscript: Harding-Forrester.

36 Critical revision of the manuscript for important intellectual content: All authors.

37 Statistical analysis: Harding-Forrester, Kalincik.

38 Obtained funding: Kalincik.

39 Administrative, technical, or material support: Kalincik.

40 Supervision: Kalincik.

44 *Funding*

45
46
47 This study was financially supported by the National Health and Medical Research Council of Australia (1129189,
48 1140766). The MSBase Foundation is a not-for-profit organization that has received support from Biogen, CSL,
49 Merck, Novartis, Roche, Sanofi, and Teva; this study was conducted independently of the sponsors.

Competing interests

Dr Roos served on scientific advisory boards for Novartis and Merck, and received conference travel support and/or speaker honoraria from Roche, Novartis, Biogen, Teva, Sanofi Genzyme, and Merck. Dr Nguyen received grants from MS Research Australia; grants, personal fees, and nonfinancial support from Biogen; grants and personal fees from Merck Serono; personal fees from Teva and Novartis; and nonfinancial support from Roche and Sanofi Genzyme. Dr Izquierdo received speaking honoraria from Biogen, Novartis, Sanofi, Merck, Roche, Almirall, and Teva. Dr Eichau received speaker honoraria and consultant fees from Biogen Idec, Novartis, Merck, Bayer, Sanofi Genzyme, Roche, and Teva. Dr Patti received speaker honoraria and advisory board fees from Almirall, Bayer, Biogen, Celgene, Merck, Novartis, Roche, Sanofi Genzyme, and Teva, and research funding from Biogen, Merck, FISM (Fondazione Italiana Sclerosi Multipla), Reload Onlus Association, and the University of Catania. Dr Horáková received speaker honoraria and consulting fees from Biogen, Merck, Teva, Roche, Sanofi Genzyme, and Novartis, and support for research activities from Biogen and the Czech Ministry of Education (project PROGRES Q27/LF1). Dr Havrdová received honoraria or research support from Biogen, Merck Serono, Novartis, Roche, and Teva; has been a member of advisory boards for Actelion, Biogen, Celgene, Merck Serono, Novartis, and Sanofi Genzyme; and received research support from the Czech Ministry of Education (project PROGRES Q27/LF1). Dr Girard received consulting fees from Teva Canada Innovation, Biogen, Novartis, and Sanofi Genzyme; lecture payments from Teva Canada Innovation, Novartis, and EMD; and research support from the Canadian Institutes of Health Research. Dr Duquette served on editorial boards for, and has been supported to attend meetings by, EMD, Biogen, Novartis, Genzyme, and Teva Neuroscience; he holds grants from the Canadian Institutes of Health Research and the MS Society of Canada, and received funding for investigator-initiated trials from Biogen, Novartis, and Genzyme. Dr Grand'Maison received honoraria or research funding from Biogen, Genzyme, Novartis, Teva Neurosciences, Mitsubishi, and ONO Pharmaceuticals. Dr Lugaresi received personal compensation for consulting, serving on a scientific advisory board, speaking, or other activities from Biogen, Merck Serono, Mylan, Novartis, Roche, Sanofi Genzyme, and Teva; her institutions have received research grants from Novartis (in the past 4 years). Dr Grammond served on advisory boards for Novartis, EMD Serono, Roche, Biogen Idec, Sanofi Genzyme, and Pendopharm; received grant support from Genzyme and Roche; and received research grants for his institution from Biogen Idec, Sanofi Genzyme, and EMD Serono. Dr Amato received honoraria as a consultant on scientific advisory boards for Biogen, Bayer Schering, Merck, Teva, and Sanofi-Aventis, and received research grants by Biogen, Bayer Schering, Merck, Teva, and Novartis. Dr Sola served on scientific advisory boards for Biogen Idec and Teva; received funding for travel and speaker honoraria from Biogen Idec, Merck, Teva, Sanofi Genzyme, Novartis, and Bayer; and received research grants for her institution from Bayer, Biogen, Merck, Novartis, Sanofi, and Teva. Dr Ferraro received travel grants and/or speaker honoraria from Merck, Teva, Novartis, Biogen, and Sanofi Genzyme. Dr Buzzard received honoraria and consulting fees from Biogen, Teva, Novartis, Sanofi Genzyme, Roche, Merck, CSL, and Grifols. Dr Lechner-Scott received travel compensation from Novartis, Biogen, Roche, and Merck; her institution received honoraria for talks and advisory board commitments, as well as research grants from Biogen, Merck, Roche, Teva, and Novartis. Dr Alroughani received honoraria as a speaker and for serving on scientific advisory boards from Bayer, Biogen, GSK, Merck, Novartis, Roche, and Sanofi Genzyme. Dr Boz received conference travel support from Biogen, Novartis, Bayer Schering, Merck, and Teva, and participated in clinical trials by Sanofi-Aventis, Roche, and Novartis. Dr van Pesch received travel grants from Merck, Biogen, Sanofi, Celgene, Almirall, and Roche; his institution received research grants and consultancy fees from Roche, Biogen, Sanofi, Celgene, Merck, and Novartis Pharma. Dr Terzi received travel grants from Novartis, Bayer Schering, Merck, and Teva, and participated in clinical trials by Sanofi-Aventis, Roche, and Novartis. Dr Maimone received speaker honoraria for advisory board service and travel grants from Almirall, Biogen, Merck, Novartis, Roche, Sanofi Genzyme, and Teva. Dr Ramo-Tello received research funding, compensation for travel, or speaker honoraria from Biogen, Novartis, Genzyme, and Almirall. Dr Spitaleri received honoraria as a consultant on scientific advisory boards from Bayer Schering, Novartis, and Sanofi-Aventis, and compensation for travel from Novartis, Biogen, Sanofi-

1
2
3 Aventis, Teva, and Merck. Dr Granella received an institutional research grant from Biogen and Sanofi Genzyme;
4 served on scientific advisory boards for Biogen, Novartis, Merck, Sanofi Genzyme, and Roche; and received funding
5 for travel and speaker honoraria from Biogen, Merck, and Sanofi-Aventis. Dr Slee participated in, but did not receive
6 honoraria for, advisory board activity for Biogen, Merck, Bayer Schering, Sanofi-Aventis, and Novartis. Dr
7 Bergamaschi received speaker honoraria from Bayer Schering, Biogen, Genzyme, Merck, Novartis, Sanofi-Aventis,
8 and Teva; research grants from Bayer Schering, Biogen, Merck, Novartis, Sanofi-Aventis, and Teva; and congress,
9 travel, and accommodation expense compensations from Almirall, Bayer Schering, Biogen, Genzyme, Merck,
10 Novartis, Sanofi-Aventis, and Teva. Dr Ampapa received conference travel support from Novartis, Teva, Biogen,
11 Bayer, and Merck, and participated in clinical trials by Biogen, Novartis, Teva, and Actelion. Dr Sánchez-Menoyo
12 received travel compensation from Novartis and Biogen; received speaking honoraria from Biogen, Novartis, Sanofi,
13 Merck, Almirall, Bayer, and Teva; and participated in a clinical trial by Biogen. Dr Prévost received travel
14 compensation from Novartis, Biogen, Genzyme, and Teva, and speaking honoraria from Biogen, Novartis, Genzyme
15 and Teva. Dr Castillo-Triviño received speaking or consulting fees and/or travel funding from Bayer, Biogen, Merck,
16 Novartis, Roche, Sanofi Genzyme, and Teva. Dr Laureys received travel and/or consultancy compensation from Sanofi
17 Genzyme, Roche, Teva, Merck, Novartis, Celgene, and Biogen. Dr Oh received research funding from the MS Society
18 of Canada, the National MS Society, Brain Canada, Biogen Idec, Roche, and EMD Serono, and personal compensation
19 for consulting or speaking from EMD Serono, Sanofi Genzyme, Biogen Idec, Roche, Celgene, and Novartis. Dr
20 Altintas received personal fees and speaker honoraria from Teva, Merck, Biogen Gen Pharma, Roche, Novartis, Bayer,
21 and Sanofi Genzyme, and received travel and registration grants from Merck, Biogen Gen Pharma, Roche, Sanofi
22 Genzyme, and Bayer. Dr Butzkueven received compensation for consulting, talks, and advisory or steering board
23 activities from Biogen, Merck, Novartis, Genzyme, Alfred Health, and Oxford Health Policy Forum, and research
24 support from Novartis, Biogen, Roche, Merck, the National Health and Medical Research Council of Australia,
25 Pennycook Foundation, and MS Research Australia. Dr Barnett served on scientific advisory boards for Biogen,
26 Novartis, and Genzyme, received conference travel support from Biogen and Novartis, and serves on steering
27 committees for trials conducted by Novartis; his institution received research support from Biogen, Merck, and
28 Novartis. Dr Cristiano received honoraria as a consultant on scientific advisory boards for Biogen, Bayer Schering,
29 Merck, Genzyme, and Novartis, and participated in clinical trials or other research projects by Merck, Roche, and
30 Novartis. Dr Hodgkinson received honoraria and consulting fees from Novartis, Bayer Schering, and Sanofi, and travel
31 grants from Novartis, Biogen Idec, and Bayer Schering. Dr Iuliano received compensation for travel, accommodations,
32 and meeting expenses from Bayer Schering, Biogen, Merck, Novartis, Sanofi-Aventis, and Teva. Dr Kappos received
33 research support from Acorda, Actelion, Allozyne, BaroFold, Bayer HealthCare, Bayer Schering, Bayhill
34 Therapeutics, Biogen, Elan, European Union, Genmab, Gianni Rubatto Foundation, GlaxoSmithKline, Glenmark,
35 MediciNova, Merck, Novartis, Novartis Research Foundation, Roche, Roche Research Foundation, Sanofi-Aventis,
36 Santhera, the Swiss MS Society, the Swiss National Research Foundation, Teva Neuroscience, UCB, and Wyeth. Dr
37 Weinstock-Guttman participated in speakers' bureaus and/or served as a consultant for Biogen, EMD Serono, Novartis,
38 Genentech, Celgene/Bristol Meyers Squibb, Sanofi Genzyme, Bayer, Janssen, and Horizon; received grant/research
39 support from these same agencies; and serves on editorial boards for BMJ Neurology, Children, CNS Drugs, MS
40 International, and Frontiers Epidemiology. Dr Van Wijmeersch received research and travel grants and honoraria for
41 advisory and speaking fees from Bayer Schering, Biogen, Sanofi Genzyme, Merck, Novartis, Roche, and Teva. Dr
42 Kalincik served on scientific advisory boards for BMS, Roche, Sanofi Genzyme, Novartis, Merck, and Biogen, and
43 the steering committee for the Brain Atrophy Initiative by Sanofi Genzyme; received conference travel support and/or
44 speaker honoraria from WebMD Global, Novartis, Biogen, Sanofi Genzyme, Teva, BioCSL, and Merck; and received
45 support for research or educational events from Biogen, Novartis, Genzyme, Roche, Celgene, and Merck. No other
46 disclosures relevant to the manuscript were reported.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Patient consent for publication*
4

5
6 Not required.
7

8 *Ethics approval*
9

10
11 MSBase is registered with WHO ICTRP (anzctr.org.au identifier ACTRN12605000455662). MSBase is approved by
12 the Melbourne Health Human Research Ethics Committee (reference 2006.044), and by ethics committees of
13 participating centers as required by local laws and regulations.
14

15
16 *Provenance and peer review*
17

18 Not commissioned; externally peer reviewed.
19

20
21 *Data availability statement*
22

23
24 Data from each center contributing to MSBase are available at the discretion of the associated MSBase Principal
25 Investigator(s). See <https://www.msbase.org>.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Tremlett H, Zhao Y, Devonshire V. Natural history of secondary-progressive multiple sclerosis. *Mult Scler*. 2008;14(3):314–324.
2. Tutuncu M, Tang J, Zeid NA, *et al*. Onset of progressive phase is an age-dependent clinical milestone in multiple sclerosis. *Mult Scler*. 2013;19(2):188–198.
3. Scalfari A, Neuhaus A, Daumer M, Muraro PA, Ebers GC. Onset of secondary progressive phase and long-term evolution of multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 2014;85(1):67–75.
4. Lublin FD, Reingold SC, Cohen JA, *et al*. Defining the clinical course of multiple sclerosis: The 2013 revisions. *Neurology*. 2014;83(3):278–286.
5. Kurtzke JF. Rating neurologic impairment in multiple sclerosis: An expanded disability status scale (EDSS). *Neurology*. 1983;33(11):1444–1452.
6. Confavreux C, Vukusic S, Adeleine P. Early clinical predictors and progression of irreversible disability in multiple sclerosis: An amnesic process. *Brain*. 2003;126(4):770–782.
7. Confavreux C, Vukusic S. Age at disability milestones in multiple sclerosis. *Brain*. 2006;129(3):595–605.
8. Confavreux C, Vukusic S. Natural history of multiple sclerosis: A unifying concept. *Brain*. 2006;129(3):606–616.
9. Kremenchutzky M, Rice GPA, Baskerville J, Wingerchuk DM, Ebers GC. The natural history of multiple sclerosis: A geographically based study. 9: Observations on the progressive phase of the disease. *Brain*. 2006;129(3):584–594.
10. Koch M, Mostert J, Heersema D, De Keyser J. Progression in multiple sclerosis: Further evidence of an age dependent process. *J Neurol Sci*. 2007;255(1–2):35–41.
11. Leray E, Yaouanq J, Le Page E, *et al*. Evidence for a two-stage disability progression in multiple sclerosis. *Brain*. 2010;133(7):1900–1913.
12. Scalfari A, Neuhaus A, Daumer M, Ebers GC, Muraro PA. Age and disability accumulation in multiple sclerosis. *Neurology*. 2011;77(13):1246–1252.
13. McGinley M, Ontaneda D. MS progression is predominantly driven by age-related mechanisms – NO. *Mult Scler*. 2019;25(7):904–906.
14. Trojano M, Liguori M, Bosco Zimatore G, *et al*. Age-related disability in multiple sclerosis. *Ann Neurol*. 2002;51(4):475–480.
15. Debouverie M, Pittion-Vouyovitch S, Louis S, Guillemin F, LORSEP Group. Natural history of multiple sclerosis in a population-based cohort. *Eur J Neurol*. 2008;15(9):916–921.
16. Tremlett H, Zhao Y, Devonshire V. Natural history comparisons of primary and secondary progressive multiple sclerosis reveals differences and similarities. *J Neurol*. 2009;256(3):374–381.
17. Manouchehrinia A, Beiki O, Hillert J. Clinical course of multiple sclerosis: A nationwide cohort study. *Mult Scler*. 2017;23(11):1488–1495.
18. Paz Soldán MM, Novotna M, Abou Zeid N, *et al*. Relapses and disability accumulation in progressive multiple sclerosis. *Neurology*. 2015;84(1):81–88.
19. Kalincik T, Vivek V, Jokubaitis V, *et al*. Sex as a determinant of relapse incidence and progressive course of multiple sclerosis. *Brain*. 2013;136(12):3609–3617.
20. Katz Sand I, Krieger S, Farrell C, Miller AE. Diagnostic uncertainty during the transition to secondary progressive multiple sclerosis. *Mult Scler*. 2014;20(12):1654–1657.
21. Cree BAC, Arnold DL, Chataway J, *et al*. Secondary progressive multiple sclerosis: New insights. *Neurology*. 2021;97(8):378–388.
22. Lorscheider J, Buzzard K, Jokubaitis V, *et al*. Defining secondary progressive multiple sclerosis. *Brain*. 2016;139(9):2395–2405.
23. Kalincik T, Butzkueven H. The MSBase registry: Informing clinical practice. *Mult Scler*. 2019;25(14):1828–1834.
24. Kalincik T, Cutter G, Spelman T, *et al*. Defining reliable disability outcomes in multiple sclerosis. *Brain*. 2015;138(11):3287–3298.
25. Zhao Q, Sun J. Generalized log-rank test for mixed interval-censored failure time data. *Stat Med*. 2004;23(10):1621–1629.
26. Koch MW, Mostert J, Repovic P, Bowen JD, Uitdehaag B, Cutter G. Reliability of outcome measures in clinical trials in secondary progressive multiple sclerosis. *Neurology*. 2021;96(1):e111–e120.
27. VanderWeele TJ. Mediation analysis: A practitioner’s guide. *Annu Rev Public Health*. 2016;37:17–32.
28. Lange T, Rasmussen M, Thygesen LC. Assessing natural direct and indirect effects through multiple

- 1
2
3 pathways. *Am J Epidemiol*. 2014;179(4):513–518.
- 4 29. UCSF MS-EPIC Team, Cree BAC, Gourraud PA, *et al*. Long-term evolution of multiple sclerosis disability
5 in the treatment era. *Ann Neurol*. 2016;80(4):499–510.
- 6 30. Brown JW, Coles A, Horakova D, *et al*. Association of initial disease-modifying therapy with later
7 conversion to secondary progressive multiple sclerosis. *JAMA*. 2019;321(2):175–187.
- 8 31. Simonsen CS, Flemmen HØ, Broch L, *et al*. The course of multiple sclerosis rewritten: A Norwegian
9 population-based study on disease demographics and progression. *J Neurol*. 2021;268(4):1330–1341.
- 10 32. Beiki O, Frumento P, Bottai M, Manouchehrinia A, Hillert J. Changes in the risk of reaching multiple
11 sclerosis disability milestones in recent decades: A nationwide population-based cohort study in Sweden.
12 *JAMA Neurol*. 2019;76(6):665–671.
- 13 33. Lizak N, Malpas CB, Sharmin S, *et al*. Association of sustained immunotherapy with disability outcomes in
14 patients with active secondary progressive multiple sclerosis. *JAMA Neurol*. 2020;77(11):1398–1407.
- 15 34. Lassmann H. Pathogenic mechanisms associated with different clinical courses of multiple sclerosis. *Front*
16 *Immunol*. 2019;9:3116.
- 17 35. Kuhlmann T, Moccia M, Coetzee T, *et al*. Multiple sclerosis progression: Time for a new mechanism-driven
18 framework. *Lancet Neurol*. 2022;22(1):78–88.
- 19 36. Scalfari A, Lederer C, Daumer M, Nicholas R, Ebers GC, Muraro PA. The relationship of age with the
20 clinical phenotype in multiple sclerosis. *Mult Scler*. 2016;22(13):1750–1758.
- 21 37. Kantarci OH, Lebrun C, Siva A, *et al*. Primary progressive multiple sclerosis evolving from radiologically
22 isolated syndrome. *Ann Neurol*. 2016;79(2):288–294.
- 23 38. UCSF MS-EPIC Team, Cree BAC, Hollenbach JA, *et al*. Silent progression in disease activity-free relapsing
24 multiple sclerosis. *Ann Neurol*. 2019;85(5):653–666.
- 25 39. Kappos L, Wolinsky JS, Giovannoni G, *et al*. Contribution of relapse-independent progression vs relapse-
26 associated worsening to overall confirmed disability accumulation in typical relapsing multiple sclerosis in a
27 pooled analysis of 2 randomized clinical trials. *JAMA Neurol*. 2020;77(9):1132–1140.
- 28 40. Tur C, Carbonell-Mirabent P, Cobo-Calvo Á, *et al*. Association of early progression independent of relapse
29 activity with long-term disability after a first demyelinating event in multiple sclerosis. *JAMA Neurol*. 2022.
- 30 41. Debouverie M, Laforest L, Van Ganse E, Guillemin F, LORSEP Group. Earlier disability of the patients
31 followed in multiple sclerosis centers compared to outpatients. *Mult Scler*. 2009;15(2):251–257.
- 32 42. Eshaghi A, Young AL, Wijeratne PA, *et al*. Identifying multiple sclerosis subtypes using unsupervised
33 machine learning and MRI data. *Nat Commun*. 2021;12:2078.
- 34 43. Filippi M, Preziosa P, Barkhof F, *et al*. Diagnosis of progressive multiple sclerosis from the imaging
35 perspective: A review. *JAMA Neurol*. 2021;78(3):351–364.
- 36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legends & tables

Figure 1. Flow diagram of patient inclusion

52062 patients with RRMS, and not meeting the operationalized diagnostic criteria for SPMS, were excluded.

Figure 2. Nelson-Aalen cumulative hazard curves for confirmed disability accrual


(A) PPMS and SPMS. (B) PPMS-N, PPMS-A, SPMS-N, and SPMS-A. The cumulative hazard indicates the expected number of confirmed disability accrual events for a patient observed for a given duration of time. Shaded regions indicate 95% confidence intervals. A-G HR = adjusted hazard ratio obtained with Andersen-Gill models (see **table 3**); CI = confidence interval.

Figure 3. Turnbull survival curves for confirmed EDSS ≥ 7 (wheelchair dependence)

(A) PPMS and SPMS. (B) PPMS-N, PPMS-A, SPMS-N, and SPMS-A. Patients who entered observation having already reached confirmed EDSS ≥ 7 are analyzed as interval-censored observations, with EDSS ≥ 7 reached in the interval (date of MS onset, date of first observation] ($n = 37$ PPMS-N, 43 PPMS-A; none with SPMS). Among patients with SPMS, 84 reached EDSS ≥ 7 at the date of onset of the progressive phase (60 SPMS-N, 24 SPMS-A); 24 reached EDSS ≥ 7 prior to the progressive phase (16 SPMS-N, 8 SPMS-A). P values (generalized log-rank tests): SPMS versus PPMS, $P = .06$; SPMS-N versus PPMS-N, $P = .38$; SPMS-A versus PPMS-A, $P = .22$; PPMS-A versus PPMS-N, $P = .002$; SPMS-A versus SPMS-N, $P = .007$. Values in the risk table indicate the number of patients at risk at each age; the number of interval-censored patients with an interval including that age (square brackets); and the cumulative number of patients having reached EDSS ≥ 7 at that age (curved brackets). Shaded regions indicate 95% confidence intervals. EDSS = Expanded Disability Status Scale.

Figure 4. Mean EDSS trajectories in PPMS-N, PPMS-A, SPMS-N, and SPMS-A



(A) Including only study-eligible EDSS scores, starting from onset of the progressive phase. (B) Including all available EDSS scores, starting from MS onset (including RRMS). Mean scores are plotted for each 2-year age interval, for groups with 10 or more patients contributing data in that interval. Error bars indicate 95% confidence intervals. EDSS = Expanded Disability Status Scale.



Flow diagram of patient inclusion

52062 patients with RRMS, and not meeting the operationalized diagnostic criteria for SPMS, were excluded.

80x78mm (600 x 600 DPI)



Number at risk


PPMS-N	24	260	384	375	315	262	163	91
PPMS-A	14	278	409	411	339	269	184	115
SPMS-N	1541	808	420	194	92	40	17	7
SPMS-A	1033	807	547	338	190	86	33	15

Cumulative exits (events)

PPMS-N	0 (0)	38 (67)	164 (274)	336 (505)	486 (657)	607 (800)	740 (904)	834 (956)
PPMS-A	0 (0)	30 (88)	139 (274)	299 (494)	448 (697)	598 (834)	719 (919)	808 (984)
SPMS-N	0 (0)	733 (420)	1121 (694)	1347 (823)	1449 (875)	1501 (903)	1524 (912)	1534 (917)
SPMS-A	0 (0)	227 (291)	487 (548)	696 (714)	844 (823)	948 (856)	1001 (874)	1019 (878)


Nelson-Aalen cumulative hazard curves for confirmed disability accrual (A) PPMS and SPMS. (B) PPMS-N, PPMS-A, SPMS-N, and SPMS-A. The cumulative hazard indicates the expected number of confirmed disability accrual events for a patient observed for a given duration of time. Shaded regions indicate 95% confidence intervals. A-G HR = adjusted hazard ratio obtained with Andersen-Gill models (see table 3); CI = confidence interval.

184x222mm (600 x 600 DPI)



Numbers: at risk [in interval] (cumulative events)

PPMS	1872 [0] (0)	1824 [13] (5)	1632 [19] (49)	1211 [26] (140)	582 [17] (245)	183 [2] (316)	86 [0] (328)
SPMS	2574 [0] (1)	2532 [0] (15)	2203 [0] (80)	1353 [0] (232)	520 [0] (351)	95 [0] (402)	7 [0] (417)




Numbers: at risk [in interval] (cumulative events)

PPMS-N	935 [0] (0)	919 [7] (2)	853 [7] (16)	670 [14] (54)	339 [9] (110)	98 [0] (150)	39 [0] (157)
PPMS-A	937 [0] (0)	905 [6] (3)	779 [12] (33)	541 [12] (86)	243 [8] (135)	85 [2] (166)	47 [0] (171)
SPMS-N	1540 [0] (1)	1516 [0] (8)	1336 [0] (39)	842 [0] (129)	353 [0] (196)	69 [0] (229)	5 [0] (240)
SPMS-A	1034 [0] (0)	1016 [0] (7)	867 [0] (41)	511 [0] (103)	167 [0] (155)	26 [0] (173)	2 [0] (177)

Turnbull survival curves for confirmed EDSS ≥ 7 (wheelchair dependence)

(A) PPMS and SPMS. (B) PPMS-N, PPMS-A, SPMS-N, and SPMS-A. Patients who entered observation having already reached confirmed EDSS ≥ 7 are analyzed as interval-censored observations, with EDSS ≥ 7 reached in the interval (date of MS onset, date of first observation] ($n = 37$ PPMS-N, 43 PPMS-A; none with SPMS). Among patients with SPMS, 84 reached EDSS ≥ 7 at the date of onset of the progressive phase (60 SPMS-N, 24 SPMS-A); 24 reached EDSS ≥ 7 prior to the progressive phase (16 SPMS-N, 8 SPMS-A). P values (generalized log-rank tests): SPMS versus PPMS, $P = .06$; SPMS-N versus PPMS-N, $P = .38$; SPMS-A versus PPMS-A, $P = .22$; PPMS-A versus PPMS-N, $P = .002$; SPMS-A versus SPMS-N, $P = .007$. Values in the risk table indicate the number of patients at risk at each age; the number of interval-censored patients with an interval including that age (square brackets); and the cumulative number of patients having reached EDSS ≥ 7 at that age (curved brackets). Shaded regions indicate 95% confidence intervals. EDSS = Expanded Disability Status Scale.

185x210mm (600 x 600 DPI)



Mean EDSS trajectories in PPMS-N, PPMS-A, SPMS-N, and SPMS-A (A) Including only study-eligible EDSS scores, starting from onset of the progressive phase. (B) Including all available EDSS scores, starting from MS onset (including RRMS). Mean scores are plotted for each 2-year age interval, for groups with 10 or more patients contributing data in that interval. Error bars indicate 95% confidence intervals. EDSS = Expanded Disability Status Scale.

185x178mm (600 x 600 DPI)

Supplement 1. Supplementary material for the main analyses

Harding-Forrester S, Roos I, Nguyen A-L, et al. Disability accrual in primary and secondary progressive multiple sclerosis.

- Supplementary methods 1.1.** Data quality procedure
- Supplementary methods 1.2.** Directed acyclic graph: Association of confirmed disability accrual and progressive MS phenotype
- Supplementary methods 1.3.** Formulation of the Andersen-Gill model for hazard of confirmed disability accrual
- Supplementary methods 1.4.** Mediation analysis
- Supplementary figure 1.1.** Dates of onset of the progressive phase and of inclusion
- Supplementary table 1.1.** Included patients and treating centers, by country
- Supplementary table 1.2.** Relapse characteristics of included patients
- Supplementary table 1.3.** Andersen-Gill model for hazard of confirmed disability accrual, adjusted for sex and age
- Supplementary table 1.4.** Andersen-Gill model for hazard of confirmed disability accrual, corrected for proportional hazards violation
- Supplementary table 1.5.** Andersen-Gill model for hazard of confirmed disability accrual, stratified by phenotype
- Supplementary table 1.6.** Andersen-Gill model for hazard of confirmed disability accrual, stratified by phenotype, with activity–DMT interaction
- Supplementary table 1.7.** Sensitivity analyses—comparison of PPMS and SPMS
- Supplementary table 1.8.** Sensitivity analyses—comparison of PPMS-N, PPMS-A, SPMS-N, and SPMS-A

Abbreviations

MS = multiple sclerosis

PPMS = primary progressive MS; PPMS-N = PPMS with no superimposed relapse activity; PPMS-A = PPMS with superimposed relapse activity

RRMS = relapsing-remitting MS

SPMS = secondary progressive MS; SPMS-N = SPMS with no superimposed relapse activity; SPMS-A = SPMS with superimposed relapse activity

Supplementary methods 1.1.

Data quality procedure

Data quality procedure for MSBase data

Duplicate patient records were removed.

Centers with < 10 patient records were excluded.

Patients with missing date of birth were excluded.

MS onset dates after the data extraction date were removed.

Patients with missing date of the first clinical presentation of MS were excluded.

The dates of MS onset and of the first recorded MS course (e.g., primary progressive MS; clinically isolated syndrome) were aligned.

Patients with age at MS onset outside the range 0–100 years were excluded.

A logical sequence of MS courses was assured (e.g., clinically isolated syndrome, then relapsing-remitting MS, then secondary progressive MS).

Visit entries with a missing visit date, or with a recorded visit date before the recorded date of MS onset or after the data extraction date, were removed.

EDSS scores outside the range of possible EDSS values were removed.

Duplicate visit entries were merged.

MS relapses with missing onset date, or with a recorded onset date after the data extraction date, were removed.

Duplicate MS relapse entries were merged.

Relapses occurring within 30 days of each other were merged.

Visits preceded by relapses were identified, and time from the last relapse was calculated for each visit.

Therapies were labelled as discontinued or continuing.

Therapies with erroneous date entries were removed (e.g., treatment commencement date after treatment termination date; treatment commencement date after the data extraction date; commencement of disease-modifying therapy before the year 1980).

MS disease-modifying therapies were identified and labelled.

Duplicate treatment entries were removed.

Where multiple disease-modifying therapies were recorded simultaneously, the termination date of the previous therapy was imputed as the commencement date of the following therapy.


“Data extraction date” refers to the date of data extraction from MSBase (January 7, 2020).

Abbreviations: EDSS = Expanded Disability Status Scale.

Supplementary methods 1.2.

Directed acyclic graph:

Association of disability accrual and progressive MS phenotype



Random effects included in the models (treating center and patient identity) are not shown.
 EDSS = Expanded Disability Status Scale; DMT = disease-modifying therapy; IST = immunosuppressant therapy.

Review Only

Supplementary methods 1.3.**Formulation of the Andersen-Gill model for hazard of confirmed disability accrual**

Each patient is represented as a series of observations in successive time intervals:

(start of follow-up, first event], (first event, second event], ..., (kth event, end of follow-up].

The intensity process $\lambda_{ijk}(t)$ driving confirmed disability accrual, for observation epoch k in patient j at center i , is:

$$\lambda_{ijk}(t) = Y_{ijk}(t) \lambda_0(t) \exp \{ \beta_1 \text{phenotype}_{ij} + \beta_2 \text{sex}_{ij} + \beta_3 \text{age}_{ijk}(t) + \beta_4 \text{EDSS}_{ijk}(t) + \beta_5 \text{DMT}_{ijk}(t) + \beta_6 \text{IST}_{ijk}(t) + \beta_7 \text{freq}_{ijk}(t) + b_1 \text{center}_i + b_2 \text{patient}_{ij} \}$$

with terms defined as:

Term	Definition
$Y_{ijk}(t)$	At-risk indicator: $Y_{ijk}(t) = 1$ during follow-up, 0 after end of follow-up
$\lambda_0(t)$	Baseline hazard (non-parametric, non-negative, time-dependent)
Independent variables, constant	
phenotype_{ij}	Progressive MS phenotype— <i>either</i> : PPMS (reference) or SPMS, <i>or</i> PPMS-N (reference), PPMS-A, SPMS-N, or SPMS-A
sex_{ij}	Female (reference) or male
Independent variables, time-dependent	
$\text{age}_{ijk}(t)$	Age in years at the start of the current epoch
$\text{EDSS}_{ijk}(t)$	EDSS score at the start of the current epoch
$\text{DMT}_{ijk}(t)$	Proportion of time on disease-modifying therapy during the current epoch
$\text{IST}_{ijk}(t)$	Proportion of time on immunosuppressant therapy during the current epoch
$\text{freq}_{ijk}(t)$	Frequency of EDSS scores during the current epoch
Random effects	
center_i	Treating center, with Gaussian distribution: $b_1 \sim G(0, \Sigma\theta)$
patient_{ij}	Patient identity, with Gaussian distribution: $b_2 \sim G(0, \Sigma\eta)$

Abbreviations: EDSS = Expanded Disability Status Scale.

Supplementary methods 1.4.

Mediation analysis

In a proportional hazards model with a common outcome, the hazard ratio after adjustment for mediators may not accurately estimate the causal direct effect of the exposure of interest.¹ To assess whether the effect of phenotype (PPMS or SPMS) on disability accrual was mediated by DMT exposure and EDSS score frequency, a mediation analysis was performed using a broadly applicable natural effects estimation procedure.^{2,3} The proportion of time receiving DMT and the annualized frequency of EDSS scores were treated as mediators M within each epoch, with phenotype (SPMS versus PPMS) as the exposure A ; sex and age at the start of each epoch as covariates C ; and treating center and patient identity (nested within treating center) as random effects Z . Given $x = 2$ levels of the exposure and $k = 2$ mediators, a counterfactually augmented dataset was constructed from the observed data, with k auxiliary exposure variables and x^k copies of each observation epoch. The auxiliary exposure variables were assigned counterfactual values of the exposure in all possible combinations. Linear regression models were fitted for each mediator, conditional on the exposure and covariates, and were used to compute a weight for each observation i in the augmented dataset:

$$w_i = \prod_{k=1}^K \frac{\Pr(M^k = M_i^k | A = A_i^k, C = C_i, Z = Z_i)}{\Pr(M^k = M_i^k | A = A_i, C = C_i, Z = Z_i)}$$


A weighted Andersen-Gill model was then fitted to the augmented dataset, conditional on the true and auxiliary exposure variables, covariates, and random effects (but not on the mediators). This yielded estimates of the natural direct effect from the true-exposure coefficient, and of the natural indirect effect via each mediator from the coefficient for its corresponding auxiliary exposure variable. Standard errors were estimated by bootstrapping (1000 resamples of complete patient records).

Mediation analysis assumes adequate control for exposure–outcome, mediator–outcome, and exposure–mediator confounding, and the absence of mediator–outcome confounders affected by the exposure.^{1–3}

References

1. VanderWeele TJ. Mediation analysis: A practitioner's guide. *Annu Rev Public Health*. 2016;37:17–32.
2. Lange T, Vansteelandt S, Bekaert M. A simple unified approach for estimating natural direct and indirect effects. *Am J Epidemiol*. 2012;176(3):190–195.
3. Lange T, Rasmussen M, Thygesen LC. Assessing natural direct and indirect effects through multiple pathways. *Am J Epidemiol*. 2014;179(4):513–518.

Supplementary figure 1.1.
Dates of onset of the progressive phase and of inclusion



White regions in histogram bars indicate patients excluded by the requirement for onset of the progressive phase since January 1, 1995, who would otherwise have been included.

Supplementary table 1.1.
Included patients and treating centers, by country

Country	Centers	Patients (n)				
		Total	PPMS-N	PPMS-A	SPMS-N	SPMS-A
IT	11	1115	146	235	383	351
CA	6	605	178	44	291	92
ES	10	602	216	60	175	151
AU	19	477	127	100	180	70
TR	7	384	36	113	124	111
CZ	2	332	5	89	107	131
NL	5	233	83	51	65	34
BE	4	136	51	32	35	18
PT	2	71	11	18	25	17
KW	1	69	7	32	28	2
CH	3	64	0	42	21	1
US	1	52	13	9	24	6
LB	1	44	3	23	16	2
TN	1	44	1	17	8	18
AR	5	42	8	8	19	7
IR	2	38	11	27	0	0
GB	3	26	14	1	8	3
CU	1	25	6	4	12	3
BR	3	15	8	3	3	1
SA	1	12	2	8	1	1
EG	2	12	0	10	0	2
HU	4	9	0	2	6	1
OM	2	9	0	3	4	2
IL	1	6	0	0	1	5
DK	1	5	1	1	0	3
IN	2	4	1	2	1	0
RO	1	4	0	1	3	0
IE	1	3	3	0	0	0
GR	1	2	1	0	0	1
MT	1	2	1	1	0	0
MY	1	2	2	0	0	0
MK	1	2	0	0	1	1
NZ	1	1	0	1	0	0
Total	107	4447	935	937	1541	1034

Each country is indicated by its ISO 3166-1 alpha-2 code.

1
2 **Supplementary table 1.2.**
3 **Relapse characteristics of included patients**
4

	PPMS-A (n = 937 patients)	SPMS-A (n = 1034 patients)
Patient characteristics		
Patients with relapse during follow-up; No. (%)	336 (36)	948 (92)
Annualized relapse rate; median [IQR]	0.26 [0.14–0.48]	0.34 [0.19–0.60]
Relapse characteristics (across patients)		
Relapses recorded	743	2117
Relapses with phenotype(s) recorded; No. (% of all relapses)	610 (82)	1759 (83)
Relapse phenotypes; number of relapses (% of relapses with phenotype(s) recorded)		
Pyramidal	450 (74)	1212 (69)
Sensory	206 (34)	575 (33)
Cerebellar	95 (16)	216 (12)
Brainstem	76 (12)	273 (16)
Bowel/Bladder	46 (8)	104 (6)
Visual	56 (9)	107 (6)
Neuropsychology	16 (3)	26 (1)

Annualized relapse rate is calculated for the subset of patients with one or more relapses during follow-up.

Abbreviations: No. = number; IQR = interquartile range.

Supplementary table 1.3.**Andersen-Gill model for hazard of confirmed disability accrual, adjusted for sex and age**

Variable	Hazard ratio (95% CI)	<i>P</i>
Phenotype, SPMS	0.75 (0.70–0.81)	< .001
Sex, male	1.16 (1.09–1.24)	< .001
Age, at start of epoch	1.00 (0.99–1.00)	.11

Treating center and patient identity were modelled as random effects, with identity nested within center.

Abbreviations: CI = confidence interval.

Supplementary table 1.4.**Andersen-Gill model for hazard of confirmed disability accrual, corrected for proportional hazards violation (EDSS score frequency)**

Variable	Hazard ratio (95% CI)	<i>P</i>	Schoenfeld residuals <i>P</i>
Phenotype, SPMS	0.82 (0.75–0.89)	< .001	.61
3Sex, male	1.19 (1.12–1.28)	< .001	.92
Age, at start of epoch	1.00 (1.00–1.00)	.99	.87
EDSS baseline, at start of epoch	0.96 (0.94–0.99)	.001	.65
DMT, % of epoch on treatment [†]	0.95 (0.93–0.98)	< .001	.64
IST, % of epoch on treatment [†]	0.93 (0.90–0.96)	< .001	.19
EDSS score frequency (weeks from onset of the progressive phase)			
0–4 weeks	2.41 (1.84–3.17)	< .001	
> 4–8 weeks	1.70 (1.55–1.86)	< .001	
> 8–24 weeks	1.35 (1.29–1.42)	< .001	.14
> 24 weeks	1.12 (1.10–1.13)	< .001	

The proportional hazards violation for EDSS score frequency was corrected by including a time-dependent coefficient for this covariate, specified by a step function. To facilitate inclusion of time-stratified coefficients, this model omits random effects, and accounts for within-patient correlations by including patient identity as a cluster term to obtain robust variances. For comparison, for the model in the main text comparing SPMS to PPMS without proportional hazards correction (**table 3**), Schoenfeld residuals *P* values are: phenotype, .61; sex, .96; age, .64; EDSS baseline, .68; DMT, .74; IST, .19; EDSS score frequency, < .001.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale; DMT = disease-modifying therapy; IST = immunosuppressant therapy.

[†] 25% increments.

Supplementary table 1.5.
Andersen-Gill model for hazard of confirmed disability accrual,
stratified by phenotype

Variable	PPMS		SPMS	
	Hazard ratio (95% CI)	<i>P</i>	Hazard ratio (95% CI)	<i>P</i>
Sex, male	1.21 (1.10–1.33)	< .001	1.12 (1.01–1.24)	.04
Age, at start of epoch	1.00 (1.00–1.00)	.97	1.00 (0.99–1.00)	.07
EDSS baseline, at start of epoch	0.92 (0.89–0.94)	< .001	0.94 (0.90–0.99)	.01
DMT, % of epoch on treatment†	0.97 (0.94–1.01)	.10	0.98 (0.95–1.01)	.21
IST, % of epoch on treatment†	0.91 (0.87–0.96)	< .001	1.00 (0.95–1.06)	.95
EDSS score freq., during epoch	1.15 (1.13–1.17)	< .001	1.15 (1.12–1.19)	< .001

Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale;
 DMT = disease-modifying therapy; IST = immunosuppressant therapy; freq. = frequency.

† 25% increments.

Supplementary table 1.6.
Andersen-Gill model for hazard of confirmed disability accrual,
stratified by phenotype, with activity–DMT interaction

Variable	PPMS		SPMS	
	Hazard ratio (95% CI)	<i>P</i>	Hazard ratio (95% CI)	<i>P</i>
Relapse activity	0.96 (0.86–1.09)	.56	0.93 (0.79–1.10)	.41
Sex, male	1.21 (1.10–1.32)	< .001	1.11 (1.00–1.23)	.05
Age, at start of epoch	1.00 (1.00–1.00)	.88	0.99 (0.99–1.00)	.02
EDSS baseline, at start of epoch	0.92 (0.89–0.94)	< .001	0.94 (0.89–0.98)	.006
DMT, % of epoch on treatment†	0.98 (0.93–1.04)	.60	1.00 (0.96–1.04)	.98
IST, % of epoch on treatment†	0.92 (0.88–0.96)	< .001	1.00 (0.95–1.06)	1.00
EDSS score freq., during epoch	1.15 (1.13–1.17)	< .001	1.16 (1.12–1.19)	< .001
Interaction, relapse activity × DMT	0.98 (0.92–1.05)	.60	0.96 (0.90–1.01)	.13

Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale;
 DMT = disease-modifying therapy; IST = immunosuppressant therapy; freq. = frequency..

† 25% increments.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Supplementary table 1.7.
Sensitivity analyses—comparison of PPMS and SPMS

Sensitivity analysis (number of patients included)	Confirmed disability accrual		Age, years (95% CI)		
	Hazard ratio (95% CI) SPMS vs. PPMS (reference)	<i>P</i>		Onset (median)	Confirmed EDSS ≥ 7 (25th percentile)
Onset since July 1, 2004 (955 PPMS, 2297 SPMS)	0.78 (0.69–0.88)	< .001	PPMS	45.3 (44.5–46.2)	66.8 (64.4–70.8)
			SPMS	47.1 (46.6–47.6)	65.8 (63.1–68.4)
EDSS score every ≤ 15 months (1602 PPMS, 2315 SPMS)	0.79 (0.70–0.88)	< .001	PPMS	43.9 (43.3–44.5)	59.8 (58.1–61.7)
			SPMS	46.6 (45.9–47.1)	60.7 (58.3–62.8)
365-day EDSS score confirmation (1734 PPMS, 2320 SPMS)	0.86 (0.78–0.94)	.002	PPMS	43.9 (43.3–44.4)	60.6 (59.4–62.8)
			SPMS	46.6 (46.0–47.2)	62.6 (60.3–64.2)
Period-specific hazard ratios					
0–5 years from onset (1096 PPMS, 2575 SPMS)	0.88 (0.77–1.01)	.08		--	--
5–10 years from onset (1285 PPMS, 967 SPMS)	0.81 (0.71–0.93)	.002		--	--
10–15 years from onset (938 PPMS, 282 SPMS)	0.78 (0.62–0.97)	.03		--	--
Disability accrual from EDSS 4–5 (876 PPMS, 1585 SPMS)	0.77 (0.69–0.85)	< .001	PPMS	44.2 (43.5–45.0)	--
			SPMS	46.6 (45.7–47.2)	--

'Onset' refers to onset of the progressive phase. Disability accrual is assessed from onset of the progressive phase, except in the fifth sensitivity analysis, in which it is assessed from EDSS 4–5. Hazard ratios for confirmed disability accrual are adjusted for sex and (as time-dependent covariates) age at the start of each epoch; EDSS score at the start of each epoch; the proportion of each epoch receiving any disease-modifying therapy; the proportion of each epoch receiving any immunosuppressant therapy; and the annualized frequency of EDSS scores during each epoch. Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency. Ages at onset were estimated using the Kaplan-Meier estimator, and ages at EDSS ≥ 7 using the Turnbull estimator, as for the main analyses.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.

Supplementary table 1.8.

Sensitivity analyses—comparison of PPMS-N, PPMS-A, SPMS-N, and SPMS-A

Sensitivity analysis (number of patients included)	Confirmed disability accrual		
		Hazard ratio (95% CI) vs. PPMS-N (reference)	<i>P</i>
Onset since July 1, 2004 (478 PPMS-N, 477 PPMS-A, 1426 SPMS-N, 871 SPMS-A)	PPMS-A	1.04 (0.89–1.21)	.64
	SPMS-N	0.85 (0.73–1.00)	.04
	SPMS-A	0.74 (0.63–0.86)	< .001
EDSS score every ≤ 15 months (773 PPMS-N, 829 PPMS-A, 1345 SPMS-N, 970 SPMS-A)	PPMS-A	1.08 (0.95–1.23)	.23
	SPMS-N	0.88 (0.75–1.02)	.08
	SPMS-A	0.79 (0.68–0.91)	.001
365-day EDSS score confirmation (855 PPMS-N, 879 PPMS-A, 1326 SPMS-N, 994 SPMS-A)	PPMS-A	0.98 (0.88–1.09)	.64
	SPMS-N	0.95 (0.84–1.08)	.46
	SPMS-A	0.77 (0.67–0.87)	< .001
Period-specific hazard ratios			
0–5 years from onset (548 PPMS-N, 548 PPMS-A, 1541 SPMS-N, 1034 SPMS-A)	PPMS-A	0.90 (0.75–1.08)	.26
	SPMS-N	0.92 (0.78–1.10)	.37
	SPMS-A	0.76 (0.63–0.91)	.002
5–10 years from onset (637 PPMS-N, 648 PPMS-A, 420 SPMS-N, 547 SPMS-A)	PPMS-A	1.04 (0.90–1.21)	.59
	SPMS-N	0.85 (0.70–1.03)	.10
	SPMS-A	0.82 (0.68–0.98)	.03
10–15 years from onset (449 PPMS-N, 489 PPMS-A, 92 SPMS-N, 190 SPMS-A)	PPMS-A	0.92 (0.77–1.08)	.30
	SPMS-N	1.03 (0.74–1.43)	.85
	SPMS-A	0.61 (0.45–0.82)	< .001
Disability accrual from EDSS 4–5 (426 PPMS-N, 450 PPMS-A, 932 SPMS-N, 653 SPMS-A)	PPMS-A	1.01 (0.87–1.17)	.91
	SPMS-N	0.81 (0.70–0.94)	0.007
	SPMS-A	0.73 (0.63–0.85)	< .001

'Onset' refers to onset of the progressive phase. Disability accrual is assessed from onset of the progressive phase, except in the fifth sensitivity analysis, in which it is assessed from EDSS 4–5. Hazard ratios for confirmed disability accrual are adjusted for sex and (as time-dependent covariates) age at the start of each epoch; EDSS score at the start of each epoch; the proportion of each epoch receiving any disease-modifying therapy; the proportion of each epoch receiving any immunosuppressant therapy; and the annualized frequency of EDSS scores during each epoch. Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.

Supplement 2. Alternative analyses: SPMS identified by physician diagnosis (since Jan 1, 1995)

Harding-Forrester S, Roos I, Nguyen A-L, et al. Disability accrual in primary and secondary progressive multiple sclerosis.

Supplementary figure 2.1. Flow diagram of patient inclusion

Supplementary figure 2.2. Dates of onset of the progressive phase and of inclusion

Supplementary table 2.1. Included patients and treating centers, by country

Supplementary table 2.2. Clinical characteristics of included patients

Supplementary table 2.3. Relapse characteristics of included patients

Supplementary table 2.4. Ages at onset of the progressive phase and at confirmed EDSS ≥ 7

Supplementary table 2.5. Andersen-Gill model for hazard of confirmed disability accrual, adjusted for sex and age

Supplementary table 2.6. Andersen-Gill models for hazard of confirmed disability accrual, fully adjusted

Supplementary table 2.7. Andersen-Gill model for hazard of confirmed disability accrual, stratified by phenotype

Supplementary table 2.8. Andersen-Gill model for hazard of confirmed disability accrual, stratified by phenotype, with activity–DMT interaction

Supplementary table 2.9. Ages at onset of the progressive phase and at confirmed EDSS scores, by time period

Supplementary figure 2.3. Nelson-Aalen cumulative hazard curves for confirmed disability accrual

Supplementary figure 2.4. Turnbull survival curves for confirmed EDSS ≥ 7 (wheelchair dependence)

Supplementary figure 2.5. Mean EDSS trajectories in PPMS-N, PPMS-A, SPMS-N, and SPMS-A

Supplementary table 2.10. Sensitivity analyses—comparison of PPMS and SPMS

Supplementary table 2.11. Sensitivity analyses—comparison of PPMS-N, PPMS-A, SPMS-N, and SPMS-A

Abbreviations


MS = multiple sclerosis

PPMS = primary progressive MS; PPMS-N = PPMS with no superimposed relapse activity; PPMS-A = PPMS with superimposed relapse activity

RRMS = relapsing-remitting MS


SPMS = secondary progressive MS; SPMS-N = SPMS with no superimposed relapse activity; SPMS-A = SPMS with superimposed relapse activity

Supplementary figure 2.1. SPMS identified by physician diagnosis.
Flow diagram of patient inclusion



49730 patients with RRMS, and without a physician diagnosis of SPMS, were excluded.

**Supplementary figure 2.2. SPMS identified by physician diagnosis.
Dates of onset of the progressive phase and of inclusion**



White regions in histogram bars indicate patients excluded by the requirement for onset of the progressive phase since January 1, 1995, who would otherwise have been included.

**Supplementary table 2.1. SPMS identified by physician diagnosis.
Included patients and treating centers, by country**

Country	Centers	Patients (n)				
		Total	PPMS-N	PPMS-A	SPMS-N	SPMS-A
IT	11	1384	146	235	407	596
CA	7	969	178	44	448	299
AU	21	793	127	100	305	261
ES	10	780	216	60	254	250
TR	7	534	36	113	106	279
NL	6	370	83	51	103	133
CZ	2	335	5	89	111	130
US	2	295	13	9	194	79
BE	3	227	51	32	72	72
KW	1	127	7	32	70	18
PT	2	111	11	18	46	36
LB	1	99	3	23	45	28
CH	3	99	0	42	37	20
TN	1	60	1	17	19	23
GB	3	53	14	1	29	9
AR	5	49	8	8	21	12
EG	3	39	0	10	8	21
IR	2	38	11	27	0	0
CU	1	27	6	4	12	5
BR	2	18	8	3	6	1
IE	1	16	3	0	10	3
SA	1	15	2	8	2	3
OM	2	9	0	3	4	2
IN	2	8	1	2	3	2
IL	1	7	0	0	2	5
DK	1	5	1	1	0	3
RO	1	4	0	1	2	1
MT	1	3	1	1	0	1
HU	1	3	0	2	1	0
MY	1	2	2	0	0	0
NZ	2	2	0	1	1	0
GR	1	1	1	0	0	0
Total	108	6482	935	937	2318	2292

Each country is indicated by its ISO 3166-1 alpha-2 code.

Supplementary table 2.2. SPMS identified by physician diagnosis. Clinical characteristics of included patients

Characteristics	PPMS (all)	PPMS-N	PPMS-A	SPMS (all)	SPMS-N	SPMS-A
Patients	1872	935	937	4610	2318	2292
Sex, male (%)	878 (47)	442 (47)	436 (47)	1485 (32)	770 (33)	715 (31)
Age, MS onset	43.5 ± 10.5	45.6 ± 9.8	41.5 ± 10.7	31.9 ± 10.2	32.9 ± 10.6	30.9 ± 9.8
Age, progressive phase	43.5 ± 10.5	45.6 ± 9.8	41.5 ± 10.7	46.4 ± 10.4	48.5 ± 10.5	44.3 ± 9.7
Age, inclusion	48.7 ± 10.8	50.7 ± 10.1	46.7 ± 11.2	48.3 ± 10.5	50.4 ± 10.6	46.2 ± 9.9
Time, MS onset to inclusion; years	4.1 [2.0–7.3]	4.1 [2.1–7.3]	4.1 [1.9–7.3]	15.0 [9.2–22.1]	16.0 [9.9–23.7]	14.0 [8.6–20.3]
Time, follow-up; years	4.2 [1.9–7.9]	3.8 [1.7–7.5]	4.4 [2.1–8.2]	5.1 [2.3–8.9]	4.1 [1.9–7.3]	6.5 [3.0–10.3]
EDSS score frequency, annualized	1.53 [1.00–2.28]	1.37 [0.92–2.10]	1.68 [1.12–2.44]	1.44 [0.95–2.10]	1.30 [0.86–1.95]	1.59 [1.06–2.27]
Disability at inclusion; EDSS	4.0 [3.0–6.0]	4.0 [3.0–6.0]	4.0 [3.0–6.0]	5.5 [4.0–6.5]	6.0 [4.0–6.5]	5.5 [4.0–6.0]
Disability at censoring; EDSS	6.0 [4.5–6.5]	6.0 [4.5–6.5]	6.0 [4.0–6.5]	6.5 [5.5–7.0]	6.5 [5.5–7.0]	6.5 [5.5–7.0]
Disability increase, annualized; EDSS	0.15 [0.00–0.38]	0.15 [0.00–0.37]	0.15 [0.00–0.39]	0.08 [0.00–0.24]	0.02 [0.00–0.24]	0.10 [0.00–0.24]
Deaths recorded, from any cause (%)	59 (3)	33 (4)	26 (3)	212 (5)	127 (5)	85 (4)
Relapses during follow-up						
Patients (%)	336 (18)	0 (0)	336 (36)	1744 (38)	0 (0)	1744 (76)
Annualized relapse rate	0.26 [0.14–0.48]	NA	0.26 [0.14–0.48]	0.27 [0.14–0.52]	NA	0.27 [0.14–0.52]
Cerebrospinal fluid oligoclonal bands; patients (%)						
Present	1149 (61)	546 (58)	603 (64)	2124 (46)	986 (43)	1138 (50)
Absent	89 (5)	46 (5)	43 (5)	102 (2)	42 (2)	60 (3)
Not assessed	634 (34)	343 (37)	291 (31)	2384 (52)	1290 (56)	1094 (48)
Disease-modifying therapy, proportion of follow-up receiving treatment; patients (%)						
0%	1104 (59)	652 (70)	452 (48)	1623 (35)	1001 (43)	622 (27)
> 0–25%	221 (12)	105 (11)	116 (12)	549 (12)	235 (10)	314 (14)
> 25–50%	144 (8)	46 (5)	98 (10)	401 (9)	161 (7)	240 (10)
> 50–75%	106 (6)	36 (4)	70 (7)	384 (8)	137 (6)	247 (11)
> 75%	297 (16)	96 (10)	201 (21)	1653 (36)	784 (34)	869 (38)
Disease-modifying therapy, exposure to specific agents during follow-up; patients (%)						
Interferon beta	318 (17)	85 (9)	233 (25)	1787 (39)	728 (31)	1059 (46)
Glatiramer acetate	154 (8)	46 (5)	108 (12)	687 (15)	260 (11)	427 (19)
Fingolimod	118 (6)	39 (4)	79 (8)	483 (10)	169 (7)	314 (14)
Teriflunomide	28 (1)	7 (1)	21 (2)	146 (3)	56 (2)	90 (4)
Dimethyl fumarate	24 (1)	8 (1)	16 (2)	162 (4)	62 (3)	100 (4)
Cladribine	2 (0)	0 (0)	2 (0)	36 (1)	10 (0)	26 (1)
Mitoxantrone	91 (5)	29 (3)	62 (7)	507 (11)	168 (7)	339 (15)
Natalizumab	83 (4)	27 (3)	56 (6)	457 (10)	182 (8)	275 (12)
Alemtuzumab	12 (1)	3 (0)	9 (1)	44 (1)	10 (0)	34 (1)
Daclizumab	1 (0)	0 (0)	1 (0)	1 (0)	1 (0)	0 (0)
Rituximab	62 (3)	16 (2)	46 (5)	108 (2)	57 (2)	51 (2)
Ocrelizumab	169 (9)	90 (10)	79 (8)	83 (2)	39 (2)	44 (2)
Siponimod	0 (0)	0 (0)	0 (0)	10 (0)	8 (0)	2 (0)
ASCT	4 (0)	0 (0)	4 (0)	22 (0)	12 (1)	10 (0)
Immunosuppressant therapy, exposure to specific agents during follow-up; patients (%)						
Azathioprine	233 (12)	61 (7)	172 (18)	493 (11)	157 (7)	336 (15)
Methotrexate	121 (6)	41 (4)	80 (9)	275 (6)	86 (4)	189 (8)
Cyclophosphamide	94 (5)	26 (3)	68 (7)	244 (5)	79 (3)	165 (7)
Mycophenolate mofetil	5 (0)	2 (0)	3 (0)	22 (0)	11 (0)	11 (0)
Pregnancy during follow-up						
Patients (%)	15 (1)	2 (0)	13 (1)	36 (1)	8 (0)	28 (1)

Values are number (%), mean ± standard deviation, or median [interquartile range]. Patients classified as having PPMS-A include all eligible patients with a recorded diagnosis of progressive-relapsing MS (n = 540 patients; 58%). Patients classified as having SPMS under physician diagnosis include 2458 patients (53%) with SPMS under the operationalized diagnostic criteria. Annualized relapse rate is calculated for the subset of patients with one or more relapses during follow-up. Abbreviations: EDSS = Expanded Disability Status Scale; ASCT = autologous stem-cell transplant, assumed effective for 5 years.

**Supplementary table 2.3. SPMS identified by physician diagnosis.
Relapse characteristics of included patients**

	PPMS-A (n = 937 patients)	SPMS-A (n = 2292 patients)
Patient characteristics		
Patients with relapse during follow-up; No. (%)	336 (36)	1744 (76)
Annualized relapse rate; median [IQR]	0.26 [0.14–0.48]	0.27 [0.14–0.52]
Relapse characteristics (across patients)		
Relapses recorded	743	4096
Relapses with phenotype(s) recorded; No. (% of all relapses)	610 (82)	3206 (78)
Relapse phenotypes; number of relapses (% of relapses with phenotype(s) recorded)		
Pyramidal	450 (74)	2200 (69)
Sensory	206 (34)	1026 (32)
Cerebellar	95 (16)	430 (13)
Brainstem	76 (12)	461 (14)
Bowel/Bladder	46 (8)	224 (7)
Visual	56 (9)	242 (8)
Neuropsychology	16 (3)	64 (2)

Annualized relapse rate is calculated for the subset of patients with one or more relapses during follow-up.

Abbreviations: No. = number; IQR = interquartile range.

**Supplementary table 2.4. SPMS identified by physician diagnosis.
Ages at onset of the progressive phase and at confirmed EDSS \geq 7**

Phenotype	All patients	Female	Male
Age at onset of the progressive phase (median), years (95% CI)			
PPMS	43.9 (43.3–44.4)	44.4 (43.8–45.2)	43.0 (42.3–44.0)
PPMS-N	46.0 (45.2–46.8)	46.5 (45.3–47.7)	45.5 (44.5–46.7)
PPMS-A	41.9 (40.7–42.6)	42.8 (42.0–43.7)	40.0 (38.6–41.8)
SPMS	46.3 (45.8–46.7)	46.8 (46.3–47.4)	45.0 (44.3–45.8)
SPMS-N	48.4 (47.8–49.1)	48.9 (48.3–49.8)	47.3 (46.2–48.4)
SPMS-A	44.2 (43.7–44.8)	44.8 (44.3–45.5)	43.0 (42.1–43.8)
Age at confirmed EDSS \geq 7 (25th percentile), years (95% CI)			
PPMS	60.3 (58.8–62.5)	63.4 (60.5–66.2)	58.1 (56.6–60.1)
PPMS-N	62.5 (60.1–65.8)	66.0 (62.5–68.7)	60.1 (57.3–62.7)
PPMS-A	57.6 (55.4–60.9)	60.9 (56.3–65.3)	55.8 (52.8–58.6)
SPMS	54.4 (53.7–55.2)	55.1 (54.1–56.1)	53.1 (50.8–54.5)
SPMS-N	55.4 (54.1–56.8)	55.7 (54.1–57.1)	54.8 (52.2–57.5)
SPMS-A	53.7 (52.4–54.8)	54.7 (53.5–55.8)	51.2 (49.2–53.4)

Ages at onset of the progressive phase (Kaplan-Meier estimator; log-rank test) were younger in PPMS versus SPMS ($P < .001$), in subgroups with activity (-A) versus those without (-N) ($P < .001$ in both PPMS and SPMS), and in males versus females ($P = .22$ in PPMS, $P = .02$ in SPMS). Ages at confirmed EDSS \geq 7 (Turnbull estimator; generalized log-rank test) were younger in SPMS versus PPMS ($P < .001$; among females, $P < .001$; among males, $P = .002$; among patients with activity, $P < .001$; without activity, $P < .001$), and younger in patients with activity than in those without (PPMS-A vs. PPMS-N, $P = .002$; SPMS-A vs. SPMS-N, $P = .004$).

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.

**Supplementary table 2.5. SPMS identified by physician diagnosis.
Andersen-Gill model for hazard of confirmed disability accrual,
adjusted for sex and age**

Variable	Hazard ratio (95% CI)	P
Phenotype, SPMS	0.77 (0.73–0.82)	< .001
Sex, male	1.12 (1.07–1.18)	< .001
Age, at start of epoch	0.99 (0.99–1.00)	< .001

Treating center and patient identity were modelled as random effects, with identity nested within center.

Abbreviations: CI = confidence interval.

**Supplementary table 2.6. SPMS identified by physician diagnosis.
Andersen-Gill models for hazard of confirmed disability accrual,
fully adjusted**

Variable	Hazard ratio (95% CI)	P
Model comparing SPMS to PPMS (reference)		
Phenotype, SPMS	0.89 (0.83–0.95)	< .001
Sex, male	1.12 (1.06–1.18)	< .001
Age, at start of epoch	1.00 (0.99–1.00)	.002
EDSS baseline, at start of epoch	0.89 (0.87–0.90)	< .001
DMT, % of epoch on treatment†	1.01 (0.99–1.02)	.56
IST, % of epoch on treatment†	0.98 (0.95–1.01)	.14
EDSS score freq., during epoch	1.07 (1.06–1.08)	< .001
Model comparing PPMS-A, SPMS-N, and SPMS-A to PPMS-N (reference)		
Phenotype		
PPMS-A	0.92 (0.83–1.02)	.11
SPMS-N	0.86 (0.79–0.94)	.001
SPMS-A	0.85 (0.78–0.93)	< .001
Sex, male	1.11 (1.05–1.18)	< .001
Age, at start of epoch	1.00 (0.99–1.00)	.001
EDSS baseline, at start of epoch	0.89 (0.87–0.90)	< .001
DMT, % of epoch on treatment†	1.01 (0.99–1.02)	.45
IST, % of epoch on treatment†	0.98 (0.95–1.01)	.17
EDSS score freq., during epoch	1.07 (1.06–1.08)	< .001

Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale; DMT = disease-modifying therapy; IST = immunosuppressant therapy; freq. = frequency.

† 25% increments.

**Supplementary table 2.7. SPMS identified by physician diagnosis.
Andersen-Gill model for hazard of confirmed disability accrual,
stratified by phenotype**

Variable	PPMS		SPMS	
	Hazard ratio (95% CI)	<i>P</i>	Hazard ratio (95% CI)	<i>P</i>
Sex, male	1.21 (1.10–1.33)	< .001	1.07 (0.99–1.14)	.07
Age, at start of epoch	1.00 (1.00–1.00)	.97	0.99 (0.99–1.00)	< .001
EDSS baseline, at start of epoch	0.92 (0.89–0.94)	< .001	0.87 (0.85–0.89)	< .001
DMT, % of epoch on treatment†	0.97 (0.94–1.01)	.10	1.02 (1.00–1.04)	.11
IST, % of epoch on treatment†	0.91 (0.87–0.96)	< .001	1.01 (0.98–1.04)	.58
EDSS score freq., during epoch	1.15 (1.13–1.17)	< .001	1.05 (1.03–1.07)	< .001

Treating center and patient identity were modelled as random effects, with identity nested within center.
The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale;
DMT = disease-modifying therapy; IST = immunosuppressant therapy; freq. = frequency.

† 25% increments.

**Supplementary table 2.8. SPMS identified by physician diagnosis.
Andersen-Gill model for hazard of confirmed disability accrual,
stratified by phenotype, with activity–DMT interaction**

Variable	PPMS		SPMS	
	Hazard ratio (95% CI)	<i>P</i>	Hazard ratio (95% CI)	<i>P</i>
Relapse activity	0.96 (0.86–1.09)	.56	1.10 (1.00–1.21)	.05
Sex, male	1.21 (1.10–1.32)	< .001	1.06 (0.99–1.14)	.10
Age, at start of epoch	1.00 (1.00–1.00)	.88	0.99 (0.99–1.00)	< .001
EDSS baseline, at start of epoch	0.92 (0.89–0.94)	< .001	0.86 (0.84–0.88)	< .001
DMT, % of epoch on treatment†	0.98 (0.93–1.04)	.60	1.06 (1.03–1.09)	< .001
IST, % of epoch on treatment†	0.92 (0.88–0.96)	< .001	1.01 (0.97–1.04)	.66
EDSS score freq., during epoch	1.15 (1.13–1.17)	< .001	1.05 (1.03–1.07)	< .001
Interaction, relapse activity × DMT	0.98 (0.92–1.05)	.60	0.93 (0.90–0.97)	< .001

Treating center and patient identity were modelled as random effects, with identity nested within center.
The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale;
DMT = disease-modifying therapy; IST = immunosuppressant therapy; freq. = frequency.

† 25% increments.

**Supplementary table 2.9. SPMS identified by physician diagnosis.
Ages at onset of the progressive phase and at confirmed EDSS scores,
by time period**


	Median age at progressive multiple sclerosis milestones, years (95% CI)			
	Onset	EDSS ≥ 4	EDSS ≥ 6	EDSS ≥ 7
Complete dataset (1995–2020)				
PPMS (<i>n</i> = 1872)	43.9 (43.3–44.4)	50.4 (49.6–51.1)	56.9 (55.8–57.4)	72.8 (70.8–76.4)
PPMS-N (<i>n</i> = 935)	46.0 (45.2–46.8)	51.9 (50.8–53.2)	57.5 (56.9–59.0)	75.4 (71.0–NA)
PPMS-A (<i>n</i> = 937)	41.9 (40.7–42.6)	47.7 (46.2–49.6)	54.8 (53.6–56.6)	70.8 (68.9–NA)
SPMS (<i>n</i> = 4610)	46.3 (45.8–46.7)	43.6 (43.0–44.2)	50.4 (50.0–50.8)	69.3 (67.5–70.6)
SPMS-N (<i>n</i> = 2318)	48.4 (47.8–49.1)	45.8 (44.9–46.5)	51.9 (51.3–52.5)	71.3 (70.0–73.3)
SPMS-A (<i>n</i> = 2292)	44.2 (43.7–44.8)	41.8 (41.2–42.5)	48.8 (47.9–49.8)	66.9 (65.4–69.3)
1995–2003				
PPMS (<i>n</i> = 125)	41.9 (39.3–44.9)	47.0 (45.8–50.9)	51.9 (50.0–55.2)	60.6 (56.0–65.0)
PPMS-N (<i>n</i> = 56)	44.5 (41.5–48.7)	50.7 (47.9–53.5)	54.3 (51.0–57.5)	62.2 (56.0–68.5)
PPMS-A (<i>n</i> = 69)	41.0 (37.5–43.7)	44.4 (43.3–50.8)	47.6 (46.2–54.5)	58.8 (55.0–65.4)
SPMS (<i>n</i> = 511)	44.9 (44.1–46.3)	42.0 (40.9–43.6)	47.1 (46.2–48.1)	56.4 (55.2–57.5)
SPMS-N (<i>n</i> = 183)	48.8 (47.1–50.6)	45.1 (42.0–47.4)	48.2 (46.1–51.5)	57.5 (57.1–60.9)
SPMS-A (<i>n</i> = 328)	43.8 (42.7–44.7)	41.2 (39.6–42.5)	46.5 (45.8–47.9)	55.4 (54.2–56.6)
2004–2011				
PPMS (<i>n</i> = 266)	44.4 (43.3–45.9)	49.3 (46.6–50.8)	53.4 (52.2–56.9)	65.1 (62.4–NA)
PPMS-N (<i>n</i> = 132)	46.5 (44.5–48.3)	50.5 (48.3–52.5)	54.8 (52.4–57.3)	67.5 (60.0–NA)
PPMS-A (<i>n</i> = 134)	42.8 (40.6–44.5)	46.4 (45.1–49.7)	53.0 (50.2–59.0)	65.1 (62.4–NA)
SPMS (<i>n</i> = 1269)	47.3 (46.6–48.1)	44.6 (43.6–45.5)	50.8 (50.2–51.7)	64.2 (62.3–66.2)
SPMS-N (<i>n</i> = 575)	49.7 (48.7–50.9)	46.6 (45.9–48.2)	52.0 (51.3–53.5)	65.9 (62.9–70.8)
SPMS-A (<i>n</i> = 694)	45.0 (44.2–46.0)	43.0 (42.1–44.1)	50.0 (48.4–50.7)	62.1 (60.8–64.6)
2012–2020 (ending January 7, 2020)				
PPMS (<i>n</i> = 250)	47.6 (45.9–49.8)	54.2 (52.6–59.1)	63.3 (61.1–70.6)	80.8 (NA–NA)
PPMS-N (<i>n</i> = 129)	48.3 (47.2–51.5)	57.8 (52.6–60.8)	64.8 (60.8–NA)	NA (71.9–NA)
PPMS-A (<i>n</i> = 121)	45.5 (42.8–50.2)	54.2 (50.3–58.7)	62.3 (58.8–NA)	80.8 (NA–NA)
SPMS (<i>n</i> = 912)	47.5 (46.6–48.4)	44.6 (43.5–45.9)	52.7 (51.5–54.1)	87.8 (NA–NA)
SPMS-N (<i>n</i> = 582)	48.8 (47.7–49.8)	46.4 (44.6–47.9)	53.9 (52.6–56.0)	87.8 (NA–NA)
SPMS-A (<i>n</i> = 330)	45.5 (43.7–46.6)	42.2 (41.0–44.3)	50.1 (48.8–52.7)	75.7 (NA–NA)

Median ages at onset of the progressive phase (Kaplan-Meier estimator) and at confirmed EDSS scores ≥ 4 , 6, and 7 (Turnbull estimator), for the complete dataset and for three constituent time periods. Analyses for each time period include only those patients with onset of PPMS or (physician-diagnosed) SPMS during that period, and only EDSS scores recorded by the end of that period (such that the combined number of patients across the three periods is less than the total number of patients in the complete dataset). Note that in some cases data were insufficient for complete Turnbull estimates of the median ages at EDSS ≥ 6 or 7.


Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.

**Supplementary figure 2.3. SPMS identified by physician diagnosis.
Nelson-Aalen cumulative hazard curves for confirmed disability accrual**

A



B



Number at risk

PPMS-N	24	260	384	375	315	262	163	91	31	10
PPMS-A	14	278	409	411	339	269	184	115	55	8
SPMS-N	934	1294	1017	720	478	311	165	84	30	7
SPMS-A	805	1494	1410	1133	813	508	282	145	52	8


Cumulative exits (events)

PPMS-N	0 (0)	38 (67)	164 (274)	336 (505)	486 (657)	607 (800)	740 (904)	834 (956)	902 (988)	925 (994)
PPMS-A	0 (0)	30 (88)	139 (274)	299 (494)	448 (697)	598 (834)	719 (919)	808 (984)	880 (1025)	929 (1039)
SPMS-N	0 (0)	487 (552)	990 (1011)	1417 (1343)	1738 (1555)	1956 (1690)	2117 (1780)	2224 (1818)	2284 (1832)	2311 (1846)
SPMS-A	0 (0)	231 (553)	584 (1190)	996 (1746)	1381 (2141)	1730 (2373)	1989 (2508)	2143 (2573)	2240 (2598)	2284 (2604)

(A) PPMS and SPMS. (B) PPMS-N, PPMS-A, SPMS-N, and SPMS-A. The cumulative hazard indicates the expected number of confirmed disability accrual events for a patient observed for a given duration of time. Shaded regions indicate 95% confidence intervals. A-G HR = adjusted hazard ratio obtained with the Andersen-Gill model (see supplementary table 2.6); CI = confidence interval.

Supplementary figure 2.4. SPMS identified by physician diagnosis.
Turnbull survival curves for confirmed EDSS ≥ 7 (wheelchair dependence)


A



Numbers: at risk [in interval] (cumulative events)

PPMS	1872 0	1824 [13](5)	1632 [19](49)	1211 [26](140)	582 [17](245)	183 [2](316)	86 [0](328)
SPMS	4563 [44](3)	4376 [174](24)	3910 [222](163)	2797 [155](523)	1474 [74](870)	587 [12](1045)	354 [0](1085)

B




Numbers: at risk [in interval] (cumulative events)

PPMS-N	935 0	919 [7](2)	853 [7](16)	670 [14](54)	339 [9](110)	98 [0](150)	39 [0](157)
PPMS-A	937 0	905 [6](3)	779 [12](33)	541 [12](86)	243 [8](135)	85 [2](166)	47 [0](171)
SPMS-N	2287 [30](1)	2178 [114](8)	1958 [151](65)	1489 [115](217)	852 [60](375)	381 [11](460)	242 [0](484)
SPMS-A	2276 [14](2)	2198 [60](16)	1952 [71](98)	1308 [40](306)	622 [14](495)	206 [1](585)	112 [0](601)

(A) PPMS and SPMS. (B) PPMS-N, PPMS-A, SPMS-N, and SPMS-A. Patients who entered observation having already reached confirmed EDSS ≥ 7 are analyzed as interval-censored observations, with EDSS ≥ 7 reached in the interval (date of MS onset, date of first observation) ($n = 37$ PPMS-N, 43 PPMS-A, 227 SPMS-N, 107 SPMS-A). Among patients with SPMS, 120 reached EDSS ≥ 7 at the date of onset of the progressive phase (93 SPMS-N, 27 SPMS-A); 75 reached EDSS ≥ 7 prior to the progressive phase (50 SPMS-N, 25 SPMS-A). P values (generalized log-rank tests): SPMS vs. PPMS, $P < .001$; SPMS-N vs. PPMS-N, $P < .001$; SPMS-A vs. PPMS-A, $P < .001$; PPMS-A vs. PPMS-N, $P = .002$; SPMS-A vs. SPMS-N, $P = .004$. Values in the risk table indicate the number of patients at risk at each age; the number of interval-censored patients with an interval including that age (square brackets); and the cumulative number of patients having reached confirmed EDSS ≥ 7 at that age (curved brackets). Shaded regions indicate 95% confidence intervals. EDSS = Expanded Disability Status Scale.

**Supplementary figure 2.5. SPMS identified by physician diagnosis.
Mean EDSS trajectories in PPMS-N, PPMS-A, SPMS-N, and SPMS-A**


A (EDSS scores in the progressive phase)



Number of patients (each 2-year interval)

PPMS-N	1	6	13	17	29	39	65	81	100	111	130	169	185	216	258	282	279	276	253	231	212	185	147	115	78	51	32	20	10	5	2
PPMS-A	12	27	43	61	77	91	111	135	151	194	216	228	237	245	253	258	241	233	221	189	145	121	100	69	49	44	23	16	12	6	1
SPMS-N	11	21	33	44	68	108	154	194	253	309	399	457	522	576	608	647	624	647	599	556	482	428	354	275	203	154	100	69	42	18	13
SPMS-A	21	34	56	86	137	205	285	370	475	581	651	720	750	768	800	805	751	697	646	555	450	364	287	212	148	89	55	31	21	10	5

B (EDSS scores from MS onset)



Number of patients (each 2-year interval)

PPMS-N	0	1	1	6	13	17	29	39	65	81	100	111	130	169	185	216	258	282	279	276	253	231	212	185	147	115	78	51	32	20	10	5	2
PPMS-A	2	6	12	27	43	61	77	91	111	135	151	194	216	228	237	245	253	258	241	233	221	189	145	121	100	69	49	44	23	16	12	6	1
SPMS-N	11	23	41	62	96	145	177	248	314	362	443	512	600	675	737	764	795	803	760	755	679	612	519	462	375	287	213	157	101	70	43	18	14
SPMS-A	26	36	67	99	150	217	297	395	496	595	693	779	855	899	920	931	907	896	812	744	675	584	468	377	293	217	151	91	55	31	21	10	5

Mean EDSS scores by age in PPMS-N, PPMS-A, SPMS-N, and SPMS-A, (A) for study-eligible EDSS scores, starting from onset of the progressive phase. (B) for all available EDSS scores, starting from MS onset (including RRMS). Mean scores are plotted for each 2-year age interval, for groups with 10 or more patients contributing data in that interval. Error bars indicate 95% confidence intervals. EDSS = Expanded Disability Status Scale.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

**Supplementary table 2.10. SPMS identified by physician diagnosis.
Sensitivity analyses—comparison of PPMS and SPMS**

Sensitivity analysis (number of patients included)	Confirmed disability accrual		Age, years (95% CI)		
	Hazard ratio (95% CI) SPMS vs. PPMS (reference)	<i>P</i>		Onset (median)	Confirmed EDSS ≥ 7 (25th percentile)
Onset since July 1, 2004 (955 PPMS, 2864 SPMS)	0.85 (0.77–0.93)	< .001	PPMS	45.3 (44.5–46.2)	66.8 (64.4–70.8)
			SPMS	47.2 (46.6–47.6)	58.3 (57.0–60.2)
EDSS score every ≤ 15 months (1602 PPMS, 3881 SPMS)	0.84 (0.77–0.91)	< .001	PPMS	43.9 (43.3–44.5)	59.8 (58.1–61.7)
			SPMS	46.1 (45.7–46.6)	54.1 (53.2–55.0)
365-day EDSS score confirmation (1734 PPMS, 4328 SPMS)	0.90 (0.84–0.97)	.005	PPMS	43.9 (43.3–44.4)	60.6 (59.4–62.8)
			SPMS	46.3 (45.8–46.8)	54.9 (54.0–55.8)
Period-specific hazard ratios					
0–5 years from onset (1096 PPMS, 4001 SPMS)	0.91 (0.85–1.08)	.51		--	--
5–10 years from onset (1285 PPMS, 2836 SPMS)	0.83 (0.73–0.92)	< .001		--	--
10–15 years from onset (938 PPMS, 1491 SPMS)	0.84 (0.73–0.96)	.003		--	--
Disability accrual from EDSS 4–5 (876 PPMS, 1639 SPMS)	0.84 (0.77–0.92)	< .001	PPMS	44.2 (43.5–45.0)	65.8 (63.7–68.2)
			SPMS	46.1 (45.4–46.8)	65.9 (64.1–70.4)

'Onset' refers to onset of the progressive phase. Disability accrual is assessed from onset of the progressive phase, except in the fifth sensitivity analysis, in which it is assessed from EDSS 4–5. Hazard ratios for confirmed disability accrual are adjusted for sex and (as time-dependent covariates) age at the start of each epoch; EDSS score at the start of each epoch; the proportion of each epoch receiving any disease-modifying therapy; the proportion of each epoch receiving any immunosuppressant therapy; and the annualized frequency of EDSS scores during each epoch. Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency. Ages at onset were estimated using the Kaplan-Meier estimator, and ages at EDSS ≥ 7 using the Turnbull estimator, as for the main analyses.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.

**Supplementary table 2.11. SPMS identified by physician diagnosis.
Sensitivity analyses—comparison of PPMS-N, PPMS-A, SPMS-N, and SPMS-A**

Sensitivity analysis (number of patients included)	Confirmed disability accrual		
		Hazard ratio (95% CI) vs. PPMS-N (reference)	<i>P</i>
Onset since July 1, 2004 (478 PPMS-N, 477 PPMS-A, 1534 SPMS-N, 1330 SPMS-A)	PPMS-A	0.98 (0.84–1.13)	.75
	SPMS-N	0.86 (0.75–0.98)	.02
	SPMS-A	0.82 (0.72–0.94)	.004
EDSS score every ≤ 15 months (773 PPMS-N, 829 PPMS-A, 1869 SPMS-N, 2012 SPMS-A)	PPMS-A	0.98 (0.87–1.11)	.80
	SPMS-N	0.85 (0.75–0.95)	.006
	SPMS-A	0.82 (0.73–0.92)	< .001
365-day EDSS score confirmation (855 PPMS-N, 879 PPMS-A, 2135 SPMS-N, 2193 SPMS-A)	PPMS-A	0.91 (0.82–1.01)	.09
	SPMS-N	0.88 (0.79–0.97)	.008
	SPMS-A	0.85 (0.77–0.94)	< .001
Period-specific hazard ratios			
0–5 years from onset (548 PPMS-N, 548 PPMS-A, 2007 SPMS-N, 1994 SPMS-A)	PPMS-A	0.84 (0.66–1.02)	.06
	SPMS-N	0.91 (0.76–1.07)	.25
	SPMS-A	0.85 (0.70–1.01)	.04
5–10 years from onset (637 PPMS-N, 648 PPMS-A, 1226 SPMS-N, 1610 SPMS-A)	PPMS-A	1.02 (0.87–1.17)	.80
	SPMS-N	0.78 (0.64–0.92)	< .001
	SPMS-A	0.87 (0.74–1.00)	.04
10–15 years from onset (449 PPMS-N, 489 PPMS-A, 580 SPMS-N, 911 SPMS-A)	PPMS-A	0.88 (0.72–1.04)	.11
	SPMS-N	0.82 (0.66–0.99)	.02
	SPMS-A	0.76 (0.61–0.92)	< .001
Disability accrual from EDSS 4–5 (426 PPMS-N, 450 PPMS-A, 750 SPMS-N, 889 SPMS-A)	PPMS-A	0.94 (0.82–1.09)	.42
	SPMS-N	0.81 (0.71–0.92)	.002
	SPMS-A	0.82 (0.72–0.93)	.002

'Onset' refers to onset of the progressive phase. Disability accrual is assessed from onset of the progressive phase, except in the fifth sensitivity analysis, in which it is assessed from EDSS 4–5. Hazard ratios for confirmed disability accrual are adjusted for sex and (as time-dependent covariates) age at the start of each epoch; EDSS score at the start of each epoch; the proportion of each epoch receiving any disease-modifying therapy; the proportion of each epoch receiving any immunosuppressant therapy; and the annualized frequency of EDSS scores during each epoch. Treating center and patient identity were modelled as random effects, with identity nested within center. The proportional hazards assumption was violated for EDSS score frequency.

Abbreviations: CI = confidence interval; EDSS = Expanded Disability Status Scale.