

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Cognitive Functions in Substance-Related and Addictive Disorders

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Bouchard, A.E., Garofalo, S., Rouillard, C., Fecteau, S. (2021). Cognitive Functions in Substance-Related and Addictive Disorders. Cham : Springer [10.1007/978-3-030-76136-3\_26].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/858863> since: 2022-02-15

*Published:*

DOI: [http://doi.org/10.1007/978-3-030-76136-3\\_26](http://doi.org/10.1007/978-3-030-76136-3_26)

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

## **Cognitive functions in substance-related and addictive disorders**

Amy E. Bouchard, MSc, Department of Psychiatry and Neurosciences, Faculty of Medicine, Université Laval; CERVO Brain Research Centre, Centre intégré universitaire en santé et services sociaux de la Capitale-Nationale, Quebec City, Quebec, Canada.

Sara Garofalo, PhD, Department of Psychology, University of Bologna, Psychometrics and Psychophysics Lab, Centre for Studies and Research in Cognitive Neuroscience, Bologna, Emilia-Romagna, Italy

Claude Rouillard, PhD, Department of Psychiatry and Neurosciences, Faculty of Medicine, Université Laval; Axe Neurosciences, Centre de Recherche du CHU de Québec, Quebec City, Quebec, Canada.

Shirley Fecteau, PhD, Department of Psychiatry and Neurosciences, Faculty of Medicine, Université Laval; CERVO Brain Research Centre, Centre intégré universitaire en santé et services sociaux de la Capitale-Nationale, Quebec City, Quebec, Canada.

## **Abstract**

Despite the availability of treatments such as medications and cognitive behavioural therapy, relapse remains high in substance-related and addictive disorders. Cognitive functions, including cognitive biases and executive functions, are significant predictors of relapse. Alleviating cognitive deficits is therefore promising to suppress craving and decrease relapse. Transcranial current stimulation holds potential to improve cognitive functioning in these patients. This chapter provides an overview of the use of transcranial current stimulation to modulate cognitive functions in substance-related and addictive disorders. We show that transcranial current stimulation can decrease cognitive biases and improve executive functions across various substance-related and addictive disorders, especially regarding decision-making and when targeting the dorsolateral prefrontal cortex bilaterally. We also discuss the relationship between cognitive functions and several processes, such as craving, mood, and stress, which should be taken into account when developing future transcranial current stimulation protocols. Lastly, we provide suggestions as to how future studies could be improved to better treat patients with these debilitating disorders.

**Keywords:** substance-related and addictive disorders, executive functions, cognitive bias, dorsolateral prefrontal cortex

## 1. Introduction

Substance-related and addictive disorders (SRADs), including alcohol, cannabis, gambling, and stimulant use disorders, are characterized by maladaptive behaviour or dysfunctional use of a substance that leads to clinically distressing consequences (e.g., craving, health issues, interference with work, school or personal life) [1]. SRADs are difficult to treat and relapse remains a big issue despite available pharmacological and behavioural treatments. Crucially, cognitive deficits (e.g., cognitive biases, deficits in executive functions) can predict relapse [2]. Hence, improving cognitive functions is a promising therapeutic option for dealing with craving and relapse [2]. Cognitive deficits can be present before the onset of SRADs and worsen with chronicity [3]. Yet, not all patients with SRADs present the same cognitive profile, as they can vary across diagnoses and as a function of comorbidities [2]. More specifically, a meta-analysis found that patients with alcohol and stimulant use disorders particularly present impaired cognitive flexibility; patients with cannabis and 3,4-methylenedioxy-methamphetamine (MDMA) use disorders predominantly display impairments in complex planning and processing speed; patients with opioid use disorder mostly demonstrate reasoning impairments, and patients with cannabis and methamphetamine use disorders mainly show memory deficits [4].

Within this context, transcranial current stimulation (tCS) over the dorsolateral prefrontal cortex (DLPFC) has been successfully used to strengthen cognitive functions [5-14] and help patients resist craving and avoid relapse. Given such evidence, an overview of which cognitive functions have been successfully improved in patients with SRADs can inform clinical practice and help develop new interventions. Hence, this chapter reviews studies that examined tCS-induced effects on cognitive functions relevant to SRADs, namely cognitive bias and executive functions. The relationship between cognitive functions and craving, mood, and stress is also discussed. All included studies are sham-controlled, randomized, blinded, and used transcranial direct current stimulation, unless otherwise stated (**Table 1**).

## 2. tCS effects in cognitive functions in SRADs

Several studies assessed the effects of tCS on cognitive functions in SRADs. These can be divided into two main categories: studies on implicit cognitive functions, e.g., cognitive bias, and studies on explicit cognitive functions, e.g., executive functions (see **Figure 1**; **Table 1**).

### 2.1 tCS effects on cognitive biases in SRADs

Some patients with SRADs are aware that their addictive behaviour is detrimental, yet they still carry it out despite the negative consequences. One way to explain this behaviour is by taking into account implicit cognitive functioning, such as cognitive biases. Cognitive biases are automatic, implicit, and favorable processing of certain stimuli (e.g., external cues) over others [15]. Two major forms are approach bias and attentional bias. Approach bias happens when patients are quicker to approach rather than avoid cues [15]. Attentional bias occurs when patients display biased attention towards cues, which can increase craving [16]. Seven studies assessed the effects of tCS on cognitive biases [5, 6, 13, 14, 17-19] in alcohol and methamphetamine users. Four of these studies found significant reductions in cognitive biases when targeting the bilateral DLPFC [5, 6, 13, 14], as well as the DLPFC and shoulder [6]. In particular, two studies found reduced approach biases in alcohol users when placing the anode over the right and cathode over the left DLPFC [5] and vice versa [13]. Also, one of these studies combined tCS with a cognitive bias modification protocol [13]. In addition, one study found decreased attentional biases in tobacco smokers when patients received real transcranial alternating current stimulation (tACS) paired with attentional bias modification as compared to sham tACS with attentional bias modification training, as shown by decreased time observing smoking-related stimuli measured with

an eye tracker [14]. Further, a single study reported decreased attentional bias towards drug cues in abstinent, treatment-seeking patients with methamphetamine use disorder [6]. Patients performed a probe detection task before and after they received two 13-minute tCS sessions. Patients were randomly assigned to one of six groups with different electrode montages: 1) anode over the left DLPFC, cathode over the right shoulder; 2) anode over the right DLPFC, cathode over the left shoulder; 3) anode over the left DLPFC, cathode over the right supraorbital ridge; 4) anode over the right DLPFC, cathode over the left supraorbital ridge; 5) anode over the left DLPFC, cathode over the right DLPFC. Sham condition consisted of electrodes over the right and left DLPFC. Of these, two groups displayed reduced attentional bias towards cues as measured by reaction times, that is, one group receiving anodal and cathodal tDCS over the left and right DLPFC, respectively, and one group receiving anodal and cathodal tDCS over the left DLPFC and the shoulder, respectively.

## 2.2 tCS effects on executive functions in SRADs

Higher order cognitive functions, such as executive functions, are believed to be impaired in SRADs [2, 3]. Some researchers purport that patients with SRADs have an imbalance between implicit and explicit processes, in which executive functions fail to control implicit urges. In line with this, a series of studies attempted to increase cognitive control to reduce addictive behaviour. Several studies have assessed the effects of tCS in SRADs on a wide range of executive functions, such as cognitive flexibility, decision-making, working memory, self-regulation, and selective attention (see **Figure 1; Table 1**).

Cognitive flexibility, also known as set-shifting, reflects the ability to adapt to different responses or situations [20]. Measuring cognitive flexibility can be a useful marker of cognitive control and possibly compulsivity [21]. Two studies reported improved cognitive flexibility following tCS over the DLPFC (anode over the left, cathode over the right DLPFC) in patients with methamphetamine use disorder [7] and gambling disorder (anode over the right, cathode over the left DLPFC) [8]. More specifically, patients showed decreased perseveration errors and/or completed categories on the Wisconsin Card Sorting Task [22].

Decision-making encompasses evaluating potential outcomes and selecting the most appropriate option [23]. This ability can be impaired in patients with SRADs, in that they show a tendency to choose immediate rewards (e.g., drug or monetary rewards) despite the possible detrimental consequences [24]. Up to now, six studies examined the effect on decision-making of tCS over the bilateral DLPFC across different SRADs, including cocaine [9], gambling [8], methamphetamine [7], tobacco [10, 11, 14], and cannabis [25] use disorders. Some studies applied the anode and cathode over the right and left DLPFC [8, 10], and vice versa [7], whereas some used both montages [9, 11, 25], and one used tACS to target both DLPFCs [14]. The first six studies reported improvements in various measures of decision-making (e.g., Balloon Analog Risk Task, Iowa Gambling Task, Game of Dice Task, Ultimatum Game, Columbia Card Task, Delay Discounting Task). The last study reported increased risky choices among patients with cannabis use disorder [25]. Nonetheless, this study demonstrated that these patients display different decision-making processes as compared to healthy individuals for the same task [26].

Working memory refers to the ability to store and use short-term information [3, 27] which can influence other processes, such as decision-making. For instance, working memory training decreases delay discounting in patients with stimulant use disorder [28]. Two studies evaluated the effects of tCS on working memory in patients with SRADs. The first one reported decreased response time and increased accuracy on the N-Back Task by applying tCS over the DLPFC (anode over the left and cathode over the right DLPFC) in methamphetamine use disorder [7]. The second study (which placed the electrodes over

the ventromedial PFC and the DLPFC with reversed polarity) did not find significant effects on the N-Back Task [29].

Self-regulation reflects the ability to maintain ideal motivational, emotional, and cognitive arousal, including inhibition and self-control [27]. Inhibition is the ability to control actions, thoughts, behaviours, and/or emotions to overcome internal (e.g., craving) or external (e.g., cue-induced) desire [27]. Self-control reflects the ability to resist temptations and hastiness [27]. Low self-control is a hallmark of SRADs [1] as it may predispose individuals to the inability to control, reduce, or stop the addictive behaviour [30]. Four studies evaluated tCS-induced effects on response inhibition [7, 29, 31] and self-control [12]. Regarding response inhibition, one study applied tCS over the DLPFC (anode over the left DLPFC, cathode over the right DLPFC) and reported significantly increased accuracy of trials and decreased reaction time on the Go/No-Go task [7]. The other two studies were conducted in patients with tobacco use disorder [29] and heavy drinkers (98.9% of individuals displayed alcohol use disorder) [31] but they did not report significant tCS-induced effects. To note, one of these studies combined tCS with mindfulness-based relapse prevention [31]. The effects of tCS on self-control was evaluated in a prospective study on patients with internet gaming disorder [12]. This was a single-arm, open-label study in which patients received 12 active tCS sessions (anode over the left and cathode over the right DLPFC) three times a week for four weeks. Patients displayed increased self-control, which correlated with decreased severity and time playing games as assessed by the Brief Self Control Scale [12]. Interestingly, the tCS regimen was followed by a partial alleviation of the asymmetry of glucose metabolism between the two DLPFCs. Although speculative, this may reflect a better communication between the two DLPFCs, which could lead to increased self-control. Despite the promising results, randomized, sham-controlled studies are necessary to draw further conclusions.

Selective attention is demonstrated by the ability to maintain attentional focus on the environment [27]. This function is closely related to working memory and attentional biases, since both require holding attention for some time [15, 27]. In SRADs, selective attention predicts the motivation to engage in treatment [32]. Work by Xu and collaborators [33] found no effect of tCS on selective attention in patients with tobacco use disorder. The study used anodal and cathodal tCS over the left DLPFC and the right supraorbital area, respectively. The authors discussed that this may be due to spurious factors, such as the fact that patients were abstinent overnight, which might influence tCS-induced effects on cortical excitability.

Two studies evaluated the effects of tCS on overall executive functions [34, 35], as assessed by the Frontal Assessment Battery, in patients with alcohol use disorder. Although the studies used different montages, neither of them found significant effects. Nevertheless, some limitations of the studies should be mentioned. For one, one study presented differences in the baseline amount of drinking between the active and sham groups [34]. Moreover, both studies had small sample sizes, which may reflect a lack of statistical power.

### **2.3 tCS effects on craving, mood, and stress in SRADs**

Craving, mood, and stress also play a major role in SRADs. They can influence cognition and can be modulated by tCS [36] (see **Figure 1**).

Craving is a complex process where individuals display a powerful urge or desire for a substance or an addictive behaviour (e.g., gambling, internet gaming) [1]. Craving can be triggered by external cues (e.g., a person, a place, or an object), as well as internal signals, such as mood or stress [37]. It is believed to play a central role in SRADs and constitutes one of the diagnostic criteria in the DSM-5 [1]. Several

clinical studies confirmed that tCS over the bilateral DLPFC can decrease craving in SRADs (for reviews, see [36, 38]). Yet, it remains to be seen whether this effect is due to a direct impact of the stimulation on craving or to an indirect effect which is secondary to an improvement of cognitive control [2].

Mood can also influence SRADs, since it can reinforce addictive behaviour [39]. For instance, anxious or depressive moods can influence cognitive functions such as self-control or decision-making and trigger craving and relapse. Therefore, improving mood might be one way to improve cognitive control to resist substances. Some evidence points to the effectiveness of tCS in improving mood in patients with SRADs [33, 40]. Two studies on tobacco use disorder found reduced negative affect following (1) anodal stimulation over the right (but not left) DLPFC and cathodal stimulation over the right DLPFC [40] and (2) anodal stimulation over the left DLPFC and cathodal stimulation over the right supraorbital area [33]. In both studies, there were differences neither in craving, nor in cigarette consumption. To note, patients were abstinent for at least six [40] or ten [33] hours, possibly suggesting the pertinence of testing in sated patients. Further, a preliminary study reported that tCS increased the perception of the quality of life in patients with online gaming disorder [12] (the details of this study are described in Table 1 as well as in a previous section about self-control).

Stress is a psychological and phenomenological experience accompanied by a specific physiological response [41]. Stress is purported to play a role in different stages of SRADs, from the initiation of the addictive behaviour to its relapse [41]. Both stress and addictive disorders are thought to share common neurophysiology, including a disrupted hypothalamic-pituitary-adrenal axis, as well as disrupted cognitive functions (e.g., selective attention, decision-making). In turn, both stress and addictive disorders may influence mood and cue reactivity, thereby increasing craving and probability of relapse. Furthermore, withdrawal symptoms themselves can cause stress for the individual. Thus, it is important to provide stress-coping strategies for patients with SRADs. Some evidence indicates that one session of active tCS over the DLPFC (anode over left DLPFC; cathode over right DLPFC), as compared to sham, can prevent a stress response (e.g., cortisol level) and decrease anxiety in healthy individuals that undergo psychosocial stress [42]. It remains to be seen whether tCS may be beneficial to stress reduction also in patients with SRADs.

### **3. Discussion**

Taken together, there are some trends that allow us to observe a general picture. Firstly, targeting the bilateral DLPFC appears to be the most effective tCS approach [5-10, 12-14], regardless of anode or cathode placement (see **Table 1**). This might suggest the importance of location and not laterality in SRADs [36], at least for cognitive functions. Secondly, decision-making was the most improved function across a variety of SRADs (tobacco, methamphetamine, gambling, cocaine use disorders, but not cannabis use disorder), which all targeted the bilateral DLPFC. Hence, there appears to be a link between targeting the DLPFCs and ameliorated decision-making. One possible explanation is that tCS modulates the interhemispheric balance between the two DLPFCs that is needed for decision-making functions [43]. It might be interesting for future studies to examine any possible underlying mechanisms (e.g., using fMRI). Also, it might be worth examining whether tCS can modulate other cognitive functions that are impaired across different SRADs (e.g., cognitive flexibility in alcohol and stimulant use disorders, reasoning in opioid use disorder, etc. [4]). Combining tCS with behavioural interventions, such as cognitive bias modification, does not appear to lead to promising results for alcohol use disorder. This might be due to several factors, such as the motivation of the subjects (some were not treatment seeking, and therefore might not be motivated to reduce their drinking), or the study design (perhaps there were too few sessions to induce changes). Interestingly, combining tACS with attentional bias modification

decreased attentional biases, as well as improving decision-making and decreasing craving in patients with tobacco use disorder. Although this was a proof of concept study [14], it nevertheless demonstrated the potential pertinence of combining these two interventions in SRADs.

Furthermore, a series of limitations of the reviewed studies should be taken into account. Firstly, patient characteristics such as age and sex knowingly influence tCS-induced effects [44-47] but were not always properly considered. In addition, the pattern of substance use disorders is different in men and women. For example, most studies included samples with a majority of men or even men-only. It would be important to include more women in studies. Importantly, it would be imperative to determine whether there are differences between sexes in tCS responses. Secondly, the majority of studies included detoxified and abstinent patients, while other stages of SRADs (e.g., sated, non-treatment seekers) remain unexplored. A recent study in non-treatment-seeking tobacco smokers suggested that sated patients responded better to tCS as compared to deprived patients, as reflected by a greater deactivation of the default mode network [29]. To support, acute nicotine in sated, as compared to abstinent, patients may present greater neural plasticity [48], thus presumably they may respond more to tCS. Thirdly, most studies did not include patients with comorbid disorders other than tobacco use disorder. Considering that comorbidities (e.g., mood disorders) are common in SRADs [49], it might be worth examining different subgroups. Fourthly, the motivation to change, which is associated with better response to tCS, remains unexplored [50]. Improving selective attention might be one way to improve motivation [32]. Also, greater motivation may relate to a better adherence to tCS regimens, which likely require several sessions in order to produce clinically meaningful improvements of symptoms [36]. Fifthly, behavioural states before stimulation and individual differences in brain morphometry on the effect of tCS treatments should be considered [51]. For instance, we previously observed that behaviours and brain morphometry impacted tDCS changes on neural substrates in patients with gambling disorder. In one study, there were positive correlations between tCS-induced changes of neurotransmitter levels in prefrontal and striatal regions and gambling related behaviours (i.e., craving, impulsivity, risk-taking) in patients with gambling disorder [52]. In another study, there were positive correlations between tDCS-induced elevations of prefrontal GABA levels and morphometry (volume and thickness) of the DLPFC in patients with gambling disorder [51]. In addition, the use of more objective and standardized outcome measures (e.g., a cue-provoked paradigm for craving) would allow more direct comparisons across studies. Lastly, future work could assess whether tCS can modulate other cognitive functions that may be relevant to SRADs, such as memory bias [53] and mindfulness [54].

#### **4. Conclusion**

In conclusion, tCS holds a strong clinical potential to improve cognitive functions when targeting the DLPFC. Further work is needed to determine the most effective protocols. One interesting therapeutic avenue might be individualized treatments based on patient characteristics such as brain morphometry, age and sex. Future studies could aim to optimize outcomes by combining tCS with medications or behavioural interventions (e.g., cognitive behavioural therapy) in order to improve outcomes even more [36].

#### **Acknowledgements**

AEB was supported by a Canadian Institutes of Health Research Frederick Banting and Charles Best award. SF was supported by the Canada Research Chair in Cognitive Neuroplasticity.

#### **Figure and table captions**

**Figure 1.** Transcranial current stimulation and current treatments might be used to target implicit as well as explicit cognitive functions in substance-related and addictive disorders. Some other processes might be worth targeting as well, such as craving, mood and stress, since they can influence cognitive functions and vice versa.

**Table 1.** Transcranial current stimulation can modulate several cognitive functions in substance-related and addictive disorders.

First author, year [ref#]	Design (N)	Addictive disorder	tDCS parameters	Anode placement	Cathode placement	Outcome measure(s)	Findings
<b>Cognitive bias</b>							
Mondino, 2020 [14]	Randomized Double-blind Sham-controlled Crossover (19)	Tobacco	1 session/condition <sup>a</sup> 10Hz, 2mA 30 min	N/A	N/A	Observation of smoking-related and neutral stimuli with eye tracking	↓ amount of time looking at smoking-related pictures
Vanderhasselt, 2020 [5]	Randomized Double-blind Sham-controlled Crossover (37)	Alcohol	1 session/condition 2 mA 20 min	R DLPFC	L DLPFC	Rewarded Go/No Go	↓ reward-triggered approach bias
Claus, 2019 [18]	Randomized Double-blind Sham-controlled 2 x 2 factorial (79)	Alcohol	4 sessions (once a week for 4 consecutive weeks) <sup>a</sup> 2 mA 2 x 10 min	R IFG	L upper arm	Approach-Avoidance-Task	No significant effect
Den Uyl, 2018 [17]	Randomized Double-blind Sham-controlled 2 x 2 factorial (83)	Alcohol	4 sessions over one week <sup>a</sup> 2mA 20 min	L DLPFC	R DLPFC	Visual probe task  Implicit Association Task	No significant effect
Shahbabaie, 2018 [6]	Randomized Double-blind Sham-controlled Parallel (90)	Methamphetamine	1 session 2mA 2 x 13 min	L DLPFC	R shoulder or R DLPFC <sup>^</sup>	Probe detection task	↓ attentional bias toward drug cues
Den Uyl, 2017 [13]	Randomized Double-blind Sham-controlled Parallel (91)	Alcohol	4 sessions over one week <sup>a</sup> 2 mA 20 min	L DLPFC	R DLPFC	Approach avoidance task	↓ approach bias
Den Uyl, 2016 [19]	Randomized Double-blind Sham-controlled 2 x 2 factorial (78)	Alcohol	3 sessions over 3 or 4 days <sup>a</sup> 1 mA 15 min	L DLPFC	R supraorbital area	Approach avoidance task  Implicit Association Task	No significant effect
<b>Cognitive flexibility</b>							
Alizadehgoradel, 2020 [7]	Randomized Double-blind Sham-controlled Parallel (39)	Methamphetamine	10 sessions over 5 weeks 2 mA 20 min	L DLPFC	R DLPFC	Wisconsin Card Sorting Task	↓ perseverative errors ↑ completed categories
Soyata, 2019 [8]	Randomized Triple-blind Sham-controlled Parallel (20)	Gambling	3 every other day sessions 2 mA 20 min	R DLPFC	L DLPFC	Wisconsin Card Sorting Task	↓ perseveration errors
<b>Decision making</b>							
Mondino, 2020 [14]	Randomized Double-blind Sham-controlled Crossover (19)	Tobacco	1 session/condition <sup>a</sup> 10Hz, 2mA 30 min	N/A	N/A	Delay Discounting Task	↓ percent of immediate choices
Alizadehgoradel, 2020 [7]	Randomized Double-blind Sham-controlled Parallel (39)	Methamphetamine	10 sessions over 5 weeks 2 mA 20 min	L DLPFC	R DLPFC	Balloon Analog Risk Task	↓ adjusted value ↓ maximum pumping
Soyata, 2019 [8]	Randomized Triple-blind Sham-controlled Parallel (20)	Gambling	3 every other day sessions 2 mA 20 min	R DLPFC	L DLPFC	Iowa Gambling Task	↑ net score

Gorini, 2014 [9]	Randomized Single-blind Sham-controlled Crossover (18)	Cocaine	1 session/condition 1.5 mA 20 min	L/R DLPFC	L/R DLPFC	Game of Dice Task  Balloon Analog Risk Task	↓ average of safe bets (anode over L DLPFC) ↑ average of safe bets (anode over R DLPFC)  ↓ adjusted average pumps
Fecteau, 2014 [10]	Randomized Quadruple-blind Sham-controlled Crossover (12)	Tobacco	5 daily sessions/condition 2 mA 30 min	R DLPFC	L DLPFC	Ultimatum Game	↑ rejected offers of cigarettes
Pripfl, 2013 [11]	Counterbalanced Sham-controlled Crossover (18)	Tobacco	1 session/condition .45 mA 15 min	L/R DLPFC	L/R DLPFC	Cold Columbia Card Task  Hot Columbia Card Task	↓ number of cards chosen in risky gamble (anode L DLPFC/cathode R DLPFC)  ↓ number of cards chosen in risky gamble (anode R DLPFC/cathode L DLPFC)
Boggio, 2010 [25]	Randomized Double-blind Sham-controlled Parallel (25)	Cannabis	1 session 2 mA 15 min	L/R DLPFC	L/R DLPFC	Risk Task	↑ choice of more risky prospects
<b>Self-regulation</b>							
Alizadehgoradel, 2020 [7]	Randomized Double-blind Sham-controlled Parallel (39)	Methamphetamine	10 sessions over 5 weeks 2 mA 20 min	L DLPFC	R DLPFC	Go/No-Go	↓ reaction time ↑ accuracy go trials ↑ accuracy no-go trials
Aronson Fischell, 2020 [29]	Randomized Double-blind Sham-controlled Crossover (15)	Tobacco	1 session/condition 2 mA 25 min	L DLPFC R VMPFC	R VMPFC L DLPFC	Flanker Task	No significant effect
Witkiewitz, 2019 [31]	Randomized Double-blind Sham-controlled Parallel (84)	Alcohol	Variable number of sessions <sup>ab</sup> 2 mA 30 min	R IFG	L upper arm	Stop Signal reaction time task	No significant effect
Lee, 2018 [12]	Open-label Single-arm (15)	Internet Gaming	3 sessions a week for 4 weeks 2 mA 30 min	L DLPFC	R DLPFC	Brief Self Control Scale	↑ self-control
<b>Selective attention</b>							
Xu, 2013 [33]	Counterbalanced Single-blind Sham-controlled Crossover (24)	Tobacco	1 session/condition 2 mA 20 min	L DLPFC	R supraorbital area	Visual attention task	No significant effect
<b>Working memory</b>							
Alizadehgoradel, 2020 [7]	Randomized Double-blind Sham-controlled Parallel (39)	Methamphetamine	10 sessions over 5 weeks 2 mA 20 min	L DLPFC	R DLPFC	N-back	↓ response time ↑ accuracy
Aronson Fischell, 2020 [29]	Randomized Double-blind Sham-controlled Crossover (15)	Tobacco	1 session/condition 2 mA 25 min	L DLPFC R VMPFC	R VMPFC L DLPFC	N-back	No significant effect
<b>Overall executive function</b>							
da Silva, 2013 [34]	Randomized Sham-controlled Parallel (13)	Alcohol	One session per week for five weeks 2 mA 20 min	L DLPFC	R supradeltoid area	Frontal Assessment Battery	No significant effect
Klauss, 2014 [35]	Randomized Double-blind Sham-controlled Parallel (33)	Alcohol	2 daily sessions for 5 consecutive days 2 mA 13 min	R DLPFC	L DLPFC	Frontal Assessment Battery	No significant effect

---

DLPFC: dorsolateral prefrontal cortex; IFG: inferior frontal gyrus; VMPFC: ventromedial prefrontal cortex; L: left hemisphere; R: right hemisphere; ↑: increase; ↓: decrease. Some articles appear more than once since they measured more than one cognitive function;<sup>a</sup>some or all subjects received a behavioural intervention as well; <sup>b</sup>subjects participated in a rolling group mindfulness-based relapse prevention while receiving either active or sham tDCS; those in the active and sham groups attended 4.32 and 3.78 sessions, respectively.

## References

1. American Psychiatric Association: Diagnostic and statistical manual of mental disorders. Fifth ed. Arlington, VA: 2013. Doi: 10.1176/appi.books.9780890425596.
2. Verdejo-Garcia A, Garcia-Fernandez G, Dom G. Cognition and addiction. *Dialogues Clin Neurosci*. 2019;21(3):281-90. doi: 10.31887/DCNS.2019.21.3/gdom.
3. Copersino ML. Cognitive mechanisms and therapeutic targets of addiction. *Curr Opin Behav Sci*. 2017;13:91-8. doi: 10.1016/j.cobeha.2016.11.005.
4. Fernandez-Serrano MJ, Perez-Garcia M, Verdejo-Garcia A. What are the specific vs. generalized effects of drugs of abuse on neuropsychological performance? *Neurosci Biobehav Rev*. 2011;35(3):377-406. doi: 10.1016/j.neubiorev.2010.04.008.
5. Vanderhasselt M-A, Allaert J, De Raedt R, Baeken C, Krebs RM, Herremans S. Bifrontal tDCS applied to the dorsolateral prefrontal cortex in heavy drinkers: Influence on reward-triggered approach bias and alcohol consumption. *Brain Cognition*. 2020;138:105512. doi: 10.1016/j.bandc.2019.105512.
6. Shahbabaie A, Hatami J, Farhoudian A, Ekhtiari H, Khatibi A, Nitsche MA. Optimizing electrode montages of transcranial direct current stimulation for attentional bias modification in early abstinent methamphetamine users. *Front Pharmacol*. 2018;9:907. doi: 10.3389/fphar.2018.00907.
7. Alizadehgoradel J, Nejati V, Sadeghi Movahed F, Imani S, Taherifard M, Mosayebi-Samani M et al. Repeated stimulation of the dorsolateral-prefrontal cortex improves executive dysfunctions and craving in drug addiction: A randomized, double-blind, parallel-group study. *Brain Stimul*. 2020;13(3):582-93. doi: 10.1016/j.brs.2019.12.028.
8. Soyata AZ, Aksu S, Woods AJ, İşçen P, Saçar KT, Karamürsel S. Effect of transcranial direct current stimulation on decision making and cognitive flexibility in gambling disorder. *Eur Arch Psychiatry Clin Neurosci*. 2019;269(3):275-84. doi: 10.1007/s00406-018-0948-5.
9. Gorini A, Lucchiari C, Russell-Edu W, Pravettoni G. Modulation of risky choices in recently abstinent dependent cocaine users: A transcranial direct-current stimulation study. *Front Hum Neurosci*. 2014;8. doi: 10.3389/fnhum.2014.00661.
10. Fecteau S, Agosta S, Hone-Blanchet A, Fregni F, Boggio P, Ciraulo D et al. Modulation of smoking and decision-making behaviors with transcranial direct current stimulation in tobacco smokers: a preliminary study. *Drug Alcohol Depend*. 2014;140:78-84. doi: 10.1016/j.drugalcdep.2014.03.036.
11. Pripfl J, Neumann R, Köhler U, Lamm C. Effects of transcranial direct current stimulation on risky decision making are mediated by 'hot' and 'cold' decisions, personality, and hemisphere. *Eur J Neurosci*. 2013;38(12):3778-85. doi: 10.1111/ejn.12375.
12. Lee SH, Im JJ, Oh JK, Choi EK, Yoon S, Bikson M et al. Transcranial direct current stimulation for online gamers: A prospective single-arm feasibility study. *J Behav Addict*. 2018;7(4):1166-70. doi: 10.1556/2006.7.2018.107.
13. Den Uyl TE, Gladwin TE, Rinck M, Lindenmeyer J, Wiers RW. A clinical trial with combined transcranial direct current stimulation and alcohol approach bias retraining. *Addict Biol*. 2017;22(6):1632-40. doi: 10.1111/adb.12463.
14. Mondino M, Lenglos C, Cinti A, Renaud E, Fecteau S. Eye tracking of smoking-related stimuli in tobacco use disorder: A proof-of-concept study combining attention bias modification with alpha-transcranial alternating current stimulation. *Drug Alcohol Depend*. 2020. In press.
15. Stacy AW, Wiers RW. Implicit cognition and addiction: a tool for explaining paradoxical behavior. *Annu Rev Clin Psychol*. 2010;6:551-75. doi: 10.1146/annurev.clinpsy.121208.131444.
16. Field M, Cox WM. Attentional bias in addictive behaviors: a review of its development, causes, and consequences. *Drug Alcohol Depend*. 2008;97(1-2):1-20. doi: 10.1016/j.drugalcdep.2008.03.030.

17. Den Uyl TE, Gladwin TE, Lindenmeyer J, Wiers RW. A clinical trial with combined transcranial direct current stimulation and attentional bias modification in alcohol-dependent patients. *Alcohol Clin Exp Res.* 2018;42(10):1961-9. doi: 10.1111/acer.13841.
18. Claus ED, Klimaj SD, Chavez R, Martinez AD, Clark VP. A randomized trial of combined tDCS over right inferior frontal cortex and cognitive bias modification: Null effects on drinking and alcohol approach bias. *Alcohol Clin Exp Res.* 2019;43(7):1591-9. doi: 10.1111/acer.14111.
19. Den Uyl TE, Gladwin TE, Wiers RW. Electrophysiological and behavioral effects of combined transcranial direct current stimulation and alcohol approach bias retraining in hazardous drinkers. *Alcohol Clin Exp Res.* 2016;40(10):2124-33. doi: 10.1111/acer.13171.
20. Jurado MB, Rosselli M. The elusive nature of executive functions: a review of our current understanding. *Neuropsychol Rev.* 2007;17(3):213-33. doi: 10.1007/s11065-007-9040-z.
21. Morris LS, Voon V. Dimensionality of cognitions in behavioral addiction. *Curr Behav Neurosci Rep.* 2016;3:49-57. doi: 10.1007/s40473-016-0068-3.
22. Berg EA. A simple objective technique for measuring flexibility in thinking. *J Gen Psychol.* 1948;39:15-22. doi: 10.1080/00221309.1948.9918159.
23. Clark L, Robbins T. Decision-making deficits in drug addiction. *Trends Cogn Sci.* 2002;6(9):361. doi: 10.1016/s1364-6613(02)01960-5.
24. Verdejo-Garcia A, Perez-Garcia M, Bechara A. Emotion, decision-making and substance dependence: a somatic-marker model of addiction. *Curr Neuropharmacol.* 2006;4(1):17-31. doi: 10.2174/157015906775203057.
25. Boggio PS, Zaghi S, Villani AB, Fecteau S, Pascual-Leone A, Fregni F. Modulation of risk-taking in marijuana users by transcranial direct current stimulation (tDCS) of the dorsolateral prefrontal cortex (DLPFC). *Drug Alcohol Depen.* 2010;112(3):220-5. doi:10.1016/j.drugalcdep.2010.06.019.
26. Fecteau S, Knoch D, Fregni F, Sultani N, Boggio P, Pascual-Leone A. Diminishing risk-taking behavior by modulating activity in the prefrontal cortex: a direct current stimulation study. *J Neurosci.* 2007;27(46):12500-5. doi: 10.1523/JNEUROSCI.3283-07.2007.
27. Diamond A. Executive functions. *Annu Rev Psychol.* 2013;64:135-68. doi: 10.1146/annurev-psych-113011-143750.
28. Bickel WK, Yi R, Landes RD, Hill PF, Baxter C. Remember the future: working memory training decreases delay discounting among stimulant addicts. *Biol Psychiatry.* 2011;69(3):260-5. doi: 10.1016/j.biopsych.2010.08.017.
29. Aronson Fischell S, Ross TJ, Deng Z-D, Salmeron BJ, Stein EA. Transcranial direct current stimulation applied to the dorsolateral and ventromedial prefrontal cortices in smokers modifies cognitive circuits implicated in the nicotine withdrawal syndrome. *Biol Psychiatry Cogn Neurosci Neuroimaging.* 2020;5(4):448-60. doi: 10.1016/j.bpsc.2019.12.020.
30. Percy A. Moderate adolescent drug use and the development of substance use self-regulation. *Int J Behav Dev.* 2008;32(5):451-8. doi: 10.1177/0165025408093664.
31. Witkiewitz K, Stein ER, Votaw VR, Wilson AD, Roos CR, Gallegos SJ et al. Mindfulness -based relapse prevention and transcranial direct current stimulation to reduce heavy drinking: A double -blind sham-controlled randomized trial. *Alcohol: Clin Exp Res.* 2019;43(6):1296-307. doi: 10.1111/acer.14053.
32. Rubenis AJ, Fitzpatrick RE, Lubman DI, Verdejo-Garcia A. Sustained attention but not effort-based decision-making predicts treatment motivation change in people with methamphetamine dependence. *J Subst Abuse Treat.* 2018;95:48-54. doi: 10.1016/j.jsat.2018.09.007.
33. Xu J, Fregni F, Brody AL, Rahman AS. Transcranial direct current stimulation reduces negative affect but not cigarette craving in overnight abstinent smokers. *Front Psychiatry.* 2013;4:112. doi: 10.3389/fpsy.2013.00112.

34. Da Silva MC, Conti CL, Klauss J, Alves LG, Do Nascimento Cavalcante HM, Fregni F et al. Behavioral effects of transcranial direct current stimulation (tDCS) induced dorsolateral prefrontal cortex plasticity in alcohol dependence. *J Physiol Paris*. 2013;107(6):493-502. doi: 10.1016/j.jphysparis.2013.07.003.
35. Klauss J, Penido Pinheiro LC, Silva Merlo BL, Correia Santos GDA, Fregni F, Nitsche MA et al. A randomized controlled trial of targeted prefrontal cortex modulation with tDCS in patients with alcohol dependence. *Int J Neuropsychoph*. 2014;17(11):1793-803. doi: 10.1017/s1461145714000984.
36. Ekhtiari H, Tavakoli H, Addolorato G, Baeken C, Bonci A, Campanella S et al. Transcranial electrical and magnetic stimulation (tES and TMS) for addiction medicine: A consensus paper on the present state of the science and the road ahead. *Neurosci Biobehav Rev*. 2019;104:118-40. doi: 10.1016/j.neubiorev.2019.06.007.
37. Sayette MA. The role of craving in substance use disorders: Theoretical and methodological issues. *Annu Rev Clin Psychol*. 2016;12:407-33. doi: 10.1146/annurev-clinpsy-021815-093351.
38. Hone-Blanchet A, Ciraulo DA, Pascual-Leone A, Fecteau S. Noninvasive brain stimulation to suppress craving in substance use disorders: Review of human evidence and methodological considerations for future work. *Neurosci Biobehav Rev*. 2015;59:184-200. doi: 10.1016/j.neubiorev.2015.10.001.
39. Koob GF. The dark side of emotion: the addiction perspective. *Eur J Pharmacol*. 2015;753:73-87. doi: 10.1016/j.ejphar.2014.11.044.
40. Pripfl J, Lamm C. Focused transcranial direct current stimulation (tDCS) over the dorsolateral prefrontal cortex modulates specific domains of self-regulation. *Neurosci Res*. 2015;91:41-7. doi: 10.1016/j.neures.2014.09.007.
41. Lemieux A, al'Absi M. Stress psychobiology in the context of addiction medicine: from drugs of abuse to behavioral addictions. *Prog Brain Res*. 2016;223:43-62. doi: 10.1016/bs.pbr.2015.08.001.
42. Carnevali L, Pattini E, Sgoifo A, Ottaviani C. Effects of prefrontal transcranial direct current stimulation on autonomic and neuroendocrine responses to psychosocial stress in healthy humans. *Stress*. 2019:1-11. doi: 10.1080/10253890.2019.1625884.
43. Fecteau S, Fregni F, Boggio PS, Camprodon JA, Pascual-Leone A. Neuromodulation of decision-making in the addictive brain. *Subst Use Misuse*. 2010;45(11):1766-86. doi: 10.3109/10826084.2010.482434.
44. Russell M, Goodman T, Wang Q, Groshong B, Lyeth BG. Gender differences in current received during transcranial electrical stimulation. *Front Psychiatry*. 2014;5:104. doi: 10.3389/fpsy.2014.00104.
45. Thomas C, Datta A, Woods A. Effect of aging on cortical current flow due to transcranial direct current stimulation: Considerations for safety. *Conf Proc IEEE Eng Med Biol Soc*. 2018;3084-7. doi: 10.1109/EMBC.2018.8513014.
46. Antonenko D, Nierhaus T, Meinzer M, Prehn K, Thielscher A, Ittermann B et al. Age-dependent effects of brain stimulation on network centrality. *Neuroimage*. 2018;176:71-82. doi: 10.1016/j.neuroimage.2018.04.038.
47. Lee S, Chung SW, Rogasch NC, Thomson CJ, Worsley RN, Kulkarni J et al. The influence of endogenous estrogen on transcranial direct current stimulation: A preliminary study. *Eur J Neurosci*. 2018;48(4):2001-12. doi: 10.1111/ejn.14085.
48. Grundey J, Thirugnanasambandam N, Kaminsky K, Drees A, Skwirba AC, Lang N et al. Neuroplasticity in cigarette smokers is altered under withdrawal and partially restituted by nicotine exposition. *J Neurosci*. 2012;32(12):4156-62. doi: 10.1523/JNEUROSCI.3660-11.2012.
49. NIDA. Common Comorbidities with Substance Use Disorders Research Report. 2020. Retrieved from <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders/>

50. Vitor de Souza Brangioni MC, Pereira DA, Thibaut A, Fregni F, Brasil-Neto JP, Boechat-Barros R. Effects of prefrontal transcranial direct current stimulation and motivation to quit in tobacco smokers: A randomized, sham controlled, double-blind Trial. *Front Pharmacol.* 2018;9:14. doi: 10.3389/fphar.2018.00014.
51. Bouchard AE, Dickler M, Renauld E, Lenglos C, Ferland F, Edden RA et al. The impact of brain morphometry on tDCS effects on GABA levels. *Brain Stimul.* 2020;13(2):284-6. doi: 10.1016/j.brs.2019.10.013.
52. Dickler M, Lenglos C, Renauld E, Ferland F, Edden RA, Leblond J et al. Online effects of transcranial direct current stimulation on prefrontal metabolites in gambling disorder. *Neuropharmacology.* 2018;131:51-7. Doi: 10.1016/j.neuropharm.2017.12.002.
53. Goldfarb EV, Fogelman N, Sinha R. Memory biases in alcohol use disorder: enhanced memory for contexts associated with alcohol prospectively predicts alcohol use outcomes. *Neuropsychopharmacology.* 2020;45(8):1297-305. doi:10.1038/s41386-020-0650-y.
54. Cavicchioli M, Movalli M, Maffei C. The clinical efficacy of mindfulness-based treatments for alcohol and drugs use disorders: A meta-analytic review of randomized and nonrandomized controlled trials. *European Addiction Research.* 2018;24(3):137-62. doi:10.1159/000490762.