

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Multi-stage sulfur and carbon mobility in fossil continental subduction zones: New insights from carbonate-bearing orogenic peridotites

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Consuma, G., Aulbach, S., Braga, R., Martin, L.A.J., Tropper, P., Gerdes, A., et al. (2021). Multi-stage sulfur and carbon mobility in fossil continental subduction zones: New insights from carbonate-bearing orogenic peridotites. *GEOCHIMICA ET COSMOCHIMICA ACTA*, 306, 143-170 [10.1016/j.gca.2021.05.008].

Availability:

This version is available at: <https://hdl.handle.net/11585/857005> since: 2022-02-12

Published:

DOI: <http://doi.org/10.1016/j.gca.2021.05.008>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Consuma G.; Aulbach S.; Braga R.; Martin L.A.J.; Tropper P.; Gerdes A.; Fiorentini M.L.: *Multi-stage sulfur and carbon mobility in fossil continental subduction zones: New insights from carbonate-bearing orogenic peridotites*

GEOCHIMICA ET COSMOCHIMICA ACTA

VOL. 306

ISSN 0016-7037

DOI: 10.1016/j.gca.2021.05.008

The final published version is available online at:

<https://dx.doi.org/10.1016/j.gca.2021.05.008>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Multi-stage sulfur and carbon mobility in fossil continental subduction zones: new insights from carbonate-bearing orogenic peridotites

Giulia Consuma^{a*}, Sonja Aulbach^{b,c}, Roberto Braga^a, Laure A.J. Martin^d, Peter Tropper^e, Axel Gerdes^{b,c} and Marco L. Fiorentini^f

^a Department of Biological, Geological and Environmental Sciences, University of Bologna, Piazza di Porta San Donato 1, 40126 Bologna, Italy

^b Institute for Geosciences, Goethe University, Altenhöferallee 1, 60438 Frankfurt, Germany

^c Frankfurt Isotope and Element Research Center (FIERCE), Goethe-Universität Frankfurt, Frankfurt am Main, Germany.

^d Centre for Microscopy, Characterisation and Analysis (CMCA), Australian Research Council Centre of Excellence for Core to Crust Fluid Systems (CCFS), The University of Western Australia, Crawley, 6009, WA, Australia

^e Institute of Mineralogy and Petrology, University of Innsbruck, Innrain52, 6020 Innsbruck

^f Centre for Exploration Targeting, ARC Centre of Excellence for Core to Crust Fluid Systems, The University of Western Australia, 35 Stirling Highway, Crawley, 6009WA, Australia.

Corresponding author: giulia.consuma@unibo.it

Keywords: sulfur isotopes; sulfide; mantle wedge; orogenic garnet peridotite

Abstract

The volatile transfer in subduction zones and the role of sulfate as a vector for the mobilization of oxidized components from down-going slabs remain hotly debated issues. Orogenic spinel and garnet peridotite lenses from the Ulten Zone (Eastern Alps, Italy), exhumed as part of felsic metamorphic terranes in continental collision zones, bear witness to mass transfer processes in these pivotal environments.

In this study, we report a multi-method investigation of mantle sulfides coexisting with four generations of carbonates, indicating coupled sulfur and carbon mobility throughout the peridotites' metamorphic evolution as part of the Variscan subduction architecture. Detailed petrography, bulk rock measurements, *in situ* chemical and geochemical analyses of sulfides as well as Sr isotope analyses of associated clinopyroxene and amphibole are combined with the aim to constrain the origin, nature and effect of multiple C-O-H-S-bearing fluids and melts the peridotite interacted with. The first, pre-peak, metasomatic pulse (Stage 1) is represented by an H₂S-CO₂-bearing melt from the subduction-modified hot mantle wedge, which formed a pyroxenite layer hosting matrix pentlandite with $\delta^{34}\text{S}$ of +2.77‰. Heavier $\delta^{34}\text{S}$ (up to +3.43‰), radiogenic Sr ($^{87}\text{Sr}/^{86}\text{Sr}_{\text{clinopyroxene}} > 0.7052$) and elevated Pb abundances recorded in coarse-grained garnet peridotite under eclogite-facies peak-conditions (Stage 2) are ascribed to interaction with isotopically heavy melts carrying recycled crustal component, permissive of, but not requiring, involvement of oxidized S species. Matrix carbonates occasionally occur in the coarse-grained peridotite. Conversely, isotopically lighter matrix pentlandite ($\delta^{34}\text{S} = -1.62$ to +0.67‰), and radiogenic Sr in amphibole ($^{87}\text{Sr}/^{86}\text{Sr} = 0.7056$) and associated dolomite (published data), from fine-grained garnet-amphibole peridotite may point to involvement of H₂S-

CO₂-bearing crustal fluids, which variably equilibrated with the mantle before interacting with the peridotites. Sulfide and amphibole in retrogressed spinel peridotite reveal the highest contents of fluid-mobile elements (As, Sb) and ⁸⁷Sr/⁸⁶Sr_{amphibole} up to 0.7074, suggesting late interaction with isotopically heavy crustal fluids at high fluid-rock ratios. The post-peak Stage 3 marks the entrapment of peridotites to form a tectonic mélange. Here, kelyphitization of garnet is catalyzed by further ingress of a S-bearing fluid ($\delta^{34}\text{S} = -0.38\text{‰}$), while carbonate veining with occasional sulfides witnesses channelized fluid flow. Textural observations indicate that, during Stage 4, serpentinization of peridotite at low $f\text{S}_2$ played an active role not only in CO₂ release by conversion of dolomite to calcite + brucite intergrowths, but also in local removal of ³²S during the final exhumation stage. Late channelized sulfur remobilization is evidenced by the serpentine + magnetite (\pm millerite \pm calcite) vein carrying > 300 ppm S.

Overall, the relatively narrow range of sulfur isotope composition ($\delta^{34}\text{S} = -1.62$ to $+3.76\text{‰}$) denotes limited interaction with isotopically heavy crustal fluids, and points to a subordinate role for subduction-derived sulfate throughout the extended evolution of UZ peridotites, first in the mantle wedge and then as part of a tectonic mélange.

1 Introduction

1 Sulfur (S) is a minor component in the mantle, yet it is thought to exert a key control on mantle redox
2 conditions as well as ore-forming processes (e.g., [Tomkins and Evans, 2015](#); [Evans et al., 2017](#); [Holwell et](#)
3 [al., 2019](#)). In xenoliths and orogenic peridotites, S is mostly stored in sulfides, such as pentlandite,
4 pyrrhotite, chalcopyrite, monosulfide-solid-solution (mss) and intermediate-solid-solution (iss) (e.g., [Lorand](#)
5 [et al., 2013](#); [Kiseeva et al., 2017](#); [Vaughan and Corkhill, 2017](#)). The speciation and mobility of S in
6 subduction zones is of particular interest because magmas in convergent-margin settings have been shown to
7 be more oxidized than those from spreading ridges (e.g., [Kelley and Cottrell, 2009](#); [Evans, 2012](#)), and with a
8 redox budget of eight electrons, S has been invoked as a potential oxidizing agent ([Evans et al., 2017](#);
9 [Bénard et al., 2018](#)).
10 Development of cutting-edge analytical techniques and improvement of thermodynamic modeling tools have
11 recently led to major advances in constraining the speciation, sources and mobility of sulfur in oceanic
12 subduction settings and associated ore deposit formation (e.g., [Schwarzenbach et al., 2018a](#), [Li et al., 2020](#);
13

14 [Walters et al., 2020](#)). However, major uncertainties persist for several reasons, such as (i) the lack of defined
15 sulfide solid solution models for thermodynamic modeling; (ii) the limited control on physicochemical
16 conditions (e.g. oxygen fugacity) determining S speciation and behavior, (iii) the lack of data from the
17 mantle wedge, prevented by the small size of mantle sulfides, which remains a less-well known domain
18 within the subduction factory.

19 Of note, the resultant recent research efforts produced conflicting lines of evidence for the sulfur behavior in
20 subduction zones and for its contribution to the oxidation state of the sub-arc mantle region. Thus, exhumed
21 mafic and ultramafic rocks from the island of Syros (Greece), interpreted as blocks detached from the
22 lithospheric slab to form a subduction channel mélange under blueschist-facies conditions, were shown to
23 have bulk-rock S isotopic signatures inherited from seawater alteration and overprinted by sediment-derived
24 fluids mobilizing S along the plate interface ([Schwarzenbach et al., 2018a](#)). [Evans et al. \(2014\)](#) find complex
25 S isotopic zoning of pyrite grains in eclogites originating as subducted oceanic crust from the Zermatt-Saas
26 zone (Western Alps) and the Pouébo terrane (New Caledonia), which they ascribe to S loss during prograde
27 metamorphism and sulfide growth at the earliest stage of exhumation. Similarly, [Walters et al. \(2019\)](#)
28 describe sulfides of metasomatic origin from a global suite of exhumed high-pressure rocks, that yield a wide
29 range of S isotopic compositions interpreted as precipitated from sulfate-bearing fluids. [Bénard et al. \(2018\)](#)
30 [also maintain that](#) oxidized slab components have the potential to oxidate the sub-arc mantle, by
31 documenting dissolved S^{6+} in spinel-hosted glass inclusions in sub-arc mantle xenoliths brought to the
32 surface by a recent volcanic activity in the Kamchatka (Russia) and West Bismarck arcs (Papua New
33 Guinea). On the other hand, [Li et al. \(2020\)](#) show most S to be present as reduced species, and argue for
34 limited S release in slab fluids. These constraints are based on a combination of petrographic observations,
35 whole-rock and *in situ* S isotope measurements and thermodynamic modelling of sulfide-bearing high-
36 pressure rocks and veins from the southwestern Tianshan (ultra-)high-pressure metamorphic belt (China).
37 Thermodynamic modeling results of [Piccoli et al. \(2019\)](#) also support the reducing nature of dehydrated slab
38 fluids at sub-arc depth. [Giacometti et al. \(2014\)](#) report little evidence from microtextural and *in situ* S
39 isotopic analysis of exhumed meta-ophiolites from the Western Alps for S mobilization during
40 metamorphism. Two-stage release of S during subduction is predicted by the thermodynamic modelling
41 results of [Walters et al. \(2020\)](#), with a small flux of reduced S at shallow depth and elevated S fluxes from

42 deep, cold oxidized slabs via release of sulfate and sulfide species. Although Tomkins and Evans (2015) also
43 predict a two-stage release process, they model anhydrite dissolution into fluids at low temperatures and the
44 release of H₂S when pyrite converts to pyrrhotite at higher temperatures.

45 Despite the importance of orogenic mantle peridotites as proxies for deep and retrograde-fluid-mediated
46 volatile exchange between crust and mantle (Scambelluri et al., 2010), few studies have investigated the S
47 speciation and isotopic composition of orogenic mantle rocks as an integral component of the global S cycle.

48 Orogenic spinel lherzolites in the North-Eastern Pyrénées yield negative values of $\delta^{34}\text{S} = -3.2\text{‰}$ to represent
49 an ancient S depletion event in the upper mantle, whereas positive values up to +3.3‰ of pyroxenite
50 intrusions were interpreted to reflect various enriched mantle sources (Chaussidon and Lorand, 1990). The
51 positive shift in the $\delta^{34}\text{S}$ of sulfides up to +5.4‰ from the garnet-bearing pyroxenite in the Western Gneiss
52 Region of Norway (Rielli et al., 2018) are ascribed to contamination from oceanic slab-derived oxidized
53 fluids, whereas negative values down to -10.0‰ are attributed to fluids sourced from the enclosing gneisses
54 or subducted oceanic sediments.

55 We focus on this little-explored facet of the S cycle, by investigating well-characterized orogenic spinel and
56 garnet peridotites from the Ulten Zone (UZ) in the Eastern Alps that originated as continental mantle that
57 became involved in continent-continent collision during the Variscan Orogeny. Displaying a diversity of
58 compositional and mineralogical features, including several generations of hydrous and carbonate minerals,
59 UZ peridotites have been the subject of multiple investigations, with a view to unraveling the element cycles
60 in convergent margin settings (Rampone and Morten, 2001; Scambelluri et al., 2006; Hermann et al., 2006;
61 Marocchi et al., 2009; Sapienza et al., 2009; Marocchi et al., 2010; Scambelluri et al., 2010; Marocchi et al.,
62 2011; Braga and Massonne, 2012; Förster et al., 2017; Ionov et al., 2017; Gudelius et al., 2019; Consuma et
63 al., 2020; Lo Pò et al., 2020). The occurrence of sulfides in the UZ peridotites and their petrological
64 implications have so far been overlooked as an important source of information on the behavior of S in
65 continental subduction zones. Taking advantage of the spatial resolution and textural control afforded by *in*
66 *situ* analyses on single sulfide grains, we use S isotopes as a sensitive tracer of the speciation and source of S
67 in subduction zones. Additional constraints on the nature of metasomatic agents can be inferred from the
68 variation in the trace element composition of texturally diverse pentlandite, which is known to reflect
69 changes in the physicochemical fluid parameters (e.g., pressure, temperature, oxygen fugacity). Dehydration

70 and partial melting generally depletes the upper mantle in fluid-mobile and magmaphile elements,
71 respectively, including the chalcophile elements As, Sb, Se, Te and Pb (Hattori et al., 2002; Hattori and
72 Guillot, 2003). As a consequence, their enrichment in the mantle wedge is usually explained by the supply of
73 these elements from slab-derived fluids and melts that percolate and metasomatize the overlying mantle.
74 These variably chalcophile elements (e.g., Barnes, 2016) would be expected to follow sulfide (e.g., as HS⁻)
75 mobilized in fluids and melts. *In-situ* determination of S isotopic and trace element compositions of
76 pentlandite, combined with whole-rock analyses, *in situ* Sr isotope compositions of amphibole and
77 clinopyroxene, and published *in situ* Sr isotope of the associated carbonates, allow to delineate a model for
78 the formation and modification of sulfide and associated carbonates, and to discuss the potential sources and
79 effects of C-O-H-S fluids percolating the mantle wedge during continental collision.

80

81 **2. Geological setting and metasomatic history**

82 The Ulten Zone (UZ) is a Variscan tectonic unit located in the Tonale Nappe (Eastern Alps) (Fig. 1). It is
83 part of the metamorphic basement of the Austroalpine domain, interpreted as a constituent of Gondwana
84 before its involvement in the Variscan collision with Laurussia (Regorda et al., 2020). In this unit, relicts of
85 high-pressure (HP) metamorphism are preserved in lenses of ultramafic rocks (mostly garnet-bearing
86 peridotites and eclogites) within migmatites and garnet-kyanite gneisses. The peridotites first resided in a
87 supra-subduction zone mantle wedge at relatively high temperature (about 1200°C), followed by re-
88 equilibration under eclogite-facies HP conditions of ~850°C and pressure of 2.2-2.7 GPa (Nimis and Morten,
89 2000; Braga and Sapienza, 2007), as a result of their movement toward deeper portions of the upper mantle
90 during continental crust subduction along a relatively cold path (Fig. 2a-b). Prior to or contemporaneously
91 with the attainment of eclogite-facies conditions, the former coarse-protogranular peridotites experienced
92 deformation and recrystallization into fine-grained peridotites, with formation of garnet and amphibole at the
93 expense of spinel and clinopyroxene. The post-collisional stage comprises the development of a crust-
94 peridotite tectonic mélange, and the onset of its exhumation is estimated to begin at around 330 Ma (Tumiati
95 et al., 2003). The three different lithologies described by Obata and Morten (1987) synthesize the different
96 stages of peridotite evolution: (1) (pre-peak) coarse-grained protogranular spinel peridotite (CS) that
97 transforms into a variety with coronitic garnet (CG), consisting of olivine + enstatite + diopside + Cr-Al

98 spinel (\pm garnet); (2) (peak) porphyroclastic (P-FGA) to equigranular fine-grained garnet-amphibole
99 peridotite (FGA) consisting of olivine + orthopyroxene + clinopyroxene + garnet + Ca-amphibole + spinel;
100 and (3) (post-peak/exhumation) fine-grained spinel (chlorite-amphibole) peridotite (FSA) with olivine +
101 orthopyroxene + Ca-amphibole + chlorite \pm spinel. All the UZ peridotites were variably serpentinized as they
102 were involved in a crust-mantle *mélange* along the exhumation path. The exceptional complexity of the
103 study area results from polyphase C-O-H-metasomatism along the Variscan continent collision (i.e. peak
104 pressure conditions and afterwards), during which inputs of metasomatic liquids percolated the overlying
105 orogenic mantle wedge inducing the formation of hydrous and carbonate phases.

106 On the basis of the high modal amphibole with a strong LILE/HSFE fractionation, [Rampone and Morten](#)
107 [\(2001\)](#) suggested that the metasomatic agents of the high-pressure hydrous metasomatism are fluids with a
108 low CO₂/H₂O ratio sourced from the neighboring crustal rocks. Nevertheless, carbonate phases related to the
109 progressive metasomatic stages are ubiquitous in Ulten Zone peridotites, the petrography of which is
110 described in detail in [Förster et al. \(2017\)](#). Inclusion of dolomite in coarse spinel, itself enclosed in garnet,
111 was ascribed as a trapped melt during the high-temperature metasomatic stage in the mantle wedge ([Stage 1](#)
112 [in Fig. 2a-b](#)), despite the radiating fracturing may point to a late metasomatic process ([Consuma et al. 2020](#)).

113 Formation of interstitial dolomite with minor magnesite has been ascribed as a peak-pressure feature, while
114 late exhumation-related serpentinization caused dolomite dissolution and formation of calcite+brucite
115 intergrowths via the reaction $\text{CaMg}(\text{CO}_3)_2 + \text{H}_2\text{O} \rightarrow \text{CaCO}_3 + \text{Mg}(\text{OH})_3 + \text{CO}_2$. This reaction implies CO₂
116 liberation from peridotites now entrapped in a crust-mantle *mélange* ([Förster et al., 2017](#)). Subsequent *in situ*
117 Sr isotope measurements in the UZ carbonates have shown multiple signatures related to their textural
118 positions ([Consuma et al., 2020](#)): relatively unradiogenic ⁸⁷Sr/⁸⁶Sr of ~ 0.705 of matrix dolomite formed at
119 eclogite-facies conditions, were taken to indicate that fluid sources that had variably equilibrated with a
120 depleted mantle reservoir mostly acted at high-pressure peak conditions. In contrast, variably radiogenic
121 ⁸⁷Sr/⁸⁶Sr (0.7036 to 0.7083) was ascribed to hybridized sources (i.e. mixing of depleted mantle and crustal
122 liquids) generated after the entrapment of peridotites into crustal rocks during the exhumation stage
123 ([Consuma et al., 2020](#)). The occurrence of these carbonates commonly in intimate association with sulfides
124 in different textural positions proves that metasomatic fluids and melts played a role in the mobility not only

125 of carbon but also of sulfur in the mantle wedge and tectonic mélange, and thus opens new questions
126 regarding their role in the global S cycle.

127

128 **3. Samples, prior work and sulfide petrography**

129 We investigated twenty orogenic peridotite samples from different localities of the Ulten Zone (Fig. 1), as
130 representative of different domains of the Variscan Ulten Zone mantle wedge. The selection was guided by
131 the motivation to include specimens that are representative of the metamorphic and metasomatic stages,
132 mineral assemblages, and presence of carbonates and sulfides.

133 Ten samples encompassing coarse spinel peridotites (UN1, UN2), fine-grained garnet-amphibole peridotites
134 (UN3, UN6, UN7, UN9, UN10, UN14) and fine-grained spinel-amphibole peridotites (UN8, UN16) were
135 previously studied in detail for their petrography, bulk-rock and mineral major-, minor- and trace-element
136 composition, as well as for garnet Fe speciation (Gudelius et al., 2019). The study shows that LILE, LREE
137 and some HFSE are co-enriched and revealed significant compositional differences between sampling
138 localities (Samerberg vs. Seefeld/Malga Masa Murada; Fig. 1), which is ascribed to different positions of the
139 peridotite bodies relative to the slab during subduction and exhumation. Despite evidence for strong
140 metasomatic overprint, $\text{Fe}^{3+}/\text{Fe}^{2+}$ in garnet remains low (<0.046), and oxygen fugacities relative to the
141 Fayalite-Magnetite-Quartz buffer ($\Delta\log f\text{O}_2$) of -2.4 to -0.32 are calculated for $P=2$ GPa, suggesting reducing
142 conditions in the mantle wedge beneath the Ulten Zone (Gudelius et al., 2019). Conversely, Malaspina et al.
143 (2009) find more oxidizing conditions (FMQ to FMQ+2 calculated at $P = 3$ GPa) in FGA peridotites from the
144 Malga Masa Murada and Samerberg areas. Thus, the UZ records heterogeneous redox conditions of various
145 mantle wedge and tectonic mélange domains (Fig. 2c).

146 A further seven samples were previously investigated by Consuma et al. (2020) (18LP1, MOL1-C, KL2.4-
147 2b, KL2.4-3, VM10A, KL1-A, MM1), integrating petrography, *in situ* major elements, Raman analyses and
148 *in situ* Sr isotopes of dolomite and calcite. Results show that high-pressure carbonation of the UZ peridotite
149 led to matrix dolomite formation from fluids variably equilibrated with a depleted mantle reservoir,
150 indicating the efficiency of the mantle wedge, as represented by UZ peridotites, in storing carbonates. During
151 exhumation, further percolation of hybridized fluids (sourced both from mantle and associated migmatites)
152 may have metasomatized the peridotites as part of a crust-mantle mélange, as suggested by the large range of

153 $^{87}\text{Sr}/^{86}\text{Sr}$ obtained for carbonate veins crosscutting the peridotite matrix. Dissolution of carbonates prevailed
154 during late exhumation-related serpentinization, as testified by pseudomorphs of calcite+brucite intergrowths
155 after dolomite first described by Förster et al. (2017).

156 The porphyroclastic fine-grained garnet-amphibole peridotite P10B was first investigated by Lo Pò et al.
157 (2020). Based on mineral chemistry and thermodynamic modelling, they interrogate the formation of
158 polymineralic inclusions (PI) in highly fractured cm-sized garnet. PI are proposed to crystallize as incoming
159 saline brines flow through fractures in garnet and locally interact with it along a post-Variscan, retrograde
160 path.

161 For the purpose of the present study, two additional peridotite samples were investigated: (i) MM1B, which
162 represents the composite serpentine vein crosscutting the matrix of FSA peridotite MM1; and (ii) coarse-
163 grained peridotite SBA2, the only carbonate-free sample (CGn). Classification of the samples with sulfide
164 assemblages, salient published data and analyses performed in this study are synthesized in Table 1.

165

166 Sulfides occur as discrete monophase and subordinately as polyphase grains with different shapes (rounded
167 grains, elongated, blocky, skeletal or irregular) and sizes (up to 150 μm). They show various stages of
168 alteration, ranging from none, to partial or strong alteration and fracturing, as illustrated in Figure 3 and
169 Electronic Annex B.

170 Six texturally distinct sulfide types are recognized linked to the different metamorphic stages of the UZ
171 peridotites (Fig. 2d):

- 172 1. *Polycrystalline Inclusions (PI) (Stage 1-4)*: sulfide inclusions in a cm-sized garnet and in spinel
173 included in mm-sized garnet occur as part of PI with grain size varying from 5 to 30 μm (Fig. 3a-b).
- 174 - *Included in cm-sized garnet* (peridotite P10B): Pyrrhotite (\pm pentlandite) is solely included in garnet
175 from this sample, whereas monophase pentlandite dominates in the matrix (Fig. 3a). PI fill a fracture
176 crosscutting the host garnet from the rim through the core. Here, PI show a complex association of
177 unusual minerals (amphibole + dolomite + chlorite + pentlandite + pyrrhotite + apatite +
178 kinoshitalite + sapphirine + garnet included in amphibole; Lo Pò et al., 2020).
- 179 - *Included in spinel (enclosed in garnet)* (peridotites KL2.4-2b and KL2.4-3): They occur as
180 assemblages of apatite + ilmenite + dolomite + pentlandite; calcite-brucite intergrowths + Cl-apatite

181 + pentlandite + ilmenite + phlogopite + chalcopyrite (Fig. 3b); calcite + ilmenite + Cl-apatite +
182 pentlandite. Also PI of native Cu + ilmenite + Cl-apatite + amphibole occur with radiating fractures,
183 linking the PI to the host spinel rim. Occasionally, PI (e.g. amphibole + ilmenite + sphalerite +
184 calcite + pentlandite) occur within, but at the edge of, coarse spinel connected with the surrounding
185 kelyphitic corona.

186 2. *Discrete sulfide inclusions (Stage 1-2)*: Orthopyroxene grains, Cr-rich spinel in garnet, Cr-poor
187 spinel in the matrix and kelyphitic corona host different types of sulfides, whereas no sulfides are
188 found in olivine. Notably, kelyphite around spinel and garnet commonly hosts sulfides (mostly
189 pentlandite) of exceptional dimensions (30-150 μm ; Fig. 3d). They commonly occur at the edge of
190 coarse spinel following the directions of the kelyphite growths or at the edge of the corona in contact
191 with the peridotite matrix (e.g. pentlandite with magnetite exsolutions and cut by late chlorite flakes;
192 pentlandite + sphalerite + native-Cu blebs associated with Cl-apatite and cut by chlorite flakes;
193 30 μm zoned chalcopyrite grain).

194 3. *Interstitial sulfides (Stage 2)*: They occur at silicate grain boundaries and commonly in close spatial
195 relationship with matrix dolomite (Fig. 3c). In the coarse-grained peridotite (CS + CG + CGn), they
196 are usually anhedral or subhedral with rounded or elongated shape and show a grain size from 10 to
197 50 μm . The carbonate-free peridotite SBA2 contains only five fractured blocky monophase
198 pentlandite grains with no alteration features. In CS and CG peridotites MOL1-C and 18LP1,
199 interstitial sulfides occur mainly as pentlandite + millerite assemblages partially replaced by
200 magnetite. In MOL1-C, pentlandite (partly replaced and rimmed by thin magnetite) + heazlewoodite
201 + dolomite + serpentine assemblage is cut by chlorite flakes (Fig. 3c). In the FGA peridotites KL1-A
202 and VM10A, pentlandite is commonly interstitial to olivine, orthopyroxene and pargasitic to
203 hornblenditic amphibole. It is locally associated with disseminated dolomite. Nickeline blebs (Ni,
204 As) are found within disseminated rounded pentlandite in association with Cl-apatite in FGA
205 peridotite VM10A. It is worth noting that this peridotite is very weakly serpentinized.

206 4. *In carbonate veins (Stage 3)*: Pentlandite + millerite cut by late chlorite flakes (with magnetite filling
207 the cleavage) are hosted in a vein predominantly formed by dolomite (and minor magnesite) +
208 tremolite + serpentine that cuts the peridotite matrix 18LP1 (Fig. 3e). Monophase pentlandite also

209 occurs not directly within the vein but in spatial proximity with it (i.e. few μm from the vein). A
210 nickeline bleb enclosed into pentlandite partially replaced by magnetite is documented.

211 5. *In matrix serpentine or serpentine vein (Stage 4)*: P-FGA peridotites KL2.4-2b and KL2.4-3 show
212 pentlandite + magnetite exsolutions contoured by chalcopyrite blebs. Exceptionally in these two
213 samples, sphalerite can also be found included in matrix serpentine, as single phase or polyphase-
214 assemblages (pentlandite + sphalerite + chalcocite). Matrix serpentine in FSA peridotite MM1 is
215 dominated by pentlandite rarely replaced by millerite or heazlewoodite. Conversely, the composite
216 serpentine + magnetite (\pm calcite \pm dolomite) vein MM1B, crosscutting the matrix of peridotite
217 MM1, hosts fractured elongated-to-rounded millerite in intimate association with Ni-oxide grains
218 with sizes of ~ 30 and $20 \mu\text{m}$ respectively. Nickeline blebs (NiAs; $3\text{-}5 \mu\text{m}$ on average) occasionally
219 occur within pentlandite (\pm magnetite) grains associated with serpentine (e.g. in peridotites MOL1-
220 C, 18LP1 and VM10A). Pentlandite enclosed in the serpentine + magnesite + tremolite vein
221 (peridotite MOL1-C) is pseudomorphously replaced by magnetite (Fig. 3e). Perpendicularly to this
222 vein, serpentine + pentlandite + magnetite fills the cleavage of the host mm-sized enstatite, rarely
223 containing small nickeline blebs.

224 6. *In halos, apophyses and sulfide-filled fractures (Stage 2-3-4)*: Diffuse sulfide material is commonly
225 observed in i) halos around discrete sulfide grains, ii) apophyses leading into sulfide-filled fractures
226 and iii) sulfide in fractures apparently unconnected to discrete sulfide grains (Fig. 3f). In megacrysts,
227 similar observations are ascribed to decrepitation of the sulfide during heating in contact with host
228 basalt (Andersen et al., 1987), a mechanism that does not apply here. It is noteworthy that such
229 features are observed both in fine-grained garnet amphibole peridotite and in spinel peridotite and
230 therefore not (exclusively) related to late-stage retrogression.

231 In addition to the reported textural occurrence of sulfides in peridotites, pentlandite grains interstitial to
232 silicates are documented occurring in a pyroxenite layer found in the coarse spinel peridotite MOL1-C,
233 which also hosts matrix dolomite. According to the Ulten Zone geodynamic scenario, these grains are related
234 to Stage 1.

235 To sum up, in order of decreasing abundance, sulfides occur mostly interstitial to silicates and associated
236 with matrix dolomite, enclosed in veins of serpentine and carbonates (dolomite, magnesite) and finally as

237 inclusions (in coarse spinel and few grains in silicates), coexisting with dolomite and calcite. Interstitial
238 pentlandite is by far the dominant sulfide phase occurring in all the UZ peridotite samples, with an average
239 grain size of 5-150 μm , followed by heazlewoodite and millerite mostly found in association with pentlandite
240 in serpentine and carbonate veins. Enclosed pentlandite is common in spinel as a discrete phase or PI, while
241 pyrrhotite (\pm pentlandite) occurs as PI in cm-sized garnet in peridotite P10B, except for few matrix pyrrhotite
242 grains in the “UN” sample suite. Finally, few sphalerite, chalcopyrite and chalcocite grains are hosted only in
243 P-FGA peridotites, mostly within coarse-grained spinel and kelyphite (KL24-2b, KI24-3, VM10A, P10B).

244

245 **4. Methods**

246 Each analytical method and specifics on quality control measures are detailed in [Electronic Annex C](#).

247 Samples were examined in polished thin section and slices of peridotite of 3 mm thickness by optical

248 transmitted and reflected light using a polarized optical microscope, and by Scanning Electron Microscopy

249 (SEM) for X-ray maps at the Department of Biological, Geological and Environmental Sciences, University

250 of Bologna, Italy.

251

252 Bulk rock major- and minor-element compositions along with selected trace elements were acquired by X-

253 Ray-Fluorescence, using a Phillips Magi XPRO spectrometer at Johannes Gutenberg-University Mainz.

254 Results are shown in [Table S1](#) in [Electronic Annex D](#). The same instrument, and a LECO-Analyzer CS 125,

255 were used to obtain bulk-rock S concentrations for a subset of samples (UN1-UN16) for which major- and

256 minor-element compositions had been previously determined ([Gudelius et al., 2019](#)). Remaining S

257 concentration measurements were carried out at Karlsruhe Institute of Technology, also using a LECO

258 instrument. Individual results are provided in [Table S2](#). Sulfide major- and minor-element concentrations

259 were obtained *in situ* by EPMA, using a JEOL JXA-8100 Superprobe electron microprobe at the Institute of

260 Mineralogy and Petrology, University of Innsbruck and a JEOL JXA-8900 at Goethe-University Frankfurt.

261 Results and summary are shown in [Table S3](#).

262

263 Trace-element abundances were determined *in situ* in sulfides following the procedure detailed in [Aulbach et](#)

264 [al. \(2012\)](#). Measurement statistics for two separate analytical campaigns and results are given in [Table S4](#)

265 [and S5 in Electronic Annex D](#). Abundances of Au cannot be quantified in a subset of samples that were Au-
266 coated prior to analysis for multiple S isotopes, as Au may have collected in small pits and cracks in the
267 sulfides.

268

269 The Rb-Sr isotopic composition of clinopyroxene and amphibole ([Table S6 in Electronic Annex D](#)) was
270 obtained *in situ* by laser ablation microprobe (LAM) multi-collector inductively-coupled plasma mass
271 spectrometry (MC-ICPMS), using a Resonetics RESolution M-50-HR laser probe linked to a
272 ThermoFinnigan Neptune mass spectrometer at Goethe-University Frankfurt, as described in [Electronic](#)
273 [Annex C](#).

274

275 Sulfur isotope compositions were measured *in situ* by secondary-ion mass spectrometry (SIMS) using a
276 Cameca IMS-1280 at the Centre for Microscopy and Microanalysis (CMCA), University of Western
277 Australia. SIMS measurements were undertaken on 30 pentlandite grains from seven peridotites and results
278 are reported in [Table 4](#). Grains with a minimum size of 20 μm were previously selected according to the
279 different textural positions and association with carbonates, which track progressive metasomatic episodes
280 during pre-peak, high-pressure-peak and retrograde conditions. All $\delta^{34}\text{S}$ values are reported relative to
281 Vienna Canyon Diablo Troilite – VCDT ([Ding et al., 2001](#)). Further details on the procedure are given in
282 [LaFlamme et al. \(2016\)](#).

283

284 **5. Results**

285 **5.1 Whole rock major elements and sulfur concentration**

286 New whole rock compositions are reported in [Table S1 in Electronic Annex D](#) and illustrated together with
287 published data (UN series) in [Figure 4](#). Loss On Ignition (LOI) ranges between 0.37 and 13.10 wt.%, in line
288 with the degree of serpentinization. The highest value of 13.10 wt.% is from the serpentine vein MM1B
289 previously hand-picked from FSA peridotite MM1. Contents of Al_2O_3 in the UZ peridotites range between
290 1.40 and 3.12 wt.%, while CaO ranges from 0.44 to 2.84 wt.%, TiO_2 from 0.02 to 0.11 wt.% and MgO from
291 37.87 to 42.79 wt.%. For comparison, primitive and depleted mantle have estimated Al_2O_3 concentrations of

292 4.45 and 4.28 wt.%, respectively, CaO of 3.55 and 3.50 wt.%, TiO₂ contents of 0.201 and 0.19 wt.% and
293 MgO contents of 37.8 and 38.22 wt.%, respectively (McDonough and Sun, 1995; Salters and Stracke, 2004).
294 Sulfur concentrations for eighteen peridotite samples are reported in Figures 4-5, and Table S2 in Electronic
295 Annex D. Overall, the S content of the orogenic peridotites ranges from as low as a few tens to hundreds
296 ppm, averaging 171 (± 45 ; n=18) ppm with a median value of 169 ppm.
297 Coarse-garnet peridotite (CG) shows an average value of 180 ± 28 ppm (n=5), fine-grained garnet amphibole
298 peridotite (FGA) an average value of 160 ± 34 ppm (n=9) and three fine-grained spinel peridotites (FSA)
299 have wide sulfur concentration ranging from 40 to 228 ppm. The serpentine + magnetite \pm sulfide (mostly
300 millerite) \pm calcite \pm dolomite vein MM1B reaches the highest value of 329 ppm S. Again for comparison,
301 primitive and depleted mantle have S concentrations of 250 and 60-119 ppm, respectively (Alt et al., 2007;
302 McDonough and Sun, 1995; Salters and Stracke, 2004).

303

304 5.2 Sulfide major element composition and low-pressure re-equilibration

305 The dataset comprises 295 sulfide grains analyzed for their major element compositions (Fig. S1 in
306 Electronic Annex A and Table S3 in Electronic Annex D); Table 2 reports the average major element
307 concentrations for pentlandite grains according to their different textural positions while Electronic Annex B
308 shows Backscatter-Electron (BSE) images for all analyzed sulfides in UN series samples.

309

310 Pentlandite is compositionally heterogeneous with 22.8-39.5 wt.% Fe, 28.8-37.8 wt.% Ni and below
311 detection limit (bdl)-16.6 wt.% Co. Here, pentlandite with Ni > 36wt.% is classified as Ni-pentlandite
312 (mostly found in serpentine vein), and that with Co > 6wt.% as Co-pentlandite (mostly found associated with
313 carbonate veins and in FSA peridotites). Data show that pentlandite enclosed in coarse spinel included in
314 garnet (occurring both as discrete grain and PI) and in silicates, as well as most of the matrix pentlandite
315 grains in P-FGA and FGA peridotites and some pentlandite enclosed in kelyphitic coronas around garnet fall
316 in the “typical” mantle sulfide range according to Wang et al. 2010 (Fig. S1 in Electronic Annex A).
317 Conversely, the other sulfides, such as those found in the matrix of CGn peridotite SBA2 and those included
318 in a serpentine vein show higher Co/(Fe+Ni+Cu) and Co/Ni ratio with respect to the mantle sulfides.

319 The Ni/(Ni+Fe) ratio of pentlandite averages 0.46 (± 0.05), ranging from 0.34 to 0.59, except for three Ni-
320 pentlandite grains with ratios >0.60 . In CGn peridotite SBA2, pentlandite shows a relatively high Co-
321 content, with an average of 1.89 wt.% (n=10). The Co-content increases in FSA peridotites MM1 and vein
322 MM1B with an average of 5.6 wt.% (n=7), while the highest value (Co = 16.6 wt.%) is reached in few
323 pentlandite grains enclosed in the dolomite vein of peridotite 18LP1. Millerite and heazlewoodite (\pm
324 pentlandite) with Ni > 63.6 wt.% are the dominant phases in peridotite MM1 and vein MM1B. Cu-enriched
325 pentlandite is found in grains enclosed in kelyphitic coronas around garnet, with Cu concentration from bdl-
326 2.6 wt.%. Nickel and Fe in pentlandite occurring in peridotites 18LP1, MM1 and MM1B are positively
327 correlated; conversely, a negative correlation is shown for pentlandite in fine, prophyroclastic, garnet-
328 amphibole peridotites (Fig. S1). The overall occurrence of pentlandite in the studied peridotites is consistent
329 with the typical sulfide component of orogenic mantle (Alard et al., 2000), and the measured Ni/(Ni+Fe)
330 ratio of about 0.46 (± 0.05) is also similar to those found in mantle xenoliths (e.g., Szabó and Bodnar, 1995;
331 Guo et al., 1999; Aulbach et al., 2019).

332 Pyrrhotite shows homogeneous chemical compositions, with a sulfur content of about 50.40 at.% (± 0.19).
333 The Ni content is always below detection limit. Minor Cu-sulfides are represented by few tiny blebs (max.
334 5 μm) of chalcopyrite and chalcocite, except for one chalcopyrite grain of 30 μm size found enclosed in
335 kelyphite. The metal/S ratio of chalcopyrite averages 0.97 (± 0.01). Only two chalcocite grains were found in
336 sample KL2.4-2b.

337 The high-temperature precursors of the UZ sulfides can be revealed by comparing their major chemical
338 compositions to experimentally determined phase relations at various temperatures in the ternary diagram S-
339 Fe-(Ni+Co) in at.% (Fig. S2a in Electronic Annex A). This indicates that minor pyrrhotite and dominant
340 pentlandite could be derived from re-equilibration of mss to low temperature (T). In addition, in the ternary
341 system Fe_9S_8 - Ni_9S_8 - Co_9S_8 , experimental studies have shown that the pentlandite composition is
342 temperature-dependent (Kaneda et al., 1986) and can reflect the condition under which it forms (e.g., Lorand
343 and Grégoire, 2006; LaFlamme et al., 2016). Most of the investigated pentlandite grains plot within the
344 entire field of stability for T of 200°C and 300°C (Fig. S2b), which are consistent with low-T assemblages
345 ($\leq 300^\circ\text{C}$) re-equilibrated from high-T-(mono)sulfides, whereby higher Co content shifts the stability field

346 for pentlandite formation to higher T (c. 300° C). Pentlandite can also originate by crystallization from
347 metasomatic Ni-Cu-rich sulfide melts with high metal/S ratios (Lorand et al., 2013).

348

349 **5.3 Sulfide trace element concentrations**

350 Trace elements were measured in pentlandite and pyrrhotite from the UN suite samples, peridotite P10B, and
351 from pentlandite previously measured by SIMS. A few millerite grains were also analyzed.

352 Pyrrhotite from PI in garnet shows Ni concentrations ranging from 119 to 8332 ppm, Cu from 22 to 569
353 ppm, Zn from 0.96 to 7.41 ppm, As from 0 to 30 ppm, Se from 2 to 24 ppm, Ag from 0.08 to 0.70 ppm, Sb
354 from 0.07 to 0.82 ppm, Te from bdl to 1.3 ppm, Pb from 1.7 to 22.6 ppm and Bi from 0.2 to 1.1 ppm.

355 Median values are reported in Table 3 and illustrated in Figure 6, while for the full trace-element dataset the
356 reader is referred to Table S5 in Electronic Annex D.

357 Siderophile and chalcophile element concentrations in pentlandite span a large range of values. Arsenic is the
358 most abundant trace element, ranging from 1 to 1213 ppm (median 514 ppm; n=86). Abundances of Zn
359 range from bdl to 535 ppm, maintaining low concentrations in matrix pentlandite from peridotite P10B (bdl-
360 1.73 ppm). Selenium and Te are always present in detectable abundances, ranging from 12 to 211 ppm
361 (median=81 ppm; n= 86) and from 1 to 89 ppm (median=14 ppm; n=80) respectively, while Ag ranges from
362 bdl to 51.67 ppm (median = 1 ppm; n=86). Concentrations of Sb are usually low (from bdl to 40 ppm;
363 median 1 ppm; n=80), except for few matrix pentlandite grains in FSA peridotites UN8 and UN16, and when
364 in association with a dolomite vein or enclosed in a serpentine vein of peridotite MOL1-C (here Sb values
365 are always > 13 ppm). By contrast, matrix pentlandite from P-FGA peridotite P10B varies from 0.07 to 0.28
366 ppm and the pentlandite in PI in spinel shows Sb concentration of 7.1 ppm. Pb is from 0.9 to 1730 ppm
367 (median=9 ppm; n=80) and Bi from 0.1 to 15.6 ppm (median=1 ppm; n=80).

368

369 **5.4 SIMS Sulfur isotope compositions of pentlandite**

370 Since the textural context of the sulfides provides crucial information on the stage during which S was
371 potentially added or removed (Fig. 2d), *in situ* acquisition of S isotopic composition of sulfides is
372 indispensable. Only a few grains showed a suitable size and composition (close to matrix-matched standards)
373 for SIMS analysis. Results are listed in Table 4 and shown in Figure 7.

374 In the coarse-grained spinel peridotite MOL1-C, a pyroxenite layer hosts a pentlandite grain with values of
375 $+2.77 \pm 0.37\%$ (n=2).

376 Matrix pentlandite in the only carbonate-free sample (SBA2), a coarse-grained garnet peridotite, consistently
377 shows relatively heavy $\delta^{34}\text{S}$, with an average of $+2.71 \pm 0.43\%$ (n=14). Except for one grain showing $+0.74$
378 $\pm 0.41\%$, the other grains are $\geq 1.52 \pm 0.45\%$, reaching a maximum value of $+3.54 \pm 0.35\%$. There is no
379 significant intra-grain isotope variability.

380 The CG peridotite 18LP1 contains a dolomite vein with pentlandite showing the highest Co values. Given
381 the restricted range of Co imposed by the sulfide standard to be measured by SIMS, only few grains could be
382 analyzed, yielding an average value of $+0.80 \pm 0.40\%$ (n=4) for pentlandite enclosed or locally associated
383 with the dolomite vein, while pentlandite included in coarse spinel gives $\delta^{34}\text{S} = -0.18 \pm 0.45\%$.

384 Pentlandite grains exhibit a wide isotopic range when occurring in serpentine + magnesite + tremolite vein,
385 with $\delta^{34}\text{S}$ values spanning between $-0.29 \pm 0.34\%$ and $+3.76 \pm 0.36\%$. These two extreme values were
386 measured in the same serpentine-enclosed pentlandite grain. Matrix pentlandite associated with dolomite
387 (Fig. 3c) averages $+3.11 \pm 0.38\%$ (n = 6).

388 The Mann-Whitney U Test (Electronic Annex C) reveals a significant difference between the $\delta^{34}\text{S}$ values of
389 pentlandite within the fine-grained and the coarse-grained peridotite lithotypes, at a level of significance of
390 1% ($p < 0.00001$). The isotopic values of pentlandite in the fine-grained peridotite are significantly lighter
391 than those in the coarse-grained peridotite.

392 The single pentlandite in PI in coarse spinel (in turn included in garnet) exhibits $\delta^{34}\text{S} = +0.46 \pm 0.34\%$. Two
393 matrix pentlandite grains in the fine-grained peridotite KL1-A, which are intimately associated with matrix
394 dolomite, show values of $-1.62 \pm 0.42\%$ and $-1.14 \pm 0.39\%$ respectively. Matrix pentlandite in sample
395 KL2.4-3 averages $+0.31 \pm 0.34\%$ (n=3) with little intra-grain isotope variability. Within the same peridotite,
396 a pentlandite grain found in kelyphite around spinel and cut by a chlorite flake averages $-0.38 \pm 0.31\%$
397 (n=3). In peridotite VM10A, matrix pentlandite $\delta^{34}\text{S}$ ranges from $-1.32 \pm 0.40\%$ to $+0.67 \pm 0.40\%$ with an
398 average of $-0.21 \pm 0.41\%$ (n=12).

399 Overall, the S isotope compositions obtained for the UZ pentlandite grains range between $-1.62 \pm 0.42\%$ and
400 $+3.76 \pm 0.36\%$. Matrix pentlandite from fine-grained peridotites is isotopically light, with an average of

401 $\delta^{34}\text{S} = -0.11 \pm 0.39\%$ (n=15), ranging from $-1.62 \pm 0.42\%$ to $+0.67 \pm 0.40\%$. Conversely, matrix pentlandite
402 from coarse-grained peridotites is isotopically heavy, with an average $\delta^{34}\text{S} = +2.48\% \pm 0.41\%$ (1σ , n=25)
403 and ranging between $+0.41 \pm 0.42\%$ to $+3.76\% \pm 0.36\%$. For comparison, $\delta^{34}\text{S}$ of the depleted mantle is
404 estimated at $-1.28 \pm 0.33\%$ (Labidi et al. 2013).

405

406 5.5 In situ Sr isotope of clinopyroxene and amphibole

407 Laser ablation MC-ICP-MS analyses were performed on 15 clinopyroxene grains from 5 UZ peridotite
408 samples (total of 25 analyses; Table S6 in Electronic Annex D) (Fig. 8). Clinopyroxene in the two fine-
409 grained garnet-amphibole peridotites P10B and VM10A sampled from Mt. Hochwart (Fig. 1) show an
410 average measured $^{87}\text{Sr}/^{86}\text{Sr}$ value of 0.70538 ± 0.000076 (1σ , n = 11). The porphyroclastic garnet-amphibole
411 peridotite sample KL2.4-2b from Klappfbergalm (Fig. 1) shows similar $^{87}\text{Sr}/^{86}\text{Sr}$ with 0.70535 ± 0.00012 (1σ ,
412 n = 4). Clinopyroxene grains from the two coarse-grained peridotite samples UN1 and UN2 from Samerberg
413 area (Fig. 1) show similarly radiogenic values of 0.70544 ± 0.00015 and 0.7055 ± 0.0020 . The lower
414 standard deviations obtained for clinopyroxene in fine-grained amphibole peridotite compared to coarse
415 protogranular peridotites (using the same method in the same laboratory) suggests a greater degree of
416 isotopic homogenization of the former, which were subjected to recrystallization and retrogression.

417

418 Furthermore, 23 amphibole grains from 8 samples were analyzed for *in situ* Sr isotope compositions (n=41
419 analyses; Table S6). Measured $^{87}\text{Sr}/^{86}\text{Sr}$ of amphibole in the P-FGA peridotites UN3, UN6 (from Samerberg)
420 and UN7 (from Seefeld) averages 0.70548 ± 0.00006 (1σ , n = 15), while peridotite UN10 from Seefeld
421 shows more radiogenic values of 0.70597 ± 0.00009 (n = 4), and the most radiogenic value is found in
422 peridotite UN14 from Malga Masa Murada with 0.70705 ± 0.00006 (n = 6). The FSA peridotites UN8 and
423 UN16, from Seefeld and Malga Masa Murada respectively, exhibit $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.70679 ± 0.00008 (n = 5)
424 and 0.70779 ± 0.00011 (n = 6), respectively. For comparison, the primitive and depleted mantle have
425 estimated $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.7045 and 0.7026, respectively (Workman and Hart, 2005). Whereas the low $^{87}\text{Rb}/^{86}\text{Sr}$
426 ratio of the clinopyroxenes (average 0.02) imply that initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios (at 330 Ma; Tumiati et al. 2003)
427 are not much different from the measured ones, this is not the case for amphiboles with an average $^{87}\text{Rb}/^{86}\text{Sr}$

428 ratio of 0.14 (Table S6). For the purpose of the following discussion, the initial values will be used for both
429 clinopyroxene and amphibole.

430

431 **6. Discussion**

432 **6.1 Fractionation of sulfur isotopes during high-temperature processes and late-stage re-equilibration**

433 Few studies have systematically examined the combined geochemical and isotopic signature of sulfides from
434 orogenic mantle domains as a probe to investigate the nature of high-pressure and exhumation processes in
435 collisional settings. The full set of metasomatic and re-equilibration processes affecting the UZ peridotites
436 throughout their evolution dictates their sulfide mineralogy as well as their chemical and sulfur isotope
437 compositions. As these processes may produce sulfur isotopic fractionations, the effects of high-temperature
438 processes and low-temperature re-equilibration of the sulfide assemblage on $\delta^{34}\text{S}$ have to be considered
439 before attempting any interpretation on the relationships between sulfur isotope signatures and source/s.
440 Pentlandite is not stable at the mantle conditions experienced by UZ peridotites during their multi-stage
441 evolution (Fig. 2a-b). It either formed together with minor amounts of pyrrhotite, now in part replaced by
442 magnetite, from a high-temperature monosulfide precursor (Fig. S2), or precipitated directly from a
443 metasomatic metal-rich fluid (Lorand and Grégoire, 2006; Lorand et al., 2013; Giuliani et al., 2016). Prior
444 modeling has demonstrated that the S isotopic fractionation between pentlandite and a precursor monosulfide
445 solid solution (mss) is limited to 0.7 ‰ for equal proportions of pentlandite and mss and low temperature of
446 pentlandite formation (300°C; Giuliani et al., 2016). Given the dominance of pentlandite in the low-
447 temperature assemblages in UZ peridotites, it is argued that the $\delta^{34}\text{S}$ signature of pentlandite largely reflects
448 that of its high-temperature precursor. Furthermore, since the pH of fluids does not induce significant isotope
449 fractionation at the metamorphic conditions of subduction zones (Giacometti et al., 2014), the measured $\delta^{34}\text{S}$
450 of pentlandite is considered broadly representative of the bulk rock that interacted with various S-bearing
451 metasomatic fluids or melts.. The sulfur isotope composition is not modified by high-pressure prograde
452 metamorphism of subducted rocks (Evans et al., 2014; Li et al. 2021), and the S isotopic fractionation
453 between solid sulfide and sulfide melt or H_2S in reduced fluids is negligible (Ohmoto and Rye, 1979; Li and
454 Liu, 2006; Marini et al., 2011). Instead, significant fractionation occurs between oxidized and reduced S
455 species (up to 4-5‰) even at mantle temperatures (i.e. $T > 900^\circ\text{C}$; Ohmoto and Rye, 1979). During the

456 Variscan subduction and consequent exhumation, the precursor sulfide to pentlandite may have interacted
457 with, or precipitated from, fluids or melts containing oxidized or reduced S species.

458 Sulfate is recycled with sediments and altered oceanic crust (Alt and Shanks, 2006; Schwarzenbach et al.,
459 2018b), and it usually preserves a positive S isotope signature close to the seawater composition (Alt et al.,
460 1995; Alt and Shanks, 2001, Li et al., 2021). The timing and mechanisms of sulfate-to-sulfide reduction
461 remain poorly constrained, since so far no sulfate grains have been found in equilibrium with mineral
462 assemblages of HP-UHP subducted metamorphic rocks (Li et al. 2021 and references therein). Of note, Li et
463 al. (2021) measure heavy $\delta^{34}\text{S}$ signature (+25‰) for pyrite in veins from subducted HP metamorphic rocks
464 from the Southwestern Tianshan HP-UHP metamorphic belt (northwestern China), which exceptionally
465 documents that the seawater sulfate-derived $\delta^{34}\text{S}$ signature may be retained at great depths (i.e. 70 km depth).
466 The authors suggest that reduction of sulfate to sulfide may have occurred during pre-subduction
467 hydrothermal alteration or contemporaneous with subduction at fore-arc depths, implying that no isotope
468 fractionation occurred (Li et al., 2021). On the other hand, a fractionation value of ~20‰ between dissolved
469 sulfate and pyrite precipitation was modeled for fluid migration throughout the slab-mantle interface at
470 300°C, reaching up to 120‰ depending on distance from the source, pathway and amount of pyrite
471 precipitation (Walters et al., 2019).

472 In the UZ peridotites, the high oxygen fugacity of FMQ to FMQ+2 estimated by Malaspina et al. (2009) at
473 3GPa approach conditions where significant proportions of oxidized S would be present (Jugo et al., 2010).
474 However, the oxygen fugacity conditions recorded by the UZ peridotites investigated in this study (<FMQ at
475 2 GPa; Gudelius et al. 2019) are far too reducing to allow the presence of significant amounts of sulfate.
476 Moreover, despite the restricted range of S isotope values in the UZ peridotite, we observe a clear distinction
477 among the lithotypes and textural occurrences. We therefore propose that multiple melt-and-fluid/rock
478 interactions occurred along and after the Variscan Orogeny, and we discuss the nature of the fluid sources in
479 Section 6.4.2. Although Rielli et al. (2018) obtained a much larger range of $\delta^{34}\text{S}$ (-10.0 to +5.4‰), it is worth
480 noting that their sample suite is composed of pyroxenite and websterite lenses in garnetite and amphibole
481 layers in pyroxenite within the main peridotite bodies from the Western Gneiss Region (Norway), whereas
482 no sulfide grain was found within the matrix of the only peridotite sample in their study. Consequently, there

483 are no sulfur isotope compositions of garnet peridotite available in the literature to make a strong comparison
484 with the data presented in our study.

485

486 **6.2 Near-quantitative sulfide extraction during SCLM stabilization and addition during refertilization**

487 The sulfur isotope composition of sub-continental lithospheric mantle (SCLM), which stabilizes by
488 extraction of variable degrees of partial melt (e.g., [Griffin et al., 2004](#)), remains poorly constrained and has
489 been mostly addressed in studies of sulfide inclusions in diamonds from mantle xenoliths, giving a $\delta^{34}\text{S}$
490 range between -4‰ and + 6‰ ([Chaussidon et al., 1987](#); [Farquhar, 2002](#); [Thomassot et al., 2009](#)). In the
491 Ulten Zone, during high-temperature pre-peak conditions (Stage 1), the coarse-grained protogranular
492 peridotites containing the assemblage olivine + enstatite + diopside + Cr-Al spinel reside in the continental
493 lithospheric mantle and equilibrate at 1200°C and 1.3-1.6 GPa ([Nimis and Morten, 2000](#)). Their study may
494 therefore provide insights into mantle wedge evolution before its entrainment into the Variscan subduction
495 architecture. Samples in this study appear to fall along a curved array with respect to bulk-rock TiO_2 and
496 Al_2O_3 , both of which are moderately incompatible (Fig.4). Such arrays result from partial melt extraction.
497 In accord with [Ionov et al. \(2017\)](#) and [Gudelius et al. \(2019\)](#), major element compositions indicate mostly
498 ~15-25% of melt extraction (Fig. 4a), although this is a minimum estimate if refertilization occurred
499 ([Gudelius et al., 2019](#)). At such melt fractions, sulfide would have been exhausted, depending on oxygen
500 fugacity and S concentration ([Keays, 1995](#); [Mungall et al., 2006](#)). For example, for Al_2O_3 contents of 2-2.3
501 wt.% recorded in some UZ peridotites (Fig. 4a), less than 100 ppm S would be expected to remain in the
502 mantle for fractional melt extraction at 3 GPa and an oxygen fugacity of FMQ-2 ([Aulbach et al., 2016](#)). If
503 melt extraction occurred at lower pressure, then S concentrations in the residue would be even lower, given
504 increasing S solubility in silicate melt at sulfide saturation with decreasing pressure ([Mavrogenes and](#)
505 [O'Neill, 1999](#)). In this light, the UZ peridotites containing >100 ppm S ([Table S2, Electronic Annex D](#))
506 almost certainly experienced S addition. Furthermore, sulfidation of peridotite has been shown to cause Fe-
507 Ni zoning in olivine ([Papike et al., 1995](#); [Bataleva et al., 2016](#)). Although such zoning is not observed in the
508 UZ olivine ([Gudelius et al., 2019](#); [Consuma et al., 2020](#)), higher NiO contents in olivine are observed in S-
509 poor peridotites relative to olivine from S-rich peridotites, consistent with Ni extraction from olivine during
510 metasomatic sulfide addition (Fig. 4c).

511 Hereafter, for each metamorphic stage, we discuss (i) the process and related metasomatic agents that may
512 have modified the sulfide component throughout the evolution of peridotites; (ii) the relevant conditions (P-
513 T-fO₂-fS₂) that may have exerted a major control on the nature of sulfides and their mobility with respect to
514 the melt/fluid phase; (iii) the textural relationship of sulfides with the coexisting carbonates and hydrous
515 minerals. We conclude with the potential implications that our findings may have for the understanding of
516 the global sulfur cycle.

517

518 **6.3 Pre-collision mantle metasomatism (Stage 1)**

519 Stage 1 is represented by rare coarse spinel peridotites containing pyroxenite layers (Nimis and Morten,
520 2000), with only one sample available for this study (MOL1-C). During this stage, percolation of a hydrous
521 basic melt induces an enrichment of the lithospheric mantle in LREE, LILE and Li at spinel-facies
522 conditions (Nimis and Morten, 2000; Scambelluri et al., 2006). Precipitation of this melt prompts the
523 formation of sulfide-bearing pyroxenite layers within the peridotite. The $\delta^{34}\text{S}$ signature ($+2.77 \pm 0.37\%$;
524 $n=2$) of matrix pentlandite in pyroxenites is heavier than that estimated for continental lithospheric mantle
525 ($+1.35 \pm 0.25 \%$; Fiorentini et al., 2018), and similar to the $\delta^{34}\text{S}$ signature (up to $+5.4\%$) of sulfides
526 contained in pyroxenite lenses within peridotite bodies from the Western Gneiss Region (Rielli et al., 2017).
527 Similarly, pentlandite associated with dolomite in the surrounding peridotite matrix is also isotopically heavy
528 ($+3.11 \pm 0.38\%$; $n = 6$) (Table 4).

529 Euhedral inclusions of dolomite in primary spinel within porphyroclastic garnet from coarse-grained UZ
530 peridotite, may represent relics that testify to the percolation of carbon-bearing liquids prior to Stage 2 garnet
531 growth (Förster et al., 2017). The aforementioned petrographic and isotopic evidence indicate the presence
532 of (possibly subduction-related) isotopically heavy S in the source of the melts involved in Stage 1
533 metasomatism. Temperature and oxygen fugacity are the main factors that dictate the solubility and
534 speciation of sulfur in silicate melts (Clemente et al. 2004; Jugo et al. 2010; Jégo and Dasgupta, 2014; Canil
535 and Fellows, 2017). Experimental data on the sulfide vs. sulfate stability in basaltic and sediment glasses
536 (Jugo et al., 2010; Canil and Fellows, 2017) suggest that the melt is saturated in sulfide (S²⁻) rather than in
537 sulfate (S⁶⁺) at oxygen fugacity lower than around the Fayalite-Magnetite-Quartz (FMQ) buffer. In this light,
538 the relatively reducing nature (<FMQ; Gudelius et al., 2019) of the investigated samples is suggested to also

539 indicate a system dominated by reduced S species. Additionally, the overall lack of identifiable reduced
540 carbon species (i.e. graphite or disordered carbonaceous material; e.g., Vitale Brovarone et al., 2020) in the
541 UZ peridotites requires carbon to be present mostly as carbonate (CO₂). We speculate that any S added from
542 melt leaving a lithospheric mantle source would have involved an H₂S-CO₂-bearing agent, consistent with
543 suggestions that mantle fluids are H₂S-dominated (Eggler and Lorand, 1993).

544

545 **6.4 High-pressure (garnet) metamorphism and recrystallization (Stage 2)**

546 *6.4.1 Coarse-grained vs Fine-grained peridotite*

547 With the onset of the Variscan Orogeny, the corner flow initiated by continental subduction dragged the UZ
548 peridotites deeper into the lithospheric mantle towards the slab-wedge interface, causing their textural
549 transition from coarse protogranular to porphyroclastic and finally fine equigranular. The peak conditions
550 were attained at ~850°C and 2.2-2.7 GPa (Nimis and Morten, 2000; Braga and Sapienza, 2007). At this
551 stage, the UZ peridotites still resided in the mantle wedge (Scambelluri et al., 2006), where matrix
552 pentlandite coexisted along with high-pressure matrix phases, such as recrystallized olivine and pyroxenes,
553 pargasitic to hornblenditic amphibole, garnet, and dolomite. Thus, the S isotopic composition of pentlandite
554 may reflect peak-garnet conditions with superposed local interaction with retrograde fluids. The Mann-
555 Whitney U Test allows to distinguish two populations of matrix sulfides with respect to δ³⁴S: one from fine-
556 grained peridotite (δ³⁴S = -0.11‰ ± 0.39; 1σ, n=15) and one from the coarse counterpart (δ³⁴S = +2.35‰ ±
557 0.43‰; 1σ, n=18). Thus, despite the limited range of measured sulfur isotope ratios for the entire data set (-
558 1.62 to +3.76 ‰), differences in the source of sulfur for texturally diverse sulfides are detected. Moreover,
559 some coupling between the δ³⁴S signature of pentlandite and trace elements becomes evident when not only
560 the textural positions (Fig. 9), but also the different sampling localities are considered. This suggests some
561 provinciality with respect the nature of metasomatism (see Fig. S3 in Electronic Annex A), as previously
562 recognized based on lithophile elements (Gudelius et al., 2019).

563

564 *Coarse-grained garnet peridotite.* The positive δ³⁴S values obtained for a population of matrix pentlandite in
565 the coarse-grained garnet peridotites are distinct from the proposed depleted mantle range of δ³⁴S of -1.28 ±
566 0.33‰ (Labidi et al. 2013), from the suggested mantle wedge composition of 0‰ (Li et al. 2020), and from

567 the proposed composition of continental lithospheric mantle ($+1.35 \pm 0.25 \text{ ‰}$; Fiorentini et al., 2018).
568 Combined with elevated S abundances ($\geq 166 \text{ ppm}$; Table 1), the measured isotope values require addition of
569 isotopically heavy S, either from an enriched mantle source as recognized by Ionov et al. (1992) or involving
570 processes inducing heavy S isotope enrichment, such as fractionation of isotopically light sulfide (Walters et
571 al., 2019). The initial Sr isotopic composition of clinopyroxene in two coarse-grained peridotites
572 ($^{87}\text{Sr}/^{86}\text{Sr}=0.70509\text{-}0.70538$; Table 1) is more radiogenic than that of the primitive or depleted mantle
573 (0.7045 and 0.7026 , respectively; Workman and Hart, 2005), and also indicates that the enrichment in
574 radiogenic Sr pre-dates the entrapment of peridotites into the tectonic *mélange*. Indeed, coarse-grained UZ
575 peridotites show evidence for interaction with crust-derived silicic melts (Gudelius et al., 2019).
576 Figure 9 shows that the isotopically heavy $\delta^{34}\text{S}$ signature of pentlandite in coarse peridotites is coupled with
577 with elevated Pb concentrations, but not with similarly elevated abundances of other fluid-mobile elements,
578 such as As and Sb, which may again point to a continental crustal contaminant carried by a melt rather than a
579 fluid. Indeed, for sulfide- rather than sulfate-saturated sediment-derived melts, Sb will be depleted in the
580 melt relative to As and Pb (Canil and Fellows, 2017), which corresponds to the systematics observed for
581 sulfide in the coarse-grained peridotite (Fig. 9).
582
583 *Fine-grained garnet-amphibole peridotite.* Following the interpretation of Obata and Morten (1987), the
584 fine-grained lithotypes record re-crystallization of the coarse-grained peridotites during downward
585 movement into the deeper portion of the supra-subduction mantle. Compared to coarse-grained peridotites,
586 the average sulfur isotope composition of matrix pentlandite from fine-grained peridotites is distinctly lower
587 ($\delta^{34}\text{S} = -0.11\text{ ‰} \pm 0.39$; 1σ , $n=15$), with few grains within the DM value of $\delta^{34}\text{S} = -1.28 \pm 0.33\text{ ‰}$ (Labidi et
588 al. 2013), as shown in Figure 7. Although the relationship between coarse- and fine-grained peridotites
589 remains unclear, the latter clearly document interaction with crust-derived hydrous fluids based on elevated
590 contents of fluid-mobile lithophile elements (Gudelius et al., 2019). This may be borne out by sulfide
591 compositions, where contents of the fluid-mobile elements As and Sb, which are highly soluble in aqueous
592 fluids (Jochum and Verma, 1996), are slightly more elevated in fine-grained than in coarse-grained
593 peridotites (Fig. 9). In addition, Se and Te abundances are markedly higher in fine-grained peridotites, and,
594 as strongly chalcophile elements (e.g., Barnes, 2016), might be expected to follow the sulfide anion. Thus,

595 formation from reduced and slightly isotopically heavy (with respect to DM) reduced S species is inferred
596 during this stage. At > 30km reduced S species carrying As(-1), Sb and Te may be sourced from the
597 subducted slab and dissolved in hot aqueous fluids (Hattori and Guillot, 2007), where they dominate over
598 oxidized species and are efficiently mobilized (Li et al., 2020).

599 Initial $^{87}\text{Sr}/^{86}\text{Sr}$ of matrix amphibole from high-pressure fine-grained peridotites is elevated, ranging from
600 0.705334 ± 0.00006 to 0.70691 ± 0.00006 , again pointing to the addition of an isotopically evolved crustal
601 component, either from the surrounding gneisses or from seawater-altered slab components (considering that
602 the $^{87}\text{Sr}/^{86}\text{Sr}$ of seawater at 330Ma was equal to 0.7078; Peterman et al., 1970). Variable equilibration of the
603 resultant fluids with unradiogenic mantle material prior to metasomatizing the UZ peridotites can explain the
604 variable $^{87}\text{Sr}/^{86}\text{Sr}$ signature. This was previously invoked to explain relatively unradiogenic *in situ* $^{87}\text{Sr}/^{86}\text{Sr}$
605 values of ~ 0.705 of matrix dolomite formed under eclogite-facies conditions (Consuma et al. 2020). The
606 heterogeneous nature of Stage 2 metasomatism is also reflected in the S concentration of fine-grained
607 peridotites, which encompasses the entire range of 40 to 288 ppm in this study. This suggests local variations
608 in terms of S concentration and/or degrees of S saturation levels due to multiple fluid infiltration, causing
609 only minor sulfide addition in some (e.g., peridotite UN-3 with a sulfide mode of 0.02 wt%) and abundant
610 sulfide addition in other samples (e.g., peridotite UN-14 with sulfide mode of 0.09 wt%). Interestingly, the
611 aforementioned peridotites were sampled from Samerberg and Malga Masa Murada, which also respectively
612 show lower and higher levels of lithophile fluid-mobile element enrichment, suggestive of more and less
613 distal positions relative to the source of the metasomatic fluids (Gudelius et al., 2019).

614

615 6.4.2. Stage 2: The sources of Ulten Zone metasomatic agents

616 The early stages of the Ulten Zone continental subduction were characterized by a continental crust possibly
617 still attached to a mature oceanic slab (Ranalli et al. 2005). In this light, possible sources capable to furnish
618 sulfur to the mantle wedge are subducted altered oceanic crust, overlying metasediments, metaserpentinites
619 and enclosing gneisses. The low $f\text{O}_2$ recorded in the investigated UZ peridotites (Fig.2c) does not preclude
620 that isotopically heavy sulfide with $\delta^{34}\text{S}$ up to +3.76 ‰ formed by reduction of oxidized S species under
621 appropriate conditions. Oxidized S could derive from seawater-altered oceanic lithosphere, which records a
622 large range of $\delta^{34}\text{S}$ values (-50 to +20 ‰) in seafloor hydrothermal sedimentary pyrite, but with a

623 predominantly negative isotopic signature (Canfield & Farquhar, 2009). In contrast, sulfates in serpentinite
624 and metabasic rocks from Syros, as analogs of subducted lithosphere, record dominantly heavier $\delta^{34}\text{S}$ of
625 +15.7 to +20.8‰ and +3.5 to +20.1‰, respectively (Schwarzenbach et al., 2018a). In exhumed high-
626 pressure rocks, metasomatic sulfides with a much larger range of $\delta^{34}\text{S}$ (-21.7 to +13.9‰) than in UZ
627 peridotites were ascribed to the combined effects of isotope fractionations due to sulfate reduction and fluid
628 evolution due to precipitation of isotopically light sulfide (Walters et al., 2019). Li et al. (2021) collate a
629 comprehensive dataset of sulfides from worldwide HP-UHP metamorphic terranes. Combined with new
630 data, they find a $\delta^{34}\text{S}$ signature of -33 to -6‰ in metasediments, $\delta^{34}\text{S}$ of -4 to +4‰ in
631 metabasite/metagabbros, and $\delta^{34}\text{S}$ of +2 to +18‰ in HP serpentinites.

632 Given the mélangé setting of Ulten Zone peridotites, the continental crust may also be a source of
633 isotopically heavy S, either by direct formation from H_2S if the lighter value applies, or by sulfate reduction
634 if the heavier value applies.

635 The availability of both S and Sr isotopic data in this study allow some constraints to be placed on the
636 sources and amount of contaminant involved in the formation of sulfides and associated clinopyroxene or
637 amphibole (Fig.10). The S isotopic composition of the garnet-kyanite gneisses of continental crust enclosing
638 the UZ peridotites is unknown, but metapelites metamorphosed under amphibolite-facies conditions from the
639 Kinzigite Formation of the Ivrea Zone ($\delta^{34}\text{S}$ values of +3.6‰ to +10.1‰; Fiorentini et al., 2018) may
640 represent reasonable analogues that can be used to model mixing with the depleted mantle. This is because
641 these metapelites are of a similar age and composition as the UZ gneisses, and, moreover, have S
642 concentrations (600 ± 10 ppm, $n=12$; Bargossi et al., 2003) similar to those employed in the mixing model.

643 Figure 10 shows the binary mixing between a depleted mantle (DM) reservoir with plausible slab-derived
644 sources in a closed system (full dataset in Electronic Annex C). The sample scatter reflects the variability of
645 the S and Sr concentrations and isotopic composition of melts/fluids and their evolution between leaving the
646 source and their interaction with the Ulten Zone peridotites. Results show that $\ll 10\%$ of a continental crust
647 component with $\delta^{34}\text{S}$ values +10.1‰ and 600ppm of S are required to explain the S-Sr isotopic composition
648 of UZ fine-grained peridotite during Stage 2. Concentrations of several 100 ppm S may be expected for a
649 crustal melt, and of several 1000 ppm S for a fluid equilibrated with crustal rock (Electronic Annex C).

650 Conversely, in accordance with trace element composition and with the lack of carbonates, the outlier CGn

651 peridotite SBA2 points to a liquid carrying isotopically heavy S, which percolated the mantle wedge in the
652 Samerberg Area, and which can be explained by invoking a third S source derived from isotopically heavy,
653 but less radiogenic serpentinite (Fig. 10). In this scenario, formation of isotopically heavy sulfide by
654 reduction of sulfate, accompanied by isotopic fractionation of several ‰ at the estimated conditions of
655 850°C (cf., Ohmoto and Rye, 1979) is not required, nor is it precluded by the data.

656

657 **6.5 Exhumation of crust-mantle mélange and retrogression (Stage 3)**

658 Shortly after the attainment of peak-P conditions, UZ peridotites became part of a crust-mantle mélange
659 (Scambelluri et al., 2006), where they were exposed to infiltration of channelized C-O-H fluids and late
660 serpentinization along a fast exhumation path in the late Carboniferous (Ranalli et al. 2005; Consuma et al.
661 2020), with conditions of 650-700 °C and P ranging between 2.2 and 1 GPa (Fig. 2a,b). Two types of
662 sulfides represent this stage and are interpreted as retrograde: (1) sulfide grains in retrograde kelyphitic
663 coronas around garnet, and (2) those included and associated with a dolomite + tremolite + chlorite vein.
664 Late pyrite in retrograde coronas reported in few studies has been interpreted as a result of late influx of S-
665 rich fluids which were produced deeper in the slab (c.f., Evans et al., 2014; Giacometti et al., 2014), possibly
666 involving redox reactions (Walters et al., 2019). Similarly, the presence of pentlandite in garnet coronas of
667 UZ peridotites suggests that partial kelyphitization was catalyzed by ingress of a S-bearing fluid. Corona-
668 associated pentlandite averages $\delta^{34}\text{S}$ of -0.38‰ (1σ , $n=3$), with negligible intra-grain variability. This grain
669 is also cut by a late chlorite flake, whose cleavage is filled with magnetite, suggesting its formation above the
670 chlorite-out reaction (i.e. at $\sim 750\text{-}800^\circ\text{C}$ and P of 1.7 GPa), that is, somewhat warmer than the conditions
671 inferred earlier (650-700°C; Consuma et al. 2020). No carbonates occur in these retrograde coronas. Low Pb,
672 As and Sb contents (Fig. 9) suggest kelyphitization at low fluid-rock ratios, consistent with incomplete
673 garnet breakdown.

674 On the basis of mineral assemblages and textural relationships, it is suggested that the pentlandite-hosting
675 dolomite vein formed prior to serpentinization at $T = 650^\circ\text{C}\text{-}700^\circ\text{C}$ and $P = 1\text{GPa}$ (Consuma et al. 2020).
676 Along this vein, enclosed pentlandite associated with millerite exhibits $\delta^{34}\text{S} = +1.13 \pm 0.38\text{‰}$, which is
677 somewhat heavier than the values of pentlandite in kelyphite and interstitial to silicates. Furthermore, a
678 distinct fluid source is otherwise required to explain the lower $\delta^{34}\text{S}$ of $0.68 \pm 0.40\text{‰}$ ($n=3$) for matrix

679 pentlandite spatially close to the dolomite vein, found in peridotite sample 18LP1. The heterogeneity of the
680 measured *in situ* Sr isotopes of this dolomite vein ($^{87}\text{Sr}/^{86}\text{Sr}=0.7035\text{-}0.7085$) was interpreted to reflect
681 retrograde hybridized C-fluids channelized into the peridotites during exhumation, when the crust-peridotite
682 mélange was already formed (Consuma et al. 2020). Coupled S and Sr isotopic ratios related to dolomite
683 veining during early exhumation (in coarse peridotite 18LP1) and even more so in serpentinite + magnesite
684 vein MOL1 -C suggest higher degrees of contamination from a continental crustal source (Fig.10). High Se
685 and Te contents, together with intermediate to high Sb and As concentrations, suggest that this was a
686 hydrous fluid carrying HS^- as the dominant S species.

687

688 **6.6 Serpentinization and local effect of fluid-rock interactions on S content and isotope composition**

689 **(Stage 4)**

690 The latest stage recording metasomatic modification of UZ peridotites is the serpentinization that occurred
691 over a span of temperatures (<600°C to 300°C) along a slow and cooling exhumation path during the
692 Permian-Triassic (300-205Ma; Ranalli et al. 2005; Consuma et al. 2020).

693 The high-T phase of Stage 4 is characterized by veining of serpentine + magnesite + tremolite crosscutting
694 the coarse spinel peridotite MOL1-C, an assemblage which indicates a temperature of formation of < 600°C
695 on the retrograde path (Consuma et al. 2020). Pentlandite grains enclosed in this vein locally show textural
696 alteration features, including (i) magnetite pseudomorphs after pentlandite and (ii) association with
697 heazlewoodite grains. The pentlandite ± heazlewoodite ± magnetite assemblages documented here and
698 previously studied in kimberlite-borne mantle xenoliths from the Kaapvaal craton (Giuliani et al. 2016;
699 Lorand and Grégoire, 2006), could originate from partial desulfidation of the low-T sulfide assemblage
700 during serpentinization, which is a common feature of mantle peridotites (e.g. Lorand, 1989).

701 Desulfurization of pentlandite to awaruite (not observed in the UZ peridotite) and/or heazlewoodite
702 represents the lowest S fugacity for a sulfide assemblage (see Fig. 8 in Lorand and Grégoire, 2006).

703 Furthermore, the relatively wide S isotopic range of these pentlandite grains ($\delta^{34}\text{S} = -0.29$ to $+3.76\text{‰}$) attests
704 to the strongly heterogeneous effect of serpentinization on the S isotope signature, and reflects minimal
705 isotopic homogenization at the vein scale. Importantly, this vein is texturally cut by late dolomite vein with
706 highly radiogenic Sr ($^{87}\text{Sr}/^{86}\text{Sr} > 0.7118$; Table 1).

707 The low-T phase of Stage 4 is characterized by dolomite dissolution via serpentinization and formation of
708 pseudomorphs of calcite + brucite intergrowths, found in several fine-grained garnet-amphibole peridotite
709 (Förster et al., 2017; Consuma et al. 2020; Gudelius et al., 2019; this study), and by the composite serpentine
710 vein MM1B hosting magnetite + millerite with minor calcite and dolomite.

711 Several PI included in coarse spinel are intimately associated with calcite + brucite intergrowths and
712 decrepitation features (i.e. radiating fractures; Fig. 3), and therefore represent Stage 4 processes.

713 Nevertheless, the $\delta^{34}\text{S}$ ($+0.46 \pm 0.34\text{‰}$) of pentlandite in PI is only moderately heavier than the suggested
714 depleted mantle with $\delta^{34}\text{S}$ of $-1.28 \pm 0.33\text{‰}$ (Labidi et al., 2013). The PI assemblages (see Section 3; Fig. 3b)
715 may point to the involvement late, C-S-bearing saline, possibly seawater-derived agents, as previously
716 suggested (Lo Pò et al., 2020). Indeed, saline, oxidizing, sulphate-bearing brines have been implicated in
717 sulfide dissolution in subduction mélanges (Schwarzenbach et al., 2018a), corresponding to the tectonic
718 setting where UZ peridotites were serpentinized. Experimental studies performed by Bataleva et al. (2018)
719 reveal that interaction of olivine with sulfate and C-bearing brines may induce mantle sulfide and carbonate
720 formation via the reaction $\text{olivine} + \text{anhydrite} + \text{C} \rightarrow \text{diopside} + \text{S}_0 + \text{CO}_2$, followed by olivine sulfidation. If
721 so, the relatively low $\delta^{34}\text{S}$ of PI sulfide requires a low-T origin when the isotopic fractionation would be
722 largest. For examples, Walters et al. (2019) model fractionations of some 20‰ for pyrite precipitation from
723 sulfate at 300 °C. Pentlandite in the PI has also moderate to high contents of the fluid-mobile elements Sb
724 and As, but low Se and Te compared to other samples (Fig. 6). This may indicate limited addition of Se and
725 Te by saline media where concentrations may be low due to limited availability of the HS^- ligand, with
726 which chalcophile elements could complex.

727 Last, in contrast to textural evidences for desulfurization and dedolomitization, high bulk S contents (> 300
728 ppm) and high As and Sb concentrations in sulfides from the serpentine vein (Fig. 9) indicates channelized
729 S-enriched fluid infiltration during Stage 4 (in the form of millerite grains) near the area of Malga Masa
730 Murada.

731 To conclude, together with the possible involvement of oxidizing H_2S -poor saline fluids in the formation of
732 PI, serpentinization appears to be associated with highly localized processes. These may involve sulfide
733 dissolution in S-undersaturated oxidizing saline fluids, perhaps sourced from metasediments
734 (Schwarzenbach et al., 2018a), sulfide precipitation when such fluids reach S saturation, and removal of

735 isotopically light S in reducing hydrous fluids, such as those associated with dehydration of oceanic mantle
736 (Piccoli et al., 2019).

737

738 **7. Conclusions and implications for coupled sulfur-carbon cycling during continent collision**

739 We provide a new perspective of sulfur and carbon mobility in continental subduction zones from the view
740 point of the previously poorly studied mantle wedge. A first-order observation is the intimate association
741 between carbonates and sulfides, which occur in different textural positions related to the multi-metamorphic
742 evolution of the Ulten Zone orogenic peridotite, as summarized in the conceptual sketch of Figure 11. Using
743 detailed petrography with bulk-rock as well as high-resolution chemical and isotopic analyses of sulfides and
744 associated metasomatic minerals (amphibole, clinopyroxene, carbonates), we demonstrate that the high-
745 pressure orogenic mantle domain from the Variscan Ulten Zone repeatedly interacted with multiple
746 metasomatic agents variably enriched in sulfur and carbon.

747 (1) The Variscan lithospheric mantle was initially depleted and sulfide-poor, and subsequently
748 inherited a sulfur component during an early stage, when hot, H₂S-bearing melts leaving a subduction-
749 modified source invaded the overlying spinel-facies peridotite in the mantle wedge, leading to precipitation
750 of sulfide-bearing pyroxenite with $\delta^{34}\text{S}_{\text{peridotite}}$ of +2.77 ‰. A reported dolomite inclusion in coarse spinel
751 indicates carbonate mobility at this early stage.

752 (2) Under peak eclogite-facies conditions, heterogeneous melt and fluid sources variably enriched in
753 carbon, isotopically heavy sulfur and radiogenic Sr were involved: (i) the positive shifts of $\delta^{34}\text{S}$ (up to
754 +3.43‰) from typical mantle values, and radiogenic $^{87}\text{Sr}/^{86}\text{Sr}_{\text{clinopyroxene}} > 0.7053$ suggest that, at high-
755 pressure conditions, coarse-grained garnet peridotites interacted with ³⁴S- and Pb-enriched crustal melts prior
756 to their becoming part of a crust-mantle mélange. Matrix carbonate is usually rare in the coarse peridotite.

757 (ii) The hydrous fluids interacting with mantle rocks during recrystallization and deformation from coarse- to
758 fine-grained peridotites added radiogenic Sr ($^{87}\text{Sr}/^{86}\text{Sr}_{\text{amphibole}} \sim 0.7061$, this study) and had a heterogeneous
759 effect on the bulk S content (58 to 288 ppm S), as S was variably added or removed. The presence of
760 dolomite with $^{87}\text{Sr}/^{86}\text{Sr} \sim 0.705$ (Consuma et al., 2020) suggests that carbon was also mobilized at this stage.
761 Given Sr isotopic evidence for a crustal fluid source, the low $\delta^{34}\text{S}$ (-1.62 to +0.67‰) compared to the coarse
762 peridotites is suggested to reflect involvement of a sulfide- rather than sulfate-bearing fluid. The observed S-

763 Sr isotopic relationship in the fine-grained peridotite can be satisfied if a depleted mantle source is
764 contaminated by <<10% of radiogenic and isotopically heavy fluids sourced from the neighboring garnet-
765 kyanite paragneisses, with minor contribution of less radiogenic metaserpentinite-derived fluid.

766 (3) Shortly after peak conditions the crust-mantle mélange interacted with hybridized fluids, leading
767 to formation of a dolomite veinlet with heterogeneous Sr isotope values ($^{87}\text{Sr}/^{86}\text{Sr} = 0.7036$ to 0.7083 ;
768 [Consuma et al., 2020](#)), which is intimately associated with (mostly) Co-pentlandite and pentlandite with $\delta^{34}\text{S}$
769 close to 0‰.

770 (4) Exhumation-related serpentinization in the crust-mantle mélange had a markedly heterogeneous
771 effect on sulfur isotopes. Pentlandite grains enclosed in a serpentine + magnesite vein crosscutting the
772 coarse-grained peridotite matrix, and texturally cut by a late dolomite vein with $^{87}\text{Sr}/^{86}\text{Sr}$ 0.711 ([Consuma et](#)
773 [al. 2020](#)), show a broad range of $\delta^{34}\text{S}$ between -0.29 and +3.76‰. Furthermore, the presence of magnetite as
774 pseudomorphs after pentlandite indicates local desulfurization with inferred preferential loss of ^{32}S , whereas
775 the S concentration of 300 ppm in a cm-sized vein of serpentine + magnetite \pm millerite \pm calcite crosscutting
776 the fine spinel peridotite indicates sulfur pathways during late exhumation stages. The presence of
777 pentlandite as part of polycrystalline inclusions in spinel associated with calcite + brucite intergrowths
778 indicates that local desulfurization may have been coupled with dissolution of the associated dolomite and
779 subsequent release of CO_2 .

780 The recorded oxidation state of the studied peridotites likely precludes stabilization of sulfate, but is
781 permissive of carbonate as opposed to graphite stability, as confirmed by petrographic observations. The
782 close association of sulfides and carbonates at all stages of mantle wedge evolution indicates that peak
783 metamorphic pervasive fluids and melts, and channelized fluids generated during late exhumation, may serve
784 as pathways for coupled sulfur and carbon transport under the investigated conditions. This indicates that
785 these volatiles should be considered together in a discussion of mantle wedge redox evolution. Despite
786 evidence for long-lasting and polyphase metasomatic processes involving slab-derived C-O-H-S fluids and
787 melts ([Fig.11](#)), the measured S isotope values of the UZ pentlandite display a narrow range ($\delta^{34}\text{S} = -1.62 \pm$
788 0.42% to $+3.76 \pm 0.36\%$), which suggests that reduction of sulfate (S^{6+}) to sulfide species (S^{2-}) had a limited
789 role throughout the Ulten Zone peridotite metamorphic evolution. Importantly, this study provides
790 petrographic and geochemical evidence in support of a dominantly reduced nature of slab-derived sulfur-

791 bearing subduction liquids in the orogenic mantle wedge under high-pressure conditions and during
792 subsequent exhumation within the tectonic mélangé, suggesting that S is not the dominant agent causing
793 oxidation of the mantle wedge in this setting.

794

795 **Acknowledgements**

796 This manuscript is part of the Ph.D. project of G.C. and has been financially supported by Almaidea-UniBO
797 and FFARB to R.B. FIERCE is financially supported by the Wilhelm and Else Heraeus Foundation and by
798 the Deutsche Forschungsgemeinschaft (DFG, INST 161/921-1 FUGG and INST 161/923-1 FUGG), which is
799 gratefully acknowledged. This is FIERCE contribution No. XX. Financial support was also provided by the
800 Australian Research Council through the Centre of Excellence for Core to Crust Fluid Systems
801 (CE1100001017), as well as the Minerals Research Institute of Western Australia.

802 M. Tribus (University of Innsbruck) is thanked for assisting G.C. during EPMA analyses at University of
803 Innsbruck. At University of Mainz, Nora Groschopf is thanked for providing top-quality XRF data, and for
804 guidance during acquisition of S concentrations, and Dejan Prelevic for facilitating the work. At Goethe-
805 University Frankfurt, the expert assistance of Alexander Schmidt in the mass spectrometry laboratory is
806 gratefully acknowledged. The authors acknowledge the technical, and the scientific assistance from the
807 Australian Microscopy & Microanalysis Research Facility at the Centre for Microscopy, Characterisation &
808 Analysis of The University of Western Australia (UWA). The School of Earth Sciences at UWA is
809 particularly thanked for hosting G.C. during her visiting research activity.

810

811 **Electronic Annexes A, B, C, D.**

812

813 **REFERENCES**

814 Alard O., Griffin W. L., Lorand J. P., Jackson S. E. and O'Reilly S. Y. (2000) Non-chondritic
815 distribution of the highly siderophile elements in mantle sulphides. *Nature* **407**, 891–894.

816 Alt J. C. (1995) Sulfur isotopic profile through the oceanic crust: Sulfur mobility and seawater-
817 crustal sulfur exchange during hydrothermal alteration. *Geology* **23**, 585.

818 Alt J. C., Shanks W. C., Bach W., Paulick H., Garrido C. J. and Beaudoin G. (2007) Hydrothermal
819 alteration and microbial sulfate reduction in peridotite and gabbro exposed by detachment
820 faulting at the Mid-Atlantic Ridge, 15°20'N (ODP Leg 209): A sulfur and oxygen isotope
821 study: HYDROTHERMAL ALTERATION PROCESSES. *Geochem. Geophys. Geosyst.* **8**.

- 822 Andersen T., Griffin W. L. and O'Reilly S. Y. (1987) Primary sulphide melt inclusions in mantle-
823 derived megacrysts and pyroxenites. *Lithos* **20**, 279–294.
- 824 Aulbach S., Stachel T., Seitz H.-M. and Brey G. P. (2012) Chalcophile and siderophile elements in
825 sulphide inclusions in eclogitic diamonds and metal cycling in a Paleoproterozoic
826 subduction zone. *Geochim. Cosmochim. Acta* **93**, 278–299.
- 827 Aulbach S., Mungall J. E. and Pearson D. G. (2016) Distribution and Processing of Highly
828 Siderophile Elements in Cratonic Mantle Lithosphere. *Rev. Mineral. Geochem.* **81**, 239–
829 304.
- 830 Aulbach S., Sun J., Tappe S. and Gerdes A. (2019) Effects of multi-stage rifting and metasomatism
831 on PGE-¹⁸⁷Os/¹⁸⁸Os systematics of the cratonic mantle beneath SW Greenland. *Contrib. Mineral.*
832 *Petrol.* **174**: 16
- 833 Bargossi G.M., Bondi M., Mordenti A., Morten L. (2003). The abundance of 55 elements and
834 petrovolumetric models of the crust in the Non and Ulten Valleys (Site 3). Accademia Nazionale
835 delle Scienze detta dei XL, Scritti e Documenti, XXXII. Published on the occasion of the 32nd
836 International Geological Congress, Florence-Italy, August 20-28, 2004, pp. 163-196.
- 837 Barnes, S.-J. (2016) Chalcophile Elements in: White, W. M. (ed.) Encyclopedia of Geochemistry.
838 Springer International Publishing, Switzerland. DOI 10.1007/978-3-319-39193-9_220-1
- 839 Bataleva Y. V., Palyanov Y. N., Borzdov Y. M. and Sobolev N. V. (2016) Sulfidation of silicate
840 mantle by reduced S-bearing metasomatic fluids and melts. *Geology* **44**, 271–274.
- 841 Bataleva Y. V., Palyanov Y. and Borzdov Y. (2018) Sulfide Formation as a Result of Sulfate
842 Subduction into Silicate Mantle (Experimental Modeling under High P,T-Parameters).
843 *Minerals* **8**, 373.
- 844 Bénard A., Klimm K., Woodland A. B., Arculus R. J., Wilke M., Botcharnikov R. E., Shimizu N.,
845 Nebel O., Rivard C. and Ionov D. A. (2018) Oxidising agents in sub-arc mantle melts link
846 slab devolatilisation and arc magmas. *Nat. Commun.* **9**.
- 847 Braga R. and Massonne H.-J. (2012) H₂O content of deep-seated orogenic continental crust: the
848 Ulten Zone, Italian Alps. *Int. Geol. Rev.* **54**, 633–641.
- 849 Braga R. and Sapienza G. T. (2007) The retrograde evolution of a dolomite-bearing hydrous
850 peridotite from the Ulten Zone (Italian Alps). *GeoActa* **6**, 37–45.
- 851 Brazhkin V. V., Popova S. V. and Voloshin R. N. (1999) Pressure–temperature phase diagram of
852 molten elements: selenium, sulfur and iodine. *Physica B: Condens. Matter* **265**, 64–71.
- 853 Canil, D., Fellows, S.A., 2017. Sulphide–sulphate stability and melting in subducted sediment and
854 its role in arc mantle redox and chalcophile cycling in space and time. *Earth Planet. Sci.*
855 *Lett.* **470**, 73–86. <https://doi.org/10.1016/j.epsl.2017.04.028>
- 856 Cannaò, E., Malaspina, N., 2018. From oceanic to continental subduction: Implications for the
857 geochemical and redox evolution of the supra-subduction mantle. *Geosphere* **14**, 2311–
858 2336. <https://doi.org/10.1130/GES01597.1>

- 859 Chaussidon M. and Lorand J.-P. (1990) Sulphur isotope composition of orogenic spinel lherzolite
860 massifs from Ariège (North-Eastern Pyrenees, France): An ion microprobe study. *Geochim.*
861 *Cosmochim. Acta* **54**, 2835–2846.
- 862 Chaussidon M., Albarède F. and Sheppard S. M. F. (1987) Sulphur isotope heterogeneity in the
863 mantle from ion microprobe measurements of sulphide inclusions in diamonds. *Nature* **330**,
864 242–244.
- 865 Consuma G., Braga R., Giovanardi T., Bersani D., Konzett J., Lugli F., Mazzucchelli M. and
866 Tropper P. (2020) In situ Sr isotope analysis of mantle carbonates: Constraints on the
867 evolution and sources of metasomatic carbon-bearing fluids in a paleo-collisional setting.
868 *Lithos* **354–355**, 105334.
- 869 Creighton S., Stachel T., Matveev S., Höfer H., McCammon C. and Luth R. W. (2009) Oxidation of
870 the Kaapvaal lithospheric mantle driven by metasomatism. *Contrib. Mineral. Petrol.* **157**,
871 491–504.
- 872 Ding T., Valkiers S., Kipphardt H., De Bièvre P., Taylor P. D. P., Gonfiantini R. and Krouse R.
873 (2001) Calibrated sulfur isotope abundance ratios of three IAEA sulfur isotope reference
874 materials and V-CDT with a reassessment of the atomic weight of sulfur. *Geochim.*
875 *Cosmochim. Acta* **65**, 2433–2437.
- 876 Eggler D. H. and Lorand J. P. (1993) Mantle sulfide geobarometry. *Geochim. Cosmochim. Acta* **57**,
877 2213–2222.
- 878 Evans B.W., Johannes, W., Oterdoom, H., Trommsdorff, V. (1976). Stability of chrysotile and
879 antigorite in the serpentinite multisystem. *Schweiz. Mineral. Petrogr. Mitt.* **56**, 79–93.
880
- 881 Evans K. A. (2012) The redox budget of subduction zones. *Earth-Sci. Rev.* **113**, 11–32.
- 882 Evans K. A., Reddy S. M., Tomkins A. G., Crossley R. J. and Frost B. R. (2017) Effects of
883 geodynamic setting on the redox state of fluids released by subducted mantle lithosphere.
884 *Lithos* **278–281**, 26–42.
- 885 Evans K. A., Tomkins A. G., Cliff J. and Fiorentini M. L. (2014) Insights into subduction zone
886 sulfur recycling from isotopic analysis of eclogite-hosted sulfides. *Chem. Geol.* **365**, 1–19.
- 887 Farquhar J. (2002) Mass-Independent Sulfur of Inclusions in Diamond and Sulfur Recycling on
888 Early Earth. *Science* **298**, 2369–2372.
- 889 Fiorentini M. L., LaFlamme C., Denyszyn S., Mole D., Maas R., Locmelis M., Caruso S. and Bui
890 T.-H. (2018) Post-collisional alkaline magmatism as gateway for metal and sulfur
891 enrichment of the continental lower crust. *Geochim. Cosmochim. Acta* **223**, 175–197.
- 892 Förster B., Braga R., Aulbach S., Pò D. L., Bargossi G. M. G. M. and Mair V. (2017) A
893 petrographic study of carbonate phases in the Ulten Zone ultramafic rocks: insights into
894 carbonation in the mantle wedge and exhumation-related decarbonation. *Ophioliti* **42**, 105–
895 127.
- 896 Fumagalli P. and Poli S. (2005) Experimentally Determined Phase Relations in Hydrous Peridotites
897 to 6.5 GPa and their Consequences on the Dynamics of Subduction Zones. *J. Petrol* **46**,
898 555–578.

- 899 Giacometti F., Evans K. A., Rebay G., Cliff J., Tomkins A. G., Rossetti P., Vaggelli G. and Adams
900 D. T. (2014) Sulfur isotope evolution in sulfide ores from Western Alps: Assessing the
901 influence of subduction-related metamorphism. *Geochem., Geophys., Geosyst.* **15**, 3808–
902 3829.
- 903 Giuliani A., Fiorentini M. L., Martin L. A. J., Farquhar J., Phillips D., Griffin W. L. and LaFlamme
904 C. (2016) Sulfur isotope composition of metasomatised mantle xenoliths from the
905 Bultfontein kimberlite (Kimberley, South Africa): Contribution from subducted sediments
906 and the effect of sulfide alteration on S isotope systematics. *Earth Planet. Sci. Lett.* **445**,
907 114–124.
- 908 Griffin W. L., Belousova E. A., Shee S. R., Pearson N. J. and O'Reilly S. Y. (2004) Archean crustal
909 evolution in the northern Yilgarn Craton: U–Pb and Hf-isotope evidence from detrital
910 zircons. *Precambrian Res.* **131**, 231–282.
- 911 Gudelius D., Aulbach S., Braga R., Höfer H. E., Woodland A. B. and Gerdes A. (2019) Element
912 Transfer and Redox Conditions in Continental Subduction Zones: New Insights from
913 Peridotites of the Ulten Zone, North Italy. *J. Petrol.* **60**, 231–268.
- 914 Guo J., Griffin W. L. and O'Reilly S. Y. (1999) Geochemistry and Origin of Sulphide Minerals in
915 Mantle Xenoliths: Qilin, Southeastern China. *J. Petrol.* **40**, 1125–1149.
- 916 Hermann J., Spandler C., Hack A. and Korsakov A. V. (2006) Aqueous fluids and hydrous melts in
917 high-pressure and ultra-high pressure rocks: Implications for element transfer in subduction
918 zones. *Lithos* **92**, 399–417.
- 919 Herzberg C. (2004) Geodynamic Information in Peridotite Petrology. *J. Petrol.* **45**, 2507–2530.
- 920 Holwell D. A., Fiorentini M., McDonald I., Lu Y., Giuliani A., Smith D. J., Keith M. and Locmelis
921 M. (2019) A metasomatized lithospheric mantle control on the metallogenic signature of
922 post-subduction magmatism. *Nat. Commun.* **10**.
- 923 de Hoog J. C. M., Mason P. R. D. and van Bergen M. J. (2001) Sulfur and chalcophile elements in
924 subduction zones: constraints from a laser ablation ICP-MS study of melt inclusions from
925 Galunggung Volcano, Indonesia. *Geochim. Cosmochim. Acta* **65**, 3147–3164.
- 926 Ionov D. A., Hoefs J., Wedepohl K. H. and Wiechert U. (1992) Content and isotopic composition
927 of sulphur in ultramafic xenoliths from central Asia. *Earth and Planetary Science Letters*
928 **111**, 269–286.
- 929 Ionov D. A., Bigot F. and Braga R. (2017) The Provenance of the Lithospheric Mantle in
930 Continental Collision Zones: Petrology and Geochemistry of Peridotites in the Ulten–
931 Nonsberg Zone (Eastern Alps). *J. Petrol.* **58**, 1451–1472.
- 932 Jenner F. E., O'Neill H. ST. C., Arculus R. J. and Mavrogenes J. A. (2010) The Magnetite Crisis in
933 the Evolution of Arc-related Magmas and the Initial Concentration of Au, Ag and Cu. *J.*
934 *Petrol.* **51**, 2445–2464.
- 935 Jochum K. P. and Verma S. P. (1996) Extreme enrichment of Sb, Tl and other trace elements in
936 altered MORB. *Chemical Geology* **130**, 289–299.
- 937 Jugo, P.J., Wilke, M., Botcharnikov, R.E., 2010. Sulfur K-edge XANES analysis of natural and
938 synthetic basaltic glasses: Implications for S speciation and S content as function of oxygen

- 939 fugacity. *Geochim. Cosmochim. Acta* 74, 5926–5938.
 940 <https://doi.org/10.1016/j.gca.2010.07.022>
- 941 Jugo P. J., Luth R. W. and Richards J. P. (2005) Experimental data on the speciation of sulfur as a
 942 function of oxygen fugacity in basaltic melts. *Geochim. Cosmochim. Acta* **69**, 497–503.
- 943 Johannes, W. and Puhon, D. (1971). The calcite-aragonite transition, reinvestigated. *Contrib.*
 944 *Mineral. Petrol.* **31**, 28–38.
- 945
 946 Kaneda H., Takenouchi S. and Shoji T. (1986) Stability of pentlandite in the Fe-Ni-Co-S system.
 947 *Miner. Deposita* **21**, 169–180.
- 948
 949 Keays R. R. (1995) The role of komatiitic and picritic magmatism and S-saturation in the formation
 950 of ore deposits. *Lithos* **34**, 1–18.
- 951 Kelley K. A. and Cottrell E. (2009) Water and the Oxidation State of Subduction Zone Magmas.
 952 *Science* **325**, 605–607.
- 953 Kiseeva E. S., Fonseca R. O. C. and Smythe D. J. (2017) Chalcophile Elements and Sulfides in the
 954 Upper Mantle. *Elements* **13**, 111–116.
- 955 Labidi J., Cartigny P. and Moreira M. (2013) Non-chondritic sulphur isotope composition of the
 956 terrestrial mantle. *Nature* **501**, 208–211.
- 957 Labidi J., Cartigny P., Hamelin C., Moreira M. and Dosso L. (2014) Sulfur isotope budget (³²S,
 958 ³³S, ³⁴S and ³⁶S) in Pacific–Antarctic ridge basalts: A record of mantle source
 959 heterogeneity and hydrothermal sulfide assimilation. *Geochim. Cosmochim. Acta* **133**, 47–
 960 67.
- 961 LaFlamme C., Martin L., Jeon H., Reddy S. M., Selvaraja V., Caruso S., Bui T. H., Roberts M. P.,
 962 Voute F., Hagemann S., Wacey D., Littman S., Wing B., Fiorentini M. and Kilburn M. R.
 963 (2016) In situ multiple sulfur isotope analysis by SIMS of pyrite, chalcopyrite, pyrrhotite,
 964 and pentlandite to refine magmatic ore genetic models. *Chem. Geol.* **444**, 1–15.
- 965 Langone, A., Braga, R., Massonne, H.-J., Tiepolo, M., 2011. Preservation of old (prograde
 966 metamorphic) U–Th–Pb ages in unshielded monazite from the high-pressure paragneisses of
 967 the Variscan Ulten Zone (Italy). *Lithos* **127**, 68–85.
- 968 Lee C.-T. y A. (2002) Platinum-group element geochemistry of peridotite xenoliths from the Sierra
 969 Nevada and the Basin and Range, California. *Geochim. Cosmochim. Acta* **66**, 3987–4005.
- 970 Li Y. and Liu J. (2006) Calculation of sulfur isotope fractionation in sulfides. *Geochimica et*
 971 *Cosmochimica Acta* **70**, 1789–1795.
- 972 Li J.-L., Schwarzenbach E. M., John T., Ague J. J., Huang F., Gao J., Klemd R., Whitehouse M. J.
 973 and Wang X.-S. (2020) Uncovering and quantifying the subduction zone sulfur cycle from
 974 the slab perspective. *Nat. Commun.* **11**, 514.
- 975 Lo Pò D., Braga R., Tropper P., Konzett J., Mair V. and Bargossi G. M. (2020) Polymineralic
 976 inclusions as tracers of multistage metasomatism in a paleo mantle wedge. *Lithos*, **364-365**,
 977 105517.

- 978 Lorand J. P. (1989) Abundance and distribution of Cu-Fe-Ni sulfides, sulfur, copper and platinum-
979 group elements in orogenic-type spinel lherzolite massifs of Arirge (northeastern Pyrenees,
980 France). *Earth Planet. Sci. Lett.* **93**, 50–64.
- 981 Lorand J.-P. and Grégoire M. (2006) Petrogenesis of base metal sulphide assemblages of some
982 peridotites from the Kaapvaal craton (South Africa). *Contrib. Miner. Petrol.* **151**, 521–538.
- 983 Lorand J.-P., Alard O., Luguët A. and Keays R. R. (2003) Sulfur and selenium systematics of the
984 subcontinental lithospheric mantle: Inferences from the Massif Central xenolith suite
985 (France). *Geochim. Cosmochim. Acta* **67**, 4137–4151.
- 986 Lorand J.-P., Luguët A. and Alard O. (2013) Platinum-group element systematics and petrogenetic
987 processing of the continental upper mantle: A review. *Lithos* **164–167**, 2–21.
- 988 Malaspina N. and Tumiati S. (2012) The role of COH and oxygen fugacity in subduction-zone
989 garnet peridotites. *Eur. J. Mineral.* **24**, 607–618.
- 990 Malaspina N., Poli S. and Fumagalli P. (2009) The oxidation state of metasomatized mantle wedge:
991 insights from C–O–H-bearing garnet peridotite. *J. Petrol.* **50**, 1533–1552.
- 992 Marini L., Moretti R. and Accornero M. (2011) Sulfur Isotopes in Magmatic-Hydrothermal
993 Systems, Melts, and Magmas. *Reviews in Mineralogy and Geochemistry* **73**, 423–492.
- 994 Marocchi M., Mair V., Tropper P. and Bargossi G. M. (2009) Metasomatic reaction bands at the
995 Mt. Hochwart gneiss-peridotite contact (Ulten Zone, Italy): insights into fluid-rock
996 interaction in subduction zones. *Mineral. Petrol.* **95**, 251.
- 997 Marocchi M., Hermann J., Tropper P., Bargossi G. M. and Mair V. (2010) Amphibole and
998 phlogopite in “hybrid” metasomatic bands monitor trace element transfer at the interface
999 between felsic and ultramafic rocks (Eastern Alps, Italy). *Lithos* **117**, 135–148.
- 1000 Marocchi M., Marschall H. R., Konzett J., Tropper P., Ludwig T., Mair V. and Bargossi G. M.
1001 (2011) Metasomatic tourmaline in hybrid contact-bands between gneiss and peridotite in the
1002 Ulten zone of the Eastern Italian Alps: chemistry and boron isotopic composition. *Can.*
1003 *Mineral.* **49**, 245–261.
- 1004 Mavrogenes J. A. and O’Neill H. St. C. (1999) The relative effects of pressure, temperature and
1005 oxygen fugacity on the solubility of sulfide in mafic magmas. *Geochim. Cosmochim. Acta*
1006 **63**, 1173–1180.
- 1007 McDonough W. F. and Sun S. (1995) The composition of the Earth. *Chem. Geol.* **120**, 223–253.
- 1008 McInnes B. I. A., Gregoire M., Binns R. A., Herzig P. M. and Hannington M. D. (2001) Hydrous
1009 metasomatism of oceanic sub-arc mantle, Lihir, Papua New Guinea: petrology and
1010 geochemistry of fluid-metasomatised mantle wedge xenoliths. *Earth Planet. Sci. Lett.* **188**,
1011 169–183.
- 1012 Métrich N., Schiano P., Clocchiatti R. and Maury R. C. (1999) Transfer of sulfur in subduction
1013 settings: an example from Batan Island (Luzon volcanic arc, Philippines). *Earth Planet. Sci.*
1014 *Lett.* **167**, 1–14.

- 1015 Mungall J. E., Hanley J. J., Arndt N. T. and Debecdelievre A. (2006) Evidence from meimechites
1016 and other low-degree mantle melts for redox controls on mantle-crust fractionation of
1017 platinum-group elements. *Proc. Natl. Acad. Sci.* **103**, 12695–12700.
- 1018 Nimis P. and Morten L. (2000) P–T evolution of ‘crustal’ garnet peridotites and included
1019 pyroxenites from Nonsberg area (upper Austroalpine), NE Italy: from the wedge to the slab.
1020 *J. Geodyn.* **30**, 93–115.
- 1021 O’Neill H. St. C. (1991) The origin of the moon and the early history of the earth—A chemical
1022 model. Part 2: The earth. *Geochim. Cosmochim. Acta* **55**, 1159–1172.
- 1023 Obata M. and Morten L. (1987) Transformation of spinel lherzolite to garnet lherzolite in ultramafic
1024 lenses of the Austridic crystalline complex, northern Italy. *J. Petrol.* **28**, 599–623.
- 1025 Ohmoto, H., & Rye, R. O. (1979). Isotopes of sulfur and carbon. In H. L. Barnes (Ed.),
1026 *Geochemistry of Hydrothermal Ore Deposits*, (pp. 517–611). New York: Wiley.
- 1027 Ono S., Keller N. S., Rouxel O. and Alt J. C. (2012) Sulfur-33 constraints on the origin of
1028 secondary pyrite in altered oceanic basement. *Geochim. Cosmochim. Acta* **87**, 323–340.
- 1029 Palme H. and O’Neill H. St. C. (2007) Cosmochemical Estimates of Mantle Composition. In
1030 *Treatise on Geochemistry* Elsevier. 1–38.
- 1031 Papike J. J., Spilde M. N., Fowler G. W., Layne G. D. and Shearer C. K. (1995) The Lodran
1032 primitive achondrite: petrogenetic insights from electron and ion microprobe analysis of
1033 olivine and orthopyroxene. *Geochim. Cosmochim. Acta* **59**, 3061–3070.
- 1034 Peterman Z. E., Hedge C. E. and Tourtelot H. A. (1970) Isotopic composition of strontium in sea
1035 water throughout Phanerozoic time. *Geochim. Cosmochim. Acta* **34**, 105–120.
- 1036 Piccoli F., Hermann J., Pettke T., Connolly J. A. D., Kempf E. D. and Vieira Duarte J. F. (2019)
1037 Subducting serpentinites release reduced, not oxidized, aqueous fluids. *Sci. Rep.* **9**, 1–7.
- 1038 Poulson S. R., Kubilius W. P. and Ohmoto H. (1991) Geochemical behavior of sulfur in granitoids
1039 during intrusion of the South Mountain batholith, Nova Scotia, Canada. *Geochim.*
1040 *Cosmochim. Acta* **55**, 3809–3830.
- 1041 Rampone E. and Morten L. (2001) Records of crustal metasomatism in the garnet peridotites of the
1042 Ulten Zone (Upper Austroalpine, Eastern Alps). *J. Petrol.* **42**, 207–219.
- 1043 Regorda A., Lardeaux J.-M., Roda M., Marotta A. M. and Spalla M. I. (2020) How many
1044 subductions in the Variscan orogeny? Insights from numerical models. *Geosci. Front.* **11**,
1045 1025–1052.
- 1046 Rielli A., Tomkins A. G., Nebel O., Raveggi M., Jeon H., Martin L. and Ávila J. N. (2018) Sulfur
1047 isotope and PGE systematics of metasomatised mantle wedge. *Earth Planet. Sci. Lett.* **497**,
1048 181–192.
- 1049 Salters V. J. and Stracke A. (2004) Composition of the depleted mantle. *Geochem. Geophys.*
1050 *Geosyst.* **5**, Q05B07.

- 1051 Sapienza G. T., Scambelluri M. and Braga R. (2009) Dolomite-bearing orogenic garnet peridotites
1052 witness fluid-mediated carbon recycling in a mantle wedge (Ulten Zone, Eastern Alps,
1053 Italy). *Contrib. Mineral. Petrol.* **158**, 401–420.
- 1054 Scambelluri M., Hermann J., Morten L. and Rampone E. (2006) Melt-versus fluid-induced
1055 metasomatism in spinel to garnet wedge peridotites (Ulten Zone, Eastern Italian Alps): clues
1056 from trace element and Li abundances. *Contrib. Mineral. Petrol.* **151**, 372.
- 1057 Scambelluri M., Rampone E., Braga R. and Malaspina N. (2010) The Variscan garnet peridotites
1058 from the Eastern Alps (Ulten Zone): records of subduction metasomatism in the mantle
1059 wedge. In: (Eds.) Marco Beltrando, Angelo Peccerillo, Massimo Mattei, Sandro Conticelli,
1060 and Carlo Doglioni, *J. Virtual Explor.* **36**. doi: 10.3809/jvirtex.2009.00259
- 1061 Schramke J.A., Kerrick D.M. and Blencoe J.G. (1982) Experimental determination of the brucite
1062 = periclase + water equilibrium with a new volumetric technique. *Am. Min.* **67**, 269–276.
- 1063 Schwarzenbach E. M., Caddick M. J., Petroff M., Gill B. C., Cooperdock E. H. G. and Barnes J. D.
1064 (2018a) Sulphur and carbon cycling in the subduction zone mélangé. *Sci. Rep.* **8**, 15517.
- 1065 Schwarzenbach E. M., Gill B. C. and Johnston D. T. (2018b) Unraveling multiple phases of sulfur
1066 cycling during the alteration of ancient ultramafic oceanic lithosphere. *Geochim.*
1067 *Cosmochim. Acta* **223**, 279–299.
- 1068 Sun S.-S. (1982) Chemical composition and origin of the earth's primitive mantle. *Geochim.*
1069 *Cosmochim. Acta* **46**, 179–192.
- 1070 Sun S.-S., Wallace D. A., Hoatson D. M., Glikson A. Y. and Keays R. R. (1991) Use of
1071 geochemistry as a guide to platinum group element potential of mafic-ultramafic rocks:
1072 examples from the west Pilbara Block and Halls Creek Mobile Zone, Western Australia.
1073 *Precambrian Res.* **50**, 1–35.
- 1074 Szabó Cs. and Bodnar R. J. (1995) Chemistry and origin of mantle sulfides in spinel peridotite
1075 xenoliths from alkaline basaltic lavas, Nógrád-Gömörr Volcanic Field, northern Hungary
1076 and southern Slovakia. *Geochim. Cosmochim. Acta* **59**, 3917–3927.
- 1077 Thomassot E., Cartigny P., Harris J. W., Lorand J. P., Rollion-Bard C. and Chaussidon M. (2009)
1078 Metasomatic diamond growth: A multi-isotope study (¹³C, ¹⁵N, ³³S, ³⁴S) of sulphide
1079 inclusions and their host diamonds from Jwaneng (Botswana). *Earth Planet. Sci. Lett.* **282**,
1080 79–90.
- 1081 Tomkins A. G. and Evans K. A. (2015) Separate zones of sulfate and sulfide release from
1082 subducted mafic oceanic crust. *Earth Planet. Sci. Lett.* **428**, 73–83.
- 1083 Tumiati S., Thöni M., Nimis P., Martin S. and Mair V. (2003) Mantle–crust interactions during
1084 Variscan subduction in the Eastern Alps (Nonsberg–Ulten zone): geochronology and new
1085 petrological constraints. *Earth Planet. Sci. Lett.* **210**, 509–526.
- 1086 Tumiati S., Fumagalli P., Tiraboschi C. and Poli S. (2013) An Experimental Study on COH-bearing
1087 Peridotite up to 3.2 GPa and Implications for Crust–Mantle Recycling. *J. Petrol.* **54**, 453–
1088 479.
- 1089 Vaughan D. J. and Corkhill C. L. (2017) Mineralogy of Sulfides. *Elements* **13**, 81–87.
1090

- 1091 Vitale Brovarone, A., Tumiati, S., Piccoli, F., Ague, J. J., Connolly, J. A. D., & Beyssac, O. (2020).
 1092 Fluid-mediated selective dissolution of subducting carbonaceous material: Implications for
 1093 carbon recycling and fluid fluxes at forearc depths. *Chemical Geology* **549**, 119682.
- 1094 Voshage H., Hofmann A. W., Mazzucchelli M., Rivalenti G., Sinigoi S., Raczek I. and Demarchi
 1095 G. (1990) Isotopic evidence from the Ivrea Zone for a hybrid lower crust formed by
 1096 magmatic underplating. *Nature* **347**, 731–736.
- 1097 Walters J. B., Cruz-Uribe A. M. and Marschall H. R. (2019) Isotopic Compositions of Sulfides in
 1098 Exhumed High-Pressure Terranes: Implications for Sulfur Cycling in Subduction Zones.
 1099 *Geochem. Geophys. Geosyst.* **20**, 3347-3374.
- 1100 Walters J. B., Cruz-Uribe A. M. and Marschall H. R. (2020) Sulfur loss from subducted altered
 1101 oceanic crust and implications for mantle oxidation. *Geochem. Perspect. Lett.* **13**, 36–41.
- 1102 Wang, K.-L., O'Reilly, S.Y., Honda, M., Matsumoto, T., Griffin, W.L., Pearson, N.J., Zhang, M.,
 1103 2010. Co-rich sulfides in mantle peridotites from Penghu Islands, Taiwan: Footprints of
 1104 Proterozoic mantle plumes under the Cathaysia Block. *J. Asian Earth Sci.* **37**, 229–245.
 1105 <https://doi.org/10.1016/j.jseae.2009.08.008>
- 1106 Workman R. K. and Hart S. R. (2005) Major and trace element composition of the depleted MORB
 1107 mantle (DMM). *Earth Planet. Sci. Lett.* **231**, 53–72.
- 1108 Zhang Z., Lentsch N. and Hirschmann M. M. (2015) Carbon-saturated monosulfide melting in the
 1109 shallow mantle: solubility and effect on solidus. *Contrib. Miner. Petrol.* **170**, 47.
 1110
 1111

Figure captions

Preference color: online only

Figure 1 (a) Geological sketch map of the central and Eastern Alps modified after [Langone et al. \(2011\)](#). (b) Geological-structural map of the Ulten Zone modified after [Tumiati et al. \(2003\)](#) and [Consuma et al. \(2020\)](#). The reader is referred to [Consuma et al. \(2020\)](#) and ([Gudelius et al., 2019](#)) for the exact location of the samples.

Figure 2. (a-b) Pressure (P)-Temperature (T) diagram modified after [Consuma et al. \(2020\)](#) showing the pressure-temperature evolution of the Ulten Zone peridotites, with respect to various metasomatic stages and phase relations for COH-bearing species. The experimentally determined melting curves of sulfur, pyrrhotite, Mss and decomposition of pyrite are taken from [Brazhkin et al. \(1999\)](#), [Zhang et al. \(2015\)](#) and [Bataleva et al. \(2018\)](#). Dolomite-in, magnesite-in, dolomite-out, garnet-in curves from [Tumiati et al. \(2013\)](#); tremolite (Tr) + magnesite (Mgs) = enstatite (En) + dolomite (Dol) are from [Malaspina and Tumiati \(2012\)](#) for $X_{\text{CO}_2}=0.5$; brucite (Brc) = periclase (Per) + water (H₂O) from [Schramke et al. \(1982\)](#). Antigorite-out curve and chlorite-out curve from ([Fumagalli and Poli, 2005](#)). Antigorite (Atg) = chrysotile (Ctl) / lizardite (Liz) are from [Evans et al. \(1976\)](#); aragonite (Arg) = calcite (Cal) from [Johannes and Puhon \(1971\)](#). P-T conditions of Stage 2 are taken from [Nimis and Morten \(2000\)](#) and [Sapienza et al. \(2009\)](#). **Sulf**—sulfide, **Po**—pyrrhotite, **Py**—pyrite, **Mss**—monosulfide solid solution.

(c) Ranges and average values of oxygen fugacity relative to the fayalite–magnetite–quartz buffer (FMQ) for the Ulten Zone peridotites for various pressures (GPa), as well as CHO speciation, modified from Cannà and Malaspina (2018) and Gudelius et al. (2019). Note the offset to higher fO_2 obtained by Malaspina et al. (2009), who, using the same method, obtained higher $Fe^{3+}/\Sigma Fe$ in garnet from their samples. Fields denote fO_2 where most S in basalt glass (Jugo et al., 2010) and in sediment glass (Canil and Fellows, 2017) is present as HS^- ($\sim FMQ$ and maximum $FMQ < +0.2$, respectively) and as $H_2SO_4^{2-}$ ($\sim FMQ + 2$ and minimum $FMQ > +0.2$, respectively), with mixed species at intermediate fO_2 in the basalt.

(d) Table showing the presence of rock-forming minerals, hydrous minerals, carbonates and sulfides as linked to distinct metamorphic stages (1-4) of the Ulten Zone peridotites. Grain size is qualitatively indicated for spinel, orthopyroxene, clinopyroxene and garnet. *In situ* Sr isotope compositions are reported for clinopyroxene, amphibole (this study) and for dolomite, calcite (Consuma et al., 2020). S isotope compositions of pentlandite measured by SIMS (this study) are also shown. **HT**—high temperature, **HP**—high pressure, **grt**—garnet.

Figure 3. Photomicrographs, back-scattered electron images, and color maps illustrating the textural settings of sulfides in Ulten Zone peridotites. (a) Scan image of thin section P10B with stars indicating locations of polyphase inclusions (PI) in garnet and matrix pentlandite (modified after Lo Pò et al. 2020) along with a BSE image of a PI and relative color mapping showing major element compositions of Po + Pn. (b) Coarse spinel enclosing PI. Note calcite + brucite intergrowths associated with pentlandite (KL24-2b). PI in spinel with ilmenite + chlorite + apatite + pentlandite (KL24-3); (c) matrix pentlandite in intimate association with dolomite (MOL1-C) and in close relationship with dolomite (VM10A). (d) kelyphite corona enclosing pentlandite cut by a chlorite flake and magnetite string; (e) dolomite vein (+ magnesite) enclosing pentlandite + millerite cut by chlorite flake (18LP1); magnesite + serpentine + tremolite vein enclosing pentlandite partially replaced by magnetite pseudomorphs (MOL1-C); (f) halos and apophyses around pentlandite (UN3) and pentlandite-fracture fillings (UN14).

Figure 4. (a) FeO and (b) TiO₂ (wt%) as a function of Al₂O₃ of whole-rock peridotite analyses from the Ulten Zone investigated in this study (data in Table S3, Obata and Morten, 1987, Ionov et al. 2017 and Gudelius et al., 2019). Shown for comparison in (a) are residues from batch melting at 2, 4 and 6 GPa (stippled blue lines) and from polybaric fractional melting of primitive mantle (**PM**—yellow star) at 2-0, 3-0, 5-1 and 7-2 GPa (thick red lines), as well as contours of % melt extracted (orange lines) (from Herzberg, 2004). Shown for comparison in (b) are paths for polybaric fractional melting of PM at 1.5-0.5 and 2.5-0.5 GPa. Low TiO₂ contents at a given Al₂O₃ in the UZ peridotites compared to the model suggest formation from a more depleted source. (c) NiO (wt.%) content of olivine grains vs bulk S concentration (ppm). Values of NiO taken from Gudelius et al. (2019) and Consuma et al. (2020). Correlation fitting lines and respective R² values are reported.

Figure 5. Sulfur contents (in ppm S) in UZ peridotites (this study) compared to those in different mantle and crustal reservoirs (references in panel).

Figure 6. Primitive-mantle normalized trace element compositions of pentlandite and pyrrhotite (primitive mantle of McDonough and Sun, 1995). For simplicity, only medians for coarse and fine peridotites, and for different textural positions are reported. Element order corresponds to decreasing compatibility during partial melting of primitive mantle (from Aulbach et al., 2012). Shown for comparison in (a) is range of abundances in pyrrhotite included in eclogitic diamond and obtained using the same method employed in this study (Aulbach et al., 2012). Shown for comparison in (c) is the range of matrix pentlandite compositions shown in (b). **CS**—Coarse-

grained spinel peridotite, **CGn**—Carbonate-free coarse-grained garnet peridotite, **CG**—Coarse-grained garnet/spinel with coronitic garnet peridotite, **(P)FGA**—Porphyroclastic and non-porphyroclastic fine-grained garnet amphibole peridotite, **FSA**—Fine-grained spinel (chlorite-amphibole) peridotite. **Po**—pyrrhotite, **Pn**—pentlandite.

Figure 7. (a) Sulfur isotope variations of this study compared to those in different reservoirs (Giuliani et al., 2016, and references therein), in high-pressure rocks reported in Li et al. (2020, and references therein), in slab fluids estimated by Walters et al. (2019) and Li et al. (2020), pelagic sediments from Schwarzenbach et al. (2018) and Li et al. 2020, and in sulfides (green line) and sulfates (orange line) from altered oceanic lithosphere and serpentinite (Schwarzenbach et al., 2018a,b, Li et al. 2020); (b) Histograms showing the sulfur isotope compositions in pentlandite in various textural settings measured by SIMS (this study). The yellow box corresponds to the depleted mantle value range, following Labidi et al. 2013 ($\delta^{34}\text{S} = -1.28 \pm 0.25\%$) and Labidi et al., (2014) ($\delta^{34}\text{S} = -1.40 \pm 0.50\%$). **CS**—Coarse-grained spinel peridotite, **CGn**—Carbonate-free coarse-grained garnet peridotite, **CG**—Coarse-grained garnet/spinel with coronitic garnet peridotite, **(P)FGA**—Porphyroclastic plus non-porphyroclastic fine-grained garnet amphibole peridotite, **FSA**—Fine-grained spinel (chlorite-amphibole) peridotite. **PI**—polycrystalline inclusions. **VCDT**—Vienna Canyon Diablo Troilite standard.

Figure 8. *In situ* Sr isotope ratios of (a) amphibole and clinopyroxene compared to As concentrations in the associated pentlandite. Values are reported as total averages for the different lithotypes. **CS**—Coarse-grained spinel peridotite, **CG**—Coarse-grained garnet/spinel with coronitic garnet peridotite, **(P)FGA**—Porphyroclastic and non-porphyroclastic fine-grained garnet amphibole peridotite, **FSA**—Fine-grained spinel (chlorite-amphibole) peridotite. (b) *In situ* Sr isotope compositions of clinopyroxene vs S isotope of pentlandite (this study) and vs *in situ* Sr isotope compositions of carbonates (Consuma et al. 2020). The small number of data points is due to the rare occurrence of clinopyroxene combined with the typically small grain size of clinopyroxene and carbonates, which precludes laser ablation sampling of large enough volumes for statistically meaningful Sr isotope analysis. Further limitations are imposed by the requirement of using matrix-matched sulfide grains for SIMS analyses.

Figure 9. Trace element concentrations (Co, Se, Te, Pb, Sb, As) versus $\delta^{34}\text{S}$ measured in pentlandite in the Ulten Zone peridotites distinguished by textural positions. When significant, correlation fitting lines and respective R^2 are reported. **CS**—Coarse-grained spinel peridotite, **CGn**—Carbonate-free coarse-grained garnet peridotite, **CG**—Coarse-grained garnet/spinel with coronitic garnet peridotite, **(P)FGA**—Porphyroclastic and non-porphyroclastic fine-grained garnet amphibole peridotite, **FSA**—Fine-grained spinel (chlorite-amphibole) peridotite. **Po**—pyrrhotite, **Pn**—pentlandite.

Figure 10. $\delta^{34}\text{S}$ in pentlandite (this study) vs. $^{87}\text{Sr}/^{86}\text{Sr}$ in coexisting carbonate, clinopyroxene and amphibole (this study and Consuma et al., 2020). Note that MOL1-C refers to the carbonates (+serpentine)-bearing vein crosscutting the coarse spinel peridotite of Figure 3e. Mixing trends between depleted mantle (DM) source and possible slab-derived melt and fluid sources. The continental crustal component may be a melt or a fluid with variable S concentrations as shown. A single Sr concentration (130 ppm) is assumed for the purpose of modeling, since in our study the Sr concentration in fluid and melt is estimated to vary less than two-fold. Light grey area and individual data for glasses dredged on the South Atlantic ridge from Labidi et al. (2013). Input data with references, as well as the effect of varying Sr concentration on the curvature of the mixing lines, are reported in Electronic Annex C.

Figure 11. Conceptual model of the metamorphic stages of the Ulten Zone peridotites, see discussion for details. **Stage 1**— Only a small residual sulfur component would have remained after ca. 15-25% of melt extraction. Concentrations of 170-210 ppm S with a heavy $\delta^{34}\text{S}$ signature may reflect percolation by reduced, H_2S -bearing hydrous basic melts from the subduction-modified deeper lithospheric mantle. **Stage 2**—At peak (garnet) conditions, when UZ peridotites still reside in the mantle wedge, sulfur is introduced into peridotites from heterogeneous melt and fluid sources, which variably equilibrated with the depleted mantle and possibly mixed with de-serpentinization fluids. **Stage 3**— The supra-depleted mantle S isotope signature again require addition of heavy sulfur. **Stage 4**—Sulfur and CO_2 are variously released from the mantle-crust mélange during late-stage serpentinization, while high S concentrations (>300 ppm) in a serpentine vein suggest efficient mobilization in channelized fluids.