

This is the final peer-reviewed accepted manuscript of:

Martina Vagnoni, Chiara Samorì, Daniele Pirini, Maria Katrina Vasquez De Paz, Dawit Gebremichael Gidey, Paola Galletti, 2022. “Lipase catalysed oxidations in a sugar-derived natural deep eutectic solvent”. *Biocatalysis and Biotransformation*, 40(6), 422-431

The final published version is available online at:
<https://doi.org/10.1080/10242422.2021.1913126>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1 **Lipase catalyzed oxidations in a sugar-derived Natural Deep Eutectic**
2 **Solvent**

3
4 Martina Vagnoni*^a, Chiara Samorì^a, Daniele Pirini^a, Maria Katrina Vasquez De Paz,
5 Dawit Gebremichael Gidey, Paola Galletti^a

6
7 ^a Department of Chemistry “Giacomo Ciamician”, University of Bologna, Ravenna
8 Campus, Via S. Alberto 163, Ravenna (Italy).

9
10 *email: martina.vagnoni3@unibo.it

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57

58 **Abstract**

59 Chemoenzymatic oxidations involving the CAL-B/H₂O₂ system was developed in a sugar derived
60 Natural Deep Eutectic Solvent (NaDES) composed by a mixture of glucose, fructose and sucrose.
61 Good to excellent conversions of substrates like cyclooctene, limonene, oleic acid and stilbene to
62 their corresponding epoxides, cyclohexanone to its corresponding lactone and 2-
63 phenylacetophenone to its corresponding ester, demonstrate the viability of the sugar NaDES as a
64 reaction medium for epoxidation and Baeyer-Villiger oxidation.

65
66 **Keywords**

67 Chemoenzymatic oxidations, green solvents, Natural Deep Eutectic Solvents, Epoxidation, Baeyer-
68 Villiger, Lipase

69
70
71 **1. Introduction**

72 Oxidation reactions have always been a major area of research due to their tremendous industrial
73 applications. However, several oxidation processes present sustainability issues from the point of
74 view of oxidants, catalysts and solvents used. (Cavani & Teles 2009) For example, a peracid is
75 often used as the oxidizing agent, (Swern 1949) but transportation and storage of organic peracids
76 leads to significant safety issues and costs; when achievable, molecular oxygen or air are for sure
77 the ideal oxidants, with hydrogen peroxide as the second-best choice, in terms of atom economy
78 and applicability to various oxidation systems. (O. Burek et al. 2019) In this context, enzymes, that
79 can work in sustainable solvents with mild oxidants, can contribute to increase the greenness of the
80 oxidation reactions. (Niku-Paavola & Viikari 2000; Constable et al. 2007; Gorke et al. 2008;
81 Kotlewska et al. 2011; Silva et al. 2011; Hollmann et al. 2011; Drożdż et al. 2015; Qin et al. 2015;
82 Yin et al. 2015; Yang & Duan 2016; García et al. 2018)

83 In fact, a very interesting system for obtaining peracids *in situ* is the chemoenzymatic system
84 lipase/H₂O₂ that continuously forms the peracids through a lipase-catalyzed perhydrolysis of
85 carboxylic acids or their esters. (Björkling et al. 1990; Yadav & Devi 2002; Busto et al. 2010) A
86 broad range of hydrolases has been investigated for the peracid formation and among them the
87 lipase B from *Candida Antarctica* (CAL-B), immobilized onto an acrylic resin (Novozyme 435) is
88 the most reactive. (Ortiz et al. 2019) This system has been successfully applied to both epoxidations
89 of alkenes (Prileshajev-epoxidation) and Baeyer-Villiger (B-V) oxidations (Scheme 1). (Lemoult et
90 al. 1995; Aouf et al. 2014)

91 Epoxides are fundamental intermediates in organic synthesis but, despite their relevance, their
92 industrial synthesis is scarcely sustainable (both environmentally and economically). Epoxidation of
93 some natural products is industrially carried out by the Prileshajev-epoxidation (epoxidation of an
94 alkene with a peracid) using either preformed or in-situ-generated short chain peroxy acids. (Rüsch
95 gen. Klaas & Warwel 1999; Hilker et al. 2001) Nevertheless, the need for a strong acid to catalyze
96 peroxy acid formation in this process can result in unsatisfactory selectivity and undesirable side
97 reactions *via* oxirane ring opening, leading to diols, hydroxyesters, and dimers. Prileshajev-
98 epoxidation can be chemoenzymatically carried out on various substrates with CAL-B, a carboxylic
99 acid as precursor of the peracid, and H₂O₂ as oxidant (Scheme 1); this method allows an
100 improvement in terms of sustainability, mild reaction conditions, limited side products and use of
101 less toxic reagents. (Niku-Paavola & Viikari 2000; Moreira & Nascimento 2007; Silva et al. 2011;
102 Hollmann et al. 2011)

103 The B-V oxidation is a very well-known and useful reaction for the synthesis of esters and lactones
104 starting from ketones, important building blocks in pharmaceutical and polymer synthesis. (Renz &
105 Meunier 1999; Brink et al. 2004; Woodruff & Hutmacher 2010) Peracids such as meta-
106 chloroperbenzoic acid or peracetic acid are used as stoichiometric reagents, but also various
107 catalytic methods that use metals have been studied. (Strukul 1998; Ma et al. 2014) Protocols based
108 on B-V monooxygenases have also been developed, but given their need for oxygen, cofactor

109 NADPH and their intrinsic low stability, they are considered unpractical. (Alphand et al. 2003;
110 Leisch et al. 2011; Balke et al. 2012) Nevertheless, the simple chemoenzymatic method based on
111 CAL-B, used for epoxidation of alkenes described above, has been also applied to this kind of
112 oxidations (Scheme 1). (Lemoult et al. 1995; Ríos et al. 2007; Rios et al. 2008)
113 Green solvents exploited in chemoenzymatic oxidation reactions can be categorized into two main
114 groups: i) water and ii) non-aqueous solvents like ionic liquids, (Moniruzzaman et al. 2010;
115 Elgharbawy et al. 2020) supercritical fluids and fluorinated solvents. (Hobbs & Thomas 2007)
116 Despite their interesting properties and application possibility, ionic liquids suffer from several
117 drawbacks like cost, toxicity, low biodegradability, complexity in preparation and handling.
118 (Samorì et al. 2015; Lei et al. 2017; B. Wang et al. 2017) Deep Eutectic Solvents (DESs), described
119 for the first time by *Abbott et al.*, (Abbott et al. 2001) are low melting mixtures based on a
120 combination of readily-available and inexpensive components, like quaternary ammonium salts as
121 hydrogen bond acceptors (HBA), and acids, amides, amines, carbohydrates and alcohols as
122 hydrogen bond donors (HDB). They are liquid at or below 100 °C, thanks to H-bond interactions
123 between the single components that create specific supramolecular structures. The number and the
124 spatial positions of hydrogen atoms in the donor and acceptor groups, available for hydrogen
125 bonding, influence the formation and stability of the DES itself. (Nkuku & LeSuer 2007; Zhang et
126 al. 2012; Paiva et al. 2014; Smith et al. 2014; Tommasi et al. 2017; Samorì et al. 2019) Dai *et al.*
127 reported numerous preparations of Natural Deep Eutectic Solvents (NaDESs) by using plant
128 metabolites. Interestingly, when water is added to the mixtures, in different proportions according to
129 the NaDES, it can be incorporated into this structure and becomes strongly bound, reducing the
130 viscosity of the NaDES while retaining its original characteristics. (Dai et al. 2013)
131 The chemoenzymatic oxidation systems described above (Scheme 1) have been studied in several
132 solvents, including DESs (T. Gorke et al. 2008; Kotlewska et al. 2011; Durand et al. 2012; Drożdż
133 et al. 2015; Yang & Duan 2016; Zhou et al. 2017; Ranganathan et al. 2017; Gotor-Fernández &
134 Paul 2019; Ma et al. 2019), in which it has been demonstrated that the enzymatic activity, stability,
135 and selectivity can be enhanced. (Zhou et al. 2017; Ülger & Takaç 2017; Oh et al. 2019; Guajardo
136 et al. 2019; Gotor-Fernández & Paul 2019; Nian et al. 2020). Among the various NaDES proposed
137 by Dai *et al.* we focused on the only one sugar-derived and chlorine-free combination, composed by
138 glucose, fructose sucrose and water (1:1:1:11), that we called with the acronym GFS. To the best of
139 our knowledge the lipase/H₂O₂ system was never reported in a solvent like GFS and, following our
140 interest in biocatalysis in sustainable reaction media, (Galletti et al. 2007) herein we report on its
141 application in the epoxidation of alkenes and B-V oxidation of ketones.

142 143 **2. Materials and methods**

144 2.1 Material: all chemicals and solvents were purchased from Sigma-Aldrich or Alfa Aesar and
145 used without any further purification.

146 CAL-B (Lipase B from *Candida antarctica*) immobilized on Immobead 150, recombinant from
147 yeast, 4000 U/g was used.

148 2.2 DESs preparation: the components were mixed with the appropriate stoichiometric ratios,
149 heated at about 80-90 °C (120 °C for GFS) and magnetically stirred until homogeneous liquid was
150 obtained; for GFS, distilled water (up to 30 wt %) was then added to get a homogeneous colorless
151 liquid phase. All the DESs were cooled to rt (20 °C) before the use and stored in the fridge (4 °C).

152 2.3 Representative procedure for enzymatic epoxidations of alkenes: in a 4-mL vial, the
153 immobilized CAL-B (amounts reported in Tables 1-4 and Scheme 4) and 400 mg of DES (200 mg
154 for **1d**, 800 mg for **1e**) were weighted, followed by the addition of 1.6 mmol of alkene (0.8 mmol
155 for **1f**), carboxylic acid (amounts reported in Tables 1-4 and Scheme 4) and 1 equivalent (eq) of
156 H₂O₂ (30% aqueous solution). For entries 17 and 18 in Table 1, H₂O₂ has been added in 4 aliquots
157 in 4 h, for substrate **1f** 1.5 eq has been used.

158 The vial was heated at 45 °C (or rt, 20 °C, for **1d**) for various reaction times, then crudes were
159 extracted with cyclohexane or ethyl acetate and analyzed by GC-MS. Extraction residues were

160 checked after derivatization by silylation for the presence of other by-products (see section 2.5).
161 Conversions were calculated as ratios between products areas and total areas. ^1H and ^{13}C NMR
162 spectra of some products have been acquired after purification of the crude by flash-column
163 chromatography (see section 2.6), some isolated yields are also reported in the Tables. All products
164 are known, they were recognized by comparison with standards or through mass spectra matching
165 to what reported in NIST database. Formation of byproducts was checked by GC-MS and NMR.

166 2.4 Representative procedure for B-V oxidation of ketones: in a 4-mL vial, the immobilized CAL-B
167 (amounts in Tables 3-4) and 400 mg of GFS for **3a**, **3b**, **3c**, (700 mg of GFS for **3d**) were weighted,
168 followed by the addition of 0.8 mmol of ketone (0.4 mmol for **3d**), carboxylic acid (amounts in
169 Tables 3-4) and 1 eq of H_2O_2 (30% aqueous solution), different amounts of H_2O_2 are reported in
170 Table 3. The vial was heated at 45 °C or at kept at rt, 20 °C, for various reaction times, then crudes
171 were extracted with ethyl acetate and analyzed by GC-MS as described above. ^1H and ^{13}C NMR
172 spectra of some purified products have been acquired after purification (see section 2.6) of the crude
173 by flash-column chromatography, some isolated yields are also reported in Tables 3-4.

174 2.5 Silylation procedure: 50 μL of silylating agent *N,O*-bis(trimethylsilyl)trifluoroacetamide and
175 1% chlorotrimethylsilane, (BSTFA + 1% TMCS), 100 μL of CH_3CN and 20 μL of pyridine were
176 added to 1–10 mg of sample into a GC-MS vial. The vial was heated at 60–80 °C for 30-40 min.
177 The sample was then diluted with CH_3CN before the injection.

178 2.6 Purification procedure of selected products: reaction mixtures were extracted with ethyl acetate
179 or cyclohexane then washed with a NaHCO_3 solution to remove the octanoic acid (OA). After
180 evaporating the solvent, the crude was purified by flash chromatography. The fractions containing
181 the product were mixed, the solvent evaporated, and the purified products were analyzed by GC-MS
182 and NMR (See spectra in supplementary information, SI).

183 2.7 Instrumentation: GC-MS analysis of epoxides and ester **4d** were performed using an Agilent HP
184 6850 gas chromatograph connected to an Agilent HP 5975 quadrupole mass spectrometer. Analytes
185 were separated on a HP-5MS fused-silica capillary column (stationary phase 5%-Phenyl)-
186 methylpolysiloxane, 30 m, 0.25 mm i.d., 0.25 μm film thickness), with helium as the carrier gas (at
187 constant pressure, 36 cm s^{-1} linear velocity at 200 °C). Mass spectra were recorded under electron
188 ionization (70 eV) at a frequency of 1 scan s^{-1} within the 12–600 m/z range. The injection port
189 temperature was 250 °C. The temperature of the column was kept at 50 °C for 5 min, then increased
190 from 50 to 250 °C at 10 °C min^{-1} and the final temperature of 250 °C was kept for 12 min.

191 GC-MS analysis of Baeyer-Villiger products (except **4d**) were performed using an Agilent 7820A
192 gas chromatograph connected to an Agilent 5977E quadrupole mass spectrometer. Analytes were
193 separated on a DB-FFAP polar column (30 m length, 0.25 mm i.d., 0.25 μm film thickness), with
194 helium flow of 1 mL min^{-1} . Mass spectra were recorded under electron ionization (70 eV) at a
195 frequency of 1 scan s^{-1} within the 29–450 m/z range. The injection port temperature was 250 °C.

196 The temperature of the column was kept at 50 °C for 5 min, then increased from 50 to 250 °C at 10
197 °C min^{-1} and the final temperature of 250 °C was kept for 15 min.

198 ^1H NMR spectra were recorded on Varian 400 (400 MHz) spectrometers. ^{13}C NMR spectra were
199 recorded on a Varian 400 (100 MHz) spectrometers. Chemical shifts were reported in ppm from
200 trimethylsilane (TMS) with the solvent resonance as the internal standard (deuteriochloroform: 7.26
201 ppm).

202

203 **3. Results and discussion**

204 3.1 Alkenes epoxidation

205 We studied the chemoenzymatic epoxidation of various alkenes (focusing for each of them on the
206 enzyme amount, peracid precursors and H_2O_2 amount and additions, and reaction time): cyclic
207 alkenes (Table 1), poorly-reactive stilbene (Table 2) and oleic acid (Scheme 2) .

208

209 3.1.1 Cyclic alkenes 1 a-d

210 Studying CAL-B mediated chemoenzymatic epoxidation in various NaDESs, Zhou *et al.* showed
211 that amine-based DESs (i.e. choline chloride-urea, 1:2 molar ratio, called Reline) significantly
212 reduced the stability of CAL-B in a wide temperature range whereas the polyol-based ones
213 increased it. (Zhou et al. 2017) For this reason, we focused our initial experiments on polyol-based
214 NaDESs. Cyclohexene **1a** (Table 1, entry 1) was epoxidized with immobilized CAL-B (100 mg per
215 1.6 mmol of alkene), octanoic acid OA (one eq respect to the alkene), and H₂O₂ (one eq respect to
216 the alkene) in choline chloride-sorbitol (ChCl-Sorb), 1:1 molar ratio (400 mg) (similarly to Zhou *et*
217 *al.*). We observed a complete conversion of the starting material but a very low selectivity towards
218 the epoxide; in fact the most of epoxide was converted into the chlorinated by-product and the diol
219 after 20h. This unexpected result prompted us to turn our attention towards chloride-free, sugar-
220 based NaDESs as the GFS. We tested both solvents (ChCl-Sorb and GFS) in the same conditions
221 with more easily detectable cyclododecene **1b** and we observed that GFS gave better conversions
222 (Table 1, entries 2 and 3); the same held true for other cyclic substrates (cyclooctene **1c** and
223 limonene **1d**) (see in SI Table S1, entries 1 and 5, and Table S2, entry 1). So, we decided to test
224 various substrates to check the viability of the system.
225 When Z/E mixtures were used in the starting alkene (as in **1b**), no diastereoselectivity was observed
226 and the final product diastereomeric ratio reflected the diastereomeric distribution in the reagent.
227 OA resulted the most reactive acid precursor under our conditions (Table 1, entries 9-12),
228 confirming the literature results, and its amount can be significantly lowered from 1 eq to 0.1 eq
229 with all the substrates (Table 1, entries 6, 8, 13, 14, 18). Considering aliphatic acids with different
230 chain lengths, butanoic acid BA (Table 1, entry 10) gave very good results on **1c** while acetic acid
231 AA (Table 1, entry 11) was poorly reactive; the biobased levulinic acid (LA) gave **2c** in good
232 conversion (Table 1, entry 12), prompting us to include 40% of LA as a component of the GFS
233 instead of water, with the aim of using it both as peracid precursor and solvent component; however
234 in this case the epoxidation of cyclododecene **1b** was not satisfactory (Table 1, entry 5). We also
235 tested GFS-LA in combination with OA as peracid precursor on **1b**; results were good but lower
236 than using GFS (Table 1, entries 3 and 4). The same happened with **1c** (see SI, Table S1, entry 4).
237 Dimethyl carbonate DMC was also tested as peracid precursor but without good results (see SI,
238 Table S1, entries 1 and 3). The amount of the enzyme could be lowered till 30-25 U/mmol without
239 significant loss of reactivity (Table 1, **1b** entries 7 and 8, **1c** entry 14, **1d** entries 18 and 19).
240 Limonene **1d** is a very important biobased substrate, whose epoxy derivative is having some
241 relevance in the field of polymer synthesis (Auriemma et al. 2015). Its internal double bond is much
242 more reactive than the terminal one, being electron-richer; in all the tested condition the product **2d**
243 was obtained. Since we initially observed the formation of the diol as by-product (Table 1, entry
244 15), milder conditions were tested: i) halving the amount of NaDES; ii) keeping the temperature
245 below 25 °C; iii) lowering addition rate of H₂O₂. All these conditions allowed to avoid the diol
246 formation (Table 1, entries 17 and 18). The use of a catalytic amount of acid precursor (Table 1,
247 entry 16) and a lower amount of the enzyme defined the best conditions to obtain **2d** in very good
248 conversion (Table 1, entry 18).
249 As expected from limonene results, terminal bonds of styrene and itaconic anhydride were not
250 reactive in the mild condition we tested for the other substrates (see SI, Table S2, entries 5 and 6).

251 252 3.1.2 *trans*-Stilbene **1e**

253 *trans*-Stilbene **1e** is a challenging substrate because its double bond is electron-poor and it is poorly
254 soluble in polar solvents like GFS. OA and other linear aliphatic carboxylic acids with shorter
255 (hexanoic HA, butanoic BA and acetic acid AA) and longer (dodecanoic acid, DA) chain lengths
256 were tested as peracid precursors (Table 2). In all cases OA resulted the most effective acid
257 precursor also in this case but 1 eq was needed to obtain an effective conversion (Table 2, entry 2).
258 A decrease of the enzyme amount was possible, but a conversion of 75% was reached only after 48
259 h (Table 2, entry 4). Longer reaction times did not increase the conversion (SI, Table 2, entries 3
260 and 4). Differently, the electron-poor, α - β double bonds of crotonic acid and methyl crotonate were

261 very difficult to be epoxidized (see SI, Table S2, entries 7 and 8) and we obtained just traces of the
262 products. We also tested substrates carrying hydroxyl groups such as 1-octen-3-ol or *trans*-2-hexen-
263 1-ol but, as expected, the main product was the ester formed by OA and the alcohol under CAL-B
264 catalysis (data not shown).

265

266 3.1.3 Oleic acid **1f**

267 Oleic acid **1f** is a very interesting substrate since its epoxide (9,10-epoxystearic acid) is a highly-
268 valuable oleochemical due to its wide range of industrial applications, including cosmetics, personal
269 care, and pharmaceutical products. The epoxidation worked very well and without the addition of
270 OA (Scheme 2), thanks to an autocatalytic mechanism that formed the peroxy acid from the oleic
271 acid itself. (Rüsch gen. Klaas & Warwel 1999) A temperature of 45 °C was required not only to
272 catalyze the reaction but also to avoid the product solidification. The condition used are the same
273 suggested and used by the recent literature (temperature at maximum 50 °C, an excess of H₂O₂,
274 short reaction time), except for the use of the solvent, which is generally toluene. (Milchert et al.
275 2015) The epoxidation can also be carried out in a solvent-free system, but the process is more
276 efficient for methyl oleate since the corresponding epoxide is liquid respect to solid 9,10-
277 epoxystearic acid. (Orellana-Coca et al. 2005)

278

279 3.2 Baeyer-Villiger oxidations

280 The first use of CAL-B as catalyst for B-V oxidations was performed in toluene with myristic acid
281 as peracid precursor. (Lemoult et al. 1995) Recent examples report ethyl acetate both as solvent and
282 peracid precursor (therefore in large excess with respect to the starting material) (Ríos et al. 2007;
283 Rios et al. 2008; Chávez et al. 2013; Drożdż et al. 2013) and combination of ionic liquids and OA
284 as solvent and peracid precursor, respectively (OA in excess with respect to the starting ketone).
285 (Kotowska et al. 2011; Drożdż et al. 2015) Urea-hydrogen peroxide is considered a milder oxidant
286 than H₂O₂ alone and it was used to reduce the formation of water in the reaction, (Ríos et al. 2007;
287 Rios et al. 2008) nevertheless other studies showed no significant improvement in product
288 conversion and enzyme recycling. (Chávez et al. 2013) Considering that water is already present in
289 our GFS, the availability and the lower cost of hydrogen peroxide, this last one was thus chosen as
290 oxidant in our study. As for reaction times, when the reaction was carried out at room temperature it
291 generally required very long reaction times (in the order of days) to reach effective conversions.
292 (Rios et al. 2007; Rios et al. 2008; Chávez et al. 2013; Drożdż et al. 2013; Drożdż et al. 2015)

293 We tested B-V oxidation on various substrates (see section 2.4), using the same chemoenzymatic
294 method in GFS previously described for epoxidation reactions: CAL-B, H₂O₂ (30% aqueous
295 solution) and OA as peracid precursor (scheme in Table 3). Also in this case, a detailed study was
296 conducted on the reaction conditions, with the aim of reducing the use of the reagents in excess and
297 to use the mildest possible conditions.

298

299 3.2.1 Cyclic ketones **3a-c**

300 Since highly reactive in B-V oxidations, cyclohexanone **3a** was the first substrate tested. By
301 carrying out the reaction at 20 °C, lactone **4a**, (ϵ -caprolactone, Table 3, entry 1) was obtained but
302 the reaction proceeded very slowly and an increase in time lead to the formation of the by-product,
303 6-hydroxyhexanoic acid **4a'**, caused by the ring-opening of **4a**. Increasing the amount of catalyst or
304 temperature did not increase the selectivity towards **4a** formation (Table 3, entries 2 and 3).
305 Differently from the epoxidation reaction of cyclic alkenes (Table 1), the use of the peracid
306 precursor in catalytic amount did not give good results (Table 3, entry 4). Indeed, ω -hydroxy acid
307 formation is the main drawback in CAL-B mediated B-V oxidation (amounts reported in Table 3).
308 (X.-P. Wang et al. 2017) Increasing the amount of H₂O₂ to 2 eq (Table 3, entries 5 and 6) gave
309 higher conversions, without the formation of any by-product, while a shorter reaction time was
310 achieved by conducting the reaction at 45 °C (Table 3, entry 6). The use of both OA and H₂O₂ in
311 excess (Table 3, entries 7 and 8) gave the best conversion: 74% at 20 °C and 58% at 45 °C. As

312 previous observed when the reaction was conducted at 45 °C, the reaction must be stopped after a
313 few hours to avoid by-product formation (Table 3, entry 8). Further increasing of both oxidant and
314 acid amounts was not effective (Table 3, entry 9).

315 The expected higher reactivity of cyclopentanone **3b** prompted us to lower the enzyme amount but
316 also tuning temperature, oxidant and OA amounts (Table 3, entries 11-13) the high reactivity of the
317 substrate caused a rapid formation of the by-product **3b'**. As expected from the literature, (Chávez
318 et al. 2013; Drożdż et al. 2013; Drożdż et al. 2015) substrates with larger rings, as cyclooctanone
319 **3c**, are unreactive in all the tested conditions (Table 3, entry 14).

320

321 3.2.2 2-phenylacetophenone **3d**

322 When using 2-phenylacetophenone **3d** the regioselectivity issue must be considered, due to the
323 formation of two possible regioisomers **4d** and **4d'** (structure in Table 4 foot) caused by the
324 migration of the phenyl group instead of the benzyl one (favored). As expected, we predominantly
325 obtained regioisomer **4d** in 50% conversion at long reaction times (Table 4, entry 1). Higher
326 temperature did not increase the conversion but significantly increased the reaction rate (Table 4,
327 entries 1 and 3). Conversions decreased by lowering the amount of OA and enzyme (Table 4,
328 entries 2 and 4). Using 2 eq of OA and H₂O₂ was not effective (Table 4, entry 5), while a great
329 excess of H₂O₂ gave 60% of **4d** (Table 4, entry 6). Linear ketones and levulinic acid were tested but
330 the reactin did not work under the developed conditions (data not shown).

331

332 Conclusions

333 We demonstrated that chemoenzymatic oxidations using lipase CAL-B to form the active oxidant
334 from carboxylic acid/H₂O₂ pair can be performed in a sugar-based NaDES composed by an
335 equimolar mixture of glucose, fructose, sucrose and water (GFS). Specific conditions to perform the
336 reaction on selected substrates in good conversion and selectivity were found. The best conditions
337 for epoxidations proved to be related to the substrate reactivity; reaction conditions were tuned and
338 catalysts amounts decreased to obtain epoxides from poorly reactive and steric-hindered double
339 bonds (as *trans*-stilbene) and to control the formation of byproducts in more reactive alkenes (like
340 internal double bond of R-limonene). Baeyer-Villiger oxidations always required at least
341 stoichiometric amount of the peracid precursor to proceed and an excess of both oxidant and acid to
342 obtain good conversions.

343

344 Acknowledgements

345 We thank the EMM ChIR (Chemical innovation and regulation) for M.K. Vasquez De Paz and D.G.
346 Gidey thesis fellowship, the University of Bologna (RFO program) and MIUR (PhD program) for
347 funding.

348

349

350 Disclosure of interest

351 The authors report no conflict of interest.

352

353

354 References

355 Abbott AP, Capper G, Davies DL, Munro HL, Rasheed RK, Tambyrajah V. 2001. Preparation of
356 novel, moisture-stable, Lewis-acidic ionic liquids containing quaternary ammonium salts with
357 functional side chains. *Chem Commun.*(19):2010–2011.

358

359 Alphand V, Carrea G, Wohlgemuth R, Furstoss R, Woodley JM. 2003. Towards large-scale
360 synthetic applications of Baeyer-Villiger monooxygenases. *Trends in Biotechnology.* 21(7):318–
361 323.

362

363 Aouf C, Durand E, Lecomte J, Figueroa-Espinoza M-C, Dubreucq E, Fulcrand H, Villeneuve P.
364 2014. The use of lipases as biocatalysts for the epoxidation of fatty acids and phenolic compounds.
365 *Green Chem.* 16(4):1740–1754.
366

367 Auriemma F, De Rosa C, Di Caprio MR, Di Girolamo R, Ellis WC, Coates GW. 2015.
368 Stereocomplexed Poly(Limonene Carbonate): A Unique Example of the Cocrystallization of
369 Amorphous Enantiomeric Polymers. *Angewandte Chemie.* 127(4):1231–1234.
370

371 Balke K, Kadow M, Mallin H, Saß S, T. Bornscheuer U. 2012. Discovery, application and protein
372 engineering of Baeyer–Villiger monooxygenases for organic synthesis. *Organic & Biomolecular*
373 *Chemistry.* 10(31):6249–6265.
374

375 Björkling F, Godtfredsen SE, Kirk O. 1990. Lipase-mediated formation of peroxycarboxylic acids
376 used in catalytic epoxidation of alkenes. *J Chem Soc, Chem Commun.*(19):1301–1303.
377

378 ten Brink G-J, Arends IWCE, Sheldon RA. 2004. The Baeyer–Villiger Reaction: New
379 Developments toward Greener Procedures. *Chem Rev.* 104(9):4105–4124.
380

381 Busto E, Gotor-Fernández V, Gotor V. 2010. Hydrolases: catalytically promiscuous enzymes for
382 non-conventional reactions in organic synthesis. *Chemical Society Reviews.* 39(11):4504–4523.
383

384 Cavani F, Teles JH. 2009. Sustainability in Catalytic Oxidation: An Alternative Approach or a
385 Structural Evolution? *ChemSusChem.* 2(6):508–534.
386

387 Chávez G, Hatti-Kaul R, Sheldon RA, Mamo G. 2013. Baeyer–Villiger oxidation with peracid
388 generated in situ by CaLB-CLEA catalyzed perhydrolysis. *Journal of Molecular Catalysis B:*
389 *Enzymatic.* 89:67–72.
390

391 Constable DJC, Dunn PJ, Hayler JD, Humphrey GR, Johnnie L, Leazer J, Linderman RJ, Lorenz K,
392 Manley J, Pearlman BA, Wells A, et al. 2007. Key green chemistry research areas—a perspective
393 from pharmaceutical manufacturers. *Green Chem.* 9(5):411–420.
394

395 Dai Y, van Spronsen J, Witkamp G-J, Verpoorte R, Choi YH. 2013. Natural deep eutectic solvents
396 as new potential media for green technology. *Analytica Chimica Acta.* 766:61–68.
397

398 Drożdż A, Chrobok A, Baj S, Szymańska K, Mrowiec-Białoń J, Jarzębski AB. 2013. The chemo-
399 enzymatic Baeyer–Villiger oxidation of cyclic ketones with an efficient silica-supported lipase as a
400 biocatalyst. *Applied Catalysis A: General.* 467:163–170.
401

402 Drożdż A, Erfurt K, Bielas R, Chrobok A. 2015. Chemo-enzymatic Baeyer–Villiger oxidation in
403 the presence of *Candida antarctica* lipase B and ionic liquids. *New J Chem.* 39(2):1315–1321.
404

405 Durand E, Lecomte J, Baréa B, Piombo G, Dubreucq E, Villeneuve P. 2012. Evaluation of deep
406 eutectic solvents as new media for *Candida antarctica* B lipase catalyzed reactions. *Process*
407 *Biochemistry.* 47(12):2081–2089.
408

409 Elgharbawy AAM, Moniruzzaman M, Goto M. 2020. Recent advances of enzymatic reactions in
410 ionic liquids: Part II. *Biochemical Engineering Journal.* 154:107426.
411

412 Galletti P, Moretti F, Samorì C, Tagliavini E. 2007. Enzymatic acylation of levoglucosan in
413 acetonitrile and ionic liquids. *Green Chem.* 9(9):987–991.

414
415 García C, Hoyos P, Hernáiz MJ. 2018. Enzymatic synthesis of carbohydrates and glycoconjugates
416 using lipases and glycosidases in green solvents. *Biocatalysis and Biotransformation*. 36(2):131–
417 140.
418
419 Gorke JT, Srienc F, Kazlauskas RJ. 2008. Hydrolase-catalyzed biotransformations in deep eutectic
420 solvents. *Chemical Communications*.(10):1235–1237.
421
422 Gotor-Fernández V, Paul CE. 2019. Deep eutectic solvents for redox biocatalysis. *Journal of*
423 *Biotechnology*. 293:24–35.
424
425 Guajardo N, Ahumada K, Domínguez de María P, Schrebler RA. 2019. Remarkable stability of
426 *Candida antarctica* lipase B immobilized *via* cross-linking aggregates (CLEA) in deep eutectic
427 solvents. *Biocatalysis and Biotransformation*. 37(2):106–114.
428
429 Hilker I, Bothe D, Prüss J, Warnecke H-J. 2001. Chemo-enzymatic epoxidation of unsaturated plant
430 oils. *Chemical Engineering Science*. 56(2):427–432.
431
432 Hobbs HR, Thomas NR. 2007. Biocatalysis in supercritical fluids, in fluoruous solvents, and under
433 solvent-free conditions. *Chem Rev*. 107(6):2786–2820.
434
435 Hollmann F, Arends IWCE, Buehler K, Schallmeyer A, Bühler B. 2011. Enzyme-mediated
436 oxidations for the chemist. *Green Chem*. 13(2):226–265.
437
438 Kotlewska AJ, van Rantwijk F, Sheldon RA, Arends IWCE. 2011. Epoxidation and Baeyer–
439 Villiger oxidation using hydrogen peroxide and a lipase dissolved in ionic liquids. *Green Chem*.
440 13(8):2154.
441
442 Lei Z, Chen B, Koo Y-M, Macfarlane DR. 2017. Introduction: Ionic Liquids. *Chemical Reviews*.
443 117(10):6633–6635.
444
445 Leisch H, Morley K, Lau PCK. 2011. Baeyer–Villiger Monooxygenases: More Than Just Green
446 Chemistry. *Chem Rev*. 111(7):4165–4222.
447
448 Lemoult SC, Richardson PF, Roberts SM. 1995. Lipase-catalysed Baeyer–Villiger reactions. *J*
449 *Chem Soc, Perkin Trans 1*.(2):89–91.
450
451 Ma Q, Xing W, Xu J, Peng X. 2014. Baeyer–Villiger oxidation of cyclic ketones with aqueous
452 hydrogen peroxide catalyzed by transition metal oxides. *Catalysis Communications*. 53:5–8.
453
454 Ma Y, Li P, Li Y, Willot SJ-P, Zhang W, Ribitsch D, Choi YH, Verpoorte R, Zhang T, Hollmann F,
455 Wang Y. 2019. Natural Deep Eutectic Solvents as Multifunctional Media for the Valorization of
456 Agricultural Wastes. *ChemSusChem*. 12(7):1310–1315.
457
458 Milchert E, Malarczyk K, Kłos M. 2015. Technological Aspects of Chemoenzymatic Epoxidation
459 of Fatty Acids, Fatty Acid Esters and Vegetable Oils: A Review. *Molecules*. 20(12):21481–21493.
460
461 Moniruzzaman M, Kamiya N, Goto M. 2010. Activation and stabilization of enzymes in ionic
462 liquids. *Organic & Biomolecular Chemistry*. 8(13):2887–2899.
463
464 Moreira MA, Nascimento MG. 2007. Chemo-enzymatic epoxidation of (+)-3-carene. *Catalysis*
Communications. 8(12):2043–2047.

465
466 Nian B, Cao C, Liu Y. 2020. How *Candida antarctica* lipase B can be activated in natural deep
467 eutectic solvents: experimental and molecular dynamics studies. *Journal of Chemical Technology &*
468 *Biotechnology*. 95(1):86–93.
469
470 Niku-Paavola M-L, Viikari L. 2000. Enzymatic oxidation of alkenes. *Journal of Molecular*
471 *Catalysis B: Enzymatic*. 10(4):435–444.
472
473 Nkuku CA, LeSuer RJ. 2007. Electrochemistry in Deep Eutectic Solvents. *J Phys Chem B*.
474 111(46):13271–13277.
475
476 O. Burek B, Bormann S, Hollmann F, Z. Bloh J, Holtmann D. 2019. Hydrogen peroxide driven
477 biocatalysis. *Green Chemistry*. 21(12):3232–3249.
478
479 Oh Y, Park S, Yoo E, Jo S, Hong J, Kim HJ, Kim KJ, Oh KK, Lee SH. 2019. Dihydrogen-bonding
480 deep eutectic solvents as reaction media for lipase-catalyzed transesterification. *Biochemical*
481 *Engineering Journal*. 142:34–40.
482
483 Orellana-Coca C, Törnvall U, Adlercreutz D, Mattiasson B, Hatti-Kaul R. 2005. Chemo-enzymatic
484 epoxidation of oleic acid and methyl oleate in solvent-free medium. *Biocatalysis and*
485 *Biotransformation*. 23(6):431–437.
486
487 Ortiz C, Luján Ferreira M, Barbosa O, Santos JCS dos, C. Rodrigues R, Berenguer-Murcia Á,
488 E. Briand L, Fernandez-Lafuente R. 2019. Novozym 435: the “perfect” lipase immobilized
489 biocatalyst? *Catalysis Science & Technology*. 9(10):2380–2420.
490
491 Paiva A, Craveiro R, Aroso I, Martins M, Reis RL, Duarte ARC. 2014. Natural Deep Eutectic
492 Solvents – Solvents for the 21st Century. *ACS Sustainable Chem Eng*. 2(5):1063–1071.
493
494 Qin Y-Z, Li Y-M, Zong M-H, Wu H, Li N. 2015. Enzyme-catalyzed selective oxidation of 5-
495 hydroxymethylfurfural (HMF) and separation of HMF and 2,5-diformylfuran using deep eutectic
496 solvents. *Green Chemistry*. 17(7):3718–3722.
497
498 Ranganathan S, Zeitlhofer S, Sieber V. 2017. Development of a lipase-mediated epoxidation
499 process for monoterpenes in choline chloride-based deep eutectic solvents. *Green Chem*.
500 19(11):2576–2586.
501
502 Renz M, Meunier B. 1999. 100 Years of Baeyer–Villiger Oxidations. *European Journal of Organic*
503 *Chemistry*. 1999(4):737–750.
504
505 Ríos MY, Salazar E, Olivo HF. 2007. Baeyer–Villiger oxidation of substituted cyclohexanones via
506 lipase-mediated perhydrolysis utilizing urea–hydrogen peroxide in ethyl acetate. *Green Chem*.
507 9(5):459–462.
508
509 Rios MY, Salazar E, Olivo HF. 2008. Chemo-enzymatic Baeyer–Villiger oxidation of
510 cyclopentanone and substituted cyclopentanones. *Journal of Molecular Catalysis B: Enzymatic*.
511 54(3–4):61–66.
512
513 Rüschen gen. Klaas M, Warwel S. 1999. Complete and partial epoxidation of plant oils by lipase-
514 catalyzed perhydrolysis | Based partly on a lecture at the International Conference of the Association
515 for the Advancement of Industrial Crops, Saltillo, Mexico, September 14–18, 1997. *Industrial*

516 Crops and Products. 9(2):125–132.
517
518 Samorì C, Campisi T, Fagnoni M, Galletti P, Pasteris A, Pezzolesi L, Protti S, Ravelli D, Tagliavini
519 E. 2015. Pyrrolidinium-based Ionic Liquids: Aquatic Ecotoxicity, Biodegradability, and Algal
520 Subinhibitory Stimulation. ACS Sustainable Chem Eng. 3(8):1860–1865.
521
522 Samorì C, Mazzei L, Ciurli S, Cravotto G, Grillo G, Guidi E, Pasteris A, Tabasso S, Galletti P.
523 2019. Urease Inhibitory Potential and Soil Ecotoxicity of Novel “Polyphenols–Deep Eutectic
524 Solvents” Formulations. ACS Sustainable Chem Eng. 7(18):15558–15567.
525
526 Silva WSD, Lapis AAM, Suarez PAZ, Neto BAD. 2011. Enzyme-mediated epoxidation of methyl
527 oleate supported by imidazolium-based ionic liquids. Journal of Molecular Catalysis B: Enzymatic.
528 68(1):98–103.
529
530 Smith EL, Abbott AP, Ryder KS. 2014. Deep Eutectic Solvents (DESs) and Their Applications.
531 Chem Rev. 114(21):11060–11082.
532
533 Strukul G. 1998. Transition Metal Catalysis in the Baeyer–Villiger Oxidation of Ketones.
534 Angewandte Chemie International Edition. 37(9):1198–1209.
535
536 Swern Daniel. 1949. Organic Peracids. Chem Rev. 45(1):1–68.
537
538 Tommasi E, Cravotto G, Galletti P, Grillo G, Mazzotti M, Sacchetti G, Samorì C, Tabasso S,
539 Tacchini M, Tagliavini E. 2017. Enhanced and Selective Lipid Extraction from the Microalga *P.*
540 *tricornutum* by Dimethyl Carbonate and Supercritical CO₂ Using Deep Eutectic Solvents and
541 Microwaves as Pretreatment. ACS Sustainable Chem Eng. 5(9):8316–8322.
542
543 Ülger C, Takaç S. 2017. Kinetics of lipase-catalysed methyl gallate production in the presence of
544 deep eutectic solvent. Biocatalysis and Biotransformation. 35(6):407–416.
545
546 Wang B, Qin L, Mu T, Xue Z, Gao G. 2017. Are Ionic Liquids Chemically Stable? Chem Rev.
547 117(10):7113–7131.
548
549 Wang X-P, Zhou P-F, Li Z-G, Yang B, Hollmann F, Wang Y-H. 2017. Engineering a lipase B from
550 *Candida antartica* with efficient perhydrolysis performance by eliminating its hydrolase activity. Sci
551 Rep. 7(1):44599.
552
553 Woodruff M, Hutmacher D. 2010. The return of a forgotten polymer - polycaprolactone in the 21st
554 century. Progress in Polymer Science. 35(10):1217–1256.
555
556 Yadav GD, Devi KM. 2002. Enzymatic synthesis of perlauric acid using Novozym 435.
557 Biochemical Engineering Journal. 10(2):93–101.
558
559 Yang S-L, Duan Z-Q. 2016. Insight into enzymatic synthesis of phosphatidylserine in deep eutectic
560 solvents. Catalysis Communications. 82:16–19.
561
562 Yin J, Wang J, Li Z, Li D, Yang G, Cui Y, Wang A, Li C. 2015. Deep desulfurization of fuels
563 based on an oxidation/extraction process with acidic deep eutectic solvents. Green Chemistry.
564 17(9):4552–4559.
565
566 Zhang Q, Vigier KDO, Royer S, Jérôme F. 2012. Deep eutectic solvents: syntheses, properties and

567 applications. *Chemical Society Reviews*. 41(21):7108–7146.
568
569 Zhou P, Wang X, Yang B, Hollmann F, Wang Y. 2017. Chemoenzymatic epoxidation of alkenes
570 with *Candida antarctica* lipase B and hydrogen peroxide in deep eutectic solvents. *RSC Advances*.
571 7(21):12518–12523.
572

Tables

Table 1. Epoxidation of cyclic alkenes with chemoenzymatic method in NaDES.


Entry	Alkene	NaDES	CAL-B (U/mmol)	Peracid precursor (eq)	Product conversion (%) ^a	By-products conversion (%) ^a
1	1a	ChCl- Sorb [1:1]	250	OA, 1	6	34 59
						-
2	1b	ChCl- Sorb [1:1]	250	OA, 1	68	
3	1b	GFS	250	OA, 1	71	
4	1b	GFS [1:1:1] -LA	250	OA, 1	64	
5	1b	GFS [1:1:1] -LA	250	LA	15	
6	1b	GFS	250	OA, 0.1	75	
7	1b	GFS	25	OA, 1	80	
8	1b	GFS	25	OA, 0.1	79 (77)	
						-
9	1c	GFS	250	OA, 1	>99	
10	1c	GFS	250	BA, 1	99	
11	1c	GFS	250	AA, 1	48	
12	1c	GFS	250	LA, 1	87	
13	1c	GFS	250	OA, 0.1	93	
14	1c	GFS	25	OA, 0.1	95 (91)	

	1d				2d	2d'
15 ^c	1d	GFS	60	OA, 1	76	14
16 ^c	1d	GFS	60	OA, 0.1	73	16
17 ^{c,d}	1d	GFS	60	OA, 1	89	-
18 ^{c,d}	1d	GFS	30	OA, 0.1	96	-
19 ^c	1d	GFS	30	OA, 0.1	53	31

^a conversion by GC-MS, isolated yield in brackets; ^b diastereomeric ratio Z/E in **1b** and **2b** is always 2:1; ^c Room temperature (20 °C); ^d H₂O₂ total amount divided into 4 portions added in 4 hours.

Acronyms: ChCl= choline chloride, Sorb = sorbitol, GFS = glucose, fructose, sucrose, H₂O (1:1:1:11), OA= octanoic acid, LA= levulinic acid, AA = acetic acid, BA=butyric acid.









Table 2. Epoxidation of *trans*- stilbene **1e** with chemoenzymatic method in GFS NaDES.

			
Entry	CAL-B (U/mmol)	Peracid precursor (eq)	2e conversion (%) ^a
1 ^b	250	OA, 1	60
2 ^b	250	OA, 0.1	traces
3	250	OA, 1	74 (70)
4	25	OA, 1	73
5	25	DA, 1	11
6	25	HA, 1	54
7	25	BA, 1	-
8	25	AA, 1	-

^a conversion by GC-MS, isolated yield in brackets; ^b time (20h)


Acronyms: GFS = glucose, fructose, sucrose, water (1:1:1:11), OA= octanoic acid, DA = dodecanoic acid, HA= Hexanoic Acid, BA=butyric acid, AA = acetic acid.

Table 3. Baeyer-Villiger oxidation of lactones with chemoenzymatic method in GFS.


$ \begin{array}{ccc} \text{R}_1-\text{C}(=\text{O})-\text{R}_2 & \xrightarrow[\text{GFS}]{\text{CAL-B, Octanoic Acid, H}_2\text{O}_2} & \text{R}_1-\text{C}(=\text{O})-\text{O}-\text{R}_2 \\ \text{3a-c} & & \text{4a-c} \\ \text{T}^\circ\text{C} & & \text{time} \end{array} $								
Entry	Ketone	CAL-B (U/mmol)	OA (eq)	H ₂ O ₂ (eq)	T (°C)	time (h)	Product conversion (%) ^a	By-product conversion (%) ^a
	 3a						 4a	 4a'
1	3a	125	1	1	20	20 70	32 23	- 53
2	3a	250	1	1	20	20	21	16
3	3a	200	1	1	45	5 15	35 36	- 28
4	3a	125	0,1	1	20	20 40	15 15	- -
5	3a	125	1	2	20	20 40	37 55	- -
6	3a	125	1	2	45	5 20	30 20	- 8
7	3a	125	2	2	20	20 40 64	49 61 74	- - -
8	3a	125	2	2	45	5 20	52 29	- 21
9	3a	125	1	3	20	20 4d	36 54	- 16
10	3a	125	3	3	20	20 4d	58 54	- 16
	 3b						 4b	 4b'
11	3b	65	1	1	20	20 40	15 7	- 50
12	3b	65	1	1	45	20	10	30
13	3b	65	0.1	1	20	40	4	15
	 3c						 4c	Not found
14	3c	65	various	various	20	various	traces	-

^a conversion by GC-MS, isolated yield in brackets;
 Acronyms: GFS = glucose, fructose, sucrose, H₂O (1:1:1:11), OA= octanoic acid.

Table 4. Baeyer-Villiger oxidation of 2-phenylacetophenone **3d** in GFS

						
Entry	Ketone	CAL-B (U/mmol)	OA (eq)	H ₂ O ₂ (eq)	time (h)	Conversion 4d (%) ^{a,c}
1 ^b	3d	250	1	1	40 50 7 days	7 17 50
2 ^b	3d	250	0.1	1	50	12
3	3d	250	1	1	40 7 days	40 (34) 50
4	3d	100	1	1	20 5 days	18 27
5	3d	250	2	2	20 50 3 days	38 42 44
6	3d	250	1	3	23 4 days	55 60


^a conversion by GC-MS, isolated yield in brackets; ^b temperature (20 °C); ^c in all the entries there are traces of the regioisomer of




Acronyms: GFS = glucose, fructose, sucrose, H₂O (1:1:1:11), OA= octanoic acid.

Schemes

Scheme 1



Scheme 2



Schemes Captions

Scheme 1. Chemoenzymatic pathway for epoxidations and Baeyer-Villiger oxidations.

Scheme 2. Epoxidation of oleic acid with chemoenzymatic method in GFS.