

NOTE

Disseminated phaeohyphomycosis in a beluga sturgeon (*Huso huso*)

L. Mandrioli¹, E. Soto², Z. Yazdi², E. Volpe¹, F. Errani¹, S. Ciulli¹, A. Camus³

¹ Department of Veterinary Medical Sciences (DIMEVET), Alma Mater Studiorum University of Bologna, Italy, ² Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California, USA (UC Davis), ³ Department of Pathology, College of Veterinary Medicine, University of Georgia, USA (UGA)

Keywords: beluga sturgeon, dematiaceous fungi, *Huso huso*, phaeohyphomycosis

<https://doi.org/10.48045/001c.37808>

Bulletin of the European Association of Fish Pathologists

Sturgeon aquaculture is an emerging worldwide industry both for human consumption as well as for conservation. Several transmissible diseases limit this industry including mycosis by dematiaceous fungi such as *Veronaea botryosa*. This study reports the first diagnosis of phaeohyphomycosis in a beluga sturgeon (*Huso huso*) in Italy.

Phaeohyphomycosis is a fungal infection caused by dematiaceous or melanised fungi (Seyedmousavi, Guillot, and de Hoog 2013). Although considered ubiquitous residents of plant material, soil, and wood, melanised fungi are likely adapted to specific niches that facilitate variable opportunistic or true pathogenic potentials. Exposure is typically associated with inoculation by minor trauma or inhalation. In mammals, infections are commonly cutaneous, subcutaneous, upper respiratory or primary cerebral, but in cold-blooded vertebrates are often disseminated and accompanied by severe tissue necrosis (Revankar, Sutton, and Rinaldi 2004; Seyedmousavi, Guillot, and de Hoog 2013).

In fish, various melanised fungi within the family *Herpotrichiellaceae* have been reported (Seyedmousavi et al. 2018). Fungi of the genus *Cladophialophora* have been only occasionally reported (Soto et al. 2017) and are considered commensals of gastrointestinal tract (Ghorbani-Choboghlo et al. 2014). In contrast, the genera *Exophiala* and *Veronaea* contain a number of well-recognized fish pathogens. Several *Exophiala* species have been associated with phaeohyphomycosis in various fish species (Seyedmousavi et al. 2018), but have never been reported in sturgeons. Conversely, *Veronaea botryosa* has been reported as the cause of systemic phaeohyphomycosis in two sturgeon species in the USA, the Siberian sturgeon (*Acipenser baerii*) and white sturgeon (*A. transmontanus*) (Steckler et al. 2014).

A juvenile beluga sturgeon (*Huso huso*), one of 11 mortalities from a group of 46 experiencing chronic losses while being grown for restocking purposes in a recirculating culture system, is the subject of this note. Diagnostic investigation of the outbreak identified varied microbial agents, including multiple bacterial species and a putative herpesvirus, as well as inconsistent pathologic findings suggestive of nutritional myopathy and this single case of phaeohyphomycosis. The mortalities were ultimately attributed to environmental stress and nutritional imbalances that likely predisposed the fish to a compromised

immune system and opportunistic infections (Ciulli et al. 2020). The influence of temperature in this case is unclear, as the fish were being grown at 19°C near the high end of their 10-20°C optimal range (Froese and Pauly 2021).

The fish was found dead in its tank and immediately transferred under refrigeration conditions to the diagnostic laboratory at the Local Operative Unit of Cesenatico (DIMEVET, Alma Mater Studiorum University of Bologna, Italy) for necropsy. Gross findings included arching of the body, multifocal reddening and erosion of ventral skin ([Figure 1a](#)), pale grey gills, sero-haemorrhagic coelomic effusion, a soft dark-brown liver, and a tan gastrointestinal tract devoid of ingesta ([Figure 1b](#)). Cytologic evaluation of coelomic fluid stained with May-Grünwald-Giemsa revealed pigmented, brown hyphae and bacteria later identified as *Aeromonas veronii* and *Citrobacter freundii* (Ciulli et al. 2020). In brief, a complete set of tissues (integument, head, gills, heart, liver, spleen, kidney, stomach, intestine) was collected, fixed in 10% neutral buffered formalin, processed routinely, and stained with haematoxylin & eosin (H&E) for histologic evaluation. Select sections were stained by the periodic acid-Schiff (PAS), Gomori methenamine silver (GMS), and Fontana-Masson (FM) methods. As the samples were collected some hours post-mortem, some autolytic findings, particularly at the level of intestinal mucosa, and some post mortem changes had occurred. However, after preservation tissues were deemed in a sufficiently good state for examination and interpretation. Pigmented, bronze hyphae, predominantly 3-5 µm (up to 10 µm) in diameter, with thin, non-parallel walls, infrequent septation, septal constrictions, and acute angle branching were widely distributed in all tissues examined. Hyphae were present transmurally in intestinal sections ([Figure 2a](#)) and were particularly abundant in the coelom adjacent to the intestinal serosa. Angioinvasion was common, with hyphae present in vascular lumens, including those of the gills and heart ([Figure 2b](#)). PAS and GMS stains highlighted the massive presence and morphologic features of hyphae in tissues ([Figure 2c, d](#)). The FM stain confirmed the presence of melanin pigment within the fungal walls ([Figure 2e](#)). Additional histological findings included severe lymphoid depletion of epicardial haematopoietic tissue, epaxial muscle pyrenophores degeneration and atrophy and multifocal ulcerative dermatitis.

In an attempt to identify the fungal agent, hyphae were excised from formalin fixed tissue sections using an MMI CellCut Plus laser capture microdissection (LCM) system (Molecular Machines & Industries, Glattbrugg, Switzerland) at the Department of Pathology, University of Georgia (Athens, GA, USA). Total DNA was extracted using a DNeasy Blood and Tissue Kit (Qiagen) after pre-treatment for formalin-fixed paraffin-embedded (FFPE) tissue following the manufacturer's instructions. Whole genome amplification was then performed using a REPLI-g mini kit (Qiagen) to improve the quantity and quality of DNA. Endpoint PCR was attempted on both FFPE and REPLI-g amplified DNA using universal primers for amplification of fungal ribosomal RNA




Figure 1. Beluga sturgeon (*Huso huso*) juvenile.

(a) Multifocal reddening and erosion of ventral skin, (b) sero-hemorrhagic coelomic effusion, a soft dark-brown liver, and a tan gastrointestinal tract devoid of ingesta.




Figure 2. Beluga sturgeon (*Huso huso*) juvenile.

(a) Hyphae were present transmurally in intestinal sections, (b) Pigmented, bronze hyphae, predominantly 3-5 μm (up to 10 μm) in diameter, with thin, non-parallel walls, infrequent septation, septal constrictions, and acute angle branching were widely distributed in the tissues and present within the heart. (c) PAS and (d) GMS stains highlighted the massive presence and morphologic features of hyphae in tissues. (e) The FM stain confirmed the presence of melanin pigment within the fungal walls.

genes (White et al. 1990; Kurtzman and Robnett 1997) and a quantitative PCR (qPCR) performed following established protocols with *V. botryosa* specific primers (Yazdi et al. 2021) at the Department of Medicine and Epidemiology, University of California (Davis, CA, USA). However, corresponding products failed to amplify, likely the result of DNA degradation associated with formalin fixation. In a further attempt to identify the fungus, frozen samples including a DNA sample (20-25 μ L) and supernatant from a tissue lysate were sent to the University of California. However, due to shipping delays, the samples arrived at ambient temperature after 10 days in transit. The tissue lysate was centrifuged to produce a small pellet, the DNA extracted (Yazdi et al. 2021) then amplified and subjected to PCR and qPCR as described above. The presence of qPCR inhibitors was assessed by including an internal positive control for each sample. Again, no *V. botryosa* specific or other fungal products were detected. Control samples produced positive results as expected.

Despite the inability to identify the fungus by molecular methods, gross and microscopic findings in this juvenile beluga sturgeon are compatible with reports of *V. botryosa*-induced phaeohyphomycosis in other sturgeon species, including angioinvasion and widespread dissemination of pigmented hyphae (Steckler et al. 2014; Coleman et al. 2018; Groff et al. 2020). Significantly, the presence of coelomic fluid and involvement of serosal surfaces is typical of the condition known to producers of white sturgeon in the United States as “fluid belly” (Coleman et al. 2018).

Systemic mycosis by *V. botryosa* was the primary diagnostic consideration in this investigation due to association of the fungus with cultured white and Siberian sturgeon (Steckler et al. 2014; Coleman et al. 2018; Groff et al. 2020). However, additional novel and known pigmented fungi are increasingly identified as pathogens of bony and cartilaginous fish. Commonly implicated genera include *Exophiala*, *Ochrochonis*, and *Phoma* (Armwood et al. 2021). Significantly, fungal identification based on morphologic features, particularly in histologic sections, is unreliable and molecular methods should be employed when fresh or frozen tissue is available. Targets for identification based on PCR amplification and sequencing include the internal-transcribed spacer region of the rDNA (ITS), large ribosomal subunit gene D1/D2 domains (LSU), and the DNA polymerase II gene (PBP2) gene. While certain PCR protocols can work on FFPE tissues (Santos et al. 2013; Steckler et al. 2014; Meason-Smith et al. 2017; Cañete-Gibas and Wiederhold 2018), molecular characterisation attempts failed to identify the agent, presumably the result of DNA degradation due to formalin fixation and delayed sample processing caused by covid-19 associated shipping interruptions of frozen sample material.

Apart from the identification of the dematiaceous fungus, it is evident that the observed infection was predisposed by the poor health condition of the subject suggested by the bacterial coinfection findings, the immune cell depletion and

the epaxial muscle pyrenophores degeneration. These detrimental effects are presumably due to nutritional imbalances and environmental stress induced by poor farming conditions. In these circumstances high pathogenic fungi such as *Exophiala* and *Veronaea* and less pathogenic ones such as *Cladophialophora* could take over the fish defence mechanisms.

Overfishing and poaching have greatly reduced wild populations of beluga sturgeon, prompting the IUCN to list the species as critically endangered (Froese and Pauly 2021). Declining numbers of wild sturgeon highlight the need for repopulation efforts based on artificial culture conditions. Despite the inability to identify the pathogen in this case, fish pathologists involved with sturgeon aquaculture should be aware of emerging fungal pathogens, particularly *V. botryosa*, as a potential threat to economic productivity and the conservation of beluga sturgeon, as well as culture conditions that may promote infection. Particularly, the setup of specific nutritional and environmental conditions are required for these ancient species even if this is actually limited by the paucity of information available (Hung 2017).

Concluding, fungal culture should be considered when sampling diseased sturgeon and fresh or frozen tissue is collected for use in molecular testing, to achieve an aetiological diagnosis and contribute significantly to the concurrent histopathological findings.

Submitted: December 15, 2021 BST, Accepted: June 01, 2022 BST



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

REFERENCES

- Armwood, Abigail R., Connie F. Cañete-Gibas, Jennifer A. Dill-Okubo, Nathan P. Wiederhold, and Alvin C. Camus. 2021. "Retrospective Study of Phaeohyphomycosis in Aquarium-Housed Fish, with First Descriptions of *Exophiala Lecanii-Corni* and *Neodevriesia Cladophorae* in Fish." *Journal of Fish Diseases* 44 (10): 1563–77. <https://doi.org/10.1111/jfd.13477>.
- Cañete-Gibas, Connie F., and Nathan P. Wiederhold. 2018. "The Black Yeasts: An Update on Species Identification and Diagnosis." *Current Fungal Infection Reports* 12 (2): 59–65. <https://doi.org/10.1007/s12281-018-0314-0>.
- Ciulli, S., E. Volpe, G. Tura, E. Zavatta, A. Renzi, T. Pretto, A. Toffan, et al. 2020. "Multiple Co-Infections and Environmental Stressors as Causes of Chronic Mortalities in Juvenile Sturgeons." *Bulletin of the European Association of Fish Pathologists* 40: 13–15.
- Coleman, Denver J., Alvin C. Camus, Beatriz Martínez-López, Susan Yun, Brittany Stevens, and Esteban Soto. 2018. "Effects of Temperature on *Veronaea Botryosa* Infections in White Sturgeon *Acipenser Transmontanus* and Fungal Induced Cytotoxicity of Fish Cell Lines." *Veterinary Research* 49 (1): 11. <https://doi.org/10.1186/s13567-018-0507-0>.
- Froese, R., and D Pauly, eds. 2021. "Fishbase." *World Wide Web Electronic Publication*.
- Ghorbani-Choboghlo, H., A.R. Khosravi, A. Sharifzadeh, M. Taghavi, S. Darvishi, Ashrafi Tamami I, and A. Erfanmanesh. 2014. "Gastrointestinal Microflora of Captured Stellate Sturgeon (*Acipenser Stellatus*, Pallas, 1771) from Southeast Caspian Sea, Iran." *Iranian Journal of Fisheries Sciences* 13: 319–29.
- Groff, Joseph M., Mai Y. Mok, Steven V. Kubiski, Adam O. Michel, Galaxia A. Cortés-Hinojosa, Barbara A. Byrne, Brian L. Wickes, E. III Scott Weber, Lori A Campbell, and Thomas B. Waltzek. 2020. "Phaeohyphomycosis Due to *Veronaea Botryosa* in Cultured White Sturgeon (*Acipenser Transmontanus* Richardson) from California USA during 2006 to 2015." *Journal of Fish Diseases* 44 (6): 793–801. <https://doi.org/10.1111/jfd.13308>.
- Hung, Silas S.O. 2017. "Recent Advances in Sturgeon Nutrition." *Animal Nutrition* 3 (3): 191–204. <https://doi.org/10.1016/j.aninu.2017.05.005>.
- Kurtzman, C P, and C J Robnett. 1997. "Identification of Clinically Important Ascomycetous Yeasts Based on Nucleotide Divergence in the 5' End of the Large-Subunit (26S) Ribosomal DNA Gene." *Journal of Clinical Microbiology* 35 (5): 1216–23. <https://doi.org/10.1128/jcm.35.5.1216-1223.1997>.
- Meason-Smith, Courtney, Erin E. Edwards, Caitlin E. Older, Mackenzie Branco, Laura K. Bryan, Sara D. Lawhon, Jan S. Suchodolski, Gabriel Gomez, Joanne Mansell, and Aline Rodrigues Hoffmann. 2017. "Panfungal Polymerase Chain Reaction for Identification of Fungal Pathogens in Formalin-Fixed Animal Tissues." *Veterinary Pathology* 54 (4): 640–48. <https://doi.org/10.1177/0300985817698207>.
- Revankar, Sanjay G., Deanna A. Sutton, and Michael G. Rinaldi. 2004. "Primary Central Nervous System Phaeohyphomycosis: A Review of 101 Cases." *Clinical Infectious Diseases* 38 (2): 206–16. <https://doi.org/10.1086/380635>.
- Santos, Daniel W. C. L., Ana Carolina B. Padovan, Analy S. A. Melo, Sarah S. Gonçalves, Viviane R. Azevedo, Marília M. Ogawa, Tainá Veras Sandes Freitas, and Arnaldo L. Colombo. 2013. "Molecular Identification of Melanised Non-Sporulating Moulds: A Useful Tool for Studying the Epidemiology of Phaeohyphomycosis." *Mycopathologia* 175 (5–6): 445–54. <https://doi.org/10.1007/s11046-012-9608-x>.

- Seyedmousavi, Seyedmojtaba, Sandra de M G Bosco, Sybren de Hoog, Frank Ebel, Daniel Elad, Renata R Gomes, Ilse D Jacobsen, et al. 2018. "Fungal Infections in Animals: A Patchwork of Different Situations." *Medical Mycology* 56 (suppl_1): S165–87. <https://doi.org/10.1093/mmy/myx104>.
- Seyedmousavi, Seyedmojtaba, J. Guillot, and G. S. de Hoog. 2013. "Phaeohyphomycoses, Emerging Opportunistic Diseases in Animals." *Clinical Microbiology Reviews* 26 (1): 19–35. <https://doi.org/10.1128/cmr.00065-12>.
- Soto, E, C Richey, SR Reichley, B Stevens, KV Kenelty, J Lewis, B Byrne, et al. 2017. "Diversity of *Veronaea Botryosa* from Different Hosts and Evaluation of Laboratory Challenge Models for Phaeohyphomycosis in *Acipenser Transmontanus*." *Diseases of Aquatic Organisms* 125 (1): 7–18. <https://doi.org/10.3354/dao03134>.
- Steckler, NK, RPE Yanong, DB Poudel, A Nyaoke, DA Sutton, JR Lindner, BL Wickes, S Jr Frasca, JC Wolf, and TB Waltzek. 2014. "New Disease Records for Hatchery-Reared Sturgeon. II. Phaeohyphomycosis Due to *Veronaea Botryosa*." *Diseases of Aquatic Organisms* 111 (3): 229–38. <https://doi.org/10.3354/dao02755>.
- White, T.J., T. Bruns, S. Lee, and J. Taylor. 1990. "Amplification and Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics." *PCR Protocols* 18: 315–22. <https://doi.org/10.1016/b978-0-12-372180-8.50042-1>.
- Yazdi, Z, MJ Griffin, F Pierezan, A Eetemadi, K Shahin, and E Soto. 2021. "Quantitative PCR for Detection and Quantification of *Veronaea Botryosa* in Fish and Environmental Samples." *Diseases of Aquatic Organisms* 144 (May): 175–85. <https://doi.org/10.3354/dao03582>.