

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Hairy black holes by gravitational decoupling

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Hairy black holes by gravitational decoupling / Ovalle, J.; Casadio, R.; Contreras, E.; Sotomayor, A.. - In: PHYSICS OF THE DARK UNIVERSE. - ISSN 2212-6864. - ELETTRONICO. - 31:(2021), pp. 100744.1-100744.10. [10.1016/j.dark.2020.100744]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/780131> since: 2020-11-12

*Published:*

DOI: <http://doi.org/10.1016/j.dark.2020.100744>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

J. Ovalle, R. Casadio, E. Contreras, A. Sotomayor, *Hairy black holes by gravitational decoupling*, Physics of the Dark Universe, Volume 31, 2021.

The final published version is available online at:  
<https://doi.org/10.1016/j.dark.2020.100744>

#### Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

# Hairy black holes by gravitational decoupling

J. Ovalle\*

*Research Centre of Theoretical Physics and Astrophysics, Institute of Physics,  
Silesian University in Opava, CZ-746 01 Opava, Czech Republic.*

R. Casadio†

*Dipartimento di Fisica e Astronomia, Alma Mater Università di Bologna, 40126 Bologna, Italy  
Istituto Nazionale di Fisica Nucleare, Sezione di Bologna, 40127 Bologna, Italy*

E. Contreras‡

*Departamento de Física, Colegio de Ciencias e Ingeniería,  
Universidad San Francisco de Quito, Quito, Ecuador.*

A. Sotomayor§

*Departamento de Matemáticas, Universidad de Antofagasta, Antofagasta, Chile.*

Black holes with hair represented by generic fields surrounding the central source of the vacuum Schwarzschild metric are examined under the minimal set of requirements consisting of i) the existence of a well defined event horizon, namely, we require the Killing horizon coincides with the causal horizon and ii) the strong or dominant energy condition for the hair outside the horizon. We develop our analysis by means of the gravitational decoupling approach. We find that trivial deformations of the seed Schwarzschild vacuum preserve the energy conditions and provide a new mechanism to evade the no-hair theorem based on a primary hair associated with the charge generating these transformations. Under the above conditions i) and ii), this charge consistently increases the entropy from the minimum value given by the Schwarzschild geometry. As a direct application, we find a non-trivial extension of the Reissner-Nordström black hole showing a surprisingly simple horizon. Finally, the non-linear electrodynamics generating this new solution is fully specified.

## I. INTRODUCTION

Black holes are among the most extraordinary objects in the Universe. The direct observations of black holes through the detection of gravitational waves [1, 2] and the reconstruction of the black hole shadow [3] have definitely raised them from being exotic solutions of general relativity, to the status of real astrophysical systems with well determined characteristics.

The original *no-hair* conjecture states that black hole solutions should not carry any other charges [4] except three fundamental parameters, namely the mass  $M$ , angular momentum  $J$  and electric charge  $Q$  [5]. However, there could exist other charges associated with inner gauge symmetries (and fields), and it is now known that black holes could also have (soft) quantum hair [6]. The *general* existence of *hairy* black hole solutions is precisely the topic under study in this article.

For a long time different scenarios have been studied for circumventing the no-hair theorem (see Refs. [7–13] for some recent works and Refs. [14–17] for earlier works). For instance, scalar fields have played a preponderant role, mainly due to their simplicity and also in analogy

with particle physics and cosmology (see also Refs. [18–20] and references therein). In this paper, following our previous work [21], instead of considering specific fundamental fields to generate hair in black holes, we shall just assume the presence of a generic source in addition to the one generating the vacuum Schwarzschild geometry. We then impose a minimal set of conditions we expect should hold for hairy black holes, namely: i) we require that the system has a well-defined event horizon and ii) the additional source is described by a conserved energy-momentum tensor  $\theta_{\mu\nu}$  which satisfies either the strong (SEC) or the dominant energy condition (DEC) in the region outside the event horizon.

From the technical point of view, we will also assume the energy-momentum tensor  $\theta_{\mu\nu}$  is *decoupled from the vacuum* by means of the extended gravitational decoupling method (EGD henceforth) of Ref. [22]. This approach was originally introduced in Ref. [23] in the form of the so-called minimal geometric deformation (MGD) [24, 25] (for some earlier works on the MGD, see for instance Refs. [26–37], and Refs. [38–56] for some recent applications). Analogously to the electro-vacuum and scalar-vacuum cases, in this work we will thus consider a Schwarzschild black hole surrounded by a spherically symmetric “tensor-vacuum”, represented by the aforementioned  $\theta_{\mu\nu}$ . Following the EGD, we can separate the complete Einstein field equations and obtain the “quasi-Einstein” equations for  $\theta_{\mu\nu}$  [See Eqs. (20)-(22) below]. These are precisely the equations of motion for the deformed Schwarzschild vacuum, which after decoupling,

---

\* jorge.ovalle@physics.slu.cz

† casadio@bo.infn.it

‡ econtreras@usfq.edu.ec

§ adrian.sotomayor@uantof.cl

contains five unknown functions that can be further analysed by imposing the two conditions discussed above. Finally, we want to emphasize that this is the first work in which we investigate black hole geometries by means of the EGD starting from general physical requirements on the horizon and black hole hair. We will show that, when we apply this approach to the Schwarzschild metric, the deformation leads to known results, namely, the Schwarzschild-de Sitter black hole or at best, the well-known Kiselev black hole [57]. The reason is that evading the no-hair theorem in a physically sensible way is a very non-trivial task.

The paper is organised as follows: in Section II, we first review the fundamentals of the EGD approach to a spherically symmetric system containing two sources; in Section III, after imposing a simple condition to guarantee a well defined horizon, and after considering the strong and dominant energy conditions on  $\theta_{\mu\nu}$ , we show a new way to evade the no-hair theorem. We apply this to generate two new families of hairy black holes containing primary hairs. As a special case, we show a non-trivial extension of the Reissner-Nordström black hole associated with a non-linear electrodynamics; finally, we summarize our conclusions in Section IV.

## II. GRAVITATIONAL DECOUPLING

In this Section, we briefly review the EGD for spherically symmetric gravitational systems described in detail in Ref. [22]. The gravitational decoupling approach and its extended version EGD are particularly attractive for at least three reasons [58–92]: a) it allows for extending known (seed) solutions of the Einstein field equations into more complex ones; b) it can be used to systematically reduce (decouple) a complex energy-momentum tensor  $T_{\mu\nu}$  into simpler components; and c) it can be used to find solutions in gravitational theories beyond Einstein's. In light of the above, it is natural to apply the EGD for the purpose of describing hairy black holes.

Let us consider the Einstein field equations <sup>1</sup>

$$G_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = k^2 \tilde{T}_{\mu\nu} , \quad (1)$$

with a total energy-momentum tensor containing two contributions,

$$\tilde{T}_{\mu\nu} = T_{\mu\nu} + \theta_{\mu\nu} , \quad (2)$$

where  $T_{\mu\nu}$  is usually associated with some known solution of general relativity, whereas  $\theta_{\mu\nu}$  may contain new fields or a new gravitational sector. Since the Einstein tensor

$G_{\mu\nu}$  satisfies the Bianchi identity, the total source must be covariantly conserved,

$$\nabla_{\mu} \tilde{T}^{\mu\nu} = 0 . \quad (3)$$

For spherically symmetric and static systems, the metric  $g_{\mu\nu}$  can be written as

$$ds^2 = e^{\nu(r)} dt^2 - e^{\lambda(r)} dr^2 - r^2 d\Omega^2 , \quad (4)$$

where  $\nu = \nu(r)$  and  $\lambda = \lambda(r)$  are functions of the areal radius  $r$  only and  $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$ . The Einstein equations (1) then read

$$k^2 (T_0^0 + \theta_0^0) = \frac{1}{r^2} - e^{-\lambda} \left( \frac{1}{r^2} - \frac{\lambda'}{r} \right) \quad (5)$$

$$k^2 (T_1^1 + \theta_1^1) = \frac{1}{r^2} - e^{-\lambda} \left( \frac{1}{r^2} + \frac{\nu'}{r} \right) \quad (6)$$

$$k^2 (T_2^2 + \theta_2^2) = -\frac{e^{-\lambda}}{4} \left( 2\nu'' + \nu'^2 - \lambda'\nu' + 2\frac{\nu' - \lambda'}{r} \right) \quad (7)$$

where  $f' \equiv \partial_r f$  and  $\tilde{T}_3^3 = \tilde{T}_2^2$  due to the spherical symmetry. By simple inspection, we can identify in Eqs. (5)-(7) an effective density

$$\tilde{\rho} = T_0^0 + \theta_0^0 , \quad (8)$$

an effective radial pressure

$$\tilde{p}_r = -T_1^1 - \theta_1^1 , \quad (9)$$

and an effective tangential pressure

$$\tilde{p}_t = -T_2^2 - \theta_2^2 . \quad (10)$$

Moreover, the anisotropy

$$\Pi \equiv \tilde{p}_t - \tilde{p}_r \quad (11)$$

usually does not vanish and the system of Eqs. (5)-(7) may be treated as an anisotropic fluid [93, 94].

We next consider a solution to the Eqs. (1) for the seed source  $T_{\mu\nu}$  alone [that is,  $\theta_{\mu\nu} = 0$ ], which we write as

$$ds^2 = e^{\xi(r)} dt^2 - e^{\mu(r)} dr^2 - r^2 d\Omega^2 , \quad (12)$$

where

$$e^{-\mu(r)} \equiv 1 - \frac{k^2}{r} \int_0^r x^2 T_0^0(x) dx = 1 - \frac{2m(r)}{r} \quad (13)$$

is the standard general relativity expression containing the Misner-Sharp mass function  $m = m(r)$ . The addition of the source  $\theta_{\mu\nu}$  can then be accounted for by the extended geometric deformation (EGD) of the seed metric (12), namely

$$\xi \rightarrow \nu = \xi + \alpha g \quad (14)$$

$$e^{-\mu} \rightarrow e^{-\lambda} = e^{-\mu} + \alpha f , \quad (15)$$

where  $f$  and  $g$  are respectively the geometric deformations for the radial and temporal metric components,

<sup>1</sup> We use units with  $c = 1$  and  $k^2 = 8\pi G_N$ , where  $G_N$  is Newton's constant.

with the parameter  $\alpha$  introduced to keep track of these deformations. We emphasize that the expressions in Eqs. (14) and (15) should not be viewed as a coordinate transformation. In fact, as previously mentioned, these expressions represent the change in the spacetime geometry (12) generated by a physical source with energy-momentum tensor  $\theta_{\mu\nu}$ . Moreover, for a given Lagrangian  $\mathcal{L}$ , we can always derive the energy-momentum  $\theta_{\mu\nu}$  producing the geometric deformation in Eqs. (14) and (15) from the standard definition

$$\theta_{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g}\mathcal{L})}{\delta g^{\mu\nu}} = 2 \frac{\delta\mathcal{L}}{\delta g^{\mu\nu}} - g_{\mu\nu}\mathcal{L}. \quad (16)$$

The opposite procedure to obtain  $\mathcal{L}$  from  $\theta_{\mu\nu}$  is usually more complicated. For example, we will show in Section III how to find the Lagrangian  $\mathcal{L}$  for a non-linear electrodynamics by the  $P$ -dual formalism [95, 96].

By means of Eqs. (14) and (15), the Einstein equations (5)-(7) are separated in two sets: A) one is given by the standard Einstein field equations with the energy-momentum tensor  $T_{\mu\nu}$ , that is

$$k^2 T_0^0 = \frac{1}{r^2} - e^{-\mu} \left( \frac{1}{r^2} - \frac{\mu'}{r} \right), \quad (17)$$

$$k^2 T_1^1 = \frac{1}{r^2} - e^{-\mu} \left( \frac{1}{r^2} + \frac{\xi'}{r} \right), \quad (18)$$

$$k^2 T_2^2 = -\frac{e^{-\mu}}{4} \left( 2\xi'' + \xi'^2 - \mu'\xi' + 2\frac{\xi' - \mu'}{r} \right), \quad (19)$$

which is assumed to be solved by the seed metric (12); B) the second set contains the source  $\theta_{\mu\nu}$  and reads

$$k^2 \theta_0^0 = -\alpha \frac{f}{r^2} - \alpha \frac{f'}{r}, \quad (20)$$

$$k^2 \theta_1^1 + \alpha Z_1 = -\alpha f \left( \frac{1}{r^2} + \frac{\nu'}{r} \right) \quad (21)$$

$$k^2 \theta_2^2 + \alpha Z_2 = -\alpha \frac{f}{4} \left( 2\nu'' + \nu'^2 + 2\frac{\nu'}{r} \right) - \alpha \frac{f'}{4} \left( \nu' + \frac{2}{r} \right), \quad (22)$$

where

$$Z_1 = \frac{e^{-\mu} g'}{r} \quad (23)$$

$$4Z_2 = e^{-\mu} \left( 2g'' + g'^2 + \frac{2g'}{r} + 2\xi'g' - \mu'g' \right). \quad (24)$$

The above equations clearly show that the tensor  $\theta_{\mu\nu}$  must vanish when the deformations vanish ( $\alpha = 0$ ). Moreover, on assuming  $g = 0$ , Eqs. (20)-(22) reduce to the simpler “quasi-Einstein” system of the MGD of Ref. [23], in which  $f$  is only determined by  $\theta_{\mu\nu}$  and the undeformed metric (12).

It is also important to discuss the conservation equa-

tion (3) which now reads

$$\begin{aligned} (T_1^1)' - \frac{\xi'}{2} (T_0^0 - T_1^1) - \frac{2}{r} (T_2^2 - T_1^1) \\ - \frac{\alpha g'}{2} (T_0^0 - T_1^1) \\ + (\theta_1^1)' - \frac{\nu'}{2} (\theta_0^0 - \theta_1^1) - \frac{2}{r} (\theta_2^2 - \theta_1^1) = 0. \end{aligned} \quad (25)$$

The first line precisely represents the divergence of  $T_{\mu\nu}$  computed with the covariant derivative  $\nabla^{(\xi,\mu)}$  for the metric (12), and is a linear combination of the Einstein field equations (17)-(19). In fact, the Einstein tensor  $G_{\mu\nu}$  for the metric (12) must satisfy its respective Bianchi identity, and the energy momentum tensor  $T_{\mu\nu}$  is therefore conserved by construction in this geometry,

$$\nabla_{\sigma}^{(\xi,\mu)} T^{\sigma}_{\nu} = 0, \quad (26)$$

Upon using the deformed metric in Eq. (4), one instead obtains

$$\nabla_{\sigma} T^{\sigma}_{\nu} = \nabla_{\sigma}^{(\xi,\mu)} T^{\sigma}_{\nu} - \alpha \frac{g'}{2} (T_0^0 - T_1^1) \delta_{\nu}^1, \quad (27)$$

which explains the origin of the term in the second line of Eq. (25). Finally, Eq. (25) becomes

$$\begin{aligned} 0 &= \nabla_{\sigma} T^{\sigma}_{\nu} + \nabla_{\sigma} \theta^{\sigma}_{\nu} \\ &= \alpha \frac{g'}{2} (T_0^0 - T_1^1) \delta_{\nu}^1 + \nabla_{\sigma} \theta^{\sigma}_{\nu} \\ &= (\theta_1^1)' - \frac{\nu'}{2} (\theta_0^0 - \theta_1^1) - \frac{2}{r} (\theta_2^2 - \theta_1^1) \\ &\quad - \alpha \frac{g'}{2} (T_0^0 - T_1^1), \end{aligned} \quad (28)$$

which is also a linear combination of the “quasi-Einstein” field equations (20)-(22) for the source  $\theta_{\mu\nu}$ . We therefore conclude that the two sources  $T_{\mu\nu}$  and  $\theta_{\mu\nu}$  can be successfully decoupled by means of the EGD. This result is particularly remarkable, since it does not require a perturbative expansion in the parameter  $\alpha$  [25].

### III. HAIRY BLACK HOLES

Conditions to evade the no-hair theorem have been investigated for a long time [7–17]. A straightforward possibility is to fill the static vacuum with some source of potentially fundamental origin, often described as a scalar field [18–20]. We recently considered a more general scenario within the MGD approach, where the Schwarzschild vacuum for  $T_{\mu\nu} = 0$  is filled with a generic static and spherically symmetric source of energy-momentum tensor  $\theta_{\mu\nu}$ , namely, a “tensor-vacuum” [21]. This leads to hairy black hole solutions with a rich geometry described by the mass  $M$  and a discrete set of charges generating primary hair. However, the MGD (15) leaves the temporal component of the metric (4) exactly equal to the Schwarzschild one,

$$e^{\nu} = e^{\xi} = 1 - \frac{2M}{r}, \quad (29)$$

which hinders the existence of stable black holes with a well-defined event horizon. Indeed, the relation (29) implies that only hairy black hole solutions with the event horizon at  $r_{\text{H}} = 2M$  can be free of pathologies. The advantage of the EGD is that the temporal component is also modified according to Eq. (14), thus yielding a potentially larger number of hairy black hole solutions with horizons other than  $r_{\text{H}} = 2M$ .

We apply the analysis in the previous Section to the particular case of  $T_{\mu\nu} = 0$ . The seed metric (12) is thus given by the Schwarzschild solution with

$$e^{\xi} = e^{-\mu} = 1 - \frac{2M}{r}, \quad (30)$$

which solves Eqs. (17)-(19) for  $T_{\mu\nu} = 0$ . In order to find hairy black holes, we then need to solve the resulting “quasi-Einstein” system (20)-(22), which contain the three components of  $\theta_{\mu\nu}$  and the two deformations  $f$  and  $g$ . Furthermore, we reduce the number of unknown quantities, so that they specify a unique solution, by prescribing the two conditions discussed in the Introduction.

First of all, in order to have black hole solutions with a well-defined horizon structure, we demand that the deformed metric (4) satisfies <sup>2</sup>

$$e^{\nu} = e^{-\lambda}. \quad (31)$$

The condition (31) defines the so-called spacetimes of the Kerr-Schild class, which has been extensively studied (see for instance Refs. [97–100]). This condition ensures that the radius  $r = r_{\text{H}}$  such that

$$e^{\nu(r_{\text{H}})} = e^{-\lambda(r_{\text{H}})} = 0 \quad (32)$$

will be both a killing horizon ( $e^{\nu} = 0$ ) and a causal horizon ( $e^{-\lambda} = 0$ ). A direct consequence of the condition (31), following from the Einstein equations (5) and (6), is the equation of state

$$\tilde{p}_r = -\tilde{\rho}. \quad (33)$$

Therefore, only negative radial pressure is allowed (for positive density). The condition (31) and the Schwarzschild solution (30) then relate the metric deformations  $f$  and  $g$  according to

$$\alpha f(r) = \left(1 - \frac{2M}{r}\right) \left(e^{\alpha g(r)} - 1\right), \quad (34)$$

so that the line element (4) becomes

$$ds^2 = \left(1 - \frac{2M}{r}\right) e^{\alpha g(r)} dt^2 - \left(1 - \frac{2M}{r}\right)^{-1} e^{-\alpha g(r)} dr^2 - r^2 d\Omega^2. \quad (35)$$

<sup>2</sup> We remark that Eq. (31) implies the condition (32) but is not necessary for it to hold, as one could also consider cases with  $e^{\nu} \neq e^{-\lambda}$  for  $r \neq r_{\text{H}}$ .

Once more we remark that Eqs. (14) and (15) are neither conformal transformations nor scale transformations, although the factor  $e^{\alpha g}$  could be interpreted as a Weyl factor related with a “partial scale transformation”. <sup>3</sup>

We are now left with the deformation  $g$  and the three components of  $\theta_{\mu\nu}$ , which must satisfy the three “quasi-Einstein” Eqs. (20)-(22). We can therefore impose some physically motivated restriction on  $g$  or a constraint on  $\theta_{\mu\nu}$ , like a reasonable equation of state. For instance, in the region  $r \geq 2M$ , we can consider the tensor-vacuum satisfies

$$\theta_0^0 = a\theta_1^1 + b\theta_2^2, \quad (36)$$

with  $a$  and  $b$  constants. In this case, Eqs. (20)-(22) yield the differential equation

$$br(r-2M)h'' + 2[(a+b-1)r - 2(a-1)M]h' + 2(a-1)h = 2(a-1), \quad (37)$$

for

$$h(r) = e^{\alpha g(r)}. \quad (38)$$

The solution can be written as

$$e^{\alpha g(r)} = 1 + \frac{1}{r-2M} \left[ \ell_0 + r \left( \frac{\ell}{r} \right)^{2(1-a)/b} \right], \quad (39)$$

where  $\ell_0$  and  $\ell$  are constants (proportional to  $\alpha$ ) with dimensions of a length. By using this expression in the line element (35), we obtain the metric functions

$$e^{\nu} = e^{-\lambda} = 1 - \frac{2\mathcal{M}}{r} + \frac{\ell^n}{r^n}, \quad (40)$$

where we defined the new mass as  $\mathcal{M} = M + \ell_0/2$  and

$$n = \frac{2}{b}(a-1), \quad (41)$$

with  $n > 1$  for a correct asymptotic behavior. <sup>4</sup> The possible horizons  $r = r_{\text{H}}$  are given by the solutions of

$$r_{\text{H}}^n - 2\mathcal{M}r_{\text{H}}^{n-1} + \ell = 0, \quad (42)$$

and the space-time represents a Kiselev black hole [57], which was analyzed in great detail by Visser in Ref. [101]. This line element is produced by the effective density

$$\tilde{\rho} = \theta_0^0 = \alpha \frac{(n-1)\ell^n}{k^2 r^{n+2}}, \quad (43)$$

<sup>3</sup> Since there is some confusion in the literature, we adhere to the convention that conformal transformations are coordinate transformations leaving angles invariant, whereas Weyl (or scale) transformations are more general transformations of the metric which leave the angles invariant but do not affect the coordinates.

<sup>4</sup> Note that a trivial deformation which leaves the Schwarzschild metric (30) unaffected is also recovered by setting  $a = 1$  ( $n = 0$ ) in Eq. (37). We will have more to say about this in the next subsections.

the effective radial pressure

$$\tilde{p}_r = -\theta_1^1 = -\tilde{\rho}, \quad (44)$$

and the effective tangential pressure

$$\tilde{p}_t = -\theta_2^2 = \frac{n}{2} \tilde{\rho}. \quad (45)$$

The anisotropy (11) is thus given by

$$\Pi = \left(\frac{n}{2} + 1\right) \tilde{\rho}. \quad (46)$$

We see that the Schwarzschild-de Sitter solution ( $n = -2$ ) is the only one which allows for an isotropic tensor-vacuum. On the other hand, the DEC, namely  $\tilde{\rho} \geq |\tilde{p}_r|$  and  $\tilde{\rho} \geq |\tilde{p}_t|$ , yields  $n \leq 2$ . Combining this with asymptotic flatness ( $n > 1$ ), we obtain the range

$$1 \leq n \leq 2. \quad (47)$$

The extreme cases  $n = 1$  and  $n = 2$  are, respectively, the Schwarzschild solution and the conformal solution with traceless  $\theta_{\mu\nu}$ , like the Maxwell case. Since the Kiselev black hole has already been studied extensively, we will abandon Eq. (36) and continue analysing the deformed metric (35) based on energy conditions.

Let us recall that the energy conditions are a set of requirements which are usually imposed on the energy-momentum tensor to avoid exotic matter sources, hence we can see them as sensible guidelines to avoid classically unphysical configurations [102, 103]. In particular, we will impose energy conditions on the source  $\theta_{\mu\nu}$  in the region of space-time accessible to an outer observer (while possibly relax them inside the event horizon).

### A. Strong energy condition

Let us start with the SEC, namely

$$\begin{aligned} \tilde{\rho} + \tilde{p}_r + 2\tilde{p}_t &\geq 0 \\ \tilde{\rho} + \tilde{p}_r &\geq 0 \\ \tilde{\rho} + \tilde{p}_t &\geq 0, \end{aligned} \quad (48)$$

which, as a consequence of Eq. (33), reduce to

$$\theta_2^2 \leq 0 \quad (49)$$

$$\theta_0^0 \geq \theta_2^2, \quad (50)$$

where we used the definitions (8) and (10).

Using Eqs. (20) and (22) we find that the conditions (49) and (50) respectively lead to the second-order linear differential inequalities

$$G_1(r) \equiv (r - 2M)h'' + 2h' \geq 0, \quad (51)$$

$$G_2(r) \equiv r(r - 2M)h'' + 4Mh' - 2h + 2 \geq 0, \quad (52)$$

where  $h$  was defined in Eq. (38). It is now useful to recall that a Grönwall's inequality of the form

$$U'(r) \leq \beta(r)U(r), \quad (53)$$

in the interval  $r_0 \leq r \leq \infty$ , admits the solution

$$U(r) \leq U(r_0) e^{\int_{r_0}^r \beta(s) ds} \equiv U_0(r), \quad (54)$$

where the bounding function  $U_0$  is obtained by saturating the differential inequality (53). For the inequality (51), we can define

$$U(r) = -h'(r) \quad (55)$$

and  $\beta(r) = -2(r - 2M)^{-1}$ , so that Eq. (54) yields

$$h'(r) \geq h'(r_0) \left(\frac{r_0 - 2M}{r - 2M}\right)^2, \quad (56)$$

and finally

$$h(r) \geq h(r_0) + (r_0 - 2M) \left(1 - \frac{r_0 - 2M}{r - 2M}\right) h'(r_0). \quad (57)$$

We therefore find that the bounding function solving  $G_1(r) = 0$  behaves as

$$h_1(r) \sim c_1 - \frac{\ell_1}{r - 2M}, \quad (58)$$

where  $c_1$  is a dimensionless constants and  $\ell_1$  a constant with dimensions of a length. However, any deformation of the form in Eq. (58) plugged into the metric (35) uniquely leads to

$$e^\nu = e^{-\lambda} = c_1 \left(1 - \frac{2c_1 M + \ell_1}{c_1 r}\right), \quad (59)$$

which becomes the Schwarzschild solution (30) by imposing asymptotic flatness (that is, setting  $c_1 = 1$ ) and rescaling the mass  $2M + \ell_1 \rightarrow 2M$ . Indeed, we notice that  $G_1(r) = 0$  equals the differential equation (37) for  $a = 1$ , whose solution yields the Schwarzschild metric. On the other hand, following the same procedure for the inequality (52), we find that the bounding function satisfying  $G_2(r) = 0$  behaves as

$$h_2(r) \sim 1 - \frac{1}{r - 2M} \left(\ell_2 - \frac{\Lambda}{3} r^3\right), \quad (60)$$

where  $\ell_2$  is again a length and  $\Lambda$  a constant with dimensions of the inverse of a squared length. Likewise, the deformation (60) plugged into the metric (35) leads to

$$e^\nu = e^{-\lambda} = 1 - \frac{2M + \ell_2}{r} + \frac{\Lambda}{3} r^2, \quad (61)$$

which is the Schwarzschild-de Sitter metric with cosmological constant  $\Lambda$  and mass  $2M + \ell_2 \rightarrow 2M$ . Again, notice that  $G_2(r) = 0$  is the differential equation (37) for  $a = 0$  and  $b = 1$  and the bounding solution  $h_2$  for the extremal case  $G_2(r) = 0$  (with  $\alpha \neq 0$ ) leads to the line element (40) with  $n = -2$ . Since both inequalities (51) and (52) must hold, the unique bounding deformation  $h_0(r)$  which solves  $G_1(r) = G_2(r) = 0$  is obtained when



Eq. (58) equals Eq. (60), that is for  $\Lambda = 0$ . We thus conclude that the bounding deformations which saturate the SEC are of the form

$$h_0(r) = 1 - \frac{\ell_0}{r - 2M} \quad (62)$$

and leave the Schwarzschild geometry unaffected. This is not at all surprising since  $G_1(r) = G_2(r) = 0$  is tantamount to  $\tilde{\rho} = \tilde{p}_r = \tilde{p}_t = 0$  and the Schwarzschild geometry cannot possibly be deformed in this case.

Before we proceed to consider deformations which do not saturate the inequalities (51) and (52), we notice that the transformation

$$h(r) \rightarrow \bar{h}(r) \equiv h(r) - \frac{\ell_0}{r - 2M} \quad (63)$$

leaves  $G_1(r)$  and  $G_2(r)$  invariant. Under this transformation, the metric functions change as

$$e^\nu = e^{-\lambda} = \left(1 - \frac{2M}{r}\right) h(r) \rightarrow \left(1 - \frac{2M}{r}\right) h(r) - \frac{\ell_0}{r}, \quad (64)$$

where it is reasonable to assume that  $h(r)$  admits an expansion in powers of  $1/r$  for a regular exterior. In particular, the effect of the metric transformation (64) will amount to the usual shift of the mass  $M \rightarrow \mathcal{M} = M + \ell_0/2$  at order  $1/r$ . This redefinition of the asymptotic mass, in turn, will introduce a new dependence on the length  $\ell_0$  in  $h(r)$  whenever the latter contains the unshifted seed mass  $M = \mathcal{M} - \ell_0/2$ , thus generating a new solution with parameters  $\mathcal{M}$  and  $\ell_0$ . Of course, for  $h(r) \sim 1$ , Eq. (63) acts like a ‘‘gauge’’ symmetry corresponding to the trivial deformations (62) of the seed Schwarzschild geometry (30).

We have just seen that the parameter  $\ell_0$  in Eq. (63) appears as a new ‘‘gauge’’ charge. The way this all works can be made more explicit by considering concrete examples. Since we are interested in solutions with a proper horizon at  $r = r_H \sim 2M$ , which also behave approximately like the Schwarzschild metric for  $r \gg 2M$  (so as to meet all experimental bounds in the weak field regime), we could consider any positive function  $G_1(r)$  satisfying the boundary conditions

$$G_1(r) = 0 \quad \text{for} \quad \begin{cases} r \sim 2M \\ r \gg M, \end{cases} \quad (65)$$

A simple example of such a function containing just the parameters  $\alpha$  and  $M$  is given by

$$G_{\alpha, M}(r) = \frac{\alpha}{M^2} (r - 2M) e^{-r/M}. \quad (66)$$

Upon solving Eq. (51) for the corresponding deformation  $h$ , we obtain

$$h(r) = c_1 - \alpha \frac{\ell - r e^{-r/M}}{r - 2M}, \quad (67)$$




FIG. 1. Metric function for different values of  $\ell$  with  $\alpha = 0.4$  and  $M = 1$ . The shift of the horizon  $r_H$  is controlled by the parameter  $\ell$ . The Schwarzschild horizon corresponds to the saturated case of the inequality (74).

where  $\alpha \ell = \ell_0$  and we can set  $c_1 = 1$  to recover the proper limit for  $\alpha = 0$  (in which  $G_{\alpha, M} \rightarrow 0$ ). The deformation in Eq. (67) must also satisfy the inequality (52), which becomes

$$\frac{\alpha}{M^2} (r^2 - 2M^2) e^{-r/M} \geq 0, \quad (68)$$

and it is satisfied for all  $r \geq \sqrt{2}M$ .

Finally, using (67) in the line element (35) yields the metric functions

$$e^\nu = e^{-\lambda} = 1 - \frac{2\mathcal{M}}{r} + \alpha e^{-r/(\mathcal{M} - \alpha \ell/2)}, \quad (69)$$

where we modded out the term proportional to  $\ell$ , corresponding to the gauge transformation (63), by introducing the mass  $\mathcal{M} = M + \alpha \ell/2$ . The new solution (69) thus asymptotically approaches the Schwarzschild geometry with a total mass  $\mathcal{M}$ . The source  $\theta_{\mu\nu}$  decays exponentially away from the center of the system, as can be seen from the effective density

$$\tilde{\rho} = \theta_0^0 = -\tilde{p}_r = \frac{\alpha e^{-r/M}}{k^2 M r^2} (r - M), \quad (70)$$

and the effective tangential pressure

$$\tilde{p}_t = -\theta_2^2 = \frac{\alpha e^{-r/M}}{2k^2 M^2 r} (r - 2M). \quad (71)$$

We can immediately see some important features of the metric (69). The first one is that the physical singularity at  $r = 0$  remains and is further reflected in the singular behaviour of the effective quantities in Eqs. (70) and (71). The second feature is that the SEC (49) is only satisfied as long as  $r \geq 2M$ , as we can see from (71). Also notice that in the limit  $M \rightarrow 0$  the source  $\theta_\mu^\nu \rightarrow 0$  for  $r > 0$ , and  $\mathcal{M} \rightarrow \alpha \ell$ . In other words, the source  $\theta_\mu^\nu$  approaches a Dirac-delta function for vanishing seed mass  $M$ .






FIG. 2. Effective source terms  $\times 10^4$  for  $\alpha = 0.2$ . The horizon is located at  $r_{\text{H}} \gtrsim 2M$ , with  $M = 1$ .

The equation determining the horizon  $r = r_{\text{H}}$  of the metric (69) is given by

$$\alpha \ell = r_{\text{H}} - 2M + \alpha r_{\text{H}} e^{-r_{\text{H}}/M}, \quad (72)$$

which allows to write the metric (69) in terms of its horizon in the form

$$e^{\nu} = e^{-\lambda} = 1 - \frac{r_{\text{H}}}{r} + \alpha \left( e^{-r/M} - \frac{r_{\text{H}}}{r} e^{-r_{\text{H}}/M} \right). \quad (73)$$

It is of course impossible to find analytical solutions to Eq. (72), except for particular values of the parameters. For example, according to our prescription for the SEC, we need  $r_{\text{H}} \geq 2M$ , or

$$\ell \geq 2M/e^2. \quad (74)$$

The extremal case  $\ell = 2M/e^2$  leads to the solution

$$e^{\nu} = e^{-\lambda} = 1 - \frac{2M}{r} + \alpha \left( e^{-r/M} - \frac{2M}{e^2 r} \right). \quad (75)$$

which has the horizon at  $r_{\text{H}} = 2M$ . The SEC (49) and (50) are both satisfied in the outer region, but the black hole should have the same thermodynamic properties of the Schwarzschild geometry.

The metric function (69) is plotted in Fig. 1 for a given value of  $\alpha$ , which shows that the horizon is shifted to larger radii when  $\ell$  increases from the minimum allowed value (74) corresponding to  $r_{\text{H}} = 2M$ . We find the same behavior when  $\alpha$  increases. In this respect, we notice that the effective density  $\tilde{\rho}$  and pressures  $\tilde{p}_r$  and  $\tilde{p}_t$  in Eqs. (70) and (71) do not depend on the parameter  $\ell$ , unlike the horizon. This allows us to choose suitable values for the parameters  $\alpha$  and  $\ell$  such that the SEC is satisfied for  $r \gtrsim r_{\text{H}}$ , as is shown in Fig. 2, where both the density and tangential pressure are positive. We conclude that the metric (69) represents a hairy black hole (having in general non-trivial thermodynamic properties) endowed with the parameters  $\{M, \alpha, \ell\}$ , where  $\ell_0 = \alpha \ell$  represents a charge associated with primary hair.

## B. Dominant energy condition

We shall next consider the DEC, which requires [102]

$$\tilde{\rho} \geq |\tilde{p}_r| \quad (76)$$

$$\tilde{\rho} \geq |\tilde{p}_t|. \quad (77)$$

In particular, we will see that these conditions allow for deforming the Schwarzschild solution into the Reissner-Nordström-de Sitter geometry with an effective charge  $Q$  and an effective cosmological constant  $\Lambda$ .

We first point out that the inequality (76) is saturated as a consequence of Eq. (33) for a positive effective density, for which Eq. (77) reduces to

$$-\tilde{\rho} \leq \tilde{p}_t \leq \tilde{\rho}. \quad (78)$$

We can again write the condition (78) in terms of the definitions (8) and (10) as

$$\theta_0^0 + \theta_2^2 \geq 0 \quad (79)$$

$$\theta_0^0 - \theta_2^2 \geq 0, \quad (80)$$

which yield respectively the differential inequalities

$$H_1(r) \equiv -r(r - 2M)h'' - 4(r - M)h' - 2h + 2 \geq 0 \quad (81)$$

$$H_2(r) \equiv r(r - 2M)h'' + 4Mh' - 2h + 2 \geq 0, \quad (82)$$

where we used Eqs. (20) and (22), and  $h$  is the same defined in Eq. (38).

We can then notice that  $H_1(r) = 0$  is the differential equation (37) for  $a = 0$  and  $b = -1$ . Hence, the bounding solution  $h_1$  for the extremal case  $H_1(r) = 0$  (with  $\alpha \neq 0$ ) leads to the line element (40) with  $n = 2$ , namely, the Reissner-Nordström solution (with an effective charge  $Q \sim \ell$ ). On the other hand,  $H_2(r) = 0$  is the differential equation (37) for  $a = 0$  and  $b = 1$ . Hence, the bounding solution  $h_2$  for the extremal case  $H_2(r) = 0$  leads to the line element in (40) with  $n = -2$ , namely, the Schwarzschild-de Sitter solution with cosmological constant  $\Lambda \sim \ell^{-2}$ . Since  $H_1(r) = H_2(r) = 0$  corresponds to vanishing  $\theta_{\mu\nu}$  like for the SEC, the unique bounding deformation  $h_0(r) = h_1(r) = h_2(r)$  is obtained for  $Q = \Lambda = 0$ , so that the only possible deformation remains again the trivial one in Eq. (62), which yields the seed Schwarzschild solution. Indeed, the functions  $H_1(r)$  and  $H_2(r)$  in Eqs. (81) and (82) are also invariant under the transformation (63).

Like in Section III A, we proceed to investigate deformations which do not saturate the inequalities (81) and (82) everywhere by considering positive functions  $H_1(r)$  or  $H_2(r)$  which saturate that inequalities only near the boundaries of the outer region, that is

$$H_1(r) = 0 \quad \text{for} \quad \begin{cases} r \sim 2M \\ r \gg M. \end{cases} \quad (83)$$

In fact, we can still employ the function in Eq. (66) and set

$$H_1 = M G_{\alpha, M}. \quad (84)$$

Upon solving (81) for the corresponding  $h$ , we obtain

$$h(r) = 1 - \frac{1}{r - 2M} \left( \alpha \ell + \alpha M e^{-r/M} - \frac{Q^2}{r} \right), \quad (85)$$

where  $\alpha \ell = \ell_0$  and  $Q$  is also a constant with dimension of a length and proportional to  $\alpha$ . A second constant of integration was adjusted to meet the proper Schwarzschild limit for  $\alpha \rightarrow 0$  (in which we remark that  $Q \sim \alpha$  vanishes as well). The deformation in Eq. (85) also has to satisfy the inequality (82), which reads

$$\frac{4Q^2}{r^2} \geq \frac{\alpha}{M} (r + 2M) e^{-r/M}. \quad (86)$$

Using (85) in the line element (35), we obtain the metric functions

$$e^\nu = e^{-\lambda} = 1 - \frac{2M + \alpha \ell}{r} + \frac{Q^2}{r^2} - \frac{\alpha M e^{-r/M}}{r}, \quad (87)$$

which is a sort of ‘‘charged’’ version of the solution (69) again with asymptotic mass  $\mathcal{M} = M + \alpha \ell/2$ . The effective density is now given by

$$\tilde{\rho} = \theta_0^0 = -\tilde{p}_r = \frac{Q^2}{k^2 r^4} - \frac{\alpha e^{-r/M}}{k^2 r^2} \quad (88)$$

and an effective tangential pressure reads

$$\tilde{p}_t = -\theta_2^2 = \frac{Q^2}{k^2 r^4} - \frac{\alpha e^{-r/M}}{2k^2 M r}. \quad (89)$$

We can see that

$$\tilde{\rho} - \tilde{p}_t = \frac{\alpha e^{-r/M}}{2k^2 M r^2} (r - 2M) \quad (90)$$

and the DEC is satisfied for  $r \geq 2M$ , as we originally required. We can also see that the physical singularity at  $r = 0$  remains.

The horizon radii  $r_H$  are given by solutions of

$$\alpha \ell = r_H - 2M + \frac{Q^2}{r_H} - \alpha M e^{-r_H/M}, \quad (91)$$

which allows us to write the metric functions (87) as

$$e^\nu = e^{-\lambda} = 1 - \frac{r_H}{r} \left( 1 + \frac{Q^2}{r_H^2} - \frac{\alpha M}{r_H} e^{-r_H/M} \right) + \frac{Q^2}{r^2} - \frac{\alpha M}{r} e^{-r/M}. \quad (92)$$

As with the SEC, it is always possible to choose suitable values for the parameters  $\alpha$ ,  $\ell$  and  $Q$  such that analytical solutions for  $r_H$  can be found. However, since the DEC requires  $r_H \geq 2M$ , the choice of these values cannot be arbitrary. We can see this by evaluating the density (88) at the horizon, in addition to using the expression (91), which yields

$$Q^2 \geq 4\alpha(M/e)^2 \quad \text{and} \quad \ell \geq M/e^2. \quad (93)$$

We remark that  $Q$  does not need to be an electric charge. It could be, for instance, a tidal charge of extra-dimensional origin or any other source. However, when  $Q$  represents an electric charge, we can say that the electrovacuum of the Reissner-Nordström geometry also contains a tensor-vacuum whose components are those explicitly proportional to  $\alpha$  in Eqs. (88) and (89). Let us recall that the Reissner-Nordström metric has two horizons: the event horizon

$$r_H = M + \sqrt{M^2 - Q^2} < 2M, \quad (94)$$

and an internal Cauchy horizon given by

$$r_{CH} \equiv M - \sqrt{M^2 - Q^2} < r_H. \quad (95)$$

For our solution (87), we can identify at least three cases for which the event horizon  $r_H$  has simple analytical expressions, and the DEC (76) and (77) are satisfied. As in the Reissner-Nordström metric, each one of these cases has an internal Cauchy horizon  $r_{CH} < r_H$ .

#### Case 1

Let us start by considering the case saturating the inequalities (93), for which the metric components (92) become

$$e^\nu = e^{-\lambda} = 1 - \frac{2M}{r} \left( 1 + \frac{\alpha}{2e^2} \right) + \frac{4\alpha M^2}{e^2 r^2} - \frac{\alpha M}{r} e^{-r/M}. \quad (96)$$

The event horizon is again precisely at  $r_H = 2M$ , which parallels the case of Eq. (75), and we can also write

$$r_H = \frac{e}{\sqrt{\alpha}} Q. \quad (97)$$

Notice that by defining

$$\mathcal{M} = M \left( 1 + \frac{\alpha}{2e^2} \right), \quad (98)$$

the metric functions (96) can be written in a more suggestive form as

$$e^\nu = e^{-\lambda} = 1 - \frac{2\mathcal{M}}{r} + \frac{Q^2}{r^2} - \frac{\sqrt{\alpha} Q}{2r} e^{1-2\sqrt{\alpha}r/eQ}, \quad (99)$$

which can be interpreted as a nonlinear electrodynamics coupled with gravity.

#### Case 2

The second case we consider is given by

$$Q^2 = \alpha \ell M \left( 2 + \alpha e^{-\alpha \ell/M} \right), \quad (100)$$

with  $\ell \neq 0$ , leading to

$$e^\nu = e^{-\lambda} = 1 - \frac{2M + \alpha \ell}{r} + \frac{2\alpha \ell M}{r^2} - \frac{\alpha M}{r^2} e^{-r/M} \left( r - \alpha \ell e^{\frac{r-\alpha \ell}{M}} \right). \quad (101)$$




FIG. 3. Metric function for the three different analytic cases with  $\alpha = 0.8$  and  $M = 1$  for  $\ell = 0.4, 4, 1$ , respectively.




FIG. 4. Effective source terms  $\times 10^4$  for  $\alpha = 0.1$ ,  $\ell = 0.4$ ,  $Q = 0.3$ , with  $M = 1$ .

The event horizon is now at  $r_H = \alpha\ell = \ell_0 \geq 2M$ . If we further assume  $\alpha\ell \sim M$ , we can express  $M = M(Q)$  from (100) and the metric functions (101) can again be interpreted as a nonlinear electrodynamics coupled with gravity.

### Case 3

Finally, we consider

$$Q^2 = \alpha M (2M + \alpha\ell) e^{-\frac{(2M + \alpha\ell)}{M}}, \quad (102)$$

so that

$$e^\nu = e^{-\lambda} = 1 - \frac{2M + \alpha\ell}{r} - \frac{\alpha M}{r^2} e^{-r/M} \times \left[ r - (2M + \alpha\ell) e^{\frac{r - (2M + \alpha\ell)}{M}} \right]. \quad (103)$$

The event horizon is at  $r_H = 2M + \alpha\ell = 2\mathcal{M} \geq 2M$ , and the Schwarzschild horizon is recovered as usual for  $\alpha = 0$ . As in the two previous cases, the interpretation in terms of nonlinear electrodynamics is obtained for  $\alpha\ell \sim M$ .

The metric functions for the three analytical cases in Eqs. (96), (101) and (103) are displayed in Fig. 3, each for a different value of the parameter  $\ell$ . In all cases the density and pressures have the same qualitative behavior as in Fig. 4, which corresponds to the general case given by Eqs. (87)-(89). We conclude that the metric (87) represents a hairy black hole endowed with the parameters  $M, Q, \alpha$  and  $\ell$ , where  $\{Q, \ell_0 = \alpha\ell\}$  represents a potential set of charges generating primary hair. Among this new family of solutions, we can identify three different cases representing hairy black holes having simple analytical horizons. All these cases can be interpreted as a nonlinear electrodynamics coupled with gravity, whose charges are  $M, \ell_0 = \alpha\ell$  and  $Q$ .<sup>5</sup>

Finally, we want to end by emphasizing a rather important result. When the horizon in (92) has the simple form  $r_H = KM$ <sup>6</sup>, with  $K \geq 2$  in order to satisfy the DEC, the metric functions become

$$e^\nu = e^{-\lambda} = 1 - \frac{2\mathcal{M}}{r} + \frac{Q^2}{r^2} - \frac{\alpha r_H}{K r} e^{-K r/r_H}, \quad (104)$$

and the event horizon takes the simple Reissner-Nordström form

$$r_H = \bar{\mathcal{M}} + \sqrt{\bar{\mathcal{M}}^2 - \bar{Q}^2}, \quad (105)$$

where  $\bar{\mathcal{M}} = \mathcal{M}/\beta$  and  $\bar{Q}^2 = Q^2/\beta$  with

$$\beta = 1 - \alpha \frac{e^{-K}}{K}. \quad (106)$$

We conclude that the metric (104) represents a black hole associated with some non-linear electrodynamics, whose horizon is related with the Reissner-Nordström one by the quite simple expression

$$r_H = \frac{r_{\text{RN}}}{\beta} \geq r_{\text{RN}}, \quad (107)$$

since  $\beta \leq 1$ .

Regarding the field theory behind the deforming energy-momentum tensor  $\theta_{\mu\nu}$ , we remark that it is in principle always possible to find a Lagrangian  $\mathcal{L}$  associated with a non-linear electrodynamics by means of the  $P$ -dual formalism [95, 96]. Of course, there is no guarantee  $\mathcal{L}$  is fundamental or motivated by first principles. In order to specify this nonlinear electrodynamics, we identify

$$\theta_{\mu\nu} = -L_F F_{\mu\alpha} F^\alpha{}_\nu - L(F) g_{\mu\nu}, \quad (108)$$

<sup>5</sup> This way of presenting the results should not overshadow the fact that  $Q^2 \sim \alpha$ , so that the Schwarzschild geometry is always recovered for  $\alpha \rightarrow 0$ .

<sup>6</sup> Notice that  $K$  can be written as  $K \geq \frac{1}{2\alpha} \left( \frac{Q_{\text{min}}}{\ell_{\text{min}}} \right)^2$ , where  $Q_{\text{min}}$  and  $\ell_{\text{min}}$  are the saturated values in Eq. (93).

where

$$F = \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \quad \text{and} \quad L_F = \frac{dL}{dF}. \quad (109)$$

In the static spherically symmetric case, we have

$$F_{\mu\nu} = E(r) (\delta_\mu^0 \delta_\nu^1 - \delta_\mu^1 \delta_\nu^0). \quad (110)$$

Using Eqs. (31), (108)-(110) in the Einstein equations (5) and (6), we obtain

$$-\frac{2}{r^2} \frac{d\tilde{m}}{dr} = 8\pi [L(F) + E^2 L_F] \quad (111)$$

$$-\frac{1}{r} \frac{d^2\tilde{m}}{dr^2} = 8\pi L(F), \quad (112)$$

where

$$\tilde{m}(r) = \frac{k^2}{2} \int_0^r x^2 \theta_0^0(x) dx \quad (113)$$

is the Misner-Sharp mass function for the deforming tensor  $\theta_{\mu\nu}$ . The corresponding conservation equation (3) now reads  $\nabla_\mu (L_F F^{\mu\nu}) = 0$ , and leads to

$$E(r) L_F = -\frac{Q}{4\pi r^2}. \quad (114)$$

Subtracting (111) from Eq. (112), we obtain

$$r \frac{d}{dr} \left( \frac{1}{r^2} \frac{d\tilde{m}}{dr} \right) = 8\pi E^2 L_F. \quad (115)$$

Finally, combining Eqs. (114) and (115) we arrive at

$$E = -\frac{r^3}{2Q} \frac{d}{dr} \left( \frac{1}{r^2} \frac{d\tilde{m}}{dr} \right). \quad (116)$$

In our case given by the metric (104), we have

$$\tilde{m}(r) = \mathcal{M} - \frac{Q^2}{2r} + \frac{\alpha r_H}{2K} e^{-Kr/r_H}, \quad (117)$$

and the explicit form of the electric field (116) generating the black hole solution described by the metric (104) reads

$$E(r) = \frac{Q}{r^2} - \alpha e^{-\frac{Kr}{r_H}} \frac{K r + 2r_H}{4r_H Q}. \quad (118)$$

In order to find the Lagrangian  $L$  of the underlying nonlinear electrodynamics, we will use the  $P$ -dual formalism [95, 96]. This formalism is based on the Legendre transformation

$$H = 2F L_F - L, \quad (119)$$

where  $H$  represents the Hamiltonian in the dual formulation. Now, defining  $P_{\mu\nu} = L_F F_{\mu\nu}$ , it is straightforward to see that  $H$  is a function of  $P = \frac{1}{4} P_{\mu\nu} P^{\mu\nu}$  so that we can write [for all the details, see Ref. [95]]

$$L = 2P H_P - H, \quad (120)$$

where  $H_P$  denotes the derivative of  $H$  with respect to its argument  $P$ . In terms of  $H$  the energy-momentum tensor reads

$$\theta_{\mu\nu} = -H_P P_{\mu\alpha} P^{\alpha\nu} - g_{\mu\nu} (2P H_P - H). \quad (121)$$

Assuming  $P_{\mu\nu} = (\delta_\mu^0 \delta_\nu^1 - \delta_\mu^1 \delta_\nu^0) D(r)$ , where  $D(r)$  is the dual electric field, the Einstein equations in terms of  $H$  reveal that

$$H = \frac{1}{4\pi r^2} \frac{d\tilde{m}}{dr}. \quad (122)$$

Since  $P = L_F^2 F = -\frac{1}{2} L_F^2 E^2$ , we obtain

$$P = -\frac{Q^2}{32\pi^2 r^4}. \quad (123)$$

Finally, using Eqs. (117), (122) and (123) in Eq. (120), we obtain the Lagrangian

$$L(P) = -4\pi P - \frac{\alpha K \sqrt[4]{-2P} e^{\mathcal{G}(P)}}{4\sqrt{\pi} \sqrt{Q} r_H}, \quad (124)$$

with

$$\mathcal{G}(P) = -\frac{K \sqrt{Q}}{2\sqrt{\pi} \sqrt[4]{-2P} r_H}. \quad (125)$$

We want to conclude emphasizing once again that  $Q^2 \sim \alpha$ , so that the Schwarzschild geometry is always recovered for  $\alpha \rightarrow 0$ . However, when  $Q$  represents an electric charge, we can say that the electro-vacuum (Reissner-Nordström geometry) is filled with a tensor-vacuum whose origin lies in the nonlinear electrodynamics with the Lagrangian (124).

#### IV. CONCLUSIONS

Using the EGD approach, we studied the emergence of hairy black holes due to matter surrounding the central source of the Schwarzschild metric. Demanding that the solution always admits a well defined horizon through Eq. (31), and that the hair satisfies the SEC or DEC through Eqs. (48) or (76) and (77), respectively, we found two new families of hairy black holes displayed in Eqs. (69) and (87). These geometries were analysed and, in particular, we found that the solutions satisfying the DEC contain the non-trivial extension of the Reissner-Nordström black hole shown in (104) which possesses a simple event horizon. The Lagrangian of the nonlinear electrodynamics, which sources this solution, was also obtained explicitly.

From the technical point of view, since aim of the present work was to study some general conditions under which hair can be added to the (spherically symmetric) vacuum black holes of general relativity, the EGD was the natural approach to employ from the onset. In fact, the EGD is precisely devised for describing deformations of known solutions of general relativity induced by adding extra sources. Moreover, the properties of the

added source were restricted in order to enforce the conditions mentioned above, rather than assuming the hair is described in terms of fundamental fields. Nonetheless, we were able to give a description in terms of a nonlinear electrodynamics at least in one specific case.

Finally, we would like to remark that the charges  $Q$  and  $\ell_0 = \alpha \ell$  associated with our hairy black holes admit simple physical interpretations. The charge  $Q$  can be viewed as an effective electric charge and is proportional to  $\alpha$ , which is the generic parameter measuring the deviation from the chosen vacuum solution (which is given by the Schwarzschild metric). A special mention deserves the parameter  $\ell_0$ , which is associated with gauge trans-

formations of the seed Schwarzschild metric and which seems to always push the event horizon to radii larger than the Schwarzschild radius. Therefore  $\ell_0 = \alpha \ell$  measures how much the entropy of the black hole increases from its minimum Schwarzschild value  $S = 4 \pi M^2$  when hair is added.

### Acknowledgments

R.C. is partially supported by the INFN grant FLAG and his work has also been carried out in the framework of activities of the National Group of Mathematical Physics (GNFM, INDAM) and COST action *Cantata*.

- 
- [1] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **116**, 061102 (2016), [arXiv:1602.03837 \[gr-qc\]](#).
  - [2] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **119**, 141101 (2017), [arXiv:1709.09660 \[gr-qc\]](#).
  - [3] K. Akiyama *et al.* (Event Horizon Telescope), *Astrophys. J.* **875**, L1 (2019).
  - [4] R. Ruffini and J. A. Wheeler, *Phys. Today* **24**, 30 (1971).
  - [5] S. Hawking, *Commun. Math. Phys.* **25**, 152 (1972).
  - [6] S. W. Hawking, M. J. Perry, and A. Strominger, *Phys. Rev. Lett.* **116**, 231301 (2016), [arXiv:1601.00921 \[hep-th\]](#).
  - [7] T. P. Sotiriou and V. Faraoni, *Phys. Rev. Lett.* **108**, 081103 (2012), [arXiv:1109.6324 \[gr-qc\]](#).
  - [8] E. Babichev and C. Charmousis, *JHEP* **08**, 106 (2014), [arXiv:1312.3204 \[gr-qc\]](#).
  - [9] A. Cisterna and C. Ericces, *Phys. Rev. D* **89**, 084038 (2014), [arXiv:1401.4479 \[gr-qc\]](#).
  - [10] T. P. Sotiriou and S.-Y. Zhou, *Phys. Rev. Lett.* **112**, 251102 (2014), [arXiv:1312.3622 \[gr-qc\]](#).
  - [11] G. Antoniou, A. Bakopoulos, and P. Kanti, *Phys. Rev. Lett.* **120**, 131102 (2018), [arXiv:1711.03390 \[hep-th\]](#).
  - [12] G. Antoniou, A. Bakopoulos, and P. Kanti, *Phys. Rev. D* **97**, 084037 (2018), [arXiv:1711.07431 \[hep-th\]](#).
  - [13] D. Grumiller, A. Pérez, M. Sheikh-Jabbari, R. Troncoso, and C. Zwickel, *Phys. Rev. Lett.* **124**, 041601 (2020), [arXiv:1908.09833 \[hep-th\]](#).
  - [14] M. Volkov and D. Galtsov, *JETP Lett.* **50**, 346 (1989).
  - [15] P. Kanti, N. Mavromatos, J. Rizos, K. Tamvakis, and E. Winstanley, *Phys. Rev. D* **54**, 5049 (1996), [arXiv:hep-th/9511071](#).
  - [16] P. Kanti, N. Mavromatos, J. Rizos, K. Tamvakis, and E. Winstanley, *Phys. Rev. D* **57**, 6255 (1998), [arXiv:hep-th/9703192](#).
  - [17] K. G. Zloshchastiev, *Phys. Rev. Lett.* **94**, 121101 (2005), [arXiv:hep-th/0408163](#).
  - [18] C. Martinez, R. Troncoso, and J. Zanelli, *Phys. Rev. D* **70**, 084035 (2004), [arXiv:hep-th/0406111](#).
  - [19] C. A. Herdeiro and E. Radu, *Int. J. Mod. Phys. D* **24**, 1542014 (2015), [arXiv:1504.08209 \[gr-qc\]](#).
  - [20] T. P. Sotiriou, *Class. Quant. Grav.* **32**, 214002 (2015), [arXiv:1505.00248 \[gr-qc\]](#).
  - [21] J. Ovalle, R. Casadio, R. da Rocha, A. Sotomayor, and Z. Stuchlik, *Eur. Phys. J.* **C78**, 960 (2018), [arXiv:1804.03468 \[gr-qc\]](#).
  - [22] J. Ovalle, *Phys. Lett.* **B788**, 213 (2019), [arXiv:1812.03000 \[gr-qc\]](#).
  - [23] J. Ovalle, *Phys. Rev.* **D95**, 104019 (2017), [arXiv:1704.05899 \[gr-qc\]](#).
  - [24] J. Ovalle, *Mod. Phys. Lett.* **A23**, 3247 (2008), [arXiv:gr-qc/0703095 \[gr-qc\]](#).
  - [25] J. Ovalle and R. Casadio, *Beyond Einstein Gravity*, SpringerBriefs in Physics (Springer Nature, Cham, 2020).
  - [26] J. Ovalle, *Int. J. Mod. Phys. D* **18**, 837 (2009), [arXiv:0809.3547 \[gr-qc\]](#).
  - [27] J. Ovalle, *Mod. Phys. Lett. A* **25**, 3323 (2010), [arXiv:1009.3674 \[gr-qc\]](#).
  - [28] R. Casadio and J. Ovalle, *Phys. Lett.* **B715**, 251 (2012), [arXiv:1201.6145 \[gr-qc\]](#).
  - [29] R. Casadio and J. Ovalle, *Gen. Rel. Grav.* **46**, 1669 (2014), [arXiv:1212.0409 \[gr-qc\]](#).
  - [30] J. Ovalle, F. Linares, A. Pasqua, and A. Sotomayor, *Class. Quant. Grav.* **30**, 175019 (2013), [arXiv:1304.5995 \[gr-qc\]](#).
  - [31] R. Casadio, J. Ovalle, and R. da Rocha, *Class. Quant. Grav.* **31**, 045016 (2014), [arXiv:1310.5853 \[gr-qc\]](#).
  - [32] J. Ovalle and F. Linares, *Phys. Rev.* **D88**, 104026 (2013), [arXiv:1311.1844 \[gr-qc\]](#).
  - [33] J. Ovalle, L. A. Gergely, and R. Casadio, *Class. Quant. Grav.* **32**, 045015 (2015), [arXiv:1405.0252 \[gr-qc\]](#).
  - [34] R. Casadio, J. Ovalle, and R. da Rocha, *EPL* **110**, 40003 (2015), [arXiv:1503.02316 \[gr-qc\]](#).
  - [35] R. Casadio, J. Ovalle, and R. da Rocha, *Class. Quant. Grav.* **32**, 215020 (2015), [arXiv:1503.02873 \[gr-qc\]](#).
  - [36] J. Ovalle, *Proceedings, 9th Alexander Friedmann International Seminar on Gravitation and Cosmology and 3rd Satellite Symposium on the Casimir Effect: St. Petersburg, Russia, June 21-27, 2015*, (2015), [10.1142/S2010194516601320](#), [*Int. J. Mod. Phys. Conf. Ser.*41,1660132(2016)], [arXiv:1510.00855 \[gr-qc\]](#).
  - [37] R. T. Cavalcanti, A. G. da Silva, and R. da Rocha, *Class. Quant. Grav.* **33**, 215007 (2016), [arXiv:1605.01271 \[gr-qc\]](#).
  - [38] R. da Rocha, *Phys. Rev.* **D95**, 124017 (2017), [arXiv:1701.00761 \[hep-ph\]](#).
  - [39] R. da Rocha, *Eur. Phys. J.* **C77**, 355 (2017), [arXiv:1703.01528 \[hep-th\]](#).



- [40] A. Fernandes-Silva and R. da Rocha, *Eur. Phys. J. C* **78**, 271 (2018), arXiv:1708.08686 [hep-th].
- [41] R. Casadio, P. Nicolini, and R. da Rocha, *Class. Quant. Grav.* **35**, 185001 (2018), arXiv:1709.09704 [hep-th].
- [42] A. Fernandes-Silva, A. J. Ferreira-Martins, and R. Da Rocha, *Eur. Phys. J. C* **78**, 631 (2018), arXiv:1803.03336 [hep-th].
- [43] E. Contreras and P. Bargueño, *Eur. Phys. J. C* **78**, 558 (2018), arXiv:1805.10565 [gr-qc].
- [44] E. Contreras, *Eur. Phys. J. C* **78**, 678 (2018), arXiv:1807.03252 [gr-qc].
- [45] E. Contreras and P. Bargueño, *Eur. Phys. J. C* **78**, 985 (2018), arXiv:1809.09820 [gr-qc].
- [46] G. Panotopoulos and A. Rincón, *Eur. Phys. J. C* **78**, 851 (2018), arXiv:1810.08830 [gr-qc].
- [47] R. Da Rocha and A. A. Tomaz, *Eur. Phys. J. C* **79**, 1035 (2019), arXiv:1905.01548 [hep-th].
- [48] C. Las Heras and P. León, *Eur. Phys. J. C* **79**, 990 (2019), arXiv:1905.02380 [gr-qc].
- [49] A. Rincón, L. Gabbanelli, E. Contreras, and F. Tello-Ortiz, *Eur. Phys. J. C* **79**, 873 (2019), arXiv:1909.00500 [gr-qc].
- [50] R. da Rocha, *Symmetry* **12**, 508 (2020), arXiv:2002.10972 [hep-th].
- [51] E. Contreras, F. Tello-Ortiz, and S. Maurya, (2020), arXiv:2002.12444 [gr-qc].
- [52] C. Arias, F. Tello Ortiz, and E. Contreras, *Eur. Phys. J. C* **80**, 463 (2020), arXiv:2003.00256 [gr-qc].
- [53] R. da Rocha, *Phys. Rev. D* **102**, 024011 (2020), arXiv:2003.12852 [hep-th].
- [54] F. Tello-Ortiz, S. Maurya, and Y. Gomez-Leyton, *Eur. Phys. J. C* **80**, 324 (2020).
- [55] R. da Rocha and A. A. Tomaz, (2020), arXiv:2005.02980 [hep-th].
- [56] P. Meert and R. da Rocha, (2020), arXiv:2006.02564 [gr-qc].
- [57] V. Kiselev, *Class. Quant. Grav.* **20**, 1187 (2003), arXiv:gr-qc/0210040.
- [58] J. Ovalle, R. Casadio, R. da Rocha, and A. Sotomayor, *Eur. Phys. J. C* **78**, 122 (2018), arXiv:1708.00407 [gr-qc].
- [59] L. Gabbanelli, A. Rincón, and C. Rubio, *Eur. Phys. J. C* **78**, 370 (2018), arXiv:1802.08000 [gr-qc].
- [60] C. L. Heras and P. Leon, *Fortsch. Phys.* **66**, 1800036 (2018), arXiv:1804.06874 [gr-qc].
- [61] M. Estrada and F. Tello-Ortiz, *Eur. Phys. J. Plus* **133**, 453 (2018), arXiv:1803.02344 [gr-qc].
- [62] M. Sharif and S. Sadiq, *Eur. Phys. J. C* **78**, 410 (2018), arXiv:1804.09616 [gr-qc].
- [63] E. Morales and F. Tello-Ortiz, *Eur. Phys. J. C* **78**, 618 (2018), arXiv:1805.00592 [gr-qc].
- [64] M. Sharif and S. Sadiq, *Eur. Phys. J. Plus* **133**, 245 (2018).
- [65] E. Morales and F. Tello-Ortiz, *Eur. Phys. J. C* **78**, 841 (2018), arXiv:1808.01699 [gr-qc].
- [66] M. Estrada and R. Prado, *Eur. Phys. J. Plus* **134**, 168 (2019), arXiv:1809.03591 [gr-qc].
- [67] M. Sharif and S. Saba, *Eur. Phys. J. C* **78**, 921 (2018), arXiv:1811.08112 [gr-qc].
- [68] J. Ovalle, R. Casadio, R. da Rocha, A. Sotomayor, and Z. Stuchlik, *EPL* **124**, 20004 (2018), arXiv:1811.08559 [gr-qc].
- [69] E. Contreras, *Class. Quant. Grav.* **36**, 095004 (2019), arXiv:1901.00231 [gr-qc].
- [70] S. K. Maurya and F. Tello-Ortiz, *Eur. Phys. J. C* **79**, 85 (2019).
- [71] E. Contreras, A. Rincón, and P. Bargueño, *Eur. Phys. J. C* **79**, 216 (2019), arXiv:1902.02033 [gr-qc].
- [72] E. Contreras and P. Bargueño, *Class. Quant. Grav.* **36**, 215009 (2019), arXiv:1902.09495 [gr-qc].
- [73] L. Gabbanelli, J. Ovalle, A. Sotomayor, Z. Stuchlik, and R. Casadio, *Eur. Phys. J. C* **79**, 486 (2019), arXiv:1905.10162 [gr-qc].
- [74] M. Estrada, *Eur. Phys. J. C* **79**, 918 (2019), arXiv:1905.12129 [gr-qc].
- [75] J. Ovalle, C. Posada, and Z. Stuchlik, *Class. Quant. Grav.* **36**, 205010 (2019), arXiv:1905.12452 [gr-qc].
- [76] S. Maurya and F. Tello-Ortiz, *Phys. Dark Univ.* **27**, 100442 (2020), arXiv:1905.13519 [gr-qc].
- [77] S. Hensh and Z. e. Stuchlik, *Eur. Phys. J. C* **79**, 834 (2019), arXiv:1906.08368 [gr-qc].
- [78] F. X. Linares Cedeño and E. Contreras, *Phys. Dark Univ.* **28**, 100543 (2020), arXiv:1907.04892 [gr-qc].
- [79] P. León and A. Sotomayor, *Fortsch. Phys.* **67**, 1900077 (2019), arXiv:1907.11763 [gr-qc].
- [80] V. Torres-Sánchez and E. Contreras, *Eur. Phys. J. C* **79**, 829 (2019), arXiv:1908.08194 [gr-qc].
- [81] R. Casadio, E. Contreras, J. Ovalle, A. Sotomayor, and Z. Stuchlick, *Eur. Phys. J. C* **79**, 826 (2019), arXiv:1909.01902 [gr-qc].
- [82] K. Singh, S. Maurya, M. Jasim, and F. Rahaman, *Eur. Phys. J. C* **79**, 851 (2019).
- [83] S. Maurya, *Eur. Phys. J. C* **79**, 958 (2019).
- [84] M. Sharif and A. Waseem, *Annals Phys.* **405**, 14 (2019).
- [85] G. Abellán, V. Torres-Sánchez, E. Fuenmayor, and E. Contreras, *Eur. Phys. J. C* **80**, 177 (2020), arXiv:2001.08573 [gr-qc].
- [86] M. Sharif and Q. Ama-Tul-Mughani, *Annals Phys.* **415**, 168122 (2020), arXiv:2004.07925 [gr-qc].
- [87] F. Tello-Ortiz, *Eur. Phys. J. C* **80**, 413 (2020).
- [88] S. Maurya, *Eur. Phys. J. C* **80**, 429 (2020).
- [89] A. Rincón, E. Contreras, F. Tello-Ortiz, P. Bargueño, and G. Abellán, *Eur. Phys. J. C* **80**, 490 (2020), arXiv:2005.10991 [gr-qc].
- [90] M. Sharif and A. Majid, *Phys. Dark Univ.* **30**, 100610 (2020), arXiv:2006.04578 [gr-qc].
- [91] S. Maurya, K. N. Singh, and B. Dayanandan, *Eur. Phys. J. C* **80**, 448 (2020).
- [92] M. Zubair and H. Azmat, *Annals Phys.* **420**, 168248 (2020), arXiv:2005.06955 [physics.gen-ph].
- [93] L. Herrera and N. O. Santos, *Phys. Rept.* **286**, 53 (1997).
- [94] M. K. Mak and T. Harko, *Proc. Roy. Soc. Lond.* **A459**, 393 (2003), arXiv:gr-qc/0110103 [gr-qc].
- [95] I. Salazar, A. Garcia, and J. Plebanski, *J. Math. Phys.* **28**, 2171 (1987).
- [96] E. Ayon-Beato and A. Garcia, *Phys. Rev. Lett.* **80**, 5056 (1998), arXiv:gr-qc/9911046.
- [97] E. Guendelman and A. Rabinowitz, *Gen. Rel. Grav.* **28**, 117 (1996).
- [98] T. Jacobson, *Class. Quant. Grav.* **24**, 5717 (2007), arXiv:0707.3222 [gr-qc].
- [99] I. G. Dymnikova, *Phys. Lett. B* **472**, 33 (2000), arXiv:gr-qc/9912116.
- [100] I. Dymnikova, *Int. J. Mod. Phys. D* **12**, 1015 (2003), arXiv:gr-qc/0304110.
- [101] M. Visser, *Class. Quant. Grav.* **37**, 045001 (2020), arXiv:1908.11058 [gr-qc].



- [102] M. Visser, *Lorentzian wormholes: From Einstein to Hawking* (1995).
- [103] E. Curiel, “A Primer on Energy Conditions,” (2017) pp. 43–104, [arXiv:1405.0403 \[physics.hist-ph\]](#).