

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Assessing the effect of dietary inulin supplementation on gastrointestinal fermentation, digestibility and growth in pigs: A meta-analysis

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Assessing the effect of dietary inulin supplementation on gastrointestinal fermentation, digestibility and growth in pigs: A meta-analysis / Metzler-Zebeli, B.U.*; Trevisi, P.; Prates, J.A.M.; Tanghe, S.; Bosi, P.; Canibe, N.; Montagne, L.; Freire, J.; Zebeli, Q. - In: ANIMAL FEED SCIENCE AND TECHNOLOGY. - ISSN 0377-8401. - STAMPA. - 233:(2017), pp. 120-132. [10.1016/j.anifeedsci.2017.05.010]

Availability:

This version is available at: https://hdl.handle.net/11585/623424 since: 2018-02-21

Published:

DOI: http://doi.org/10.1016/j.anifeedsci.2017.05.010

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

Manuscript Details

Manuscript number	ANIFEE_2016_564	
Title	Assessing the effect of dietary inulin supplementation on gastrointestinal fermentation, digestibility and growth in pigs: a meta-analysis	
Article type	Research Paper	

Abstract

Inulin has been reported to improve the homeostasis in the gastrointestinal tract (GIT) of pigs by stimulating modulating the intestinal microbiota and fermentation. The aim of this study was to quantify the relationship between dietary inulin and microbial response variables in digesta from the GIT and feces of weaned, growing and finishing pigs using a meta-analytical approach. We further examined the effect of dietary inulin on the coefficients of ileal (CIAD) and total tract apparent digestibility (CTTAD) of nutrients and ADG. Pig's starting body weight was considered the main inclusion criterion. Missing information about explanatory variables and few values available for response variables reduced the number of studies included. From the 33 included articles published between 2000 and 2016, individual sub-datasets for fermentation metabolites, bacterial abundances, CIAD, CTTAD and performance were built. Prediction models on the effect on inulin were computed accounting for inter- and intra-study variability. Dietary inulin levels ranged from 0.1 to 25.8 %, whereby the median and mean inulin levels were 0.1 to 2% and 3 to 4 %, respectively. Few of the investigated fermentation response variables were influenced by dietary inulin. Strong negative relationships were found between dietary inulin and gastric pH in weaned pigs (R2 = 0.81; P < 0.001; n = 12), colonic enterobacteria (R2 = 0.50; P < 0.001; n = 19) and fecal lactobacilli (R2 = 0.41; P < 0.001; n = 26) throughout all production phases, whereas observed negative relationships between inulin and colonic bifidobacteria and fecal enterobacteria and Escherichia coli were of minor physiological relevance (P < 0.05). Moreover, increasing inulin levels negatively correlated with the CTTAD of crude protein (R2 = 0.83; P < 0.001; n = 15), but they did not influence average daily gain of pigs. Best-fit models indicated that dietary crude protein amplified the effect of inulin on CTTAD of crude protein and gastric pH, but counteracted the inulin effect on fecal E. coli (P < 0.05). Accordingly, both pig's body weight and inulin decreased gastric pH and fecal lactobacilli but counteracted the inulin effect on colonic bifidobacteria and fecal E. coli (P < 0.05). In conclusion, this study supported that dietary inulin can stimulate gastric acid secretion which may be favorable GIT health in weaned pigs. However, meta-regressions did not support that inulin promotes the bacterial groups previously associated with porcine GIT health, such as lactobacilli and bifidobacteria.

Keywords	Inulin; gastro-intestinal tract; Fermentation; microbiota; meta-Analysis; pig
Taxonomy	Domestic Animals, Non-Ruminant Nutrition, Animal Dietary Supplement
Manuscript category	Non-ruminant
Corresponding Author	Barbara Metzler-Zebeli
Order of Authors	Barbara Metzler-Zebeli, Paolo Trevisi, Jose Prates, Sofie Tanghe, Paolo Bosi, Nuria Canibe, Lucile Montagne, Joao Freire, Qendrim Zebeli
Suggested reviewers	Jos Houdijk

Submission Files Included in this PDF

File Name [File Type]

Cover letter_ANIFEE_2016_564_R1.docx [Cover Letter]

Inulin Meta_Declaration of interest.docx [Conflict of Interest]

ANIFEE_2016_564_Author reply.docx [Response to Reviewers]

ANIFEE_2016_564_revision_1.pdf.docx [Manuscript File]

Inulin Meta_Highlights.docx [Highlights]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.



Dr. Barbara Metzler-Zebeli University Clinic for Swine

Department for Farm Animals and Veterinary Public Health University of Veterinary Medicine Vienna Veterinaerplatz 1, 1210 Vienna, Austria Email: barbara.metzler@vetmeduni.ac.at Tel: +43 1 25077-3209; Fax: +43 1 25077-3290

Vienna, 23rd of March, 2017

Re: Submission of revised manuscript ANIFEE_2016_564

Dear Dr. de Blas,

We are grateful for the opportunity to revise our manuscript ANIFEE_2016_564 entitled "Assessing the effect of dietary inulin supplementation on gastrointestinal fermentation, digestibility and growth in pigs: a meta-analysis".

All reviewer comments were addressed point-by-point in the author reply and line numbers in the revised manuscript were provided where changes can be found. Changes made in the manuscript text are highlighted in yellow. We hope that we could improve our manuscript to meet the high standards of Animal Feed Science and Technology.

Sincerely yours,

Barbara Metzler-Zebeli

Declaration of interests

•

The authors declare no conflict of interest. Moreover, all authors listed have contributed to the work, all authors have read and approved the final manuscript submitted to Animal Feed Science and Technology. No part of the work has been published before and is not under consideration for publication elsewhere.

ANIFEE_2016_564

AUTHORS: Dear Dr. de Blas,

We are grateful for the opportunity to revise our manuscript. All reviewer comments were addressed point-by-point in the author reply and line numbers in the revised manuscript were provided where changes can be found. Changes made in the manuscript text are highlighted in yellow. We hope that we could improve our manuscript to meet the high standards of Animal Feed Science and Technology.

Sincerely yours, Barbara Metzler-Zebeli

<u>Reviewer 1:</u> <u>General comments:</u> This is an interesting paper, comprehensively statistically analysed, examining the trans-study effects of added inulin in diets for pigs on a number of different variables.

AUTHORS: We would like to thank the reviewer for the helpful comments.

Major comments are as follows:

The main weakness of the paper is the lack of statistical accountability reflecting endogenous fructans levels in the diets used. Whilst the authors' duly acknowledge this (L353-355), strong doubt remains concerning the veracity of the data derived without taking endogenous fructans levels into account. The authors have made other assumptions (L181-183) in the paper; hence, the authors should calculate (based on 'best-bet' book or feed matrix values) endogenous fructans levels in the diets used in the studies analysed, and redo the analyses.

AUTHORS: Thank you for this comment. The strength of a meta-analysis is that it takes into account effects within treatments of a study and between studies. Each study had a 'control treatment' and the 'supplemented inulin treatment'. Our meta-analytical approach weighted the treatment to the corresponding control within the individual studies. So, the fructan content of the basal diet did not matter as each treatment effect was weighted to the respective control. In this way, the 'native' fructan content of the diet was balanced as it is clear that the native fructan content of the basal diet in the different studies varied. We modified the sentence in the Discussion to clarify that we could not distinguish the effect of the supplemented inulin from that of the naturally occurring fructan levels in the basal diet due to the fact that insufficient information was provided in the original studies (New Line 350-354)

As alluded to below (L119), the authors only searched for 'microbiota' whereas the terms flora and microbiome are often used interchangeably with microbiota. The authors need to ensure that no publications were omitted because of this.

AUTHORS: The literature was searched using other terms for "microbiota" such as microflora, and microbiome. In addition, literature was searched again using only the

search terms inulin and pig. According to this search, our original datasets comprised all relevant papers. This additional search was amended in the Materials and Methods section 2.1 Literature search (New Line 120).

As alluded to below (L151), the authors don't provide sound justification for using dietary crude protein content as a major criterion.

AUTHORS: A justification for using dietary crude protein content as a major criterion was provided (New Line 151-153). The main reason was that studies used different protein levels despite similar production stages and starting BW. Since the dietary protein content can influence the intestinal fermentation, digestibility and growth performance, it was one of the few prediction variables that were available for the studies included.

Specific comments:

1. L32-L33: '..appears to improve..', and then '...inconsistent results..'. These two parts of the sentence are in congruence, i.e., one cannot say that inulin improves but then that improvement is inconsistent. This needs rewriting.

AUTHORS: Thank you for pointing this out. The sentence was modified (New Line 32-33).

2. L32, and throughout paper: the authors need to use a consistent terminology for gastrointestinal tract/gut/intestinal etc. There are numerous 'versions' used in the paper. Suggest use gastrointestinal tract (GIT), abbreviating to GIT thereafter, unless there's more specific detail provided.

AUTHORS: Gastrointestinal tract (GIT) was used as consistent term throughout the revised manuscript.

3. L67, L86 etc.: what is meant by 'beneficial'? This is a term used loosely in the scientific literature, and is generally unhelpful and sometimes meaningless. The authors are strongly encouraged to define this more precisely, or use different language altogether.

AUTHORS: The term 'beneficial bacteria' has been removed.

4. L84: isn't it the fructans that are measured, rather than inulin?

AUTHORS: Indeed fructans are measured. However, in the studies used in this metaanalysis, the inulin level was reported, but in most cases not the fructan content or chain length of the inulin used. This is why we used 'inulin level' and not 'fructan level'.

5. L119: did the authors also use the terms flora and microbiome? These terms are often used interchangeably with microbiota. I think this is a key issue, and one that may require the authors to research the literature.

AUTHORS: Please see our comment above. We double-checked the literature with other search terms for microbiota and bacteria to ensure that we did not miss an original article.

6. L146, L159 etc., and throughout: the authors (in L131) define CIAD and CTTAD, but then use nutrient digestibility. Moreover, DM is not a nutrient. The authors should use consistent terminology throughput the paper.

AUTHORS: Consistent terminology using CIAD and CTTAD was used throughout the manuscript. CIAD and CTTAD of DM was mentioned in addition to those of nutrients.

7. L151: why was dietary crude protein content chosen? This seemingly is a major determinant of the outcomes and conclusions (e.g., L459) yet the reader is provided with little reasoning for its inclusion.

AUTHORS: Please see our reply under "major comments".

8. L176: what serotype(s) of *E. coli*? Please define.

AUTHORS: In most studies used in this meat-analysis, culturing was done using selective media for the enumeration of *E. coli* but no further culturing on strain/serotype-selective media was performed.

9. L181-183: evidence must be provided by the authors' to substantiate the assumption that CFUs and gene copy numbers 'approximately correspond to each other'.

AUTHORS: Evidence was provided (New Line 182-184).

10. L204: write as, ' MEANS procedure of..'

AUTHORS: Done as suggested (New Line 204-205).

11. L238-239: further to previous comments; I am unconvinced that just because dietary crude protein content was available for all response variables, then it should be used. Where is the hypothesis justifying its inclusion? A physiological basis for its inclusion needs to be provided.

AUTHORS: Please see our reply under "major comments". The dietary CP level varied within production phases despite similar starting BW. As the CP level can influence all of the dependent variables investigated in the present meta-analysis, the dietary CP level was considered.

12. L353-355: see General comments.

13. L429: pathogenic serotypes?

AUTHORS: Please see our comment above. *Escherichia coli* was enumerated on selective medium but no further differentiation using selective media for the various

serotypes was done in the studies used in this meta-analysis. Also, only 'total *E. coli* counts were used in this meta-analysis if specific serotypes were determined.

14. Table 1: in line with previous comments, the (base) diet formulation(s) may have a profound impact on the variables' responses. Perhaps total fructans levels should be used rather than added fructans levels?

AUTHORS: It would have been great if the fructan contents were reported in the studies used in this meta-analysis. As this was not the case for many studies, we focused on the supplemented inulin level. Please see also our reply under the "major comments" as the meta-analysis accounted for variation in the basal fructan content of the diets by taking into account the control treatment and the supplemented inulin content when estimating the relationship between a dependent variable and the inulin level.

Reviewer 2

The authors report a meta-analysis designed to establish the benefits of inulin supplementation for pigs on a number of performance and fermentation characteristics. The study is well executed, including the critical approach of paper quality to be included, and the resulting overall insight into inulin benefits is relevant for publication. However, a number of queries have arisen during my review, which I list below for the authors to consider.

AUTHORS: Thank you for the helpful comments to improve the quality of our manuscript.

General

The text contains a number of grammar inconsistencies arising from translation challenges. I consider it not to be the task of a reviewer to list these but encourage the authors to have the manuscript read by native English speakers, with experience in publication in this area of research.

AUTHORS: Thank you for pointing out grammar inconsistencies. The manuscript was checked by a native American English speaker and we hope that this person did not oversee typos and grammar issues.

Major comments

2. L218-242. Two approaches were used to assess impact of start BW, diet CP and inulin level, i.e. by inclusion as random effects and through backward elimination. It is not clear what the benefit is of doing both, especially because outcomes seem to vary. Would the conclusions be any less if only one of them is chosen and presented?

AUTHORS: The rationale behind assessing the impact of the start BW was to consider maturational changes from weaned to finisher pigs. Because studies used different crude protein (CP) levels within the same production stage and for similar starting BW, the dietary CP content was included as another predictor variable as the dietary CP content can influence the intestinal fermentation, digestibility and growth performance. Taking both effects into account in the modeling leads to a more powerful estimation of the response variables. We used the VIF in order to avoid potential multi-collinarity among predictor variables, start BW, dietary CP, dietary inulin level and quadratic inulin level.

Also, the backward elimination analysis demonstrated quadratic relationships between inulin level and a number of read outs. What is the biological meaning of this quadratic relation ((L312, 318)?

AUTHORS: The biological meaning of a quadratic relation is that there is a maximum supplementation level above which no further increase or decrease in the response variable can be expected (asymptotic approximation).

4. Where significant relations were found between level of inulin and read outs, the prediction of the read out at 3% inulin was presented. What was the rational of choosing 3%, given that the vast majority of studies as you indicated had levels of inulin between 0.1 and 2%?

AUTHORS: It is true that median levels of inulin ranged between 0.1 and 2 %. The mean (which was about 3 to 4 % inulin) rather than median supplementation level was used for the prediction as very low levels of inulin may be insufficient to produce an effect.

6. Related to the above, where a significant slope was observed, it would be useful to speculate whether the magnitude of effect is also biologically relevant. For example, would a reduction of 0.19 log units in faecal E. coli be of relevance?

AUTHORS: It was indicated in the Discussion if the magnitude was of significance relevance (New Line 380 and 429-433).

8. You state that "an insufficient level of inulin supplementation to modulate gastrointestinal fermentation and bacterial abundances was probably a crucial factor determining the results obtained". Whilst this is not disputed, could your data indicate what level would be sufficient to observe responses?

AUTHORS: A sentence was added to discuss at which inulin level responses could be expected (New Line 346-350).

10. Inulin consists of variable numbers of fructose units, as clearly indicated. This would also include the shorter chain fructo-oligosaccharides. Some of the outcomes of the effects of inulin reported and discussed here accord with effects of fructo-oligosaccharides, including effects on gastric pH and ileal lactic acid. It would be advisable to bring this up during the discussion.

AUTHORS: Thank you for this comment. Indeed, there are some consistencies when comparing the present relationships for inulin and reports for short-chain fructooligosaccharides. However, as the main focus of the present meta-analysis is the longchain fructan inulin, we prefer mainly comparing the present results with those previously reported for inulin in individual studies.

12. The conclusion that inulin is more effective in younger pigs compared to older ones (L365) is acceptable but it would be appropriate for the authors to speculate why this would be.

AUTHORS: These sentences were modified and some reasoning was provided why younger pigs may benefit more from inulin feeding than older pigs (New Line 355-359).

14. L440. The authors state here that no relation between dietary inulin and ADG could be established. Is it not better to state that the data support the view that "ADG is not sensitive to dietary inulin levels".

AUTHORS: Modified as suggested (New Line 437-438).

Minor comments

Title

L1. The order in which the results are reported are not in line with the order of the parameters in the title. As a consequence, based on the title one would expect greater emphasis on performance, then digestibility and lastly fermentation characteristics. I would suggest to amend the title to reflect this, moreover because the outcome is that inulin did not really affect growth performance in the first place.

AUTHORS: The title was modified accordingly.

Abstract

L32. Please amend abstract as per suggestions from main text where needed.

AUTHORS: The abstract was amended after revising the main text.

Introduction

L62. It is of interest to note that the ban on AGP in Europe is supported by references from Canada and Western Australia. Can a similar reference supporting this position be added coming from an EU-led review?

AUTHORS: References were replaced by references from European authors (New Line 65).

L89. Please include relevant references after "weaned pigs"; the way how this sentence is constructed requests it to be referenced.

AUTHORS: Relevant references were added (New Line 88-89).

Materials and methods

L133. It might be useful to consider how the inulin was included in the test diets, i.e. whether it was exchanged against some ingredients, or diluted a basal diet, or otherwise. This may add some insight into its variation in response.

AUTHORS: The dietary inulin most often replaced one of the main energy feedstuffs, but did not dilute the diets.

L156. A comment or observation rather than anything else but I was rather surprised to learn that 25% of articles had to be excluded as they would not report initial body weight. Could you have considered using these studies for the analyses that did not rely on BW?

AUTHORS: All dependent variables investigated in the present meta-analysis depended either on the age or the BW of the pigs. Therefore, it was obligatory that this information was provided.

L182. What do you mean with "approximately correspond"?

AUTHORS: This sentence was modified for clarification (New Line 182-184)

L198. Is it not better to say that Breed and Sex were not included in the model due to inconsistency of reporting, rather than assuming responses were not affected by Breed and Sex?

AUTHORS: This was modified accordingly (New Line 200-201).

L218. Predictor variables tested were study ? Something missing here.

AUTHORS: This was corrected and the other predictor variables tested were added (New Line 220).

Discussion

L336. Please split the list of authors into those relevant for the small intestine statement and those relevant for the large intestine statement.

AUTHORS: As suggested, the list of authors was split into those relevant for the small intestine and those relevant for the large intestine (New Line 333-335).

L356. I think the first line is not needed, as it destracts from what you want to say; start with "It was".

AUTHORS: The beginning of this sentence has been modified (New Line 360).

L388. This line is not clear. What does "effects of BW and diet CP in conjunction with inulin" mean?

AUTHORS: This sentence has been modified for clarification (New Line 383)

L396. Here you seem to underpin the outcome of the meta-analyses with that of individual studies, which likely were part of the meta-analysis. Is that acceptable? Or were there other reasons to refer to these specific studies?

AUTHORS: It is acceptable to compare the outcome of the meta-analysis with the individual studies included.

L424. In Table 8 and L319, the relationship between faecal lactobacilli is negative. Please check for consistency.

AUTHORS: Thank you. The sentence was corrected (New Line 417).

L421. To what extent it the effect of "maturation" confounded by the effect of increased feed intake?

AUTHORS: The increased feed intake but also the changing dietary composition will have contributed to the maturation of the microbiota. Because the feed intake level was not provided in many studies, we prefer not to speculate about the impact of the feed intake.

L432. Replace "If" with "Whether".

AUTHORS: Changed as suggested (New Line 425).

L435. This is consistent with a large body of evidence from across the world that CP levels can modify faecal coli counts. Perhaps worth highlighting.

AUTHORS: Due the rather low biological relevance of the changes observed, this finding should not be overemphasized.

L441. Something is missing here ". between increasing dietary CIAD of DM......"; something seems missing after "increasing" and before "dietary", as dietary CIAD does not make sense?

AUTHORS: Thank you. "dietary" was deleted to correct the sentence (New Line 439-440).

1	Assessing the effect of dietary inulin supplementation on gastrointestinal
2	fermentation, digestibility and growth in pigs: a meta-analysis
3	
4	B. U. Metzler-Zebeli ^{a*} , P. Trevisi ^b , J. A. M. Prates ^c , S. Tanghe ^d , P. Bosi ^b , N. Canibe ^e , L.
5	Montagne ^f , J. Freire ^g , Q. Zebeli ^h
6	^a University Clinic for Swine, Department for Farm Animals and Veterinary Public Health,
7	Vetmeduni Vienna, Veterinärplatz 1, 1210 Vienna, Austria
8	^b Department of Agricultural and Food Science (DISTAL) - University of Bologna, 40127
9	Bologna, Italy
10	^c CIISA, Faculty of Veterinary Medicine, University of Lisbon, Avenida da Universidade Técnica,
11	Alto da Ajuda, 1300-477 Lisbon, Portugal
12	^d Nuscience Belgium, 9031 Ghent, Belgium
13	^e Department of Animal Science Aarhus University, 8830 Tjele, Denmark
14	^f PEGASE, Agrocampus Ouest, INRA, 35590, Saint-Gilles, France
15	^g LEAF, Instituto Superior de Agronomia, Universidade de Lisboa, Tapada da Ajuda, 1349-017 Lisboa,
16	Portugal
17	^h Institute of Animal Nutrition and Functional Plant Compounds, Department for Farm Animals
18	and Veterinary Public Health, Vetmeduni Vienna, Veterinaerplatz 1, 1210 Vienna, Austria
19	
20	Abbreviations: ADG, average daily gain; BW, body weight; CFU, colony forming units; CIAD,
21	coefficient of ileal apparent digestibility; CP, crude protein; CTTAD, coefficient of total tract
22	apparent digestibility; DM, dry matter; FISH, fluorescence-in-situ-hybridization; GIT,

gastrointestinal tract; NDF, neutral detergent fiber; RMSE, root mean square error; SAS, 23

24	statistical analysis system; SE, standard error; VIF, variance inflation factor; VFA, volatile fatty
25	acids.
26	
27	* Corresponding author. Tel.: (+43) 250 77-3209; Fax: (+43) 250 77-5260.
28	Email address: barbara.metzler@vetmeduni.ac.at
29	
30	

31 ABSTRACT (400 words)

32 Inulin has been reported to improve the homeostasis in the gastrointestinal tract (GIT) of pigs by stimulating modulating the intestinal microbiota and fermentation. The aim of this study was to 33 quantify the relationship between dietary inulin and microbial response variables in digesta from 34 the GIT and feces of weaned, growing and finishing pigs using a meta-analytical approach. We 35 further examined the effect of dietary inulin on the coefficients of ileal (CIAD) and total tract 36 apparent digestibility (CTTAD) of nutrients and ADG. Pig's starting body weight was 37 considered the main inclusion criterion. Missing information about explanatory variables and 38 39 few values available for response variables reduced the number of studies included. From the 33 40 included articles published between 2000 and 2016, individual sub-datasets for fermentation metabolites, bacterial abundances, CIAD, CTTAD and performance were built. Prediction 41 models on the effect on inulin were computed accounting for inter- and intra-study variability. 42 Dietary inulin levels ranged from 0.1 to 25.8 %, whereby the median and mean inulin levels 43 were 0.1 to 2% and 3 to 4%, respectively. Few of the investigated fermentation response 44 variables were influenced by dietary inulin. Strong negative relationships were found between 45 dietary inulin and gastric pH in weaned pigs ($R^2 = 0.81$; P < 0.001; n = 12), colonic 46 enterobacteria ($R^2 = 0.50$; P < 0.001; n = 19) and fecal lactobacilli ($R^2 = 0.41$; P < 0.001; n = 26) 47 throughout all production phases, whereas observed negative relationships between inulin and 48 colonic bifidobacteria and fecal enterobacteria and *Escherichia coli* were of minor physiological 49 relevance (P < 0.05). Moreover, increasing inulin levels negatively correlated with the CTTAD 50 of crude protein $(R^2 = 0.83; P < 0.001; n = 15)$, but they did not influence average daily gain of 51 pigs. Best-fit models indicated that dietary crude protein amplified the effect of inulin on 52 CTTAD of crude protein and gastric pH, but counteracted the inulin effect on fecal E. coli (P < 53 54 0.05). Accordingly, both pig's body weight and inulin decreased gastric pH and fecal lactobacilli

- but counteracted the inulin effect on colonic bifidobacteria and fecal *E. coli* (P < 0.05). In
- 56 conclusion, this study supported that dietary inulin can stimulate gastric acid secretion which
- 57 may be favorable GIT health in weaned pigs. However, meta-regressions did not support that
- 58 inulin promotes the bacterial groups previously associated with porcine GIT health, such as
- 59 lactobacilli and bifidobacteria.
- 60
- 61 *Keywords:* inulin, gastro-intestinal tract, fermentation, microbiota, meta-analysis, pig

62 **1. Introduction**

The ban of antimicrobial growth promoters in the EU has caused an overall high interest in 63 alternative feeding concepts and products to enhance disease resistance and support growth 64 performance in pig production (Metzler et al., 2005; Gallois et al., 2009). Especially, dietary 65 inclusion of functional ingredients and supplements, such as prebiotics, are of persistent interest 66 to maintain production efficiency in pigs (de Lange et al., 2010; Pluske, 2013). Among others, 67 68 considerable attention has been paid to the non-digestible oligosaccharide inulin for which health benefits around weaning have been reported (Modesto et al., 2009; Jensen et al., 2011). Inulin 69 encompasses all β -(2,1)-linear fructans of varying chain lengths (Roberfroid, 2007) and can be 70 71 found in several fruits and vegetables, like asparagus, leek, onions, banana, wheat and garlic, and 72 in higher concentrations in chicory (Compositae family) and Jerusalem artichoke (Helianthus tuberosus). Industrially, inulin is predominantly extracted from chicory (Roberfroid, 2005; 73 Kleessen et al., 2007; Ramnani et al., 2010). Inulin-type fructans are resistant to hydrolysis by 74 enzymes in the small intestine, but are rapidly fermented by saccharolytic bacteria including 75 bifidobacteria and lactobacilli (Konstantinov et al. 2004; Kleessen et al., 2007; Kolida and 76 Gibson, 2007; Liu et al., 2016). Promotion of these bacterial genera by dietary inulin may 77 78 suppress the growth of enterotoxigenic *Escherichia coli*, thereby lowering the risk for post-79 weaning diarrhoea in piglets (Halas et al., 2009). Although inulin has been consistently shown to exert prebiotic functions in the human hindgut from infants to the elderly (Kelly, 2008; Stiverson 80 et al., 2014; Liu et al., 2016), the reported effects in pigs were more contradictory (e.g., Verdonk 81 82 et al., 2005; Loh et al., 2006). Analysis of digesta from various segments of the small and large intestines revealed measureable inulin concentrations in the jejunum and ileum, but not in the 83 cecum and colon of pigs (Branner et al., 2004; Böhmer et al., 2005), which may indicate a 84 85 reduced capacity of inulin to modify porcine hindgut fermentation. Yet, beneficial effects on the

86 microbial composition in the colon or feces were found (e.g., Janczyk et al., 2010; Gao et al.,

87 2015). Likewise, modulation of the gastrointestinal tract (GIT) microbiota by dietary inulin has

been assumed to be most effective in newly weaned pigs (Konstantinov et al., 2004; Janczyk et

89 al., 2010); however, enhanced hindgut fermentation was lately reported for finishing pigs

90 receiving a diet with 5% inulin (Gao et al., 2015).

In general, qualitative reviews on alternative feed additives have repeatedly addressed the 91 effect of dietary supplementation of inulin on GIT health in weaned and growing pigs (e.g., 92 Verdonk et al., 2005; de Lange et al., 2010). Changes in direct (type and dose) and indirect 93 factors (e.g., age of the animal) can cause varying results across research studies which cannot 94 95 be considered in qualitative reviews (Sales, 2014). Also, it is difficult to examine all potential influencing factors in one single experiment. To address this complexity, a meta-analysis of 96 published studies is an efficient way to evaluate different factors by generalizing the overall 97 treatment effect (Charbonneau et al., 2006). So far, results for inulin research in pigs were not 98 investigated using a meta-analytical approach to summarize results across individual 99 experiments and therefore across a wide range of experimental conditions. With the 100 inconsistency obtained in empirical studies on the effects of inulin on GIT fermentation, the 101 current meta-analysis was designed to quantify the effect of dietary inulin supplementation on 102 fermentation metabolites and bacterial abundances in the GIT of weaned, growing and finishing 103 pigs. Additionally, effects of inulin on growth performance and coefficients of ileal (CIAD) and 104 total tract apparent digestibility (CTTAD) of nutrients and dry matter (DM) were assessed using 105 data from the studies included in the datasets for microbial fermentation and abundances. 106 107

- 108 **2.** Materials and methods
- 109 *2.1. Literature Search*

A literature search was conducted using the public search generators Pubmed, Google 110 111 Scholar, Web of Science, and Scopus. The main aim of the present study was the impact of dietary inulin supplementation on microbial abundances and fermentation metabolites in the GIT 112 of pigs. For that reason, research articles in scientific journals on controlled experiments 113 114 investigating the effect of inulin supplementation from purified or natural sources on intestinal fermentation and bacterial abundance that appeared between the years 2000 and January 2016 115 116 were primarily considered for data extraction. The following search terms in different 117 combinations were applied to identify adequate articles: inulin, chicory, chicory root, Jerusalem 118 artichokes, pig, piglet, swine, gut, large intestine or individual segments, small intestine or 119 individual segments, stomach, fermentation, microbial metabolites, volatile fatty acids (VFA) and short-chain fatty acids, lactate, bacteria, microbiota, microflora, and microbiome. 120

121

122 *2.2. Selection of studies*

Stringent criteria were in place whether published experiments were included or excluded in 123 this study. Quality assessment criteria included information about dietary composition, inulin 124 level and source (purified concentrate or natural source), type of pigs, body weight (BW) and age 125 of the pigs, number of pigs within treatment groups, duration of the experimental period, 126 127 experimental design including randomization of treatment groups, description of statistical 128 analysis, and intra-study error (if standard deviation was provided, it was converted into standard error), as well as fermentation metabolites (i.e., volatile fatty acids (VFA) and lactate), pH, and 129 130 bacterial abundances in digesta of stomach, ileum, cecum, proximal, mid and distal colon and rectum or feces. Studies were also included that investigated the combined effects of inulin with 131 another treatments on the search parameters. From those studies, data for the control without any 132 133 treatment and the sole inulin treatment were considered, or, if the basal (control) diet already

134	contained the other alternative feed additive, data for this basal diet without inulin and with inulin
135	were included. Published research studies on in-vitro experiments were excluded.
136	
137	2.3. Construction of Database
138	Our search found 45 articles that were eligible for the present meta-analysis by meeting a
139	sufficient number of above mentioned eligibility criteria. Beside the dietary inulin level as main
140	prediction variable and dependent variables microbial abundance, pH and fermentation
141	metabolites in the various GIT segments, given details on pig (breed, age, BW, gender,
142	production stage), experimental design, housing condition, dietary ingredients and chemical
143	composition of diets were extracted from the 45 articles to be considered as probable additional
144	prediction variables in the regression analysis. If provided, average daily feed intake, average
145	daily weight gain (ADG) and CIAD and CTTAD of DM and nutrients were extracted as well.
146	Careful examination and quality assessment of the dataset, however, showed that predictor
147	variables and dependent variables of interest were not always available across all studies or ill-
148	defined, leading to a large number of missing data. The main criterion to be considered in this
149	meta-analysis was "age" and "start BW" at the beginning of the experiment in order to use these
150	variables as additional predictor variable to consider maturational changes from weaned to
151	finisher pigs. Studies used different crude protein (CP) levels within the same production stage
152	and for similar starting BW. Since the dietary CP content can influence the intestinal
153	fermentation, digestibility and growth performance, this variable was included as further
154	predictor variable. Because "age" and "start BW" of pigs were not provided in all studies but
155	more often "start BW" than "age" was given, we decided to set "start BW" as the required
156	information needed to be provided in the study to remain in the dataset for analysis. Studies that
157	did not provide "start BW" at the beginning of the experimental period were removed from the

dataset. Due to this, twelve articles had to be excluded and a total of 33 studies formed the
"filtered" dataset which was used to compile the sub-datasets for GIT fermentation metabolites
and digesta characteristics (pH and DM), absolute bacterial abundances, growth performance,
CIAD and CTTAD of DM and nutrients with data from weaned, growing and finishing pigs. The
list of publications from which sub-dataset were built is provided in Table 1. The sub-datasets for
GIT fermentation metabolites, pH and absolute bacterial abundances were divided further; one
sub-dataset was created for each GIT site.

As minimum, three studies were set as requirement to quantify a combined effect size 165 (Lipsey and Wilson, 2001). In addition, a minimum of single observations (treatment means) of 166 167 10 per dependent variable as well as the respective standard error (SE) of each variable were set as further requirement to measure the combined effect size. According to this requirement, 168 169 sufficient numbers of studies and observations were available to evaluate ADG, and CIAD of DM and CP, and CTTAD of DM, CP, ash and neutral detergent fiber (NDF) as dependent 170 variables for performance. Luminal pH, total and individual VFA (i.e., acetate, propionate, 171 butyrate, isobutyrate, valerate and isovalerate) and lactate in gastric, ileal, cecal and colonic 172 digesta and feces were response variables related to microbial action. Data of fermentation 173 metabolites in digesta from proximal and distal colon were also extracted. However, they did not 174 175 fulfill the minimum requirement of 10 single observations (treatment means). As dependent 176 variables for absolute bacterial abundances, sufficient numbers of observations were only available for lactobacilli, bifidobacteria, enterobacteria and E. coli; however, not for all GIT 177 178 segments. Although studies using quantitative PCR and fluorescence-in-situ-hybridization (FISH) quantified the abundances of other bacterial groups which are difficult to culture, the number of 179 observations was often too small and primer and probe sets not equivalent, thereby hampering the 180 181 comparison of data. Results of bacterial abundances originated from both culturing and

quantitative molecular approaches. Although these are different methods of quantification, it was 182 183 assumed that colony forming units (CFU) correlate to gene copy numbers per gram of sample (Hein et al., 2001). Bacterial data were expressed as CFU/g digesta or fecal sample. If provided 184 on DM basis, fermentation metabolites and bacterial abundances were converted to fresh matter 185 186 basis. Data reported for the rectum were included in the "feces dataset" for fermentation metabolites, pH, digesta DM and absolute bacterial abundances. Fermentation and bacterial data 187 reported to be collected from the colon were allocated to the "mid-colon dataset" for 188 fermentation, pH and digesta. 189

190 Taken together, the recorded information from the research articles that matched the 191 inclusion criteria included authors, year of publication, dietary inulin level and source (i.e., purified or natural source), experimental design, sex, type and start BW of pig, breed, housing 192 (individual or pen), number of pigs per treatment, duration of feeding period, number of 193 experimental periods, and dietary main cereals and protein feedstuffs, and dietary CP level as 194 well as the dependent variables. The chain length of inulin was not provided in most research 195 articles and could therefore not be considered. Other dietary fibrous components might interact in 196 the inulin effects on fermentation and bacterial variables and should be considered. However, the 197 dietary level of fibrous components and the fructan content of the basal diet were not provided in 198 all studies or different fiber analytical methodologies were applied (e.g., crude fiber, total dietary 199 fiber, neutral-detergent fiber), thereby hindering comparisons among studies. Moreover, due to 200 the inconsistencies in reporting, breed and sex were also not included in the analysis. 201 202

202

203 *2.4. Data Analysis*

Descriptive statistics on predictor and dependent variables was performed using the MEANS
 procedure of SAS (SAS Inst. Inc., version 9.4). Microbial, CIAD, CTTAD and performance data

were subjected to mixed modeling analysis using the MIXED procedure according to the
following algorithm (St-Pierre, 2001):

208
$$Y_{ij} = \alpha_0 + \beta_1 X_{ij} + s_i + b_i X_{ij} + e_{ij}$$

where Y_{ij} = expected outcome for the dependent variable *Y* observed at level *j* (j = 2,..., n) of the predictor variable *X* in the study *i*, whereas *n* is the number of treatment means in study *i*, α_0 = overall intercept across all studies (fixed effect), β_1 = overall regression coefficient of Y on X across all studies (fixed effect), X_{ij} = the value *j* of continuous variable X in study *i*, s_i = random effect of the study *i* (*i* = 1,...), b_i = the random effect of study *i* on the regression coefficient of *Y* on *X* in study *i*, and e_{ij} = the unexplained error. Thus, the random effect components of the model include $s_i + b_i X_{ij} + e_{ij}$, and the distributions are shown below:

216
$$e_{ij} \sim iid N(0, \sigma_e^2)$$
 and $\begin{bmatrix} s_i \\ b_i \end{bmatrix} \sim iid N\left[(\frac{0}{0}), \Sigma\right]$, which assumes that e_{ij} is normally distributed

with a mean of 0 and constant variance, and that s_i and b_i are normally distributed, have means of

218 0 and
$$\Sigma$$
 is their variance-covariance matrix: $\Sigma = \begin{bmatrix} \sigma_s^2 & \sigma_{sb} \\ \sigma_{sb} & \sigma_b^2 \end{bmatrix}$.

219 Predictor variables tested were study, start BW, the dietary CP and inulin level. The slope and intercept by study, start BW, the dietary CP and inulin level were initially included as random 220 effects and an unstructured variance-covariance matrix (type = UN) was used to avoid a positive 221 222 correlation between intercepts and slopes (St-Pierre, 2001). To take the unequal variance among studies into account, the dependent variable was weighted by the inverse of its squared SE (SE of 223 treatment means were taken directly from studies). When a predictor variable was significant (P 224 < 0.05), its squared term was included in the model to test any quadratic relationship. In this case, 225 the variance-covariance matrix was modeled as variance components (TYPE = VC). Significant 226

quadratic relationships did not exist, only linear relationships between predictor and response variables for the present datasets. Trends were discussed at $0.05 \le P \le 0.10$. Data were visualized using the GPLOT procedure. Estimates, root mean square error (RMSE) and R^2 were computed and used to evaluate the goodness of fit. For established relationships, changes in the quantity of dependent variables caused by dietary inulin supplementation were illustrated for an assumed dietary inulin level of 3%.

To obtain a more precise prediction of influencing factors on dependent variables that 233 were influenced by the dietary inulin level, we used backward elimination analysis (Zebeli et al., 234 2008). In doing so, we concurrently evaluated the effects of the predictor variables dietary inulin 235 236 level, squared dietary inulin level, start BW as well as the dietary CP level on the response variables. Start BW was used as indicative for maturational changes from weaner to finishing 237 period. Changes in the dietary composition, such as the dietary CP content, affect gastrointestinal 238 microbial action. The dietary CP content varied among studies within one production phase and 239 for similar starting BW. Therefore, the effect of dietary CP was taken into consideration. Model 240 overparameterization was limited by considering a variance inflation factor less than 10 (which 241 assumes no significant multicollinearity among predictor variables tested) for every continuous 242 independent variable tested (Neter et al., 1996). 243

244

245 **3. Results**

246 *3.1. Database description*

Table 1 presents the main characteristics of the 33 studies between the years 2000 and January 2016 included in this meta-analysis. In eight studies the inulin supplement originated from chicory root (extract, powder or fiber) or Jerusalem artichokes, and in 25 studies from commercially available purified inulin concentrates (Table 1). The experimental diets were

251	mainly composed of wheat, barley, and corn, with soybean meal, fish meal, skimmed milk
252	powder, whey protein and soy protein concentrate as protein feedstuffs (Table 1).
253	Results of the descriptive statistics for the response variables of fermentation metabolites,
254	pH and digesta DM, bacterial abundances as well as ADG, CIAD and CTTAD of DM and
255	nutrients are presented in Table 2, 3 and 4. Inulin supplementation levels ranged from 0.1 to
256	25.8% with means for the various categories of response variables averaging around 3 to 4%
257	inulin (as-fed; Tables 2, 3 and 4). Only in one study the effect of a very high dietary
258	supplementation level of 25.8% inulin was investigated; this study was included in the
259	performance, CIAD and CTTAD sub-dataset. In the other sub-datasets, maximum inulin levels
260	were 15 to 20% (Tables 3 and 4). Median values, however, showed that most data were available
261	for low dietary inulin levels of 0.1 to 2% (Tables 2, 3 and 4). Body weight of pigs ranged from
262	5.9 to 112 kg (Tables 2, 3 and 4). With regards to the response variables for ADG, CIAD and
263	CTTAD, cecal and mid-colonic fermentation metabolites as well as for colonic and fecal
264	bacterial abundances, minimum and maximum BW values indicated that all production phases
265	were covered in this study (Tables 2, 3 and 4), whereas the influence of inulin on gastric pH only
266	included data from weaned pigs (Table 3). Ileal and fecal fermentation metabolites and ileal
267	bacterial abundances encompassed data from weaned and growing pigs (Table 2 and 3).
268	According to the means and median values throughout all categories of response variables, the
269	data originated mostly from weaned and growing pigs. The dietary CP levels ranged from 13.7 to
270	24.5% DM and had mean and median values of about 20% DM (Tables 2, 3 and 4).
271	
272	3.2. Inulin Effects on Digesta pH and Fermentation Metabolites along the GIT
273	In weaned pigs, gastric pH showed a negative linear relationship with increasing dietary

inulin levels ($R^2 = 0.81$; P < 0.001; Table 5). Accordingly, a dietary inclusion level of 3% would

275	decrease gastric pH by 0.12 units, whereas the digesta pH in ileum, cecum, colon and feces was
276	not affected by the dietary inulin level. The ileal lactate concentration tended to increase with
277	more inulin in the diet ($R^2 = 0.28$; $P = 0.062$), which would amount to 4.5 mmol/kg with 3%
278	inulin. In contrast, there was a small negative relationship between the cecal concentration of
279	acetate and increasing dietary inulin levels ($R^2 = 0.13$; $P = 0.080$). Fermentation metabolites in
280	colonic digesta and feces, in turn, were not influenced by the dietary inulin level.

281

282 *3.3. Inulin Effects on Ileal, Colonic and Fecal Bacterial Abundances*

Ileal abundances of lactobacilli and bifidobacteria were independent of the dietary inulin 283 284 level (Table 6). Likewise, an increasing dietary inulin level from 0 to 20% did not modify the absolute abundance of lactobacilli in colonic digesta. By contrast, higher dietary inulin levels 285 lowered the colonic abundance of bifidobacteria and enterobacteria, whereby the inhibiting 286 effect of inulin was twice as strong for enterobacteria (-0.55 log units with a dietary inulin level 287 of 3%; $R^2 = 0.50$; P < 0.001) as for bifidobacteria (-0.29 log units with a dietary inulin level of 288 3%; $R^2 = 0.37$; P = 0.022). In feces, increasing inulin levels reduced the abundance of 289 lactobacilli ($R^2 = 0.61$; P < 0.001) which amounted to a reduction in lactobacilli numbers of 1.69 290 log units with a dietary inulin level of 3%, whereas bifidobacteria tended to be slightly enhanced 291 by dietary inulin ($R^2 = 0.29$; P = 0.086). Increasing dietary inulin levels reduced the absolute 292 enterobacteria abundance ($R^2 = 0.23$; P = 0.006) and the abundance of E. coli in feces ($R^2 = 0.55$; 293 P < 0.001). Accordingly, a dietary inulin supplementation of 3% reduced the enterobacteria and 294 *E. coli* numbers by 0.32 and 0.19 log units, respectively. 295 296

297 *3.4. Inulin Effects on ADG, CIAD and CTTAD*

Average daily weight gain was not affected by dietary inulin when inulin was supplemented in the range from 0 to 25.8% ($R^2 = 0.05$; P = 0.311; Table 7). There was a tendency for a linear effect that increasing dietary inulin levels decreased CIAD of DM ($R^2 = 0.24$; P = 0.091) which corresponded to a decrease in CIAD of 7.4% with a dietary inulin level of 3%. The CTTAD of CP also linearly decreased with increasing inulin levels ($R^2 = 0.83$; P < 0.001), amounting to a 1.3%-decrease with a dietary supplementation level of inulin of 3%.

304

305 *3.5. Effects of pig's BW, dietary CP and inulin level*

Including the dietary CP and starting BW of pigs in the same model as the dietary inulin 306 307 level enhanced the prediction accuracy for several variables when compared to the analysis of the dietary inulin level alone (Tables 8). As such, gastric pH linearly decreased with increasing 308 BW and dietary inulin level, but it was also negatively correlated with increasing dietary CP 309 levels ($R^2 = 0.98$; $P \le 0.003$). According to the equation derived from these associations, the 310 effect of BW was the strongest on gastric pH. Ileal lactate concentration showed a square effect 311 for dietary inulin level, thereby indicating that the positive relationship was asymptotic. As a 312 313 tendency, backward elimination showed that cecal acetate was mainly negatively associated with dietary inulin level ($R^2 = 0.35$; P = 0.071) but not influenced by other predictor variables tested. 314 Backward elimination further showed that increasing BW positively affected the colonic 315 316 abundance of bifidobacteria which was stronger and opposite to the dietary inulin effect ($R^2 =$ 0.74; P < 0.01). Colonic enterobacteria were only affected by inulin ($R^2 = 0.57$; P < 0.001); 317 318 however, the square effect of inulin indicated an asymptotic approximation. Fecal abundance of 319 lactobacilli was negatively associated with both increasing BW and increasing dietary inulin 320 level ($R^2 = 0.77$; P < 0.01). Moreover, backward elimination analysis indicated that increasing 321 BW and dietary inulin were the main factors influencing bifidobacteria abundance in feces ($R^2 =$

3220.54; P < 0.05). Likewise, *E. coli* abundance in feces was not only negatively affected by323increasing dietary inulin levels, but it was positively correlated to the dietary protein level and324pig's BW ($R^2 = 0.80; P < 0.05$). Furthermore, backward elimination showed a strong positive325relation between the CIAD of DM and increasing dietary CP levels as well ($R^2 = 0.89; P <$ 3260.001), whereas the CTTAD of CP was not only negatively related with the dietary inulin level327(P < 0.001) but also with the dietary CP level (P = 0.029).

328

329 **4. Discussion**

Published research showed inconsistent results for the microbiota-modulating abilities of 330 inulin in the GIT of pigs (e.g., Böhmer et al., 2005; Gao et al., 2015). Similarly, the GIT region 331 where dietary inulin would be most effective was not clear since some authors reported 332 alterations in the small intestine (ileum; Böhmer et al., 2005; Loh et al., 2006), whereas others 333 observed inulin-related changes in the large intestine (Loh et al., 2006; Janczyk et al., 2010; Gao 334 et al., 2015). Therefore, the present meta-analysis aimed at investigating and quantifying the 335 effects of dietary inulin supplementation on fermentation metabolites and absolute bacterial 336 abundances along the GIT and feces, together with effects on performance and CIAD and 337 **CTTAD** of DM and nutrients. Data of the included studies covered a wide range of experimental 338 339 conditions; therefore, models derived from these data may yield relevant predictions to assist in the conclusion of effects of the target factors (Sauvant et al., 2008). Overall, the current results 340 provided insights into the discussion of the usefulness of inulin supplementation in pig diets and 341 342 confirm that inulin can be effective along the GIT. The level of inclusion is one of the critical factors for measurable inulin effects in the GIT of 343 pigs. The medians of the dietary inulin levels of 0.1 to 2% inulin showed that the dietary 344

345 supplementation level may explain the small effect of inulin on fermentation metabolites

- 346 observed in the individual studies and in the present meta-analysis. According to the established
- 347 relationships between the dietary inulin supplementation and dependent variables (e.g., gastric
- ³⁴⁸ pH, ileal lactate, and colonic and fecal enterobacteria) a minimum supplementation level of
- 349 inulin of 3 to 5% may be advisable to modulate physiological and microbial parameters in the
- 350 GIT of pigs. Most studies used wheat and barley as main cereals in the diet. These cereals
- naturally contain fructans in a range of 0.2 to 4% in wheat and 0.5 to 1% in barley (Moshfegh et
- al., 1999). As the endogenous fructan levels of the basal diets were not provided in most studies,
- 353 the effect of them on the observed effects of the supplemented inulin could not be distinguished
- 354 in the present meta-analysis. Moreover, median values for the BW at the start of the experiment
- 355 indicate that most data originated from studies in weaned and growing pigs. This is consistent
- with the general assumption that alternative feed additives, such as prebiotics, are more effective
- 357 in young pigs due to their immature GIT functions and microbial community (e.g., de Lange et
- al., 2010). Relationships established in the present meta-analysis may be therefore more
- 359 applicable to young pigs.
- 360 Increasing dietary inulin levels negatively affected CTTAD of CP, gastric pH, bifidobacteria
- and enterobacteria in colonic digesta, and lactobacilli and *E. coli* in feces, whereas VFA
- 362 concentrations along the GIT and feces appeared to be mostly unaffected by inulin. A specific
- 363 stimulation of lactic acid producing bacteria in the small intestine, such as lactobacilli and
- 364 bifidobacteria (Van Loo, 2004) may have been indicated by the positive relationship between
- 365 dietary inulin level and ileal lactate. In humans, a general positive relationship between the daily
- consumed amount of inulin and the abundance of bifidobacteria in stool exists (Van Loo, 2004).
- 367 Since pigs have a higher microbial activity in the small intestine (Jensen and Jørgensen, 1994)
- and a lower abundance of bifidobacteria in the GIT than humans (Loh et al., 2006), the proposed
- 369 prebiotic effect of inulin in humans cannot be extrapolated to pigs. Unfortunately, CIAD and

370	CTTAD of inulin were not adequately provided in the research papers to link inulin availability
371	in digesta and microbial numbers along the GIT. It is generally estimated that about 50% of
372	inulin are prececally fermented in pigs (Graham and Åman, 1986; Böhmer et al., 2005), whereby
373	the reported range of prececal CIAD of inulin ranges from 50 to 98% in the literature (e.g.,
374	Branner et al., 2004; Böhmer et al., 2005). Differences in the degree of fermentation may be
375	associated with the source of inulin (natural versus purified), degree of polymerization and the
376	maturation of the porcine GIT microbiota. Theoretically, inulin should lead to a decrease in the
377	pH along the GIT due to stimulation of fermentation and hence VFA and lactate production
378	(e.g., Böhmer et al., 2005). This assumption may be supported by the present negative and
379	positive relationships between dietary inulin level and gastric pH and, as trend, ileal lactate,
380	respectively, whereby at least 3% inulin should be supplemented to achieve physiological
381	changes. Aside from fermentation, higher water-holding properties of non-absorbable sugars
382	such as inulin may have reduced the gastric passage, thereby enhancing the acidification of the
383	gastric content (Wiggins, 1984). However, it needs to be considered that the data for gastric pH
384	originated only from three studies in weaned pigs.
385	Lactic acid producing <i>Bifidobacterium</i> and <i>Lactobacillus</i> strains encode β-
386	fructofuranosidases with different activities towards short- and long-chain fructans (e.g., Janer et
387	al., 2004; Ryan et al., 2005; Saulnier et al., 2007). Present regression models, however, did not
388	show an enhancing effect on absolute abundances of bifidobacteria and lactobacilli in ileal
389	digesta of weaned and early growing pigs, thereby confirming findings from individual research
390	studies (e.g., Böhmer et al., 2005; Janczyk et al., 2010). This was contrary to our findings for
391	ileal lactate and may indicate that changes were more at metabolic level. In addition, a certain
392	bias from the combination of the different methodologies used for bacterial quantification (i.e.,
393	culturing versus PCR-based approaches) cannot be excluded. Moreover, multiple regression

394	models demonstrated that dietary inulin can modulate the abundance of lactobacilli and
395	bifidobacteria in the large intestine; thereby supporting the observations of many individual
396	studies (e.g., Loh et al., 2006; Halas et al., 2009; Janczyk et al., 2010; Yan et al., 2013). In
397	contrast to the general assumption but in conformity with some literature findings (e.g., Vhile et
398	al., 2012), colonic and fecal lactobacilli and bifidobacteria abundances were not always
399	positively correlated with the dietary inulin level. In fact, increasing inulin levels decreased
400	colonic bifidobacteria and fecal lactobacilli numbers. This raises the question if the present
401	observations were direct effects of inulin or related to changes in substrate availability in digesta
402	of the large intestine, microbial cross-feeding and other microbe-to-microbe interactions (Flint et
403	al., 2012). Accordingly, cross-feeding of lactate produced by bacterial inulin fructan
404	fermentation has been reported to increase Megasphaera elsdenii in the colon of growing pigs
405	(Mølback et al., 2007). Aside from lactobacilli and bifidobacteria strains, inulin-degraders are
406	widespread among other bacterial genera, such as Roseburia and Blautia within Clostridium
407	cluster XIVa (Eckburg et al., 2005; Manderson et al., 2005). Also, Catenibacterium and
408	Bacteroides appear to have growth advantages in cecal digesta of pigs fed inulin supplemented
409	diets (Yan et al., 2013). Since most inulin entering the large intestine would be available to the
410	cecal bacterial community, the cecal bacterial responses to inulin might provide the link between
411	small and large intestines. In spite of the fact that about <10 to 50% of the ingested inulin
412	reaches the cecum, we could only establish a small trend for a negative relationship between
413	dietary inulin and acetate concentration in cecal digesta, indicating alterations in the substrate
414	available for microbial fermentation. Unfortunately, regression models for bacterial abundances
415	in cecal digesta could not be developed due to ill-definition of variables in the respective
416	literature. Maturational changes in the abundances of bifidobacteria in colonic digesta, and
417	lactobacilli and <i>E. coli</i> in feces were indicated by their relationships with increasing BW and

thus the age of the pigs which either counteracted or strengthened the observed inulin effect. 418 419 Although the family *Enterobacteriaceae* belongs to the commensal microbiota in the porcine GIT (Mach et al., 2015; Metzler-Zebeli et al., 2015; Zhao et al., 2015), it contains some common 420 etiological agents of diarrhea including enterotoxigenic *E. coli* (Fairbrother et al., 2005). 421 422 Nutritional attempts to control the intestinal numbers of *E. coli*, especially in the early postweaning period, have therefore received considerable attention (de Lange et al., 2010; 423 Pluske, 2013). The current regression models indicated that dietary inulin might have the ability 424 to control colonic and fecal numbers of enterobacteria and fecal *E. coli* numbers. Whether this 425 reduction can be linked to the increased abundance of bifidobacteria as often presumed remains 426 427 open due to the complexity of the fecal bacterial community (e.g., Mach et al., 2015). However, meta-regression results also showed that the dietary CP level should be concurrently controlled 428 in order not to counteract a potential inhibiting effect of inulin on *E. coli* in feces. Finally, it 429 should be considered that only changes in bacterial abundances of more than 0.5 log units may 430 be of physiological relevance. Therefore, higher dietary supplementation levels of more than 6 to 431 8% are necessary to impair colonic bifidobacteria and to be effective against E. coli in feces 432 according to the present meta-regressions. 433 One selection criterion for alternative feed additives is their effect on growth performance 434 435 and feed efficiency. Although non-digestible carbohydrates may have a negative impact on pig's performance (Grieshop et al., 2001), this may not be applicable for most of the studies included 436 in this meta-analysis due to the low dietary inulin level. Accordingly, ADG appeared not be 437 438 sensitive to dietary inulin levels as no relation between dietary inulin and ADG could be established. This was despite the fact that negative relationships between increasing CIAD of 439 DM (as trend) and CTTAD of CP were found. Reduced CIAD of DM, but not of CP, with 440 increasing dietary inulin levels may indicate inulin residuals in ileal digesta. According to the 441

442	best-fit models, higher dietary CP levels could counteract the inulin effect on CIAD of DM. This
443	may be related to the fact that dietary CP can elongate the retention time of the feed in the
444	stomach, thereby allowing luminal bacteria more time to ferment dietary components (Wiggins,
445	1984). The CTTAD of CP was reduced with more inulin in the diet and, in contrast to the CIAD
446	of DM, this effect was greater with increasing dietary CP levels and can likely be associated with
447	enhanced microbial protein synthesis in the large intestine due to greater substrate availability.
448	
449	5. Conclusion
450	This meta-analysis showed that dietary inulin supplementation may have the ability to lower
451	gastric pH in weaned pigs. Together with the trend for higher ileal lactate with increasing dietary
452	inulin levels, this may support an increased microbial activity in the upper GIT. Despite the
453	negative relation between dietary inulin and bifidobacteria in the colon and lactobacilli in feces,
454	the observed inhibition of enterobacteria numbers in feces with higher dietary inulin levels may
455	be favorable for porcine GIT health postweaning. However, pig's BW and the dietary CP level
456	were other sources of variation which may act synergistically and counteract the inulin effect.
457	Finally, since some results were based on low numbers of observations and often low dietary
458	inulin levels were tested, established relationships should be regarded as universal trends and
459	may be more applicable for weaned and early growing pigs.
460	
461	Acknowledgements
462	This article is based upon work from COST Action FA1401 PiGutNet, supported by COST
463	(European Cooperation in Science and Technology).
464	
465	References

466	Aufreiter, S., Kim, J. H., O'Connor, D.L., 2011. Dietary oligosaccharides increase colonic weight
467	and the amount but not concentration of bacterially synthesized folate in the colon of piglets.
468	J. Nutr. 141, 366-372.
469	Brambillasca, S., Zunino, P. and Cajarville C. 2015. Addition of inulin, alfalfa and citrus pulp in
470	diets for piglets: Influence on nutritional and faecal parameters, intestinal organs, and colonic
471	fermentation and bacterial populations. Livest. Sci. 178, 243-250.
472	Branner, G.R., Böhmer, B.M., Erhardt, W., Henke, J., Roth-Maier, D.A., 2004. Investigation on
473	the precaecal and faecal digestibility of lactulose and inulin and their influence on nutrient
474	digestibility and microbial characteristics. Arch. Anim. Nutr. 58, 353-366.
475	Böhmer. B.M., Branner, G.R., Roth-Maier, D.A., 2005. Precaecal and faecal digestibility of
476	inulin (DP 10-12) or an inulin/Enterococcus faecium mix and effects on nutrient digestibility
477	and microbial gut flora. J. Anim. Physiol. Anim. Nutr. (Berl.) 89, 388-396.
478	Boudry, C., Poelaert, C., Portetelle, D., Thewis, A., Bindelle, J. 2012. Discrepancies in
479	microbiota composition along the pig gastrointestinal tract between in vivo observations and
480	an in vitro batch fermentation model. J. Anim. Sci. 90 (Suppl 4), 393-396.
481	Charbonneau, E., Pellerin, D., Oetzel, G.R., 2006. Impact of lowering dietary cation-anion
482	difference in nonlactating dairy cows: a meta-analysis. J. Dairy Sci. 89, 537-548.
483	de Lange, C.F.M., Pluske, J.R., Gong, J., Nyachoti, C.M., 2010. Strategic use of feed ingredients
484	and feed additives to stimulate gut health and development in young pigs. Livest. Sci. 134,
485	124–134.
486	Eckburg, P.B., Bik, E.M., Bernstein, C.N., Purdom, E., Dethlefsen, L., Sargent, M., Gill, S.R.,
487	Nelson, K.E., Relman, D.A., 2005. Diversity of the human intestinal microbial flora. Science
488	308, 1635–1638.

489	Fairbrother, J.M., Nadeau, E., Gyles, C.L., 2005. Escherichia coli in postweaning diarrhea in
490	pigs: an update on bacterial types, pathogenesis, and prevention strategies. Anim. Health
491	Res. Rev. 6, 17-39.

- 492 Flint, H.J., Scott, K.P., Duncan, S.H., Louis, P., Forano, E., 2012. Microbial degradation of
- 493 complex carbohydrates in the gut. Gut Microbes 3, 289–306.
- Gallois, M., Rothkötter, H.J., Bailey, M., Stokes, C.R., Oswald, I.P. 2009. Natural alternatives to
 in-feed antibiotics in pig production: can immunomodulators play a role? Animal 3, 1644-
- 496 <mark>1661.</mark>
- 497 Gao, L., Chen, L., Huang, Q., Meng, L., Zhong, R., Liu, C., Tang, X., Zhang, H., 2015. Effect of
- dietary fiber type on intestinal nutrient digestibility and hindgut fermentation of diets fed to
 finishing pigs. Livest. Sci.174, 53–58.
- Graham, H., Åman, P., 1986. Composition and digestion in the pig gastrointestinal tract of
 Jerusalem artichoke tubers. Food Chem. 22, 67–76.
- 502 Grela, E.R., Sobolewska, S., Rozinski, T., 2014. Effect of inulin extracts or inulin-containing
- plant supplement on blood lipid indices and fatty acid profile in fattener tissues. Polish J.
 Vet.Sci. 1, 93-98.
- 505 Grieshop, C.M., Reese, D.E., Fahey Jr., G.C., 2001. Nonstarch polysaccharides and
- 506 oligosaccharides in swine nutrition, in: Lewis, A.J., Southern, L.L., (Eds.), Swine Nutrition,
- second ed. CRC Press, Boca Raton, FL, pp. 187–212.
- Halas, D., Hansen, C.F., Hampson, D.J., Mullan, B.P., Wilson, R.H., Pluske, J.R., 2009. Effect of
- 509 dietary supplementation with inulin and/or benzoic acid on the incidence and severity of
- 510 post-weaning diarrhoea in weaner pigs after experimental challenge with enterotoxigenic
- 511 Escherichia coli. Arch. Anim. Nutr. 63, 267–280.

- 512 Hedemann MS, Bach Knudsen KE. 2010. Dried chicory root has minor effects on the
- 513 digestibility of nutrients and the composition of the microflora at the terminal ileum and in
- faeces of growing pigs. Livest. Sci. 134, 53-55.
- 515 Hein, I., Lehner, A., Rieck, P., Klein, K., Brandl, E., Wagner, M. 2001. Comparison of different
- 516 approaches to quantify *Staphylococcus aureus* cells by real-time quantitative PCR and
- application of this technique for examination of cheese. Appl. Environ. Microbiol. 67, 31223126.
- 519 Ivarsson, E., Liu, H.Y., Dicksved, J., Roos, S., Lindberg, J.E., 2012. Impact of chicory inclusion
- 520 in a cereal-based diet on digestibility, organ size and faecal microbiota in growing pigs.
- 521 Animal 4, 1077–1085.
- Janczyk, P., Pieper, R., Smidt, H., Souffrant, W.B., 2010. Effect of alginate and inulin on
 intestinal microbial ecology of weanling pigs reared under different husbandry conditions.
 FEMS Microbiol. Ecol. 72, 132-142.
- Janer, C., Rohr, L.M., Peláez, C., Laloi, M., Cleusix, V., Requena, T., Meile, L., 2004.
- Hydrolysis of oligofructoses by the recombinant beta-fructofuranosidase from ifidobacterium
 lactis. Syst. Appl. Microbiol. 27, 279-285.
- Jensen, B.B., Jørgensen, H., 1994. Effect of dietary fiber on microbial activity and microbial gas
 production in various regions of the gastrointestinal tract of pigs. Appl. Environ. Microbiol.
 60, 1897-1904.
- Jensen, A.N., Mejer, H., Mølbak, L., Langkjær, M., Jensen, T.K., Angen, Ø., Martinussen, T.,
- 532 Klitgaard, K., Baggesen, D.L., Thamsborg, S.M., Roepstorff, A., 2011. The effect of a diet
- with fructan-rich chicory roots on intestinal helminths and microbiota with special focus on
- 534 Bifidobacteria and Campylobacter in piglets around weaning. Animal 5, 851–860.
- Kelly, G., 2008. Inulin-type prebiotics-a review: part 1. Altern. Med. Rev. 13, 315-329.

536	Kjos, N.P., Øverland, M., Fauske, A.K., Sørum, H., 2010. Feeding chicory inulin to entire male
537	pigs during the last period before slaughter reduces skatole in digesta and backfat. Livest.
538	Sci. 134, 143–145.

539 Kleessen, B., Schwarz, S., Boehm, A., Fuhrmann, H., Richter, A., Henle, T., Krueger, M., 2007.

Jerusalem artichoke and chicory inulin in bakery products affect faecal microbiota of healthy
volunteers. Br. J. Nutr. 98, 540–549.

- Kolida, S., Gibson, G.R., 2007. Prebiotic capacity of inulin-type fructans. J. Nutr. 137 (Suppl.1),
 2503S-2506S.
- 544 Konstantinov, S.R., Awati, A., Smidt, H., Williams, B.A., Akkermans, A.D., de Vos, W.M.,
- 545 2004. Specific response of a novel and abundant Lactobacillus amylovorus-like phylotype to
 546 dietary prebiotics in the guts of weaning piglets. Appl. Environ. Microbiol. 70, 3821-3830.
- Lipsey, M.W., Wilson, D.B., 2001. The way in which intervention studies have "personality" and
 why it is important to meta-analysis. Eval. Health Prof. 24, 236-254.
- 549 Liu H, Ivarsson E, Dicksved J, Lundh T, Lindberg JE., 2012. Inclusion of chicory (Cichorium
- 550 intybus L.) in pigs' diets affects the intestinal microenvironment and the gut icrobiota Appl.
- 551 Environ. Microbiol. 78, 4102-4109.
- Liu, Y., Gibson, G.R., Walton, G.E., 2016. An in vitro approach to study effects of prebiotics and

553 probiotics on the faecal microbiota and selected immune parameters relevant to the elderly.

- 554 PLoS One 11(9):e0162604.
- Loh, G., Eberhard, M., Brunner, R.M., Hennig, U., Kuhla, S., Kleessen, B., Metges, C.C., 2006.
- Inulin alters the intestinal microbiota and short-chain fatty acid concentrations in growing
 pigs regardless of their basal diet. J. Nutr. 136, 1198-1202.
- 558 Lynch, M.B., Sweeney, T., Callan, J.J., Flynn, B., O'Doherty, J.V., 2007. The effect of high and
- low dietary crude protein and inulin supplementation on nutrient digestibility, nitrogen

560

561

562

excretion, intestinal microflora and manure ammonia emissions from finisher pigs. Animal 1, 1112-1121.

Lynch, B., Callan, J.J., O'Doherty, J.V., 2009. The interaction between dietary crude protein and

- fermentable carbohydrate source on piglet post weaning performance, diet digestibility and 563 564 selected faecal microbial populations and volatile fatty acid concentration. Livest. Sci. 124, 93–100. 565 Mach, N., Berri, M., Estellé, J., Levenez, F., Lemonnier, G., Denis, C., Leplat, J.J., Chevaleyre, 566 C., Billon, Y., Doré, J., Rogel-Gaillard, C., Lepage, P., 2015. Early-life establishment of the 567 swine gut microbiome and impact on host phenotypes. Environ. Microbiol. Rep. 7, 554-569. 568 569 Jolliff, J.S., Mahan, D.C., 2012. Effect of dietary inulin and phytase on mineral digestibility and tissue retention in weanling and growing swine. J. Anim. Sci. 90, 3012-3022. 570 Mair, C., Plitzner, C., Domig, K.J., Schedle, K., Windisch, W., 2010. Impact of inulin and a 571 572 multispecies probiotic formulation on performance, microbial ecology and concomitant fermentation patterns in newly weaned piglets. J. Anim. Physiol. Anim. Nutr. (Berl.) 94, 573 164-177. 574 Manderson, K., Pinart, M., Tuohy, K.M., Grace, W.E., Hotchkiss, A.T., Widmer, W., Yadhav, 575 M.P., Gibson, G.R., Rastall, R.A., 2005. In vitro determination of prebiotic properties of 576 oligosaccharides derived from an orange juice manufacturing by-product stream. Appl. 577 Environ. Microbiol. 71, 8383-8389. 578 Metzler, B., Bauer, E. and R. Mosenthin. 2005. Microflora management in piglets. Asian-Austr. 579 J. Anim. Sci. 18, 1353-1362. 580 Metzler-Zebeli, B.U., Ratrivanto, A., Jezierny, D., Sauer, N., Eklund, M., Mosenthin, R., 2009. 581
- 582 Effects of betaine, organic acids and inulin as single feed additives or in combination on

microbial populations in the gastrointestinal tract of weaned pigs. Arch. Anim. Nutr. 63, 427441.

Metzler-Zebeli, B.U., Schmitz-Esser, S., Mann, E., Grüll, D., Molnar, T., Zebeli Q., 2015.

- Adaptation of the cecal bacterial microbiome of growing pigs in response to resistant starch 586 type 4. Appl. Environ. Microbiol. 81, 8489-8499. 587 Modesto, M., Rosaria D'Aimmo, M., Stefanini, I., Trevisi, P., De Filippi, S., Casini, L., Mazzoni, 588 M., Bosi, P., Biavati, B., 2009. A novel strategy to select Bifidobacterium strains and 589 prebiotics as natural growth promoters in newly weaned pigs. Livest. Sci. 122, 248-258. 590 Mølbak, L., Thomsen, L.E., Jensen, T.K., Bach Knudsen, K.E., Boye, M., 2007. Increased 591 592 amount of Bifidobacterium thermacidophilum and Megasphaera elsdenii in the colonic microbiota of pigs fed a swine dysentery preventive diet containing chicory roots and sweet 593 lupine. J. Appl. Microbiol. 103, 1853–1867. 594 Moshfegh, A.J., Friday, J.E., Goldman, J.P., Ahuja, J.K., 1999. Presence of inulin and 595 oligofructose in the diets of Americans. J. Nutr. 129 (Suppl. 7), 1407S-1411S. 596 Neter, J., Kutner, M.H., Nachtsheim, C.J., Wasserman, W., 1996. Applied Linear Statistical 597 Models, forth ed. McGrow-Hill, Boston, MA. 598 O'Shea, C.J., Sweeney, T., Bahar, B., Ryan, M.T., Thornton, K., O'Doherty, J.V., 2012. Indices 599 600 of gastrointestinal fermentation and manure emissions of growing finishing pigs as influenced through singular or combined consumption of Lactobacillus plantarum and inulin. 601 J. Anim. Sci. 90, 3848-3857. 602 603 Øverland, M., Kjos, N.K., Fauske, A.K., Teige, J., Sørum, H., 2011. Easily fermentable
- 604 carbohydrates reduce skatole formation in the distal intestine of entire male pigs. Livest. Sci.
 - 605 140, 206-217.

606	Patterson, J.K., Lei, X.G., Miller, D.D., 2008. The pig as an experimental model for elucidating
607	the mechanisms governing dietary influence on mineral absorption. Exp. Biol. Med. 233,
608	651-664.

Patterson, J.K., Yasuda, K., Welch, R.M., Miller, D.D., Lei, X.G., 2010. Supplemental dietary

610 inulin of variable chain lengths alters intestinal bacterial populations in young pigs. J. Nutr.611 140, 2158-61.

Pierce, K.M., Callan, J.J., McCarthy, P., O'Doherty, J.V., 2005. Performance of weaning pigs
offered low or high lactose diets supplemented with avilamycin or inulin. Anim. Sci. 80,
313–318.

Pluske, J.R. 2013. Feed- and feed additives-related aspects of gut health and development in
weanling pigs. J. Anim. Sci. Biotechnol. 4, 1.

Ramnani, P., Gaudier, E., Bingham, M., van Bruggen, P., Tuohy, K.M., Gibson, G.R., 2010.

618 Prebiotic effect of fruit and vegetable shots containing Jerusalem artichoke inulin: a human
619 intervention study. Br. J. Nutr. 104, 233-240.

620 Ratriyanto, A., Mosenthin, R., Jezierny, D., Sauer, N., Eklund, M., 2009. Betaine, organic acids

and inulin do not affect ileal and total tract nutrient digestibility or microbial fermentation in

622 piglets. J. Anim. Feed Sci. 18, 453-364.

Rideout, T.C., Fan, M.Z., Cant, J.P., Wagner-Riddle, C., Stonehouse, P., 2004. Excretion of

624 major odor-causing and acidifying compounds in response to dietary supplementation of

625 chicory inulin in growing pigs J. Anim. Sci. 82, 1678-1684.

Roberfroid, M.B., 2005. Introducing inulin-type fructans. Br. J. Nutr. 93, S13-25.

Roberfroid, M., 2007. Prebiotics: the concept revisited. J. Nutr. 137, 830S-837S.

- 628 Rodrigues, M., Pozza, P.C., Pozza, M.S., Possamai, M., Bruno, L.G., Richart, E., Wochner,
- M.O., Júnior, J.M., 2013. Effects of inulin and a probiotic mixture on nutrient digestibility
 and nitrogen balance in piglets. Arch. Zootec. 62, 255-264.
- 631 Ryan, S.M., Fitzgerald, G.F. van Sinderen, D., 2005. Transcriptional regulation and
- 632 characterization of a novel bfructofuranosidase-encoding gene from Bifidobacterium breve.
- 633 Appl. Environ. Microbiol. 71, 3475–3482.
- Sales, J., 2014. Effects of access to pasture on performance, carcass composition, and meat
 quality in broilers: A meta-analysis. Poult. Sci. 93, 1523-1533.
- 636 Saulnier, L., Sado, P.-E., Branlard, G., Charmet, G., Guillon, F., 2007. Wheat arabinoxylans:
- Exploiting variation in amount and composition to develop enhanced varieties. J. Cereal Sci.
 46, 261–281.
- Sauvant, D., Schmidely, P., Daudin, J.J., St-Pierre, N.R., 2008. Meta-analyses of experimental
 data in animal nutrition. Animal 2, 1203–1214.
- 641 Sobolewska, S., Grela, E.R., 2015. Effect of inulin extract or inulin-containing plant supplement
- on short-chain fatty acids in the large intestine and jejunum histology of fatteners. Med.
- 643 Weter. 71, 432-435.
- 644 Stiverson, J., Williams, T., Chen, J., Adams, S., Hustead, D., Price, P., Guerrieri, J., Deacon, J.,
- Yu, Z., 2014. Prebiotic Oligosaccharides: Comparative Evaluation Using In Vitro Cultures
 of Infants' Fecal Microbiomes. Appl. Environ. Microbiol. 80, 7388-7397.
- 647 St-Pierre, N. R., 2001. Integrating quantitative finding from multiple studies using mixed model
 648 methodology. J. Dairy Sci. 84, 741–755.
- Tako, E., Glahn, R.P., Welch, R.M., Lei, X., Yasuda, K., Miller, D.D., 2008. Dietary inulin
- affects the expression of intestinal enterocyte iron transporters, receptors and storage protein
- and alters the microbiota in the pig intestine. Br. J. Nutr. 99, 472-480.

- Van Loo, J., 2004. The specificity of the interaction with intestinal bacterial fermentation by
 prebiotics determines their physiological efficacy. Nutr. Res. Rev. 17, 89-98.
- Varley, P.F., McCarney, C., Callan, J.J., O'Doherty, J.V., 2010. Effect of dietary mineral level
- and inulin inclusion on phosphorus, calcium and nitrogen utilisation, intestinal microflora
- and bone development. J. Sci. Food Agric. 90, 2447-2454.
- Verdonk, J.M., Shim, S.B., van Leeuwen, P., Verstegen, M.W., 2005. Application of inulin-type
 fructans in animal feed and pet food. Br. J. Nutr. 93, S125-S138.
- Vhile, S.G., Kjos, N.P., Sørum, H., Overland, M., 2012. Feeding Jerusalem artichoke reduced
- skatole level and changed intestinal microbiota in the gut of entire male pigs. Animal 6, 807-814.
- 662 Wellock, I.J., Fortomaris, P.D., Houdijk, J.G.M., Wiseman, J., Kyriazakis, I., 2008. The
- 663 consequences of non-starch polysaccharide solubility and inclusion level on the health and
- performance of weaned pigs challenged with enterotoxigenic Escherichia coli. Br. J. Nutr. 4,
 520–530.
- Wiggins, H.S., 1984: Nutritional value of sugars and related compounds undigested in the small
 gut. Proc. Nutr. Soc. 43, 69–75.
- 668 Yan, H., Potu, R., Lu, H., Vezzoni de Almeida, V., Stewart, T., Ragland, D., Armstrong, A.,
- Adeola, O., Nakatsu, C.H., Ajuwon, K.M., 2013. Dietary fat content and fiber type modulate
 hind gut microbial community and metabolic markers in the pig. PLoS One 8, e59581.
- 471 Yasuda, K., Roneker, K.R., Miller, D.D., Welch, R.M., Lei, X.G., 2006. Supplemental dietary
- inulin affects the bioavailability of iron in corn and soybean meal to young pigs. J. Nutr. 136,3033-3038.

674	Yasuda, K., Dawson, H.D., Wasmuth, E.V., Roneker, C.A., Chen, C., Urban, J.F., Welch, R.M.,
675	Miller, D.D., Lei, X.G., 2009. Supplemental dietary inulin influences expression of iron and
676	inflammation related genes in young pigs. J. Nutr. 139, 2018-2023.
677	Zhao, W., Wang, Y., Liu, S., Huang, J., Zhai, Z., He, C., Ding, J., Wang, J., Wang, H., Fan, W.,

- Zhao, J., Meng, H., 2015. The dynamic distribution of porcine microbiota across different 678
- ages and gastrointestinal tract segments. PLoS ONE 10, e0117441. 679
- Zebeli, Q., Dijkstra, J., Tafaj, M., Steingass, H., Ametaj, B.N., Drochner, W., 2008. Modeling the 680
- adequacy of dietary fiber in dairy cow based on responses of ruminal pH and milk fat 681
- production to composition of the diet. J. Dairy Sci. 91, 2046–2066. 682

683

List of references and the respective experimental variables included in the meta-analysis.

						Gastro	-intestin	al sites ^d			Variables	e
Reference	Inulin ^a	Pig	Feeding level ^b	Basal diet ^c	Sto Ile		Cec	Col	Fec	Ferm- metab	Bact- abund	Perf./ Dig.
Branner et al., 2004	р	grower	restr	corn, wheat, barley, soybean meal		х			х	х	х	х
Rideout et al., 2004	р	grower	s-ad lib	corn, corn starch, soybean meal					х	х		х
Böhmer et al., 2005	р	grower	restr	corn, wheat, barley, soybean meal		х			х	х	х	х
Pierce et al., 2005	р	weaner	ad lib	wheat, soybean meal	х		х	х	х	х	х	х
Yasuda et al., 2006	р	weaner	ad lib	corn, soybean meal					х	х		х
Lynch et al., 2007	р	finisher	restr	wheat, soybean meal			х	х		х		х
Mølback et al., 2007	na	grower	restr	triticale, barley, potato protein								
Tako et al., 2008	р	weaner	ad lib	corn, soybean meal								
Wellock et al., 2008	р	weaner	ad lib	oat, wheat, fish meal	х	Х	х	х		х	х	х
Lynch et al., 2009	р	weaner	ad lib	wheat, soybean meal					х	х	х	х
Metzler-Zebeli et al., 2009	р	weaner	restr	barley, wheat, soybean meal		х			х		х	
Ratriyanto et al., 2009	р	weaner	restr	barley, wheat, soybean meal		х			х	х		х
Yasuda et al., 2009	р	grower	ad lib	corn, soybean meal								х
Patterson et al., 2010	р	weaner	ad lib	corn, soybean meal								
Halas et al., 2010 Hedemann and Bach	р	weaner	ad lib	wheat, soybean meal			х	х		Х		
Knudsen, 2010	na	grower	restr	wheat, soybean meal								х
Kjøs et al., 2010	na	finisher	restr	barley, wheat, soybean meal					х		х	
Mair et al., 2010	р	weaner	ad lib	barley, corn, soy concentrate	х	х		х		х	х	
Varley et al., 2010	р	finisher	restr	wheat, soybean meal skim milk powder, whey powder, soy		х	х	х		Х	Х	Х
Aufreiter et al., 2011	р	weaner	restr	oil				х			х	
Øverland et al., 2011	na	grower	restr	barley, wheat, soybean meal			Х	х	х	х	х	х
Boudry et al., 2012	na	grower	restr	wheat, soybean meal								
Ivarsson et al., 2012	na	grower	ad lib	wheat, barley, potato					х		х	Х
Jolliff and Mahan, 2012	р	weaner	ad lib	corn, soybean meal								х
Liu et al., 2012	na	grower	ad lib	wheat, barley, potato		х		х	х	х		х

O'Shea et al., 2012	р	grower	ad lib	wheat, soybean meal				Х		х	Х
Vhile et al., 2012	na	finisher	restr	wheat, soybean meal			х	х	х	х	
Rodrigues et al., 2013	р	weaner	restr	corn, soybean meal							х
Yan et al., 2013	р	grower	-	corn, soybean meal		х			х		х
Grela et al., 2014	р	finisher	ad lib	wheat, barley, corn soybean meal							
Brambillasca et al., 2015	р	weaner	ad lib	corn, soybean meal	Х		х	х	х		х
Gao et al., 2015	р	grower	ad lib	corn, soybean meal		х		х	х		х
Sobolewska and Grela, 2015	р	grower	ad lib	wheat, barley soybean meal		х	х		х		

^a Inulin source: p, purified; na, natural.

^b Feeding level: restr, restrictive feeding; ad lib, ad libitum feeding; s-ad lib, semi-ad libitum feeding.

^c Main energy and protein feedstuffs of basal diet.

^d Intestinal sites: Sto, stomach; Ile, ileum; Cec, cecum; Col, colon; Fec, feces.

690 e Response variables: Ferm met, fermentation metabolites; Bact-abund, absolute bacterial abundances; Perf./Dig., performance and digestibility variables.

692 Descriptive statistics for dietary inulin and crude protein, start body weight, pH and fermentation

693	metabolites in	gastric, ileal,	cecal and colonic	digesta and	feces of pigs in t	he respective datasets.

Item ^a	$n_{\text{Treat}}^{b,c}$	Mean	SEd	Min.	Max.	Median
Gastric digesta						
Start BW (kg)	12	9.1	0.13	8.1	9.3	9.3
Dietary inulin (%)	12	3.5	1.64	0.0	15.0	0.2
Dietary CP (%)	12	21.9	0.72	17.8	24.0	23.2
pH	12	3.4	0.10	2.8	3.9	3.4
Ileal digesta						
Start BW (kg)	27	17.3	2.63	6.7	51.0	9.3
Dietary inulin (%)	27	4.3	1.22	0.0	20.0	0.4
Dietary CP (%)	27	20.8	0.55	16.6	24.0	22.2
рН	25	6.7	0.07	6.1	7.4	6.6
Total VFA (mmol/kg)	17	24.0	4.48	6.3	78.8	16.7
Acetate (mmol/kg)	15	15.5	2.15	5.0	39.4	14.2
Butyrate (mmol/kg)	13	2.1	0.69	0.0	8.2	1.1
Lactate (mmol/kg)	13	35.3	8.87	0.3	96.2	25.1
Cecal digesta						
Start BW (kg)	35	22.9	3.73	5.9	74.5	10.2
Dietary inulin (%)	35	3.6	0.82	0.0	20.0	2.0
Dietary CP (%)	20	21.2	0.55	16.6	24.0	21.8
pН	16	5.8	0.08	5.1	6.2	5.8
Total VFA (mmol/kg)	27	110.4	12.63	19.0	229.0	109.5
Acetate (mmol/kg)	25	51.0	7.80	10.7	138.8	48.1
Propionate (mmol/kg)	25	23.1	2.74	5.2	50.4	24.5
Butyrate (mmol/kg)	25	11.3	1.87	0.5	31.4	10.8
Iso-butyrate (mmol/kg)	17	1.0	0.16	0.2	2.3	0.8
Valerate (mmol/kg)	21	2.4	0.38	0.5	5.9	2.2
Iso-valerate (mmol/kg)	12	2.0	0.33	0.5	4.0	1.8
Lactate (mmol/kg)	10	6.2	1.87	0.0	15.1	6.7
Colonic digesta						
Start BW (kg)	26	43.2	8.02	5.9	112.0	30.0
Dietary inulin (%)	26	2.9	0.73	0.0	16.0	1.8
Dietary CP (%)	18	20.3	0.54	16.6	23.6	19.7
рН	18	5.8	0.08	5.2	6.3	5.8
Total VFA (mmol/kg)	19	133.9	15.28	45.1	243.6	146.6
Acetate (mmol/kg)	19	68 7	9 56	171	140.5	64.6

Propionate (mmol/kg)	19	31.8	3.37	11.3	52.6	26.5
Butyrate (mmol/kg)	19	14.1	2.62	1.1	38.2	11.5
Iso-butyrate (mmol/kg)	17	1.7	0.23	0.6	3.2	1.4
Valerate (mmol/kg)	19	3.2	0.33	1.6	5.9	2.7
Iso-valerate (mmol/kg)	14	3.2	0.59	0.4	6.1	3.5
Feces						
Start BW (kg)	26	17.0	2.16	6.0	36.5	12.7
Dietary inulin (%)	26	2.5	0.74	0.0	16.0	1.5
Dietary CP (%)	26	19.9	0.50	16.6	24.5	19.3
pН	20	6.8	0.13	6.0	8.1	6.6
Dry matter content (%)	16	24.2	0.97	17.1	33.0	23.8
Total VFA (mmol/kg)	16	88.4	15.98	7.3	181.7	92.1
Acetate (mmol/kg)	16	52.6	9.53	1.9	108.8	54.6
Propionate (mmol/kg)	16	19.1	3.12	3.5	35.7	19.9
Butyrate (mmol/kg)	16	10.1	1.94	0.8	19.6	10.1
Valerate (mmol/kg)	12	4.5	0.65	0.7	6.9	5.4

 a BW, body weight; CP; crude protein; n_{Treat} = number of treatment means included; SE = standard error; VFA,

695 volatile fatty acids.

^b Separate datasets for response variables in gastric, ileal, cecal and colonic digesta and feces were built.

- 699 Descriptive statistics for dietary inulin and crude protein, start body weight, and absolute
- abundances of bacterial groups in ileal and colonic digesta and feces of pigs in the respective
- 701 datasets.

Item ^a	n _{Treat} ^b	Mean	SE	Min.	Max.	Median
Ileal digesta						
Start BW (kg)	14	22.6	4.65	6.7	51.0	9.3
Dietary inulin (%)	14	3.2	1.68	0.0	20.0	0.1
Dietary CP (%)	14	19.4	0.78	16.6	24.0	18.0
Lactobacilli (CFU/g)	12	8.5	0.14	7.7	9.3	8.5
Bifidobacteria (CFU/g)	10	7.3	0.70	3.3	9.3	8.2
Colonic digesta						
Start BW (kg)	39	27.9	4.14	8.1	74.5	9.3
Dietary inulin (%)	39	3.7	0.86	0	20	1.25
Dietary CP (%)	39	21.0	0.44	16.6	24	22.8
Lactobacilli (CFU/g)	30	8.6	0.13	6.7	9.5	8.9
Bifidobacteria (CFU/g)	14	8.1	0.23	6.3	8.8	8.5
Enterobacteria (CFU/g)	19	6.5	0.30	2.7	8.0	6.7
Feces						
Start BW (kg)	39	25.2	3.04	2.4	67.0	21.9
Dietary inulin (%)	39	3.1	0.68	0.0	16.0	1.5
Dietary CP (%)	39	19.6	0.45	16.5	24.5	18.7
Lactobacilli (CFU/g)	26	8.7	0.31	4.2	10.7	9.2
Bifidobacteria (CFU/g)	13	8.2	0.42	5.0	9.6	8.6
Enterobacteria (CFU/g)	20	7.4	0.31	5.2	9.8	7.0
Escherichia coli (CFU/g)	19	6.6	0.12	5.8	8.0	6.6

702 ^a BW, body weight; CFU, colony forming units; CP, crude protein; n_{Treat} = number of treatment means included; SE

703 = standard error.

^b Separate datasets for response variables in ileal, and colonic digesta and feces were built.

706	Descriptive statistics	for dietary inulin	and crude protein,	start body weight	, average daily gain,
		2	1 /	, , , , , , , , , , , , , , , , , , , ,	, , , , , , , ,

and coefficients of apparent ileal and total tract digestibility of pigs in the respective dataset.

Item ^a	n _{Treat}	Mean	SE	Min.	Max.	Median
Start BW (kg)	61	21.8	2.28	6.0	74.5	15.5
Dietary inulin (%)	61	4.2	0.80	0	25.8	1.5
Dietary CP (%)	61	20.4	0.34	13.7	24.5	19.9
ADG (g)	25	611	63.3	75.0	981.5	623
CIAD of DM (%)	13	79.2	2.05	67.9	87.8	76.0
CIAD of CP (%)	11	73.8	0.58	71.0	78.5	73.5
CTTAD of DM (%)	27	87.5	0.41	83.6	91.0	87.0
CTTAD of CP (%)	15	77.4	5.15	74.3	86.0	83.5
CTTAD of ash (%)	15	59.8	1.94	46.1	68.7	62.9
CTTAD of NDF (%)	19	59.2	1.71	39.1	69.8	61.2

708 ^a ADG, average daily gain; BW, body weight; CIAD, coefficient of ileal apparent digestibility; CP, crude protein;

⁷⁰⁹ CTTAD, coefficient of total tract apparent digestibility; DM, dry matter; NDF, neutral detergent fiber; n_{Treat} =

⁷¹⁰ number of treatment means included; SE = standard error.

712 Prediction of pH and fermentation metabolites in gastric, ileal, cecal and colonic digesta and

feces as affected by supplementation dose of inulin (%) in pig diets for all production phases.

		Parameter estimates				Model statistics			
Response variable (Y) ^a	n _{Treat}	Intercept	SEIntercept	Slope	SE _{Slope}	RMSE	R^2	P-value	
Gastric digesta					·				
pH	12	3.51	0.039	-0.040	0.006	0.115	0.81	< 0.001	
Ileal digesta									
pН	25	6.62	0.073	0.015	0.009	0.293	0.01	0.116	
Total VFA (mmol/kg)	17	22.94	4.086	-0.130	0.520	14.018	0.00	0.806	
Acetate (mmol/kg)	15	16.27	2.093	-0.343	0.318	6.783	0.08	0.300	
Butyrate (mmol/kg)	13	1.87	0.561	-0.061	0.079	1.641	0.05	0.456	
Lactate (mmol/kg)	13	21.09	5.051	1.486	0.715	14.926	0.28	0.062	
Cecal digesta									
pН	16	5.82	0.077	-0.017	0.010	0.262	0.16	0.130	
Total VFA (mmol/kg)	27	109.46	15.887	0.111	2.988	66.103	0.00	0.971	
Acetate (mmol/kg)	25	63.80	10.227	-4.920	2.683	36.892	0.13	0.080	
Propionate (mmol/kg)	25	26.40	3.688	-1.296	0.967	13.302	0.07	0.193	
Butyrate (mmol/kg)	25	13.76	2.432	-1.005	0.638	8.772	0.10	0.129	
Valerate (mmol/kg)	21	2.12	0.516	0.096	0.130	1.699	0.03	0.470	
Mid colonic digesta									
pН	18	5.83	0.092	-0.023	0.019	0.321	0.09	0.234	
Total VFA (mmol/kg)	19	135.20	19.083	-0.542	3.941	68.374	0.00	0.892	
Acetate (mmol/kg)	19	71.25	11.885	-0.974	2.454	42.592	0.01	0.696	
Propionate (mmol/kg)	19	31.60	4.186	0.018	0.864	0.983	0.00	0.983	
Butyrate (mmol/kg)	19	15.06	3.249	-0.338	0.671	11.642	0.02	0.621	
Iso-butyrate (mmol/kg)	17	1.84	0.269	-0.059	0.052	0.888	0.08	0.276	
Valerate (mmol/kg)	19	3.10	0.403	0.029	0.083	1.445	0.01	0.736	
Iso-valerate (mmol/kg)	14	2.83	0.707	0.121	0.131	2.234	0.07	0.375	
Feces									
pН	20	6.82	0.151	-0.024	0.034	0.574	0.03	0.487	
Dry matter content (%)	16	23.74	1.251	0.217	0.389	3.959	0.02	0.586	
Total VFA (mmol/kg)	16	88.18	20.433	0.131	6.391	65.969	0.00	0.984	
Acetate (mmol/kg)	16	52.82	12.179	-0.209	3.809	39.320	0.00	0.957	
Propionate (mmol/kg)	16	18.36	3.978	0.406	1.244	12.844	0.01	0.749	
Butyrate (mmol/kg)	16	9.47	2.474	0.327	0.774	7.989	0.01	0.679	
Valerate (mmol/kg)	12	4.30	0.834	0.087	0.232	2.296	0.01	0.716	

714 ^a VFA, volatile fatty acids; n_{Treat} = number of treatment means included; RMSE = root mean square error.

717 Prediction of absolute abundances of bacterial groups in ileal and colonic digesta and feces as

718	affected by	supplementation	dose of inulin	(%) in	nig diets	for all	production	classes
10	anceled by	supprementation	uose or munn	(/0) m	pig ulous	ioi an	production	classes.

			Parameter est	Mo	Model statistics				
Response variable (Y) ^a	n _{Treat}	Intercept	SEIntercept	Slope	SE _{Slope}	RMSE	R^2	<i>P</i> -value	
Ileal digesta									
Lactobacilli (CFU/g)	12	8.53	0.155	-0.010	0.033	0.483	0.01	0.771	
Bifidobacteria (CFU/g)	10	7.38	0.801	-0.017	0.125	2.339	0.00	0.893	
Colonic digesta									
Lactobacilli (CFU/g)	30	8.62	0.140	0.005	0.020	0.652	0.00	0.821	
Bifidobacteria (CFU/g)	14	8.28	0.195	-0.095	0.036	0.679	0.37	0.022	
Enterobacteria (CFU/g)	19	7.07	0.256	-0.184	0.044	0.923	0.50	< 0.001	
Feces									
Lactobacilli (CFU/g)	26	9.33	0.262	-0.562	0.102	1.0317	0.61	< 0.001	
Bifidobacteria (CFU/g)	13	7.47	0.558	0.140	0.073	1.455	0.29	0.086	
Enterobacteria (CFU/g)	20	7.18	0.193	-0.108	0.037	0.889	0.23	0.006	
Escherichia coli (CFU/g)	19	6.86	0.096	-0.063	0.014	0.303	0.55	< 0.001	

^a CFU, colony forming units; n_{Treat} = number of treatment means included; RMSE = root mean square error.

- 722 Prediction of growth performance and coefficients of apparent ileal and total tract digestibility as
- affected by supplementation dose of inulin (%) in pig diets for all production classes.

			Parameter es	Model statistics				
Response variable $(Y)^a$	n _{Treat}	Intercept	SEIntercept	Slope	SE _{Slope}	RMSE	R^2	P-value
ADG (g)	25	606.99	76.404	-9.017	8.695	256.122	0.05	0.311
CIAD of DM (%)	13	81.18	2.146	-2.473	1.334	6.691	0.24	0.091
CIAD of CP (%)	11	73.68	0.658	0.024	0.067	1.808	0.01	0.727
CTTAD of DM (%)	27	87.29	0.421	0.145	0.102	2.017	0.08	0.168
CTTAD of CP (%)	15	84.51	0.467	-0.443	0.055	1.555	0.83	< 0.001
CTTAD of ash (%)	15	58.52	1.332	0.320	0.247	4.732	0.11	0.218
CTTAD of NDF (%)	19	58.73	1.850	0.234	0.382	7.396	0.02	0.548

^a ADG, average daily gain; CIAD, coefficient of ileal apparent digestibility; CP, crude protein; CTTAD, coefficient

725 of total tract apparent digestibility; DM, dry matter; NDF, neutral detergent fiber; n_{Treat} = number of treatment means

726 included; RMSE = root mean square error.

729 Best-fit equations showing the coefficients of apparent ileal and total tract digestibility and microbial response variables in relation to

race increasing dietary inulin and crude protein level, and pig's start body weight using backward elimination technique.

				Parameter e	Model statistics					
Response variable (<i>Y</i>) ^a	Predictor (X)	n _{Treat}	Intercept	SE _{Intercept}	Slope	SE _{Slope}	RMSE	R^2	VIF	P-value
Gastric pH		15	5.28	0.228			0.037	0.98		
	BW (kg)				-0.140	0.030			1.52	0.002
	dietary CP (%)				-0.024	0.006			1.52	0.003
	Inulin (%)				-0.034	0.002			1.09	< 0.001
Ileal lactate (mmol/kg)		13	20.24	4.201			13.274	0.61		
	Inulin-square (%)				0.1674	0.041			1.00	0.002
Cecal acetate (mmol/kg)		25	92.64	18.476			46.434	0.35		
	Inulin (%)				-10.620	5.100			1.00	0.071
Colonic bifidobacteria (CFU/g)		14	7.39	0.253			0.455	0.74		
	BW (kg)				0.0163	0.004			1.00	0.003
	Inulin (%)				-0.005	0.001			1.00	0.002
Colonic enterobacteria (CFU/g)		19	6.82	0.210			0.857	0.57		
	Inulin-square (%)				-0.011	0.002			1.00	< 0.001
Fecal Lactobacilli (CFU/g)		26	10.30	0.247			0.764	0.79		
	BW (kg)				-0.045	0.008			1.22	< 0.001
	Inulin (%)				-0.255	0.078			1.22	0.004
Fecal bifidobacteria (CFU/g)		13	5.81	0.837			1.113	0.54		
	BW (kg)				0.068	0.0278			1.150	0.034
	Inulin (%)				0.181	0.0578			1.150	0.011
Fecal Escherichia coli (CFU/g)		19	2.59	1.249			0.216	0.80		
	BW (kg)				0.063	0.0149			6.10	< 0.001
	dietary CP (%)				0.134	0.0472			6.31	0.012
	Inulin (%)				-0.044	0.013			1.67	0.004

	AID of DM (%)		13	24.43	5.918			2.587	0.89		
		dietary CP (%)				3.098	0.333			1.00	< 0.001
	ATTD of CP (%)		15	89.59	2.149			1.238	0.89		
		dietary CP (%)				-0.284	0.115			1.02	0.029
_		Inulin (%)				-0.420	0.046			1.02	<.0001

^a BW, body weight; CIAD, coefficient of ileal apparent digestibility; CFU, colony forming units; CP, crude protein; CTTAD, coefficient of total tract apparent

732 digestibility; DM, dry matter; n_{Treat} = number of treatment means included; RMSE, root mean square error; SE, standard error; VIF, variance inflation factor.

Highlights

•

Meta-regressions showed potential of dietary inulin to lower gastric pH in weaned pigs.

Meta-regressions indicated an inhibitory effect of dietary inulin on Escherichia coli in feces.

Meta-regressions did not confirm a stimulatory effect of dietary inulin on intestinal lactobacilli and bifidobacteria throughout the intestinal tract.