



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Modelling facilitated transport in Polyvinyl amine membranes for CO<sub>2</sub> capture: insights from Molecular Dynamics and PC-SAFT EoS.

This is the submitted version (pre peer-review, preprint) of the following publication:

*Published Version:*

Modelling facilitated transport in Polyvinyl amine membranes for CO<sub>2</sub> capture: insights from Molecular Dynamics and PC-SAFT EoS / Rea Riccardo; Fayon Pierre; Kvam Odin; Sarkisov Lev; Venturi Davide; Giacinti Baschetti, Marco; De Angelis, Maria Grazia;. - STAMPA. - (2019), pp. 146-146. (Intervento presentato al convegno EWM 2019 tenutosi a Bastad nel 8-10 April, 2019).

*Availability:*

This version is available at: <https://hdl.handle.net/11585/701918> since: 2020-02-16

*Published:*

DOI: <http://doi.org/>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

*“This is a post-peer-review, pre-copyedit version of an abstract published in “EWM 2019: Book of Abstract”.*

© [2019] *Engineering With Membranes 2019*

# Modelling facilitated transport in Polyvinyl amine membranes for CO<sub>2</sub> capture: insights from Molecular Dynamics and PC-SAFT EoS.

R. Rea, P. Fayon, O. Kvam, L. Sarkisov, D. Venturi, M. Giacinti Baschetti, M. G. De Angelis

## **Background**

In the context of CO<sub>2</sub> removal from gas streams, the project NANOMEMC<sup>2</sup> ([www.nanomemc2.eu](http://www.nanomemc2.eu)) focuses on Facilitated Transport (FT) membranes based on Polyvinyl amine (PVAm). Such materials bear amine groups that, in presence of humidity, promote reactions that boost the transport of CO<sub>2</sub> while not affecting the other gases. A possible reaction route is shown in **Figure 1**.

Very few modelling studies are present in the literature concerning these fixed sites FT membranes despite their selectivity comparable to the most common absorption processes. Aim of the present work is to provide a detailed deep investigation on the transport properties of PVAm, to partially fill this lack, for the ternary system of CO<sub>2</sub>/H<sub>2</sub>O/PVAm. Molecular Dynamics (MD) and PC-SAFT<sup>1</sup> Equation of State (EoS) were used to achieve a reliable interpretation of the physical sorption process of CO<sub>2</sub> in such a complex, strongly polar environment.

## **Modelling and Simulating Strategies**

**PC-SAFT EoS:** H<sub>2</sub>O has been treated as 2B self-associating molecule (Huang and Radosz<sup>2</sup>); the induced (by H<sub>2</sub>O) association has been considered for the PVAm with 2 and 3 sites (2B and 3B scheme) while the CO<sub>2</sub> studied as 0 or 2 (2B) association sites, not allowing for the self association even in the latter case.

**MD simulations:** molecular models of solvated PVAm membranes were constructed by explicitly considering a binary mixture of short chains (20 monomers) of PVAm and H<sub>2</sub>O. Structural characteristics of these systems, such as pore size distribution and pore limiting diameter were obtained at different levels of hydration and correlated to self-diffusion of water in these structures. Chemical reaction(s) not explicitly included in the calculations.

## **Results**

**Figure 2** shows the main results achieved in terms of physical solubility of CO<sub>2</sub> in the swollen polymer while the self diffusion coefficient of H<sub>2</sub>O in the polymer matrix, from MD simulations as well as from the scaling model of Phillis (1989) and experimental data, is depicted in **Figure 3**.

## **Conclusions**

Physical sorption of CO<sub>2</sub> in swollen by water PVAm has been predicted by PC-SAFT EoS by two associating approach. The 2B scheme for CO<sub>2</sub> allows to elucidate the role of relative on the adsorption process. MD prediction on self diffusion of H<sub>2</sub>O in PVAm agree with both experimental data and Phillis scaling model.

NANOMEMC<sup>2</sup> is a project funded by the European Commission, Horizon 2020, G.A. 727734.

