

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Extended effective field theory of inflation

This is the submitted version (pre peer-review, preprint) of the following publication:

Published Version:

Extended effective field theory of inflation / Ashoorioon, Amjad; Casadio, Roberto; Cicoli, Michele; Geshnizjani, Ghazal; Kim, Hyung J.. - In: JOURNAL OF HIGH ENERGY PHYSICS. - ISSN 1029-8479. - ELETTRONICO. - 2018:2(2018), pp. 172.1-172.25. [10.1007/JHEP02(2018)172]

Availability:

This version is available at: <https://hdl.handle.net/11585/629536> since: 2020-01-17

Published:

DOI: [http://doi.org/10.1007/JHEP02\(2018\)172](http://doi.org/10.1007/JHEP02(2018)172)

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Extended Effective Field Theory of Inflation

**Amjad Ashoorioon,^{a,b} Roberto Casadio,^{a,c} Michele Cicoli,^{a,c,d}
Ghazal Geshnizjani,^{e,f} Hyung J. Kim^e**

^aINFN, Sezione di Bologna, viale Berti Pichat 6/2, 40127 Bologna, Italy

^bSchool of Physics, Institute for Research in Fundamental Sciences (IPM),
P.O.Box 19395-5531, Tehran, Iran

^cDipartimento di Fisica e Astronomia, Università di Bologna,
via Irnerio 46, 40126 Bologna, Italy

^dAbdus Salam ICTP, Strada Costiera 11, Trieste 34151, Italy

^eDepartment of Applied Mathematics, University of Waterloo
Waterloo, Ontario, N2L 3G1, Canada

^fPerimeter Institute for Theoretical Physics
31 Caroline St. N., Waterloo, ON, N2L 2Y5, Canada

E-mail: amjad.ashoorioon@bo.infn.it, roberto.casadio@bo.infn.it, mcicoli@ictp.it,
ggeshniz@uwaterloo.ca, h268kim@uwaterloo.ca

Abstract. We present a general framework where the effective field theory of single field inflation is extended by the inclusion of operators with mass dimension 3 and 4 in the unitary gauge. These higher dimensional operators introduce quartic and sextic corrections to the dispersion relation. We study the regime of validity of this extended effective field theory of inflation and the effect of these higher dimensional operators on CMB observables associated with scalar perturbations, such as the speed of sound, the amplitude of the power spectrum and the tensor-to-scalar ratio. Tensor perturbations remain instead, unaltered.

Keywords: Inflation, Effective field theory, Tensor perturbations, Modified dispersion relation, Consistency relation

Contents

1	Introduction	1
2	Extended EFT of inflation	3
2.1	Overview of the EFT of inflation	3
2.2	Lagrangian of the Extended EFT of inflation	5
3	Perturbations in the extended EFT of inflation	5
3.1	Tensor perturbations	5
3.2	Scalar perturbations	7
3.3	Scalar power spectrum	12
3.4	Tensor-to-scalar ratio and consistency relation	15
4	Cut-off and non-gaussianity	18
5	Conclusions	21

1 Introduction

The Effective Field Theory of Inflation (EFToI) [1] provides an alternative picture to understand perturbations in single field driven inflationary models. In this picture, one first fixes the gauge by taking constant time hypersurfaces to coincide with inflaton constant surfaces. In other words, the perturbations of the inflaton field are absorbed into the fluctuations of the metric. The inflaton fluctuations transform non-linearly under the time diffeomorphism but the non-linear sigma model describing it can always be UV completed into the theory of inflation with a linear representation of time diffeomorphism. The Goldstone mode, π , corresponding to breaking time diffeomorphism symmetry, describes the scalar perturbations around the FRW background. Therefore, the most generic action for inflation can be constructed out of the quantities that respect the remaining symmetries, *i.e.* the spatial diffeomorphisms. The coefficients of these terms can be time-dependent in principle.

This means including terms such as the metric component g^{00} and the extrinsic curvature of constant time hypersurfaces, $K_{\mu\nu}$, which are only spatially diffeomorphism invariant, in addition to standard 4-diffeomorphism invariant terms. It was shown in [1] that, around a FRW background, this reduces to including functions of time, g^{00} and perturbations of the extrinsic curvature, $\delta K_{\mu\nu}$.

One can then make the Goldstone mode explicit by applying the Stückelberg technique and study the physics of the Goldstone mode at energies where the mixing with gravitons can be neglected. In this framework, the standard slow-roll inflationary model corresponds to adding only operators with time dependence and at most linear in terms of g^{00} (mass dimension zero), before performing the Stückelberg trick,. Higher dimensional operators quantify deviations from the slow-roll inflation. On the other hand, in terms of π and in the decoupling limit, the slow-roll Lagrangian will include dimension 4 operators from the kinetic terms of the canonically normalised π . With higher order or mass dimensional operators in the unitary gauge, dimension 6 operators from cubic interactions of the Goldstone modes are generated. Due to the non-linear nature of the time diffeomorphism, once it is restored, the spatially

diffeomorphism invariant operators will lead to a non-linear dependence on the Goldstone mode. This in turn leads to non-vanishing higher point correlation functions. Therefore, the coefficients of these operators can be constrained by measuring the corresponding correlation functions.

The above approach is borrowed from particle physics since also the Standard Model is described through the lowest dimension operators that are compatible with the symmetries of the system. Then higher dimensional operators describe deviations from the vanilla case and quantify the emergence of new physics. Similarly, in the EFToI, higher dimensional operators can reproduce various interesting inflationary models, such as DBI inflation [2–4], with speed of sound different from one, or Ghost Inflation [5], with leading spatial gradient term proportional to $(\nabla^2 \pi)^2$ [1]. In particular, as shown in [1], reducing the speed of sound naturally strengthens the cubic interaction, which in turn enhances the non-gaussianity.

In the EFToI, these higher dimensional operators can also change the dispersion relation. For example, the inclusion of $\delta K_{\mu\nu}^2$ operators modifies the dispersion relation to $\omega^2 \sim k^4$ (similar to Ghost Inflation). Including additional higher dimensional operators of the form $(\nabla^{(n-2)} K_{\mu\nu})^2$ with $n \geq 3$, when they are dominant, would modify the dispersion relation even further to $\omega^2 \sim k^{2n}$.

However, the authors of [1] did not include operators beyond mass dimension 2, since they claimed that these new terms would not be compatible with a sensible EFT description. The reason is the fact that in such theories, when $E \rightarrow sE$, the energy scaling dimension for π would be $\pi \rightarrow s^{-1/2+3/2n} \pi$ and so the scaling behaviour of cubic operators like $\dot{\pi}(\nabla\pi)^2$ would be $(7-3n)/(2n)$. This would imply that for $n \geq 3$ this operator would become *relevant*, whereas for $n < 3$ it is *irrelevant* at low energies. Hence a theory with a dispersion relation of the form $\omega^2 \sim k^6$ would have an IR strong coupling cut-off Λ_6^{IR} which would in general make this theory not a controllable EFT.

However, we argue here that this may not always be the case. Our reason is that a dispersion relation of the form $\omega^2 \sim k^6$ would not hold up all the way to low energy scales. It is true that the scaling power of $\dot{\pi}(\nabla\pi)^2$ is $-1/3$, but the scaling power of $(\nabla^2 \pi)^2$ and $(\nabla\pi)^2$ are respectively $-2/3$ and $-4/3$. This implies that the dispersion relation would also change from $\omega^2 \sim k^6$ to $\omega^2 \sim k^4$ at some energy scale $\Lambda_{\text{dis}}^{4-6}$. Hence, if the coefficients of the higher dimensional operators are tuned such that $\Lambda_6^{\text{IR}} \ll \Lambda_{\text{dis}}^{4-6}$, the $\omega^2 \sim k^6$ theory remains weakly coupled throughout its evolution down to low energies. On top of this, one has to require that the scale Λ_4 , where the $\omega^2 \sim k^4$ theory becomes strongly coupled, is not below $\Lambda_{\text{dis}}^{4-6}$. This is guaranteed if $\Lambda_6^{\text{IR}} \ll \Lambda_{\text{dis}}^{4-6} \ll \Lambda_4$. Similarly, one has to require that the standard $\omega^2 \sim k^2$ theory does not become strongly coupled at a scale Λ_2 which is below the scale $\Lambda_{\text{dis}}^{2-4}$ where the dispersion relation becomes dominated by the quartic term: $\Lambda_{\text{dis}}^{2-4} \ll \Lambda_2$ [6].

Under these conditions, the EFT for the perturbations remains weakly coupled all the way from low energy to the $\omega^2 \sim k^6$ regime. Note that in addition, this framework is only valid on energy scales below Λ_b , where time diffeomorphisms get spontaneously broken by the inflaton background. Thus, we also need to require $\Lambda_{\text{dis}}^{4-6} \ll \Lambda_b$. This is because at Λ_b the scalar perturbations become of the same order of magnitude of the background which, therefore, cannot be integrated out to leave an EFT for the fluctuations only. The cut-off of the EFT for the fluctuations Λ_{cutoff} is therefore the minimum between Λ_b and the UV strong coupling scale of the $\omega^2 \sim k^6$ theory, Λ_6^{UV} , associated with some higher derivative operators which remain irrelevant.

Strictly speaking, the cosmological experiment is done at energies of order the Hubble scale H where horizon exit takes place. Hence, in order for the EFT approach to be under

control, one would need just to require $H \ll \Lambda_{\text{cutoff}}$ and a horizon crossing that occurs in the region where the dispersion relation takes the standard quadratic form. However, the goal of this paper is to provide a consistent theoretical framework to motivate the emergence of modified dispersion relations which, as previously shown, can lead to (super-excited) non-Bunch-Davis initial conditions with interesting implications for CMB observables [7]. Thus we need to control the EFT also in the high energy $\omega^2 \sim k^6$ regime which describes the behavior of the perturbations deep inside the horizon. These perturbations are born in the vacuum of k^6 theory but then become excited states due to the transitions to the quartic and finally to the standard quadratic regime for the dispersion relation which happen before horizon crossing. Therefore, in order to be able to describe consistently the whole evolution of these modes from deep inside the horizon to super-horizon scales, we need to be able to trust the EFT for the fluctuations up to the $\omega^2 \sim k^6$ regime.

In this paper we shall therefore include higher order corrections which in the unitary gauge correspond to operators with mass dimension 3 and 4 that were so far neglected in the literature. Hence we shall call our framework Extended Effective Field Theory of Inflation (EEFToI). As explained above, these higher dimensional terms modify the dispersion relation and, depending on the sign and magnitude of their coefficients, can have important implications for scalar perturbations. We shall focus only on modifications of the equation of motion for the Goldstone boson π by higher order terms, and so we shall consider operators which include at least one factor of $\delta K_{\mu\nu}$ in their expression. One can write down other operators with the same mass dimensions as the one we focus on below, which would not affect the two point function for scalar perturbations. We also drop operators of the form $(1 + g^{00})^n$ with $n \geq 3$ since, contrary to $(1 + g^{00})^2$, they do not change the speed of sound for π .

The outline of this paper is as follows. In Sec. 2, we first briefly review the formalism of the EFToI and then we present the EEFToI by writing down all operators with mass dimension three and four that can add a correction proportional to k^6 in the dispersion relation. In Sec. 3, we discuss the implications of these higher dimensional operators for tensor and scalar perturbations. In particular, after identifying the allowed operators, we focus on the healthy region of parameter space for π and explore various scenarios that can arise. We study the scalar power spectrum in these scenarios, depending on the sign of the coefficients of the quartic and sextic corrections to the dispersion relation. We also analyse the implications for the tensor-to-scalar ratio. In Sec. 4, we comment on the cut-off of the EEFToI and the size of non-gaussianity, while in Sec. 5, we conclude and give directions for future research.

2 Extended EFT of inflation

2.1 Overview of the EFT of inflation

We start by reviewing the EFToI framework which allows one to write down the most generic action for single scalar field models on a quasi de Sitter background. The idea is that even though the action is invariant under all diffeomorphisms (4-diffs), the background solution provides a preferred time foliation. By restricting time hypersurfaces to constant inflaton surfaces (the unitary gauge), the action has to satisfy only 3-dimensional spatial diffeomorphism (3-diff) invariance. Therefore, one can start by writing the most general 3-diff invariant action around the FRW metric and the effect of different terms. This implies, in addition to standard 4-diff invariant metric terms, including g^{00} and pure functions of time $f(t)$ that are

also scalars under 3-diffs. We can also include terms depending on the extrinsic curvature of the constant time hypersurfaces, $K_{\mu\nu}$, as this is a tensor under spatial diffs. It turns out that any additional 3-diff invariant object, such as the 3-dimensional covariant derivative, can be rewritten in terms of the quantities already included. It can be shown that the resulting Lagrangian around flat FRW reduces to [1]

$$\mathcal{L} = M_{\text{Pl}}^2 \left[\frac{1}{2} R + \dot{H} g^{00} - \left(3 H^2 + \dot{H} \right) \right] + \sum_{m \geq 2} \mathcal{L}_m(g^{00} + 1, \delta K_{\mu\nu}, \delta R_{\mu\nu\rho\sigma}, \nabla_\mu; t), \quad (2.1)$$

where \mathcal{L}_m represent functions of order m in $g^{00} + 1$, $\delta K_{\mu\nu}$ and $\delta R_{\mu\nu\rho\sigma}$. Furthermore, the pure time dependent term and coefficient of g^{00} have become functions of H and \dot{H} by satisfying the Friedmann equations in the FRW limit.¹

Once the generic action in unitary gauge has been written, the gauge condition can be relaxed by allowing the time transformation $t \rightarrow t + \xi^0(x^\mu)$. Since the action is no more restricted to a particular time slicing, time diffeomorphism invariance has to be restored again. This can be achieved by substituting $\xi_0(x^\mu)$ with a field $-\pi(x^\mu)$ and requiring that it shifts as $\pi(x^\mu) \rightarrow \pi(x^\mu) - \xi_0(x^\mu)$ under time diffeomorphisms. Note that when we perform $t \rightarrow t + \xi^0(x^\mu)$, we no longer expect the perturbations in inflaton field, ϕ , to be zero. In fact, by introducing the Goldstone boson we are representing these perturbations, since $\delta\phi = -\dot{\phi}_0 \xi^0$ and therefore $\pi = \delta\phi/\dot{\phi}_0$.

This action is very complicated in general. However, the advantage of using this approach is that, when the generalised slow roll approximation is valid, one can ignore all the metric perturbations in the action. Basically, if the typical scale of the time dependence of the coefficients in the unitary gauge is much longer than the Hubble time, the theory effectively decouples from gravity and it is possible to compute the power spectrum neglecting metric perturbations in the action.

For instance, if we implement this procedure for the action (2.1), assuming $\mathcal{L}_m = 0 \forall m$, we obtain

$$\mathcal{L}_{\text{slow-roll}} = -M_{\text{Pl}}^2 \dot{H} \left(\dot{\pi}^2 - \frac{(\partial\pi)^2}{a^2} \right). \quad (2.2)$$

One can show, using proper gauge transformations, that π is related to the conserved quantity ζ by $\zeta = -H\pi$. Substituting ζ in the above action reduces it to the standard slow-roll inflationary action for ζ ². Since we are interested in deviations from the standard slow-roll model, we will turn on the coefficients of the operators in \mathcal{L}_m . In the EFToI, focusing only on the terms that contribute to the quadratic action of π and can change the dispersion relation up to quartic order, the unitary gauge action is

$$\mathcal{L}_{\text{EFToI}} = \mathcal{L}_{\text{slow-roll}} + \mathcal{L}_2, \quad (2.3)$$

with:

$$\mathcal{L}_2 = \frac{M_2^4}{2!} (g^{00} + 1)^2 + \frac{\bar{M}_1^3}{2} (g^{00} + 1) \delta K_\mu^\mu - \frac{\bar{M}_2^2}{2} (\delta K_\mu^\mu)^2 - \frac{\bar{M}_3^2}{2} \delta K_\nu^\mu \delta K_\mu^\nu. \quad (2.4)$$

¹In the original EFToI formalism [1], only the quadratic operators built out of $\delta K_{\mu\nu} = K_{\mu\nu} - K_{\mu\nu}^{\text{FRW}}$ were studied. As we will see, dropping higher order operators is not always necessary and one can extend the EFToI correspondingly.

²The correspondence is up to a mass correction in higher orders of slow roll parameters. In quasi de Sitter space, π gets a small mass due to time-dependent background, which has to be taken into account to get the exact correspondence

The first term, $(g^{00} + 1)^2$, is the operator that modifies the speed of sound for scalar perturbations from the speed of light. Noting that $(g^{00} + 1)$ has zero mass dimension, powers of $(g^{00} + 1)^n$ with $n \geq 3$ could also be included which result to more general K-inflationary models. In this work, without loss of generality, we only include the quadratic term in $(g^{00} + 1)$ as it explicitly modifies the speed of sound for scalar perturbations. The mass dimension 1 term, $(g^{00} + 1)\delta K^\mu_\mu$, is not symmetric under time reversal, and was already analysed in [1, 8]. Finally $\frac{\bar{M}_2^2}{2}(\delta K^\mu_\mu)^2$ and $\frac{\bar{M}_3^2}{2}\delta K^\mu_\nu\delta K^\nu_\mu$ are the operators that lead to generalized Ghost Inflation with a quartic correction to the dispersion relation [1, 5].

2.2 Lagrangian of the Extended EFT of inflation

We now proceed to introduce the Lagrangian of the EEFToI. We supplement the unitary gauge action of the EFToI with operators of mass dimension 3 and 4 constructed out of δK^ν_μ , ∇_μ and g^{00} so that the Lagrangian of the EEFToI (again keeping only terms which contribute to quadratic action for π) becomes

$$\mathcal{L}_{\text{EEFToI}} = \mathcal{L}_{\text{EFToI}} + \mathcal{L}_{2,d3} + \mathcal{L}_{2,d4}, \quad (2.5)$$

with:

$$\begin{aligned} \mathcal{L}_{2,d3} + \mathcal{L}_{2,d4} = & \frac{\bar{M}_4}{2} \nabla^\mu g^{00} \nabla^\nu \delta K_{\mu\nu} - \frac{\delta_1}{2} (\nabla_\mu \delta K^{\nu\gamma})(\nabla^\mu \delta K_{\nu\gamma}) - \frac{\delta_2}{2} (\nabla_\mu \delta K^\nu_\nu)^2 \\ & - \frac{\delta_3}{2} (\nabla_\mu \delta K^\mu_\nu)(\nabla_\gamma \delta K^{\gamma\nu}) - \frac{\delta_4}{2} \nabla^\mu \delta K_{\nu\mu} \nabla^\nu \delta K^\sigma_\sigma, \end{aligned} \quad (2.6)$$

where M_i , \bar{M}_i and δ_i 's are free (time dependent) coefficients and the sign of each term is also a priori free. We remark that the mass dimension 3 operator $\nabla^\mu g^{00} \nabla^\nu \delta K_{\mu\nu}$ is not symmetric under time reversal and δ_i 's are the coefficients of dimension 4 operators.

Our motivation for adding such operators is that they lead to sixth order corrections to the dispersion relation. The reason advocated in [1] for discarding these operators is that sixth order corrections to the dispersion relation would make higher derivative operators relevant, signaling the presence of an IR strong coupling regime. However, as explained in Sec. 1 and as we will discuss in more depth in Sec. 4, one can still have a sensible EFT description if the IR strong coupling scale of the $\omega^2 \sim k^6$ theory is below the scale $\Lambda_{\text{dis}}^{4-6}$ where the dispersion relation becomes dominated by the quartic term.

We emphasise here that in this EFT approach to inflation, the Lorentz symmetry is not preserved. In particular, the mass dimension and energy scaling dimension of the operators depending on δK do not match. We should also add that we have only kept terms that can modify the action for scalar perturbations at second order. There are other operators, e.g. proportional to $\bar{M} \delta K^3$, that do not contribute to the linear equation of motion of the Goldstone boson π . For reasons that will become clear momentarily, we first focus on the implications of the mass dimension 3 and 4 operators on the action for tensor perturbations. Then we will get back to the action for the Goldstone mode π derived from the unitary gauge action in the EEFToI.

3 Perturbations in the extended EFT of inflation

3.1 Tensor perturbations

To study the tensor perturbations, we perturb the spatial part of the metric as follows

$$g_{ij} = a^2(\delta_{ij} + \gamma_{ij}), \quad (3.1)$$

where γ_{ij} is transverse and traceless (summation over the repeated index i is assumed),

$$\gamma_{ii} = 0, \quad \partial_i \gamma_{ij} = 0, \quad (3.2)$$

and we will expand the action (2.5) up to second order in γ_{ij} . As pointed out in the original EFToI paper [1], from the mass dimension zero up to mass squared operators, only $-\frac{\bar{M}_3^2}{2} \delta K^\mu_\nu \delta K^\nu_\mu$ affects the equation of motion for γ_{ij} , and adds the following contribution to the action for tensor perturbations

$$S^{\bar{M}_3^2} = -\frac{\bar{M}_3^2}{8} \int a^3 dt d^3x \partial_0 \gamma_{ij} \partial_0 \gamma_{ij}. \quad (3.3)$$

The action (3.3), together with the terms describing the usual massless scalar field contribution for tensor perturbations,

$$S_{\text{EH}} = \frac{M_{\text{Pl}}^2}{8} \int dt d^3x a^3 (\partial_0 \gamma_{ij} \partial_0 \gamma_{ij} - a^{-2} \partial_l \gamma_{ij} \partial_l \gamma_{ij}), \quad (3.4)$$

produces the following modified equation of motion for tensor perturbations,

$$p_k'' + \left(c_{\text{T}}^2 k^2 - \frac{a''}{a} \right) p_k = 0, \quad (3.5)$$

where $p_k \equiv a \gamma_k$, the subscript k refers to the Fourier transform of the two helicities of the gravitons, and

$$c_{\text{T}}^2 = \left(1 - \frac{\bar{M}_3^2}{M_{\text{Pl}}^2} \right)^{-1}. \quad (3.6)$$

In order to avoid superluminal propagation for tensor perturbations, one must then have $\bar{M}_3^2 \leq 0$. Larger values of $|\bar{M}_3|$ correspond to smaller speed of sound and, for $|\bar{M}_3| \rightarrow M_{\text{Pl}}$, the speed of gravitational waves $c_{\text{T}} \rightarrow 1/\sqrt{2}$.

For $\bar{M}_3^2 > 0$, the speed of gravitation waves is superluminal. Lorentz invariant EFTs with this property have been argued to be non-local and not embeddable in a local quantum field theory or string theory [9]. However, [10] argues that such models, despite having superluminal propagation, do not lead to any violation of causality. If $\bar{M}_3^2 > 0$ is taken as a legitimate choice, larger values of \bar{M}_3^2 increase the value of speed of sound and in fact this value diverges for $\bar{M}_3^2 \rightarrow M_{\text{Pl}}^2$.

Now let us focus on the new operators with mass dimension 3 and 4. The term $\frac{\delta_1}{2} (\nabla_\mu \delta K^{\nu\gamma}) (\nabla^\mu \delta K_{\nu\gamma})$, up to second order in γ_{ij} , yields

$$\frac{\delta_1}{2} \left[-\frac{H^2}{2} (\partial_0 \gamma_{ij})^2 + \frac{1}{4a^2} (\partial_{0m}^2 \gamma_{ij})^2 - \frac{1}{4} (\partial_0^2 \gamma_{ij})^2 \right], \quad (3.7)$$

where i, j and m are spatial indices and summation over m is assumed. Other operators do not contribute to the Lagrangian of tensor perturbations once the transverse traceless condition on γ_{ij} is imposed. The appearance of $\partial_0^2 \gamma_{ij}$ in the Lagrangian density (3.7) leads to ghost instability for the tensor perturbations. Thus the requirement of having no ghosts in the tensor sector of the theory translates into

$$\delta_1 = 0. \quad (3.8)$$

Therefore, dispersion relation of the primordial tensor fluctuations remains resilient to changes in the scalar and gravitational sector in the EEFToI, like it was shown originally in [11].

3.2 Scalar perturbations

The Goldstone boson can explicitly appear in the action via the Stückelberg trick. In this procedure, the variation of the metric with broken time diffeomorphism, $t \rightarrow \tilde{t} = t + \xi^0(x)$, $x \rightarrow \tilde{x} = \vec{x}$, is obtained and $\xi^0(x(\tilde{x}))$ is everywhere replaced by $-\tilde{\pi}(\tilde{x})$ in the transformed action. Then $\pi(x)$ is assumed to transform non-linearly to $\tilde{\pi}(\tilde{x}) = \pi(x) - \xi^0(x)$, which guarantees that the action remains invariant under diffs to all orders [1]. Evaluating the action explicitly for π in Fourier space, Ref. [8] found out that, in the decoupling limit corresponding to $\dot{H} \rightarrow 0$ while $M_p^2 \dot{H}$ is fixed, the EFToI Lagrangian (2.3) in the unitary gauge leads to the following second order Lagrangian in the π -gauge:

$$\begin{aligned} \mathcal{L}_{\text{EFToI}}^{(\pi)} = & M_p^2 \dot{H} (\partial_\mu \pi)^2 + 2M_2^4 \dot{\pi}^2 - \bar{M}_1^3 H \left(3\dot{\pi}^2 - \frac{(\partial_i \pi)^2}{2a^2} \right) - \frac{\bar{M}_2^2}{2} \left(9H^2 \dot{\pi}^2 - 3H^2 \frac{(\partial_i \pi)^2}{a^2} \right. \\ & \left. + \frac{(\partial_i^2 \pi)^2}{a^4} \right) - \frac{\bar{M}_3^2}{2} \left(3H^2 \dot{\pi}^2 - H^2 \frac{(\partial_i \pi)^2}{a^2} + \frac{(\partial_j^2 \pi)^2}{a^4} \right), \end{aligned} \quad (3.9)$$

whereas, following the same procedure, the mass dimension 3 parity-violating operator in (2.6) in the same limit $\dot{H} \rightarrow 0$ yields

$$\mathcal{L}_{2,d_3}^{(\pi)} = \frac{\bar{M}_4}{2} \left(\frac{k^4 H \pi^2}{a^4} + \frac{k^2 H^3 \pi^2}{a^2} - 9H^3 \dot{\pi}^2 \right). \quad (3.10)$$

This operator only yields a quartic correction to the dispersion relation. On the other hand, the mass dimension 4 operators in the EEFToI action (2.5) give ³

$$\begin{aligned} \mathcal{L}_{2,d_4}^{(\pi)} = & -\frac{1}{2} \delta_1 \left(\frac{k^6 \pi^2}{a^6} - \frac{3H^2 k^4 \pi^2}{a^4} - \frac{k^4 \dot{\pi}^2}{a^4} + \frac{4H^4 k^2 \pi^2}{a^2} - 6H^4 \dot{\pi}^2 - 3H^2 \ddot{\pi}^2 \right) \\ & -\frac{1}{2} \delta_2 \left(\frac{k^6 \pi^2}{a^6} + \frac{H^2 k^4 \pi^2}{a^4} - \frac{k^4 \dot{\pi}^2}{a^4} + \frac{6H^4 k^2 \pi^2}{a^2} - 9H^2 \ddot{\pi}^2 \right) \\ & -\frac{1}{2} \delta_3 \left(\frac{k^6 \pi^2}{a^6} + \frac{3H^2 k^4 \pi^2}{a^4} + \frac{H^2 k^2 \dot{\pi}^2}{a^2} - 9H^4 \dot{\pi}^2 \right) \\ & -\frac{1}{2} \delta_4 \left(\frac{k^6 \pi^2}{a^6} + \frac{H^2 k^4 \pi^2}{2a^4} + \frac{9H^4 k^2 \pi^2}{2a^2} + \frac{3H^2 k^2 \dot{\pi}^2}{a^2} + \frac{27}{2} H^4 \dot{\pi}^2 \right). \end{aligned} \quad (3.11)$$

The terms $-\frac{\delta_1}{2} (\nabla_\mu \delta K^{\nu\gamma}) (\nabla^\mu \delta K_{\nu\gamma})$ and $-\frac{\delta_2}{2} (\nabla_\mu \delta K^\nu{}_\nu)^2$ produce $\ddot{\pi}^2$. According to Ostrogradski's theorem [12], higher time derivatives usually lead to ghost instabilities. In principle such a ghost term could be avoided if $\delta_1 = -3\delta_2$. However, as noticed in the previous section, one has to set $\delta_1 = 0$ in order to avoid ghosts in the tensor sector, which enforces

$$\delta_1 = \delta_2 = 0. \quad (3.12)$$

³In [7], in the definition for $\delta K_{\mu\nu} \equiv K_{\mu\nu} - K_{\mu\nu}^{(0)}$, we had mistakenly subtracted the FRW result $K_{\mu\nu}^{(0)} = -a^2 H \delta_{ij} \delta_\mu^i \delta_\nu^j$ from the extrinsic curvature of constant t spatial surfaces. However, this would not produce a tensorial structure for $\delta K_{\mu\nu}$. Following [1], one has to instead subtract $K_{\mu\nu}^{(0)} = -H h_{\mu\nu}$, which reduces to the FRW result when the perturbations vanish and guarantees that $\delta K_{\mu\nu}$ is covariant. The difference in the obtained results for the action of Goldstone boson is due to this. We thank P. Creminelli for clarification of this issue to us. We take the opportunity to correct the mistake/typo in [1] in defining $K_{\mu\nu}^{(0)}$ as $a^2 H h_{\mu\nu}$. There seems to be an extra factor of $-a^2$ in their expression.

Therefore, only the terms $\frac{\delta_3}{2} \nabla_\mu \delta K^\mu_\nu \nabla_\gamma \delta K^{\gamma\nu}$ and $\frac{\delta_4}{2} \nabla^\mu \delta K_{\nu\mu} \nabla^\nu \delta K^\sigma_\sigma$ are allowed.

We shall now derive the equation of motion for π using the total EFTToI Lagrangian in the π language and in the decoupling limit, which is the sum of (3.9), (3.10) and (3.11),

$$\mathcal{L}_{\text{EFTToI}}^{(\pi)} = \mathcal{L}_{\text{EFTToI}}^{(\pi)} + \mathcal{L}_{2,d_3}^{(\pi)} + \mathcal{L}_{2,d_4}^{(\pi)} \quad (3.13)$$

Before that, we briefly discuss the equation of motion corresponding to the EFTToI Lagrangian. Variation of $\mathcal{L}_{\text{EFTToI}}^{(\pi)}$, yields the same equation of motion previously derived in [8],

$$A_0 \ddot{\pi}_k + (B_0 + 3HC_0) \dot{\pi}_k + \left(E_0 \frac{k^4}{a^4} + D_0 \frac{k^2}{a^2} \right) \pi_k = 0 \quad (3.14)$$

with

$$\begin{aligned} A_0 &= -2M_{\text{Pl}}^2 \dot{H} + 4M_2^4 - 6\bar{M}_1^3 H - 9H^2 \bar{M}_2^2 - 3H^2 \bar{M}_3^2 \\ B_0 &= -6\bar{M}_1^3 \dot{H} - 18\dot{H} H \bar{M}_2^2 - 6\dot{H} H \bar{M}_3^2 \\ C_0 &= -2M_{\text{Pl}}^2 \dot{H} + 4M_2^4 - 6\bar{M}_1^3 H - 9H^2 \bar{M}_2^2 - 3H^2 \bar{M}_3^2 \\ D_0 &= -2M_{\text{Pl}}^2 \dot{H} - \bar{M}_1^3 H - 3H^2 \bar{M}_2^2 - \bar{M}_3^2 H^2 \\ E_0 &= \bar{M}_2^2 + \bar{M}_3^2, \end{aligned} \quad (3.15)$$

where the mixing with gravity has been neglected. Noting that the canonical $\pi_c \sim \sqrt{A_0} \pi$ and $\delta g_c^{00} \sim M_{\text{Pl}} \delta g^{00}$, the mixing energy between gravity and the Goldstone mode π can be neglected at energies

$$E > E_{\text{mix}} \sim \frac{\sqrt{A_0}}{M_{\text{Pl}}}, \quad (3.16)$$

which is known as the equivalence theorem. Namely, the perturbations of the longitudinal mode decouple from the perturbations of the metric at energies higher than E_{mix} . Being able to compute the correlation functions at horizon crossing corresponds to assuming $H > E_{\text{mix}}$. In the regime where $M_{\text{Pl}}^2 \dot{H}$ is bigger than the other contributions to A_0 , $E_{\text{mix}} \sim \epsilon^{1/2} H$, and so the decoupling limit is guaranteed by the slow-roll condition $\epsilon \ll 1$. On the other hand if M_2^4 is bigger than the other parameters in A_0 , $E_{\text{mix}} \sim M_2^2/M_{\text{Pl}}$, and decoupling happens if $M_2^2 < M_{\text{Pl}} H$. These conditions were obtained in [1] too. Other limiting cases suggest $\bar{M}_1^{3/2} < M_{\text{Pl}} H^{1/2}$, $\bar{M}_2 < M_{\text{Pl}}$ and $\bar{M}_3 < M_{\text{Pl}}$.

In terms of the conformal time, $d\tau \equiv dt/a$ and through a change of variable $u_k = a\pi_k$, the equation of motion (3.14) takes the following form

$$u_k'' + \frac{B_0 + 3HC_0 - 3A_0 H}{A_0} a u_k' + \left(\frac{D_0}{A_0} k^2 + \frac{E_0}{A_0} k^4 a^2 - \frac{a''}{a} \right) u_k = 0. \quad (3.17)$$

Noting that $B_0 + 3HC_0 - 3A_0 H = -6\dot{H}(\bar{M}_1^3 + 3H^2 \bar{M}_2^2 + H \bar{M}_3^2)$, in the limit $\dot{H} \rightarrow 0$, this equation is reduced to

$$u_k'' + \left(\frac{D_0}{A_0} k^2 + \frac{E_0}{A_0} k^4 a^2 - \frac{a''}{a} \right) u_k = 0. \quad (3.18)$$

which has the same form derived in [8].

If we now include the higher dimensional corrections $\mathcal{L}_{4,d_3}^{(\pi)}$ and $\mathcal{L}_{4,d_4}^{(\pi)}$ to the total Lagrangian (3.13), the equation of motion for π becomes

$$A_1 \ddot{\pi}_k + B_1 \dot{\pi}_k + \left(C_1 \frac{k^6}{a^6} + D_1 \frac{k^4}{a^4} + F_1 \frac{k^2}{a^2} \right) \pi_k = 0. \quad (3.19)$$

Defining

$$F_0(k, \tau) = \frac{9}{2}\delta_3 - \frac{27}{4}\delta_4 - \frac{k^2}{2a^2H^2}(\delta_3 + 3\delta_4) - \frac{9}{2H}\bar{M}_4, \quad (3.20)$$

the coefficients of the equation of motion (3.19) read

$$\begin{aligned} A_1 &= A_0 + 2H^4 F_0(k, \tau), \\ B_1 &= B_0 + 3HC_0 + H^5 \left[6F_0(k, \tau) + \frac{2k^2}{a^2H^2}(\delta_3 + 3\delta_4) \right], \\ C_1 &= \delta_3 + \delta_4, \\ D_1 &= E_0 + H^2 \frac{\delta_4}{2} + 3H^2\delta_3 - \bar{M}_4H, \\ F_1 &= D_0 + 3H^4 \left(\delta_3 + \frac{3}{2}\delta_4 \right) - \bar{M}_4H^3, \end{aligned} \quad (3.21)$$

where we have assumed that the time-dependence of the coefficients is slow compared to the Hubble time. This means that the terms coming from the Taylor expansion of the coefficients are small.

Expressing the equation of motion for $u_k = a \pi_k$, in conformal time and again in the limit $\dot{H} \rightarrow 0$, we obtain

$$u_k'' + \frac{2k^2H^3(\delta_3 + 3\delta_4)}{aA_1} u_k' + u_k \left(\frac{C_1 k^6}{A_1 a^4} + \frac{D_1 k^4}{A_1 a^2} + \frac{F_1}{A_1} k^2 - \frac{a''}{a} \right) = 0. \quad (3.22)$$

In a de Sitter space, where $a = -1/(H\tau)$, the above equation reads

$$\begin{aligned} u_k'' + \frac{G_3 \tau k^2 \delta_4}{G_1 + G_2 \tau^2 k^2} u_k' + u_k \left(\frac{F_2}{G_1 + G_2 \tau^2 k^2} k^2 + \frac{D_2 k^2}{G_1 + G_2 k^2 \tau^2} k^2 \tau^2 \right. \\ \left. + \frac{C_2 k^2}{G_1 + G_2 \tau^2 k^2} k^4 \tau^4 - \frac{2}{\tau^2} \right) = 0 \end{aligned} \quad (3.23)$$

where

$$\begin{aligned} G_1 &\equiv A_0 + 9H^4 \left(\delta_3 - \frac{3}{2}\delta_4 \right) - 9\bar{M}_4H^3 \\ G_2 &\equiv -H^4 (\delta_3 + 3\delta_4) \\ G_3 &\equiv -2H^4 (\delta_3 + 3\delta_4) = 2G_2 \\ F_2 &\equiv F_1 \\ D_2 &\equiv D_1 H^2 \\ C_2 &\equiv C_1 H^4 \end{aligned} \quad (3.24)$$

and

$$A_1(k, \tau) = G_1 + G_2 k^2 \tau^2. \quad (3.25)$$

The speed of propagation of the perturbations in the intermediate IR, i.e. for $1 < \kappa\tau < \sqrt{G_1/G_2}$, can now be read out from eq. (3.23) to be

$$c_s^2 = \frac{F_2}{G_1}. \quad (3.26)$$

In terms of a dimensionless variable, $x \equiv k\tau$, the equation of motion (3.23) takes the form

$$\frac{d^2 u_k}{dx^2} + \frac{G_3 x}{G_1 + G_2 x^2} \frac{du_k}{dx} + \left(\frac{F_2}{G_1 + G_2 x^2} + \frac{D_2 x^2}{G_1 + G_2 x^2} + \frac{C_2 x^4}{G_1 + G_2 x^2} - \frac{2}{x^2} \right) u_k = 0. \quad (3.27)$$

In general, due to non-polynomial scale dependence of the dispersion relation, the sound wave analogy does not necessarily apply. A variety of scenarios can in principle arise, depending on the size and sign of various coefficients. For example, if G_1 and G_2 are positive, the modes start with $c_s^2 \sim 0$ in infinite past and gradually, achieve a constant c_s^2 given by F_2/G_1 as they finally exit the horizon for $|k\tau| \ll 1$. If one of G_1 or G_2 (but not both) is negative, there could be a singular time in the evolution of each mode from inside the horizon when the speed of sound becomes infinite. It is also possible that both the speed of sound and the coefficient of the quartic correction to the dispersion relation are negative. These instabilities can in principle be avoided in the UV by a positive sextic correction to the dispersion relation. Even though such scenarios are all interesting, we will not focus on them in this paper.

In the following, we set $\delta_3 = -3\delta_4$, so that the equation of motion (3.27) simplifies to ⁴

$$\frac{d^2 u_k}{dx^2} + \left(\frac{F_2}{G_1} + \frac{D_2}{G_1} x^2 + \frac{C_2}{G_1} x^4 - \frac{2}{x^2} \right) u_k = 0. \quad (3.28)$$

If $F_2/G_1 > 0$, as assumed here, our definition of the speed of sound for scalar perturbations (3.26) is real and well-defined:

$$c_s^2 = \frac{F_2}{G_1}. \quad (3.29)$$

If $F_2/G_1 < 0$, one deals with the sort of dispersion relation that comes up in Ghost Inflation-like scenarios [5, 13]. This will not be something that we focus on in this investigation. Moreover, in order to avoid potential issues related to superluminal propagation in the IR and a unitary UV completion, we assume that [9]

$$0 \leq c_s^2 \leq 1. \quad (3.30)$$

The first inequality could be satisfied if F_2 and G_1 have the same sign. Note that it is also possible to have $c_s^2 = 1$ if

$$4M_2^4 - 5\bar{M}_1^3 H - 6H^2 \bar{M}_2^2 - 2H^2 \bar{M}_3^2 - 9H^4 \delta_4 - 8\bar{M}_4 H^3 = 0 \quad (3.31)$$

eventhough, higher-point interactions will not be zero. This shows that, although in the EFToI one tries to quantify the deviations from the standard slow-roll model with the operators \mathcal{L}_m in (2.1), there is still the possibility that one flows back close to the regime of slow-roll inflation with $c_s = 1$ via appropriate tuning of the coefficients of the various higher dimensional operators. This result was also true in the EFToI with δK^2 terms in the unitary gauge Lagrangian. Although one expects to see an enhancement in the non-gaussianity amplitudes due to the modified dispersion relation.

⁴With $\delta_3 \neq -3\delta_4$, the mixing between gravity and the Goldstone boson becomes time-dependent for each mode. Even though at horizon crossing, one can tune the parameters to make $H > E_{\text{mix}}$, when the physical momentum of the mode $k/a > M_{\text{Pl}}(\delta_3 + 3\delta_4)^{-1/2}$, the mixing with gravity cannot be neglected. At such momenta the theory of General Relativity also breaks down and in absence of a quantum theory of gravity the predictions of the theory becomes unreliable.

Defining $x' \equiv c_s x$, the equation of motion (3.28) reduces further to

$$\frac{d^2 u_k}{dx'^2} + \left(1 + \alpha_0 x'^2 + \beta_0 x'^4 - \frac{2}{x'^2} \right) u_k = 0 , \quad (3.32)$$

where

$$\alpha_0 \equiv \frac{D_2}{G_1 c_s^4} = \frac{D_2 G_1}{F_2^2} \quad \beta_0 \equiv \frac{C_2}{G_1 c_s^6} = \frac{C_2 G_1^2}{F_2^3} . \quad (3.33)$$

In order to have a stable vacuum in the deep UV, where the dispersion relation is dominated by the k^6 , we consider $\beta_0 > 0$ ⁵. If Hubble crossing takes place when $\omega^2 \sim k^2$, transitions between regions with a different dispersion relation can lead to excited initial states [7]. In this case, if $\alpha_0 < 0$, one can have relatively large corrections, much bigger than one, compared to the standard inflationary predictions for the two point function. This can happen even if the dispersion relation has a single turning point, *i.e.* a single horizon crossing is assumed to occur.⁶ The latter scenario is possible if $D_2/G_1 < 0$ and $C_2/G_1 > 0$ ⁷.

One should note that the coefficients \bar{M}_i 's, with $i = 1, 2, 3$, in (2.4) can appear with either sign. In fact, as noticed in the previous section about tensor perturbations, one has to assume $\bar{M}_3^2 < 0$ in order to avoid superluminal tensor perturbations. Noting that

$$D_2 = (\bar{M}_2^2 + \bar{M}_3^2 + H^2 \frac{\delta_4}{2} - \bar{M}_4 H) H^2 , \quad (3.34)$$

using the definition (3.33), the condition $\delta_3 = -3\delta_4$ and the expressions (3.24) for C_2 and G_1 , it is not difficult to show that

$$D_2 = -\frac{2\alpha_0}{\beta_0 c_s^2} \delta_4 H^4 . \quad (3.35)$$

As stated before, to achieve the stability in the UV, one has to assume that $\beta_0 > 0$, and hence, D_2 and $\alpha_0 \delta_4$ must have opposite signs. If $\delta_4 > 0$, noting that $C_2 = -2\delta_4 H^4$ will be negative, one has to assume $G_1 < 0$ to obtain a positive β_0 . Assuming $\dot{H} < 0$, that puts an upper bound on how large the first slow-roll parameter ϵ can be in this scenario. Then in order to achieve $\alpha_0 < 0$, one has to assume that $D_2 > 0$. This combined with that $\bar{M}_3^2 < 0$ for tensor perturbations to be subluminal, means

$$|\bar{M}_3|^2 < \bar{M}_2^2 + \frac{\delta_4}{2} H^2 - \bar{M}_4 H . \quad (3.36)$$

On the other hand, if $\delta_4 < 0$, one has to assume $D_2 < 0$ and the reverse of the above equality should hold and, conversely, a lower bound on ϵ is found under the assumption that $\dot{H} < 0$. One should emphasize that the value of α_0 and β_0 are not solely determined by c_s . There are other variables involved in these parameters and even for the choice $c_s \approx 1$, α_0 and β_0 can be substantial. Of course, with $c_s \ll 1$, the values of these parameters will get larger, but a

⁵Dispersion relations dominated by the k^6 term can also be motivated from studies in the context of condensed matter physics applied to black holes [14].

⁶We are referring to $1 + \alpha_0 x'^2 + \beta_0 x'^4 \sim \frac{2}{x'^2}$ as horizon crossing.

⁷In solid state physics, the perturbations that appear close to the minimum of the dispersion relations at large k 's are known as *rotons*, the ones that are close to the maximum of the dispersion relation are called *maxons* and the ones that are in the linear regime of the dispersion relation are *phonons*. In the cosmological setup, starting from deep inside the horizon, each mode in principle can undergo all three phases until it exits the horizon.

reduced sound speed is not necessary to have large corrections to the dispersion relation. For $c_s \approx 1$, since non-gaussianities are usually determined by factors $1/c_s^2$, one expects at least not to get an enhancement from speed of sound. Nonetheless the non-linear evolution of the mode inside the horizon due to the non-linear dispersion relations, may enhance the value of the non-gaussianity and bring it potentially to a value that can be observed. In fact, if, as in [7], the evolution of the modes inside the horizon can be mapped into excited states with large occupation numbers at horizon crossing, it is expected that, even for $c_s = 1$, equilateral [15, 16] and flattened non-gaussianity configurations [17, 18] will get enhanced. Whether such an equivalence holds and how higher dimensional operators in the EFTofI contribute to the bispectrum shapes and amplitude is something that we plan to investigate in near future.

3.3 Scalar power spectrum

In this section we investigate the effect of the modified mode equation (3.32) on the scalar power spectrum for different scenarios, corresponding to different signs and magnitudes of the coefficients of the quartic and sextic corrections to the dispersion relation. In general, overall correction to standard the Bunch-Davies power spectrum can be expressed as a multiplicative factor

$$P_s = \gamma_s P_s^{\text{B.D.}} = \gamma_s \frac{H^2}{8\pi^2 c_s \epsilon} . \quad (3.37)$$

If both $\alpha_0 > 0$ and $\beta_0 > 0$, the mode evolution inside the Hubble patch remains oscillatory throughout, and unless the parameters α_0 and β_0 are of order one, one does not expect to see a significant modification to the Bunch-Davies power spectrum. We solve the mode equation (3.32) numerically, from the positive frequency WKB mode in the infinite past,

$$u_k(x' \rightarrow -\infty) \simeq \frac{1}{2} \left(-\frac{\pi}{3} x' \right)^{1/2} H_{\frac{1}{6}}^{(1)} \left(-\frac{\sqrt{\beta_0}}{3} x'^3 \right) , \quad (3.38)$$




Figure 1. Plot of γ_s as a function of β_0 when α_0 and β_0 are both positive and $0.01 \leq \alpha_0 < 1$ with $0 \leq \beta_0 \leq 2$.




Figure 2. Plot of γ_s as a function of β_0 when α_0 and β_0 are both positive and $1 \leq \alpha_0 \leq 2$ with $0 \leq \beta_0 \leq 2$.

and then read off the power spectrum when the mode exits the horizon, $x' \rightarrow 0$. We have computed the factor γ_s for several values of α_0 and β_0 . For example, for $\alpha_0 = \beta_0 = 0.2$, one has $\gamma_s \simeq 0.717$, for $\alpha_0 = \beta_0 = 0.5$, $\gamma_s \simeq 0.53$ and for $\alpha_0 = \beta_0 = 1$, $\gamma_s = 0.4$. It seems that in this case, $\gamma_s < 1$ in most of the parameter space. In fact, solving and plotting the power spectrum for different values of α_0 and $0.01 < \beta_0 < 2$ confirms this conjecture. The larger the values of α_0 and β_0 , the more suppressed the power spectrum. However, this suppression is at most of order a few tenths. For $\alpha_0 = \beta_0 \sim 2$, the modulation factor $\gamma_s \approx 0.3$ (see Figs. 1 and 2). Noting that, in the case of positive quartic correction to the dispersion relation, enhancement of the coefficient of the quartic term suppresses the power spectrum too [19], one is tempted to deduce that positive higher order corrections to the dispersion relation will in general reduce the amplitude of the power spectrum.

On the other hand, if $\alpha_0 < 0$, there could be large modulation factors on the power spectrum. In [7], we focused on the range of parameters for α_0 and β_0 such that

$$z \equiv \frac{\beta_0}{\alpha_0^2} \geq \frac{1}{4} . \quad (3.39)$$

This requirement was coming from the fact that the dispersion relation should not become tachyonic for any physical momentum in flat spacetime. If we allow for $z < 1/4$, the quantum modes will get amplified while they are still inside the horizon, which will increase the resulting power spectrum exponentially. However, such a large enhancement for the scalar power spectrum will probably come about at the price of enhancing the size of non-gaussianity, and may thus clash with experimental bounds on them. The issue of non-gaussianity with such modified dispersion relations is something that we plan to return to in the next work. As for the power spectrum itself, we examined this regime numerically. We have plotted in Figs. 3 and 4 the factor γ_s , with β_0 in the range $\frac{\alpha_0^2}{5} < \beta_0 < \frac{\alpha_0^2}{4}$, for several values of α_0 . As can be readily seen, when β_0 goes below $\alpha_0^2/4$, the power spectrum grows substantially. With the




Figure 3. Log plot of γ_s as a function of β_0/α_0^2 when α_0 is big and negative ($0.1 \leq |\alpha_0| \leq 2$) and $\frac{\alpha_0^2}{5} \leq \beta_0 \leq \frac{\alpha_0^2}{4}$.

decrease of the parameter α_0 , the interval of time the mode spends under the influence of negative quartic contribution and possibly becomes tachyonic while still inside the horizon, increases and one sees a larger amount of enhancement for γ_s . In fact, for $\alpha_0 = 0.01$, and $\beta_0 = \alpha_0^2/5$, γ_s can reach values as large as 1000 (see Fig. 4). One other point that may be worth highlighting is that for large values of β_0 for each α_0 , the maximum enhancement in this case seem to occur at $\alpha_0 \sim 0.2$. As we will see, in the interval $1/4 < z < 1/3$, the same




Figure 4. Log plot of γ_s as a function of β_0/α_0^2 when α_0 is small and negative ($|\alpha_0| = 0.01$) and $\frac{\alpha_0^2}{5} \leq \beta_0 \leq \frac{\alpha_0^2}{4}$.

pattern repeats itself.

In [7], we had also assumed that the mode never becomes lighter than the Hubble parameter up until the last horizon crossing. That would mean that the equation $\omega^2 = 2H^2$ has only one solution. For

$$z > \frac{1}{3}, \quad (3.40)$$

this equation automatically has only one solution. For values of α_0 and β_0 such that

$$\frac{1}{4} \leq z \leq \frac{1}{3}, \quad (3.41)$$

having only one solution imposes one of the following conditions

$$\begin{aligned} \alpha_0 &\leq \frac{9z - 2 - 2(1 - 3z)^{3/2}}{54z^2} \\ \text{or } \alpha_0 &\geq \frac{9z - 2 + 2(1 - 3z)^{3/2}}{54z^2}. \end{aligned} \quad (3.42)$$

The requirement of having one solution was to show that the gluing technique becomes ineffective in capturing the modification to the power spectrum from the sextic dispersion relation. In that case, since $\omega^2 = 2H^2$ had only one solution, calculating the power spectrum analytically required the gluing procedure only once. In this paper, we do not distinguish between multiple and single horizon crossings. What matters to us is the amplitude of the power spectrum when the wavelength of the mode is much larger than the Hubble length or, more quantitatively, when $x' \ll 1$. Therefore, when investigating the $z > 1/4$ region, we allow for $z > 1/3$ as well. Plots of γ_s for different values of α_0 and β_0 for $z > 1/4$ are presented in Figs. 5, 6 and 7. In all cases, the factor γ_s is bigger than one. As one moves away from $\alpha_0 = 0.2$ in both directions, the amount of enhancement in the power spectrum decreases.

Finally, in the case $z = 1/4$, as noted in [7], for $\alpha_0 \sim 0.2$ and $\beta_0 = \alpha_0^2/4$, one can achieve $\gamma_s \simeq 14.774$.

3.4 Tensor-to-scalar ratio and consistency relation

Having modified spectra for both tensor and scalar fluctuations, it is expected that the tensor-to-scalar ratio be modified as well. For general modified equations of motion for scalar and tensor perturbations, the tensor-to-scalar ratio is given by

$$r = 16\epsilon \frac{c_s}{c_T} \frac{1}{\gamma_s}. \quad (3.43)$$

Therefore, a departure of any of the parameters, c_s , c_T and γ_s from one, will bring about changes in the tensor-to-scalar ratio in comparison with the slow-roll counterpart. It is well known that the majority of kinetic term dominated inflationary models, known as K-inflation [20], would lower r . However, the possibility of enhancing r by allowing for c_s to be bigger than one has also been entertained in [21]. As mentioned before, whether superluminal propagation in EFTs is necessarily an indication of any pathology is still a topic of debate in the community [22].

Depending on the sign of the operator $-\frac{\bar{M}_3^2}{2} \delta K^\mu_\nu \delta K^\nu_\mu$, which contributes to the speed of sound for gravity waves, one encounters two branches. In the *superluminal* branch, with




Figure 5. γ_s as a function of β_0/α_0^2 when $\alpha_0 < 0$ and $\beta_0 > 0$ and $0.01 \leq |\alpha_0| \leq 2$ and $\frac{\alpha^2}{4} \leq \beta_0 \leq \frac{\alpha^2}{3}$.




Figure 6. γ_s for big values of α_0 (0.5, 1, 2) as a function of β_0 when $\alpha_0 < 0$ and $\beta_0 > 0$ and $\frac{\alpha^2}{3} \leq \beta_0 \leq 2$.

$\bar{M}_3^2 > 0$, gravitons propagate faster than light during inflation, $c_T > 1$. In the *subluminal* branch, with $\bar{M}_3^2 < 0$, the speed of gravity waves is less than the speed of light. Were one to drop the superluminal branch because of superluminality and causality considerations, the effect of departure from the speed of light is always to enhance r . As \bar{M}_3 cannot be pushed beyond M_{Pl} in an EFT treatment, the gravity wave speed during inflation, is bounded from below by $1/\sqrt{2}$, and thus one at most would get an enhancement of $\sqrt{2}$ in r in the subluminal branch. Otherwise, in the superluminal branch, r can be significantly suppressed as \bar{M}_3^2 gets




Figure 7. γ_s for small values of α_0 (0.01, 0.1, 0.2) as a function of β_0 when $\alpha_0 < 0$ and $\beta_0 > 0$ and $\frac{\alpha^2}{3} \leq \beta_0 \leq 2$.

larger.

To summarize, in this section we showed that the effect of the EEFToI on scalar perturbations is to change their dispersion relation, not only by modifying the speed of sound but also by including higher order corrections. If the coefficients of the dispersion relation are all positive, the amplitude of the scalar power spectrum gets suppressed by a factor $\gamma_s \lesssim 1$. This would enhance r by a factor of order one. However, if the quartic term of the dispersion relation is negative, the evolution of the modes involves an intermediate phase with negative group velocity, or tachyonic phase, which can in fact enhance the amplitude of scalar perturbations considerably with respect to the Bunch-Davies result, picking up a factor of $\gamma_s \gg 1$. If one allows for the tachyonic evolution of the mode inside the horizon, r can get suppressed by a large factor. On the other hand, if only one allows for having a negative group velocity without a tachyonic phase, the maximum one can achieve is $\gamma_s \approx 14.7$, which suppresses r by a factor of ~ 0.06 . This is achieved by $\alpha_0 \sim 0.2$ and $\beta_0 = \alpha_0^2/4$ which was studied in [7] as a dispersion relation that realises super-excited states with large occupation numbers. The mode equation has one turning point for such values of parameters, corresponding to a single horizon crossing.

One can also easily verify that the consistency relation [23, 24]

$$n_T = -\frac{r}{8} \quad (3.44)$$

is modified to

$$n_T = -\frac{c_T \gamma_s}{c_s} \frac{r}{8}. \quad (3.45)$$

This implies that for a given r , we will have a much redder tensor spectrum for $\gamma_s \gg 1$.

4 Cut-off and non-gaussianity

As explained in the introduction, the EEFToI can provide a controlled description of the perturbations around a quasi de Sitter background from low energy up to the UV regime where the dispersion relation becomes $\omega^2 \sim k^6$, provided the coefficients of the higher dimensional operators are tuned such that the theory remains always weakly coupled in this energy range. This means that the scattering of Goldstone bosons does not violate perturbative unitarity at any scale below the point where the dispersion relation becomes dominated by the sextic term.

The strong coupling scale can be easily estimated in the π language from the size of the coupling of the interactions described by higher dimensional operators [1]. In order to perform this estimate, let us first note that the canonically normalised Goldstone boson π_c has a time kinetic term of the form

$$\int dt d^3x \frac{1}{2} \dot{\pi}_c^2,$$

and so the dimension of π_c , denoted as $[\pi_c]$, is

$$[\pi_c] = k^{3/2} \omega^{-1/2}. \quad (4.1)$$

The energy dimension of the coefficient c of a generic operator \mathcal{O} ,

$$\int dt d^3x c \mathcal{O},$$

can be inferred by imposing

$$\frac{c[\mathcal{O}]}{\omega k^3} = 1, \quad (4.2)$$

where $[\mathcal{O}] = k^s \omega^m [\pi_c]^p$ is the dimension of \mathcal{O} . Combining (4.1) and (4.2), we end up with

$$[c] = \omega^a k^b \quad \text{where} \quad a = 1 - m + \frac{p}{2} \quad \text{and} \quad b = 3 - s - \frac{3p}{2}.$$

If we now use the dispersion relation $\omega^2 \sim k^{2n}$ we find

$$[c] = \omega^q \quad \text{with} \quad q = 1 - m + \frac{1}{n}(3 - s) + \frac{p}{2} \left(1 - \frac{3}{n}\right). \quad (4.3)$$

Let us now consider, as illustrative examples, three different operators: $\mathcal{O}_1 = \dot{\pi}_c (\partial_i \pi_c)^2$, $\mathcal{O}_2 = (\partial_i \pi_c)^4$ and $\mathcal{O}_3 = \dot{\pi}_c^2 (\partial_i \pi_c)^2$ whose coefficients c_i have energy dimensions $[c_i] = \omega^{q_i}$ with $i = 1, 2, 3$. Different forms of the dispersion relation then give different energy scalings of these coefficients:

• $n = 1$	\Rightarrow	$q_1 = -2$	$q_2 = -4$	$q_3 = -3$
• $n = 2$	\Rightarrow	$q_1 = -\frac{1}{4}$	$q_2 = -\frac{1}{2}$	$q_3 = -\frac{5}{4}$
• $n = 3$	\Rightarrow	$q_1 = \frac{1}{3}$	$q_2 = \frac{2}{3}$	$q_3 = -\frac{2}{3}$

Interestingly, for $n < 3$ we get $q_i < 0 \forall i = 1, 2, 3$, while q_1 and q_2 become positive for $n = 3$. This implies that all three operators are *irrelevant* when the dispersion relation is at most dominated by the quartic term. This behaviour can be shown to hold for all higher derivative operators. Hence the $\omega^2 \sim k^2$ and $\omega^2 \sim k^4$ theories have only UV strong coupling scales which we will denote respectively Λ_2 and Λ_4 . On the other hand, for $\omega^2 \sim k^6$, \mathcal{O}_1 and \mathcal{O}_2 become *relevant*, while \mathcal{O}_3 remains *irrelevant*. This implies that the $\omega^2 \sim k^6$ theory features both a UV and an IR strong coupling scale which we will indicate respectively with Λ_6^{UV} and Λ_6^{IR} . The scale Λ_6^{UV} is associated with operators like \mathcal{O}_3 whose coupling grows when the energy increases, whereas Λ_6^{IR} is related to operators like \mathcal{O}_1 and \mathcal{O}_2 whose strength grows when the energy decreases. In general, this behaviour is a signal of an inconsistent EFT description.

However, one can still have a sensible EFT for the fluctuations if Λ_6^{IR} is below the transition from the $\omega^2 \sim k^4$ to the $\omega^2 \sim k^6$ regime that happens at the scale $\Lambda_{\text{dis}}^{4-6}$, provided the $\omega^2 \sim k^4$ theory is always weakly coupled in its regime of validity. Thus we need to tune the coefficients of the higher dimensional operators such that $\Lambda_6^{\text{IR}} \ll \Lambda_{\text{dis}}^{4-6} \ll \Lambda_4$. On top of this, the theory should not become strongly coupled at lower energies when the dispersion relation still takes the standard quadratic form. Hence we need also to require that Λ_2 is above the scale $\Lambda_{\text{dis}}^{2-4}$ which denotes the transition from the $\omega^2 \sim k^2$ to the $\omega^2 \sim k^4$ regime: $\Lambda_{\text{dis}}^{2-4} \ll \Lambda_2$.

The strong coupling scale Λ_2 can be estimated from the coefficient of the four-leg interaction term \mathcal{O}_2 . Given that $q_2 = -4$ in the deep IR where the dispersion relation is $\omega^2 = c_s^2 k^2$, the coefficient c_2 scales as $c_2 \sim 1/\Lambda_2^4$, where it can be shown that [1]

$$\Lambda_2^4 = c_s^7 \Lambda^4 \quad \text{with} \quad \Lambda^4 = \frac{2M_p^2 \dot{H}}{c_s^2 (1 - c_s^2)}. \quad (4.4)$$

This strong coupling scale goes to infinity for $c_s \rightarrow 1$ but for $c_s \ll 1$, which is for example a regime interesting for large non-gaussianities, it can become very small. However, as noted in [6], similarly to particle physics, new physics is expected to appear before reaching the strong coupling regime. In turn, this new physics pushes the strong coupling region to higher energies. As explained above, in our case, new physics corresponds to a change in the dispersion relation from linear, $\omega^2 = c_s^2 k^2$, to non-linear, $\omega^2 = k^4/\rho^2$, at the energy scale $\Lambda_{\text{dis}}^{2-4} = c_s^2 \rho$. This happens before the $\omega^2 = c_s^2 k^2$ theory becomes strongly coupled if $c_s \rho^4 \ll \Lambda_4$. Moreover, in the region where the dispersion relation is quartic, the new strong coupling scale becomes $\Lambda_4^4 = \Lambda^4 (\Lambda/\rho)^{28}$ [6]. Hence the condition $c_s \rho^4 \ll \Lambda^4$ guarantees also that this strong coupling scale is indeed larger than the one in the case with a linear dispersion relation.

As noted in [1], the strong coupling scale can also be quantified in terms of the size of non-gaussianities, since

$$\frac{\mathcal{L}_3}{\mathcal{L}_2} \equiv f_{\text{NL}} \zeta. \quad (4.5)$$

Thus perturbation theory breaks down when this ratio becomes of order one. Given that $\zeta \sim 10^{-4}$, this happens when $f_{\text{NL}} \sim 10^4$. Notice however that CMB observations set a stronger upper bound on equilateral non-gaussianities of the order $f_{\text{NL}} = -42 \pm 75$ [25]. The relation between non-gaussianities and strong couplings becomes even more manifest by noting that (4.5) can also be rewritten as [6]

$$\frac{\mathcal{L}_3}{\mathcal{L}_2} \sim \left(\frac{H}{\Lambda_2} \right)^2, \quad (4.6)$$

and so $f_{\text{NL}} \sim 10^4$ when horizon exit takes place in the strong coupling region, i.e. $H \sim \Lambda_2$. Given that $f_{\text{NL}} \sim c_s^{-2}$, large non-gaussianities emerge when $c_s \ll 1$. From (4.4) it is clear that small values of the speed of sound imply also a low strong coupling scale Λ_2 . However if horizon exit occurs in a region where the dispersion relation is dominated by the quartic term, i.e. when $\Lambda_{\text{dis}}^{2-4} \ll H \sim \Lambda_2 \ll \Lambda_4$, the theory should be weakly coupled. In fact, the ratio (4.5) takes an expression different from (4.6) when evaluated in the $\omega^2 = k^4/\rho^2$ region

$$\frac{\mathcal{L}_3}{\mathcal{L}_2} \sim \left(\frac{H}{\Lambda_4} \right)^{3/4}. \quad (4.7)$$

Given that $H \sim \Lambda_2 \ll \Lambda_4$, the theory is still weakly coupled even if $c_s \ll 1$. This implies also that f_{NL} should be smaller. In fact, the expression for f_{NL} in the quartic regime gets modified to $f_{\text{NL}} \sim (\Lambda_{\text{dis}}^{2-4}/H) c_s^{-2}$ which is suppressed with respect to its expression in the quadratic regime since $\Lambda_{\text{dis}}^{2-4} \ll H$. Hence we conclude that even a theory with small speed of sound and large non-gaussianities close to the edge of detectability can remain weakly coupled if the dispersion relation changes before hitting its strong coupling regime.

Notice also that, as explained in Sec. 3.3, large non-gaussianities might arise also in the case where the coefficients of the higher dimensional operators are chosen such that $c_s \simeq 1$. In fact, in this case the transitions inside the horizon between regions with modified dispersion relations would give $f_{\text{NL}} \sim \gamma_n/c_s^2$ with γ_n which could potentially be large. This case might also be interesting to keep the EEFToI weakly coupled since both Λ_2 and Λ_4 tend to infinity for $c_s \rightarrow 1$ (one would however still need to ensure that $\Lambda_6^{\text{IR}} \ll \Lambda_{\text{dis}}^{4-6}$).

One can insist on pushing the strong coupling scale to even higher energies via the appearance of new physics again before reaching Λ_4 . In this case, the new physics would correspond to a change in the dispersion relation from quartic to sextic at the scale $\Lambda_{\text{dis}}^{4-6}$ but we argued above that the $\omega^2 \sim k^6$ regime features both an UV and IR strong coupling scale. Hence a sensible EFT description all the way up to the sextic region requires to have also Λ_6^{IR} below $\Lambda_{\text{dis}}^{4-6}$.

In addition to the scale where perturbative unitarity is lost, another fundamental scale for the EFT of the perturbations around a quasi de Sitter background is the scale Λ_b where time diffeomorphisms are spontaneously broken by the background. Ref. [6] showed that, for the standard slow-roll case with $c_s = 1$, $\Lambda_b = \dot{\phi}^{1/2} \ll \Lambda_2$, while for $c_s \ll 1$, $\Lambda_2 \sim c_s \Lambda_b \ll \Lambda_b$. Moreover Λ_b controls the size of the perturbations with respect to the background since

$$k^3 P_\zeta(k) = \frac{1}{4} \frac{H^2}{M_{\text{Pl}}^2 \epsilon c_s} \sim \left(\frac{H}{\Lambda_b} \right)^4, \quad (4.8)$$

and so at energies around the symmetry breaking scale the background cannot be integrated out anymore to leave a theory for the fluctuations only. Hence we conclude that the cut-off of the EEFToI should be identified as the minimum between Λ_6^{UV} and Λ_b . As an illustrative example, a possible hierarchy of scales which would make the EEFToI valid for the entire energy range up to the $\omega^2 \sim k^6$ region would be

$$\Lambda_{\text{dis}}^{2-4} \ll \Lambda_2 \ll \Lambda_6^{\text{IR}} \ll \Lambda_{\text{dis}}^{4-6} \ll \Lambda_4 \ll \Lambda_6^{\text{UV}} \ll \Lambda_b. \quad (4.9)$$

We finally stress that cosmological observations are performed only at horizon crossing which occurs at energies of order H where the dispersion relation might take the standard linear expression. Hence, strictly speaking, the only requirement to have a sensible EFT is $H \ll \Lambda_{\text{cutoff}}$. However, we are demanding more since we want to provide a consistent theoretical

framework for the picture described in [7]. In fact, we want to be able to trust our EEFToI also in the high energy region which determines the behaviour of modes deep inside the horizon. We want these modes to be originally in the adiabatic vacuum of the $\omega^2 \sim k^6$ theory, so that the standard $\omega^2 \sim k^2$ theory features non-Bunch-Davis initial conditions due to the transitions between regions characterised by a different dispersion relation. In turn, these excited initial states can have very interesting implications for cosmological observables.

5 Conclusions

In this work we extended the formalism of the Effective Field Theory of Inflation, proposed earlier in [1], by including in the unitary gauge Lagrangian operators with mass dimensions three and four. There is a healthy combination of operators in this extended theory that allows one to avoid ghosts and imparts a sixth order correction to the dispersion relation for scalar perturbations. Modified dispersion relations are known to give rise to interesting effects on CMB observables due to non-Bunch-Davies initial conditions [7].

In this paper we provided a theoretical framework to motivate the emergence of modified dispersion relations. In particular, we discussed the conditions under which the Extended Effective Field Theory of Inflation is weakly coupled all the way from low energy to the UV region where the dispersion relation becomes dominated by the sextic term. In this situation, the evolution of the modes can be consistently followed from deep inside the horizon to super-horizon scales.

We also analysed the phenomenological implications of our EEFToI for the fluctuations around a quasi de Sitter background. We found that the tensor perturbations are untouched. On the other hand, various scenarios for the scalar perturbations can occur, depending on the parameters of the dispersion relation. If the coefficients of the quartic and the sextic dispersion relation are positive, the scalar power spectrum is suppressed. However, if the quartic coefficient is negative, even if for the case where the mode does not become tachyonic before crossing the only turning point, the scalar power spectrum can be enhanced. This in turn leads to a suppression of r with respect to models with Lorentzian dispersion relation.

In this paper we focused on the implications for the two point function but in the future it would be interesting to study also the behaviour of the three point function, in particular, because the size of non-gaussianities control the strong coupling regime of our EFT. Hence a careful analysis of non-gaussianities should provide a clear understanding on the possibility to tune the coefficients of the higher dimensional operators in order to have a sensible EFT description. This would correspond to non-gaussianities which are not larger than present observational constraints [25]. Another issue we plan to return to in the future, is the investigation of the possibility to obtain further extensions of the EFToI with terms higher than k^6 in the dispersion relation.

Acknowledgement

We are thankful to N. Bartolo, C.P. Burgess, F. Pedro, L. Senatore, M. M. Sheikh-Jabbari, G. P. Vacca, M. Zaldarriaga and in particular Paolo Creminelli for many helpful discussions. A.A. is thankful to the hospitality of Perimeter Institute where part of this work was completed. A.A. and R.C. are partly supported by INFN-FLAG.

References

- [1] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan, and L. Senatore, *The Effective Field Theory of Inflation*, JHEP **03** (2008) 014, [[0709.0293](#)].
- [2] M. Alishahiha, E. Silverstein, and D. Tong, *DBI in the sky*, Phys. Rev. **D70** (2004) 123505, [[hep-th/0404084](#)].
- [3] G. Shiu and B. Underwood, *Observing the Geometry of Warped Compactification via Cosmic Inflation*, Phys. Rev. Lett. **98** (2007) 051301, [[hep-th/0610151](#)].
- [4] S. E. Shandera and S. H. H. Tye, *Observing brane inflation*, JCAP **0605** (2006) 007, [[hep-th/0601099](#)].
- [5] N. Arkani-Hamed, P. Creminelli, S. Mukohyama, and M. Zaldarriaga, *Ghost inflation*, JCAP **0404** (2004) 001, [[hep-th/0312100](#)].
- [6] D. Baumann and D. Green, *Equilateral Non-Gaussianity and New Physics on the Horizon*, JCAP **1109** (2011) 014, [[1102.5343](#)].
- [7] A. Ashoorioon, R. Casadio, G. Geshnizjani, and H. J. Kim, *Getting Super-Excited with Modified Dispersion Relations*, JCAP **1709** (2017), no. 09 008, [[1702.06101](#)].
- [8] N. Bartolo, M. Fasiello, S. Matarrese, and A. Riotto, *Large non-Gaussianities in the Effective Field Theory Approach to Single-Field Inflation: the Bispectrum*, JCAP **1008** (2010) 008, [[1004.0893](#)].
- [9] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis, and R. Rattazzi, *Causality, analyticity and an IR obstruction to UV completion*, JHEP **10** (2006) 014, [[hep-th/0602178](#)].
- [10] E. Babichev, V. Mukhanov, and A. Vikman, *k-Essence, superluminal propagation, causality and emergent geometry*, JHEP **02** (2008) 101, [[0708.0561](#)].
- [11] P. Creminelli, J. Gleyzes, J. Norea, and F. Vernizzi, *Resilience of the standard predictions for primordial tensor modes*, Phys. Rev. Lett. **113** (2014), no. 23 231301, [[1407.8439](#)].
- [12] M. Ostrogradski, *Memoires sur les equations differentielles relatives au probleme des isoperimetre*, Mem. Ac. St. Petersburg **VI** 4 (1850) 385.
- [13] L. Senatore, K. M. Smith, and M. Zaldarriaga, *Non-Gaussianities in Single Field Inflation and their Optimal Limits from the WMAP 5-year Data*, JCAP **1001** (2010) 028, [[0905.3746](#)].
- [14] T. Jacobson, *Black hole evaporation and ultrashort distances*, Phys. Rev. **D44** (1991) 1731–1739.
- [15] I. Agullo and L. Parker, *Non-gaussianities and the Stimulated creation of quanta in the inflationary universe*, Phys. Rev. **D83** (2011) 063526, [[1010.5766](#)].
- [16] A. Ashoorioon, K. Dimopoulos, M. M. Sheikh-Jabbari, and G. Shiu, *Reconciliation of High Energy Scale Models of Inflation with Planck*, JCAP **1402** (2014) 025, [[1306.4914](#)].
- [17] R. Holman and A. J. Tolley, *Enhanced Non-Gaussianity from Excited Initial States*, JCAP **0805** (2008) 001, [[0710.1302](#)].
- [18] A. Ashoorioon and G. Shiu, *A Note on Calm Excited States of Inflation*, JCAP **1103** (2011) 025, [[1012.3392](#)].
- [19] A. Ashoorioon, D. Chialva, and U. Danielsson, *Effects of Nonlinear Dispersion Relations on Non-Gaussianities*, JCAP **1106** (2011) 034, [[1104.2338](#)].
- [20] C. Armendariz-Picon, T. Damour, and V. F. Mukhanov, *k - inflation*, Phys. Lett. **B458** (1999) 209–218, [[hep-th/9904075](#)].
- [21] V. F. Mukhanov and A. Vikman, *Enhancing the tensor-to-scalar ratio in simple inflation*, JCAP **0602** (2006) 004, [[astro-ph/0512066](#)].

- [22] G. Goon and K. Hinterbichler, *Superluminality, black holes and EFT*, JHEP **02** (2017) 134, [[1609.00723](#)].
- [23] A. A. Starobinsky, *Cosmic Background Anisotropy Induced by Isotropic Flat-Spectrum Gravitational-Wave Perturbations*, Sov. Astron. Lett. **11** (1985) 133.
- [24] A. R. Liddle and D. H. Lyth, *COBE, gravitational waves, inflation and extended inflation*, Phys. Lett. **B291** (1992) 391–398, [[astro-ph/9208007](#)].
- [25] **Planck** Collaboration, P. A. R. Ade et. al., *Planck 2013 Results. XXIV. Constraints on primordial non-Gaussianity*, Astron. Astrophys. **571** (2014) A24, [[1303.5084](#)].