

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Detection of Carcinogenic Polycyclic Aromatic Hydrocarbons in Stranded Caspian Seals (*Pusa caspica*)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Detection of Carcinogenic Polycyclic Aromatic Hydrocarbons in Stranded Caspian Seals (*Pusa caspica*) / Seyedeh Malihe Hoseini, Somayeh Namroodi, Annalisa Zaccaroni, Amir Sayad-Shirazi, Marcos Pérez-López, Francisco Soler-Rodríguez. - In: AQUATIC MAMMALS. - ISSN 0167-5427. - ELETTRONICO. - 46:1(2020), pp. 58-66. [10.1578/AM.46.1.2020.58]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/713461> since: 2020-01-14

*Published:*

DOI: <http://doi.org/10.1578/AM.46.1.2020.58>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of: S. M. Hoseini, S. Namroodi, A. Zaccaroni, A. Sayad-Shirazi, M. Pérez-López, and F. Soler-Rodríguez: Detection of Carcinogenic Polycyclic Aromatic Hydrocarbons in Stranded Caspian Seals (*Pusa caspica*). AQUATIC MAMMALS Volume 46 (1) Pages 58-66. The final published version is available online at: <https://doi.org/10.1578/AM.46.1.2020.58>

**Detection of Carcinogenic Polycyclic Aromatic Hydrocarbons in Stranded Caspian Seals**  
**(*Pusa caspica*)**

Seyedeh Malihe Hoseini,<sup>1</sup> Somayeh Namroodi,<sup>1</sup> Annalisa Zaccaroni,<sup>2,3</sup> Amir Sayad-Shirazi,<sup>4</sup> and  
Marcos Pérez-López<sup>5</sup> and Francisco Soler-Rodríguez,<sup>5</sup>

<sup>1</sup>*Gorgan University of Agricultural Sciences & Natural Resources, Department of  
Environmental sciences, Faculty of fisheries and environment, Gorgan, Iran  
e-mail: namroodi@gau.ac.ir*

<sup>2</sup>*University of Bologna, Department Veterinary Medical Sciences, University of Bologna, Viale  
Vespucci 2, 47042 Cesenatico (FC), Italy*

<sup>3</sup>*MarLab, Place du Chateau 7, 06250 Mougins, France*

<sup>4</sup>*Caspian Seal Conservation Center, Ashooradeh Island, Golestan Province, Iran*

<sup>5</sup>*University of Extremadura, Toxicology Unit, Veterinary School, Avda. de la Universidad s/n  
10003; Cáceres, Spain.*

**Abstract**

Polycyclic aromatic hydrocarbons (PAHs), which contain many carcinogenic compounds, are a major ingredient of petroleum/oil. PAH pollution of the Caspian Sea, the world's largest lake, is rapidly occurring and may be affecting the endangered Caspian seal (*Pusa caspica*), the only marine mammal in this lake. To analyze the entrance of PAHs into the Caspian Sea food chain and the health status of Caspian seals, we measured 16 carcinogenic PAHs in the blubber, kidney, and liver tissues of 10 Caspian seal carcasses from the coastal region of northeastern Iran, using gas chromatography-mass spectrophotometry. Of the 16 PAHs investigated, anthracene, phenanthrene, and naphthalene only were identified in nine sampled Caspian seals. Concentrations of anthracene ( $\bar{x}=84.83\pm79.86$  ppb wet weight [w.w.]), phenanthrene ( $\bar{x}=31.75\pm52.22$  ppb w.w.), and naphthalene ( $\bar{x}=25.1\pm31.57$  ppb w.w.) in blubber tissues were higher than in kidney and liver tissues. The concentration of PAHs in tissues was significantly higher in male than in female seals

and we found an inverse relationship between seal age and PAH concentration in tissues. Although no data exist concerning toxic effect concentrations of PAHs in Caspian seals, PAH detection in seal carcasses highlights a potentially stressful condition that may impact the health of Caspian seals and other sea life in this lake. Appropriate strategies for the control of PAH entrance into the Caspian Sea should be sought and studies for the determination of pathogenic and lethal doses of PAHs in Caspian seals should be pursued.

**Key words:** PAHs, Iran, tissues, *Pusa caspica*

## **Introduction**

The Caspian Sea, the largest closed lake in the world, is surrounded by five countries including Russia, Ghazakhstan, Azerbaijan, Turkmenistan and Iran (Shinsuke et al., 2003). This multinational access, in addition to the high density of passenger and commercial vessels and the influx of contaminated water from the numerous rivers that flow through industrial and agricultural areas, have resulted in the Caspian Sea becoming increasingly polluted (Effimoff, 2000; Kaplin, 1995). Extraction of oil from the lake, especially from the central and northern parts, and the entry of petroleum-based fuels in contaminated water have also contributed to pollution of the lake (Mille, 2007; Tolosa, 2004).

Polycyclic aromatic hydrocarbons (PAHs) with benzene rings are a highly toxic component of oil. Sixteen compounds of PAHs from oil have been identified as highly toxic and mutagenic for animals (Freeman et al., 1990). Generally, PAHs enter the environment in two ways: (a) human activities (pyrolytic origin) such as extraction, refining, transfer and export of crude oil, anthropogenic oil spills and other products, and (b) natural entrance (petrogenic origin) of biological resources through PAH synthesis by plankton, bacteria, algae, and the decomposition

of plants (Nasrollahzadeh Saravi et al., 2013). Fortunately, PAHs break down quickly when exposed to light and oxygen (e.g., photo-oxidation) and some PAHs are biodegraded by naturally-occurring bacteria (Kanaly & Harayama, 2000; Mrozik et al., 2003). PAHs divided in low and high molecular weight based on number of their benzene rings. Oil spills represent one of major threats for marine mammals, both in the short and long term. However, high molecular weight (HMW) PAHs have higher tendency to accumulate in animals' tissues than low molecular weight (LMW) PAHs. In aquatic ecosystem HMW PAHs precipitate faster than LMW PAHs and have lower chance to accumulate in body of marine animals (Lawal, 2017). Limited studies have shown PAH pollution in Caspian Sea coastal areas of Baku (due to oil extraction), northern Russia and Kazakhstan, and southern Iran (Kardovani, 1995; Tolosa, 2004).

The Caspian seal (*Pusa caspica*), the only species of marine mammal in the Caspian Sea, is at the top of the food chain in this ecosystem. This species has migratory life style and due to various factors such as death in fishing nets and reduced reproductive rates in male and female seals because of exposure to pollutants, the health status of this species is poor (Harkonen et al. 2008, ;Watanabe et al., 2002). Mass mortalities of these seals along the Caspian Sea coast during 1997-2000, along with the factors cited above, have resulted in placement of the Caspian seal on the IUCN red list (Goodman & Dmitrieva, 2016). Considering the increase in Caspian Sea oil pollution in recent years and the migratory lifestyle of Caspian seals, we monitored PAH pollution in the Caspian Sea and their health status by measuring and assessing concentrations of 16 mutagenic PAHs in stranded Caspian seal tissues.

## **Methods**

### *Sampling*

After approval of the study by the Ethics Committee of the Deputy of Natural Environment of Golestan Province (permit number: 125/7894), 10 stranded Caspian seals along the coasts of Golestan and Mazandaran provinces, Iran, were collected during 2012-2015. Stranded Caspian seals were trapped in fishing nets and were found fresh by fisheries. Following biometric analysis and clinical examination, the seals were transferred to the laboratory and necropsied. Liver, kidney, and blubber tissues were sampled to measure PAH concentrations and a lower canine tooth was extracted to determine seal age (Amano et al., 2000).

#### *Measurement of PAH concentration*

This analysis was performed at the Toxicology Lab, Faculty of Veterinary Medicine, University of Extremadura, Caceres, Spain. To measure tissue concentrations of PAHs, the standard method of Lucas and Zhao (2015) using gas chromatography-mass spectrometry (GC-MS/MS) was applied. The samples were shredded with glass slurry, freeze dried for 72 h, and homogenized using a mortar and pestle. A portion of each freeze-dried sample was weighed into 50 mL centrifuge tubes and MilliQ water was added to obtain the equivalent to 5 g of fresh sample. Ten ml of acetonitrile were added to the tube and the sample was shaken vigorously by hand for 2 min and centrifuged for 5 min at 5,000 rpm. The supernatant was transferred to a 15-ml centrifuge tube containing 1 g of Agilent Enhanced Matrix Removal-Lipid sorbent (Agilent Technologies, Santa Clara, California, USA) and placed on a vortex mixer for 60 s. The solution was then centrifuged at 5,000 rpm for 3 min. The supernatant was transferred to a second 15-ml tube containing 2 g salt (1:4 NaCl: MgSO<sub>4</sub>) and immediately vortexed and centrifuged for 3 min. The top layer containing acetonitrile was evaporated to dryness in a rotatory concentrator, reconstituted in ethyl acetate:n-hexane (1:4) and filtered (0.45 micron) into a 2-ml GC vial.

Obtained extracts (1 µl) were injected into a gas chromatograph-mass spectrometer (SCION GC-MS-Triple Quad MS/MS; Bruker Scientific Instruments, Billerica Massachusetts, USA) and PAHs were separated on a 30-m Agilent J&W DB-5ms (5% diphenyl/dimethylpolysiloxane) capillary column (250 µm id. x 0.25 µm film thickness). For quality control and for quantitative purposes, one sample from each matrix (blubber, liver, and kidney) was pre-spiked at 40 ng/g with a certificate standard (DE-PROM 16 EPA Priority PAHs in Toluene from LGC Standards, 100 µg/ml) containing the 16 PAHs considered to be of primary concern by the United States Environmental Protection Agency (USEPA). Qualification and quantification ions were monitored for each compound at  $m/z = 102+126+127$  for naphthalene,  $126+150+151$  for acenaphthylene,  $127+151+152$  for acenaphthene,  $139+163+164$  for fluorene,  $152+176+177$  for phenanthrene, anthracene, 200 for fluoranthene,  $151+200$  for pyrene,  $202+226+227$  for benzo[a]anthracene, chrysene, 250 for benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[a]pyrene, 274 for benzo[g,h,i]-perylene, indeno[1,2,3-cd]pyrene, and  $276+277$  for dibenzo[a,h]anthracene.

Results of each spiked matrix were used for quantitation of matrix-related samples. These matrix-matched samples were chosen in order to compensate for any signal suppression/enhancement compared to their relative response in pure solvent. Procedural blanks containing reagents only were screened during the analysis of each batch (one per tissue) in order to ensure that solvents and all the used material were free from PAHs residues. To evaluate the linearity of the method calibration curves were built with concentrations ranging from 4 to 400 µg/L with  $r^2 > 0.98$ . The limit of quantitation (LOQ) was determined as the lowest concentration of the compounds which can be reliably detected with the signal-to-noise (S/N) ratio higher than

10. In our study, the LOQ for individual PAHs in samples ranged from 0.5 to 5 ng/g, on a wet weight [w.w.] basis.

#### *Data analysis*

Normality and lack of data were evaluated using Kolmogorov-Smirnov test. The result of this test was higher than 0.05, indicating that the data was normal. The effect of sex on the PAH concentration in sampled tissues was investigated using Student's t-tests. Linear regression test was used to survey impact of age on PAH concentration. Also ANOVA one way test was used to survey difference of PAH concentrations in sampled tissues.

### **Results**

Of the 16 PAHs studied, three (i.e. anthracene ( $\text{Log } K_{\text{OW}}=4.54$ , octanol-air partition coefficient ( $K_{\text{AO}}=7.55$ , number of benzene rings ( $\text{NB}=3$ ), phenanthrene ( $\text{Log } K_{\text{OW}}=4.57$ ,  $K_{\text{AO}}=7.57$ ,  $\text{NB}=3$ ) and naphthalene ( $\text{Log } K_{\text{OW}}=3.37$ ,  $K_{\text{AO}}=5.19$ ,  $\text{NB}=2$ ) were identified in the tissues of sampled Caspian seals (Table 1) (National Center for Biotechnology Information, 2019). We detected PAHs in nine of the 10 sampled seals at different concentrations and the accumulation pattern of PAHs differed among tissue types. Among blubber tissues, 70% were contaminated with anthracene, 50% with naphthalene, and 40% with phenanthrene. For liver tissues, 80% of samples also were contaminated with anthracene, 80% contained phenanthrene, and 50% contained naphthalene. For kidney tissues 90% were contaminated with anthracene and 40% with naphthalene. Phenanthrene contamination was not detected in kidney tissues (Table 1). The concentrations of anthracene ( $\bar{x}=84.83\pm79.86$  ppb w.w.), phenanthrene ( $\bar{x}=31.75\pm52.22$  ppb w.w.), and naphthalene ( $\bar{x}=25.1\pm31.57$  ppb w.w.) in blubber tissues were significantly ( $p=0.001$ ) higher than in kidney and liver (Table 2). The concentration of anthracene in all sampled tissues



was significantly ( $p=0.001$ ) higher than the concentrations of phenanthrene and naphthalene (Table 2). Concentrations of PAHs in liver, kidney, and blubber were significantly higher in male than in female seals (Table 3 & Figure 1). Linear regression test indicated a negative relationship between Caspian seal age (using body length as a surrogate for age) and the concentration of PAHs in the tissues (Figure 2).

Place table 1, 2, and 3 here, Place figures 1&2 here

## Discussion

Detection of PAHs in sampled Caspian seals indicated that PAHs entered the Caspian Sea food chain. Also, previous studies have indicated pollution with just anthracene, phenanthrene, Dibenzo(a,h) anthracene, fluoranthene, fluorene, and naphthalene in the Caspian Sea (Eskandarpour et al., 2014; Nasrallahzadeh Saravi et al., 2012). Absence of other 13 surveyed PAHs in Caspian seals' tissues can be due to elimination of them in Caspian seals by metabolism or absence of them in Caspian seals' habitat (Meador et al., 1995b). Nasrallahzadeh Saravi et al. (2012) studied the sediment and muscle tissue of fish in the southern Caspian Sea near Astara, White River, Tonekabon, and Amir Abad, Iran. They measured an anthracene concentration of  $7.6 \pm 4$  ppb w.w and dibenzene concentration of  $66.6 \pm 75$  ppb w.w in sediment. Benzo fluoranthene, and benzo pyrene were detected in muscle tissues of several Caspian Sea fish species, including the Caspian kutum (*Rutilus frisii kutum*) with concentrations of 530 ppb w.w. and 96.6 ppb w.w. and the leaping mullet (*Chelon saliens*) at 80 ppb w.w. and 176 ppb w.w., respectively. In another study on fish flour (fish meal) of Caspian Sea sprat (*Clupeonella cultriventris caspia*), naphthalene, fluorine, and anthracene concentrations were measured at  $24.66 \pm 15.52$  ppb w.w.,  $1.32 \pm 1.54$  ppb w.w., and  $1.1 \pm 1.92$  ppb w.w., respectively (Eskandarpour et al., 2014). Absence of the other

studied PAHs in sampled Caspian seals can be due to short time presence of Caspian seal in sea shore of Iran (Harkonen et al. 2008). As surveyed tissues in previous mentioned studies on fish are not similar with tested tissues in our study, we cannot compare obtained results. However, Compared to the PAH concentration in some of these fish species, the higher concentrations in sampled Caspian seals may be due to higher concentrations of fat in Caspian seal tissue and their greater longevity, allowing for longer bioaccumulation.

Unfortunately, there are few similar studies on carcinogenic PAH contamination in marine mammals. Marsili et al. (2001) measured total PAHs in blubber tissues of fin whales (*Balaenoptera physalus*) and striped dolphins (*Stenella coeruleoalba*) and found high levels of contamination in these Mediterranean cetaceans. Marsili et al. (1997) surveyed PAH concentration in liver tissues of South American sea lions (*Otaria flavescens*) in the Plata Sea, Argentina. They detected naphthalene ( $\bar{x}=194\pm54.07$  ppb w.w), anthracene ( $\bar{x}= 0.30\pm2.270$  ppb w.w.), and phenanthrene ( $\bar{x}=27.1\pm21.75$  ppb w.w.) in sampled tissues. Many factors such as level and time of animal PAHs exposure, sex and age of sampled animals, species ability to metabolize PAHs and difference in volume of PAHs entrance in animals' habitat can result in detection of higher concentrations of naphthalene and phenanthrene and lower concentration of anthracene in sampled sea lions than sampled Caspian seals (Meador et al., 1995b). Also, Hellou et al. (1991) examined PAH concentration in muscle tissue of harp seals (*Phoca groenlandica*) in the northwestern Atlantic Ocean. Those specimens were contaminated with phenanthrene, anthracene, and fluorene, and PAH concentration was measured between 10 and 31 ppb lipid w.w. As we did not examine muscle tissues in sampled Caspian seals, we cannot compare our results with results of Hellou et al. (1991) on harp seals.

Of the three PAHs detected (naphthalene, phenanthrene, and anthracene), anthracene occurred at the highest concentrations in sampled tissues while phenanthrene had the lowest concentrations. A different trend in PAH concentrations has been documented in fish species of the Caspian Sea; for example, Eskandarpour et al. (2014) reported higher concentrations of naphthalene than anthracene in Caspian Sea sprat (*Clupeonella cultriventris caspia*). Similarly, Kannan & Perrotta (2008) detected naphthalene in kidney tissues in higher concentrations compared to other PAHs in 81 adult female sea otters (*Enhydra lutris*) from the California coast. Many factors, such as differences in the source of PAHs in sampling areas, chemical condition of the sampling area, number of PAH benzene rings, and tendency for bioaccumulation of PAHs in different species could lead to these differences in findings (Meador et al., 1995b).

As mentioned, tissue accumulation patterns of detected PAHs differed among Caspian seal tissue types. The number of PAH benzene rings is an important variable associated with bioaccumulation of PAHs in Caspian seal tissues. With increases in the number of rings, the hydrophobic and lipophilic properties of PAHs increase. When PAHs enter Caspian seal bodies, they can accumulate in several tissues, but especially in blubber. Microbial degradation of PAHs decreases with increases in the number of PAH benzene rings, so PAHs with more benzene rings can accumulate to higher concentrations in tissues (Landrum, 1989; Landrum & Robbins, 1990; Meador et al., 1995a; Meador, 2003). It follows that higher tissue concentration of anthracene than naphthalene could be explained by the higher number benzene rings of anthracene. Higher anthracene tissue concentration than phenanthrene, which has the same number of benzene rings, might be explained by a higher rate of entry of anthracene into the Caspian Sea.

The higher PAH concentrations in blubber tissues may be explained by the hydrophobic and lipophilic characteristics of PAHs. Eskandarpour et al. (2014) found that bioaccumulation of

PAHs in Caspian Sea sprat was directly related to the concentration of fat in sampled tissues, with higher concentrations detected in tissues with higher levels of fats (Eskandarpour et al., 2014). In contrast, there was no correlation between tissue PAH concentrations and the amount of fat in tissues of Harp seals (Hellou et al., 1991).

We found the lowest PAH concentrations in liver tissues. Most of the PAHs are absorbed through the digestive system and transferred to the liver for detoxification by enzymes, including the P450 family and microsomal enzymes (Engelhardt, 1982; Addison & Brodie, 1984; Marsili et al., 1997; Lee et al., 2005). This detoxification process could lead to lower concentrations of PAHs in liver tissues than blubber or kidney.

A negative correlation between PAH concentration in tissues with the age of sampled Caspian seals was found, indicating the influence of age as a confounding factor on PAH concentrations in our seals. Similarly, studies of Northern pike (*Esox lucius*) and Caspian Sea sprat revealed that animal age can affect PAH bioaccumulation rate with similar trends (Ghorbani et al., 2010; Eskandarpour et al., 2014). As age increases, PAH concentration in tissues can decrease due to an increase in PAH metabolism. Hellou et al. (1991) showed that PAH concentration was higher in younger Harp seals than older seals. They also found that the activity of the metabolizing enzymes, including P450 family enzymes, on PAHs increases with age and can lead to a decrease in PAH concentrations in tissues by age (Hellou et al., 1991). However, Harris et al., (2011) who surveyed Hydrocarbon concentrations and patterns in blood samples of 29 live-captured sea otters (*Enhydra lutris*) from British Columbia, Canada, reported similar hydrocarbon concentrations among different age classes (Harris et al., 2011).

The higher average age sampled females (18.2 y old) than males (14.5 y old), may be one explanation for our finding showing lower PAH concentrations in tissues of female Caspian seals, highlighting the putative biotransformation of PAHs by P450 enzymes. This result supports findings of Addison et al. (1973) and Moyer et al. (1988) in harp seals and ringed seals (*Phoca hispida*), respectively. Those authors suggested that this difference could be due to higher activity of P450 family enzymes in females resulting in a lower PAH concentration in female tissues. Eskandarpour et al. (2014) explained that such results can occur due to maternal transfer of PAHs in fish through egg laying. However, studies by Hellou et al. (1991) and Marsili et al. (2001) on harp seals and cetaceans (fin whales and striped dolphins), respectively, showed no relation between sex and PAH tissue concentrations.

Elimination of organochlorine pesticides in female Caspian seals through giving birth has been described, but there is no information about offloading of PAHs in seals via that mechanism (i.e. maternal transfer) (Tanabe et al., 1978; Subramanian et al., 1987; Nakata et al., 1995). Thus, additional studies are needed on this topic.

Molecular ratios of PAHs have been used to identify their sources in the environment. According to Hajizadeh et al. (2010), if the ratio of phenanthrene to anthracene is  $>10$ , the origin of these pollutants is considered petrogenic, and but if the ratio is  $<10$ , their origin is considered pyrolytic. Nasrollahzadeh Saravi et al. (2012), who detected PAHs in leaping mullet and Caspian kutum in the southern part of the Caspian Sea (i.e. Mazandaran and Golestan), reported that the origin of PAHs in that region of the Caspian Sea is more pyrolytic. Nemati Varnosfaderany et al. (2014) and Baniemam et al. (2017) reported both pyrolitic and petrogenic origins of PAHs in the south Caspian Sea (Iranian coastal regions). In the present study, the ratio of PAHs with low molecular weight (phenanthrene) to those with a high molecular weight (anthracene) also indicated

that the PAH compounds in sampled Caspian seals may have pyrolytic sources. Nevertheless, Kannan & Perrotta (2008) concluded that the detection of PAHs with a predominance of di- and tri-cyclic PAHs over tetra- and penta-cyclic PAHs (as we also detected in Caspian seals) suggest petrogenic sources.

In conclusion, it appears that Caspian seals are exposed to PAHs of both petrogenic and pyrolytic origins in the Caspian Sea. Our results show that PAHs eventually accumulate in the lake's food chain and enter Caspian seal tissues. Because PAHs retention rate time are low in the environment (i.e. high elimination rate), their accumulation in Caspian seals indicates that the emission or inflow rate of PAH entering into the Caspian Sea ecosystem is much higher than their rate of removal (e.g., burial rate). Of course, given the migratory lifestyle of Caspian seals, it is not possible to identify the exact route and location of PAH contamination of Caspian seals (Anyakora et al., 2005).

No ecotoxicological risk assessments have been conducted to derive the toxic effect concentration and/or safe level thresholds of PAHs on Caspian seals. PAH accumulation in tissues could have negative effects on the health status of this species, even at very low concentrations. Appropriate strategies for limitation of PAH entrance into the Caspian Sea should be considered in countries bordering the Caspian Sea. Further studies are needed to determine pathogenic and lethal doses of PAHs in Caspian seals.

### **Acknowledgments**

This study was supported by Gorgan University of Agricultural Sciences and Natural Resources. The author thanks Dr. James Norman Milles for edition of the earlier manuscript that greatly

improved it. We also show our gratitude to Seyyed Reza Kahlegi for her help in obtaining some sampled Caspian seals.

### Literature Cited

Addison, R.F., & Brodie, P.F. (1984). Characterization of ethoxyresorufin O-de-ethylase in grey seal *Halichoerus grypus*. *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology*, 79, 261-263.

Addison, R.F., Kerr, S.R., Dale, J., & Sergeant, D.E. (1973). Variation of organochlorine residue levels with age in Gulf of St. Lawrence Harp seals (*Pagophilus groenlandicus*). *Fisher Board of Canada*, 30, 595-600.

Amano, M., Miyazaki, N., & Petrov, E.A. (2000). Age determination and growth of Baikal seals (*Phoca sibirica*). In: Rossiter A, Kawanabe H, editors. *Advances in Ecological Research, Ancient Lakes: Biodiversity. Biodiversity Ecology Evolution*, 31, 449–462.

Anyakora, C., Ogbeche, A., Palmer, P., & Coker, H. (2005). Determination of polynuclear aromatic hydrocarbons in marine samples of Siokolo Fishing Settlement. *Chromatography*, 10, 323-330.

Baniemam, M., Moradi, A.M., Bakhtiari, A.R., Fatemi, M.R., & Khanghah, K.E. (2017). Seasonal variation of polycyclic aromatic hydrocarbons in the surface sediments of the southern Caspian Sea. *Marine Pollution Bulletin*, 15, 478-485.

Effimoff, I. (2000). The oil and gas resource base of the Caspian region. *Petroleum Science and Engineering*, 28, 157–159.

- 309 Engelhardt, F.R. (1982). Hydrocarbon metabolism and cortisol balance in oil-exposed ringed seals,  
310 *Phoca hispida*. Comparative Biochemistry and Physiology Part C: *Comparative*  
311 *Pharmacology*, 72, 133-136.
- 312 Eskandarpour, A., Aryanashja, A., & Salimi, L. (2014). Qualitative and quantitative analyze of  
313 PAHs (sixteen of important PAHs in EPA list) in tissues and meal of *Clupeonella*  
314 *cultriventris caspia* of Guilan Province Caspian waters (Bandar Anzali). *Aquatic Physiology*  
315 *Biotechnology*, 105, 51-56. (In Persian)
- 316 Freeman, D.J., & Cattell, F.C.R. 1990). Wood burning as a source of atmospheric polycyclic  
317 aromatic hydrocarbons. *Environmental Science and Technology*, 24, 24-39.
- 318 Ghorbani, S., Salimi, L., & Matlabi, A. (2010). Quantitative and qualitative study of PAHs (EPA  
319 index list) in fish duck muscle tissue *Ezox lucius* Anzali wetland, Master thesis, Islamic Azad  
320 University, North Tehran Branch.
- 321 Goodman, S., & Dmitrieva, L. (2016). Pusa caspica. The IUCN Red List of Threatened Species.
- 322 Hajizadeh, N., AbbasI, A., Ramini, S., Moghaddam, M., Rahmani, I. (2010). The concentration  
323 and origin of petroleum hydrocarbons in sediments of Anzali port. *Ecology*, 6, 99-106.
- 324 Harkonen, T., Jüssi, M., Baimukanov, M., Bignert, A., Dmitrieva, L., Kasimbekov, Y., Verevkin,  
325 M., Wilson, S. & Goodman, S. J. (2008). Pup production and breeding distribution of the  
326 Caspian seal (*Phoca caspica*) in relation to human impacts. *AMBIO: A Journal of the Human*  
327 *Environment*, 37(5), 356-361.
- 328 Harris, K.A., Nichol, L.M., & Ross, P.S. (2011). Hydrocarbon concentrations and patterns in free-  
329 ranging sea otters from British Columbia, Canada. *Environmental Toxicology and*  
330 *Chemistry*, 30, 2184-2193.



- 331 Hellou, J., Upshall, C., Ni, I.H., Payne, J.F., & Huang, Y.S. (1991). Polycyclic aromatic  
332 hydrocarbons in harp seals (*Phoca groenlandica*) from the Northwest Atlantic. *Archives of*  
333 *Environmental Contamination and toxicology*, 21, 135-140.
- 334 Kanaly, R. A., & Harayama, S. (2000). Biodegradation of high-molecular-weight polycyclic  
335 aromatic hydrocarbons by bacteria. *Journal of bacteriology*, 182(8), 2059-2067.
- 336 Kannan, K., & Perrotta, E. (2008). Polycyclic aromatic hydrocarbons (PAHs) in livers of  
337 California sea otters. *Chemosphere*, 7, 649–655.
- 338 Kaplin, P. (1995). Caspian Sea environmental situation in the condition of water level rises. *Water*  
339 *Develop*, 1, 123-150.
- 340 Kardovani, P. (1995). Iran marine ecosystem (Caspian Sea). Tehran: Ghomse Publication.
- 341 Landrum, P.F. (1989). Bioavailability and toxicokinetics of polycyclic aromatic hydrocarbons  
342 sorbed to sediments for the amphipod *Pontoporeia hoyi*. *Environmental Science Technology*,  
343 23, 588-595.
- 344 Landrum, P.F., & Robbins, J.A. (1990). Bioavailability of sediment associated contaminants: a  
345 review and simulation model, sediments. of in-place pollutants, Lewis, Chelsea. *Chemistry*  
346 *Toxicity*, 237-263.
- 347 Lawal, A. T. (2017). Polycyclic aromatic hydrocarbons. A review. *Cogent Environmental*  
348 *Science*, 3(1), 1339841.
- 349 Lee, R. F., & Anderson, J. W. (2005). Significance of cytochrome P450 system responses and  
350 levels of bile fluorescent aromatic compounds in marine wildlife following oil spills. *Marine*  
351 *Pollution Bulletin*, 50, 705-723.

- 352 Lucas, D., & Zhao, L. (2015). PAH analysis in salmon with enhanced matrix removal. *Agilent*  
353 *Technologies Application*, pp. 5991-6088.
- 354 Marsili, L., Caruso, A., Fossi, M.C., Zanardelli, M., Politi, E., & Focardi, S. (2001). Polycyclic  
355 aromatic hydrocarbons (PAHs) in subcutaneous biopsies of Mediterranean cetaceans.  
356 *Chemosphere*, 44, 147-154.
- 357 Marsili, L., Fossi, M.C., Casini, S., Savelli, C., Jimenez, B., Junin, M., & Castello, H. (1997).  
358 Fingerprint of polycyclic aromatic hydrocarbons in two populations of southern sea lions  
359 (*Otaria flavescens*). *Chemosphere*, 34, 759-770.
- 360 Meador, J.P. (2003). Bioaccumulation of PAHs in marine invertebrates; PAHs: An  
361 ecotoxicological perspective; John Willey and Sons Ltd. *Ecotoxicological*, 147-171.
- 362 Meador, J.P., Casillas, E., Sloan, C.A., & Varanasi, U. (1995a). Comparative bioaccumulation of  
363 polycyclic aromatic hydrocarbons from sediment by two infaunal invertebrates. *Marine*  
364 *Ecology Progress Series*, 107-124.
- 365 Meador, J.P., Stein, J.E., Reichert, W.L., & Vatanasi, U. (1995b). Bioaccumulation of polycyclic  
366 aromatic hydrocarbons by marine organisms. *Reviews of Environmental Contamination and*  
367 *Toxicology*, 143,79-165.
- 368 Mille, G., Asia, L., Guiliano, M., Malleret, L., & Doumenq, P. (2007). Hydrocarbons in Coastal  
369 Sediments from the Mediterranean Sea (*Gulf of Fos Area, France*). *Marine Pollution*  
370 *Bulletin*, 54, 566-575.
- 371 Mrozik, A., Piotrowska-Seget, Z., & Labuzek, S. (2003). Bacterial degradation and bioremediation  
372 of polycyclic aromatic hydrocarbons. *Polish Journal of Environmental Studies*, 12(1): 11-  
373 15.

- 374 Moyer, D.C., Norstrom, R.J., & Simon, M. (1988). Organochlorine contaminants in Arctic marine  
375 food chains: accumulation of specific polychlorinated biphenyls and chlordane-related  
376 compounds. *Environmental Science technology*, 22, 1071-1079.
- 377 Nakata, H., Tanabe, S., Tatsukawa, R., Amano, M., Miyazaki, N., & Petrov, EA. (1995). Persistent  
378 organochlorine residues and their accumulation kinetics in Baikal seal (*Phoca sibirica*) from  
379 Lake Baikal, Russia. *Environmental Science Technology*, 29, 2877-2885.
- 380 Nasrollahzadeh Saravi, H., Pourgholam, R., Unesipour, H., & Makhloogh, A. (2012).  
381 Polyaromatic hydrocarbons (16 PAHs) at the sediments and edible tissue of *Chelon saliens*  
382 and *Rutilus frisii kutum* in Caspian Sea. *Journal of Mazandaran university of medical*  
383 *sciences*, 22(94), 79-90. (In Persian)
- 384 National Center for Biotechnology Information. (2019). <https://pubchem.ncbi.nlm.nih.gov/>  
385 compound
- 386 Nemati Varnosfaderany, M., Riyahi Bakhtiari, A., & Guoqiang Chu, Z.H. (2014). Vertical  
387 distribution and source identification of polycyclic aromatic hydrocarbons (PAHs) in  
388 southwest of the Caspian Sea: Most petrogenic events during the late Little Ice Age. *Marine*  
389 *Pollution Bulletin*, 15, 152-163.
- 390 Shinsuke, T., Satoko, N., Tu Binh, M., Nobuyuki, M., & Evgeny, A. (2003). Petrov Temporal  
391 Trends of Persistent Organochlorine Contamination in Russia: A Case Study of Baikal and  
392 Caspian Seal. *Arch Environmental Contaminate Toxicology*, 44, 533–545.
- 393 Subramanian, A., Tanabe, S., & Tatsukawa, R. (1987). Age and size trends and male female  
394 differences of PCBs and DDE in Dalli-type Dall's porpoises, *Phocoenoides dalli* of  
395 northwestern North Pacific. *Proc NIPR Symp Polar Biol*, 1, 205–216.

- 396 Tanabe, S., Loganathan, B.G., Subramanian, A.N., & Tatsukawa, R. (1987). Organochlorine  
397 residues in short-finned pilot whale; possible use as tracers of biological parameters. *Marine*  
398 *Pollution Bulletin*, 18, 561-563.
- 399 Tolosa, I., Mora, S., Sheikholeslami, M.R., Villeneuve, J.P., Bartocci, J., & Cattini, C. (2004).  
400 Aliphatic and aromatic hydrocarbons in coastal Caspian Sea sediments. *Marine Pollution*  
401 *Bulletin*, 48, 44–60.
- 402 Watanabe, Kunito, T., Tanabe, S., Amano, M., Koyama, Y., Miyazaki, N., Petrov, E.A., &  
403 Tatsukawa, R. (2002). Accumulation of Heavy Metals in Caspian Seals (*Phoca caspica*).  
404 *Environmental Contamination Toxicology*, 43, 109–120.
- 405
- 406

407 **Table 1:** PAHs concentration (ppb w.w.) in the tissues of Caspian seals based on age and sex.

Sample number	1	2	3	4	5	6	7	8	9	10	Average
Sex	F	F	M	F	F	M	M	M	M	M	
Age	26	17	16	11	19	9	10	20	6	26	
A(BT)	0	56.93	82.44	95.86	44.22	194.55	174.69	0	199.68	0	84.83±79.86
A(LT)	0	6.68	7.34	8.56	6.35	10.42	10.05	7.15	14.27	0	7.08±4.40
A(KT)	0	12.04	12.49	12.24	9.33	17.93	13.06	12.01	27.30	10.24	11.64±7.91
PH(BT)	0	0	0	47.93	0	87.34	27.48	0	154.75	0	31.75±52.22
PH(LT)	0	5.83	7.13	8.64	5.71	9.56	8.31	5.72	12.68	0	6.35±3.96
PH(KT)	0	0	0	0	0	0	0	0	0	0	0
N(BT)	0	26.53	0	47.93	0	58.23	28.46	0	89.85	0	25.1±31.57
N(LT)	0	0	0	7.42	0	8.88	0	0	10.49	0	2.67±4.37
N(KT)	0	0	0	4.08	3.90	8.16	3.58	0	8.53	0	2.82±3.40

408 Anthracene: A, Phenanthrene: PH, Naphthalene: N, L: Liver, K: Kidney, B: Blubber, F: Female, M: Male

**Table 2:** Results of ANOVA test on concentration of PAHs (ppb w.w.) in different tissue samples

	Blubber	P	Liver	Kidney
Anthracene	84.83±79.86**	0.001	7.08±4.40*	11.64±7.91*
Phenanthrene	31.75±52.22**	0.001	6.35±3.96*	0 <sup>b</sup> *
Naphthalene	25.1±31.57**	0.01	2.67±4.37*	2.82±3.40*


\* Unsignificant difference, \*\* Significant difference (p<0.05)

**Table 3:** Results of Student's t-tests between sex of sampled seals and concentration of PAHs (ppb w.w.) in tissue samples

PAHs concentration in liver, kidney and blubber tissues	P	t-value
Concentration of anthracene in blubber tissue	0.00*	0.46
Concentration of anthracene in liver tissue	0.03*	0.08
Concentration of anthracene in kidney tissue	0.04*	3.70
Concentration of phenanthrene in blubber tissue	0.00*	3.58
Concentration of phenanthrene in liver tissue	0.04*	2.28
Concentration of naphthalene in blubber tissue	0.00*	5.95
Concentration of naphthalene in liver tissue	0.01*	3.65
Concentration of naphthalene in kidney tissue	0.03*	8.83

\* Significant difference (p<0.05)

**Fig 1:** Mean (SD) PAHs concentration (ppb w.w.) in male and female Caspian seals



421


422 F: female, M: male

**Fig 2:** Results of liner regression test on relationship between PAHs concentration (ppb w.w.) and age (year) of male and female

424 Caspian seals

425

# *PAHs exposure in Caspian seals*



a

A: anthracene, N: naphthalene, PH: phenanthrene, BT: blubber tissue, LT: liver tissue, KT: kidney tissue