

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Backdating systematic shell ornament making in Europe to 45,000 years ago

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Backdating systematic shell ornament making in Europe to 45,000 years ago / Simona Arrighi, Eugenio Bortolini, Laura Tassoni, Andrea Benocci, Giuseppe Manganelli, Vincenzo Spagnolo, Luca Maria Foresi, Anna Maria Bambini, Federico Lugli, Federica Badino, Daniele Aureli, Francesco Boschin, Carla Figus, Giulia Marciani, Gregorio Oxilia, Sara Silvestrini, Anna Cipriani, Matteo Romandini, Marco Peresani, Annamaria Ronchitelli, Adriana Moroni, Stefano Benazzi. - In: ARCHAEOLOGICAL AND ANTHROPOLOGICAL ACHINGTRY: ISSN 1866-9557. - ELETTRONICO. - 12:(2020), pp. 59.1-59.22. [10.1007/s12520-019-00985-3] This version is available at: https://hdl.handle.net/11585/724188 since: 2020-03-04

Published:

DOI: http://doi.org/10.1007/s12520-019-00985-3

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

1	
2	This is the final peer-reviewed accepted manuscript of:
3	Simona Arrighi, Eugenio Bortolini, Laura Tassoni, Andrea Benocci, Giuseppe
4	Manganelli, Vincenzo Spagnolo, Luca Maria Foresi, Anna Maria Bambini, Federico
5	Lugli, Federica Badino, Daniele Aureli, Francesco Boschin, Carla Figus, Giulia
6	Marciani, Gregorio Oxilia, Sara Silvestrini, Anna Cipriani, Matteo Romandini, Marco
7	Peresani, Annamaria Ronchitelli, Adriana Moroni, Stefano Benazzi
8	Backdating systematic shell ornament making in Europe to 45,000 years ago
9	Archaeological and Anthropological Sciences
10	The final published version is available online at:
11	https://doi.org/10.1007/s12520-019-00985-3
12	
13	
14	
15	Rights / License:
16 17	The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.
18	
19	
20	
21	
22	
23	

24	Simona Arrighi <sup>a,b,c*</sup> , Eugenio Bortolini <sup>a</sup> , Laura Tassoni <sup>d</sup> , Andrea Benocci <sup>b</sup> , Giuseppe Manganelli <sup>b</sup> ,
25	Vincenzo Spagnolo <sup>b,c</sup> , Luca Maria Foresi <sup>b</sup> , Anna Maria Bambini <sup>b</sup> , Federico Lugli <sup>a,e</sup> , Federica
26	Badino <sup>a,f</sup> , Daniele Aureli <sup>b,c,g</sup> , Francesco Boschin <sup>b,c</sup> , Carla Figus <sup>a</sup> , Giulia Marciani <sup>a,b</sup> , Gregorio
27	Oxilia <sup>a</sup> , Sara Silvestrini <sup>a</sup> , Anna Cipriani <sup>e,h</sup> , Matteo Romandini <sup>a,d</sup> , Marco Peresani <sup>d</sup> , Annamaria
28	Ronchitelli <sup>b</sup> & Adriana Moroni <sup>b,c,i</sup> , Stefano Benazzi <sup>a,j</sup>
29	
30	Backdating systematic shell ornament making in Europe to 45,000 years ago
31	
32	<sup>a</sup> Dipartimento di Beni Culturali, Università di Bologna,48 121 Ravenna, Italy, <sup>b</sup> Dipartimento di
33	Scienze Fisiche della Terra e dell'Ambiente, Università di Siena, 53100 Siena, Italy <sup>c</sup> Centro Studi
34	sul Quaternario, 52037 Sansepolcro, Italy, <sup>d</sup> Dipartimento di Studi Umanistici, Università di Ferrara,
35	44100 Ferrara, Italy, <sup>e</sup> Dipartimento di Scienze Chimiche e Geologiche, Università di Modena e
36	Reggio Emilia, 41125 Modena, <sup>f</sup> C.N.R Istituto per la Dinamica dei Processi Ambientali, 20126
37	Milano, Italy, <sup>g</sup> UMR 7041 ArScAn Equipe An-TET, Université Parigi Ouest Nanterre La Défense,
38	F-92023 Nanterre, France, h Lamont-Doherty Earth Observatory, Columbia University, Palisades,
39	New York, 10964, USA, <sup>i</sup> Istituto Italiano di Paleontologia Umana, 03012 Anagni (FR), Italy, <sup>j</sup>
40	Department of Human Evolution Max Planck Institute for Evolutionary Anthropology, 04103
41	Leipzig, Germany.
42	*Corresponding author (e-mail: simona.arrighi@unibo.it)
43	
44	

## 45 Abstract

46 Personal ornaments are commonly linked to the emergence of symbolic behaviour. Although their47 presence in Africa dates back to the Middle Stone Age, evidence of ornament manufacturing in

Eurasia are sporadically observed in Middle Palaeolithic contexts, and until now large scale 48 diffusion has been until now well documented only since the Upper Palaeolithic. 49 Nevertheless, little is known during the period between ca.50,000 and 40,000 years ago (ka), when 50 51 modern humans colonized Eurasia replacing existing hominin populations such as the Neandertals, and a variety of so called "transitional" and/or early Upper Palaeolithic cultures emerged. Here we 52 present shell ornaments from the Uluzzian site of Grotta del Cavallo in Italy, southern Europe. Our 53 results show evidence of a local production of shell beads for ornamental purposes as well as a 54 trend toward higher homogeneity in tusk bead shape and size over time. The temporal interval of 55 the layers of interest (45-40 ka) makes Cavallo the earliest known shell ornament making context in 56 Europe. 57

58 Key words: Uluzzian, Italy, Grotta del Cavallo, shell ornaments

59

#### 60 1.Introduction

61

The use of personal ornaments in human history is a key issue to understand the evolutionary processes that led to modern humankind beyond a biological perspective. Over the past decades various hypotheses have been expressed on this topic, based on the different perspectives of anthropology, ethnography, sociology, and linguistics (White 1989; d'Errico et al. 2003a; McElreath et al. 2003; Abadia and Nowell 2015).

Personal adornments are interpreted as markers of population structure from a cultural/linguistic and social point of view since the beginning of the Upper Palaeolithic (Vanhaeren and d'Errico 2006; Stiner et al. 2013), but they are also proof of an initial aesthetic perception (Leroi-Gourhan 1964) and their presence is the best archaeological evidence of shared symbolic language linked to individual appearance (White 1989; d'Errico et al. 2003a). Unfortunately, the vast majority of deliberate body modifications (including tattoos, scarifications, ear piercing, lip and neck stretching and other numerous kinds of alterations aimed to modify one's own physical appearance) are
embedded in perishable materials and cannot be retrieved from Palaeolithic contexts. This makes
high concentrations of non-functional elements – such as the one presented here - even more
relevant and informative on the emergence of socially recognized symbolic expressions (which may
or may not have involved other kinds of body alterations).

The earliest known evidence of ornaments is associated with Modern Humans (MHs), and dates 78 back to 135-70 ka in Africa and in the Levant (Vanhaeren et al. 2006; d'Errico et al. 2009; 79 Vanhaeren et al. 2013; d'Errico and Backwell 2016). In the latter region, evidence of personal 80 ornaments is still consistently found in the period ranging from the Levantine Middle to Upper 81 82 Palaeolithic (Bosch et al. 2019), while there is broad consensus on the fact that, in Europe, the 83 systematic production of ornaments became a substantial component of material culture since the Protoaurignacian, an Upper Palaeolithic culture dated between 42-36 ka and confidently attributed 84 to MH (Benazzi et al. 2015). There is no doubt, however, that during the Middle to Upper 85 Palaeolithic transition (50-40 ka) ornaments were already becoming more common in the region. 86 Such a trend documents a change in human social behavior compared to the scanter evidence of 87 ornaments derived from animals and seashells attested in Middle Palaeolithic contexts attributed to 88 Neandertals (Peresani et al. 2011; Peresani et al. 2013; Romandini et al. 2014; Radovčić et al. 2015; 89 90 Hoffmann et al. 2018).

In this timeframe late Neandertals and the newly arrived modern human groups coexisted in 91 Europe, and the archaeological record shows the emergence of new techno-complexes (the so-92 93 called transitional complexes) which exhibit a number of innovative features (such as formal bone tools manufacturing, the systematic use of ornaments and pigments, etc...) compared to late 94 Mousterian assemblages (d'Errico et al. 2003a, 2003b; Zilhão 2006; Flas 2011; Boscato and 95 Crezzini 2012; d'Errico and Banks 2013; Moroni et al. 2013 and 2018; Škardla, 2013; Hublin 96 2015). Attribution of transitional industries to either Neandertals or modern humans is often 97 difficult because of the scarcity of human fossils found in association with elements of the material 98

culture. Labelling these industries as "transitional" entails that they express admixed Middle and 99 100 Upper Palaeolithic features and are the product of late Neandertal groups which interacted with modern humans or independently acquired modern behaviours (Hublin 2015). 101 More recent research, however, questioned (Bar-Yosef 2007; Bar-Yosef and Bordes 2010; Benazzi 102 et al. 2011; Moroni et al. 2013 and 2018; Gravina et al. 2018) the association of Neandertals with 103 both Uluzzian (Zilhão et al. 2015, Villa et al. 2018) and Châtelperronian material cultures (Lévêque 104 and Vandermeersch 1980; Hublin et al. 1996; Welker et al. 2016). In particular, the Uluzzian is now 105 considered to be made by modern humans (Benazzi et al. 2011, Moroni et al. 2013 and 2018; Sano 106 et al. 2019 but see contra Zilhao et al. 2015, Villa et al. 2018) on the basis of two deciduous teeth 107 recovered at Grotta del Cavallo and attributed to *Homo sapiens*. Uluzzian assemblages display 108 109 explicitly modern elements, such as the systematic presence of bone tools, colouring substances, 110 ornaments, and a lithic technology that does not show any evidence of continuity with the Mousterian technological tradition (Ronchitelli et al. 2009; d'Errico et al. 2012; Peresani et al. 111 2016; Moroni et al. 2018; Peresani et al. 2019a; Sano et al. 2019). 112 Archaeological assemblages of continental Europe dated to the Middle to Upper Palaeolithic shift 113 yielded sporadic evidence of ornaments and ornament manufacturing on animal bone and teeth 114 (d'Errico et al. 1998; Zilhão 2007), with the exception of Châtelperronian contexts in France which 115 also included a few seashells (Zilhão 2007) (Fig.1). On the other hand, personal ornaments 116 uncovered in Uluzzian assemblages (~45-40 ka) of Mediterranean Europe only consist of seashells 117 (Palma di Cesnola 1989; Fiocchi 1998; Stiner 2010). Findings of this kind are present and 118 sometimes abundant at Klissoura (Greece; n=32), Grotta della Cala (southern Italy; n=30), and 119 Riparo Broion (northern Italy; n=6). Grotta del Cavallo (southern Italy) stands absolutely out with 120 531 seashells distributed across the entire Uluzzian sequence, dated between 45.5 - 39.9 ka 121 (Benazzi et al. 2011; Douka et al. 2014; Moroni et al. 2018). 122



Fig. 1 Distribution of ornaments dated between ca. 45 and 39 ka across transitional contexts in 124 Europe (sites with debated cultural attribution are represented here, see "Discussion" section). Pie 125 radius is proportional to the total size of ornamental assemblages, while colors indicate the relative 126 proportion contributed by different classes of items. (1) Saint-Césaire, (2) Quinçay, (3) Roc de 127 Combe, (4) Caune de Belvis, (5) Cova Foradada, (6) Roche au Loup, (7) Grotte du Trilobite, (8) 128 Grotte des Fées, (9) Grotte du Renne, (10) Trou Magrite, (11) Ilsenhöle Ranis, (12) Willendorf II, 129 (13) Riparo Broion, (14) Grotta della Cala, (15) Grotta del Cavallo, (16) Klissoura Cave, (17) 130 Franchthi Cave, (18) Bacho Kiro, (19) Brynzeny I 131

132

Nonetheless, a systematic study of the Uluzzian shell assemblage was never undertaken. The abundance of shell beads found at Grotta del Cavallo offers a privileged perspective for a systematic reassessment of ornament making in this and other Uluzzian assemblages. The key question is therefore to understand whether systematic manufacturing and personal use of shell beads took place at the site, and therefore if Cavallo back dates to ~45 ka the beginning of one of the most salient traits of Upper Palaeolithic Europe.

Here we present a thorough analysis of the entire sequence of shell assemblages found in theUluzzian layers at Grotta del Cavallo. At this site the Uluzzian can be followed from the archaic

(layer EIII) to the evolved-late phase (layer EII-I and D), sandwiched by the tephra Y-6 at  $45.5 \pm 1.0$ 141 ka (Zanchetta et al. 2018) and Y-5 (Campanian Ignimbrite) at  $39.85 \pm 0.14$  ka (Giaccio et al. 2017; 142 Zanchetta et al. 2018). All dates obtained from shells (5 Antalis sp. from EII-I, spit E-D, DII and 143 DIIb, 1 Lembulus pella from DIIb - Benazzi et al. 2011) are consistent with the chronological frame 144 established by the above mentioned tephra layers. Previous works maintained that the presence in 145 the Uluzzian deposit of important post-depositional disturbances might undermine the association 146 between Uluzzian assemblage and modern humans (Gioia 1990; Zilhão 2007; Zilhão et al. 2015). 147 Nevertheless, more recent research has definitely confirmed the integrity of the archaeological 148 deposit in which the teeth attributed to MHs were recovered and therefore the reliability of their 149 150 association with the Uluzzian materials (Moroni et al. 2018; Zanchetta et al. 2018; see 151 Supplementary Information Appendix, Section I).

Here, we ascertain the presence of anthropogenic intervention on scaphopods (also known as tusk shells), gastropods, and bivalves as well as their use as personal ornaments through the combined use of quantitative analysis of morphology, experimental tests, use-wear, Sr isotopes, and micropalaeontological examination. Results, which are compared with the broader picture emerging from other Uluzzian contexts, show evidence of an increasing regularity in shape and size of the beads, as well as of their local production for ornamental purposes.

158

159

### 160 2.Materials - The shell assemblage from Grotta del Cavallo

The Uluzzian layers of Cavallo yielded 618 shell remains (531 NISP), mostly retrieved in the uppermost layers DII and DI. Among them, 285 can be assigned to 32 *taxa* (Table 1 and Fig. 2) while the majority is still undetermined due to their bad state of preservation. Among identified classes, scaphopods are the most represented and ubiquitous one across the entire sequence, while gastropods and bivalves are considerably less abundant and clustered in the upper layers DI and DII (n=124, of which 28 are pierced). The lowermost Uluzzian layer yielded, in addition to 67

- scaphopods, 7 bivalves and 33 gastropods, one of which consists of a pierced *Tritia neritea*
- 168 (Fig.2b). Species richness increases over time and reaches its maximum in the uppermost layers
- 169 with 22 identified species encompassing both edible and not edible species. The most represented
- 170 species are *Tritia neritea*, *Lembulus pella* and *Glycymeris* sp.
- 171

Taxa	E III	EII-I	E-D	DII	DI	D (DI+DII)	Total
Gasteropode undet.	21	-	-	-	-	-	21
Patella rustica *	2	-	-	-	-	-	2
Jujubinus striatus	-	-	-	-	2	-	2
Phorcus sp.	-	-	-	-	1	-	1
Phorcus turbinatus *	-	-	-	1 (1)	-	-	1 (1)
Homalopoma sanguineum	-	-	-	2	8 (1)	1 (1)	11 (2)
Bittium reticulatum	1	-	-	-	-	-	1
Cerithium sp.	-	-	-	-	1	2	3
Cerithium vulgatum *	3	-	-	-	1	1	5
<i>Turritella</i> sp.	3	-	-	-	2	2	7
Melaraphe neritoides	1	-	-	-	-	-	1
Trivia pulex	-	-	-	-	1	-	1
Naticarius hebraeus *	1	1	-	-	-	-	2
Euspira catena	-	-	1	-	1	-	2
Aporrhais pespelecani*	-	-	-	1	-	-	1
Tritia sp.	-	-	-	-	-	1	1
Tritia cuvierii/unifasciata	-	-	-	-	2	-	2
Tritia neritea	1 (1)	-	-	5 (4)	22 (5)	4 (2)	32 (12)
Tritia nitida	-	-	-	1 (1)	-	-	1 (1)
Columbella rustica *	-	-	-	-	1	2 (1)	3 (1)
Total gastropods	33 (1)	1	1	10 (6)	42 (6)	13 (4)	100 (17)
Lembulus pella	-	-	-	-	10 (10)	-	10 (10)
Mytilus galloprovincialis *	-	-	-	-	27	-	27

Glycymeris sp.	1	-	-	2	13 (2)	1	17 (2)
Glycymeris nummaria *	-	-	-	-	-	1	1
Pecten jacobaeus *	-	-	-	-	1	-	1
Spondylus gaederopus *	-	-	1	-	-	-	1
Acanthocardia sp. *	6	-	-	-	1	-	7
Callista chione *	-	1	-	-	2	-	3
Corbula gibba	-	-	-	-	1	-	1
Total bivalves	7	1	1	2	55 (12)	2	68 (12)
Antalis sp.	10	11	13	63	72	77	246
Antalis vulgaris	32	1	2	23	15	6	79
Antalis dentalis/inaequicostata	25	6	2	1	-	3	37
Fissidentalium rectum	-	-	-	-	-	1	1
Total scaphopods	67	18	17	87	87	87	363
Total NISP	107	20	19	99	184	102	531
Undetermined	8	4	-	-	52	23	87
Total NR	115	24	19	99	236	125	618

Table 1 Malacological assemblage from Grotta del Cavallo. Edible species are marked with an asterisk. Numbers in brackets are for pierced specimens. The taxonomic analysis is based on the updated datasets available online on the World Register of Marine Species (WoRMS; <u>www.marinespecies.org</u>), also the number of rests (NR) and the number of identified specimens (NISP) have been used in order to define the right amount of the assemblage

178

179

Shells from layer EIII were mostly found in an area of about 4 square meters (squares E 13, F11,
F12 and G 11) (94.1%) while only 5.9% comes from the 1963-66 excavation trench corresponding
to an area of about 2.5-3 x 3.5 m (Moroni et al., 2018) (Table 2; see *SI Appendix, Section I* for
details and Fig.S1).

Shells from layer EII-I were retrieved in square H11 and in the 1963-1966 excavation trench (Table
2). Shells from layers DII and DI come almost exclusively from two square meters (H11 and H7 Fig. S1). Only very few specimens from both layers were found in the 1963-66 excavation trench
(Table 2).

188

	EIII	EII-I	E-D	DII	DI	D whole	Total
E-F-G8*; E-F9*; E-F10* (1963-66 excavation trench)	5 (4)	5 (5)	-	21 (17)	3	-	34 (26)
E13*	17 (1)	-	-	-	-	-	17 (1)
E11	2	-	-	-	-	-	2
F13*	-	-	-	-	-	-	-
F12*	12 (4)	-	-	-	-	-	12 (4)
F11*	18 (12)	-	-	-	-	-	18 (12)
G11*	59 (46)	-	-	-	-	-	59 (46)
G10*	-	-	-	-	-	-	-
G7	-	-	-	-	-	-	-
H11	-	17 (13)	17 (15)	78 (70)	233 (87)	102 (74)	447 (259)
H8*	-	-	-	-	-	-	-
H7*	2	2	2 (2)			23 (13)	29 (15)
Total	115 (67)	24 (18)	19 (17)	99 (87)	236 (87)	125 (87)	618 (363)

190 Table 2 Spatial distribution of the shells by layer. For the squares marked with an asterisk only the

- 191 portion occupied by undisturbed deposit has been considered (for further details see *SI Appendix*,
- 192 *Section I* and Fig.S1). Numbers in brackets are the scaphopods



- 194 Fig. 2 Selection of shells from Grotta del Cavallo. a Antalis sp. b Tritia neritea. c Antalis sp. d
- 195 Antalis sp. e Tritia neritea. f Homalopoma sanguineum. g Columbella rustica. h Phorcus
- 196 turbinatus. i Lembulus pella. J Glycymeris sp.

#### 199 **3. Methods**

The present research mostly focuses on tusk shells due to their abundance across all layers. Their bad state of preservation, however, required the design of experimental and quantitative analyses to: a) understand whether these shells were collected from fossiliferous deposits rather than from thanatocoenoses (namely 'gathered on the beach); b) ascertain the presence and the effects of anthropogenic breakage and post-depositional processes; c) analyse non-macroscopic morphological change over time.

206

#### 207 **3.1 Micropalaeontological and isotope analysis**

The micropalaentological content of several samples of sediment contained in the tusks from Grotta del Cavallo and in the Uluzzian layers of the cave was examined to establish if the shells were collected from nearby sedimentary outcrops or from beach deposits contemporaneous to the Uluzzian activity at the site. The material was examined through classic micropalaentological techniques, which include the preparation of washed samples for the analysis of microfossils and of smear slides for calcareous nannofossils, as well as observation under a microscope (*SI appendix, Section II*).

In situ trace element analysis was carried out by means of LA-ICP-MS on eight samples of tusk
 specimens from layers E and D of Grotta del Cavallo to test their diagenetic preservation, and
 <sup>87</sup>Sr/<sup>86</sup>Sr ratio was analyzed by MC-ICP-MS to test the non-fossil origin of the archaeological
 samples (see Lugli et al. 2017) (*SI appendix, Section III*). Elemental and isotope analyses were
 conducted at the Centro Interdipartimentale Grandi Strumenti of the University of Modena and
 Reggio Emilia.

221

#### 222 **3.2 Morphological analysis**

All the archeological tusk shells were measured (length, maximum and minimum diameters) in accordance with normalized zoological parameters: the apex of the shell is anatomically dorsal, the large aperture is ventral and anterior, and the concave side is anatomically dorsal (Shimek and Steiner 1997).

- The archeological and experimental tusk shells were classified according to the type of apical andbasal fractures. Seven different types of fractures were described (Fig. 3 and Fig. S5):
- a)Rectilinear: regular and straight fracture, perpendicular to the longest axis of the shell;
- b)Oblique: regular and straight fracture, oblique to the longest axis of the shell;
- c)Symmetric-irregular: uneven fracture forming a symmetric profile;
- d)Asymmetric-irregular: uneven fracture;
- e)Flute-mouth: uneven fracture taking the shape of a flute mouth;
- f)Notch: fracture forming a notch;
- g)Rectangular notch.



Fig. 3 Type of fractures. Classification according to the type of fracture on the experimental

- material: rectilinear (a), oblique (b), symmetric irregular (c), asymmetric irregular (d), flute-mouth
  (e), and notch (f). The bar is 1 mm
- 240

241 The degree of the fracture's invasiveness and of the discontinuity of the fractured edge were also

described. Rectilinear and oblique fractures, which usually have a straight edge, are degree 0.

Jagged edges have a degree value included between 1 and 3, according to invasiveness of the

alteration (Fig. 4).



245

Fig. 4 Fracture degrees. Flute-mouth fracture with degree 1 (a). Flute-mouth fracture with degree 2
(b). Flute-mouth fracture with degree 3 (c). The bar is 2 mm

249 Specimens were classified according to their remnant portion (entire, apical, central and basal).

Given the general bad preservation of the archaeological shells, this classification is not based on

251 zoological parameters. We consider entire or nearly entire the specimens where all the

morphological features can be recognized. The apical (or distal) elements are those in which the

253 posterior end is recognizable. Central (or mesial) portions are usually considered those elements

showing scarce difference between maximum and minimum diameter. In the basal (or proximal)

elements the base is present.

The possibility that a thread could go through the apical hole was considered as a critical

discriminant factor. In this case, apical hole refers to the actual empty space within the shell, and

258 may or may not coincide with the minimum diameter. A minimum threshold of 0.5 mm was

established for this condition to be met, considering that sturdy horsehairs, the smallest suitable

strings documented in ethnographic studies (Orchard 1929), present with diameters ranging from

261 0.08 to 0.4 mm (Craveri 1931). Of course, other materials, such as vegetal fibers or sinews, cannot

262 be excluded as potential strings for suspension.

#### 263 **3.3 Experimental tests**

Experimental tests for shell beads production and post-depositional damage was carried out only for tusk shells, due to their relevance among the whole assemblage and to the difficulty in evaluating the origin of the fracturing on this kind of shell.

267 The experimental program involved the gathering of current specimens (mostly *Antalis vulgaris*) in

the surroundings of Grotta del Cavallo, their breaking by bending, sawing, crushing (direct

269 percussion) and trampling (for further details see SI Appendix, Section IV). The experimental shells

were measured and classified according to the same parameters used for the archaeological ones,

271 before and after each experimental test.

Possible technological attributes of gastropods and bivalves were instead evaluated by comparison
with literature data and available reference collections (Taborin 1993; Benghiat et al. 2009; Stiner et al. 2013; Vanhaeren et al. 2013; Tátá et al. 2014).

275

#### 276 **3.4 Exploratory data analysis**

Significant differences in unique variables such as length, minimum diameter, and maximum 277 diameter across layers were assessed through non-parametric tests due to small sample size and to 278 the presence of violations of the assumptions of parametric tests. More specifically, two-tailed 279 280 Mann-Whitney U test for independent study design was performed when only two groups were compared, while Kruskal-Wallis test was used when more than two groups were to be compared 281 against each other. Associations between pairs of categorical variables were explored on 282 283 contingency tables via Correspondence Analysis using the function ca of the package ca (Nenadic and Greenacre 2007) and through Pearson's Chi-square test for independence. All analyses were run 284 in R version 3.4.4 (R Core Team 2018). 285

286

#### 287 **3.5 Diversity and similarity**

Intra-layer diversity was measured through Gini-Simpson index of diversity and its numbers 288 289 equivalent (i.e. the exponential of Gini-Simpson index, following Jost 2006 and 2007 based on the relative frequency of proximal fracture types, distal fracture types, and remnant portions obtained 290 for each layer. Results were obtained using respectively the functions H and d contained in the 291 package vegetarian (with order of diversity q=2; Charney and Record, 2012). Similarity between 292 layers was measured through Morisita-Horn index of overlap using the function sim.table of the 293 package vegetarian (with order of diversity q=2). Clustering of layers was obtained using 294 hierarchical clustering with Ward's clustering criterion (Murtagh and Legendre 2014), i.e. 295 dissimilarities were squared before cluster updating. All analyses were run in R version 3.4.4. 296

297

#### 298 **3,6 Taphonomic analysis**

The taphonomic study of the malacological assemblage was performed on the NISP and it focused on three main kinds of alterations: pre-depositional alterations (e.g. marine abrasion, predation by other molluscs and bioerosion); intentional/unintentional anthropogenic alterations (e.g. thermic alterations, ochre traces and human modifications due to consumption or ornamental purpose); and post-depositional transformations (e.g. fragmentations and decalcification).

304

#### **305 3.7 Use-wear analysis**

The shells were analyzed by means of a Hirox KH 7700 3D digital microscope (Arrighi and Borgia 2009) using two different optics: a MX-G 5040Z zoom lens equipped with an AD-5040 Lows objective lens (20-50x) and a coaxial vertical lighting MXG-10C zoom lens and an OL-140II objective lens (140-560x). The analytical criteria (surface polishing, rounding, faceting) for the functional interpretation of the ornaments build upon previous analyses of prehistoric shells described in literature (e.g. Taborin 1993; Vanhaeren and d'Errico 2001; Bonnardin 2007).

313 **4. Results** 

### 314 **4.1 Taphonomic processes**

315 The shell assemblage from Grotta del Cavallo exhibits traces of decalcification (Table 3), in

particular in the case of tusk shells. Among the latter, some specimens exhibit partially exfoliated

surfaces that probably facilitated post-depositional fragmentation. Sea wash is also documented.

318 Only a few specimens are affected by alterations due to thermic effects, predator activity or

319 reducing conditions in soil.

Layers/ Spit	Classes	NISP	Sea washed	Boring sponge	Predators	Burned	Decalc.	Post-dep. cracks
	Gastr.	33	25	_	1	1	28	14
E III	Biv.	7	7	-	-	-	6	5
	Scap.	67	7	-	-	-	37	3
	Gastr.	1	-	-	-	-	1	1
E II-I	Biv.	1	1	-	-	-	1	-
	Scap.	18	-	-	-	-	17	2
	Gastr.	1	1	-	-	-	1	-
E-D	Biv.	1	1	-	-	-	1	1
	Scap.	17	-	-	1	-	16	-
	Gastr.	10	10	-	-	-	6	1
D II	Biv.	2	2	-	-	-	2	2
	Scap.	87	-	-	-	1?	79	3
	Gastr.	42	37	1	6	5	19	12
DI	Biv.	45	31	-	10	-	26	17
-	Scap.	87	5	-	-	-	85	-
D whole	Gastr.	13	11	_	-	1	9	7

	Biv.	2	2	1	-	-	2	1
	Scap.	87	1	-	2	-	73	1
Total %	-	521	27.1%	0.4%	3.6%	1.2%	78.5%	13.4%
Total		521	141	2	20	7	409	70

Table 3 Taphonomy of the malacological assemblage from Grotta del Cavallo. Anthropic traces are not included in this table, as they are discussed in the following chapters

324

#### 325 4.2 Scaphopods

326 <u>4.2.1 Fossil and non-fossil origin of the archaeological specimens.</u>

327 Since *Fissidentalium rectum* is an extinct species, this specimen was probably collected at a

fossiliferous Pleistocene formation cropping in the Salento region (Largiolli et al. 1969).

329 Both micropaleontological analyses (SI Appendix, Section II) carried out on the sediment preserved

inside the tusks, and Sr isotopic analysis (*SI Appendix, Section III*), excluded a fossil origin of all

the other *taxa*.

332

#### 333 <u>4.2.2 Experimental study</u>

334 Manufacturing was experimentally tested on complete and almost complete specimens collected on

the present-day beach in the surroundings of Grotta del Cavallo. Naturally broken specimens

usually lack the apical portion. Fractures are frequently asymmetric-irregular (~60%) (see

paragraph *Methods* and the *SI Appendix, Section IV* for description of the methodology applied).

Rectilinear (~24%) and oblique (~8%) fractures are less frequent. Many elements have their natural

basis preserved. The fracturing degree is predominantly  $1 (\sim 46\%)$  and, to a lesser extent,  $2 (\sim 24\%)$ .

340 Experimental manufacturing revealed an association (see *SI Appendix, Section V*) of rectilinear and

oblique fractures with bending (~91%) and sawing (~46%), as well as an association between

irregular fractures and crushing (~60%). Unlike other actions, sawing can produce a tiny
rectangular notch due to the pressure exerted on the starting point.

Notch fractures, which can be considered lesser developed flute-mouth fractures, cannot be clearly related to a specific bead making technique. These kinds of fractures rarely occur, although their frequency is a bit higher in crushing tests. Vanhaeren and d'Errico (2001) suggest that notch and flute-mouth fractures originate when a needle is used for stringing a thread into a shell. Comparable notch fractures, however, were obtained from our experimental trampling tests and from some explorative suspension tests when one tusk was inserted into the other, as also observed in other research (Álvarez Fernández 2006).

The occurrence of clear cut-marks on the shells is consistently associated with sawing. Bending is mostly related to fractures of degree 0 (85%) and to a lesser extent degree 1 (~14%), sawing almost exclusively to degree 0 (~46%) and 1 (~50%), whereas the effect of crushing is more variable, including degrees 1 (~36%), 2 (50%) and 3 (3%).

Experimental trampling did not produce substantial evidence of breakage, although it generally results in a more conspicuous presence of chipping at the base of the shell. Fractures associated with trampling are mostly asymmetric-irregular and to a lesser extent flute-mouth and notch usually with degree 2 or 3.

359

#### 360 <u>4.2.3 Morphological and use-wear analysis</u>

Scaphopods are the most recorded class in all the Uluzzian layers of Grotta del Cavallo. When
identified at the species level, they are *Antalis dentalis/inaequicostata* and *Antalis vulgaris* (Table
1). Noteworthy is the presence of a single fossil specimen, probable *Fissidentalium rectum* in the
entire macro-layer D.

Central and apical portions are the most common ones in all the layers (*SI Appendix, Section VI* Table S14). In the uppermost layers (D), short apical portions are particularly abundant (32 out of 261), usually truncated with rectilinear fractures at their base (Fig. 5). This evidence suggests the

368 systematic removal of the shell apical portions by flexing, in order to obtain segments with a

diameter as wide as possible to allow the passage of the string. For this reason, these short apical

370 fragments could be interpreted as waste products.



Fig. 5 Waste products. Small apical portions interpreted as waste products

373

371

374 Entire or almost entire specimens are considerably less frequent (SI Appendix, Section VI Table S14). Among these, specimens preserving the apical portions (tot=91; nEIII=20, nEII-I=5, nD=66) 375 are not suitable for suspension, since their apical holes are too narrow to be stringed. 376 Rectilinear fractures, followed by irregular asymmetric ones, are the dominant types in all layers 377 (Table 4, *SI Appendix, Section VI* Table S15-16). In layer EII-I there is a slight predominance of 378 flute-mouth fracture (~25 %) and rectilinear fracture (25%) compared to and asymmetric irregular 379 ones (~16%; SI Appendix, Section VI Tables S15 and S16; Fig S4). Rectangular notch fractures 380 related to sawing are attested on 1 specimen from layer EII-I and on 2 specimens respectively from 381 DII and D whole (SI Appendix, Section VI Fig. S5). None of these shells exhibit cut-marks. 382 Scratches comparable to those experimentally produced by cutting with a flint tool (SI Appendix, 383 Section VI Fig. S5 and Fig 6) are recorded only on one tusk (layer DII), which displays rectilinear 384 fractures at both ends and some sort of marks overlapping with the basal one. 385 386

Layer/	Entire	Rectilinear	Oblique	Asymmetric-	Symmetric-	Flute-	Notch	Rect.	Unid.
Spit				irregular	irregular	mouth		notch	
EIII	12.6%	38.8 %	6.7 %	23.1%	7.4%	2.9%	8.2%	0%	0%
	(17)	(52)	(9)	(31)	(10)	(4)	(11)	(0)	(0)

EII-I	8.3%	25%	5.5%	16.6%	8.3%	25%	8.3%	2.7%	0%
	(3)	(9)	(2)	(6)	(3)	(9)	(3)	(1)	(0)
E-D	5.8%	32.3%	5.8%	23.5%	0%	11.7%	17.6%	0%	2.6%
	(2)	(11)	(2)	(8)	(0)	(4)	(6)	(0)	(1)
DII	12.6%	37.3%	5.1%	24.7%	2.8%	7.4%	8%	0,5%	1.1%
	(22)	(65)	(9)	(43)	(5)	(13)	(14)	(1)	(2)
DI	13.7%	40.8%	4%	29.8%	1.1%	5.7%	2.8%	0%	1.7%
	(24)	(71)	(7)	(52)	(2)	(10)	(5)	(0)	(3)
D	10.9%	43.1%	2.8%	28.7%	2.2%	8%	2.8%	0.5%	0.5%
whole	(19)	(75)	(5)	(50)	(4)	(14)	(5)	(1)	(1)

387	Table 4 - Frequencies of the fractures identified on tusk shells by layer. Real data are reported in

388 brackets

389

The majority of fractures exhibit degree 0 (50.8%), and frequency tends to decrease with increasing fracture degree (degree 1=22.8%; degree 2=19.8%; degree 3=6.0%) *SI Appendix, Section VI* Tables

392 S17 – S22). A different trend is registered in layer EII-I, where degree 2 is the predominant one.



Fig. 6 Traces of anthropogenic manipulation. Grotta del Cavallo. Layer DII—tusk shell with cutmarks (a), this kind of marks is very similar to those obtained experimentally by cutting tusk shells
with a flint tool (b). Layer EIII—tusk shell with a notch fracture showing well rounded (c) and

polished edges (d). Layer EIII—tusk shell showing a notch fracture with flattened (e) and polished
areas (f). Layer EIII—weak polishes inside a notch fracture (g) and weak polishes inside a notch
fracture produced during suspension experiment with a leather string (h)

The retrieval of two shells inserted into each other is documented in three cases (layers EII-I, E-D, 401 and DI). This eventuality can be due either to human intervention or taphonomic causes (Fig. 7). 402 Intentional insertion of tusk shells into each other, especially for the making of necklaces, is well 403 documented in both archaeological and ethnographic evidence (Ruppert and Bernet 2001; 404 Dimitrijević 2014). Otherwise, this occurrence was also observed during the gathering of present-405 day tusks used for experimental tests, even if in the whole collected sample (1908 tusks) only one 406 case of insertion was reported. Since the tusks retrieved in layer DI could be easily separated, the 407 parts of the surface originally located inside the external tusk were analyzed. This portion appears 408 considerably better preserved and presents some polished areas, while the exposed surface exhibits 409 410 the usual altered appearance (Fig. 7).



411

400

Fig. 7 Tusk shells inserted one into another from Grotta del Cavallo. Tusk shells inserted one

- 414 into another from layer EII-I (a), spit E-D (b), and layer DII (c), while the external surface of the
- 415 tusks looks weathered (d), and the inner, protected trait shows its original aspect (e)

- 416 Consequently, the insertion can be considered intentional and not due to post-depositional
- 417 processes. Unexpectedly, the same pattern was detectable on the surface of the external shell. In this

418 case two scenarios can be envisaged:

- 419 1. The external shell was in its turn inserted into another shell, which is now lost.
- 420 2.This differential surface preservation could be attributed to post-depositional phenomena acting
  421 with different intensity on the two shells (partial burial for some time?)
- 421 with different intensity on the two shells (partial burial for some time?)
- 422 According to the first hypothesis, we can imagine an adornment object composed of several tusks,

inserted into each other, as largely reported in archaeological and ethnographic contexts (Ruppert

424 and Bernet 2001; Dimitrijević 2014).

- 425
- 426 A number of tusks (11) from layer EIII are coated with red pigment, usually on the whole outer and

427 inner surfaces (Fig 8). Other items (14) preserve only a few residues of red coloring. They were all

- retrieved in two adjacent square meters of the excavation grid (SI Appendix, Section VI Fig. S1) and
- 429 20 out of 25 ochered items are suitable for suspension, possibly suggesting that the shells were
- 430 colored in an ornamental perspective.
- 431 Traces of red pigment were also recovered on a specimen from layer EII-I. The coloring substance432 is distributed inside the tusk and spreads longitudinally (Fig 8).



433

434 Fig. 8 Ochered shells from Grotta del Cavallo. Layer EIII—ocher associated to a notch fracture (a).

435 Ocher and black residues on the external surface of a specimen (b). Ocher on the apical end of a

tusk. The edge is rounded and polished (c). Layer EII-I—ocher located inside the shell and
longitudinally spread (d)

438

Usually, scaphopods are not well-preserved due to decalcification. Only in layer EIII a good number
(30) of tusks exhibits well-preserved surfaces. About half of them (18) show weak traces of
polishing on the outer surface of one or both ends (Fig. 6), often clustered on a limited portion of
the rim (Fig 6). Such traces appear more frequently in central (9) and apical (7) portions, and to a
lesser extent in basal ones (1) and almost complete (1) shells. Polishing traces are associated with
rectilinear, asymmetric, symmetric irregular, and notch fractures.

In the uppermost layers the state of preservation of tusks did not allow for a reliable evaluation of suspension traces. Nevertheless, when the pristine surface of the shell is preserved, slightly rounded edges on one or both extremities can be observed, although it is difficult to identify actual patterns and therefore to discriminate between usage (suspension) and post-depositional processes.
Specimens displaying such traces are 3 in layer EII-I, 3 in split E-D, and 25 in unit D. It usually

450 consists of central portions mainly associated with rectilinear or notch fractures.

451

#### 452 <u>4.2.4 Analysis of variability over time</u>

Quantitative analyses were run on a controlled subset (n=255) obtained excluding observations
uncovered in the E-D split and shells generically attributed either to layer DI or DII. Records with
missing information were also not considered (n=4). In this case, sampled shells are almost equally
distributed across layers with the exception of EII-I (nEIII=67, nEII-I=17, nDII=87, nDI=84).

457 Results show that shell minimum and maximum diameter are significantly wider in *Antalis sp.* than

458 in Antalis vulgaris and Antalis dentalis/inaequicostata (SI Appendix, Section VII Tables S23, S26

and S27). At the same time, maximum diameter increases over time for all the examined portions

460 including apical ones (SI Appendix, Section VII Fig. S6; Tables S30-32), and such an increase is

461 mirrored by significant increase in the minimum diameter of medial pieces (SI Appendix, Section

VII Fig. S6; Table S33). Length of entire shells also increases over time (SI Appendix, Section VII 462 463 Tables S28 and S35, Fig. S6). However, no difference in length in any layers between likely suspended apical pieces and pieces that are less likely to have been used in suspension was recorded 464 (SI Appendix, Section VII Table S34). 465 The distribution of both distal and proximal fracture types significantly differs across layers (Tables 466 S24-25 and S38-40). More in detail, rectilinear and irregular asymmetric fractures are the most 467 frequent types among proximal fractures, and their presence tends to increase over time (SI 468 Appendix, Section VII Table S36). Layer EII-I is associated with "notch", "rectilinear cut", and 469 "flute-mouth" fractures and is segregated from all other layers in multivariate analyses. 470 Correspondence Analysis also suggests a shift over time from irregular/symmetric and oblique 471 fractures to irregular/asymmetric and rectilinear fractures in both distal and proximal fractures. 472 The distribution of remnant portions and that of potentially suspended pieces does not significantly 473 differ from one layer to the other (SI Appendix, Section VII Table S35 and Tables S38-39). The 474 475 distribution of distal fractures is significantly associated with the possibility of being worn in suspension in the whole of the stratigraphic sequence (SI Appendix, Section VII Table S41-43). 476 More specifically, rectilinear fractures are more frequent in potentially suspended pieces, while non-477 478 suspended pieces exhibit no distal fractures in any layers with the exception of two instances of rectilinear fracture in layer DI (SI Appendix, Section VII Tables S44-46). The relative proportion of 479 different fracture types visibly changes over time (SI Appendix, Section VII Tables S47-53). 480 Diversity in proximal and distal fracture types steadily decreases over time, while diversity in 481 remnant portion types tends to increase over time (SI Appendix, Section VII Fig. 9). Frequency 482 483 seriation (SI Appendix, Section VII Fig. S8) suggests the progressive emergence of irregular asymmetric fractures become more abundant over time while irregular symmetric, notch, oblique, 484 and flute-mouth fractures decrease in both apex and base. Proximal portions and entire shells are the 485 least represented in the assemblage (SI Appendix, Section VII Fig. S8). It is interesting to note that, 486







Fig. 9 Diversity over time graphs representing Gini-Simpson's diversity index (a) and their 490

- equivalent numbers (b; following Jost 2006) across the sequence of examined layers at Grotta del 491 Cavallo
- 492

When layers are clustered based on measures of pairwise similarity (SI Appendix, Section VII Table 494 S54; Fig. S9) EII-I always emerges as the outlier. 495

496

#### 4.3 Gastropods and bivalves 497

- In the archaic Uluzzian layer EIII gastropods and bivalves are scarce (n=40) and affected by high 498
- fragmentation (87.5%) and decalcification (Table 3). 499

Layers/	Classes	NISP			Anthrop	pogenic proc	cesses	
Spit			ochre	tot_hole	broken	abr_crack	anthropic	anthropized
ЕШ	Gastr.	33	-	2	-	-	1	-
	Biv.	7	-	-	-	-	-	-

	Gastr.	1	-	-	-	-	-	-
E-D	Biv.	1	-	_	_	_	-	-
DII	Gastr.	10	1	6	-	-	6	-
	Biv.	2	-	-	-	-	-	-
DI	Gastr.	42	2	17	1	6	5	-
	Biv.	45	1	12	2	2	-	8
D whole	Gastr.	13	1	5	3	1	1	-
	Biv.	2	-	1	-	1	-	-
Total %	-	158	3.1%	27.84%	3.7%	6.9%	8.2%	5.0%
Total	-	158	5	44	6	11	13	8

501 Table 5 Anthropogenic traces on the bivalve and gastropod assemblage by layer

Among all uncovered specimens the most outstanding one consists of a complete pierced specimen of *Tritia neritea* (Table 5 and Fig. 10a). The cross-section exhibits traces of flacking on the outside surface (defined "interior wedging" by Tátá et al. 2014) that are consistent with perforation from the inside of the shell. A notch found on the edge of the shell (Fig. 10a) and the rounded rims of the pierced hole are compatible with repeater contact with a rope used to thread the bead.



507

Fig. 10 Technological traces on gastropods from Cavallo. Layer EIII— perforated *Tritia neritea*showing a notch consistent with suspension (a). Layer DII—close-up of use-wear on the edge of the
hole on a perforated *Phorcus turbinatus* (b). Layer DII—rounded edges on perforated specimens of *Tritia neritea* (c, d). Layer DI—perforation on a *Columbella rustica*, showing a notch and rounded
edge (e). Layer DI— subcircular hole on a *Homalopoma sanguineum* (f)

A decalcified fragment of *Callista chione*, a *Naticarius hebraeus* covered with concretions and two

unidentified fragments are the only shells found in layer EII-I. The three holes identified on the

specimen of *N. hebraeus* present with irregular contours and potential recent breakage of the edges,

all of which point to post-depositional processes (Fig. 11a).



518

Fig. 11 Layer EII-I—postdepositional holes on a *Naticarius hebraeus* (a). Layer DI—traces of
red pigment near the hole of a *Lembulus pella* (b). Layer DI—residues of red pigment located
inside the hole of a *Tritia neritea* (c). Layer DI—heat cracks on a *Tritia neritea* (d)

523 Only two identified specimens have been retrieved in the transitional split E-D: a fragment of 524 *Spondylus gaederopus* and a small shell of *Euspira catena*. These remains no longer show the 525 periostracum and their sculptural features due to decalcification.

Layers DII and I (Upper Uluzzian) stand out for the large number of gastropods and bivalves. In

527 particular *Tritia neritea* (n=31) and *Homalopoma sanguineum* (n=11) are the most abundant species

528 in the assemblage. The taphonomic study (Table 3) identified traces of beach weathering on 90% of

529 gastropods and traces of predation by other carnivores on ~9% of them. These results suggest that

the vast majority of gastropods were collected dead from beach shores, and they were not taken to
the cave to be consumed as food. The most represented species such as *T. neritea*. *H. sanguineum*, *C. rustica* and *Ph. turbinatus* present with regular sub-circular holes located near the shell aperture.
Notches were detected on hole rims, and edges were abraded probably due to use-wear (Fig.10 b-f).
The majority of notches are located in the IV (31%) and I (38%) quarters (Fig.12), and are
consistent with the use of *T. neritea* as suspended ornaments.



537

Fig. 12 Graphic representation of the location of notches on edges of pierced shell found at Grotta
del Cavallo during the Upper Uluzzian (DII-DI) Ochre was found inside three specimens (Fig. 11b
and c), and five gastropods show heat cracks on their external surface (Fig 11d).

541

Bivalves mostly consist of Mytilis galloprovincialis, Glycymeris sp., and Lembulus pella. The first 542 species is represented by fragments and hinges. Two broken hinges of *Glycymeris* sp. exhibit partial 543 holes on the umbos although fragmentation makes it impossible to discriminate between anthropic 544 and natural taphonomic processes. Specimens of L. pella consist instead of entire valves. 545 Taphonomic analysis based on all fragments and the small amount of complete specimens support 546 the conclusion that almost 70% of bivalve remains were abraded by sea waves on the shore. 547 Furthermore, all valves of L. pella show perfectly round and bevelled holes, both signs of drilling 548 by mollusc predators. Nevertheless, notches and irregular flaking are once again visible on the 549 edges of drilled holes (Fig. 13a-c), both of which are consistent with use-wear (Fig. 13b-c). These 550 traces, however, are not coherent with drilling perforation operated by other molluscs (Fig.13d). 551

The notches are located near the umbo (quarter IV, 32%) and on the edge above it (quarter III, 27%) (Fig.12). We therefore suggest that these naturally pierced shells were deliberately chosen to be used as beads by humans. In addition, some red dusty substance (possibly ochre) was found on the external surface of one of the valves (Fig.11d).



556

Fig. 13 Layer DI—*L. pella* with a round drilled hole showing a notch in quarter IV (a). Layer DI— *L. pella* with drilled holes showing notches in quarter IV and irregular flaking (b, c). Layer EIII—
hole due to predation on a *Turritella sp.* (d)

560

### 561 **5. Discussion**

562 The ornamental shell assemblage of Grotta del Cavallo is mostly characterised by a large amount of

- scaphopods which are present across the entire archaeological sequence and abound in the
- uppermost layers, where gastropods and bivalves are also recorded. Among gastropods *Tritia*
- *neritea* is the most frequently recorded species, while *Lembulus pella* dominants among bivalves.
- 566

#### 567 5.1 Bead making at Grotta del Cavallo

As far as tusk shells are concerned, the results of our multidisciplinary study point to an 568 anthropogenic action both in terms of their collection and manufacture. As attested by directly 569 dating results (Benazzi et al. 2011), and micropaleontological and Sr isotope analyses, the 570 archaeological specimens presented here are not fossil shells deriving from sedimentary outcrops 571 located in site surroundings, but rather consist of shells collected from beach shores at the same 572 time as the Uluzzian occupation of Grotta del Cavallo. The distance between Grotta del Cavallo and 573 the coastline was then about 12 km, enough to exclude any possible natural origin for the 574 occurrence of tusk shells inside the cave (Appendix, Section VIII and Fig S10). 575 These molluscs live in the sandy seabed below the intertidal zone, down to relatively great depths 576 577 (Appendix, Section IX). Therefore, they were most likely collected by the Uluzzian groups on the 578 beaches of shores that today are submerged. The collection of scaphopods by prehistoric people for feeding purposes is generally rejected, considering their small size and their very low nutritional 579

value (Bar-Yosef Mayer 2008).

Data resulting from our experimental and statistical tests support a human origin also for 581 manufacture, and experimental results reveal a possible association between scaphopod portioning 582 and type of fracture. Nevertheless, we agree with Vanhaeren and d'Errico (2001) and Taborin 583 (1993) that no specific fracture can be univocally associated with anthropogenic action. Although 584 585 intentionality cannot be definitely proven for any rectilinear fracture uncovered in archaeological assemblages, this kind of fracture is abundant in the dataset from Grotta del Cavallo, while it is 586 considerably rarer among experimental samples generated through trampling or in shells collected 587 588 on the present-day beaches, as already pointed out by previous research (Vanhaeren and d'Errico 2001). The only traces that surely correlate with human intervention are sawing cut-marks. In the 589 assemblage of Cavallo cut-marks were identified only on one specimen from layer D because of the 590 general bad state of preservation of all shells. In both layers E and D at Cavallo, however, a 591 particular notch-shaped fracture is documented, which, based on our experimental results and on 592 593 data from the literature (Vanhaeren and d'Errico 2001), can still be related to sawing. The other

types of fractures (in particular asymmetric and symmetric irregular fractures) are less diagnostic
because they can be also produced by trampling and other post-depositional processes. Notch and
flute-mouth fractures can also occur during suspension, due to the contact among beads (Álvarez
Fernández, 2006). In this light, the presence of red coloring substances on the shells exhibiting
flute-mouth and notch fractures becomes more informative. These characteristics are reminiscent of
the ochred shell from Riparo Broion, also displaying a notch fracture (*Appendix, Section X*, Fig.
S12).

Results also support change over time in the most represented distal and proximal fracture types. 601 Irregular asymmetric and rectilinear fractures increase their presence over time and become the 602 603 most represented ones. In earlier phases, notch, oblique, and irregular symmetric fractures are more 604 frequent. Diversity is generally high in the whole of the sequence (especially in distal fractures). Nevertheless, a tendency towards higher homogeneity of fracture types can be ascertained over time 605 (lowest diversity in DI), with a preference for rectilinear ones. This evidence may be also due to a 606 change in manufacturing techniques (i.e. percussion technique in earliest and bending in recent 607 phases respectively) or to different post-depositional processes. Dimensional analysis shows a 608 generalised increase in complete shell size and maximum diameter of all portions over time. This 609 could be due to a change in catchment preference guided by a selective pressure for bigger shells; or 610 611 it could relate to change in environmental conditions, despite data are still scarce to support this hypothesis. Relations between some environmental factors and morphological variations (e.g. body 612 size) are known for the most common benthic molluscs in the Mediterranean (e.g. Mamouridis et al. 613 2011; Peharda et al. 2017), but currently are not available for Antalis specimens. 614 Despite the increase in the size of entire shells over time, length of distal pieces does not change 615 and may reflect a choice aimed to obtain homogeneous beads. Although the level of regularity 616 exhibited by the Magdalenian parure of La Madeleine in France (Vanhaeren and d'Errico 2001) is 617 never reached. In the assemblage found at Cavallo there are also entire tusks that are suitable for 618

suspension, suggesting that this use might have involved beads of different size. The type of tusk

remnant portions tends to become more diverse over time. Only layer EII-I exhibits a different
distribution, displaying a predominance of medial portions consistently associated with a large
variety of fracture types (in both apex and base).

623 Furthermore, the presence of short apical portions in unit D often associated with rectilinear

624 fractures at the base of the apical portion, could suggest a manufacturing process *in situ*. They seem

to illustrate the systematic breakage of the apexes by bending to obtain beads easily and quickly.

626

As far as gastropods and bivalves are concerned, 13.2% of the total NISP exhibits traces of
anthropic manipulation for ornamental purposes. Nevertheless, shell collection for dietary purpose
can be confidently excluded for most gastropods and bivalves, because of their small size and
because they were collected dead on beaches, as suggested by taphonomic analysis (Stiner 1999;
Vanhaeren and d'Errico 2006; Colonese et al. 2011; Stiner et al. 2013; Stiner 2014; Bosch et al.
2015).

*Tritia neritea* is the most frequent species among gastropods (37.5% of which exhibits signs of
 perforation) followed by *Homalopoma sanguineum* (18.2% of which is perforated).

Pierced specimens show a high degree of s regularity in perforation techniques. Holes are consistently located near the shell aperture, anthropogenic-flaking is always found on the outer surface of the shell, and use-wear abrasion is consistently found on the lips near the aperture. These data are consistent with those observed in our experimental collection, even though further tests are required to better understand the exact method of perforation.

Bivalves seem to be less frequently chosen as adornment objects, evidence of which can be only

found in the latest Uluzzian phase (layer DI). This might be due to the great incidence of alimentary

642 species among bivalves, such as *Mytilus galloprovincialis*, *Glycymeris nummaria* and *Callista* 

643 *chione*. Anthropogenic traces can be detected on the 10 remains attributed to *Lembulus pella*, even

if no intentional perforation can be identified. All specimens present with circular holes with

bevelled edges that can be attributed to predator gastropods. At the same time, the best preserved
shells also exhibit notches located near the umbo that are consistent with deliberate suspension.
The opportunistic use of natural holes is commonly documented in archaeological contexts (BarYosef Mayer et al. 2009; Cabral and Martins 2016).

649

All the above mentioned results support human involvement in collection, transportation, and 650 modification (the latter for a part of the assemblage) of the shells uncovered at Grotta del Cavallo. If 651 pierced specimens and intentionally broken or used tusks can be confidently associated with an 652 ornamental role, unmodified shells can be interpreted as raw materials waiting to be used in bead 653 production. Since natural accumulation of shells can be excluded, the occurrence of unpierced 654 items which are not suitable for consumption, tusk shells which are not suitable for suspension, and 655 waste products suggests that a systematic manufacturing process was taking place at the site. 656 657 Considering the clustered distribution of shells illustrated above, the presence of an adornment manufacturing area could be proposed for the most recent phase (layer DI). This area was likely 658 located in square H11 and, to a lesser extent, in square H7, where shells have been retrieved. In the 659 oldest layers, shells were found in a different area of the cave. Here the retrieval of tusk distal 660 portions not suitable for suspension is also documented, and is also possible to hypothesize 661 breakage in situ, even if with lower levels of intensity and dimensional homogeneity. 662

663

#### **5.2 Bead making in the Uluzzian**

When looking at the other main Uluzzian sites found in Italy and Greece (e.g., Grotta della Cala,
Grotta di Castelcivita, Klissoura Cave, Riparo Broion), we observe a uniform scenario (except for
Castelcivita) exclusively characterized by shell ornaments, among which tusk shells are usually
prevalent (Fiocchi 1998; Stiner 2010; Peresani et al. 2019a). Such a composition of ornamental
assemblages testifies to the existence of a shared and widespread tradition across the Mediterranean.
Clearly recurrent features and marked similarity between sites imply either common ancestry or

cumulative mechanisms of cultural transmission and diffusion, processes that are commonly 671 accepted for the Protoaurignacian (Vanhaeren and d'Errico 2006; Stiner et al. 2013). At Riparo 672 Broion (Northern Italy), where the Uluzzian layers (SU 1f and 1g) have been recently dated to 44.4 673 - 42.8 ka cal BP, a few finished shell beads have been retrieved (Appendix, Section X, Fig. S12-13). 674 These consist of five scaphopods (4 Antalis vulgaris and 1 Antalis dentalis/inaequicostata) and a 675 pierced gastropod (Teodoxus fluvialis; Peresani et al. 2019a). All tusk beads show clear signs of 676 anthropogenic manipulation, documented by the occurrence of cut-marks on a specimen and well-677 developed suspension wear-traces on the whole of the sample (Peresani et al. 2019a). These beads 678 are notably constrained in their size range, when compared with those from Cavallo. Noteworthy is 679 680 the presence of notch fractures that can be related to suspension (Appendix, Section X, Fig. S13). In 681 addition, the gastropod shows traces of intentional perforation and suspension (Peresani et al. 2019a). The distance of Riparo Broion from possible fossil outcrops and/or the coast suggests that 682 the shells were brought to the site as finished products. 683

Moving to southern Italy, several mollusks (n=78) were found in the Uluzzian layers of Grotta della 684 Cala. These specimens belong to 25 different taxa (classes Gasteropoda, Bivalvia, Scaphopoda; 685 Fiocchi 1998). Among them there are six pierced gastropods (H. sanguineum and T. pellucida), two 686 perforated bivalves (Glycimeris nummaria) and 24 scaphopods. At a preliminary analysis, tusk 687 688 shells show rounded edges and seem to be all finished beads with probable suspension traces. Klissoura Cave I (Peloponnesus, Greece) yielded 11tusk shells and several gastropods, especially 689 *Tritia neritea* (n=7), and a few bivalves. The predominance of tusks and *Tritia neritea* draws an 690 691 interesting parallel with the uppermost layers of Grotta del Cavallo, especially considering the temporal overlap between the two contexts (Douka et al. 2014). 692

The cave site of Castelcivita, located at the foot of the Alburni massif (southern Italy), stands out from this framework as it yielded, so far, only two fragmentary shells (bivalves) (Gambassini 1997). 694 However, investigations at this site are still ongoing and the ornamental record might be enriched 695 696 by further items.

693

It is worth noting the marked resemblance between Uluzzian contexts and the Preaurignacian levels of Franchthi Cave (Peloponnesus, Greece) sealed by CI tephra (Farrand 2000; Fitzsimmons et al. 2013). Here a number of shells has been retrieved, but their poor state of preservation did not allow researchers to ascertain their use or the presence of perforation (Perlès 2019). The assemblage at Frachthi Cave mostly consists of inedible tusk shells and *Tritia neritea*, like in the case of Klissoura and Grotta del Cavallo DII-DI, both of which are used for ornament making in the following phases (Perlès 2019).

When ornaments of all Uluzzian contexts are considered at once a general trend over time can be 704 identified in assemblage composition. The earliest phases (Cavallo layers EIII and EII-I, Broion) 705 706 are characterized by scaphopods and rare pierced gastropods, whilst later phases (Cavallo layers DII 707 and DI, Klissoura, Cala and maybe also Franchthi) show an increase in richness of species used for ornament production but still with a higher percentage of tusk shells, followed by gastropods 708 709 (especially Tritia neritea). Among them H. sanguineum is marginally represented. In contrast, subsequent Upper Palaeolithic contexts display a marked preference for this species, which was 710 consistently included in personal ornamental assemblages and exchanged over long distance across 711 the Europe (Alvarez-Fernández 2006; Vanhaeren and d'Errico 2006; Peresani et al. 2019b). As far as 712 bivalves are concerned, they seem to be considered less relevant until the latest Uluzzian phases. 713 714 Ornament assemblage size in Uluzzian sites is also relevant, when compared with the coeval Châtelperronian, the only European techno-complex to have yielded conspicuous ornamental 715 assemblages, composed of bone and ivory pendants as well as very sporadic shells (Zilhão 2007). 716 717 Their size, however, is considerably smaller than that of the Uluzzian assemblages. The bulk of the Châtelperronian ornaments are from Grotte du Renne (n=35, d'Errico et al. 1998), where uncertain 718 integrity of the stratigraphic sequence made it difficult to accurately reconstruct the cultural 719 provenance of some items (White 2002; Higham et al. 2010; but see *contra* Caron et al. 2011; 720 721 Hublin et al. 2012).

The other Châtelperronian sites yielded only a few ornaments (Zilhão 2007), but many of these 722 723 deposits, such as Grotte des Fées, Roc de Combe and Trou Magrite, suffered from the intrusion of Aurignacian materials (Bordes and Labrot 1967; Rigaud 2001; Moreau 2003; Mellars et al. 2007; 724 Zilhão et al. 2007). The occurrence of several tusk shells at Saint-Césaire is certainly of great 725 interest (d'Errico et al.1998), but, regretfully, no further information is available on this material. 726 The other European transitional techno-complexes are characterized by sporadic findings 727 (Kozlowski 1982; Hülle 1977; Zilhão 2007) which do not seem to support a pivotal role of 728 nonperishable personal ornaments in the local material cultures. 729 Also worthy of note is the complete lack of tusks in the Initial Upper Palaeolithic (Emiran) 730 731 assemblages of the Levant, where gastropods and bivalves are the only species used for ornamental 732 purposes. To date, the most ancient tusk beads of this region are from layer E-E2 (ca 42. ka cal BP; Kuhn et al. 2009) of Üçağızlı Cave (Turkey; Stiner et al. 2002; Campbell 2017), containing an early 733 Ahmarian assemblage (Stiner et al. 2013). In this area other scaphopods have been recovered at 734 Manot Cave and Kebara, inside contexts with mixed Ahmarian and Aurignacian materials. The 735 presence of such shells is currently interpreted as the result of influence by groups bearing 736 European Aurignacian cultural elements (Bar Yosef-Mayer 2019). This evidence confirms the 737 differences existing between Uluzzian and IUP contexts and is consistent with the lack of Uluzzian-738

rage like techno-complexes in the Near East (Moroni et al. 2013).

In ethnographic instances ornaments connote and, in some cases, identify the human groups who 740 produce and wear them, thanks to the social, symbolic, cognitive and artistic values they embody. 741 742 In this perspective adornment objects (including but not limited to shell beads) are elements of visual communication. For these reasons their widespread presence in the Uluzzian suggests a high 743 degree of cohesion within and possibly between human groups. What the use of ornaments exactly 744 symbolized in the Uluzzian mental template is difficult to say, even if we can hypothesize that 745 these objects played a key role in defining and communicating the identity of the individual and/or 746 747 the group to which the individual belonged - thus suggesting, in some way, the occurrence of a

shared symbolic background. The systematic production and use of ornaments, in addition to other
characteristics, markedly differentiate the Uluzzian from the Mousterian, as the occurrence of
ornaments in the Middle Palaeolithic remains sporadic, despite some recent discoveries and
researchers' increased interest for this debated issue (Zilhão et al. 2009; Peresani et al. 2011;
Peresani et al. 2013; Romandini et al. 2014).

753

#### 754 **6.** Conclusion

The widespread occurrence of shell beads in Uluzzian contexts supports the emergence of a well-755 established technology of bead making in Mediterranean Europe early as 45 ka, well before the 756 onset of Aurignacian ornament productions across Europe. Uluzzian groups in Italy and Greece 757 758 shared the same ornament traditions, and Grotta del Cavallo is the archaeological context that best explains this phenomenon. The Uluzzian sequence at this site yielded the most conspicuous amount 759 of shell beads ever found in European transitional contexts. Analysing change over time in this 760 761 assemblage allow us to document the presence of finished products and unfinished by-products, 762 attesting to an increasing seek for dimensional regularity and serial production, as well as to the evidence for systematic ornament use. These features indicate that the Uluzzian bead making is 763 fully comparable with the ornament productions typical of the Upper Palaeolithic, and push back 764 the date for the beginning of systematic ornament use and manufacture in Europe to 45 ka 765

#### 766 Data availability

Authors can confirm that all data and source codes for analyses are currently included assupplementary information files.

### 769 Acknowledgements

Acknowlodgements: We thank the Soprintendenza Archeologia, Belle Arti e Paesaggio per le
Province di Brindisi, Lecce e Taranto for supporting our research in Apulia over the years. We

express gratitude to prof. Palma di Cesnola and prof. Paolo Gambassini for giving permission tostudy the Uluzzian materials from their excavations at Grotta del Cavallo.

We are grateful to Luigi Romani for his help in taxonomic classification of scaphopods, to Lavinia 774 Corbo who contributed to measuring specimens, and to Stefano Ricci who made photographic 775 documentation of the shells from Grotta del Cavallo. We also acknowledge all the students who 776 helped us in experimental tests (Francesco Antonini, Giulia Cappelletti, Margherita Gramigni, 777 Simone Marzocca and Irene Mazza). A special thanks to Sem Scaramucci for his contribution in 778 collecting tusks in Salento and for his precious advice. We are also grateful to the two anonymous 779 reviewers for their careful reading of the manuscript and constructive comments. This project has 780 781 received funding from the European Research Council (ERC) under the European Union's Horizon 782 2020 research and innovation program (grant agreement No 724046 – SUCCESS - http://www.erc-783 success.eu).

Author contributions: S.A., E.B., S.B and A.M. designed research; S.A. performed morphological 784 and use-wear analysis. E.B. carried out statistical analysis. L.T carried out taphonomic and 785 technological analysis. A.B., G.M., and L.T. made the taxonomic classification of the shells. A.M.B 786 and L.M.F. carried out micropaleontological studies and F.L. and A.C. realized isotope analysis. 787 Experimental tests were performed by S.A., D.A., F. Bo, C.F., G.M., G.O. and S.S. Paleoclimate 788 and paleoecological conditions were described by F.Ba. V.S. collected present day tusk shells and 789 reconstructed the palaeo-geographic conditions of the Uluzzian coastline. A.M., M.P., M.R. and 790 A.R. provided archaeological data and contributed to the revision of the manuscript. S.A., E.B., A. 791 M., A.B., V.S., L.M.F., A.M.B., M.R., F.L., F.Ba., L. T and S.B. wrote the paper with inputs from all 792 co-authors. 793

794 **Competing Interests:** The authors declare no competing interests.

795

#### 796 **References**

797	Abadia OM, Nowell A (2015) Palaeolithic Personal Ornaments: Historical Development and
798	Epistemological Challenges. J Archaeol Method Th 22: 952-979. https://doi.org/10.1007/s10816-
799	014-9213-z
800	
801	Álvarez Fernández E (2006) Los objetos de adorno-colgantes del Paleolítico superior y del
802	Mesolítico en la cornisa cantábrica y en el valle del Ebro: una visión europea. Unpublished PhD
803	thesis, Universidad de Salamanca
804	
805	Arrighi S, Borgia V (2009) Surface Modifications of Flint Tools and Their Functional Meaning.
806	Mater Manuf Process 24: 922-927. https://doi.org/10.1080/10426910902987150
807	
808	Bar-Yosef Mayer D (2008) Dentalium Shells Used by Hunter-Gatherers and Pastoralists in the
809	Levant. Archaeofauna 17:103-110
810	
811	Bar-Yosef Mayer D (2019) Upper Paleolithic Explorers: The Geographic Sources of Shell Beads
812	in Early Upper Paleolithic Assemblages in Israel. PaleoAnthropology 105–115.
813	doi:10.4207/PA.2019.ART126
814	
815	Bar-Yosef Mayer D E, Vandermeersch B, Bar-Yosef O (2009) Shells and ochre in Middle
816	Paleolithic Qafzeh Cave, Israel: indications for modern behavior. J Hum Evol 56 (3):307–314.
817	doi:10.1016/j.jhevol.2008.10.005
818	
819	Bar-Yosef O (2007) The archaeological framework of the Upper Paleolithic revolution. Diogenes
820	54(2): 3–18
821	
822	Bar-Yosef O, Bordes JG (2010) Who were the makers of the Châtelperronian culture? J Hum Evol

**59**(5): 586–93. https://doi.org/10.1016/j.jhevol.2010.06.009

825	Benazzi S, Douka K, Fornai C, Bauer CC, Kullmer O, Svoboda J, Pap I, Mallegni F, Bayle P,
826	Coquerelle M, Condemi S, Ronchitelli A, Harvati K, Weber GW (2011) Early dispersal of modern
827	humans in Europe and implications for Neanderthal behaviour. Nature 479: 525-528.
828	https://doi.org /10.1038/nature10617
829	
830	Benazzi S, Slon V, Talamo S, Negrino F, Peresani M, Bailey SE, Sawyer S, Panetta D, Vicino G,
831	Starnini E, Mannino MA, Salvadori PA, Meyer M, Pääbo S, Hublin JJ (2015) The Makers of the
832	Protoaurignacian and Implications for Neandertal Extinction. Science 348: 793-6
833	
834	Benghiat S, Komso D, Miracle PT (2009) An experimental analysis of perforated shells from the
835	site of Sebrn Abri, Istria, Croatia. In: McCartan S, Schulting R, Warren G, Woodman P (eds)
836	Mesolithic Horizons. Oxbow books, Oxford, pp 730-736
837	
838	Bonnardin S (2007) From traces to function of ornaments: some Neolithic examples. In: Longo L,
839	Skakun N (eds) "Prehistoric Technology" 40 Years Later. BAR Int. Ser. 1783, Archaeopress,
840	Oxford, pp 297-308
841	
842	Bordes F, Labrot J (1967) La stratigraphie du gisement de Roc de Combe (Lot) et ses implications.
843	Bull Soc Préhist Fr LXIV (1): 15–28
844	
845	Boscato P, Crezzini J (2012) Middle-Upper Palaeolithic transition in Southern Italy: Uluzzian
846	macromammals from Grotta del Cavallo (Apulia). Quat Int 252: 90–98.
847	https://doi.org/10.1016/j.quaint.2011.03.028
848	

849	Bosch MD, Buck L, Strauss A (2019) Location, location, location: Investigating Perforation
850	Locations in Tritia gibbosula Shells at Ksar 'Akil (Lebanon) Using Micro-CT Data.
851	PaleoAnthropology 52-63. https://doi.org/10.4207/PA.2019.ART123
852	
853	Cabral JP, Martins JMS (2016) Archaeological Glycymeris glycymeris shells perforated at the
854	umbo: Natural or man-made holes? J Archaeol Sci Rep 10: 474-482.
855	https://doi.org/:10.1016/j.jasrep.2016.11.008
856	
857	Campbell G (2017) The Reproduction of Small Prehistoric Tusk Shell Beads. In: Bar-Yosef Mayer
858	D, Bonsall C, Choyke AM (eds) Not Just for Show: The Archaeology of Beads, Beadwork and
859	Personal Ornaments, Oxbow Books, pp 168-224
860	
861	Caron F, d'Errico F, Del Moral P, Santos F., Zilhão J (2011) The reality of Neandertal symbolic
862	behavior at the Grotte du Renne, Arcy-sur-Cure, France. PLoS One 6(6): e21545.
863	https://doi.org/10.1371/journal.pone.0021545
864	
865	Charney N., Record S (2012) vegetarian: Jost Diversity Measures for Community Data. R package
866	version 1.2. https://CRAN.R-project.org/package=vegetarian
867	
868	Colonese A C, Mannino M A, Bar-Yosef Mayer D E, Fa DA. Finlayson JC, Lubell D, Stiner MC
869	(2011) Marine mollusc exploitation in Mediterranean prehistory: An overview. Quat Int 239 (1–2):
870	86–103. doi:10.1016/j.quaint.2010.09.001
871	
872	Craveri M (1931) Crine. In: Enciclopedia Italiana, Istituto Enciclopedia Italiana, Roma
873	

874	d'Errico F, Zilhão J, Julien M, Baffier D, Pelegrin J (1998) Neanderthal Acculturation in Western
875	Europe? A Critical Review of the Evidence and Its Interpretation. Curr Anthropol 39: 1-44
876	
877	d'Errico F, Henshilwood C, Lawson G, Vanhaeren M, Tillier AM, Soressi M, Bresson F, Maureille
878	B, Nowell A, Lakarra J, Backwell L, Julien M (2003a) Archaeological evidence for the emergence
879	of language, symbolism, and music: an alternative multidisciplinary perspective. J World Prehist
880	17: 1-70
881	
882	d'Errico F, Julien M, Liolios D, Vanhaeren M, Baffier D (2003b) Many awls in our argument.
883	Bone tool manufacture and use in the Châtelperronian and Aurignacian levels of the Grotte du
884	Renne at Arcy-sur-Cure. In: Zilhão J, d'Errico F (Eds) The Chronology of the Aurignacian and of
885	the Transitional Technocomplexes. Dating, Stratigraphies, Cultural Implications. Proceedings of
886	Symposium 6.1 of the XIV Congress of the UISPP, Trabalhos de Arqueologia, vol. 33. Instituto
887	Português de Arqueologia, Lisboa, pp 247-271
888	
889	d'Errico F, Vanhaeren M, Barton N, Bouzouggar A, Mienis H, Richter D, Hublin JJ, McPherron
890	SP, Lozouet P (2009) Additional evidence on the use of personal ornaments in the Middle
891	Palaeolithic of North Africa. Proc Natl Acad Sci 106: 16501-16056.
892	https://doi.org/10.1073/pnas.0903532106
893	
894	d'Errico F, Borgia V, Ronchitelli A (2012) Uluzzian bone technology and its implications for the
895	origin of behavioural modernity. Quat Int 259: 59-71 https://doi.org/10.1016/j.quaint.2011.03.039
896	
897	d'Errico F, Banks WE (2013) Identifying Mechanisms behind Middle Paleolithic and Middle
898	Stone Age Cultural Trajectories. Curr Anthropol 54 (8):371-387. https://doi.org/10.1086/673388
899	

900	d'Errico F, Backwell L (2016). Earliest evidence of personal ornaments associated with burial: the
901	Conus shells from Border Cave. J Hum Evol 93: 91–108
902	
903	Dimitrijević V (2014) The provenance and use of fossil scaphopod shells at the Late
904	Neolithic/Eneolithic site Vinča – Belo Brdo, Serbia. In: Szabó K, Dupont C, Dimitrijević V,
905	Gómez Gastélum L, Serrand N (eds) Archaeomalacology: Shells in the Archaeological Record
906	BAR International Series 2666, Archaeopress, Oxford, pp 33-41
907	
908	Douka K, Higham, TF, Wood R, Boscato P, Gambassini P, Karkanas P, Peresani M, Ronchitelli A
909	(2014) On the chronology of the Uluzzian. J Hum Evol 68: 1-13.
910	https://doi.org/10.1016/j.jhevol.2013.12.007
911	
912	Farrand WR (2000) Depositional History of Franchthi Cave, Sediments, Stratigraphy and
913	Chronology. With a Report on the Background of the Franchthi project by T.W. Jacobsen.
914	Excavations at Franchthi Cave, Greece, fasc. 12, Indiana University Press, Bloomington &
915	Indianapolis
916	
917	Fiocchi C (1998) Contributo alla conoscenza del comportamento simbolico di Homo sapiens
918	sapiens. Le conchiglie marine nei siti del Paleolitico superiore europeo: strategie di
919	approvvigionamento, reti di scambio, utilizzo. Unpublished PhD thesis Scienze Antropologiche,
920	Consorzio Universitario di Bologna, Ferrara, Parma
921	
922	Fitzsimmons KE., Hambach U, Veres D, Iovita R (2013) The Campanian Ignimbrite eruption: new
923	data on volcanic ash dispersal and its potential impact on human evolution. PLoS One 8(6):
924	e65839. https://doi.org/10.1371/ journal.pone.0065839
925	

926	Flas D (2011) The Middle to Upper Paleolithic transition in Northern Europe: the Lincombian-
927	Ranisian-Jerzmanowician and the issue of acculturation of the last Neanderthals. World Archaeol.
928	43: 605-627
929	
930	Gambassini P (1997) Le industrie paleolitiche di Castelcivita. In: Gambassini P (ed) Il Paleolitico
931	di Castelcivita culture e ambiente, Electa, Napoli, pp 75-145
932	
933	Giaccio B, Hajdas I, Isaia R, Deino A, Nomade S (2017) High-precision 14C and 40Ar/39Ar
934	dating of the Campanian Ignimbrite (Y-5) reconciles the time-scales of climatic-cultural processes
935	at 40 ka. Sci Rep 7: 45940
936	
937	Gioia P (1990) An aspect of the transition between Middle and Upper Paleolithic in Italy: the
938	Uluzzian. In: Farizy C (ed) Paléolithique moyen récent et Paléolithique supérieur ancien en
939	Europe Ruptures et transitions: examen critique des documents archéologiques, APRAIF,
940	Nemours, pp 241–250
941	
942	Gravina B, Bachellerie F, Caux S, Discamps E, Faivre JP, Galland A, Michel A, Teyssandier N,
943	Bordes JJ (2018) No Reliable Evidence for a Neanderthal-Châtelperronian Association at La
944	Roche-à-Pierrot, Saint-Césaire. Sci Rep 8: 15134. https://doi.org/10.1038/s41598-018-33084-9
945	
946	Higham T, Jacobi R, Julien M, David F, Basell L, Wood R, Davies W, Ramsey CB (2010)
947	Chronology of the Grotte du Renne (France) and implications for the context of ornaments and
948	human remains within the Châtelperronian. Proc Natl Acad Sci 107(47): 20234-9. https://doi.org/
949	10.1073/pnas.1007963107
950	

951	Hoffmann DL, Angelucci DE, Villaverde V, Zapata J, Zilhão J (2018) Symbolic use of marine
952	shells and mineral pigments by Iberian Neandertals 115,000 years ago. Sci Adv 4(2).
953	https://doi.org / 10.1126/sciadv.aar5255
954	
955	Hublin JJ (2015) The modern human colonization of western Eurasia: when and
956	where? Quat Sci Rev 118: 194-210. http://dx.doi.org/10.1016/j.quascirev.2014.08.011
957	
958	Hublin JJ, Spoor F, Braun M, Zonneveld F, Condemi S (1996) A late Neanderthal associated with
959	Upper Palaeolithic artefacts. Nature 381: 224–226. https://doi.org/10.1038/381224a0
960	
961	Hublin JJ, Talamo S, Julien M, David F, Connet N, Bodu P, Vandermeersch B, Richards MP
962	(2012) Radiocarbon dates from the Grotte du Renne and Saint-Césaire support a Neandertal origin
963	for the Châtelperronian, Proc Natl Acad Sci 109(46): 18743-18748;
964	https://doi.org/10.1073/pnas.1212924109
965	
966	Hülle W M (1977) Die Ilsenhöhle unter Burg Ranis/Thüringen. Eine paläolitische Jägerstation.
967	Gustav Fischer, Stuttgart
968	
969	Jost L (2006) Entropy and diversity. Oikos 113(2): 363-375. https://doi.org/10.1111/j.2006.0030-
970	1299.14714.x
971	
972	Jost L (2007) Partitioning diversity into independent alpha and beta components. Ecology 88(10):
973	2427-2439. https://doi.org/10.1890/06-1736.1
974	
975	Kozłowski J (ed) (1982) Excavation in the Bacho Kiro Cave (Bulgaria): Final Report. Polish
976	Scientific Publishers, Warsaw

977	Kuhn SL, Stiner MC, Güleç E, Özer I, Yılmaz H, Baykara I, Açıkkol A., Goldberg P, Martínez
978	Molina K, Ünay E, Suata-Alpaslan F (2009) The early Upper Paleolithic occupations at Üçağızlı
979	Cave (Hatay, Turkey). J Hum Evol 56: 87-113. https://doi.org/10.1016/j.jhevol.2008.07.014
980	
981	Largiolli T, Martinis B, Mozzi G, Nardin M, Rossi D, Ungaro S (1969) Gallipoli. Note illustrative
982	della Carta Geologica d'Italia. Poligrafica e Cartevalori, Ercolano (Napoli)
983	
984	Lévêque F, Vandermeersch B (1980) Les découvertes de restes humains dans un horizon
985	castelperronien de Saint-Césaire (Charente-Maritime). Bull. de la Soc. Préhist. de Française 77 :
986	187–189
987	
988	Leroi-Gourhan A (1964) Le geste et la parole (Vol 2). Albin Michel, Paris
989	
990	Lugli F, Cipriani A, Peretto C, Mazzucchelli M, Brunelli D (2017) In situ high spatial resolution
991	87Sr/86Sr ratio determination of two Middle Pleistocene (ca 580 ka) Stephanorhinus
992	hundsheimensis teeth by LA-MC-ICP-MS. Int J Mass Spectrom 412: 38-48.
993	https://doi.org/10.1016/j.ijms.2016.12.012
994	
995	Mamouridis V, Cartes JE, Parra S, Fanelli E, Salinas JS (2011) A temporal analysis on the
996	dynamics of deep-sea macrofauna: influence of environmental variability off Catalonia coasts
997	(western Mediterranean). Deep Sea Res Part 1 Oceanogr Res Pap 58 (4): 323-337
998	
999	McElreath R, Boyd R, Richerson PJ (2003) Shared norms and the evolution of ethnic markers.
1000	Curr Anthropol 44: 122-129. http://dx.doi.org/10.1086/345689
1001	

1002	Mellars P, Gravina B, Bronk-Ramsey C (2007) Confirmation of Neanderthal/Modern Human
1003	interstratification at the Châtelperronian type-site. Proc Natl Acad Sci 104: 3657–3662.
1004	https://doi.org/10.1073/pnas.0608053104
1005	
1006	Moreau L (2003) Les éléments de parure au Paléolithique supérieur en Belgique. L'Anthropologie
1007	107: 603–614
1008	
1009	Moroni A, Boscato P, Ronchitelli A (2013) What roots for the Uluzzian? Modern behaviour in
1010	Central-Southern Italy and hypotheses on AMH dispersal routes. Quat Int 316: 27-44.
1011	http://dx.doi.org/10.1016/j.quaint.2012.10.051
1012	
1013	Moroni A, Ronchitelli A, Arrighi S, Aureli D, Bailey SE, Boscato P, Boschin F, Capecchi G,
1014	Crezzini J, Douka K, Marciani G, Panetta D, Ranaldo F, Ricci S, Scaramucci S, Spagnolo V,
1015	Benazzi S, Gambassini P (2018) Grotta del Cavallo (Apulia – Southern Italy). The Uluzzian in the
1016	mirror. J Anthropol Sci 96: 1-36. http://dx.doi.org/ 10.4436/jass.96004
1017	
1018	Murtagh F, Legendre P (2014) Ward's hierarchical agglomerative clustering method: which
1019	algorithms implement Ward's criterion? J Classif 31. http://dx.doi.org /10.1007/s00357-014-
1020	9161-z
1021	
1022	Nenadic O, Greenacre M (2007) Correspondence Analysis in R, with two- and three-dimensional
1023	graphics: The ca package. J Stat Softw 20 (3): 1-13. http://dx.doi.org / 10.18637/jss.v020.i03
1024	
1025	Orchard WC (1929) Beads and beadwork of the American Indians. Salzwasser-verlag Gmbh,
1026	Paderborn
1027	

1028	Palma di Cesnola A (1965) Notizie preliminari sulla terza campagna di scavi nella Grotta del
1029	Cavallo (Lecce). Riv Sci Preistoriche 20: 291-302
1030	
1031	Palma di Cesnola A (1966) Gli scavi nella Grotta del Cavallo (Lecce) durante il 1966. Riv. Sci.
1032	Preistoriche 21: 289-301
1033	
1034	Palma di Cesnola A (1989) L'Uluzzien: Faciès italien du leptolithique archaïque. L'Anthropologie
1035	93 (4): 783–812
1036	
1037	Peharda M, Thébault J, Markulin K, Schöne BR, Janeković I, Chauvaud L (2017) Contrasting
1038	shell growth strategies in two Mediterranean bivalves revealed by oxygen-isotope ratio
1039	geochemistry: The case of Pecten jacobaeus and Glycymeris pilosa. Chem Geol.
1040	http://dx.doi.org/10.1016/j.chemgeo.2017.09.029
1041	
1042	Peresani M, Fiore I, Gala M, Romandini M, Tagliacozzo A (2011) Late Neandertals and the
1043	intentional removal of feathers as evidenced from bird bone taphonomy at Fumane Cave 44 ky
1044	B.P., Italy. Proc Natl Acad Sci 108 (10): 3888-3893. https://doi.org/10.1073/pnas.1016212108
1045	
1046	Peresani M, Vanhaeren M, Quaggiotto E, Queffelec A, d'Errico F (2013) An ochered fossil marine
1047	shell from the Mousterian of Fumane Cave, Italy. PLoS One 8 (7): e68572. http://dx.doi.org
1048	/10.1371/journal.pone.0068572
1049	
1050	Peresani M, Cristiani E, Romandini M, 2016 The Uluzzian technology of Grotta di Fumane and its
1051	implication for reconstructing cultural dynamics in the Middle-Upper Palaeolithic transition of
1052	Western Eurasia. J Hum Evol 91: 36-56. https://doi.org/10.1016/j.jhevol.2015.10.012
1053	

1054	
1055	Peresani M, Bertola S, Del Piano D, Benazzi S, Romandini M (2019a) The Uluzzian in the North
1056	of Italy. Insights around the new evidence at Riparo Broion Rockshelter. Archaeol Anthrop Sci.
1057	http:// doi.org/10.1007/s12520-018-0770-z
1058	
1059	Peresani M, Forte M, Quaggiotto E, Colonese AC, Romandini M, Cilli C, Giacobini G (2019b)
1060	Marine shell exploitation during the Aurignacian at Fumane Cave. PaleoAnthropology 64-81.
1061	doi:10.4207/PA.2019.ART124
1062	
1063	Perlès C (2019) Cultural Implications of Uniformity in Ornament Assemblages: Paleolithic and
1064	Mesolithic Ornaments From Franchthi Cave, Greece. PaleoAnthropology 196-207.
1065	doi:10.4207/PA.2019.ART131
1066	
1067	Radovčić D, Sršen AO, Radovčić J, Frayer DW (2015) Evidence for Neandertal Jewelry:
1068	Modified White-Tailed Eagle Claws at Krapina. PLoSOne 10 (3): e0119802.
1069	https://doi.org/10.1371/journal.pone.0119802
1070	
1071	R Core Team (2018) R: A language and environment for statistical computing. R Foundation for
1072	Statistical Computing, Vienna, Austria URL https://www.R-project.org/
1073	
1074	Romandini M, Peresani M, Laroulandie V, Metz L, Pastoors A, Vaquero M, Slimak L (2014)
1075	Convergent Evidence of Eagle Talons Used by Late Neanderthals in Europe: A Further Assessment
1076	on Symbolism. PlosOne 9 (7): e101278. https://doi.org/10.1371/journal.pone.0101278
1077	

1078	Ronchitelli A, Boscato P, Gambassini P (2009) Gli ultimi Neandertaliani in Italia: aspetti culturali.
1079	In: Facchini F, Belcastro G (eds) La Storia di Neandertal. Biologia e Comportamento. Jaka Book,
1080	Bologna, pp 257–288
1081	
1082	Rigaud JP (2001) A propos de la contemporanéité du Castelperronien et de l'Aurignacien dans le
1083	nord-est de l'Aquitaine: une révision des données et ses implications. In: Zilhâo J, Aubry T,
1084	Carvalho A (eds) Les premières hommes modernes de la Péninsule Ibérique. Actes du Colloque de
1085	la Commission VIII de l'UISPP, Instituto Português de Arqueologia, Lisboa, pp 61-68
1086	
1087	Ruppert J, Bernet JW (2001) Our Voices: Native Stories of Alaska and the Yukon. University of
1088	Toronto Press, Toronto
1089	
1090	Sano K, Arrighi S, Stani C, Aureli D, Boschin F, Fiore I, Spagnolo V, Ricci S, Crezzini J, Boscato
1091	P, Gala M, Tagliacozzo A, Birarda G, Vaccari L, Ronchitelli A, Moroni A, Benazzi S (2019) The
1092	earliest evidence for mechanically delivered projectile weapons in Europe, Nat Ecol Evol 3:140-
1093	1414.
1094	
1095	Shimek R, Steiner G (1997) Chapter 6. Microscopic anatomy of invertebrates. Volume 6B:
1096	Mollusca II, Wiley-Liss, Inc
1097	
1098	Škardla P (2013) The Bohunician in Moravia and Adjoining Regions. Archaeology, Ethnology
1099	and Anthropology of Eurasia 41 (3): 2-18- https://doi.org/10.1016/j.aeae.2014.03.002
1100	
1101	

1102	Stiner MC (1999) Palaeolithic mollusc exploitation at Riparo Mochi (Balzi Rossi, Italy): Food and
1103	ornaments from the Aurignacian through Epigravettian. Antiquity 73: 735–754.
1104	https://doi.org/10.1017/S0003598X00065492
1105	
1106	Stiner MC (2010) Shell ornaments from the Upper Paleolithic through Mesolithic layers of
1107	Klissoura Cave 1 by Prosymna (Peloponnese, Greece). Eurasian Prehistory 7 (2): 287–308
1108	
1109	Stiner MC (2014) Finding a Common Bandwidth: Causes of Convergence and Diversity in
1110	Paleolithic Beads. Biol Theory 9(1): 51-64. doi:10.1007/s13752-013-0157-4
1111	
1112	Stiner MC, Pehlevan C, Sağir M Özer I (2002) Zooarchaeological studies at Üçağızlı Cave:
1113	Preliminary results on Paleolithic subsistence and shell ornaments. Araşturma Sonuçları Toplantısı
1114	17: 29-36
1115	
1116	Stiner MC, Kuhn SL, Güleç E (2013) Early Upper Paleolithic shell beads at Üçaĝizli Cave I
1117	(Turkey): Technology and the socioeconomic context of ornament life-histories. J Hum Evol
1118	64(5): 380–398. doi:10.1016/j.jhevol.2013.01.008
1119	
1120	Taborin Y (1993) La parure en coquillage au Paléolithique. CNRS Éditions, Paris
1121	
1122	Tátá F, Cascalheira J, Marreiros J, Pereira T, Bicho N (2014) Shell bead production in the Upper
1123	Paleolithic of Vale Boi (SW Portugal): An experimental perspective. J Archaeol Sci 42(1): 29-41.
1124	doi:10.1016/j.jas.2013.10.029
1125	
1126	Vanhaeren M, d'Errico F (2001) La parure de l'enfant de La Madeleine (fouilles Peyrony). Un
1127	nouveau regard sur l'enfance au Paléolithique supérieur. Paléo 13 : 201-240

1129	
1130	Vanhaeren M, d'Errico F (2006) Aurignacian ethno-linguistic geography of Europe revealed by
1131	personal ornaments. J Archaeol Sci 33(8): 1105-1128. doi:10.1016/j.jas.2005.11.017
1132	
1133	Vanhaeren M d'Errico F, Stringer C, James SL, Todd JA, Mienis HK (2006) Middle Palaeolithic
1134	shell beads in Israel and Algeria. Science 312: 1785-1787. https://doi.org/10.1126/science.1128139
1135	
1136	Vanhaeren M, d'Errico F, van Niekerk KL, Henshilwood CS, Erasmus RM (2013) Thinking
1137	strings: additional evidence for personal ornament use in the Middle Stone Age at Blombos Cave,
1138	South Africa. J Hum Evol 64: 500-517
1139	
1140	Villa P, Pollarolo L, Conforti J, Marra F, Biagioni C, Degano I, Lucejko JJ, Tozzi C, Pennacchioni
1141	M, Zanchetta G, Nicosia C, Martini M, Sibilia E, Panzeri L (2018). From Neandertals to modern
1142	humans: New data on the Uluzzian. PLoS One 13: e0196786. doi:10.1371/journal.pone.0196786
1143	
1144	Welker F, Hajdinjak M, Talamo S, Jaouen K, Dannemann M, David F, Julien M, Meyer M, Kelso J,
1145	Barnes I, Brace S, Kamminga P, Fischer R, Kessler BM, Stewart JR, Pääbo S, Collins MJ, Hublin JJ
1146	(2016) Palaeoproteomic evidence identifies archaic hominins associated with the Châtelperronian
1147	at the Grotte du Renne. Proc Natl Acad Sci 113(40): 11162–11167.
1148	https://doi.org/10.1073/pnas.1605834113
1149	
1150	White R (1989) Visual thinking in the Ice Age. Sci Am 261(1): 92–99
1151	

1152	White R (2002) Observations technologiques sur les objets de parure. In Schmider B (ed)
1153	L'Aurignacien de la Grotte du Renne. Les fouilles d'André Leroi-Gourhan à Arcy-sur-Cure,
1154	(Yonne), XXXIVe Supplément à Gallia Préhistoire, CNRS, Paris, pp 257-266
1155	
1156	Zanchetta G, Giaccio B, Bini M, Sarti L (2018) Tephrostratigraphy of Grotta del Cavallo,
1157	Southern Italy: Insights on the chronology of Middle to Upper Palaeolithic transition in the
1158	Mediterranean. Quat Sci Rev 182: 65-77. https://doi.org/10.1016/j.quascirev.2017.12.014
1159 1160	Zilhão J, 2006 Neandertals and moderns mixed, and it matters. Evol Anthropol 15: 183-195.
1161	https://doi.org/10.1002/evan.20110
1162	
1163	Zilhão J (2007) The emergence of Ornaments and Art: An Archaeological Perspective on the
1164	Origins of "Behavioral Modernity". J Archaeol Res 15: 1-54
1165	
1166	Zilhão J, d'Errico F, Bordes JG, Lenoble A, Texier JP, Rigaud JP (2007) Grotte des Fées
1167	(Châtelperron, Allier) ou une interstratification "Châtelperronien-Aurignacien" illusoire. Histoire
1168	des fouilles, stratigraphie et datations. Paléo 19: 391-432
1169	
1170	Zilhão J, Banks WE, d'Errico F, Gioia P (2015) Analysis of Site Formation and Assemblage
1171	Integrity Does Not Support Attribution of the Uluzzian to Modern Humans at Grotta del Cavallo.
1172	PLoS One 10: e0131181
1173	List of captions
1174	Fig. 1 Distribution of ornaments dated between ca. 45-39 ka across transitional contexts in Europe
1175	(sites with debated cultural attribution are represented here, see Discussion chapter). Pie radius is

1176 proportional to the total size of ornamental assemblages, while colours indicate the relative

1177	proportion contributed by different classes of items. 1) Saint-Césaire, 2) Quinçay, 3) Roc de Combe,
1178	4) Caune de Belvis, 5) Cova Foradada, 6) Roche au Loup, 7) Grotte du Trilobite, 8) Grotte des Fées,
1179	9) Grotte du Renne, 10) Trou Magrite, 11) Ilsenhöle Ranis, 12) Willendorf II, 13) Riparo Broion,
1180	14) Grotta della Cala, 15) Grotta del Cavallo, 16) Klissoura Cave, 17) Franchthi Cave, 18) Bacho
1181	Kiro, 19) Brynzeny I
1182	
1183	Fig. 2 Selection of shells from Grotta del Cavallo. a) Antalis sp. b) Tritia neritea. c) Antalis sp. d)
1184	Antalis sp. e) Tritia neritea. f) Homalopoma sanguineum. g) Columbella rustica. h) Phorcus
1185	turbinatus. i) Lembulus pella. j) Glycymeris sp.
1186	
1187	Fig. 3 Type of fractures. Classification according to the type of fracture on the experimental
1188	material: rectilinear (a); oblique (b); symmetric irregular (c); asymmetric irregular (d); flute-mouth
1189	(e); notch (f). The bar is 1 mm
1190	
1191	<b>Fig 4</b> Fracture degrees. Flute-mouth fracture with degree 1(a); flute-mouth fracture with degree 2
1192	(b); flute-mouth fracture with degree 3 (c). The bar is 2 mm
1193	
1194	Fig 5 Waste products. Small apical portions interpreted as waste products
1195	
1196	Fig.6 Traces of anthropogenic manipulation. Grotta del Cavallo. Layer DII – Tusk shell with cut
1197	marks (a) this kind of marks are very similar to those obtained experimentally by cutting tusk shells
1198	with a flint tool (b); Layer EIII - Tusk shell with a notch fracture showing well rounded (c) and
1199	polished edges (d); Layer EIII- Tusk shell showing a notch fracture with flattened (e) and polished

1200	areas (f); Layer EIII. Polishes inside a notch fracture (g); polishes inside a notch fracture produced
1201	during suspension experiment with a leather string (h)
1202	
1203	Fig.7 Tusk shells inserted one into another from Grotta del Cavallo. Tusk shells inserted one into
1204	another from layer EII-I (a), split E-D (b) and layer DII (c). While the external surface of the tusks
1205	looks weathered (d), the inner, protected trait shows its original aspect (e)
1206	
1207	Fig.8 Ochred shells from Grotta del Cavallo. Layer EIII- Ochre associated to a notch fracture (a).
1208	Ochre and black residues on the external surface of a specimen (b). Ochre on the apical end of a
1209	tusk. The edge is rounded and polished (c). Layer EII-I- Ochre located inside the shell and
1210	longitudinally spread (d)
1211	
1212	Fig. 9 Diversity over time Graphs representing Gini-Simpson's Diversity index (a) and their
1213	Numbers equivalent (b; following Jost 2006) across the sequence of examined layers at Grotta del
1214	Cavallo
1215	
1216	Fig. 10 Technological traces on gastropods from Cavallo. Layer EIII – Perforated Tritia neritea
1217	showing a notch consistent with suspension (a). Layer DII - Close-up of use-wear on the edge of
1218	the hole on a perforated Phorcus turbinatus (b). Layer DII - Rounded edges on perforated
1219	specimens of Tritia neritea (c-d). Layer DI - Perforation on a Columbella rustica, showing a notch
1220	and rounded edge (e). Layer DI - Sub-circular hole on a Homalopoma sanguineum (f)

1222	Fig 11 Layer EII-I - Post-depositional holes on a <i>Naticarius hebraeus</i> (a). Layer DI -Traces of red
1223	pigment near the hole of a Lembulus pella (b). Layer DI - Residues of red pigment located inside
1224	the hole of a Tritia neritea (c). Layer DI - Heat cracks on a Tritia neritea (d)
1225	
1226	Fig.12 Graphic representation of notches location on edges of pierced shell found at Grotta del
1227	Cavallo during the Upper Uluzzian (DII-DI)
1228	

- 1229 Fig 13 Layer DI *L. pella* with a round drilled hole showing a notch in quarter IV (a). Layer DI *L*.
- 1230 *pella* with drilled holes showing notches in quarter IV and irregular flaking (b-c). Layer EIII Hole
- 1231 due to predation on a *Turritella* sp. (d)