

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Identification and geometric characterization of Lie triple screw systems and their exponential images

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Identification and geometric characterization of Lie triple screw systems and their exponential images / Wu, Yuanqing; Carricato, Marco. - In: MECHANISM AND MACHINE THEORY. - ISSN 0094-114X. - STAMPA. - 107:(2017), pp. 305-323. [10.1016/j.mechmachtheory.2016.09.020]

Availability:

This version is available at: <https://hdl.handle.net/11585/573023> since: 2020-04-22

Published:

DOI: <http://doi.org/10.1016/j.mechmachtheory.2016.09.020>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Wu, Y., and Carricato, M. 2017. Identification and geometric characterization of Lie triple screw systems and their exponential images. *Mechanism and Machine Theory*, 107, pp. 305-323.

ISSN 0094-114X

The final published version is available online at:

<http://dx.doi.org/10.1016/j.mechmachtheory.2016.09.020>

© 2016. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Highlights of
*Identification and Geometric Characterization of Lie
Triple Screw Systems and Their Exponential Images*

Yuanqing Wu, Marco Carricato

- a geometric characterization of the Lie product and the Lie triple product of a generic screw system.
- a systematic identification of all Lie triple screw systems of $\mathfrak{se}(3)$, by an approach based on both algebraic Lie group theory and descriptive screw theory.
- derivation of the exponential motion manifolds of the Lie triple screw systems in dual quaternion representation.

Identification and Geometric Characterization of Lie Triple Screw Systems and Their Exponential Images[☆]

Yuanqing, Wu^{a,*}, Marco, Carricato^a

^a*DIN—Dept. of Industrial Engineering, University of Bologna, Viale del Risorgimento 2, Bologna, Italy*

Abstract

The twist space of a plunging constant-velocity (CV) coupling with intersecting shafts consists, in all configurations, of a planar field of zero-pitch screws. Recently, we reported an important discovery about this screw system: it is closed under two consecutive Lie bracket operations, thus being referred to as a Lie triple system; taking the exponential of all its twists generates the motion manifold of the coupling. In this paper, we first give a geometric characterization of the Lie product and the Lie triple product of a generic screw system. Then, we present a systematic identification of *all* Lie triple screw systems of $\mathfrak{se}(3)$, by an approach based on both algebraic Lie group theory and descriptive screw theory. We also derive the exponential motion manifolds of the Lie triple screw systems in dual quaternion representation. Finally, several important applications of Lie triple screw systems in mechanism and machine theory are highlighted in the conclusions.




Keywords: screw system, Lie triple system, constant-velocity coupling, parallel mechanism, type synthesis

[☆]A preliminary version of this paper was presented at the 14th IFTOMM World Congress, Taiwan, Oct. 25 – 30, 2015 [1].

*Corresponding author

Email addresses: yuanqing.wu@unibo.it, troy.woo@gmail.com (Yuanqing, Wu), marco.carricato@unibo.it (Marco, Carricato)

Nomenclature

\mathcal{R}	revolute joint
\mathcal{S}	spherical-joint-equivalent subchain
\mathcal{E}	planar-joint-equivalent subchain
	zero-pitch screw
	infinite-pitch screw
	finite-non-zero-pitch screw
$\text{SE}(3)$	the special Euclidean group
$\mathfrak{se}(3)$	the Lie algebra of $\text{SE}(3)$
ξ, ξ_1, ξ_2, \dots	twists or screws
S	generic screw system with unspecified type
$S_{i,j}$	the j th special i -system (see [2, Ch. 12])
$S_{i,j}^0$	j th special i -system whose finite-pitch screws all have zero pitch
$L, L_{i,j}, L_{i,j}^0$	Lie product of $S, S_{i,j}$ and $S_{i,j}^0$ respectively: $L = [S, S]$
$T, T_{i,j}, T_{i,j}^0$	triple product of $S, S_{i,j}$ and $S_{i,j}^0$ respectively: $T = [[S, S], S]$

1. Introduction

Methods for inferring finite motions from infinitesimal ones are of crucial importance in kinematics of mechanisms and robotics [2, 3]. For example, a one degree-of-freedom (1-DoF) finite motion is a parameterized curve in the displacement group $\text{SE}(3)$, which may be integrated from its axode surface, a parameterized curve in the Lie algebra $\mathfrak{se}(3)$ of $\text{SE}(3)$ [3, 4]. From mechanism synthesis point of view, it is important to consider multi-DoF finite motions or *parameterized submanifolds* of $\text{SE}(3)$ that may be generated by the end-effector of an open-chain or closed-chain mechanism [5, 6, 7, 8, 9]. Several researchers characterize the end-effector motion set of mechanisms by the more general notion of algebraic varieties [10, 11]. In practice, we may always assume that such varieties are submanifolds by restricting them to an open subset about the identity element of $\text{SE}(3)$.

Inferring finite end-effector motions of mechanisms from their *twist space*¹ has been extensively studied in the literature, in particular in type synthesis of parallel mechanisms [13, 5, 6, 7, 14, 8, 15, 9, 16, 17, 12, 1]. The methods of inference involved in such works are mainly heuristic and developed on a case-by-case basis. Hunt pointed out that the full-cycle mobility of a kinematic chain may be inferred from its twist space if the latter remains invariant for all configurations [2]. The inferred finite motions are simply the connected subgroups of $\text{SE}(3)$, with their twist spaces being the corresponding Lie subalgebras of $\mathfrak{se}(3)$ [18, 5, 19]. For this reason, Lie subalgebras of $\mathfrak{se}(3)$ are sometimes referred to as *invariant screw system* (ISS) [12]. More generally, when the twist space of a mechanism remains invariant *up to a rigid displacement* under arbitrary

¹The twist space of a mechanism is the vector space of all possible end-effector twists at a given configuration. In other words, it is a vector subspace of the Lie algebra $\mathfrak{se}(3)$ of the special Euclidean group $\text{SE}(3)$.




Figure 1: Examples of PSS serial generators [12]: (a) a \mathcal{ES} -type serial chain; (b) \mathcal{EE} -type serial chain.




Figure 2: Screw systems of a CV coupling with intersecting shafts, where \mathbf{w}_1 and \mathbf{w}_2 are the input and the output twist, respectively: (a) non-plunging coupling; (b) plunging coupling. $S_{2,1}^0$ comprises a planar pencil of instantaneous rotations converging into a point. $S_{3,4}^0$ comprises a planar field of instantaneous rotations as well as an instantaneous translation perpendicular to the plane.

finite motions away from singularities, thus having a constant *class* [20], the mechanism is said to have a *persistent screw system* (PSS) of the end-effector (see Fig. 1 for examples) [12]. The notion of PSS is important as it allows us to expand the theoretical framework under which the inference of finite motion properties from instantaneous ones is guaranteed. The exhaustive derivation and classification of all *serial* chains with an m -dimensional (m -D) PSS of the end-effector is complete up to $m \leq 4$ [21, 22, 23].

However, not all persistent motions are generated by serial chains. In a seminal paper on the analysis and synthesis of CV couplings with parallel-kinematics structure [24], Hunt identified the twist space of a CV coupling with intersecting shafts to be either the fourth special three-system (with zero-pitch screws), denoted $S_{3,4}^0$, or the first special two-system (with zero-pitch screws), denoted $S_{2,1}^0$, depending on whether plunging is allowed or not (see Fig. 2 and also [2,




Figure 3: Typical connecting chains of plunging CV coupling with intersecting shafts: (a) a \mathcal{RSR} chain; (b) a \mathcal{RER} chain. Each pair of revolute joint twists (ξ_i^+, ξ_i^-) remains mirror symmetric about the bisecting plane.

Ch. 13.5)). Focusing, without loss of generality, on the plunging CV coupling, CV transmission is provided for any input and articulation angles if the twist space always remains a $S_{3,4}^0$ system, with all its screws lying on the bisecting plane [24]. The persistence property of the coupling is enforced by a parallel-kinematics architecture with prescribed types of connecting chains with five (in some cases four) DoF. All of the latter respect mirror symmetry about the bisecting plane (see Fig. 3 and also [24, 15]). The twist space of a CV coupling with intersecting shafts, although not fixed at a particular location, remains congruent to itself under arbitrary configuration changes, thus being persistent. It is briefly stated in [24] and later proved in [21] that there exists no serial chain whose twist space remains $S_{3,4}^0$ for all its configurations. The serial 5-DoF connecting chains in a plunging CV coupling are not persistent either. For example, the twist space of the $5\mathcal{R}$ chain shown in Fig. 3(b) has a reciprocal screw with zero pitch at a mirror symmetric configuration [24]; however, the chain may also have configurations where the reciprocal screw has infinite pitch [25]. This shows that the persistent behavior of $S_{3,4}^0$ in a CV coupling is not the result of the in-parallel connection of higher-dimensional-PSS serial generators.

Bonev et al. [26] studied CV-like generators of $S_{3,4}^0$ from a finite-motion perspective, by the so-called tilt-and-torsion orientation parameters. They pointed out that the end-effector motion of a 3- \mathcal{RSR} reflected tripod, which is a parallel mechanism equivalent to a plunging CV coupling (see [2, 15, pp. 397]), has zero torsion for all its configurations. In other words, the reflected tripod has a zero-torsion motion type² [27]. It is further shown in [27] that the motion

²The motion type [16] or motion pattern [9] of a mechanism is characterized by a submanifold Q of $SE(3)$, with the twist spaces of the mechanism being tangent spaces of Q at corresponding configurations. Once the motion type is given, the twist spaces can be derived

type of a plunging CV coupling is exactly the image $\exp(S_{3,4}^0)$ of $S_{3,4}^0$ under the exponential map \exp ³. The fact that its tangent spaces are all copies of $S_{3,4}^0$ can be readily verified by the half-angle property [28, 29].

Through a series of recent studies, we discovered that the many geometric properties of $\exp(S_{3,4}^0)$ are best understood from a symmetric space viewpoint [28, 29]. A symmetric space is a smooth manifold which has an inversion symmetry about every point [30]. The inversion symmetry condition for a screw system S is [28, 29]:

$$\forall \xi_1, \xi_2 \in S : \quad e^{\xi_1} e^{\xi_2} e^{\xi_1} \in \exp(S) \quad (1)$$

Both $S_{2,1}^0$ and $S_{3,4}^0$ satisfy condition (1), turning $\exp(S_{2,1}^0)$ and $\exp(S_{3,4}^0)$ into symmetric spaces. We showed in [29] that the inversion symmetry of S leads to the mirror symmetric arrangement of joint axes in a CV coupling. The necessary and sufficient condition for a screw system S to satisfy (1) is being closed under two consecutive *Lie bracket* or *commutator* [19] operations, namely [30]

$$\forall \xi_1, \xi_2, \xi_3 \in S : \quad [[\xi_1, \xi_2], \xi_3] \in S \quad (2)$$

The double commutator operation $[[\cdot, \cdot], \cdot]$ is sometimes referred to as a *Lie triple product* [31]. A screw system S that satisfies (2) is called a *Lie triple system* (LTS, see [30, pp. 78]). LTSs can be considered a generalization of Lie subalgebras (or ISSs) of $\mathfrak{se}(3)$, since *taking exponentials of these screw systems generates the corresponding motion manifolds*. However, as shown by Hunt's general theory of CV couplings, LTSs admit distinct and more complex type synthesis rules compared to ISSs. Hence, it is important to identify all LTSs of $\mathfrak{se}(3)$ and investigate their type synthesis.

A full list of the LTSs of $\mathfrak{se}(3)$, which all share the many geometric properties of $S_{2,1}^0$ and $S_{3,4}^0$, is given in [28, 29]⁴. However, details of their identification is not reported therein. In this paper, accordingly, we provide the full derivation and classification. Our main contribution here is to offer a more complete view of the geometry of LTSs from both a Lie algebraic and screw system point of view. So far, as the authors are aware of, this is the first time that two derived screw systems of a screw system S , namely the *Lie product* $[S, S]$

$$[S, S] := \{ [\xi_1, \xi_2] \mid \forall \xi_1, \xi_2 \in S \}$$

by differentiation.

³In this paper, we use \exp to denote the exponential map from $\mathfrak{se}(3)$ to $SE(3)$ [19]. The exponential of a twist ξ is simply denoted by e^ξ .

⁴It is worth pointing out that, after our early work in [28] and the presentation of the preliminary version of this paper in [1], Selig proposed a slightly different classification of LTSs for addressing optimal motion planning problems [32].

and the *Lie triple product*⁵ $[[S, S], S]$

$$[[S, S], S] := \{ [[\xi_1, \xi_2], \xi_3] \mid \forall \xi_1, \xi_2, \xi_3 \in S \}$$

are *systematically* studied from a screw theoretic point of view. In addition,
75 by using a dual-vector representation for the screws in S , we may study the
exponential manifolds of the LTSs as sub-varieties of the Study quadric, thus
offering more insight into the geometry of LTSs.

Without loss of generality, we base our identification process on the classi-
fication of screw systems provided in [2]. The classification of screw systems
80 provided by Hunt in [2] is exhaustive, even though Gibson and Hunt [20, 18]
later refined such classification by distinguishing several sub cases. While on
the one hand these refinements increase the description complexity, on the other
they are not relevant for the purpose of our work. Accordingly, Hunt's original
(and simpler) classification is used here.

85 The paper outline is as follows. In Sec. 2, we review concepts of Lie group
theory and screw geometry with the help of dual-vector representation. In
Sec. 3, we give a geometric characterization of Lie products and Lie triple
products, and introduce the concept of LTSs. In Sec. 4, we provide a systematic
identification of the LTSs of $\mathfrak{se}(3)$, by analyzing Hunt's complete list of screw
90 systems. Finally, in Sec. 5, we derive the dual quaternion representation of the
motion manifolds corresponding to the exponentials of the LTSs.

2. Definitions and notations

Denote the scalar and vector product of two vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^3$ by $\mathbf{u} \cdot \mathbf{v}$ and
 $\mathbf{u} \times \mathbf{v}$, respectively. The dual-vector representation of the instantaneous *twist*
of a rigid body motion is given by [34, Ch. 12]:

$$\begin{aligned} \xi &= \mu(\mathbf{w} + \varepsilon \mathbf{v}) \\ &= \mu(\mathbf{w} + \varepsilon(\mathbf{q} \times \mathbf{w} + h\mathbf{w})) \\ &= \mu(1 + \varepsilon h)(\mathbf{w} + \varepsilon \mathbf{q} \times \mathbf{w}) \end{aligned} \quad \left\{ \begin{array}{l} \mu, h \in \mathbb{R} \\ \mathbf{w}, \mathbf{v} \in \mathbb{R}^3 \end{array} \right. \quad \|\mathbf{w}\| = 1$$

where the dual number ε satisfies $\varepsilon^2 = 0$. ξ corresponds to the instantaneous
twist of a screw motion along the line with Plücker coordinate $\xi^\circ \triangleq \mathbf{w} + \varepsilon \mathbf{q} \times \mathbf{w}$,
95 where \mathbf{q} is any point on the line, usually chosen to be $\mathbf{w} \times \mathbf{v}$. The *pitch*, *mag-*
nitude and *dual magnitude* are given by $h = \mathbf{w} \cdot \mathbf{v}$, μ and $\mu(1 + \varepsilon h)$ respectively
[34]. We shall refer to ξ° as the *axis* of ξ . For the special case $\|\mathbf{w}\| = 0$, ξ
defines an infinite-pitch screw along direction \mathbf{v} , namely $\xi = \varepsilon \mathbf{v}$ with $\|\mathbf{v}\| = 1$.
The collection of all twists forms the 6-D vector space $\mathfrak{se}(3)$, i.e. the Lie algebra
100 of $\text{SE}(3)$ [19]. A *screw system* is a vector subspace of $\mathfrak{se}(3)$.

⁵The Lie product and Lie triple product have the same definition as the second and the
third term in the lower central series of a Lie algebra [33]. However, we find the former
locutions more easily intelligible for a non-specialized audience, since they may be related to
the familiar concepts of vector product $\mathbf{u} \times \mathbf{v}$ and vector triple product $(\mathbf{u} \times \mathbf{v}) \times \mathbf{w}$ of 3-D
vectors $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^3$.




Figure 4: Cylindroid of the general two-system $S_{2,g}$ with basis pitches α and β , $\alpha > \beta$. The length of the cylindroid is given by $\alpha - \beta$, with the plane \mathbf{ij} of the basis screws located in the middle. The variation of the pitch along the screws of the cylindroid is illustrated by the solid curve.

In particular, a two-system is spanned by two linearly independent screws, and it may either have a general form $S_{2,g}$ or one of five special forms $S_{2,j}$, $j = 1, \dots, 5$ [2, pp. 344]. The screw axes of a general two-system $S_{2,g}$ envelope a cubic surface known as the cylindroid (see Fig. 4). $S_{2,g}$ admits two concurrent and perpendicular screws with maximal pitch α and minimal pitch β , which we may use as *basis screws* of the system. For the purpose of our exposition, the basis screws of a screw subspace are chosen, without loss of generality, by reason of computational convenience.

Similarly, a three-system is spanned by three linearly independent screws, and it may have either a general form $S_{3,g}$ or one of ten special forms $S_{3,j}$, $j = 1, \dots, 10$ [2, pp. 356]. The screw axes of a general three-system $S_{3,g}$ envelope a pencil of ∞^1 concentric hyperboloids as shown in Fig. 5. $S_{3,g}$ admits three concurrent and perpendicular screws, which we use as basis screws (one has maximal pitch α , another has minimal pitch γ , and the last has an intermediate pitch β).

The commutator $[\xi_1, \xi_2] := \frac{1}{2}(\xi_1 \xi_2 - \xi_2 \xi_1)$ ([19, pp. 212]) of two dual vectors $\xi_i = \mathbf{w}_i + \varepsilon \mathbf{v}_i$, $i = 1, 2$ is simply equal to the vector product [19, 5]:

$$\begin{aligned} [\xi_1, \xi_2] &= \xi_1 \times \xi_2 = -\xi_2 \times \xi_1 \\ &= \mu_1 \mu_2 (\mathbf{w}_1 \times \mathbf{w}_2 + \varepsilon (\mathbf{w}_1 \times \mathbf{v}_2 - \mathbf{w}_2 \times \mathbf{v}_1)) \\ &= \mu_1 \mu_2 (1 + \varepsilon h_1)(1 + \varepsilon h_2) \xi_1^\circ \times \xi_2^\circ \end{aligned}$$

It is clear that the commutator is a bilinear operator satisfying, respectively, skew-symmetry and the Jacobi identity [19, 5], i.e.

$$\xi_1 \times \xi_2 = -\xi_2 \times \xi_1$$

and

$$(\xi_1 \times \xi_2) \times \xi_3 + (\xi_2 \times \xi_3) \times \xi_1 + (\xi_3 \times \xi_1) \times \xi_2 = 0 \quad (3)$$

In particular, $[\xi, \xi] = 0, \forall \xi \in \mathfrak{se}(3)$. The scalar product of two dual vectors ξ_1, ξ_2 gives

$$\xi_1 \cdot \xi_2 = \mu_1 \mu_2 (\mathbf{w}_1 \cdot \mathbf{w}_2 + \varepsilon (\mathbf{w}_1 \cdot \mathbf{v}_2 + \mathbf{w}_2 \cdot \mathbf{v}_1))$$

Given a unit twist $\xi = (1 + \varepsilon h)\xi^\circ$, the screw motion about ξ with angle θ , in dual quaternion representation (see [34]-12.4.4), is given by the exponential




Figure 5: Left: a general three-system $S_{3,g}$ with basis pitches α, β and γ , $\alpha > \beta > \gamma$; equal-pitch screws with $h \neq \beta$ form a family of ∞^1 reguli lying on concentric hyperboloids; right: the screws of pitch β lie on a degenerate regulus consisting of two line pencils.

map

$$e^{\theta \xi/2} = \cos(\hat{\theta}/2) + \sin(\hat{\theta}/2) \xi^\circ \quad (4)$$

where the dual angle $\hat{\theta}/2$ is defined as $\theta(1 + \varepsilon h)/2$ and

$$\begin{cases} \cos(\hat{\theta}/2) = \cos(\theta/2) - \varepsilon \frac{h\theta}{2} \sin(\theta/2) \\ \sin(\hat{\theta}/2) = \sin(\theta/2) + \varepsilon \frac{h\theta}{2} \cos(\theta/2) \end{cases}$$

or, for a unit infinite-pitch screw $\xi = \varepsilon \xi^\circ$,

$$e^{\theta \xi/2} = 1 + \varepsilon \frac{\theta}{2} \xi^\circ \quad (5)$$

Here, we adopt a half dual angle in order to be consistent with the usual matrix exponential $e^{\theta \xi}$ [34, Ch. 15.2.2]. For simplicity, we shall hereafter denote $\sin(\theta)$ and $\cos(\theta)$ by s_θ and c_θ , respectively.

Given the canonical basis $\mathbf{i}, \mathbf{j}, \mathbf{k}$ of \mathbb{R}^3 and a reference point $\mathbf{o} \in \mathbb{R}^3$, we define the following basis for $\mathfrak{se}(3)$:

$$\begin{cases} \mathbf{I} = \mathbf{i} + \varepsilon \mathbf{o} \times \mathbf{i} \\ \mathbf{J} = \mathbf{j} + \varepsilon \mathbf{o} \times \mathbf{j} \\ \mathbf{K} = \mathbf{k} + \varepsilon \mathbf{o} \times \mathbf{k} \end{cases} \quad \begin{cases} \varepsilon \mathbf{I} = \varepsilon \mathbf{i} \\ \varepsilon \mathbf{J} = \varepsilon \mathbf{j} \\ \varepsilon \mathbf{K} = \varepsilon \mathbf{k} \end{cases} \quad (6)$$

with commutator relations given by:

$$\begin{cases} \mathbf{I} \times \mathbf{J} = \mathbf{K} \\ \mathbf{J} \times \mathbf{K} = \mathbf{I} \\ \mathbf{K} \times \mathbf{I} = \mathbf{J} \end{cases} \quad \begin{cases} \varepsilon \mathbf{I} \times \mathbf{J} = \mathbf{I} \times \varepsilon \mathbf{J} = \varepsilon \mathbf{K} \\ \varepsilon \mathbf{J} \times \mathbf{K} = \mathbf{J} \times \varepsilon \mathbf{K} = \varepsilon \mathbf{I} \\ \varepsilon \mathbf{K} \times \mathbf{I} = \mathbf{K} \times \varepsilon \mathbf{I} = \varepsilon \mathbf{J} \end{cases} \quad \begin{cases} \varepsilon \mathbf{I} \times \varepsilon \mathbf{J} = \mathbf{0} \\ \varepsilon \mathbf{J} \times \varepsilon \mathbf{K} = \mathbf{0} \\ \varepsilon \mathbf{K} \times \varepsilon \mathbf{I} = \mathbf{0} \end{cases}$$




Figure 6: (a) Commutator of two zero-pitch screws ξ_1° and ξ_2° . \mathbf{N} is a unit line along the common perpendicular. (b): Commutator for two perpendicularly intersecting screws ξ_1, ξ_2 with pitches h_1, h_2 . In the figures, the commutator operator is denoted by \otimes .

The commutator of two unit twists ξ_1, ξ_2 with pitch $h_1, h_2 \neq \infty$ is given by

$$\begin{aligned}\xi_1 \times \xi_2 &= (1 + \varepsilon h_1)(1 + \varepsilon h_2)\xi_1^\circ \times \xi_2^\circ \\ &= (1 + \varepsilon(h_1 + h_2))\xi_1^\circ \times \xi_2^\circ.\end{aligned}$$

Since the commutator for two lines (see Fig. 6(a)) is given by [34]

$$\xi_1^\circ \times \xi_2^\circ = s_{\hat{\theta}}\mathbf{N}$$

where $\hat{\theta} = \theta + \varepsilon d$, and \mathbf{N} is the common perpendicular line between ξ_1° and ξ_2° , we have

$$\begin{aligned}\xi_1 \times \xi_2 &= (1 + \varepsilon(h_1 + h_2))s_{\hat{\theta}}\mathbf{N} \\ &= s_{\theta}(1 + \varepsilon(d \cot(\theta) + (h_1 + h_2)))\mathbf{N}.\end{aligned}\tag{7}$$

Hence, $\xi_1 \times \xi_2$ is a screw along \mathbf{N} , with pitch $d \cot(\theta) + (h_1 + h_2)$ and magnitude s_{θ} (see also [35]). In particular, when $d = 0, \theta = \frac{\pi}{2}$, the above equation simplifies to $\xi_1 \times \xi_2 = (1 + \varepsilon(h_1 + h_2))\mathbf{N}$ (see Fig. 6(b)). When $h_2 = \infty$ or $h_1 = \infty$, one has

$$\xi_1 \times \xi_2 = s_{\theta}\varepsilon\mathbf{N}$$

The commutator vanishes if both h_1 and h_2 are infinite.

120 3. Lie Product, Lie Triple Product and Lie Triple System

Definition 1. The *Lie product* L of a screw system S is the vector space spanned by the commutator of all twists in S :

$$L \triangleq [S, S] = \{\xi_1 \times \xi_2 \mid \forall \xi_1, \xi_2 \in S\}.$$

By bilinearity of the commutator, if ξ_1, \dots, ξ_k is a basis of S , then the twists $\xi_i \times \xi_j, i, j \in \{1, \dots, k\}$ (though not necessarily linearly independent) span L .

Definition 2. The *Lie triple product* T of a screw system S is the vector space spanned by the triple products of all twists in S :

$$\begin{aligned} T &\triangleq [[S, S], S] = [L, S] = [S, L] \\ &= \{ (\xi_1 \times \xi_2) \times \xi_3 \mid \forall \xi_1, \xi_2, \xi_3 \in S \}. \end{aligned}$$

By bilinearity of the commutator, if ξ_1, \dots, ξ_k is a basis of S , then the twists $(\xi_i \times \xi_j) \times \xi_l, i, j, l \in \{1, \dots, k\}$ (though not necessarily linearly independent) span T . We say that S is a **Lie triple system** (LTS) if:

$$T = [[S, S], S] \subseteq S. \quad (8)$$

In general, it is shown in [28, 29] that the exponential $\exp(S)$ of a screw system S can be generated as in (1) if and only if S is a LTS. We shall give in the next section an exhaustive identification of all such screw systems. Note that Lie subalgebras are automatically closed under the triple product, and therefore are *trivial* LTSs. There are ten classes of Lie subalgebras, which correspond to ten classes of connected Lie subgroups of $\text{SE}(3)$ [19, 5]. The Lie subalgebras are well known and, therefore, they will not be considered under the LTS framework. The following theorem concerning LTSs holds [30, 36]. The proof is reported here for illustrative purpose.

Theorem 1. If S is a LTS, then $L = [S, S]$ is a Lie subalgebra of $\mathfrak{se}(3)$, i.e.:

$$\forall \xi_1 \times \xi_2, \xi_3 \times \xi_4 \in L \quad \Rightarrow \quad (\xi_1 \times \xi_2) \times (\xi_3 \times \xi_4) \in L$$

Proof. Since S is a LTS, we have $T = [[S, S], S] \subseteq S$. Given two general elements $\xi_1 \times \xi_2, \xi_3 \times \xi_4$ of L , by skew-symmetry (2) and Jacobi identity (3), we have:

$$\begin{aligned} (\xi_1 \times \xi_2) \times (\xi_3 \times \xi_4) &= \underbrace{((\xi_3 \times \xi_4) \times \xi_2)}_{\in S} \times \xi_1 \\ &\quad - \underbrace{((\xi_3 \times \xi_4) \times \xi_1)}_{\in S} \times \xi_2 \in L. \end{aligned}$$

Therefore, $L = [S, S]$ is closed under Lie product and is a Lie subalgebra. \square

Note that Theorem 1 is relevant for screw systems with dimension three or higher, since the Lie product of a two-system is, in general, a one-system and, thus, is automatically a Lie subalgebra. We also emphasize that Theorem 1 provides a *necessary* condition for a screw system to be LTS. Hence, it may be used to simplify the identification process for 3 to 5-D LTSs, by allowing us to rule out screw systems that are not LTSs without computing their triple product.

4. Identification of Lie Triple Screw Systems

In this section, we shall verify the triple product closure condition (2) for all screw systems of $\mathfrak{se}(3)$ using Hunt's classification (up to conjugation) of general and special screw systems [2], thereby exhaustively identifying all Lie triple systems.




Figure 7: Lie product and triple product of a general two-system $S_{2,g}$.




Figure 8: Lie product and triple product of the first special two-system $S_{2,1}$.

145 4.1. Two-systems

4.1.1. $S_{2,g}$ and $S_{2,1}$

In order to investigate whether a general two-system $S_{2,g}$ with basis pitches α and β , $\alpha \geq \beta$, is a LTS, we choose the reference point \mathbf{o} to be the center of the system, and the basis (6) so that the basis screws lie on \mathbf{I} and \mathbf{J} (denoted \mathbf{I}_α and \mathbf{J}_β , respectively). It is clear from (7) that the Lie product $L_{2,g}$ is spanned by a single screw along \mathbf{K} with pitch $\alpha + \beta$, namely

$$\mathbf{I}_\alpha \times \mathbf{J}_\beta = (1 + \varepsilon(\alpha + \beta))\mathbf{K} = \mathbf{K}_{\alpha+\beta}.$$

The triple product $T_{2,g}$ is the following two-system:

$$T_{2,g} = \text{span}(\mathbf{I}_{\alpha+2\beta}, \mathbf{J}_{2\alpha+\beta})$$

where

$$\mathbf{I}_{\alpha+2\beta} = \mathbf{J}_\beta \times \mathbf{K}_{\alpha+\beta}, \quad \mathbf{J}_{2\alpha+\beta} = \mathbf{K}_{\alpha+\beta} \times \mathbf{I}_\alpha.$$

$T_{2,g}$ shares the same center and basis lines as $S_{2,g}$ (see Fig. 7). Accordingly, the LTS condition (8), namely the former is contained in the latter, only when their basis pitches are identical, namely

$$\begin{cases} \alpha + 2\beta = \alpha \\ 2\alpha + \beta = \beta \end{cases} \Leftrightarrow \alpha = \beta = 0$$

which leads to the first special two-system $S_{2,1}^0$ with two zero basis pitches (see Fig. 8).




Figure 9: Lie product and triple product of the second special two-system $S_{2,2}$.




Figure 10: Lie product and triple product of the fourth special two-system $S_{2,4}$.

The above computation illustrates the efficacy of Hunt's classification in comparison to Gibson's subtypes, which may be collectively verified by the same equation. It is interesting to note that the cylindroid of $T_{2,g}$ has the same length as that of $S_{2,g}$, but it is rotated 90° counter-clockwise about the \mathbf{K} axis. By this geometric reasoning alone, we may see that $S_{2,g}$ is not a LTS.

4.1.2. $S_{2,2}$ and $S_{2,3}$

When $\alpha = \infty, \beta \neq \infty$, then $S_{2,2} = \text{span}(\epsilon \mathbf{I}, \mathbf{J}_\beta)$, and it is easy to verify that:

$$L_{2,2} = \text{span}(\epsilon \mathbf{K}), \quad T_{2,2} = \text{span}(\epsilon \mathbf{I}) \subset S_{2,2}.$$

Therefore, $S_{2,2}$ is a LTS (see Fig. 9)⁶. When $\alpha = \beta = \infty$, we end up with the third special two-system $S_{2,3} = \text{span}(\epsilon \mathbf{I}, \epsilon \mathbf{J})$, namely the planar translation algebra.

4.1.3. $S_{2,4}$ and $S_{2,5}$

Let $\beta = -k^2\alpha$ for some positive real number $k \neq 1$ as $\alpha \rightarrow \infty$. Then, shift the reference point \mathbf{o} as shown in [2], and choose an appropriate basis $\mathbf{i}, \mathbf{j}, \mathbf{k}$ so that $S_{2,4}$ is given by

$$S_{2,4} = \text{span}(\mathbf{I}, c_\zeta \epsilon \mathbf{I} + s_\zeta \epsilon \mathbf{J})$$

⁶Note that, here, β can either be 0 or take a finite non-zero real value, thus giving rise to two classes of LTSs if the Gibson-Hunt classification [20, 18] is adopted.

	$S_{2,j}$		$L_{2,j}$		$T_{2,j}$		
	basis	condition	basis	type	basis	type	
$S_{2,g}$	$\mathbf{I}_\alpha, \mathbf{J}_\beta$	$\alpha \neq \beta$	$\mathbf{K}_{\alpha+\beta}$	$S_{1,g}$	$\mathbf{I}_{\alpha+2\beta}, \mathbf{J}_{2\alpha+\beta}$	$S_{2,g}$	
$S_{2,1}$	$\mathbf{I}_\alpha, \mathbf{J}_\alpha$	$\alpha \neq 0$	$\mathbf{K}_{2\alpha}$	$S_{1,g}$	$\mathbf{I}_{3\alpha}, \mathbf{J}_{3\alpha}$	$S_{2,1}$	
$S_{2,1}^0$	\mathbf{I}, \mathbf{J}		\mathbf{K}	$S_{1,g}$	\mathbf{I}, \mathbf{J}	$S_{2,1}^0$	LTS
$S_{2,2}$	$\varepsilon \mathbf{I}, \mathbf{J}_\beta$		$\varepsilon \mathbf{K}$	$S_{1,s}$	$\varepsilon \mathbf{I}$	$S_{1,s}$	LTS
$S_{2,3}$	$\varepsilon \mathbf{I}, \varepsilon \mathbf{J}$		N/A	N/A	N/A	N/A	subalg.
$S_{2,4}$	$\mathbf{I}, c_\zeta \varepsilon \mathbf{I} + s_\zeta \varepsilon \mathbf{J}$	$\zeta \notin \{0, \pi/2\}$	$\varepsilon \mathbf{I}$	$S_{1,s}$	$\varepsilon \mathbf{K}$	$S_{1,s}$	
$S_{2,5}$	$\mathbf{I}, \varepsilon \mathbf{I}$		N/A	N/A	N/A	N/A	subalg.

Table 1: Lie products and Lie triple products of 2-systems. All subalgebras are highlighted with background shading.

with $\cot \zeta = 2k/(k^2 - 1), \zeta \notin \{0, \pi/2\}$. It is straightforward to see that (Fig. 10):

$$L_{2,4} = \text{span}(\varepsilon \mathbf{K}), \quad T_{2,4} = \text{span}(\varepsilon \mathbf{J}).$$

$S_{2,4}$ is not a LTS since $c_\zeta \varepsilon \mathbf{I} + s_\zeta \varepsilon \mathbf{J}$ is the only infinite-pitch screw in $S_{2,4}$ but is not aligned with $\varepsilon \mathbf{J}$, and therefore $T_{2,4} \not\subseteq S_{2,4}$.

The fifth special two-system emerges by setting $k = 1$, namely $\zeta = 0$. In this case, an appropriate basis (6) may be chosen so that:

$$S_{2,5} = \text{span}(\mathbf{I}, \varepsilon \mathbf{I}).$$

$S_{2,5}$ is the cylindrical algebra, and it comprises screws of all pitches along \mathbf{I} .

In summary, we have identified two two-systems as nontrivial LTSs, namely the first special two-system $S_{2,1}^0 = \text{span}(\mathbf{I}, \mathbf{J})$, comprising a pencil of converging zero-pitch screws, and the second special two-system $S_{2,2} = \text{span}(\varepsilon \mathbf{I}, \mathbf{J}_\beta)$, comprising a planar pencil of parallel equal-pitch screws and an infinite-pitch screw perpendicular to the pencil plane. All two-systems, along with their Lie product and Lie triple product, are listed in Table 1 for further reference.

4.2. Three-systems

4.2.1. Lie subalgebras

It is well known that the following three-systems are Lie subalgebras of $\mathfrak{se}(3)$:

- $S_{3,2}^0$ with three zero-pitch basis screws, i.e. the spherical algebra, comprising all zero-pitch screws passing through a point (see Fig. 12 with $\alpha = 0$);
- $S_{3,5}$, i.e. the planar (or helicoidal planar) algebra of all screws of equal pitch parallel to a common direction, as well as infinite-pitch screws perpendicular to this direction (see Fig. 13(b));
- $S_{3,6}$, i.e. the spatial translation algebra of infinite-pitch screws along all directions in space.




Figure 11: The first special three-system $S_{3,1}$ and its Lie product $L_{3,1}$.




Figure 12: The second special three-system $S_{3,2}$ and its Lie product $L_{3,2}$.

4.2.2. $S_{3,g}$, $S_{3,1}$ and $S_{3,2}$

Given a general three-system $S_{3,g}$ (see Fig. 5) with finite basis pitches α, β and γ , $\alpha \geq \beta \geq \gamma$, we choose the reference point \mathbf{o} to be the center of the system, and the basis (6) so that the basis screws lie on $\mathbf{K}, \mathbf{I}, \mathbf{J}$ respectively (denoted $\mathbf{K}_\alpha, \mathbf{I}_\beta, \mathbf{J}_\gamma$ respectively).

If $\alpha = \beta$ or $\beta = \gamma$, $S_{3,g}$ becomes a first special three-system $S_{3,1}$, as illustrated in Fig. 11. Equal-pitch screws with $h \neq \alpha$ form ∞^1 reguli lying on a pencil of coaxial hyperboloids of revolution. Screws with pitch α form a planar pencil located at the reference point \mathbf{o} in the \mathbf{ki} plane. The Lie product $L_{3,1}$ is also a $S_{3,1}$ system, with basis pitches 2α and $\alpha + \gamma$. In this case, equal-pitch screws with $h \neq \alpha + \gamma$ lie on the same hyperboloids as those of the original $S_{3,1}$ system, but on the complementary reguli.

If $\alpha = \beta = \gamma$, $S_{3,g}$ becomes a second special three-system $S_{3,2}$, as illustrated in Fig. 12. $S_{3,2}$ comprises a bundle of equal-pitch screws. Its Lie product $L_{3,2}$ is also a $S_{3,2}$ system with a bundle of equal-pitch screws with pitch twice as that of the original system.

The Lie product $L_{3,g}$ is also a 3-system sharing the same center and basis lines with $S_{3,g}$, but with finite basis pitches $\beta + \gamma, \alpha + \gamma, \alpha + \beta$:

$$L_{3,g} = \text{span}(\mathbf{K}_{\beta+\gamma}, \mathbf{I}_{\alpha+\gamma}, \mathbf{J}_{\alpha+\beta}).$$

According to Theorem 1, $S_{3,g}$ is a LTS only if (meaning necessarily but not




Figure 13: The fourth special three-system $S_{3,4}$ and its Lie product $L_{3,4}$.

sufficiently) $L_{3,g}$ is one of the Lie subalgebras listed in Sec. 4.2.1. Since both $S_{3,5}$ and $S_{3,6}$ have one or more infinite basis pitches, the only possible case is $L_{3,g} = S_{3,2}^0$ which is true if and only if

$$\beta + \gamma = \alpha + \gamma = \alpha + \beta = 0 \quad \Leftrightarrow \quad \alpha = \beta = \gamma = 0$$

195 namely, if all basis pitches of the original system $S_{3,g}$ are zero. Therefore, $S_{3,g}$ is a LTS only if it coincides with the spherical algebra $S_{3,2}^0$, thus being a trivial LTS. The same reasoning applies to $S_{3,1}$.

4.2.3. $S_{3,3}$ and $S_{3,4}$

When one basis pitch is infinite, say $\alpha = \infty$, i.e.

$$S_{3,3} = \text{span}(\epsilon \mathbf{K}, \mathbf{I}_\beta, \mathbf{J}_\gamma),$$

we have

$$L_{3,3} = \text{span}(\mathbf{K}_{\beta+\gamma}, \epsilon \mathbf{I}, \epsilon \mathbf{J})$$

which is the Lie subalgebra $S_{3,5}$. The triple product is given by

$$T_{3,3} = \text{span}(\epsilon \mathbf{K}, \mathbf{I}_{\beta+2\gamma}, \mathbf{J}_{2\beta+\gamma}).$$

The three-system is a LTS if and only if:

$$\begin{cases} \beta + 2\gamma = \beta, \\ 2\beta + \gamma = \gamma. \end{cases} \quad \Leftrightarrow \quad \beta = \gamma = 0.$$

200 Therefore, only $S_{3,4}^0$ (with $\beta = \gamma = 0$) is a LTS. The fourth special three-system $S_{3,4}$ and its Lie product $L_{3,4}$ are shown in Fig. 13. In particular, $S_{3,4}^0$ consists of zero-pitch screws lying on a field of lines and an infinite-pitch screw perpendicular to the field plane. Its Lie product $L_{3,4}$ is the planar algebra $S_{3,5}^0$.




Figure 16: The ninth special three-system $S_{3,9}$, its Lie product $L_{3,9}$ and triple product $T_{3,9}$.




Figure 17: The tenth special three-system $S_{3,10}$, its Lie product $L_{3,10}$ and triple product $T_{3,10}$.

210 Equal-pitch screws with $h \neq \beta$ form ∞^1 reguli lying on a pencil of hyperbolic paraboloids, which have one line in common. The screws of pitch β lie on a degenerate regulus consisting of two perpendicular line pencils, one consisting of concurrent lines and the other consisting of coplanar parallel lines.

In this case, $L_{3,8}$ and $T_{3,8}$ are given by

$$\begin{cases} L_{3,8} = \text{span}(\mathbf{J}_{\beta}, \epsilon \mathbf{J}) = \text{span}(\mathbf{J}, \epsilon \mathbf{J}) \\ T_{3,8} = \text{span}(\mathbf{K}_{2\beta}, \epsilon \mathbf{K}, \mathbf{I}, \epsilon \mathbf{I}) = \text{span}(\mathbf{K}, \epsilon \mathbf{K}, \mathbf{I}, \epsilon \mathbf{I}) \end{cases}$$

$T_{3,8}$ is the fifth special four-system $S_{4,5}$, with dimension exceeding that of $S_{3,8}$.

215 Thus $T_{3,8} \not\subseteq S_{3,8}$ and $S_{3,8}$ is not a LTS.

4.2.5. $S_{3,9}$ and $S_{3,10}$

Using the same bases adopted for $S_{3,7}$ and $S_{3,8}$, we let $\beta = \infty$. When $\zeta \neq 0$, we have $S_{3,9}$, which is given by (see Fig. 16):

$$S_{3,9} = \text{span}(\mathbf{K}, c_{\zeta}\epsilon \mathbf{K} + s_{\zeta}\epsilon \mathbf{J}, \epsilon \mathbf{I})$$

Aside from a planar pencil of infinite-pitch screws spanned by $\epsilon \mathbf{I}$ and $c_{\zeta}\epsilon \mathbf{K} + s_{\zeta}\epsilon \mathbf{J}$, finite equal-pitch screws of $S_{3,9}$ form ∞^1 parallel planar pencils lying on

	$S_{3,j}$		$L_{3,j}$		$T_{3,j}$		
	basis	condition	basis	type	basis	type	
$S_{3,g}$	$\mathbf{K}_\alpha, \mathbf{I}_\beta, \mathbf{J}_\gamma$	$\alpha > \beta > \gamma$	$\mathbf{K}_{\beta+\gamma}, \mathbf{I}_{\alpha+\gamma}, \mathbf{J}_{\alpha+\beta}$	$S_{3,g}$	$\mathbf{K}, \mathbf{I}, \mathbf{J}, \varepsilon\mathbf{K}, \varepsilon\mathbf{I}, \varepsilon\mathbf{J}$	$\mathfrak{se}(3)$	
$S_{3,1}$	$\mathbf{K}_\alpha, \mathbf{I}_\alpha, \mathbf{J}_\gamma$	$\alpha \neq \gamma$	$\mathbf{K}_{\alpha+\gamma}, \mathbf{I}_{\alpha+\gamma}, \mathbf{J}_{2\alpha}$	$S_{3,1}$	$\mathbf{K}, \mathbf{I}, \mathbf{J}_{2\alpha+\gamma}, \varepsilon\mathbf{K}, \varepsilon\mathbf{I}$	$S_{5,g}$	
$S_{3,2}$	$\mathbf{K}_\alpha, \mathbf{I}_\alpha, \mathbf{J}_\alpha$	$\alpha \neq 0$	$\mathbf{K}_{2\alpha}, \mathbf{I}_{2\alpha}, \mathbf{J}_{2\alpha}$	$S_{3,1}$	$\mathbf{K}_{3\alpha}, \mathbf{I}_{3\alpha}, \mathbf{J}_{3\alpha}$	$S_{3,2}$	
$S_{3,2}^0$	$\mathbf{K}, \mathbf{I}, \mathbf{J}$		$\mathbf{K}, \mathbf{I}, \mathbf{J}$	$S_{3,2}^0$	$\mathbf{K}, \mathbf{I}, \mathbf{J}$	$S_{3,2}^0$	subalg.
$S_{3,3}$	$\varepsilon\mathbf{K}, \mathbf{I}_\beta, \mathbf{J}_\gamma$	$\beta \neq \gamma$	$\mathbf{K}_{\beta+\gamma}, \varepsilon\mathbf{I}, \varepsilon\mathbf{J}$	$S_{3,5}$	$\varepsilon\mathbf{K}, \mathbf{I}_{\beta+2\gamma}, \mathbf{J}_{2\beta+\gamma}$	$S_{3,3}$	
$S_{3,4}$	$\varepsilon\mathbf{K}, \mathbf{I}_\beta, \mathbf{J}_\beta$		$\mathbf{K}_{2\beta}, \varepsilon\mathbf{I}, \varepsilon\mathbf{J}$	$S_{3,5}$	$\varepsilon\mathbf{K}, \mathbf{I}_{3\beta}, \mathbf{J}_{3\beta}$	$S_{3,4}$	
$S_{3,4}^0$	$\varepsilon\mathbf{K}, \mathbf{I}, \mathbf{J}$		$\mathbf{K}, \varepsilon\mathbf{I}, \varepsilon\mathbf{J}$	$S_{3,5}$	$\varepsilon\mathbf{K}, \mathbf{I}, \mathbf{J}$	$S_{3,4}^0$	LTS
$S_{3,5}$	$\varepsilon\mathbf{K}, \varepsilon\mathbf{I}, \mathbf{J}_\gamma$		$\varepsilon\mathbf{K}, \varepsilon\mathbf{I}$	$S_{2,3}$	$\varepsilon\mathbf{K}, \varepsilon\mathbf{I}$	$S_{2,3}$	subalg.
$S_{3,6}$	$\varepsilon\mathbf{K}, \varepsilon\mathbf{I}, \varepsilon\mathbf{J}$		N/A	N/A	N/A	N/A	subalg.
$S_{3,7}$	$\mathbf{K}, c_\zeta\varepsilon\mathbf{K} + s_\zeta\varepsilon\mathbf{J}, \mathbf{I}_\beta$		$s_\zeta\varepsilon\mathbf{K} - c_\zeta\varepsilon\mathbf{J}, \varepsilon\mathbf{I}, \mathbf{J}_\beta$	$S_{3,9}$	$\mathbf{K}, \mathbf{I}, \varepsilon\mathbf{K}, \varepsilon\mathbf{I}, \varepsilon\mathbf{J}$	$S_{5,s}$	
$S_{3,8}$	$\mathbf{K}, \varepsilon\mathbf{K}, \mathbf{I}_\beta$		$\mathbf{J}, \varepsilon\mathbf{J}$	$S_{2,5}$	$\mathbf{K}, \mathbf{I}, \varepsilon\mathbf{K}, \varepsilon\mathbf{I}$	$S_{4,5}$	
$S_{3,9}$	$\mathbf{K}, c_\zeta\varepsilon\mathbf{K} + s_\zeta\varepsilon\mathbf{J}, \varepsilon\mathbf{I}$		$\varepsilon\mathbf{I}, \varepsilon\mathbf{J}$	$S_{2,3}$	$\varepsilon\mathbf{I}, \varepsilon\mathbf{J}$	$S_{2,3}$	
$S_{3,10}$	$\mathbf{K}, \varepsilon\mathbf{K}, \varepsilon\mathbf{I}$		$\varepsilon\mathbf{J}$	$S_{1,s}$	$\varepsilon\mathbf{I}$	$S_{1,s}$	LTS

Table 2: Lie products and Lie triple products of 3-systems. All subalgebras are highlighted with background shading.

∞^1 parallel planes perpendicular to \mathbf{i} . It is easy to verify that:

$$L_{3,9} = \text{span}(\varepsilon\mathbf{I}, \varepsilon\mathbf{J}), \quad T_{3,9} = \text{span}(\varepsilon\mathbf{I}, \varepsilon\mathbf{J}) \not\subseteq S_{3,9}$$

so that $S_{3,9}$ is not a LTS.

The tenth special three-system arises by setting $k = 1$ ($\zeta = 0$; see Fig. 17):

$$S_{3,10} = \text{span}(\mathbf{K}, \varepsilon\mathbf{K}, \varepsilon\mathbf{I}).$$

Aside from a planar pencil of infinite-pitch screws spanned by $\varepsilon\mathbf{I}$ and $\varepsilon\mathbf{K}$, $S_{3,10}$ comprises screws of arbitrary finite pitch lying on parallel lines of a planar pencil perpendicular to \mathbf{i} . It is easy to verify that:

$$L_{3,10} = \{\varepsilon\mathbf{J}\}, \quad T_{3,10} = \{\varepsilon\mathbf{I}\} \subset S_{3,10}.$$

and therefore $S_{3,10}$ is a LTS.

In summary, we have identified two three-systems as nontrivial LTSs, namely the fourth special three-system $S_{3,4}^0 = \text{span}(\varepsilon\mathbf{K}, \mathbf{I}, \mathbf{J})$, comprising a field of zero-pitch screws, and the tenth special three-system $S_{3,10} = \text{span}(\varepsilon\mathbf{K}, \mathbf{K}, \varepsilon\mathbf{I})$, comprising a planar pencil of parallel screws with arbitrary pitches. All the three-systems along with their Lie product and triple product are listed in Table 2.

4.3. Four-systems

4.3.1. Lie subalgebras

Only one four-system, namely $S_{4,3}$:

$$S_{4,3} = \text{span}(\varepsilon\mathbf{I}, \varepsilon\mathbf{J}, \mathbf{K}, \varepsilon\mathbf{K})$$

is a Lie subalgebra of $\mathfrak{se}(3)$, known as the Schönflies algebra [19, 5]. It consists of all finite-pitch screws along lines parallel to a common direction and infinite-pitch screws along any direction in space.




Figure 18: The general four-system $S_{4,g}$ and its Lie product $L_{4,g}$.

4.3.2. $S_{4,g}$ and $S_{4,1}$

Any four-system is characterized by a reciprocal two-system, whose reciprocal basis screws have pitches $-\alpha$ and $-\beta$. In the case of $S_{4,g}$ and $S_{4,1}$, we choose the reference point \mathbf{o} to be the center point of the corresponding reciprocal two-system, and an appropriate basis (6) such that $S_{4,g}$ is given by (see Fig. 18):

$$S_{4,g} = \text{span}(\mathbf{I}_\alpha, \mathbf{J}_\beta, \mathbf{K}, \varepsilon \mathbf{K}).$$

As long as α, β are finite, both in the case of $\alpha \neq \beta$ and in the case $\alpha = \beta$, the Lie product:

$$\begin{aligned} L_{4,g} &= \text{span}(\mathbf{I}_\beta, \varepsilon \mathbf{I}, \mathbf{J}_\alpha, \varepsilon \mathbf{J}, \mathbf{K}_{\alpha+\beta}) \\ &= \text{span}(\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \mathbf{K}_{\alpha+\beta}) \end{aligned}$$

230 is a 5-D vector subspace and cannot be a Lie subalgebra. Therefore, neither $S_{4,g}$ nor $S_{4,1}$ is a LTS.

4.3.3. $S_{4,2}$

When the basis pitch $\alpha = \infty$, we have the second special four-system $S_{4,2}$ (see Fig. 19):

$$S_{4,2} = \text{span}(\varepsilon \mathbf{I}, \mathbf{J}_\beta, \mathbf{K}, \varepsilon \mathbf{K}).$$

The Lie product:

$$L_{4,2} = \text{span}(\mathbf{I}_\beta, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}) = \text{span}(\mathbf{I}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K})$$

is the Schönflies algebra. However, the triple product

$$T_{4,2} = \text{span}(\varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \mathbf{K}_\beta, \varepsilon \mathbf{K}) = \text{span}(\varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{K})$$

is a special five-system; then $T_{4,2} \not\subseteq S_{4,2}$. Therefore, $S_{4,2}$ is not a LTS.

4.3.4. $S_{4,4}$ and $S_{4,5}$

Let the basis pitches satisfy $\beta = -k^2\alpha$ for some positive number $k \neq 1$ as $\alpha \rightarrow \infty$. Shift the reference point \mathbf{o} and choose an appropriate basis $\mathbf{i}, \mathbf{j}, \mathbf{k}$ such that $S_{4,4}$ is given by (see Fig. 20)

$$S_{4,4} = \text{span}(\mathbf{I}, c_\zeta \varepsilon \mathbf{I} + s_\zeta \varepsilon \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{K})$$




Figure 19: The second special four-system $S_{4,2}$ and its Lie product $L_{4,2}$.




Figure 20: The fourth special four-system $S_{4,4}$ and its Lie product $L_{4,4}$.

with $\cot \zeta = 2k/(k^2 - 1)$, $\zeta \neq 0$. We have:

$$\begin{aligned} L_{4,4} &= \text{span}(s_\zeta \varepsilon \mathbf{I} - c_\zeta \varepsilon \mathbf{J}, \mathbf{J}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}) \\ &= \text{span}(\varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}) \end{aligned}$$

which is a Schönflies algebra, and

$$T_{4,4} = \text{span}(\mathbf{I}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{K})$$

235 which is a special five-system. Since $T_{4,4} \not\subseteq S_{4,4}$, $S_{4,4}$ is not a LTS.

The fifth special four-system arises by letting $k = 1$ (and therefore $\zeta = 0$; see Fig. 21)

$$S_{4,5} = \text{span}(\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J})$$

$$L_{4,5} = \text{span}(\mathbf{K}, \varepsilon \mathbf{K})$$

and

$$T_{4,5} = \text{span}(\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}) = S_{4,5}.$$

Therefore $S_{4,5}$ is a LTS.

All four-systems along with their Lie product and triple product are listed in Table 3.




Figure 21: The fifth special four-system $S_{4,5}$ and its Lie product $L_{4,5}$.

	$S_{4,j}$		$L_{4,j}$		$T_{4,j}$		
	basis	condition	basis	type	basis	type	
$S_{4,g}$	$\mathbf{I}_\alpha, \mathbf{J}_\beta, \mathbf{K}, \varepsilon \mathbf{K}$	$\alpha \neq \beta$	$\mathbf{I}, \mathbf{J}, \mathbf{K}_{\alpha+\beta}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}$	$S_{5,g}$	$\mathbf{I}, \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$	$\mathfrak{se}(3)$	
$S_{4,1}$	$\mathbf{I}_\alpha, \mathbf{J}_\alpha, \mathbf{K}, \varepsilon \mathbf{K}$		$\mathbf{I}, \mathbf{J}, \mathbf{K}_{2\alpha}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}$	$S_{5,g}$	$\mathbf{I}, \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$	$\mathfrak{se}(3)$	
$S_{4,2}$	$\varepsilon \mathbf{I}, \mathbf{J}_\beta, \mathbf{K}, \varepsilon \mathbf{K}$		$\mathbf{I}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$	$S_{4,3}$	$\mathbf{J}, \mathbf{K}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$	$S_{5,s}$	
$S_{4,3}$	$\varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{K}$		$\varepsilon \mathbf{I}, \varepsilon \mathbf{J}$	$S_{2,3}$	$\varepsilon \mathbf{I}, \varepsilon \mathbf{J}$	$S_{2,3}$	subalg.
$S_{4,4}$	$\mathbf{I}, c_\zeta \varepsilon \mathbf{I} + s_\zeta \varepsilon \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{K}$	$\zeta \notin \{0, \pi/2\}$	$\mathbf{J}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$	$S_{4,3}$	$\mathbf{I}, \mathbf{K}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$	$S_{5,s}$	
$S_{4,5}$	$\mathbf{I}, \mathbf{J}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}$		$\mathbf{K}, \varepsilon \mathbf{K}$	$S_{2,5}$	$\mathbf{I}, \mathbf{J}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}$	$S_{4,5}$	LTS

Table 3: Lie products and Lie triple products of 4-systems. All subalgebras are highlighted with background shading.

4.4. Five-systems

240 4.4.1. $S_{5,g}$

A general five-system $S_{5,g}$ is reciprocal to a one-system with pitch $-\alpha$. Choose the reference point \mathbf{o} to be a point on the axis of the system, and an appropriate basis $\mathbf{i}, \mathbf{j}, \mathbf{k}$ such that $S_{5,g}$ is given by:

$$S_{5,g} = \text{span}(\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \mathbf{K}_\alpha).$$

It is straightforward to verify that:

$$L_{5,g} = \text{span}(\mathbf{I}_\alpha, \varepsilon \mathbf{J}, \mathbf{J}_\alpha, \varepsilon \mathbf{I}, \mathbf{K}, \varepsilon \mathbf{K}) = \mathfrak{se}(3).$$

However, $T_{5,g} = [[S_{5,g}, S_{5,g}], S_{5,g}] = [\mathfrak{se}(3), S_{5,g}] = \mathfrak{se}(3) \not\subseteq S_{5,g}$. Accordingly, $S_{5,g}$ is not a LTS.

4.4.2. $S_{5,s}$

The special five-system $S_{5,s}$ is given by

$$S_{5,s} = \text{span}(\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}).$$

It is straightforward to verify that

$$L_{5,s} = \text{span}(\varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{K})$$

which is a Schönflies algebra, and:

$$T_{5,s} = S_{5,s}.$$

i	j	$S_{i,j}$	$L_{i,j}$	$T_{i,j}$
2	1	\mathbf{I}, \mathbf{J}	\mathbf{K}	\mathbf{I}, \mathbf{J}
	2	$\varepsilon \mathbf{I}, \mathbf{J}_\beta$	$\varepsilon \mathbf{K}$	$\varepsilon \mathbf{I}$
3	4	$\mathbf{I}, \mathbf{J}, \varepsilon \mathbf{K}$	$\varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \mathbf{K}$	$\mathbf{I}, \mathbf{J}, \varepsilon \mathbf{K}$
	10	$\varepsilon \mathbf{I}, \mathbf{K}, \varepsilon \mathbf{K}$	$\varepsilon \mathbf{J}$	$\varepsilon \mathbf{I}$
4	5	$\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}$	$\mathbf{K}, \varepsilon \mathbf{K}$	$\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}$
5	1	$\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$	$\varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{K}$	$\mathbf{I}, \varepsilon \mathbf{I}, \mathbf{J}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$

Table 4: Lie triple screw systems of $\mathfrak{se}(3)$ which are not Lie subalgebras (represented by basis screws).

Therefore, $S_{5,s}$ is a LTS.

245 In this section, we have identified all Lie triple systems from Hunt's complete classification of screw systems [2]. Altogether, there are six classes of screw systems that fall in the category of Lie triple screw systems and, at the same time, are not Lie subalgebras. Our results are summarized in Table 4.

5. Dual Quaternion Characterization of the Exponential Images of LTSs

250

The exponential images of the LTSs identified in Sec. 4 correspond to symmetric subspaces of $\text{SE}(3)$ [29]. A full analysis of their geometric properties and applications is developed in [29]. In this paper, we investigate symmetric subspaces from a different perspective, namely their characterization by using dual quaternions.

255

Elements of $\text{SE}(3)$ can be represented as points in \mathbb{RP}^7 under the dual quaternion representation [19, Ch. 9]:

$$\mathbf{g} = a_0 + a_1 \mathbf{I} + a_2 \mathbf{J} + a_3 \mathbf{K} + b_0 \varepsilon + b_1 \varepsilon \mathbf{I} + b_2 \varepsilon \mathbf{J} + b_3 \varepsilon \mathbf{K} \in \text{SE}(3)$$

where $\mathbf{I}, \mathbf{J}, \mathbf{K}, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$ are as defined in (6). The homogeneous coordinate $(a_0 : a_1 : a_2 : a_3 : b_0 : b_1 : b_2 : b_3)$ satisfies the homogeneous quadratic constraint $a_0 b_0 + a_1 b_1 + a_2 b_2 + a_3 b_3 = 0$, which defines a 6-D quadric in \mathbb{RP}^7 known as the *Study quadric* (denoted Q_s ; see Fig. 22(a)). In particular, a pure rotation about an unit vector $\mathbf{w} = (w_1, w_2, w_3)$ with magnitude θ is given by:

$$e^{\theta \mathbf{w}/2} = c_{\theta/2} + s_{\theta/2} \mathbf{w} = c_{\theta/2} + s_{\theta/2} w_1 \mathbf{I} + s_{\theta/2} w_2 \mathbf{J} + s_{\theta/2} w_3 \mathbf{K}$$

and a pure translation along a unit vector $\mathbf{v} = (v_1, v_2, v_3)$ with magnitude t is given by:

$$e^{\varepsilon t \mathbf{v}/2} = 1 + \frac{t}{2} \varepsilon \mathbf{v} = 1 + \frac{t}{2} v_1 \varepsilon \mathbf{I} + \frac{t}{2} v_2 \varepsilon \mathbf{J} + \frac{t}{2} v_3 \varepsilon \mathbf{K}$$

Note that the ideal three-plane defined by:

$$A_\infty := \{(a_0 : a_1 : a_2 : a_3 : b_0 : b_1 : b_2 : b_3) \mid a_0 = a_1 = a_2 = a_3 = 0\}$$

is completely contained in Q_s , and it comprises the only elements on Q_s that do not represent elements of $\text{SE}(3)$ (see Fig. 22(b)). We shall refer to $Q_s - A_\infty \simeq$




Figure 22: (a) A 2-D analogue of the 6-D Study quadric Q_s in \mathbb{RP}^7 ; (b) the subset of Q_s that corresponds to $SE(3)$

$SE(3)$ as the set of proper dual quaternions. We shall also denote the ideal plane by $\varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J} \wedge \varepsilon \mathbf{K}$, where \wedge is the projective span operator and $\varepsilon, \varepsilon \mathbf{I}, \varepsilon \mathbf{J}, \varepsilon \mathbf{K}$ are the corresponding basis vectors.

Closely related to our work is a series of studies on sub-varieties of the Study quadric conducted by Selig [19, 37, 38]. Such sub-varieties arise either in motions satisfying natural geometric constraints (such as the point-plane constraint) [37], or in the dual quaternion representation of Lie subgroups of $SE(3)$ [38].

5.1. LTSs containing only zero- or infinite-pitch screws

From (4) and (5), we see that the exponential image of a one-system spanned by a zero-pitch or infinite-pitch screw ξ is given by:

$$\left\{ e^{\theta \xi^\circ / 2} \mid \forall \theta \in \mathbb{R} \right\} = \left\{ c_{\theta/2} + s_{\theta/2} \xi \mid \forall \theta \in \mathbb{R} \right\} = 1 \wedge \xi^\circ$$

or

$$\left\{ e^{\varepsilon \theta \xi^\circ / 2} \mid \forall \theta \in \mathbb{R} \right\} = \left\{ 1 + \varepsilon \frac{\theta}{2} \xi^\circ \mid \forall \theta \in \mathbb{R} \right\} = 1 \wedge \varepsilon \xi^\circ$$

It is a one-parameter subgroup of $SE(3)$, and corresponds to a line spanned by 1 and ξ° (or $\varepsilon \xi^\circ$) that lies completely on Q_s [19]. More generally, if a k -D screw system S comprises only zero- or infinite-pitch screws, its exponential image is given by:

$$\exp S = 1 \wedge S$$

which is a k -plane completely contained in Q_s . Based on this observation, we can immediately deduce that the exponential image of $S_{2,1}^0$, $S_{2,2}^0$ and $S_{3,4}^0$ are given by:

$$\begin{cases} \exp(S_{2,1}^0) = 1 \wedge S_{2,1}^0 = 1 \wedge \mathbf{I} \wedge \mathbf{J} \\ \exp(S_{2,2}^0) = 1 \wedge S_{2,2}^0 = 1 \wedge \mathbf{J} \wedge \varepsilon \mathbf{I} - \varepsilon \mathbf{I} \\ \exp(S_{3,4}^0) = 1 \wedge S_{3,4}^0 = 1 \wedge \mathbf{I} \wedge \mathbf{J} \wedge \varepsilon \mathbf{K} - \varepsilon \mathbf{K} \end{cases}$$

	normal form	proper submanifolds	general form
A-plane	$A_0 = 1 \wedge \mathbf{I} \wedge \mathbf{J} \wedge \mathbf{K}$	spatial rotation group	cosets of conjugates
	$A_2 = 1 \wedge \mathbf{K} \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J}$	planar Euclidean group	cosets of conjugates
	$A_\infty = \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J} \wedge \varepsilon \mathbf{K}$	N/A	invariant
B-plane	$B_1 = 1 \wedge \mathbf{I} \wedge \mathbf{J} \wedge \varepsilon \mathbf{K}$	$\exp(S_{3,4}^0)$ (symmetric subspace)	symmetric subspace
	$B_3 = 1 \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J} \wedge \varepsilon \mathbf{K}$	spatial translation group	cosets

Table 5: 3-planes of the Study quadric. Note that left/right transformation of a symmetric subspace is still a symmetric subspace.

where, in the latter two cases, the ideal point $\varepsilon \mathbf{I}$ or $\varepsilon \mathbf{K}$ (which does not correspond to proper rigid transformations) is excluded. In general, the exponential image of a one-system spanned by a finite non-zero pitch screw is not an algebraic sub-variety under dual quaternion coordinates. Consequently, $S_{2,2}$ with $\beta \notin \{0, \infty\}$ is not an algebraic sub-variety of Q_s either.

It is also interesting to note that according to Selig's earlier study [19, 39], $\exp(S_{3,4}^0)$ (without removing the ideal point) is exactly the B -plane that meets A_∞ at a single point (called B_1 -plane in [39]), while $\exp(S_{2,1}^0)$ and $\exp(S_{2,2}^0)$ are two 2-planes contained in $\exp(S_{3,4}^0)$. For convenience, we recall Selig's complete classification of 3-planes of Q_s in Table 5 along with their proper submanifolds. It should be clear that this classification is up to multiplication on the left and/or right by proper dual quaternions [39]: such proper transformations are linear transformations sending planes in Q_s to planes of the same dimension, and they leave both A_∞ and $Q_s - A_\infty \simeq \text{SE}(3)$ invariant. This should give rise to all 3-planes in Q_s from the normal forms listed in Table 5 (see also Fig. 22(a) for an analogue of the two families of 3-planes in Q_s).

5.2. LTSs comprising cylindrical algebras

Consider now the exponential image of the fifth special two-system $S_{2,5} = \text{span}(\mathbf{I}, \varepsilon \mathbf{I})$ (the cylindrical algebra):

$$\exp\left(\frac{\theta}{2}\mathbf{I} + \frac{\lambda}{2}\varepsilon\mathbf{I}\right) = c_{\theta/2} + s_{\theta/2}\mathbf{I} + \frac{\lambda}{2}\varepsilon(-s_{\theta/2} + c_{\theta/2}\mathbf{I}), \quad \forall \theta, \lambda \in \mathbb{R}$$

It is shown in [19] that this is exactly the intersection of Q_s with a three-space $1 \wedge \mathbf{I} \wedge \varepsilon \wedge \varepsilon \mathbf{I}$, a quadric prescribed by:

$$\begin{cases} a_0 b_0 + a_1 b_1 = 0 \\ a_2 = a_3 = b_2 = b_3 = 0 \end{cases} \quad (9)$$

This quadric contains an ideal line $\varepsilon \wedge \varepsilon \mathbf{I}$ which does not correspond to proper rigid transformations. By using this observation, we can deduce the exponential image of $S_{3,10}$, $S_{4,5}$ and $S_{5,s}$.

First, it is clear from Fig. 17 that $S_{3,10}$ comprises a parallel pencil of cylindrical algebras:

$$\exp(S_{3,10}) = \bigcup_{l \in \mathbb{R}} \exp(\text{span}(\mathbf{K} + l\mathbf{I}, \varepsilon \mathbf{K}))$$




Figure 23: (a) A 1-D analogue of an intersection quadric Q_s with a subspace in \mathbb{RP}^7 ; (b) a 1-D analogue of a tangential intersection of Q_s with a subspace of \mathbb{RP}^7 , resulting in a quadratic cone with the vertex being the point of tangency (denoted by the blue dot).

where $\text{span}(\mathbf{K} + \varepsilon l\mathbf{I}, \varepsilon \mathbf{K})$ is the cylindrical algebra along the line $\mathbf{K} + \varepsilon l\mathbf{I} = \mathbf{k} + \varepsilon(l\mathbf{j}) \times \mathbf{k}$. It is easy to verify that the exponential formula for this general cylindrical algebra is:

$$\exp(\text{span}(\mathbf{K} + \varepsilon l\mathbf{I}, \varepsilon \mathbf{K})) = (1 \wedge (\mathbf{K} + \varepsilon l\mathbf{I}) \wedge \varepsilon \wedge \varepsilon \mathbf{K}) \cap Q_s - \varepsilon \wedge \varepsilon \mathbf{K}$$

Consequently, we have:

$$\begin{aligned} \exp(S_{3,10}) &= \bigcup_{l \in \mathbb{R}} \exp(\text{span}(\mathbf{K} + \varepsilon l\mathbf{I}, \varepsilon \mathbf{K})) \\ &= \bigcup_{l \in \mathbb{R}} (1 \wedge (\mathbf{K} + \varepsilon l\mathbf{I}) \wedge \varepsilon \wedge \varepsilon \mathbf{K}) \cap Q_s - \varepsilon \wedge \varepsilon \mathbf{K} \\ &= (1 \wedge \mathbf{K} \wedge \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{K} - \varepsilon \mathbf{I}) \cap Q_s - \varepsilon \wedge \varepsilon \mathbf{K} \\ &= (1 \wedge \mathbf{K} \wedge \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{K}) \cap Q_s - \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{K} \end{aligned}$$

We have shown that $\exp(S_{3,10})$ corresponds to a quadric prescribed by (see Fig. 23(a) for a 1-D analogue):

$$\begin{cases} a_0 b_0 + a_3 b_3 = 0 \\ a_1 = a_2 = b_2 = 0 \end{cases}$$

which is a quadratic cone [40] in a 3-plane, with base quadric

$$\begin{cases} a_0 b_0 + a_3 b_3 = 0 \\ a_1 = a_2 = b_1 = b_2 = 0 \end{cases}$$

which is exactly a conjugate of the cylindrical group prescribed by (9), and vertex $\varepsilon \mathbf{I}$ (see Fig. 23(b) for a 1-D analogue).

By similar arguments, the exponential image of $S_{4,5}$ is given by:

$$\exp(S_{4,5}) = (1 \wedge \mathbf{I} \wedge \mathbf{J} \wedge \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J}) \cap Q_s - \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J}$$

which corresponds to a quadric in a 5-plane prescribed by (see Fig. 23(a) for a 1-D analogue)

$$\begin{cases} a_0b_0 + a_1b_1 + a_2b_2 = 0 \\ a_3 = b_3 = 0 \end{cases}$$

The exponential image of $S_{5,s}$ is given by:

$$\exp(S_{5,s}) = (1 \wedge \mathbf{I} \wedge \mathbf{J} \wedge \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J} \wedge \varepsilon \mathbf{K}) \cap Q_s - \varepsilon \wedge \varepsilon \mathbf{I} \wedge \varepsilon \mathbf{J} \wedge \varepsilon \mathbf{K}$$

which corresponds to a quadratic cone in a 6-plane, with base quadric $\exp(S_{4,5})$ and vertex $\varepsilon \mathbf{K}$ (see Fig. 23(b) for a 1-D analogue). Referring to [39], $\exp(S_{4,5})$ is exactly the space of all line symmetric motions (with a fixed base line).

In summary, all symmetric subspaces except $\exp(S_{2,2})$ (with $\beta \neq 0$) are either projective subspaces (excluding a subspace of ideal points) that are completely contained in Q_s , or quadrics (including quadratic cones) of Q_s (i.e. intersection of Q_s with a projective subspace of \mathbb{RP}^7 ; see [40, Ch. 1] for the concept of quadric and quadratic cone). Some of these spaces turn out to be certain sub-varieties of Q_s that have been discovered earlier. However, the connection to LTSs and symmetric subspaces was never made. Hopefully, such connections may be useful for analysis and design of over-constrained linkages and special motion generators [37, 39, 41, 42, 11, 29].

6. Conclusions

In this paper, we identified all six non-trivial Lie triple systems from Hunt's complete classification of screw systems [2], as shown in Table 4. The presented derivation was based on both algebraic Lie-group theory and screw geometry. We paid special attention to the geometric features characterizing each Lie triple system, as well as its associated Lie product and Lie triple product. Lie triple screw systems have been playing, unknowingly, an important role in the type synthesis of constant-velocity couplings and torsion-free parallel manipulators for many years. We have expanded the known portfolio of screw systems that have the same special algebraic and geometric properties as the screw systems characterizing the latter mechanisms, thus possibly opening the way to new designs.

For example, we illustrated in [43] that the 2-D LTS $S_{2,1}^0$ can be used to provide an elegant kinematic analysis of linkage-based 2-D CV couplings such as the Thompson CV joint [44] and Kocabas's CV joint [45]. In [43], we implemented the same kinematic principles in the design of a novel 2-DoF parallel wrist with both an ample rotation range and a fixed center of rotation compared to Omni-wrist III [46]. In [29], we developed a systematic type synthesis method for kinematic chains persistently generating Lie triple screw systems. A systematic type synthesis for parallel mechanisms generating LTSs is currently being developed.

Aside from applications in mechanism analysis and synthesis, the symmetric spaces corresponding to the Lie triple screw systems that we identified may

325 serve as important motion types for robotic applications. For example, $\exp(S_{2,1}^0)$ serves as the 2-D orientation workspace for robotic wrists (excluding the final roll axis, see [47]), quadcopters (roll and pitch), and the spindle of a five-axis machine [48]. The computational advantage of $\exp(S_{2,1}^0)$ in comparison to the traditional unit sphere model [47] may allow one to conduct spline interpolation on the orientation workspace using the exponential map [49, 50].

330 Acknowledgments

The authors gratefully acknowledge the financial support of the Italian Ministry of Education, Universities and Research through the PRIN grant No. 20124SMZ88. This work is also in partial fulfillment to China National Natural Science Foundation Grant No.51375413 and Shenzhen Municipal Science and Technology Planning Program for Basic Research No.JCYJ20120618150931822, 335 which supported Dr. Wu when he was working at HKUST.

References

- [1] Y. Wu, M. Carricato, Identification and geometric characterization of Lie triple screw systems, in: 14th IFToMM World Congress, Taipei, Taiwan, 340 October 25–30, 2015.
- [2] K. H. Hunt, Kinematic geometry of mechanisms, Clarendon Press, 1978.
- [3] O. Bottema, B. Roth, Theoretical kinematics, Dover Publication, 1978.
- [4] J. M. Selig, M. Carricato, Persistent rigid-body motions and Study’s “Ribaucour” problem, Journal of Geometry (2016) 1–21, DOI:10.1007/s00022-016-0331-5. 345
- [5] J. Hervé, The Lie group of rigid body displacements, a fundamental tool for mechanism design, Mechanism and Machine theory 34 (5) (1999) 719–730.
- [6] M. Carricato, V. Parenti-Castelli, Singularity-free fully-isotropic translational parallel mechanisms, The International Journal of Robotics Research 21 (2) (2002) 161–174. 350
- [7] Z. Huang, Q. Li, Type synthesis of symmetrical lower-mobility parallel mechanisms using the constraint-synthesis method, The International Journal of Robotics Research 22 (1) (2003) 59–79.
- [8] M. Carricato, Fully isotropic four-degrees-of-freedom parallel mechanisms for schoenflies motion, The International Journal of Robotics Research 24 (5) (2005) 397–414. 355
- [9] X. Kong, C. M. Gosselin, Type synthesis of parallel mechanisms, Springer, 2007.

- [10] M. L. Husty, M. Pfurner, H.-P. Schröcker, K. Brunnthaler, Algebraic methods in mechanism analysis and synthesis, *Robotica* 25 (06) (2007) 661–675.
- [11] J. Selig, Some remarks on the RRR linkage, in: *Advances in Robot Kinematics*, Springer, 2014, pp. 77–85.
- [12] M. Carricato, D. Zlatanov, Persistent screw systems, *Mechanism and Machine Theory* 73 (2014) 296–313.
- [13] P. Fanghella, C. Galletti, Metric relations and displacement groups in mechanism and robot kinematics, *Journal of Mechanical Design* 117 (3) (1995) 470–478.
- [14] J. Angeles, The qualitative synthesis of parallel manipulators, *Journal of Mechanical Design* 126 (4) (2004) 617–624.
- [15] M. Carricato, Decoupled and homokinetic transmission of rotational motion via constant-velocity joints in closed-chain orientational manipulators, *Journal of Mechanisms and Robotics* 1 (4) (2009) 041008.
- [16] J. Meng, G. Liu, Z. Li, A geometric theory for analysis and synthesis of sub-6 dof parallel manipulators, *IEEE Transactions on Robotics* 23 (4) (2007) 625–649.
- [17] J. M. Rico, J. J. Cervantes-Sánchez, A. Tadeo-Chávez, G. I. Pérez-Soto, J. Rocha-Chavarría, New considerations on the theory of type synthesis of fully parallel platforms, *Journal of Mechanical Design* 130 (11) (2008) 112302.
- [18] C. G. Gibson, K. H. Hunt, Geometry of screw systems—2, Classification of screw systems, *Mechanism and Machine Theory* 25 (1) (1990) 11–27.
- [19] J. M. Selig, *Geometric fundamentals of robotics*, Springer, 2005.
- [20] C. G. Gibson, K. H. Hunt, Geometry of screw systems—1, Screws: genesis and geometry, *Mechanism and Machine Theory* 25 (1) (1990) 1–10.
- [21] M. Carricato, J. M. Rico-Martínez, Persistent screw systems of dimension three, in: *Proc. of 13th World Congress in Mechanism and Machine Science*, Guanajuato, Mexico, 2011, pp. 1–12.
- [22] M. Carricato, Persistent screw systems of dimension four, in: *Latest Advances in Robot Kinematics*, Springer, 2012, pp. 147–156.
- [23] M. Carricato, Four-dimensional persistent screw systems of the general type, in: *Computational Kinematics*, Springer, 2014, pp. 299–306.
- [24] K. H. Hunt, Constant-velocity shaft couplings: a general theory, *Journal of Manufacturing Science and Engineering* 95 (2) (1973) 455–464.

- 395 [25] D. Zlatanov, I. A. Bonev, C. M. Gosselin, Constraint singularities as c-space singularities, in: *Advances in Robot Kinematics*, Springer, 2002, pp. 183–192.
- [26] I. A. Bonev, D. Zlatanov, C. M. Gosselin, Advantages of the modified Euler angles in the design and control of PKMs, in: *2002 Parallel Kinematic Machines International Conference*, 2002, pp. 171–188.
- 400 [27] Y. Wu, Z. Li, J. Shi, Geometric properties of zero-torsion parallel kinematics machines, in: *2010 IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS 2010)*, 2010, pp. 2307–2312.
- [28] Y. Wu, G. Liu, H. Löwe, Z. Li, Exponential submanifolds: a new kinematic model for mechanism analysis and synthesis, in: *2013 IEEE International Conference on Robotics and Automation (ICRA 2013)*, 2013, pp. 4177–4182.
- 405 [29] Y. Wu, H. Löwe, M. Carricato, Z. Li, Inversion symmetry of the euclidean group: Theory and application to robot kinematics, *IEEE Transactions on Robotics* 32 (2) (2016) 312–326.
- [30] O. Loos, *Symmetric spaces*, Vol. 1–2, W.A. Benjamin, 1969.
- 410 [31] W. Bertram, *The geometry of Jordan and Lie structures*, Springer, 2000.
- [32] J. Selig, A class of explicitly solvable vehicle motion problems, *IEEE Transactions on Robotics* 31 (3) (2015) 766–777.
- [33] J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Springer, 1972.
- 415 [34] J. M. McCarthy, G. S. Soh, *Geometric design of linkages*, Springer, 2010.
- [35] J. J. Cervantes-Sánchez, M. A. Moreno-Báez, J. M. Rico-Martínez, E. J. González-Galván, A novel geometrical derivation of the Lie product, *Mechanism and machine theory* 39 (10) (2004) 1067–1079.
- 420 [36] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Academic press, 1979.
- [37] J. Selig, Quadratic constraints on rigid-body displacements, *Journal of Mechanisms and Robotics* 2 (4) (2010) 041009.
- [38] J. Selig, On the geometry of the homogeneous representation for the group of proper rigid-body displacements, *Romanian Journal of Technical Sciences – Applied Mechanics* 58 (1–2) (2013) 153–176.
- 425 [39] J. Selig, M. Husty, Half-turns and line symmetric motions, *Mechanism and Machine Theory* 46 (2) (2011) 156–167.
- [40] H. Pottmann, J. Wallner, *Computational line geometry*, Springer, 2009.

- 430 [41] G. Hegedüs, J. Schicho, H.-P. Schröcker, Factorization of rational curves in the study quadric, *Mechanism and Machine Theory* 69 (2013) 142–152.
- [42] G. Hegedüs, J. Schicho, H.-P. Schröcker, The theory of bonds: A new method for the analysis of linkages, *Mechanism and Machine Theory* 70 (2013) 407–424.
- 435 [43] Y. Wu, M. Carricato, Design of a novel 3-dof serial-parallel robotic wrist: A symmetric space approach, in: *Proceedings of International Symposium on Robotics Research*, September 12 – 15, Sestri Levante, Italy, 2015.
- [44] G. A. Thompson, Constant velocity coupling and control system therefor, US Patent 7,144,326 (Dec. 5 2006).
- 440 [45] H. Kocabas, Design and analysis of a spherical constant velocity coupling mechanism, *ASME Journal of Mechanical Design* 129 (9) (2007) 991–998.
- [46] M. Rosheim, G. Sauter, New high-angulation omni-directional sensor mount, in: *International Symposium on Optical Science and Technology*, Seattle, WA, USA, July 7, 2002, pp. 163–174.
- 445 [47] R. P. Paul, C. N. Stevenson, Kinematics of robot wrists, *The International Journal of Robotics Research* 2 (1) (1983) 31–38.
- [48] Y. Wu, H. Wang, Z. Li, Quotient kinematics machines: concept, analysis, and synthesis, *Journal of mechanisms and robotics* 3 (4) (2011) 041004.
- 450 [49] F. C. Park, B. Ravani, Smooth invariant interpolation of rotations, *ACM Transactions on Graphics* 16 (3) (1997) 277–295.
- [50] Y. Wu, A. Müller, M. Carricato, The 2D orientation interpolation problem: A symmetric space approach, in: *Advances in Robot Kinematics*, Springer, 2016, to appear.