



Processes controlling the seasonal variations in ^{210}Pb and ^7Be at the Mt. Cimone WMO-GAW global station, Italy: a model analysis

Erika Brattich¹, Hongyu Liu², Laura Tositti¹, David B. Considine³, and James H. Crawford⁴

¹Department of Chemistry “G Ciamician”, Alma Mater Studiorum University of Bologna, 40126 Bologna (BO), Italy

²National Institute of Aerospace, Hampton, VA 23666, USA

³NASA Headquarters, Washington, DC 20546, USA

⁴NASA Langley Research Center, Hampton, VA 23681, USA

Correspondence to: Hongyu Liu (hongyu.liu-1@nasa.gov)

Received: 30 June 2016 – Published in Atmos. Chem. Phys. Discuss.: 4 August 2016

Revised: 20 December 2016 – Accepted: 22 December 2016 – Published: 24 January 2017

Abstract. We apply the Global Modeling Initiative (GMI) chemistry and transport model driven by NASA’s MERRA assimilated meteorological data to simulate the seasonal variations in two radionuclide aerosol tracers (terrestrial ^{210}Pb and cosmogenic ^7Be) at the WMO-GAW station of Mt. Cimone (44°12′ N, 10°42′ E; 2165 m a.s.l.; Italy), which is representative of free-tropospheric conditions most of the year, during 2005 with an aim to understand the roles of transport and precipitation scavenging processes in controlling their seasonality. The total precipitation field in the MERRA data set is evaluated with the Global Precipitation Climatology Project (GPCP) observations, and generally good agreement is found. The model reproduces reasonably the observed seasonal pattern of ^{210}Pb concentrations, characterized by a wintertime minimum due to lower ^{222}Rn emissions and weaker uplift from the boundary layer and summertime maxima resulting from strong convection over the continent. The observed seasonal behavior of ^7Be concentrations shows a winter minimum, a summer maximum, and a secondary spring maximum. The model captures the observed ^7Be pattern in winter–spring, which is linked to the larger stratospheric influence during spring. However, the model tends to underestimate the observed ^7Be concentrations in summer, partially due to the sensitivity to spatial sampling in the model. Model sensitivity experiments indicate a dominant role of precipitation scavenging (vs. dry deposition and convection) in controlling the seasonality of ^{210}Pb and ^7Be concentrations at Mt. Cimone.

1 Introduction

The use of atmospheric radionuclides to understand atmospheric dynamics, pollution transport and removal processes has a long history (e.g., Junge, 1963; Reiter et al., 1971; Gäggeler, 1995; Arimoto et al., 1999; Turekian and Graustein, 2003; WMO-GAW, 2004; Dibb, 2007; Rastogi and Sarin, 2008; Froehlich and Masarik, 2010; Lozano et al., 2012). It has been recognized that natural radionuclides are useful in a global monitoring network for atmospheric composition to support global climate change and air quality research, and therefore they are measured at many of the regional, global, and contributing-partner stations in the Global Atmosphere Watch (GAW) network of the World Meteorological Organization (WMO) (WMO-GAW, 2004). In particular, terrestrial ^{210}Pb and cosmogenic ^7Be natural radionuclides are helpful in the understanding of the roles of transport and/or scavenging in controlling the behaviors of radiatively active trace gases and aerosols (Feichter et al., 1991; Balkanski et al., 1993; Koch et al., 1996), as well as their anthropogenic (vs. natural) origin (e.g., Graustein and Turekian, 1996; Arimoto et al., 1999; Liu et al., 2004; Cuevas et al., 2013). They are routinely monitored at WMO-GAW stations around the world (Lee et al., 2004). Although ^{210}Pb and ^7Be have long (1998–2011) been measured at the Global WMO-GAW station of Mt. Cimone (Italy), their seasonal behavior has not been thoroughly elucidated (Lee et al., 2007; Tositti et al., 2014). Here we apply a state-of-the-art global chemistry and transport model (CTM) to the simulation of ^{210}Pb and ^7Be , with an objective to better understand the

roles of transport and precipitation scavenging processes in controlling their seasonal variations at Mt. Cimone.

Because of their contrasting natural origins, ^{210}Pb and ^7Be have been used as a pair to study the vertical transport and scavenging of aerosols (Koch et al., 1996). ^{210}Pb (half-life $\tau_{1/2} = 22.3$ years) is the decay daughter of ^{222}Rn ($\tau_{1/2} = 3.8$ days), which is emitted from soils by decay of ^{226}Ra . The oceanic input of ^{222}Rn is about 2 orders of magnitude less than the continental input and, because of the continental origin of ^{222}Rn , ^{210}Pb is considered a tracer of air masses with continental origin (Baskaran, 2011). ^7Be ($\tau_{1/2} = 53.3$ days) is a cosmogenic radionuclide generated by cosmic ray spallation reactions with nitrogen and oxygen (Lal et al., 1958). Most ($\sim 67\%$) of ^7Be is produced in the stratosphere and the remaining ($\sim 33\%$) is generated in the troposphere, particularly in the upper troposphere (Johnson and Viezee, 1981; Usoskin and Kovaltsov, 2008). ^7Be is thus considered a tracer of stratospheric influence (Viezee and Singh, 1980; Dibb et al., 1992, 1994; Liu et al., 2004, 2016) and subsidence (Feely et al., 1989; Koch et al., 1996; Liu et al., 2004). Once produced, both radionuclides rapidly attach to ubiquitous submicron aerosol particles in the ambient air (Papastefanou and Ioannidou, 1995; Winkler et al., 1998; Gaffney et al., 2004; Ioannidou et al., 2005) and are removed from the atmosphere mainly by wet and secondarily dry deposition (Kulan et al., 2006). The concentrations of these radionuclides in surface air thus depend on their sources, transport, wet and dry removal, and radioactive decay (in the case of ^7Be). Rainfall scavenging processes are generally more effective on ^{210}Pb than on ^7Be concentrations (Koch et al., 1996; Caillet et al., 2001; Likuku, 2006; Dueñas et al., 2009; Lozano et al., 2012).

Observational studies have previously been conducted to examine the factors influencing surface ^{210}Pb and ^7Be concentrations in Europe, the Middle East, and northern Africa. Different synoptic and mesoscale patterns are associated with the ranges of ^{210}Pb and ^7Be activity concentrations (Lozano et al., 2012, 2013). In southwestern Spain (El Arenosillo), for instance, low ^{210}Pb values are strongly linked to air masses from the Atlantic Ocean, whereas the highest values are associated with air masses clearly under the influence of continents, such as the Iberian Peninsula and North of Africa (Lozano et al., 2012). As for ^7Be , the highest ^7Be activity concentrations over the southwestern Iberian Peninsula are related to the arrival of air masses from middle latitudes, and in particular from the Canary Islands, the western Mediterranean Basin, and the north of Africa (Dueñas et al., 2011; Lozano et al., 2012).

With respect to ^{210}Pb and ^7Be spatial variability, ^{210}Pb concentrations in surface air are strongly dependent on whether it is located over land or ocean, whereas ^7Be concentration is mainly latitudinally dependent, due to their different production mechanisms. Generally speaking, in the Northern Hemisphere, higher ^7Be concentrations are present at middle latitudes ($20\text{--}50^\circ\text{N}$) because of the mix-

ing of stratospheric air into the upper troposphere along the tropopause discontinuity in midlatitude regions and subsequent convective mixing within the troposphere, which brings ^7Be -rich air masses into the planetary boundary layer and to the Earth's surface (Kulan et al., 2006). Lower ^7Be concentrations are towards the pole and towards the Equator (Kulan et al., 2006; Steinmann et al., 2013).

Many studies examined the seasonal behavior of ^{210}Pb and ^7Be at European midlatitude surface sites (e.g., Cannizzaro et al., 2004; Ioannidou et al., 2005; Daish et al., 2005; Todorovic et al., 2005; Likuku, 2006; Dueñas et al., 2009; Pham et al., 2011; Carvalho et al., 2013; Steinmann et al., 2013). High levels of ^{210}Pb during summer and low levels in winter were found, reflecting the differing rates of ^{222}Rn emanation from soil above the European land mass during winter (wet or snow-covered soil) and summer (dry soil) (Hötzl and Winkler, 1987; Caillet et al., 2001; Daish et al., 2005; Ioannidou et al., 2005). At low-elevation sites, monthly ^7Be averages are characterized by a well-defined annual cycle with lower values during winter and higher values during summer. Generally, the increase in ^7Be in ground level air from March to May is ascribed to the more efficient and higher frequency stratosphere–troposphere exchange (STE), whereas the further increase in ^7Be during summer is due to the stronger convective mixing and higher tropopause (Ioannidou et al., 2014). The higher tropopause height is associated with anticyclonic conditions, which results in downward transport from the upper troposphere and reduced wet scavenging during these conditions (Gerasopoulos et al., 2001, 2005; Ioannidou et al., 2014). In fact, compensating subsidence associated with convective mixing enhances downward transport of ^7Be from the upper troposphere (rather than direct input of stratospheric air) down to the lower troposphere and ground level (Zanis et al., 1999; Gerasopoulos et al., 2001, 2005; Ioannidou et al., 2005; Likuku, 2006; Steinmann et al., 2013).

High-elevation sites such as Jungfrauoch (Switzerland), Zugspitze (Germany), and Mt. Cimone (Italy), typically lying above the planetary boundary layer (PBL), are characterized by lower ^{210}Pb concentrations and higher ^7Be due to direct influences of air masses from the free troposphere (Zanis et al., 2000). The observed seasonal ^{210}Pb pattern at the high-altitude sites of Puy de Dôme (1465 m a.s.l., France) and Opme (660 m a.s.l., France) is characterized by maximum concentrations in spring and autumn and minimum concentrations in winter. This is due to higher radon emissions during the dry season (summer) than during the wet season (winter), and lower PBL height during winter (Bourcier et al., 2011). The latter results in weaker upward transport of ^{222}Rn and ^{210}Pb at high-altitude sites. Similar to low-elevation sites, higher ^7Be values are observed in summer due to convection-forced exchange with the upper troposphere and the higher tropopause height, which leads to more efficient vertical transport from the upper to lower troposphere (Reiter et al., 1983; Gerasopoulos et al., 2001;

Bourcier et al., 2011). At high-altitude sites a secondary maximum of ^7Be during cold months (December–March) is generally observed and attributed to the increase in stratosphere-to-troposphere events during this season (e.g., James et al., 2003; Stohl et al., 2003; Trickl et al., 2010). The higher frequency of rapid subsidence in winter at Northern Hemisphere midlatitudes can be ascribed to the intensity of baroclinic systems, which is greatest in the wintertime. In fact, well-developed tropopause folds and rapid deep intrusions are most likely to occur in the wake of intense cyclogenesis, usually limited to the wintertime storm track regions (James et al., 2003).

Numerical models have been used to analyze ^{210}Pb and ^7Be observations at high-elevation sites. One-dimensional model simulations of surface ^7Be have shown higher concentrations at high-elevation sites (Jasiulionis and Wershofen, 2005; Simon et al., 2009) but have also suggested that the diffusion of ^7Be was affected by the seasonal variation in meteorological conditions. Balkanski et al. (1993) examined the transport of ^{210}Pb in a global 3-D model and reported a weak decrease in ^{210}Pb concentrations between the continental mixed layer and the free troposphere: simulated concentrations at 6 km altitude were about 50 % of those in the continental mixed layer over much of the Northern Hemisphere in summer, and over large areas of the tropics year around, a result consistent with the few observations available for the free troposphere at that time (Moore et al., 1973). Rehfeld and Heimann (1995) compared the 3-D model simulated seasonal pattern of surface ^{210}Pb and ^7Be concentrations with the observations at several sites in both hemispheres. At Mauna Loa (19.47° N, 155.6° W; 3400 m a.s.l.; Hawaii) ^{210}Pb seasonality was characterized by high concentrations in spring and summer and lower ones in winter, as opposed to the seasonal pattern found at higher latitudes, where the ^{210}Pb maximum concentrations in winter are attributed to the advective transport of ^{210}Pb aerosols from midlatitudes. This behavior is due to the elevation of the site, representative of the conditions of the free troposphere rather than those of the PBL. As for ^7Be , the comparison between the model and the observations at Rexburg (43.8° N, 111.83° W; 1483 m a.s.l.; USA) showed systematically lower model values, due to the much higher precipitation rates in the model.

Previous studies have examined surface ^7Be observations at Mt. Cimone with respect to the role of STE in surface ozone increases (Bonasoni et al., 1999, 2000a, b; Cristofanelli et al., 2003, 2006, 2009a, 2015; Lee et al., 2007) within the framework of European projects such as VOTALP (Vertical Ozone Transport in the Alps) and STACCATO (influence of Stratosphere-Troposphere exchange in A Changing Climate on Atmospheric Transport and Oxidation capacity). These studies led to the assessment of a higher incidence of STE events during the period from October to February relative to the warm season, when thermal convection and the rising of the tropopause promote vertical mixing, which acts as a confounding factor in STE detection. Lee et

al. (2007) and Tositti et al. (2014) reported the seasonal patterns and frequency distributions of ^{210}Pb and ^7Be measured at Mt. Cimone, and highlighted higher concentrations of both radionuclides in the summertime due to the higher mixing height and horizontal transport by regional airflows. During winter, a general increase in ^7Be is associated with a decrease in ^{210}Pb , due to the dominant effect of STE and subsidence in the free troposphere. At the time of this work, no model analyses of ^{210}Pb and ^7Be observations at the site have been conducted.

In this paper, we conduct simulations of ^{210}Pb and ^7Be at Mt. Cimone with a state-of-the-art global 3-D chemistry and transport model (GMI CTM) driven by assimilated meteorological fields for the year of 2005. Our objectives are a better elucidation of the seasonal variations in ^{210}Pb and ^7Be concentrations and an improved understanding of the roles of transport and precipitation scavenging processes in their seasonalities at Mt. Cimone.

The remainder of this paper is organized as follows. Section 2 describes the measurement site, the radioactivity measurements at Mt. Cimone, and the GMI CTM. Section 3 evaluates the model performance in reproducing the observed wind and precipitation fields. Section 4 evaluates the seasonal ^{210}Pb and ^7Be concentrations in the model with those observed. Section 5 examines the sources and seasonal variations in the simulated radionuclide activities, followed by a summary and conclusions in Sect. 6.

2 Data and methods

2.1 Radionuclide measurements at Mt. Cimone

Mt. Cimone station (44°12' N, 10°42' E, 2165 m a.s.l.) is a global WMO-GAW station managed by the Meteorological Office of the Italian Air Force, which hosts the research platform “Ottavio Vittori” of the Institute of Atmospheric and Climate Science of the National Council of Research (ISAC-CNR). The station is located on top of the highest peak of the Italian Northern Apennines, with a 360° free horizon and an elevation such that the station lies above the PBL during most of the year: the Mt. Cimone measurements are considered representative of the southern Europe/Mediterranean free troposphere (Bonasoni et al., 2000a; Fischer et al., 2003; Cristofanelli et al., 2007), although during the warmer months an influence of PBL air can be detected due both to convective processes and to mountain/valley breeze regimes (Fischer et al., 2003; van Dingenen et al., 2005; Tositti et al., 2013). Note in this framework that southern Europe and the Mediterranean Basin are considered a hot-spot region in terms of both climate change (e.g., Forster et al., 2007) and air quality (Monks et al., 2009), as well as a major cross-road of different air mass transport processes (Li et al., 2001; Lelieveld et al., 2002; Millàn et al., 2006; Duncan et al., 2008; Tositti et al., 2013).

At Mt. Cimone station, ^{210}Pb , ^7Be , and aerosol mass load in the form of PM_{10} have been regularly measured in the period of 1998–2011 with a Thermo Environmental PM_{10} high-volume sampler. PM_{10} is sampled on rectangular glass fiber filters (Whatman, $20.3\text{ cm} \times 25.4\text{ cm}$, with an effective exposure area of about 407 cm^2), which were manually changed every 2–3 days, depending on weather conditions, failures of the sampling equipment, and/or of the power supply and personnel on site. The average flow rate was about $1.13\text{ m}^3\text{ min}^{-1}$ at standard temperature and pressure, with an average volume of air collected on each filter equal to 3000–4000 m^3 (about 48 h of sampling, 115–175 samples per year).

Airborne radionuclides travel attached to particulate matters and, as a consequence of their physical origin, tend to populate the fine fraction ($< 1.0\text{ }\mu\text{m}$) (Winkler et al., 1998; Gaffney et al., 2004). The PM_{10} samples were subjected to non-destructive high-resolution γ spectrometry for the determination of airborne radiotracers ^{210}Pb and ^7Be . The characteristics of the two hyper-pure germanium crystal detectors (HPGe) detectors are as follows: one p-type coaxial detector by Ortec/Ametek with a relative efficiency of 32.5 % and full width at half maximum (FWHM) 1.8 at 1332 keV and one planar DSG detector with an active surface of 1500 mm^2 and FWHM of 0.73 at 122 keV, for higher and lower energy ranges (100–2000 and 0–900 keV), respectively.

Spectra were accumulated for at least 1 day to optimize peak analysis and then processed with a specific software package (GammaVision-32, version 6.07, Ortec). Efficiency calibration was determined on both detectors with a blank glass fiber filter traced with accurately weighted aliquots of a standard solution of mixed radionuclides (QCY48, Amersham) supplemented with ^{210}Pb , homogeneously dispersed dropwise over the filter surface. Once dried under a hood under ambient conditions, the calibration filter was folded into a polystyrene container in the same geometry as the unknown samples. Quantitative analysis on samples was carried out by subtracting the spectrum of a blank filter in the same geometry, while uncertainty on peaks ($k = 1$, 68 % level of confidence) was calculated propagating the combined error over the efficiency fit previously determined with the counting error. Minimum detectable activity was calculated making use of the traditional ORTEC method (ORTEC, 2003) with a peak cut-off limit of 40 %. Activity data were corrected to the midpoint of the time interval of collection and for the decay during spectrum acquisition. For our analysis, we used monthly averages of ^{210}Pb and ^7Be data at Mt. Cimone in 2005.

2.2 GMI model

The Global Modeling Initiative (GMI, <http://gmi.gsfc.nasa.gov>) is a NASA-funded project aiming at improving assessments of anthropogenic perturbations to the Earth system; in this framework, a CTM appropriate for stratospheric assessments was developed (Rotman et al., 2001). It was firstly

used to evaluate the potential effects of stratospheric aircraft on the global stratosphere (Kinnison et al., 2001) and on the Antarctic lower stratosphere (Considine et al., 2000). The recent version of the GMI CTM includes a full treatment of both stratospheric and tropospheric photochemical and physical processes and is also capable of simulating atmospheric radionuclides ^{222}Rn , ^{210}Pb , ^7Be , and ^{10}Be throughout the troposphere and stratosphere (Considine et al., 2004, 2005; Rodriguez et al., 2004; Liu et al., 2016). Details of the model are described in Duncan et al. (2007, 2008), Strahan et al. (2007), and Considine et al. (2008).

In this work, we simulate ^{222}Rn , ^{210}Pb , ^7Be , and ^{10}Be using a version of the GMI model with the same basic structure as described by Considine et al. (2005) and Liu et al. (2016), including parameterizations of the important tropospheric physical processes such as convection, wet scavenging, dry deposition, and planetary boundary layer mixing. Meteorological data used to drive the CTM at 2° latitude by 2.5° longitude resolution, e.g., horizontal winds, convective mass fluxes, and precipitation fields, are the Modern-Era Retrospective analysis for Research and Applications (MERRA) assimilated data set from the NASA Global Modeling and Assimilation Office (GMAO) (Rienecker et al., 2011).

The flux-form semi-Lagrangian advection scheme and a convective transport algorithm from the CONVTRAN routine in NCAR CCM3 physics package are used in the model. The wet deposition scheme is that of Liu et al. (2001): it includes scavenging in wet convective updrafts, as well as first-order rainout and washout from both convective anvils and large-scale precipitations. The gravitational settling effect of cloud ice particles included in Liu et al. (2001) is not considered here. Dry deposition of aerosols is computed using the resistance-in-series approach. For the simulations of radionuclides, each simulation was run for 6 years, recycling the MERRA meteorological data for 2005, to equilibrate the lower stratosphere as well as the troposphere (Liu et al., 2001). The sixth-year output was used for analysis.

A uniform ^{222}Rn emission of $1.0\text{ atom cm}^{-2}\text{ s}^{-1}$ from land under nonfreezing conditions is assumed (Liu et al., 2001). Following Jacob and Prather (1990), the flux is reduced by a factor of 3 under freezing conditions. The flux from oceans and ice is null. Although a large variability in ^{222}Rn emission from land is observed, the above emission estimate is thought to be accurate to within 25 % globally (Turekian et al., 1977) and to within a factor of 2 regionally (Wilkening et al., 1975; Schery et al., 1989; Graustein and Turekian, 1990; Nazaroff, 1992; Liu et al., 2001).

Following Brost et al. (1991) and Koch et al. (1996), we used the Lal and Peters (1967) ^7Be source for 1958 (solar maximum year), as it best simulated stratospheric ^7Be concentrations measured from aircraft (Liu et al., 2001). The rates of ^7Be production reported more recently by Usoskin and Kovaltsov (2008) broadly agree with those of Lal and Peters (1967) with slightly (about 25 %) lower global production rate and will be tested in a separate model study. The

Lal and Peters (1967) source is represented as a function of latitude and altitude (pressure) and does not vary with season (see Fig. 1 of Koch et al., 1996). No interannual variability in the ^7Be source is considered in the model (Liu et al., 2001). This may lead to an underestimate of tropospheric ^7Be concentrations, especially at high latitudes during a solar minimum (or near minimum) year. Lal and Peters (1967) reported that the relative amplitude of the ^7Be production rate over a 11-year solar cycle is about 13 % below 300 hPa at latitudes above 45° .

Because of the coarse horizontal resolution of the model (2° latitude by 2.5° longitude), the model representation of the topography at the site is poor. The elevation of Mt. Cimone in the model is only 298 m, whereas in reality the mountain is 2165 m (a.s.l.) high (Fig. 1). For this reason, the model output was not sampled at ground level but instead at the grid box corresponding to the elevation of the site. In order to see the sensitivity of model–observation comparisons to spatial sampling, the model was sampled not only for the grid corresponding to the latitude and longitude of Mt. Cimone but also for the eight adjacent grids. To better understand the sources and seasonality of radiotracers in the model, we examine model output not only for ^{210}Pb , ^7Be , and their ratio ($^7\text{Be}/^{210}\text{Pb}$, an indicator of vertical transport; Koch et al., 1996), which can be directly compared to the measurements taken at Mt. Cimone, but also for other radiotracers and quantities, e.g., ^{222}Rn and $^{10}\text{Be}/^7\text{Be}$ (a STE tracer; Zanis et al., 2003).

Year 2005 was chosen for analysis because of the availability of the observational data and model output at the time of this work. As discussed later, the seasonal behavior of ^{210}Pb and ^7Be radionuclides during year 2005 was “typical” for Mt. Cimone. Monthly averages of ^{210}Pb and ^7Be data at Mt. Cimone were calculated for comparison with model results. To better compare the seasonalities of ^{210}Pb and ^7Be between the model and the observations, monthly percentage deviations from the annual mean concentration were also calculated.

3 Seasonal variations in transport and precipitation at Mt. Cimone: observations vs. model simulations

Mt. Cimone is the windiest meteorological station in Italy and the prevailing local winds blow from SSW and NNE directions (Ciattaglia, 1983; Ciattaglia et al., 1987; Colombo et al., 2000). The wind observations at Mt. Cimone during the period of 1998–2011, when radionuclide measurements were performed at the station (Tositti et al., 2014), agree with the climatology of local wind intensity and direction during the period of 1946–1999 as reported by the Italian Air Force (Colombo et al., 2000). NNE directions are more significant during the cold period, and fluxes from SW are more typical of the warm period. While winds blowing from the SSW sector generate a sea air inflow, a continental air inflow is ob-




Figure 1. Surface elevations (km) in the model. The white dot indicates the location of Mt. Cimone ($44^\circ 12' \text{N}$, $10^\circ 42' \text{E}$; 2165 m a.s.l.).

served when winds come from the NNE sector (Ciattaglia et al., 1987).

However, when considering the lifetimes of ^{210}Pb (about one week) and ^7Be (about 3 weeks) aerosols (Liu et al., 2001), it is apparent that regional and long-range transport has a much more important role than local transport. On a large scale, about 70 % of background air masses reaching Mt. Cimone in the period of 1996–1998 came from Atlantic and Arctic areas, with a smaller contribution from the Mediterranean Basin and the eastern area, as estimated by Bonasoni et al. (2000b). A more recent and extended study of advection patterns at Mt. Cimone (Brattich et al., 2017), analyzing clusters of 4-day kinematic back-trajectories calculated for the period of 1998–2011 with the HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) model driven by the NCEP/NCAR (National Centers for Environmental Prediction/National Center for Atmospheric Research) meteorological reanalysis, shows that the air masses advected to Mt. Cimone (55 %) arrive from the western Atlantic–North America sector, while the remaining air masses (from the Arctic, eastern Mediterranean, and Mediterranean Basin–northern Africa) together represent 45 % of trajectories. Seasonal transport to Mt. Cimone in the model is shown in Fig. 2, representing winds at the elevation of Mt. Cimone (winds are weaker at the model bottom layer). In agreement with the description of advection patterns at the site, prevailing model winds (Fig. 2) blow from the western Atlantic sector. Slow summer winds suggest the stronger influence of regional/local transport at Mt. Cimone during the period (e.g., Lee et al., 2007; Marinoni et al., 2008; Tositti et al., 2013, 2014; Brattich et al., 2015).




Figure 2. Simulated monthly mean (a) ^{210}Pb concentrations and (b) ^7Be concentrations, at the elevation of Mt. Cimone. Arrows represent the seasonality of winds in the MERRA meteorological data. The white dot indicates the location of Mt. Cimone ($44^{\circ}12' \text{ N}$, $10^{\circ}42' \text{ E}$; 2165 m a.s.l.).




Figure 3. Comparison of the MERRA total precipitation ($0\text{--}75^\circ\text{N}$, $90^\circ\text{W}\text{--}90^\circ\text{E}$) during January and July 2005 with that in the GPCP observations. The white dot indicates the location of Mt. Cimone ($44^\circ 12'\text{N}$, $10^\circ 42'\text{E}$; 2165 m a.s.l.).

In the model, Mt. Cimone appears to be in a location where there is a large horizontal gradient of wind (transport) during 2005. Long-range transport from western Europe, North America, and the Arctic region prevail during the cold period, while regional transport appears more important in summer. The model is able to capture relevant features of pressure systems and seasonal circulation patterns of the North Atlantic, Mediterranean, and African regions, such as the semi-permanent high-pressure system located in the North Atlantic with different positions during different seasons (Bermuda/Azores High), a semi-permanent system of high pressure centered in northeastern Siberia during the colder half of the year (Siberian High), and the Intertropical Convergence Zone (ITCZ) in the summer/autumn season. However, due to the coarse resolution of the global meteorological reanalysis that we use to construct the model winds, the more than 50 local-scale wind systems present in the Mediterranean and surrounding regions are not resolved (Burlando, 2009). In northern Europe, in fact, there are approximately two main states for the atmosphere, the westerly or zonal flows modulated by the advection of Atlantic lows, and the long-lived blocking anticyclonic configurations over North Sea or Scandinavia (easterly) (Burlando et al., 2008).

In the Mediterranean region, the main cyclones during winter are essentially sub-synoptic lows triggered by the major North Atlantic synoptic systems affected by the local topography of the northern Mediterranean coast (Trigo et al., 2002), whereas in summer cyclones develop because of thermal effects, orography (e.g., the Atlas Mountains), and increase in low-level thermal gradients (Trigo et al., 2002; Campins et al., 2006). Again, due to the coarse resolution of the meteorological data we use, these sub-synoptic

processes are not resolved. For instance, northern African lows and Sahara depressions (also referred to as Atlas lee depressions) and the resulting SSW wind (sirocco) (Reiter, 1975), potentially linked to ^{210}Pb variations at Mt. Cimone, appear to be an important feature missing in the degraded MERRA data, where they appear only during October/November. However, MERRA is able to capture the summertime north-northeasterly winds in the eastern Mediterranean (Aegean Sea), known as the Etesians. The Etesians are the most persistent localized wind system in the world as a result of a sharp east–west pressure gradient manifested by large-scale circulation features (i.e., low pressure over the eastern Mediterranean/Middle East and high pressure over central and southeastern Europe) (Dafka et al., 2016).

We evaluate the MERRA precipitation with that from the GPCP (Global Precipitation Climatology Project, <https://precip.gsfc.nasa.gov/>) satellite and surface observations in 2005. Figure 3 shows the MERRA and GPCP monthly precipitation for the region defined by $0\text{--}75^\circ\text{N}$ and $90^\circ\text{W}\text{--}90^\circ\text{E}$. Good agreement between the MERRA and the GPCP precipitations averaged over the region was found. In particular, summer precipitation patterns are very similar. The geographical distribution of precipitation in MERRA shows some important features in agreement with the observed climatology precipitations: the desert climate in North Africa, with very low precipitation all year long; the ITCZ, with high precipitation during the summer and autumn seasons; the North Atlantic region, with high precipitation especially during the winter and autumn seasons; and Europe, where the seasonal pattern of precipitation is similar to that in the North Atlantic region but precipitation is lower.




Figure 4. Comparison of the seasonal precipitation at Mt. Cimone in the MERRA meteorological data set with that in the GPCP observations for (a) the model grid box (“ij”) corresponding to the location of Mt. Cimone, (b) the model grid box (“ij – 1”) to the west of “ij”, (c) the model grid box (“i – 1j – 1”) to the southwest of “ij”, and (d) the model grid box (“i + 1j + 1”) to the northeast of “ij”.

Figure 4 shows the comparison of the GPCP and MERRA precipitation seasonality at Mt. Cimone. Since Mt. Cimone is located in a region with a large horizontal gradient in precipitation, we also show in the figure the comparisons for three adjacent grid boxes. The MERRA precipitation is generally lower than that of GPCP at two grid boxes (except for summer, Fig. 4a, b), but in good agreement at the other two grid boxes (Fig. 4c, d). The agreement between the MERRA and GPCP precipitation seasonality is reasonable, with the squared correlation coefficient R^2 varying between 0.56 (at the grid to the northwest of “ij”) and 0.89 (at the grid to the southeast of “ij”). Large differences between the MERRA precipitation and that locally observed at the station are instead present. While the daily mean observed 2005 precipitation is 0.81 mm, which is close to the corresponding precipitation (0.73 mm) in MERRA at the “ij” grid (i.e., a negative bias of -0.08 mm), the model bias is positive and much higher (0.31–1.28 mm) at adjacent grids. This bias may very well reflect again the fact that the observed surface precipitation is localized, whereas the satellite and MERRA precipitations correspond to a much larger scale (about 200 km). Moreover, as Colombo et al. (2000) previously pointed out, different from the surrounding area, where the climate is defined as temperate-continental, the climate at the mountain-top is classified as alpine because of the high elevation. In fact, in agreement with the GPCP precipitation in 2005, the observed climatology in the region shows a maximum during November (secondary maximum in spring) and an absolute minimum in July (secondary minimum in January),

whereas on the top of the mountain the precipitation is maximal during summer. The MERRA precipitation shows increased amounts during April and August–December, with minimum in June–July. As the local precipitation at the site is important to the scavenging of radionuclide aerosol tracers, this difference between the local and regional precipitation could contribute to any biases in our simulations. However, as we will show below, the ratio $^7\text{Be} / ^{210}\text{Pb}$ may cancel out the errors associated with precipitation scavenging (Koch et al., 1996).

Low ^{210}Pb concentrations are seen over the Atlantic Ocean, due to the negligible emissions of ^{222}Rn from the oceans and strong precipitation scavenging, and in northern and western Europe, especially during the cold season (Fig. 2a). High ^{210}Pb concentrations appear over the Sahara and northern Africa, as a result of low precipitation in this area, and also over the Middle East and southern Asia. ^{210}Pb concentrations over southern Europe appear higher during the transition seasons, especially autumn, and peak during summer when the minimum precipitation and slow winds from west are observed in the region. Low ^7Be concentrations are simulated along the equator where convective scavenging is strongest (Fig. 2b). High ^7Be concentrations are seen over the Sahara due to a combination of low precipitation and subsidence in this region. Elevated values also occur over the Middle East, North America, and Greenland. ^7Be concentrations over southern Europe appear higher during spring and peak during winter, when model winds are stronger and transport ^7Be aerosols from North America and




Figure 5. Comparison of GMI-simulated (black dotted line) monthly (a) ^{210}Pb and (b) ^7Be activities with those observed at Mt. Cimone (solid lines) in 2005. Also shown are GMI-simulated monthly activities of (c) ^{222}Rn , (d) $^{10}\text{Be}/^7\text{Be}$ ratios, and (e) stratospheric $^7\text{Be}/\text{total } ^7\text{Be}$ ratios. Model values are for the “ij” grid box corresponding to the location of Mt. Cimone. Vertical bars indicate the uncertainty in observed activities.

Greenland regions, where ^7Be production is highest (Beer et al., 2012).

4 Seasonal variations in ^{210}Pb and ^7Be at Mt. Cimone: observations vs. model simulations

The seasonality and frequency distributions of ^{210}Pb and ^7Be concentrations measured at the Mt. Cimone station were previously examined by Lee et al. (2007), while more recent analyses of the 12-year record were presented in Tositti et al. (2014) and Brattich et al. (2016). Generally, both radionuclides show a marked seasonal maximum in the summertime, a behavior shared by PM_{10} (Tositti et al., 2013) and O_3 (Bonasoni et al., 2000b). The ^{210}Pb summer maximum is mainly due to the higher mixing height and enhanced uplift from the boundary layer as a result of thermal convection. The seasonal fluctuation of ^7Be is more complex and characterized by two relative maxima, one during the cold season associated with stratosphere-to-troposphere transport, and the other during the warm season mainly associated

with tropospheric subsidence balancing lower-tropospheric air masses ascent occasionally accompanied by STE (Tositti et al., 2014). The ^{210}Pb and ^7Be measurements in 2005 are consistent with this description (Fig. 5): ^{210}Pb concentrations are characterized by two maxima during the warm period (July and September); ^7Be concentrations are characterized by one absolute maximum during summer (July) and one secondary maximum during spring (March).

Figure 5a and b compare the simulated monthly ^{210}Pb and ^7Be activities with the observations at Mt. Cimone in 2005. The comparisons for the monthly percentage deviations from the annual mean concentration are available in the Supplement (Figs. S1–S2). The seasonality of ^{210}Pb is well captured by the model. The model reproduces the presence of two seasonal maxima in the ^{210}Pb observations, with the maximum observed in July shifted to June in the simulation. The squared correlation coefficient R^2 between observed and simulated ^{210}Pb activities is equal to 0.83 at the “ij” grid and varies between 0.42 and 0.82 for adjacent grid boxes (to the north and to the west of “ij”, respectively), confirming the




Figure 6. Same as Fig. 5a and b but for the “ij – 1” to the south of “ij” (left column) and “i – 1j – 1” to the southwest of “ij” (right column) grids, respectively.

good performance of the model in reproducing the ^{210}Pb seasonal pattern.

As for ^7Be , the model well captures the March maximum (i.e., secondary maximum in the observations) and the month-to-month variation during the cold and transition seasons (January–April, October–December). However, during the warm period, the simulated ^7Be concentrations are lower by a factor of 2 than the observed. A better agreement was found at some adjacent model grid boxes (e.g., to the south and to the southwest of “ij”; Fig. 6 vs. Fig. 5). The correlation between observed and simulated monthly ^7Be activities also increases from $R^2 = 0.03$ at “ij” to $R^2 = 0.11$ – 0.60 at adjacent model grid boxes. The largest value of $R^2 = 0.6$ was obtained at the “ij – 1” grid box to the south of “ij” (Fig. 6). This improvement is due to the large horizontal gradient in the simulated ^7Be concentrations near the site (Fig. 2).

5 Sources and seasonality of ^{210}Pb and ^7Be at Mt. Cimone: a model analysis

In this section, we quantify the sources of ^{210}Pb and ^7Be and determine the processes governing their seasonality in the GMI model. Additional tracers as simulated by the model are used to aid in the interpretation. Model sensitivity experiments are conducted to examine the roles of transport and precipitation scavenging in the seasonality.

As discussed in Sect. 4, the model reproduces the ^{210}Pb seasonality well, with minimum in the cold period and maximum in the warm period. The ^{210}Pb seasonality (Fig. 5a) can be linked with the seasonal pattern of its precursor ^{222}Rn (Fig. 5c). It is seen that the summer ^{210}Pb maximum is due to stronger (thermal) convection, which uplifts more ^{222}Rn

out of the boundary layer (e.g., Lee et al., 2007; Tositti et al., 2014; Brattich et al., 2015). This uplift of ^{222}Rn from the boundary layer is minimum in the cold period, and the minimal level of ^{210}Pb in this period can be considered representative of the free troposphere. The ^{210}Pb summer increase appears to be associated with short-range and regional transport, as suggested by the model simulations (Fig. 2a). As expected, long-range transport is more typical of the winter/spring seasons because of stronger horizontal winds, while regional effects are more important during summer, when convection gets stronger.

In a similar manner, the source of the ^7Be March maximum can be investigated with model tracer simulations. Figure 5d and e also show the simulated seasonal patterns of the $^{10}\text{Be}/^7\text{Be}$ activity ratio and of the fraction of ^7Be originating from the stratosphere (strat $^7\text{Be}/\text{total } ^7\text{Be}$). The simulated seasonal pattern of the $^{10}\text{Be}/^7\text{Be}$ ratio is very similar to the observations at Jungfraujoch (Switzerland, 3580 m a.s.l.) (Zanis et al., 2003), characterized by a clear seasonal cycle with peak ratios in spring. The usefulness of $^{10}\text{Be}/^7\text{Be}$ ratio as a stratospheric tracer is due to the fact that both ^{10}Be and ^7Be cosmogenic radionuclides attach to the same aerosols and share therefore the same removal mechanism. Moreover, due to the much longer physical half-life of ^{10}Be ($\tau_{1/2} = 1.5 \times 10^6$ years) compared to ^7Be ($\tau_{1/2} = 53.3$ days), their concentration ratios in the stratosphere (about 3–4) are much higher than in the troposphere (about 2 or even less) (Koch and Rind, 1998). The simulated $^{10}\text{Be}/^7\text{Be}$ ratio behavior indicates that deep stratosphere-to-troposphere (STT) peaks during winter, while shallower STT has a spring maximum, consistent with previous analyses of stratospheric intrusions at Mt. Cimone (Cristofanelli et al., 2006, 2009a), and




Figure 7. Comparison between GMI-simulated monthly $^7\text{Be}/^{210}\text{Pb}$ ratios at the “ij” and “ij–1” grids (black dotted line) and those from the observations at Mt. Cimone (green solid line). Vertical bars indicate the uncertainty in observed activities.

more generally with the climatology of STE at the Northern Hemisphere midlatitudes (James et al., 2003). Altogether the simulated high strat $^7\text{Be}/\text{total } ^7\text{Be}$, high $^7\text{Be}/^{210}\text{Pb}$ (Fig. 7), and low $^{10}\text{Be}/^7\text{Be}$ ratios during December–January indicate strongest STE during this period, followed by spring with slightly weaker stratospheric influence on surface ^7Be . However, the model tends to overestimate the observed ^7Be concentrations and $^7\text{Be}/^{210}\text{Pb}$ ratios during December–February, suggesting that stratospheric influence and/or subsidence in the model is probably too strong in this region at this time of the year. It is noted that globally integrated STT mass fluxes in the MERRA reanalysis are actually smaller than in some other reanalyses, e.g., ERA-Interim, JRA-55, and MERRA-2 (Boothe and Homeyer, 2016).

The use of the ^7Be production rate of Lal and Peters (1967) for a solar maximum year (1958) may partly explain the lower annual mean ^7Be in the model (3.4 mBq m^{-3} annual mean at the “ij” grid) than in the observations (4.2 mBq m^{-3}). In fact, the sunspot number in 2005 (29.8) was quite low (slowly decreasing from 2000, a solar maximum year, and reaching minimum in 2008), especially compared to the 1958 value of 184.8. Sunspot number data are available from the World Data Center for the production, preservation and dissemination of the international sunspot number (Sunspot Index and Long-term Solar Observation, SILSO, Royal Observatory of Belgium, Brussels, <http://sidc.oma.be/sunspot-data/>, 2016).

During the winter period, associated with the simulated and observed ^7Be increases (Figs. 5–6), strong long-range




Figure 8. Comparison of GMI-simulated monthly ^{210}Pb and ^7Be activities at Mt. Cimone between the standard (black dotted line) and the sensitivity runs (“ij–1” grid). The sensitivity runs are those without convective transport/scavenging (red dotted line), without dry deposition (blue dotted line), and without scavenging (orange dotted line; y axis on the right). The observations are shown as a green solid line. Vertical bars indicate the uncertainty in observed activities.

transport was dominant in the European region (Fig. 2b). Transport from higher-latitude regions (Arctic, northern Europe, and North America) appears particularly important during this period (Fig. 2b); such transport from high-latitude regions, where the ^7Be production rate is highest (Beer et al., 2012), has typically been observed during STE events at Mt. Cimone in many studies (e.g., Bonasoni et al., 1999, 2000a, b).

The discrepancy between the simulated and the observed ^7Be concentrations during the warm period is partly due to the sensitivity to spatial sampling in the model. As seen from the map plots of ^{210}Pb and ^7Be concentrations at the elevation of Mt. Cimone (Fig. 2), the sampling site appears to be located in a region where the N–S gradient of concentrations is large (especially for ^7Be). An elevated gradient in




Figure 9. GMI-simulated differences of ^{210}Pb concentrations at the elevation of Mt. Cimone between a sensitivity run without convection (i.e., without transport and scavenging in convective updrafts) and the standard run. Arrows denote MERRA winds. The white dot indicates the location of Mt. Cimone ($44^{\circ}12' \text{ N}$, $10^{\circ}42' \text{ E}$; 2165 m a.s.l.).

the region surrounding Mt. Cimone was also seen for winds, as transport plays a critical role in determining the distributions of these tracers. The sensitivity to spatial sampling in the model is therefore ascribed to this observed strong gradient in the N–S direction. In fact, while the grids to the south and southwest of “ij” are better for summer ^7Be comparisons (Fig. 6), the grids to the northeast, north, and northwest of “ij” are better for winter (not shown).

The model underestimate of ^7Be levels in the warm months may also suggest the mixing of air masses between the PBL and the lower free troposphere is likely too weak. Previous observational analyses indicated that such mixing is higher in summer at Mt. Cimone due to enhanced convection and mountain wind breeze (e.g., Fischer et al., 2003; Cristofanelli et al., 2007). Weaker entrainment of free-tropospheric air into the PBL would result in lower ^7Be concentrations at the surface.

The model annual average biases are about 8 % for ^{210}Pb and about 19 % for ^7Be , respectively. By contrast, the model average bias for $^7\text{Be}/^{210}\text{Pb}$ ratios is about -13% (Fig. 7). The smaller model bias for $^7\text{Be}/^{210}\text{Pb}$ ratios than for ^7Be concentrations reflects the fact that the ratio cancels out the errors in precipitation scavenging (Koch et al., 1996) that contribute to the underestimate of ^{210}Pb and ^7Be activities. On the other hand, the negative model bias for the $^7\text{Be}/^{210}\text{Pb}$ ratio again points to weak downward mixing from the free troposphere.

If one compares the month-to-month variation in ^{210}Pb and ^7Be (Figs. 5 and 6) and precipitation in the model (Fig. 4), the maxima/minima of precipitation appear to be in phase with those of both radionuclides’ activities. This reflects the effects of precipitation scavenging on radionuclide aerosols.

We conducted model sensitivity experiments where either convection (transport and scavenging), wet scavenging due to both large-scale and convective precipitation, or dry deposition processes are turned off to examine the roles of these processes in controlling the seasonality of ^{210}Pb and ^7Be at Mt. Cimone. Figure 8 shows the results for the standard and sensitivity runs at the “grid to the south of “ij”, for which the simulated tracer seasonal variations are similar to those observed, while the monthly percentage deviations from the annual mean concentrations are shown in Fig. S3. Figures 9–12 show maps of simulated changes in ^{210}Pb and ^7Be concentrations when convection or wet scavenging is turned off.

Turning off dry deposition does not significantly change the simulated ^{210}Pb and ^7Be concentrations, partly due to sampling the higher vertical grid box in the model (larger effects are seen at the bottom model layer). With convection turned off (i.e., with neither convective transport nor convective scavenging), the simulated ^7Be seasonality also remains nearly the same. This suggests the compensating effects between subsidence (increasing ^7Be) associated with convective transport and scavenging (decreasing ^7Be) due to con-




Figure 10. Same as Fig. 9 but for a sensitivity simulation where wet scavenging is turned off.




Figure 11. GMI-simulated differences of ^7Be concentrations at the elevation of Mt. Cimone between a sensitivity run without convection and the standard run. Arrows denote MERRA winds. The white dot indicates the location of Mt. Cimone ($44^{\circ}12' \text{ N}$, $10^{\circ}42' \text{ E}$; 2165 m a.s.l.).




Figure 12. Same as Fig. 11 but for the difference between a sensitivity run without wet scavenging and the standard run.

vective precipitation. In the case of ^{210}Pb , turning off convection does not change the seasonal pattern but generally results in larger ^{210}Pb concentrations and particularly during summer/autumn, when convective transport is more important at the site. In fact, no convective transport of ^{222}Rn (Fig. S5) results in less ^{222}Rn (and ^{210}Pb) being transported to the free troposphere, as well as more ^{210}Pb being available in PBL lifted to the free troposphere by large-scale vertical transport; on the other hand, the lack of convective scavenging of ^{210}Pb increases its concentration in the free troposphere. Turning off convection therefore results in an increase in ^{210}Pb concentrations in the free troposphere. Both surface ^{222}Rn concentrations at the elevation of Mt. Cimone (Fig. S4), as well as a map of changes in ^{210}Pb concentrations due to convection in the model (Fig. 9) show that convection in the region is more important during summer and autumn but is not negligible during spring, possibly due to thermal inertia.

The model run without scavenging suggests that, apart from downward transport from the upper troposphere and lower stratosphere, wet scavenging is mainly responsible for the seasonal variation in ^7Be (Fig. 8, bottom panel). None of our simulations is able to describe the observed ^7Be summertime peak, suggesting that local and regional circulations in this region with complex topography may not be resolved by the coarse-resolution model. For ^{210}Pb (Fig. 8, top panel), it appears that wet scavenging plays a more important role during August–December than during January–

July. This appears to be associated with the seasonality of precipitation, which shows prolonged elevated values during August–December, as well as a maximum during April, as previously discussed (Fig. 5). A plot of changes in ^{210}Pb concentrations due to scavenging in the model (Fig. 10) confirms that the scavenging effect is larger during autumn and, to a lesser extent, during summer. At Mt. Cimone, the scavenging effect is not minimal during July (month of minimum precipitation, Fig. 4), suggesting the influence of precipitation scavenging elsewhere in the region on the site.

6 Summary and conclusions

We have used a global 3-D model (GMI CTM) driven by the MERRA assimilated meteorological data from NASA's GMAO to simulate the ^{210}Pb and ^7Be observations from the Mt. Cimone ($44^{\circ}12' \text{N}$, $10^{\circ}42' \text{E}$; 2165 m a.s.l.; Italy) WMO-GAW station in 2005. The two natural atmospheric radionuclides originate from contrasting source regions (lower troposphere and upper troposphere/lower stratosphere, respectively), attach to submicron particles, and are removed from the troposphere mainly by wet deposition. Our objective was to examine the roles of horizontal advection, vertical transport (large-scale and convection), and wet scavenging in determining the seasonality of ^{210}Pb and ^7Be at Mt. Cimone. The observed ^{210}Pb concentrations are characterized by maxima in summer and minima during the cold period. The seasonality of ^7Be is more complex, with a major peak

in summer, a secondary peak in spring and a minimum in winter. This is the first modeling study of ^{210}Pb and ^7Be observations at Mt. Cimone. This site is representative of free-tropospheric southern Europe/Mediterranean conditions most of the year, and thus the comparison between measurements and simulations can serve as an indication of shortcomings in the model or in the meteorological data.

Precipitation and wind fields are important to the model's performance in representing the transport and scavenging processes. We evaluated the MERRA precipitation field used by GMI CTM against the GPCP satellite and surface observations, and generally good agreement was found. The seasonality of precipitation at Mt. Cimone shows increased amounts during April and the period of August–December, and minimum in June–July. The MERRA assimilated winds at the low-resolution version we used captured the main circulation patterns (e.g., location of the Azores high-pressure system, location of the ITCZ) in the Northern Hemisphere. However, some local-scale winds and pressure systems, which are important for transport to the sampling site, were likely not well resolved at the coarse resolution we used. Generally good agreement was found between the MERRA assimilated wind fields and the main advection patterns at the site (e.g., prevalence of long-range transport from western Europe, North America, and the Arctic region during the cold season, as opposed to the prevailing regional transport during the warm season).

The model reproduced the observed ^{210}Pb seasonality well: ^{210}Pb maxima during the warm period were attributed to the stronger (thermal) convection, which uplifts more ^{222}Rn (and ^{210}Pb) from the boundary layer. The model is less successful in reproducing the observed ^7Be seasonality. ^7Be was better represented during the cold period, while the observed summer ^7Be maximum was underestimated by the model. The model underestimate of ^7Be levels in the warm months is partly due to the sensitivity to spatial sampling in the model, but also suggests that the mixing of air masses between the PBL and the lower free troposphere (e.g., via convection and compensating subsidence) is likely too weak during summer, when the Mt. Cimone station is located within the PBL. This suggests that additional work comparing the model results with more surface observations is needed in order to better understand this effect. The simulated lower annual average ^7Be concentration relative to the observation is also partly attributed to the fact that the model used the ^7Be production rate for a solar maximum year, while in 2005 (our simulation year) the solar activity was rather low.

By examining the wind fields and horizontal distribution of radiotracers in the model, we noted that the sampling site is in a location where there is a large gradient, especially in the north–south direction. Accordingly, we investigated the sensitivity of model results to spatial sampling. A better agreement between the model and the observations at some adjacent grid boxes was found. The ^7Be March maximum was linked to the large stratospheric influence during win-

ter/spring. The model tends to underestimate the summertime ^{210}Pb and ^7Be but better simulates the $^7\text{Be}/^{210}\text{Pb}$ ratio because the model errors due to precipitation scavenging appear to be canceled out in the ratio.

We have conducted a series of model sensitivity experiments to further examine and quantify the roles of wet scavenging, dry deposition, and convection (transport and scavenging) in controlling the seasonality of ^{210}Pb and ^7Be at Mt. Cimone. Dry deposition does not have a significant effect on the magnitude and seasonality of ^{210}Pb and ^7Be concentrations at the site. The relatively weak combined effects of convective transport and convective scavenging on the radiotracer seasonality were attributed to the compensating effects of convective transport and convective scavenging on tracer concentrations in the lower free troposphere (at the elevation of Mt. Cimone). Convection appears to be more important to the regional distribution of both radiotracers during summer and autumn, although it is also significant during spring. Finally, scavenging is found to be the most important process controlling the seasonal variations in ^{210}Pb and ^7Be at Mt. Cimone. For ^{210}Pb , precipitation plays a more important role during August–December than during January–July. This was attributed to the seasonality of local and regional precipitation, which shows prolonged elevated values in the period of August–December.

While our simulations demonstrated some capabilities of the model to reproduce the seasonality of ^{210}Pb and ^7Be , they highlight the weaknesses of the model in reproducing local features, presumably due to its coarse resolution. Model simulations at a higher resolution would improve this model analysis of ^{210}Pb and ^7Be observations at Mt. Cimone, a high-elevation site. The understanding of downward transport associated with convection during summer also requires improving. Therefore, ^{210}Pb and ^7Be tracers will prove to be very useful in our understanding of seasonal behaviors of other environmentally important trace gases and aerosols at Mt. Cimone. Since other aerosols and trace gases (e.g., black carbon, CO , O_3) are also measured at the station, we plan to conduct comparisons between model simulations and those measurements to corroborate or contrast with the radionuclide results.

7 Data availability

A description of the observational data and model output used in this paper can be found in Sect. 2 and they are available upon request by contacting Laura Tositti (laura.tositti@unibo.it) and Hongyu Liu (hongyu.liu-1@nasa.gov), respectively.

The Supplement related to this article is available online at doi:10.5194/acp-17-1061-2017-supplement.

Acknowledgements. Italian Air Force Meteorological Office (IAFMS) and ISAC-CNR are gratefully acknowledged for their precious technical support at the Mt. Cimone station. In particular, ISAC-CNR is gratefully acknowledged for providing infrastructural access at the WMO-GAW Global Station Italian Climate Observatory “O. Vittori” at Mt. Cimone. IAFMS is gratefully acknowledged for providing meteorological observations at Mt. Cimone station. The Italian Climate Observatory “O. Vittori” is supported by MIUR and DTA-CNR throughout the Project of National Interest “NextData”. Erika Brattich thanks the National Institute of Aerospace (NIA) Visitor Program for hosting her one-month visit, and the Department of Biological, Geological and Earth Sciences of the University of Bologna for grant support during her PhD study. Hongyu Liu is supported by NASA Modeling and Analysis Program (MAP), NASA Atmospheric Composition Modeling and Analysis Program (ACMAP), and NASA Atmospheric Composition Campaign Data Analysis and Modeling (ACCDAM) program. The GMI activity is managed by José Rodriguez and Susan Strahan (NASA GSFC). Stephen Steenrod, Megan Damon, and Jules Kouatchou (GSFC) are acknowledged for programming support. NASA Center for Computational Sciences (NCCS) provided supercomputing resources. We thank the two anonymous reviewers for their comments, which improved the quality of our work.

Edited by: J. Ma

Reviewed by: two anonymous referees

References

- Arimoto, R., Snow, J. A., Graustein, W. C., Moody, J. L., Ray, B. J., Duce, R. A., Turekian, K. K., and Maring, H. B.: Influences of atmospheric transport pathways on radionuclide activities in aerosol particles from over the North Atlantic, *J. Geophys. Res.*, 104, 301–321, 1999.
- Balkanski, Y., Jacob, D. J., Gardner, G. M., Graustein, W., and Turekian, K. K.: Transport and residence times of tropospheric aerosols inferred from a global three-dimensional simulation of ^{210}Pb , *J. Geophys. Res.*, 98, 20573–20586, 1993.
- Baskaran, M.: Po-210 and Pb-210 as atmospheric tracers and global atmospheric Pb-210 fallout: a review, *J. Environ. Radioactiv.*, 102, 500–513, 2011.
- Beer, J., McCracken, K., and von Steiger, R.: *Cosmogenic radionuclides*, Springer, Heidelberg, Germany, 2012.
- Bonasoni, P., Evangelisti, F., Bonafé, U., Feldmann, H., Memmesheimer, M., Stohl, A., and Tositti, L.: Stratosphere-troposphere exchanges: case studies recorded at Mt. Cimone during VOTALP project, *Phys. Chem. Earth C*, 24, 443–446, 1999.
- Bonasoni, P., Evangelisti, F., Bonafé, U., Ravegnani, F., Calzolari, F., Stohl, A., Tositti, L., Tubertini, O., and Colombo, T.: Stratospheric ozone intrusion episodes recorded at Mt. Cimone during VOTALP project: Case studies, *Atmos. Environ.*, 34, 1355–1365, 2000a.
- Bonasoni, P., Stohl, A., Cristofanelli, P., Calzolari, F., Colombo, T., and Evangelisti, F.: Background ozone variations at Mt Cimone station, *Atmos. Environ.*, 34, 5183–5189, 2000b.
- Boothe, A. C. and Homeyer, C. R.: Global large-scale stratosphere-troposphere exchange in modern reanalyses, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-788, in review, 2016.
- Bourcier, L., Masson, O., Laj, P., Pichon, J. M., Paulat, P., Freney, E., and Sellegri, K.: Comparative trends and seasonal variation of ^7Be , ^{210}Pb and ^{137}Cs at two altitude sites in the central part of France, *J. Environ. Radioactiv.*, 102, 294–301, 2011.
- Brattich, E., Hernández-Ceballos, M. A., Cinelli, G., and Tositti, L.: Analysis of peak ^{210}Pb values at Mt. Cimone (1998–2011), *Atmos. Environ.*, 112, 136–147, 2015.
- Brattich, E., Hernández-Ceballos, M. A., Orza, J. A. G., Bolívar, J. P., and Tositti, L.: The western Mediterranean basin as an aged aerosols reservoir: Insights from an old-fashioned but efficient radiotracer, *Atmos. Environ.*, 141, 481–493, 2016.
- Brattich, E., Orza, J. A. G., and Tositti, L.: Advection patterns at the WMO-GAW station of Mt. Cimone: seasonality, trends, and influence on atmospheric composition, in preparation, 2017.
- Brost, R. A., Feichter, J., and Heimann, H.: Three-dimensional simulation of ^7Be in a global climate model, *J. Geophys. Res.*, 96, 22423–22445, 1991.
- Burlando, M.: The synoptic-scale surface wind climate regimes of the Mediterranean Sea according to the cluster analysis of ERA-40 wind fields, *Theor. Appl. Climatol.*, 96, 69–83, 2009.
- Burlando, M., Antonelli, M., and Ratto, C. F.: Mesoscale wind climate analysis: identification of anemological regions and wind regimes, *Int. J. Climatol.*, 28, 629–641, 2008.
- Caillet, S., Arpagaus, P., Monna, F., and Dominik, J.: Factors controlling ^7Be and ^{210}Pb atmospheric deposition as revealed by sampling individual rain events in the region of Geneva, Switzerland, *J. Environ. Radioactiv.*, 53, 241–256, 2001.
- Campins, J., Jansà, A., and Genovés, A.: Three-dimensional structure of western Mediterranean cyclones, *Int. J. Climatol.*, 26, 323–343, 2006.
- Cannizzaro, F., Greco, G., Raneli, M., Spitale, M. C., and Tomarchio, E.: Concentration measurements of ^7Be at ground level air at Palermo, Italy – comparison with solar activity over a period of 21 years, *J. Environ. Radioactiv.*, 84, 457–467, 2004.
- Carvalho, A. C., Reis, M., Silva, L., and Madruga, M. J.: A decade of ^7Be and ^{210}Pb activity in surface aerosols measured over the Western Iberian Peninsula, *Atmos. Environ.*, 67, 193–202, 2013.
- Ciattaglia, L.: Interpretation of atmospheric CO_2 measurements at Mt. Cimone (Italy) related to wind data, *J. Geophys. Res.*, 88, 1331–1338, 1983.
- Ciattaglia, L., Cundari, V., and Colombo, T.: Further measurements of atmospheric carbon dioxide at Mt. Cimone, Italy, 1979–1985, *Tellus B*, 39, 13–20, 1987.
- Colombo, T., Santaguida, R., Capasso, A., Calzolari, F., Evangelisti, F., and Bonasoni, P.: Biospheric influence on carbon dioxide measurements in Italy, *Atmos. Environ.*, 34, 4963–4969, 2000.
- Considine, D. B., Douglass, A. R., Connell, P. S., Kinnison, D. E., and Rotman, D. A.: A polar stratospheric cloud parameterization for the global modeling initiative three-dimensional model and its response to stratospheric aircraft, *J. Geophys. Res.*, 105, 3955–3973, 2000.
- Considine, D. B., Connell, P. S., and Logan, J. A.: Simulating ozone in the near tropopause region with a new combined model of the stratosphere and troposphere, in: *Quadrennial Ozone Symposium QOS 2004*, edited by: Zerefos, C., International Ozone Commission, Kos, Greece, 739–740, 2004.

- Considine, D. B., Bergmann, D. J., and Liu, H.: Sensitivity of Global Modeling Initiative chemistry and transport model simulations of radon-222 and lead-210 to input meteorological data, *Atmos. Chem. Phys.*, 5, 3389–3406, doi:10.5194/acp-5-3389-2005, 2005.
- Considine, D. B., Logan, J. A., and Olsen, M. A.: Evaluation of near-tropopause ozone distributions in the Global Modeling Initiative combined stratosphere/troposphere model with ozonesonde data, *Atmos. Chem. Phys.*, 8, 2365–2385, doi:10.5194/acp-8-2365-2008, 2008.
- Cristofanelli, P., Bonasoni, P., Collins, W., Feichter, J., Forster, C., James, P., Kentarchos, A., Kubik, P.W., Land, C., Meloen, J., Roelofs, G.J., Siegmund, P., Sprenger, M., Schnabel, C., Stohl, A., Tobler, L., Tositti, L., Trickl, T., and Zanis, P.: Stratosphere-to-troposphere transport: A model and method evaluation, *J. Geophys. Res.*, 108, 8525, doi:10.1029/2002JD002600, 2003.
- Cristofanelli, P., Bonasoni, P., Tositti, L., Bonafé, U., Calzolari, F., Evangelisti, F., Sandrini, S., and Stohl, A.: A 6-year analysis of stratospheric intrusions and their influence on ozone at Mt. Cimone (2165 m above sea level), *J. Geophys. Res.*, 111, D03306, doi:10.1029/2005JD006553, 2006.
- Cristofanelli, P., Bonasoni, P., Carboni, G., Calzolari, F., Casarola, L., Zauli Sajani, S., and Santaguida, R.: Anomalous high ozone concentrations recorded at a high mountain station in Italy in summer 2003, *Atmos. Environ.*, 41, 1383–1394, 2007.
- Cristofanelli, P., Calzolari, F., Bonafé, U., Duchi, R., Marinoni, A., Roccato, F., Tositti, L., and Bonasoni P.: Stratospheric intrusion index (SI^2) from baseline measurement data, *Theor. Appl. Climatol.*, 97, 317–325, 2009a.
- Cristofanelli, P., Marinoni, A., Arduini, J., Bonafè, U., Calzolari, F., Colombo, T., Decesari, S., Duchi, R., Facchini, M. C., Fierli, F., Finessi, E., Maione, M., Chiari, M., Calzolari, G., Messina, P., Orlandi, E., Roccato, F., and Bonasoni, P.: Significant variations of trace gas composition and aerosol properties at Mt. Cimone during air mass transport from North Africa – contributions from wildfire emissions and mineral dust, *Atmos. Chem. Phys.*, 9, 4603–4619, doi:10.5194/acp-9-4603-2009, 2009b.
- Cristofanelli, P., Scheel, H.-E., Steinbacher, M., Saliba, M., Azopardi, F., Ellul, R., Fröhlich, M., Tositti, L., Brattich, E., Maione, M., Calzolari, F., Duchi, R., Landi, T. C., Marinoni, A., and Bonasoni, P.: Long-term surface ozone variability at Mt. Cimone WMO/GAW global station (2165 m a.s.l., Italy), *Atmos. Environ.*, 101, 23–33, 2015.
- Cuevas, E., González, Y., Rodríguez, S., Guerra, J. C., Gómez-Peláez, A. J., Alonso-Pérez, S., Bustos, J., and Milford, C.: Assessment of atmospheric processes driving ozone variations in the subtropical North Atlantic free troposphere, *Atmos. Chem. Phys.*, 13, 1973–1998, doi:10.5194/acp-13-1973-2013, 2013.
- Dafka, S., Xoplaki, E., Toreti, A., Zanis, P., Tyrlis, E., Zerefos, C., and Luterbacher, J.: The Etesians; from observations to reanalysis, *Clim. Dynam.*, 47, 1569–1585, doi:10.1007/s00382-015-2920-7, 2016.
- Daish, S. R., Dale, A. A., Dale, C. J., May, R., and Rowe, J. E.: The temporal variations of ^7Be , ^{210}Pb and ^{210}Po in England, *J. Environ. Radioactiv.*, 84, 457–467, 2005.
- Dibb, J. E.: Vertical mixing above Summit, Greenland: insights into seasonal and high frequency variability from the radionuclide tracers ^7Be and ^{210}Pb , *Atmos. Environ.*, 41, 5020–5030, 2007.
- Dibb, J. E., Talbot, R. W., and Gregory, G. L.: Beryllium 7 and lead 210 in the western hemisphere Arctic atmosphere: Observations from three recent aircraft-based sampling programs, *J. Geophys. Res.*, 97, 16709–16715, 1992.
- Dibb, J. E., Meeker, L. D., Finkel, R. C., Southon, J. R., Caffee, M. W., and Barrie, L. A.: Estimation of stratospheric input to the Arctic troposphere: ^7Be and ^{10}Be in aerosols at Alert, Canada, *J. Geophys. Res.*, 99, 12855–12864, 1994.
- Dueñas, C., Fernández, M. C., Cañete, S., and Pérez, M.: ^7Be to ^{210}Pb concentration ratio in ground level air in Málaga (36.7° N, 4.5° W), *Atmos. Res.*, 92, 49–57, 2009.
- Dueñas, C., Orza, J. A. G., Cabello, M., Fernández, M. C., Cañete, S., Pérez, M., and Gordo, E.: Air mass origin and its influence on radionuclide activities (^7Be and ^{210}Pb) in aerosol particles at a coastal site in the western Mediterranean, *Atmos. Res.*, 101, 205–214, 2011.
- Duncan, B. N., Strahan, S. E., Yoshida, Y., Steenrod, S. D., and Livesey, N.: Model study of the cross-tropopause transport of biomass burning pollution, *Atmos. Chem. Phys.*, 7, 3713–3736, doi:10.5194/acp-7-3713-2007, 2007.
- Duncan, B. N., West, J. J., Yoshida, Y., Fiore, A. M., and Ziemke, J. R.: The influence of European pollution on ozone in the Near East and northern Africa, *Atmos. Chem. Phys.*, 8, 2267–2283, doi:10.5194/acp-8-2267-2008, 2008.
- Feely, H. W., Larsen, R. J., and Sanderson, C. G.: Factors that cause seasonal variations in beryllium-7 concentrations in surface air, *J. Environ. Radioactiv.*, 9, 223–249, 1989.
- Feichter, J., Brost, R. A., and Heimann, M.: Three-dimensional modeling of the concentration and deposition of ^{210}Pb aerosols, *J. Geophys. Res.*, 96, 22447–22469, 1991.
- Fischer, H., Kormann, R., Klüpfel, T., Gurk, Ch., Königstedt, R., Parchatka, U., Mühle, J., Rhee, T. S., Brenninkmeijer, C. A. M., Bonasoni, P., and Stohl, A.: Ozone production and trace gas correlations during the June 2000 MINATROC intensive measurement campaign at Mt. Cimone, *Atmos. Chem. Phys.*, 3, 725–738, doi:10.5194/acp-3-725-2003, 2003.
- Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Changes in Atmospheric Constituents and in Radiative Forcing, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.
- Froehlich, K. and Masarik, J.: Radionuclides as tracers and timers of processes in the continental environment – Basic concepts and methodologies, in: *Radioactivity in the Environment*, 16, Chapter 2, 27–50, *Environmental Radionuclides: Tracers and Timers of Terrestrial Processes*, Elsevier, doi:10.1016/S1569-4860(09)01602-7, 2010.
- Gaffney, J. S., Marley, N., and Cunningham, M. M.: Natural radionuclides in fine aerosols in the Pittsburgh area, *Atmos. Environ.*, 38, 3191–3200, 2004.
- Gäggeler, H. W.: Radioactivity in the atmosphere, *Radiochim. Acta*, 70–71, 345–353, 1995.
- Gerasopoulos, E., Zanis, P., Stohl, A., Zerefos, C. S., Papastefanou, C., Ringer, W., Tobler, L., Hübener, S., Gäggeler, H. W., Kanter,

- H. J., Tositti, L., and Sandrini, S.: A climatology of ^7Be at four high-altitude stations at the Alps and the Northern Apennines, *Atmos. Environ.*, 35, 6347–6360, 2001.
- Gerasopoulos, E., Zanis, P., Zerefos, C. S., Papastefanou, C., Ringer, W., Gäggeler, H. W., Tobler, L., and Kanter, H. J.: Factors and processes controlling the concentration of the cosmogenic radionuclide ^7Be at high-altitude Alpine stations, in: *Radioactivity in the Environment*, Vol. 7, 863–870, Elsevier Ltd., doi:10.1016/S1569-4860(04)07108-6, 2005.
- Graustein, W. C. and Turekian, K. K.: Radon fluxes from soils to the atmosphere measured by ^{210}Pb - ^{226}Ra disequilibrium in soils, *Geophys. Res. Lett.*, 17, 841–844, 1990.
- Graustein, W. C. and Turekian, K. K.: ^7Be and ^{210}Pb indicate an upper troposphere source for elevated ozone in the summertime subtropical free troposphere of the eastern North Atlantic, *Geophys. Res. Lett.*, 23, 539–542, 1996.
- Hötzl, H. and Winkler, R.: Activity concentrations of ^{226}Ra , ^{228}Ra , ^{210}Pb , ^{40}K and ^7Be and their temporal variations in surface air, *J. Environ. Radioactiv.*, 5, 445–458, 1987.
- Ioannidou, A., Manolopoulou, M., and Papastefanou, C.: Temporal changes of ^7Be and ^{210}Pb concentrations in surface air at temperate latitudes (40°), *Appl. Radiat. Isotopes*, 63, 277–284, 2005.
- Ioannidou, A., Vasileiadis, A., and Melas, D.: Time lag between the tropopause height and ^7Be activity concentrations in surface air, *J. Environ. Radioactiv.*, 129, 80–85, 2014.
- Jacob, D. J. and Prather, M. J.: Radon-222 as a test of boundary layer convection in a general circulation model, *Tellus B*, 42, 118–134, 1990.
- James, P., Stohl, A., Forster, C., Eckhardt, S., Seibert, P., and Frank, A.: A 15-year climatology of stratosphere-troposphere exchange with a Lagrangian particle dispersion model 2. Mean climate and seasonal variability, *J. Geophys. Res.*, 108, 8522, doi:10.1029/2002JD002639, 2003.
- Jasiulionis, R. and Wershofen, H.: A study of the vertical diffusion of the cosmogenic radionuclides, ^7Be and ^{22}Na in the atmosphere, *J. Environ. Radioactiv.*, 79, 157–169, 2005.
- Johnson, W. and Viezee, W.: Stratospheric ozone in the lower troposphere: i. Presentation and interpretation of aircraft measurements, *Atmos. Environ.*, 15, 1309–1323, 1981.
- Junge, C. E.: *Air chemistry and radioactivity*, Academic Press, New York, USA, and London, UK, 382 pp., 1963.
- Kinnison, D. E., Connell, P. S., Rodriguez, J. M., Rotman, D. A., Considine, D. B., Tannahill, J., Ramarosan, R., Rasch, P. J., Douglass, A. R., Baughcum, S. L., Coy, L., Waugh, D. W., Kawa, S. R., and Prather, M. J.: The Global Modeling Initiative assessment model: Application to high-speed civil transport perturbation, *J. Geophys. Res.*, 106, 1693–1711, 2001.
- Koch, D. and Rind, D.: Beryllium10/beryllium7 as a tracer of stratospheric transport, *J. Geophys. Res.*, 103, 3907–3917, 1998.
- Koch, D. M., Jacob, D. J., and Graustein, W. C.: Vertical transport of tropospheric aerosols as indicated by ^7Be and ^{210}Pb in a chemical tracer model, *J. Geophys. Res.*, 101, 18651–18666, 1996.
- Kulan, A., Aldahan, A., Possnert, G., and Vintersved, I.: Distribution of ^7Be in surface air of Europe, *Atmos. Environ.*, 40, 3855–3868, 2006.
- Lal, D. and Peters, B.: Cosmic ray produced radioactivity on the Earth, in: *Handbuch der Physik*, 46/2, edited by: Sitte, K., Springer-Verlag, New York, USA, 551–561, 1967.
- Lal, D., Malhotra, P. K., and Peters, B.: On the production of radioisotopes in the atmosphere by cosmic radiation and their application to meteorology, *J. Atmos. Terr. Phys.*, 12, 306–328, 1958.
- Lee, H. N., Wan, G., Zheng, X., Sanderson, C. G., Josse, B., Wang, S., Yang, W., Tang, J., and Wang, C.: Measurements of ^{210}Pb and ^7Be in China and their analysis accompanied with global model calculations of ^{210}Pb , *J. Geophys. Res.*, 109, D22203, doi:10.1029/2004JD005061, 2004.
- Lee, H. N., Tositti, L., Zheng, X., and Bonasoni, P.: Analyses and comparisons of variations of ^7Be , ^{210}Pb and $^7\text{Be}/^{210}\text{Pb}$ with ozone observations at two Global Atmosphere Watch stations from high mountains, *J. Geophys. Res.*, 112, D05303, doi:10.1029/2006JD007421, 2007.
- Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J., Heland, J., Holzinger, R., Korrmann, R., Lawrence, M. G., Levin, Z., Markowicz, K. M., Mihopoulos, N., Minikin, A., Ramanathan, V., de Reus, M., Roelofs, G. J., Scheeren, H. A., Sciare, J., Schlager, H., Schultz, M., Siegmund, P., Steil, B., Stephanou, E. G., Stier, P., Traub, M., Warneke, C., Williams, J., and Ziereis, H.: Global air pollution crossroads over the Mediterranean, *Science*, 298, 794–799, 2002.
- Li, Q., Jacob, D. J., Logan, J. A., Bey, I., Yantosca, R. M., Liu, H., Martin, R. V., Fiore, A. M., Field, B. D., Duncan, B. N.: A Tropospheric Ozone Maximum Over the Middle East, *Geophys. Res. Lett.*, 28, 3235–3238, 2001.
- Likuku, A. S.: Factors influencing ambient concentrations of ^{210}Pb and ^7Be over the city of Edinburgh (55.9°N , 03.2°W), *J. Environ. Radioactiv.*, 87, 289–304, 2006.
- Liu, H., Jacob, D. J., Bey, I., and Yantosca, R. M.: Constraints from the ^{210}Pb and ^7Be on wet deposition and transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, *J. Geophys. Res.*, 106, 12109–12128, 2001.
- Liu, H., Jacob, D. J., Dibb, J. E., Fiore, A. M., and Yantosca, R. M.: Constraints on the sources of tropospheric ozone from ^{210}Pb - ^7Be - O_3 correlations, *J. Geophys. Res.*, 109, D07306, doi:10.1029/2003JD003988, 2004.
- Liu, H., Considine, D. B., Horowitz, L. W., Crawford, J. H., Rodriguez, J. M., Strahan, S. E., Damon, M. R., Steenrod, S. D., Xu, X., Kouatchou, J., Carouge, C., and Yantosca, R. M.: Using beryllium-7 to assess cross-tropopause transport in global models, *Atmos. Chem. Phys.*, 16, 4641–4659, doi:10.5194/acp-16-4641-2016, 2016.
- Lozano, R. L., Hernández-Ceballos, M. A., San Miguel, E. G., Adame, J. A., and Bolívar, J. P.: Meteorological factors influencing the ^7Be and ^{210}Pb concentrations in surface air from the southwestern Iberian Peninsula, *Atmos. Environ.*, 63, 168–178, 2012.
- Lozano, R. L., Hernández-Ceballos, M. A., Rodrigo, J. F., San Miguel, E. G., Casas-Ruiz, M., García-Tenorio, R., and Bolívar, J. P.: Mesoscale behavior of ^7Be and ^{210}Pb in superficial air along the Gulf of Cadiz (south of Iberian peninsula), *Atmos. Environ.*, 80, 75–84, 2013.
- Marinoni, A., Cristofanelli, P., Calzolari, F., Roccató, F., Bonafé, U., and Bonasoni, P.: Continuous measurements of aerosol physical parameters at the Mt. Cimone GAW Station (2165 m asl, Italy), *Sci. Total Environ.*, 391, 241–251, 2008.

- Millàn, M., Sanz, J., Salvador, R., and Mantilla, E.: Atmospheric dynamics and ozone cycles related to nitrogen deposition in the western Mediterranean, *Environ. Pollut.*, 118, 167–186, 2006.
- Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Aki-moto, H., Amann, M., Baklanov, A., Baltensperger, U., Bey, I., Blake, N., Blake, R. S., Carslaw, K., Cooper, O. R., Dentener, F., Fowler, D., Fragkou, E., Frost, G. J., Generoso, S., Ginoux, P., Grewe, V., Guenther, A., Hansson, H. C., Henne, S., Hjorth, J., Hofzumahaus, A., Huntrieser, H., Isaksen, I. S. A., Jenkin, M. E., Kaiser, J., Kanakidou, M., Klimont, Z., Kulmala, M., Laj, P., Lawrence, M. G., Lee, J. D., and Liousse, C.: Atmospheric composition change – global and regional air quality, *Atmos. Environ.*, 43, 5268–5350, 2009.
- Moore, H. E., Poet, S. E., and Martell, E. A.: ^{222}Rn , ^{210}Pb , ^{210}Bi , and ^{210}Po , profiles and aerosol residence times versus altitude, *J. Geophys. Res.*, 78, 7065–7075, 1973.
- Nazaroff, W. W.: Radon transport from soil to air, *Rev. Geophys.*, 30, 137–160, 1992.
- ORTEC: Gamma-Vision 32 A66-B32 user's manual, ORTEC USA, Part No. 783620, Manual Revision D, 2003.
- Papastefanou, C. and Ioannidou, A.: Aerodynamic size association of ^7Be in ambient aerosols, *J. Environ. Radioactiv.*, 26, 273–282, 1995.
- Pham, M. K., Betti, M., Nies, H., and Povinec, P. P.: Temporal changes of ^7Be , ^{137}Cs and ^{210}Pb activity concentrations in surface air at Monaco and their correlation with meteorological parameters, *J. Environ. Radioactiv.*, 102, 1045–1054, 2011.
- Rastogi, N. and Sarin, M. M.: Atmospheric ^{210}Pb and ^7Be in ambient aerosols over low- and high-altitude sites in 34 semiarid region: Temporal variability and transport processes, *J. Geophys. Res.*, 113, D11103, doi:10.1029/2007JD009298, 2008.
- Rehfeld, S. and Heimann, M.: Three dimensional atmospheric transport simulation of the radioactive tracers ^{210}Pb , ^7Be , ^{10}Be , and ^{90}Sr , *J. Geophys. Res.*, 100, 26141–26161, 1995.
- Reiter, E. R.: Weather phenomena of the Mediterranean basin. Part 1. General description of the meteorological processes, in: Handbook for forecasters in the Mediterranean basin, Environment Prediction Research Facility, Naval Postgraduate School, Monterey, California, U.S. Department of Commerce, available at: <http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA024271> (last access: 15 March 2016), 1975.
- Reiter, R., Sladkovich, K., Pötzl, K., Carnuth, W., and Kanter, H. J.: Studies on the influx of stratospheric air into the lower troposphere using cosmic-ray produced radionuclides and fallout, *Arch. Meteor. Geophys. A*, 20, 211–246, 1971.
- Reiter, R., Munzert, K., Kanter, H.-J., and Pötzl, K.: Cosmogenic radionuclides and ozone at a mountain station at 3.0 km a.s.l., *Arch. Meteor. Geophys. B*, 32, 131–160, 1983.
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648, 2011.
- Rodriguez, J. M., Logan, J. A., Bergmann, D., Megretskaja, I., Jacob, D. J., Xie, H., Das, B., and Strahan, S. E.: The impact of meteorological fields on tropospheric ozone distributions calculated by the Global Modeling Initiative (GMI) chemical-transport model, in: Quadrennial Ozone Symposium QOS 2004, edited by: Zerefos, C., 147 pp., International Ozone Commission, Kos, Greece, 2004.
- Rotman, D. A., Tannahill, J. R., Kinnison, D. E., Connell, P. S., Bergmann, D., Proctor, D., Rodriguez, J. M., Lin, S. J., Rood, R. B., Prather, M. J., Rasch, P. J., Considine, D. B., Ramaroson, R., and Kawa, S. R.: Global Modeling Initiative assessment model: Model description, integration, and testing of the transport shell, *J. Geophys. Res.*, 106, 1669–1691, 2001.
- Schery, S. D., Whittlestone, S., Hart, K. P., and Hill, S. E.: The flux of radon and thoron from Australian soils, *J. Geophys. Res.*, 100, 26141–26161, 1989.
- SILSO (Sunspot Index and Long-term Solar Observation): World Data Center – Sunspot Number and Long-Term Solar Observations, Royal Observatory of Belgium, on-line Sunspot Number catalogue, available at: <http://sidc.oma.be/silso/>, last access: 15 March 2016.
- Simon, J., Meresova, J., Sykora, I., Jeskovsky, M., and Holy, K.: Modeling of temporal variations of vertical concentration profile of ^7Be in the atmosphere, *Atmos. Environ.*, 43, 2000–2004, 2009.
- Steinmann, P., Zeller, M., Beuret, P., Ferreri, G., and Estier, S.: Cosmogenic ^7Be and ^{22}Na in ground level air in Switzerland (1994–2011), *J. Environ. Radioactiv.*, 124, 68–73, 2013.
- Stohl, A., Wernli, H., James, P., Borqui, M., Forster, C., Liniger, M. A., Seibert, P., and Sprenger, M.: A new perspective of stratosphere-troposphere exchange, *B. Am. Meteorol. Soc.*, 84, 1565–1573, doi:10.1175/BAMS-84-11-1565, 2003.
- Strahan, S. E., Duncan, B. N., and Hoor, P.: Observationally derived transport diagnostics for the lowermost stratosphere and their application to the GMI chemistry and transport model, *Atmos. Chem. Phys.*, 7, 2435–2445, doi:10.5194/acp-7-2435-2007, 2007.
- Todorovic, D., Popovic, D., Djuric, G., and Radenkovic, M.: ^7Be to ^{210}Pb concentration ratio in ground level air in Belgrade area, *J. Environ. Radioactiv.*, 79, 297–307, 2005.
- Tositti, L., Riccio, A., Sandrini, S., Brattich, E., Baldacci, D., Parmeggiani, S., Cristofanelli, P., and Bonasoni, P.: Short-term climatology of PM_{10} at a high altitude background station in southern Europe, *Atmos. Environ.*, 65, 145–152, 2013.
- Tositti, L., Brattich, E., Cinelli, G., and Baldacci, D.: 12 years of ^7Be and ^{210}Pb data in Mt. Cimone, and their correlation with meteorological parameters, *Atmos. Environ.*, 87, 108–122, doi:10.1016/j.atmosenv.2014.01.014, 2014.
- Trickl, T., Feldmann, H., Kanter, H.-J., Scheel, H.-E., Sprenger, M., Stohl, A., and Wernli, H.: Forecasted deep stratospheric intrusions over Central Europe: case studies and climatologies, *Atmos. Chem. Phys.*, 10, 499–524, doi:10.5194/acp-10-499-2010, 2010.
- Trigo, I. F., Bigg, G. R., and Davies, T. D.: Climatology of cyclogenesis mechanisms in the Mediterranean, *Mon. Weather Rev.*, 130, 549–569, 2002.
- Turekian, K. K. and Graustein, W. C.: Natural Radionuclides in the Atmosphere, in: Treatise on Geochemistry, Vol. 4, edited by: Keeling, R. F., Holland, H. D., and Turekian, K. K., 261–279, doi:10.1016/B0-08-043751-6/04042-1, Elsevier, 2003.

- Turekian, K. K., Nozaki, Y., and Benninger, L. K.: Geochemistry of atmospheric radon and radon products, *Annu. Rev. Earth Planet. Sc.*, 5, 227–255, 1977.
- Usoskin, I. and Kovaltsov, G.: Production of cosmogenic ^7Be isotope in the atmosphere: full 3D modelling, *J. Geophys. Res.*, 113, D12107, doi:10.1029/2007JD009725, 2008.
- Van Dingenen, R., Putaud, J.-P., Martins-Dos Santos, S., and Raes, F.: Physical aerosol properties and their relation to air mass origin at Monte Cimone (Italy) during the first MINATROC campaign, *Atmos. Chem. Phys.*, 5, 2203–2226, doi:10.5194/acp-5-2203-2005, 2005.
- Viezee, W. and Singh, H. B.: The distribution of beryllium-7 in the troposphere. Implications on stratosphere/tropospheric air exchange, *Geophys. Res. Lett.*, 7, 805–808, 1980.
- Wilkening, M. H., Clements, W. E., and Stanley, D.: Radon222 flux measurements in widely separated regions, in: *The Natural Radiation Environment II*, 717–730, U.S. Energy and Res. Dev. Admin., Oak Ridge, Tenn, USA, 1975.
- Winkler, R., Dietl, F., Frank, G., and Thiersch, J.: Temporal variation of ^7Be and ^{210}Pb size distributions in ambient aerosols, *Atmos. Environ.*, 32, 983–991, 1998.
- WMO-GAW (World Meteorological Organization – Global Atmosphere Watch): 1st International Expert Meeting on Sources and Measurements of Natural Radionuclides Applied to Climate and Air Quality Studies. WMO TD No. 1201, Report No. 155, Global Atmos. Watch, Geneva, Switzerland, 2004.
- Zanis, P., Schuepbach, E., Gäggeler, H. W., Huebener, S., and Tobler, L.: Factors controlling Beryllium-7 at Jungfrauoch in Switzerland, *Tellus*, 51, 789–805, 1999.
- Zanis, P., Monks, P. S., Schuepbach, R., Carpenter, L. J., Green, T. J., Mills, G. P., Bauguitte, S., and Penkett, S. A.: In situ ozone production under free tropospheric conditions during FREETEX '98 in the Swiss Alps, *J. Geophys. Res.*, 105, 24223–24234, 2000.
- Zanis, P., Gerasopoulos, E., Priller, A., Schnabel, C., Stohl, A., Zerefos, C., Gäggeler, H. W., Tobler, L., Kubik, P. W., Kanter, H. J., Scheel, H. E., Luterbacher, J., and Berger, M.: An estimate of the impact of stratosphere-to-troposphere transport (STT) on the lower free tropospheric ozone over the Alps using ^{10}Be and ^7Be measurements, *J. Geophys. Res.*, 108, 8520, doi:10.1029/2002JD002604, 2003.