

An Entry to Enantioenriched 3,3-Disubstituted Phthalides through Asymmetric Phase-Transfer-Catalyzed γ -Alkylation

Marina Sicignano, Rosaria Schettini, Giovanni Pierri, Maria Leda Marino, Irene Izzo, Francesco De Riccardis, Luca Bernardi, and Giorgio Della Sala*

Cite This: *J. Org. Chem.* 2020, 85, 7476–7484

Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: A novel asymmetric phase-transfer-catalyzed γ -alkylation of phthalide 3-carboxylic esters has been developed, giving access to 3,3-disubstituted phthalide derivatives, which present a chiral quaternary γ -carbon in good to excellent yields and good enantioselectivities (74–88% ee). The enantiomeric purity could be substantially enhanced to 94–95% ee by recrystallization. Both electron-withdrawing and electron-releasing substituents are well tolerated on the phthalide core as well as on the aromatic moiety of the alkylating agent. This methodology, enabling the introduction of an unfunctionalized group at the phthalide γ -position, fully complements previously reported organocatalytic strategies involving functionalized electrophiles, thus expanding the scope of accessible 3,3-disubstituted products. The high synthetic value of this asymmetric reaction has been proven by the formal synthesis of the naturally occurring alkaloid (+)-(9*S*,13*R*)-13-hydroxyisocyclocelabenzine.



INTRODUCTION

The isobenzofuran-1(3*H*)-one core is a ubiquitous pharmacophore incorporated in the structure of a large family of natural products and synthetic analogues, known collectively as phthalides, which display a considerably wide range of useful biological activities.^{1,2} In addition, phthalides are also valuable intermediates for the synthesis of several drugs and naturally occurring compounds such as anthracyclines and other antibacterial and anticancer quinones,³ phthalanes,⁴ isocoumarins,⁵ phthalazines,⁶ and others.^{5,7} Despite the conspicuous efforts devoted to the asymmetric syntheses of phthalides, typically performed through the stereocontrolled formation of the lactone ring, the vast majority of methods furnish 3-monosubstituted lactones, whereas enantioenriched 3,3-disubstituted derivatives containing a chiral γ -quaternary carbon atom, also widely represented in nature (Figure 1), remain elusive.⁸

To this end, the stereocontrolled direct introduction of an electrophilic group at the γ -position of a 3-substituted phthalide has recently emerged as a viable alternative (Figure 2).^{9–14} In this context, γ -activated phthalide esters and nitriles proved to be ideal substrates in reactions catalyzed by bifunctional and polyfunctional base organocatalysts containing a thiourea moiety. Such catalysts, which are tailor-made for functionalized electrophiles capable to interact with hydrogen-bond donor groups, furnished excellent results with imines,¹⁰ Michael acceptors,¹¹ and Morita–Baylis–Hillman carbonates.^{12,13} Remarkably, reactions with unfunctionalized alkylating agents, such as alkyl halides, which lack hydrogen-bond





Figure 1. Representative natural 3,3-disubstituted phthalides with a chiral γ -quaternary carbon atom.



Previous works:
Bifunctional base organocatalyst;
Ei = Functionalized electrophile (Michael acceptor, Imine, MBH carbonate)

This work:
Phase-transfer catalyst; Ei = Alkylating agent

Figure 2. Enantioselective synthesis of 3,3-disubstituted phthalides through direct group insertion at the γ -position.

Received: April 9, 2020

Published: May 14, 2020



acceptor groups, have never been reported to date in asymmetric nor racemic version. For this transformation, which would significantly expand the scope of accessible 3,3-disubstituted products, the abovementioned bifunctional base organocatalysts do not appear suitable.

It is well known that the most effective organocatalytic strategy to achieve enantioselective alkylation of weakly acidic substrates is asymmetric phase-transfer catalysis.¹⁵ Although phthalide 3-carboxylic esters meet all the requirements, their application in phase-transfer catalysis has not yet been explored. In this article, we report the development of a novel phase-transfer-catalyzed alkylation of phthalide 3-carboxylic esters, demonstrating the utility of such a process in asymmetric synthesis.

RESULTS AND DISCUSSION

Our investigation began with a preliminary screening of chiral phase-transfer catalysts (Figure 3) in the reaction of substrate




Figure 3. Phase-transfer catalysts screened in the benzylation of **15a**.

15a with benzyl bromide **16a** (Table 1). In the beginning, various cinchonidinium salts were surveyed in toluene/KOH 50% aq (Table 1, entries 1–7), but although the anticipated product **17aa** was obtained in good yields and very short reaction times in most cases, the enantioselectivities were disappointing, with ee values not exceeding 24%.¹⁶

Lygo's chiral biphenyl azepinium salts **8–11**¹⁷ also led smoothly to the desired product with low ee values (entries 8–11). A small improvement was observed with derivative **8**. Finally, a striking enhancement of enantioselectivity was achieved with Maruoka's *N*-spiro C₂-symmetric catalysts **12** and **13**, even when applied at lower (5 mol %) loadings (entries 12–13). A higher reaction rate and enantiomeric excess were achieved with the latter derivative. A poor result was instead observed with *N,N*-dibutyl ammonium salt **14** (entry 14). A smaller catalyst amount led to the decline of the enantiomeric excess (entry 15). Attempts to use the corresponding benzyl and ethyl phthalide esters under

Table 1. Screening of Catalysts^a

entry	cat (mol %)	t (h)	yield (%) ^b	ee (%) (optical rotation) ^c
1	1 (10)	0.5	99	20 (–)
2	2 (10)	1	87	22 (–)
3	3 (10)	3	41	2 (–)
4	4 (10)	1	52	6 (–)
5	5 (10)	0.5	50	18 (+)
6	6 (10)	0.5	63	22 (+)
7	7 (10)	1	70	24 (–)
8	8 (10)	1	70	34 (–)
9	9 (10)	1	64	16 (–)
10	10 (10)	1	50	12 (–)
11	11 (10)	19	70	4 (–)
12	12 (5)	1	72	56 (+)
13	13 (5)	0.25	81	70 (+)
14	14 (5)	0.5	95	24 (+)
15	13 (2)	0.25	62	43 (+)


^aReaction conditions: **15a** (0.05 mmol), **16a** (0.06 mmol), catalyst (*x* mmol), KOH 50% aq (0.3 mL), toluene (0.5 mL). ^bIsolated yields. ^cDetermined by chiral HPLC.

conditions described in Table 1, entry 13, resulted only in decomposition products.

The X-ray analysis of the major dextrorotatory enantiomer, produced in the reaction catalyzed by (*R,R*)-configured catalyst **13**, made it possible to determine its absolute configuration as (*R*).¹⁸

The effects of the aqueous base and the reaction medium were next studied (Table 2). Bases other than KOH required longer reaction times but led to improved ee values (entries 2–5). Cesium aqueous bases, especially Cs₂CO₃ (entry 3), led to the best combination of ee and yield in reasonable reaction times. Cs₂CO₃ 50% aq was therefore used in the following runs. Both the reaction rate and stereoselectivity were reduced in CH₂Cl₂ (entry 6), whereas good results were observed in ethereal solvents (entries 7–8). However, the best results were generally achieved in aromatic nonpolar solvents (entries 3 and 9–11). Both *p*- and *o*-xylene gave results almost identical to toluene (cf. entries 10–11 with entry 3), suggesting their use as alternative reaction media. However, toluene is easier to be removed. Lower ee values were obtained in more polar solvents such as chlorobenzene and fluorobenzene (entries 12–13). Conducting the reaction in toluene at 0 °C did not yield any enhancement of enantioselectivity (entry 14), whereas at –20 °C, no traces of product were detected, even after 24 h (entry 15).

With the optimized reaction conditions displayed in Table 2, entry 3, we undertook the study of process scope (Scheme 1). Gratefully, uniformly good yields (65–96%) and enantioselectivities (74–88% ee) were achieved with diversely substituted phthalide esters **15a–15e** and benzyl bromides **16a–16h**. Both electron-withdrawing and electron-releasing substituents were well tolerated at the 5- or 6-position of the phthalide (products **17ba–17ea**) as well as on the benzyl moiety (products **17ab–17ah**). The presence of an *ortho*-methyl group on the benzyl bromide partner (product **17ad**) did not affect the reactivity or the enantioselectivity. A good

Table 2. Effect of Base and Solvent^a


entry	aq base	solvent	t (h)	yield (%) ^b	ee (%) ^c
1	KOH	toluene	0.25	81	70
2	CsOH	toluene	1	65	78
3	Cs ₂ CO ₃	toluene	4	82	80
4	K ₂ CO ₃	toluene	72	40	78
5	K ₃ PO ₄	toluene	72	15	70
6	Cs ₂ CO ₃	CH ₂ Cl ₂	72	73	60
7	Cs ₂ CO ₃	Et ₂ O	3	78	73
8	Cs ₂ CO ₃	MTBE	6	62	71
9	Cs ₂ CO ₃	mesitylene	4	73	76
10	Cs ₂ CO ₃	<i>p</i> -xylene	4	83	80
11	Cs ₂ CO ₃	<i>o</i> -xylene	4	80	80
12	Cs ₂ CO ₃	chlorobenzene	20	75	72
13	Cs ₂ CO ₃	fluorobenzene	4	63	60
14 ^d	Cs ₂ CO ₃	toluene	20	88	78
15 ^e	Cs ₂ CO ₃	toluene	20		


^aReaction conditions: **15a** (0.05 mmol), **16a** (0.06 mmol), **13** (0.0025 mmol), aqueous base (50% w/w, 0.3 mL), toluene (0.5 mL).
^bIsolated yields. ^cDetermined by chiral HPLC. ^dReaction performed at 0 °C. ^eReaction performed at -20 °C.

result was also achieved in the reaction with allyl bromide (**17ai**, 54% yield, 84% ee).¹⁹ It is worth noting that no traces of hydrolysis byproducts were detected in all the cases examined. In addition, we were delighted to find that products could be easily enantioenriched up to >90% ee through recrystallization from *n*-hexane. As examples, **17aa** was enantioenriched from 80 to 94% ee, and **17ai** was enantioenriched from 84 to 95% ee, with acceptable overall yields in both cases (66 and 51%, respectively).

During the scale-up optimization of this process we found that good yield and comparable enantioselectivity can be achieved at a 1 mmol scale by using only 2 mol % catalyst **13** (62%, 78% ee for product **17aa**).

It is reasonable to assume that the observed enantioselectivity comes from a preferential orientation of the substrate anion, generated upon deprotonation of phthalide ester **15**, within the ion pair formed with the *N*-spiro C₂-symmetric cation. Transition state models, based on DFT calculations, have been previously reported for the benzylation of *N*-(diphenylmethylene)glycine *tert*-butyl esters catalyzed by these Maruoka's ammonium salts. In these models, coulombic interactions as well as nonclassical hydrogen bonds between both reactants and the two catalysts' enantiotopic fluorinated aromatic groups play a key role in dictating the mutual orientation of the alkylating agent and the substrate anion.²⁰ Thus, building on these computed catalyst conformation and interactions with the substrates, it is possible to tentatively sketch a transition state for the alkylation of *tert*-butyl phthalide 3-carboxylate **15a** with benzyl bromide **16a**, catalyzed by (*R,R*)-**13** (Figure 4). In this model, reaction occurs at the *Si*-face of the substrate anion, accounting for the high selectivity observed toward the (*R*)-**17aa** product.

Then, to further stress the synthetic significance of the present methodology in gaining access to new enantioenriched phthalide compounds, we focused on the enantioselective

Scheme 1. Scope of the Phase-Transfer-Catalyzed Alkylation of Phthalide Esters^{a-c}

^aThe following reaction conditions were applied unless otherwise specified: **15** (0.10 mmol), **16** (0.12 mmol), **13** (0.005) mmol, Cs₂CO₃ 50% aq (0.6 mL), toluene (1.0 mL). ^bIsolated yields. ^cee determined by chiral HPLC. ^dValues in parenthesis refer to the product after recrystallization. ^eAllyl bromide (0.50 mmol) was used.





Figure 4. Tentative transition state model for the reaction between phthalide 3-carboxylic ester **15a** and benzyl bromide **16a**, catalyzed by (*R,R*)-**13**.

preparation of nitrile (*R*)-**18**, a key intermediate for the synthesis of the naturally occurring spermidine alkaloid

Scheme 2. Formal Synthesis of (+)-(9*S*,13*R*)-Hydroxyisocyclocelabenzine

(+)-(9*S*,13*R*)-13-hydroxyisocyclocelabenzine (Scheme 2).^{7a,21} It should be noted that the control of the (*R*)-configuration in **18** is an essential requirement to achieve the (13*R*)-configuration in the dihydroisoquinoline moiety of the target natural product. The original synthesis involved the preparation and use of **18** as a racemate, leading to the target natural product as a mixture of (9*S*,13*R*) and (9*S*,13*S*) diastereomers, which were separated only by tedious repeated flash chromatography.^{7a} Our first synthesis of (*R*)-**18** was accomplished in six steps starting from commercially available parent phthalide, with an overall 28% yield and 95% ee (Scheme 2). The enantioselective allylation of **15a** followed by fractional crystallization afforded **17ai** in 51% yield and 95% ee, as described before. Then, de-*tert*-butylation, synthesis of primary amide **19**, and subsequent dehydration²² afforded (*R*)-**18**, a formal precursor of (+)-(9*S*,13*R*)-13-hydroxyisocyclocelabenzine.²³

CONCLUSIONS

In conclusion, the first asymmetric γ -alkylation of phthalide 3-carboxylic esters has been herein described, affording 3,3-disubstituted products incorporating benzyl and allyl groups with generally high yields and good enantioselectivity. Excellent enantiomeric excesses could be achieved by recrystallization. The present and previous studies¹⁰ demonstrated that the carboxylic group can be readily manipulated giving access to functionalized 3,3-disubstituted phthalides that were previously inaccessible in enantioenriched form. For example, we developed the enantioselective synthesis of a precursor of (+)-(9*S*,13*R*)-13-hydroxyisocyclocelabenzine. Further synthetic applications of this methodology will be reported by us in due course.

EXPERIMENTAL SECTION

General Remarks. All the chemicals and solvents were purchased from commercial suppliers and used without further purification. Catalysts **1–4** and **12–14** are commercially available, whereas catalysts **5–7**²⁴ and **8–11**^{17a} were prepared following the general procedure described in the literature. 5-Chloro-, 6-chloro-, 5-methoxy-, and 6-methoxyisobenzofuran-1(3*H*)-ones were prepared as described in the literature.²⁵ Reactions were monitored by analytical thin-layer chromatography (TLC) on precoated silica gel plates (0.25 mm) and visualized by UV light or by spraying KMnO₄/ethanol or ninhydrin/ethanol solutions and heating on a hot plate. Flash chromatography was performed on silica gel 60 (particle size: 0.040–0.063 mm). ¹H and ¹³C NMR spectra were recorded on a Bruker Avance-600 MHz spectrometer and a Bruker Avance-400 MHz spectrometer at room temperature in CDCl₃, respectively. All

the NMR spectra were referenced to residual CHCl₃ (7.26 ppm, ¹H; 77.16 ppm, ¹³C). The following abbreviations are used to indicate the multiplicity in NMR spectra: s = singlet; d = doublet; dd = double doublet; t = triplet; bs = broad signal; and m = multiplet. Coupling constants (*J*) are quoted in Hertz. Optical rotations were measured on a Jasco P-2000 digital polarimeter using WI (tungsten-halogen) lamp ($\lambda = 589$ nm). Enantiomeric excesses were determined using a CHIRALPAK AS-H column (ϕ 0.46 cm \times 25 cm) on a JASCO PU-4180 instrument equipped with a photodiode array detector MD-4015 set at 220 nm. High-resolution mass spectra (HRMS) were acquired using a Bruker solariX XR Fourier transform ion cyclotron resonance mass spectrometer (Bruker Daltonik GmbH, Bremen, Germany) equipped with a 7 T refrigerated actively shielded superconducting magnet and with a LTQ Orbitrap XL Thermo Scientific. The samples were ionized in positive-ion mode using a MALDI or ESI ionization source.

Synthesis of Substrates. *tert*-Butyl 3-Oxo-1,3-dihydroisobenzofuran-1-carboxylate (15a**).** LDA was freshly prepared by adding a 2.5 M butyllithium solution in hexanes (11 mL, 27.5 mmol) to an anhydrous solution of 0.5 M isopropylamine in THF (55 mL, 27.5 mmol) (28.0 mL) at -78 °C under a nitrogen atmosphere. The mixture was stirred for 30 min at the same temperature, and then a solution of isobenzofuran-1(3*H*)-one (2.49 g, 18.6 mmol) in anhydrous THF (3.8 mL) was slowly added. The resulting mixture was stirred for 50 min at -78 °C under a nitrogen atmosphere. Then, the reaction vessel was saturated with carbon dioxide (three freeze-pump-thaw cycles followed by connection with a carbon dioxide balloon), and stirring was kept at -78 °C for 2 h. Once reaction was complete, NH₄Cl saturated aqueous solution (15 mL) was added dropwise, and then THF was removed under reduced pressure. The mixture was basified with Na₂CO₃ saturated aqueous solution until pH 9 and washed with AcOEt (2 \times 20 mL). The aqueous phase was acidified with concd HCl solution until pH 1, and the product was extracted with AcOEt (3 \times 50 mL). The combined organic phases were dried over Na₂SO₄ and concentrated under reduced pressure. The resulting crude phthalide 3-carboxylic acid (3.31 g, 18.6 mmol) was dissolved in anhydrous CH₂Cl₂ (70 mL), and *tert*-butyl 2,2,2-trichloroacetimidate (3.3 mL, 18.6 mmol) was added. The reaction mixture was stirred for 48 h under a nitrogen atmosphere, then diluted with CH₂Cl₂, and centrifuged. The supernatant solution was concentrated under reduced pressure, and the crude residue was purified by flash chromatography (silica gel; petroleum ether/ethyl acetate, 95:5 to 80:20), affording **15a** as a white solid (4.36 g, 99% yield). The characterization data matched those previously reported.^{12a} ¹H NMR (400 MHz, CDCl₃) δ 7.92 (d, *J* = 7.6 Hz, 1H), 7.71 (t, *J* = 7.6 Hz, 1H), 7.64 (d, *J* = 7.6 Hz, 1H), 7.58 (t, *J* = 7.6 Hz, 1H), 5.76 (s, 1H), 1.48 (s, 9H).

***tert*-Butyl 5-Methoxy-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (**15b**).** A mixture of 3-methoxybenzoic acid (0.76 g, 5.0 mmol), glyoxylic acid monohydrate (0.92 g, 10 mmol), concd H₂SO₄ (0.55 mL, 10 mmol), and glacial acetic acid (20.0 mL) was stirred at 80 °C. After 48 h, the reaction mixture was cooled to room temperature and extracted with AcOEt (3 \times 50 mL). The combined organic phases

were dried over Na_2SO_4 and concentrated under reduced pressure, affording crude phthalide 3-carboxylic acid (0.56 g, 2.7 mmol). This solid was dissolved in anhydrous CH_2Cl_2 (10 mL) and treated with *tert*-butyl 2,2,2-trichloroacetimidate (0.59 g, 2.7 mmol) under a nitrogen atmosphere. After stirring for 48 h, the reaction mixture was diluted with CH_2Cl_2 and centrifuged. The supernatant solution was concentrated under reduced pressure, and the crude residue was purified by flash chromatography (silica gel; petroleum ether/ethyl acetate, 95:5 to 80:20), affording **15b** as a white solid (0.30 g, 23% yield). mp 89–90 °C. ^1H NMR (400 MHz, CDCl_3) δ 7.49 (d, J = 8.4 Hz, 1H), 7.30 (bs, 1H), 7.23 (d, J = 8.4, 1H), 5.68 (s, 1H), 3.85 (s, 3H), 1.46 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 169.7, 165.6, 161.3, 136.8, 126.4, 123.2, 123.2, 107.7, 83.7, 77.4, 55.8, 27.8. HRMS (MALDI) m/z : $[\text{M} + \text{Na}^+]$ calcd for $\text{C}_{14}\text{H}_{16}\text{NaO}_5^+$, 287.0890; found, 287.0892.

tert-Butyl 5-Chloro-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (15c). Compound **15c** was prepared following the procedure described for **15a**, starting from 5-chloroisobenzofuran-1(3H)-one (3.14 g, 18.6 mmol). Purification of the crude product by flash chromatography (silica gel; petroleum ether/ethyl acetate, 95:5 to 80:20) afforded **15c** as a white solid. Yield: 1.85 g (37%). mp 87–88 °C. ^1H NMR (400 MHz, CDCl_3) δ 7.80 (d, J = 1.4 Hz, 1H), 7.64 (dd, J = 8.2, 1.9 Hz, 1H) 7.57, (d, J = 8.2 Hz, 1H), 5.73 (s, 1H), 1.45 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 168.2, 164.9, 142.7, 136.4, 134.8, 126.8, 125.7, 123.9, 84.4, 77.5, 27.8. HRMS (MALDI) m/z : $[\text{M} + \text{H}^+]$ calcd for $\text{C}_{13}\text{H}_{14}\text{ClO}_4^+$, 269.0575; found, 269.0586.

tert-Butyl 6-Methoxy-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (15d). Compound **15d** was prepared following the procedure described for **15a**, starting from 6-methoxyisobenzofuran-1(3H)-one (3.05 g, 18.6 mmol). Purification of the crude product by flash chromatography (silica gel; petroleum ether/ethyl acetate, 95:5 to 80:20) afforded **15d** as a white solid. Yield: 2.36 g (48%). mp 82–81 °C. ^1H NMR (600 MHz, CDCl_3) δ 7.81 (d, J = 8.6 Hz, 1H), 7.09–7.06 (m, 2H), 5.68 (s, 1H), 3.90 (s, 3H), 1.49 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CDCl_3) δ 169.4, 165.6, 164.9, 147.3, 127.4, 117.3, 117.2, 106.6, 84.0, 77.2, 55.9, 27.9. HRMS (MALDI) m/z : $[\text{M} + \text{H}^+]$ calcd for $\text{C}_{14}\text{H}_{17}\text{O}_5^+$, 265.1071; found, 265.1076.

tert-Butyl 6-Chloro-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (15e). Compound **15e** was prepared following the procedure described for **15a**, starting from 6-chloroisobenzofuran-1(3H)-one (3.14 g, 18.6 mmol). Purification of the crude product by flash chromatography (silica gel; petroleum ether/ethyl acetate, 95:5 to 80:20) afforded **15e** as a white solid. Yield: 1.80 g (36%). mp 94–95 °C. ^1H NMR (400 MHz, CDCl_3) δ 7.83 (d, J = 8.2 Hz, 1H), 7.62 (d, J = 1.2 Hz, 1H), 7.55 (dd, J = 8.2, 1.2 Hz, 1H), 5.72 (s, 1H), 1.49 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 168.4, 164.8, 146.1, 141.1, 130.8, 127.0, 123.6, 123.0, 84.5, 77.1, 27.9. HRMS (MALDI) m/z : $[\text{M} + \text{H}^+]$ calcd for $\text{C}_{13}\text{H}_{14}\text{ClO}_4^+$, 269.0575; found, 269.0582.

Ethyl 3-Oxo-1,3-dihydroisobenzofuran-1-carboxylate (15f). To a solution of 3-oxo-1,3-dihydroisobenzofuran-1-carboxylic acid, prepared as described above (0.2 g, 1.1 mmol), in anhydrous ethanol (80 mL, 1.4 mmol), thionyl chloride (96 μL , 1.3 mmol) was added. The resulting solution was stirred for 2 h at room temperature under a nitrogen atmosphere. Next, the excess alcohol was evaporated, and water (10 mL) was added to the residue. The mixture was extracted with AcOEt (3 \times 20 mL), and the combined organic phases were dried over Na_2SO_4 , filtered, and concentrated under reduced pressure, affording **15f** that was used without further purification (0.22 g, 96% yield). The characterization data of compounds **15f** matched those previously reported.²⁶ ^1H NMR (400 MHz, CDCl_3) δ 7.92 (d, J = 7.6 Hz, 1H), 7.75–7.64 (m, 2H), 7.59 (t, J = 7.7 Hz, 1H), 5.87 (s, 1H), 4.28 (m, 2H), 1.30 (m, 3H).

Benzyl 3-Oxo-1,3-dihydroisobenzofuran-1-carboxylate (15g). Compound **15g** was prepared following the procedure described for **15f**, employing benzyl alcohol (140 mL, 1.4 mmol) that was used without further purification. Yield: 0.26 g (90% yield). The characterization data of compounds **15g** matched those previously reported.²⁷ ^1H NMR (400 MHz, CDCl_3) δ 7.93 (d, J = 7.5 Hz, 1H), 7.68 (m, 1H), 7.63–7.58 (m, 2H), 7.44–7.28 (m, 5H), 5.92 (s, 1H), 5.28 (d, J = 12.1 Hz, 1H), 5.22 (d, J = 12.1 Hz, 1H).

General Procedure for the Enantioselective Alkylation of Phthalide 3-Carboxylic *tert*-Butyl Esters 15a–15e. To a solution of phthalide ester **15** (1.0 equiv, 0.1 mmol) in toluene (1 mL), contained in a 4 mL vial, (*R,R*)-**13** (0.05 equiv, 0.005 mmol), alkyl bromide **16** (1.2 equiv, 0.12 mmol), and 50 % aqueous Cs_2CO_3 (0.6 mL) were added. The reaction mixture was vigorously stirred (900 rpm) at room temperature for the time specified. After completion (TLC), it was diluted with 1 M HCl (1 mL) and extracted with CH_2Cl_2 (3 \times 5 mL). The combined organic phases were dried over Na_2SO_4 and concentrated under reduced pressure. The crude residue was purified by flash chromatography, affording products **17** as white solids.

tert-Butyl (*R*)-1-Benzyl-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17aa). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 26.5 mg (82%). mp 88–89 °C. $[\alpha]_{\text{D}}^{22} +31.4$ (c 0.20, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 7.75 (d, J = 7.5 Hz, 1H), 7.73–7.66 (2H, m), 7.50 (1H, m), 7.20–7.12 (5H, m), 3.64 (1H, d, J = 14.3), 3.35 (1H, d, J = 14.3), 1.37 (9H, s). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 169.1, 167.2, 147.7, 134.1, 133.4, 130.4, 129.9, 128.1, 127.2, 125.7, 125.6, 122.5, 87.6, 83.8, 43.1, 27.7. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 16.4$ min, $\tau_{\text{major}} = 21.4$ min (80% ee). HRMS (MALDI) m/z : $[\text{M} + \text{H}^+]$ calcd for $\text{C}_{20}\text{H}_{21}\text{O}_4^+$, 325.1434; found, 325.1437.

tert-Butyl (*R*)-1-(4-(*tert*-Butyl)benzyl)-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ab). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 36.5 mg (96%). mp 99–100 °C. $[\alpha]_{\text{D}20} +63.5$ (c 0.20, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 7.78 (d, J = 7.6 Hz, 1H), 7.75–7.65 (m, 2H), 7.56–7.47 (m, 1H), 7.20 (d, J = 8.2 Hz, 2H), 7.11 (d, J = 8.2 Hz, 2H), 3.62 (d, J = 14.3 Hz, 1H), 3.28 (d, J = 14.3 Hz, 1H), 1.35 (s, 9H), 1.24 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 169.1, 167.1, 150.0, 147.8, 134.1, 130.4, 130.0, 129.8, 125.7, 125.6, 124.9, 122.5, 87.8, 83.6, 42.8, 34.3, 31.2, 27.6. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.3 mLmin⁻¹): $\tau_{\text{minor}} = 28.1$ min, $\tau_{\text{major}} = 31.4$ min (88% ee). HRMS (MALDI) m/z : $[\text{M} + \text{Na}^+]$ calcd for $\text{C}_{24}\text{H}_{28}\text{NaO}_4^+$, 403.1880; found, 403.1882.

tert-Butyl (*R*)-1-(4-Methylbenzyl)-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ac). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 32.1 mg, (95%). mp 93–94 °C. $[\alpha]_{\text{D}20} +24.4$ (c 0.20, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 7.75 (d, J = 7.6 Hz, 1H), 7.72–7.64 (m, 2H), 7.56–7.45 (m, 1H), 7.01 (d, J = 7.9 Hz, 2H), 6.96 (d, J = 7.9 Hz, 2H), 3.60 (d, J = 14.3 Hz, 1H), 3.32 (d, J = 14.3 Hz, 1H), 2.23 (s, 3H), 1.39 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 169.1, 167.2, 147.6, 136.7, 134.0, 130.2, 130.1, 129.8, 128.7, 125.7, 125.6, 122.5, 87.7, 83.7, 42.6, 27.7, 21.0. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 22.3$ min, $\tau_{\text{major}} = 32.0$ min (82% ee). HRMS (MALDI) m/z : $[\text{M} + \text{K}^+]$ calcd for $\text{C}_{21}\text{H}_{22}\text{KO}_4^+$, 377.1150; found, 377.1163.

tert-Butyl (*R*)-1-(2-Methylbenzyl)-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ad). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 32.5 mg (96%). mp 90–91 °C. $[\alpha]_{\text{D}20} +40.8$ (c 0.40, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 7.81–7.66 (m, 3H), 7.57–7.49 (m, 1H), 7.11–7.01 (m, 3H), 6.95 (m, 1H), 3.75 (d, J = 14.7 Hz, 1H), 3.35 (d, J = 14.7 Hz, 1H), 2.33 (s, 3H), 1.38 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 168.9, 167.4, 147.9, 137.6, 134.0, 131.9, 130.4 (2C), 129.9, 127.2, 125.6 (2C), 125.4, 122.6, 88.1, 83.6, 39.4, 27.7, 20.1. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 15.0$ min, $\tau_{\text{major}} = 17.8$ min (84% ee). HRMS (MALDI) m/z : $[\text{M} + \text{K}^+]$ calcd for $\text{C}_{21}\text{H}_{22}\text{NaO}_4^+$, 361.1410; found, 361.1421.

tert-Butyl (*R*)-1-(4-Methoxybenzyl)-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ae). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate 98:2 to 80:20). Yield: 32.6 mg (92%). mp 97–98 °C. $[\alpha]_{\text{D}20} +11.9$ (c 0.80, CHCl_3). ^1H NMR

(600 MHz, CDCl₃) δ 7.75 (d, J = 7.7 Hz, 1H), 7.71–7.67 (m, 2H), 7.50 (m, 1H), 7.05 (d, J = 8.2 Hz), 6.69 (d, J = 8.2 Hz), 3.73 (s, 3H), 3.58 (d, J = 14.3 Hz), 3.30 (d, J = 14.3 Hz), 1.39 (s, 9H). ¹³C{¹H} NMR (150 MHz, CDCl₃) δ 169.1, 167.3, 158.7, 147.7, 134.1, 131.5, 129.8, 125.9, 125.6, 125.3, 122.5, 113.5, 87.8, 83.7, 55.1, 42.3, 27.8. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 26.8$ min, $\tau_{\text{major}} = 42.9$ min (80% ee). HRMS (ESI) m/z : [M + Na⁺] calcd for C₂₁H₂₂NaO₅⁺, 377.1359; found, 377.1359.

tert-Butyl (R)-1-(4-Chlorobenzyl)-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17af). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 33.7 mg (94%). mp 110–111 °C. [α]_{D20} +14.4 (c 0.20, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.82–7.74 (m, 1H), 7.74–7.66 (m, 2H), 7.57–7.48 (m, 1H), 7.15 (d, J = 8.3 Hz, 2H), 7.09 (d, J = 8.3 Hz, 2H), 3.62 (d, J = 14.3 Hz, 1H), 3.31 (d, J = 14.3 Hz, 1H), 1.38 (s, 9H). ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 168.8, 167.0, 147.3, 134.2, 133.2, 131.8, 131.7, 130.0, 128.2, 125.8, 125.6, 122.3, 87.3, 83.9, 42.2, 27.7. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 21.3$ min, $\tau_{\text{major}} = 28.8$ min (80% ee). HRMS (MALDI) m/z : [M + K⁺] calcd for C₂₀H₁₉ClKO₄⁺, 397.0603; found, 397.0609.

tert-Butyl (R)-1-(4-Nitrobenzyl)-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ag). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 33.6 mg (91%). mp 96–97 °C. [α]_{D20} +37.4 (c 0.80, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.05 (d, J = 8.6 Hz, 2H), 7.82–7.69 (m, 3H), 7.60–7.49 (m, 1H), 7.56 (d, J = 8.6 Hz, 2H), 3.79 (d, J = 14.2 Hz, 1H), 3.43 (d, J = 14.2 Hz, 1H), 1.37 (s, 9H). ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 168.6, 166.6, 147.1, 147.0, 141.1, 134.5, 131.3, 130.32, 126.0, 125.4, 123.2, 122.1, 86.7, 84.3, 42.4, 27.6. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 80:20, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 35.2$ min, $\tau_{\text{major}} = 45.7$ min (82% ee). HRMS (MALDI) m/z : [M + K⁺] calcd for C₂₀H₁₉NKO₆⁺, 408.0844; found, 408.0844.

tert-Butyl (R)-1-(Naphthalen-2-ylmethyl)-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ah). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 24.3 mg (65%). mp 128–129 °C. [α]_{D20} +37.1 (c 0.20, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.80–7.68 (m, 5H), 7.66 (d, J = 8.4 Hz, 1H), 7.63 (s, 1H), 7.53–7.46 (m, 1H), 7.44–7.38 (m, 2H), 7.30 (d, J = 8.4 Hz, 1H), 3.82 (d, J = 14.3 Hz, 1H), 3.51 (d, J = 14.3 Hz, 1H), 1.36 (s, 9H). ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 169.0, 167.1, 147.6, 134.1, 133.0, 132.4, 131.0, 129.9, 129.3, 128.3, 127.6 (2C), 127.4, 125.9, 125.7, 125.7 (2C), 122.5, 87.7, 83.8, 43.2, 27.7. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 26.2$ min, $\tau_{\text{major}} = 36.7$ min (74% ee). HRMS (MALDI) m/z : [M + K⁺] calcd for C₂₄H₂₂KO₄⁺, 413.1150; found, 413.1157.

tert-Butyl (R)-1-Allyl-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ai). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 14.8 mg (54%). mp 90–91 °C. [α]_{D20} +44.3 (c 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.88 (d, J = 7.6 Hz, 1H), 7.73–7.66 (m, 1H), 7.64–7.52 (m, 2H), 5.62 (m, 1H), 5.20–5.05 (m, 2H), 3.10 (dd, J = 14.4, 7.7 Hz, 1H), 2.77 (dd, J = 14.4, 6.5 Hz, 1H), 1.43 (s, 9H). ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 169.2, 167.0, 147.7, 134.3, 129.9, 129.8, 125.7, 125.6, 122.3, 120.7, 87.1, 83.7, 41.3, 27.7. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 18.2$ min, $\tau_{\text{major}} = 23.5$ min (84% ee). HRMS (MALDI) m/z : [M + K⁺] calcd for C₁₆H₁₈KO₄⁺, 313.0837; found, 313.0842.

tert-Butyl (R)-1-Benzyl-5-methoxy-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ba). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 34.0 mg (96%). mp 109–110 °C. [α]_{D20} +70.4 (c 0.80, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.57 (d, J = 8.4 Hz, 1H), 7.23 (dd, J = 8.4, 2.3 Hz, 1H), 7.20–7.10 (m, 6H), 3.83 (s, 3H), 3.60 (d, J = 14.3 Hz, 1H), 3.33 (d, J = 14.3 Hz, 1H), 1.37 (s, 9H). ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 169.0, 167.3, 161.1, 140.0, 133.4, 130.4, 128.01, 127.2,

127.1, 123.4, 122.9, 107.3, 87.4, 83.6, 55.7, 43.0, 27.7. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 98:2, 1.0 mLmin⁻¹): $\tau_{\text{minor}} = 29.1$ min, $\tau_{\text{major}} = 32.3$ min (80% ee). HRMS (MALDI) m/z : [M + K⁺] calcd for C₂₁H₂₂KO₅⁺, 393.1099; found, 393.1110.

tert-Butyl (R)-1-Benzyl-5-chloro-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ca). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 34.4 mg (96%). mp 98–99 °C. [α]_{D20} +16.7 (c 0.68, CHCl₃). ¹H NMR (600 MHz, CDCl₃) δ 7.70 (m, 1H), 7.65–7.63 (m, 2H), 7.20–7.16 (m, 3H), 7.14–7.11 (m, 2H), 3.62 (d, J = 14.4 Hz, 1H), 3.36 (d, J = 14.4 Hz, 1H), 1.39 (s, 9H). ¹³C{¹H} NMR (150 MHz, CDCl₃) δ 167.5, 166.8, 145.8, 136.3, 134.4, 133.0, 130.4, 128.2, 127.6, 127.4, 125.5, 123.9, 87.6, 84.1, 43.0, 27.7. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{major}} = 12.1$ min, $\tau_{\text{minor}} = 14.1$ min (82% ee). HRMS (MALDI) m/z : [M + K⁺] calcd for C₂₀H₁₉ClKO₄⁺, 397.0603; found, 397.0604.

tert-Butyl (R)-1-Benzyl-6-methoxy-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17da). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 34.0 mg (96%). mp 88–89 °C. [α]_{D20} +5.6 (c 0.80, CHCl₃). ¹H NMR (600 MHz, CDCl₃) δ 7.66 (d, J = 8.7 Hz, 1H), 7.21–7.16 (m, 5H), 7.13 (d, J = 2.1 Hz, 1H), 7.01 (dd, J = 8.7, 2.1 Hz, 1H), 3.93 (s, 3H), 3.61 (d, J = 13.8 Hz, 1H), 3.32 (d, J = 13.8 Hz, 1H), 1.37 (s, 9H). ¹³C{¹H} NMR (150 MHz, CDCl₃) δ 168.8, 167.4, 164.7, 150.5, 133.6, 130.5, 128.1, 127.2, 127.1, 118.2, 117.1, 106.8, 87.0, 83.7, 55.9, 43.4, 27.8. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 1.0 mLmin⁻¹): $\tau_{\text{minor}} = 29.9$ min, $\tau_{\text{major}} = 61.4$ min (78% ee). HRMS (MALDI) m/z : [M + H⁺] calcd for C₂₁H₂₃O₅⁺, 355.1540; found, 355.1543.

tert-Butyl (R)-1-Benzyl-6-chloro-3-oxo-1,3-dihydroisobenzofuran-1-carboxylate (17ea). White solid after flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20). Yield: 34.1 mg (95%). mp 95–96 °C. [α]_{D20} +32.2 (c 0.85, CHCl₃). ¹H NMR (600 MHz, CDCl₃) δ 7.70 (d, J = 1.6 Hz, 1H), 7.68 (d, J = 8.0 Hz, 1H), 7.48 (dd, J = 8.0, 1.6 Hz, 1H), 7.20–7.17 (m, 3H), 7.16–7.13 (m, 2H), 3.62 (d, J = 13.6 Hz, 1H), 3.33 (d, J = 13.6 Hz, 1H), 1.39 (s, 9H). ¹³C{¹H} NMR (150 MHz, CDCl₃) δ 167.8, 166.7, 149.3, 140.8, 132.9, 130.6, 130.4, 128.2, 127.4, 126.7, 124.3, 123.1, 87.2, 84.3, 43.2, 27.7. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 13.2$ min, $\tau_{\text{major}} = 18.4$ min (80% ee). HRMS (MALDI) m/z : [M + K⁺] calcd for C₂₀H₁₉ClKO₄⁺, 397.0603; found, 397.0603.

Scaled-up Procedure for the Enantioselective Alkylation of Phthalide Ester 15a. To a solution of phthalide ester 15a (234 mg, 1.00 mmol) in toluene (10 mL), contained in a 50 mL round-bottom flask, (*R,R*)-13 (22 mg, 0.020 mmol), benzyl bromide (205 mg, 1.20 mmol), and 50% aqueous Cs₂CO₃ (6.6 mL) were added. The reaction mixture was vigorously stirred (900 rpm) at room temperature for 20 h followed by addition of 1 M HCl (20 mL) and extraction with CH₂Cl₂ (3 × 50 mL). The combined organic phases were dried over Na₂SO₄ and concentrated under reduced pressure. The crude residue was purified by flash chromatography (silica gel; petroleum ether/ethyl acetate, 98:2 to 80:20), affording 17aa as a white solid (201 mg, 62% yield). The % ee was determined by chiral HPLC, as described above (78% ee).

Recrystallization of Compound 17aa. Compound 17aa (26.5 mg, 82% yield, 80% ee) was dissolved in hot hexane (1 mL), and the resulting solution was cooled down at 4 °C. After 20 h, crystals of racemate were formed. The supernatant solution was separated and concentrated under reduced pressure, affording enantioenriched 17aa (21.9 mg, 66% yield). [α]_{D20} = +37.2 (c 0.20, CHCl₃). The % ee was determined by chiral HPLC, as described above (94% ee).

Recrystallization of Compound 17ai. Compound 17ai (14.8 mg, 54% yield, 84% ee) was dissolved in hot hexane (1 mL), and the resulting solution was cooled down at 4 °C. After 20 h, crystals of racemate were formed. The supernatant solution was separated and concentrated under reduced pressure, affording enantioenriched 17ai

(14.0 mg, 51% yield). $[\alpha]_{D20} +55.6$ (c 1.0, CHCl_3). The % ee was determined by chiral HPLC, as described above (95% ee).

Formal Synthesis of (+)-(9S,13R)-13-Hydroxyisocycloclabenzene. (*R*)-1-Allyl-3-oxo-1,3-dihydroisobenzofuran-1-carboxamide (**19**). To a solution of **17ai** (26.1 mg, 0.12 mmol, 95% ee after recrystallization, see above) in anhydrous CH_2Cl_2 (0.65 mL), trifluoroacetic acid (0.12 mL) was added dropwise. After stirring for 8 h, the mixture was concentrated under reduced pressure, affording carboxylic acid **25** as a white solid (28.0 mg), which was used in the next step without further purification.

Carboxylic acid **25** was dissolved in dioxane (0.1 mL), and aqueous ammonia (0.2 mL) was added dropwise at 0 °C. Once the addition was complete, the mixture was allowed to warm to room temperature, and H_2O (0.2 mL) was added. The resulting reaction mixture was stirred for 2 h and then extracted with CH_2Cl_2 (3×5 mL). The combined organic phases were dried over Na_2SO_4 and concentrated under reduced pressure. The crude residue was passed through a silica gel short path by eluting with ethyl acetate, affording **19** as a colorless oil (21.0 mg, 74% yield). $[\alpha]_{D20} +93.2$ (c 1.0, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 7.86–7.82 (m, 2H), 7.71 (t, $J = 7.4$ Hz, 1H), 7.56 (t, $J = 7.4$ Hz, 1H), 6.57 (bs, 1H), 5.84 (bs, 1H), 5.54 (m, 1H), 5.18–5.01 (m, 2H), 3.11 (dd, $J = 14.2, 7.4$ Hz, 1H), 2.80 (dd, $J = 14.2, 6.8$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 170.9, 170.0, 148.0, 135.0, 130.0, 129.4, 125.5, 124.4, 123.5, 121.2, 87.9, 42.0. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 18.6$ min, $\tau_{\text{major}} = 46.6$ min (95% ee). HRMS (ESI) m/z : $[\text{M} + \text{H}^+]$ calcd for $\text{C}_{12}\text{H}_{12}\text{O}_3\text{N}^+$, 218.0812; found, 218.0814.

(*R*)-1-Allyl-3-oxo-1,3-dihydroisobenzofuran-1-carbonitrile (**18**). To a solution of **19** (19.3 mg, 0.097 mmol) in anhydrous CH_3CN (1.0 mL), Ph_3PO (2.7 mg, 9.7 μmol) and NEt_3 (40 μL , 29 mg, 0.29 mmol) were added, and then $(\text{COCl})_2$ (17 μL , 25 mg, 0.19 mmol) was injected dropwise. After stirring for 2 h, the reaction mixture was diluted with CH_2Cl_2 , filtered on a small pad of celite, and then concentrated under reduced pressure. The crude residue was purified by flash chromatography (silica gel; petroleum ether/ethyl acetate, 90:10 to 60:40), affording **18** as a white solid (14.5 mg, 75% yield). mp 77–78 °C. $[\alpha]_{D20} -11.1$ (c 1.0, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 7.96 (d, $J = 7.9$ Hz, 1H), 7.82 (m, 1H), 7.71–7.64 (m, 2H), 5.72 (m, 1H), 5.33–5.27 (m, 2H), 3.03–2.90 (m, 2H). ^{13}C NMR (100 MHz, CDCl_3) δ 167.1, 145.4, 135.4, 131.3, 127.3, 126.5, 124.7, 123.4, 122.3, 115.8, 77.0, 43.4. The % ee was determined by chiral HPLC (CHIRALPAK AS-H column, *n*-hexane/*i*-PrOH = 90:10, 0.5 mLmin⁻¹): $\tau_{\text{minor}} = 29.2$ min, $\tau_{\text{major}} = 32.2$ min (95% ee). HRMS (ESI) m/z : $[\text{M} + \text{H}^+]$ calcd for $\text{C}_{12}\text{H}_{10}\text{O}_2\text{N}^+$, 200.0706; found, 200.0706.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.0c00880>.

Table of optimization, synthesis of ethyl and benzyl substrates, copies of NMR spectra, HPLC traces, and crystallographic data for **17aa** (PDF)

Crystallographic data for **17aa** (CIF)

■ AUTHOR INFORMATION

Corresponding Author

Giorgio Della Sala – Dipartimento di Chimica e Biologia “A. Zambelli”, Università degli Studi di Salerno, 84084 Fisciano, SA, Italy; orcid.org/0000-0001-5020-8502; Email: gdsala@unisa.it

Authors

Marina Sicignano – Dipartimento di Chimica e Biologia “A. Zambelli”, Università degli Studi di Salerno, 84084 Fisciano, SA, Italy; orcid.org/0000-0002-9382-2622

Rosaria Schettini – Dipartimento di Chimica e Biologia “A. Zambelli”, Università degli Studi di Salerno, 84084 Fisciano, SA, Italy; orcid.org/0000-0002-0515-424X

Giovanni Pierri – Dipartimento di Chimica e Biologia “A. Zambelli”, Università degli Studi di Salerno, 84084 Fisciano, SA, Italy; orcid.org/0000-0001-5433-6077

Maria Leda Marino – Dipartimento di Chimica e Biologia “A. Zambelli”, Università degli Studi di Salerno, 84084 Fisciano, SA, Italy

Irene Izzo – Dipartimento di Chimica e Biologia “A. Zambelli”, Università degli Studi di Salerno, 84084 Fisciano, SA, Italy; orcid.org/0000-0002-0369-0102

Francesco De Riccardis – Dipartimento di Chimica e Biologia “A. Zambelli”, Università degli Studi di Salerno, 84084 Fisciano, SA, Italy; orcid.org/0000-0002-8121-9463

Luca Bernardi – Department of Industrial Chemistry “Toso Montanari” & INSTM RU Bologna, Alma Mater Studiorum University of Bologna, 40136 Bologna, Italy; orcid.org/0000-0002-7840-3200

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.joc.0c00880>

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to the University of Salerno for financial support (FARB). We thank Francesca Di Riso for her valuable experimental contribution.

■ REFERENCES

- (a) León, A.; Del-Ángel, M.; Ávila, J. L.; Delgado, G. Phthalides: Distribution in Nature, Chemical Reactivity, Synthesis, and Biological Activity. In *Progress in the Chemistry of Organic Natural Products*; Vol. 104; Kinghorn, A. D.; Falk, H.; Gibbons, S.; Kobayashi, J., Eds.; Springer: Cham, 2017; pp. 127–246. (b) Lin, G.; Chan, S. S.-K.; Chung, H.-S.; Li, S.-L. Chemistry and Biological Activity of Naturally Occurring Phthalides. In *Studies in Natural Products Chemistry: Bioactive Natural Products*; Vol. 32, Part L; Atta-ur-Rahman, Ed.; Elsevier: Amsterdam, 2005; pp. 611–669. (c) Beck, J. J.; Chou, S.-C. The Structural Diversity of Phthalides from the Apiaceae. *J. Nat. Prod.* **2007**, *70*, 891–900.
- Karmakar, R.; Pahari, P.; Mal, D. Phthalides and Phthalans: Synthetic Methodologies and Their Applications in the Total Synthesis. *Chem. Rev.* **2014**, *114*, 6213–6284.
- (a) Mal, D. Applications of $[4 + 2]$ Anionic Annulations in the Total Synthesis. In *Anionic Annulations in Organic Synthesis*; Mal, D.; Elsevier, 2018; pp. 337–372. (b) Rathwell, K.; Brimble, M. A. Use of Stabilized Phthalide Anion Annulation Reactions in Synthesis: An Update. *Synthesis* **2007**, 643–662. (c) Cox, C.; Danishefsky, S. J. Synthesis of the Functionalized Tricyclic Core of Lactonamycin by Oxidative Dearomatization. *Org. Lett.* **2000**, *2*, 3493–3496. (d) Mkhize, S.; Suzuki, N.; Kurosawa, A.; Fujinami, M.; Chaicharoenpong, C.; Ishikawa, T. New Synthetic Procedure for 2-Aryl-1,4-naphthoquinone-1-oxime Methyl Ethers with Potent Antitumor Activity. *Synlett* **2014**, 25, 2059–2063. (e) Nicolaou, K. C.; Wang, Y.; Lu, M.; Mandal, D.; Pattanayak, M. R.; Yu, R.; Shah, A. A.; Chen, J. S.; Zhang, H.; Crawford, J. J.; Pasunoori, L.; Poudel, Y. B.; Chowdari, N. S.; Pan, C.; Nazeer, A.; Gangwar, S.; Vite, G.; Pitsinos, E. N. Streamlined Total Synthesis of Uncialamycin and Its Application to the Synthesis of Designed Analogues for Biological

Investigations. *J. Am. Chem. Soc.* **2016**, *138*, 8235–8246. (f) Nicolaou, K. C.; Lu, M.; Chen, P.; Shah, A. A. Practical Synthesis of *p*- and *o*-Amino- and Methoxyphenolic Anthraquinones. *Angew. Chem., Int. Ed.* **2015**, *54*, 12687–12691.

(4) (a) Vedantham, R.; Vetukuri, V. P. R.; Boini, A.; Khagga, M.; Bandichhor, R. Improved One-Pot Synthesis of Citalopram Diol and Its Conversion to Citalopram. *Org. Process Res. Dev.* **2013**, *17*, 798–805. (b) Eildal, J. N. N.; Andersen, J.; Kristensen, A. S.; Jørgensen, A. M.; Bang-Andersen, B.; Jørgensen, M.; Strømgaard, K. From the Selective Serotonin Transporter Inhibitor Citalopram to the Selective Norepinephrine Transporter Inhibitor Talopram: Synthesis and Structure-Activity Relationship Studies. *J. Med. Chem.* **2008**, *51*, 3045–3048. (c) Karmakar, R.; Pahari, P.; Mal, D. A synthetic route to 1,3-dihydroisobenzofuran natural products: the synthesis of methyl ethers of pestacin. *Tetrahedron Lett.* **2009**, *50*, 4042–4045. (d) Aggarwal, S.; Ghosha, N. N.; Anejab, R.; Joshi, H.; Chandra, R. A Convenient Synthesis of Aryl-Substituted *N*-Carbamoyl/*N*-Thiocarbamoyl Narcotic and Related Compounds. *Helv. Chim. Acta* **2002**, *85*, 2458–2462. (e) Huang, L.; Zhu, J.; Jiao, G.; Wang, Z.; Yu, X.; Deng, W. P.; Tang, W. Highly Enantioselective Rhodium-Catalyzed Addition of Arylboroxines to Simple Aryl Ketones: Efficient Synthesis of Escitalopram. *Angew. Chem., Int. Ed.* **2016**, *55*, 4527–4531.

(5) (a) Gadakh, S. K.; Sudalai, A. A simple and efficient synthesis of isocoumarins and alkylidenephthalides from 3-(1-hydroxycarboxy/alkyl)phthalides with a DEAD/PPH₃/TBHP system. *RSC Adv.* **2014**, *4*, 57658–57661. (b) Butin, A. V.; Dmitriev, A. S.; Uchuskin, M. G.; Abaev, V. T.; Trushkov, I. V. Simple and Convenient Synthesis of 4-Unsubstituted-3-(3-Oxoalkyl)isocoumarins. *Synth. Commun.* **2008**, *38*, 1569–1578. (c) Mali, R. S.; Babu, K. N. Reactions of 3-(1-Hydroxyalkyl)phthalides with Acids: Synthesis of (*Z*)-3-Alkylidenephthalides and 3-Alkyl-8-hydroxyisocoumarins. *J. Org. Chem.* **1998**, *63*, 2488–2492.

(6) (a) Cockcroft, X. I.; Dillon, K. J.; Dixon, L.; Drzewiecki, J.; Kerrigan, F.; Loh, V. M., Jr.; Martin, N. M. B.; Menear, K. A.; Smith, G. C. M. Phthalazinones 2: Optimisation and synthesis of novel potent inhibitors of poly(ADP-ribose)polymerase. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 1040–1044. (b) Napoletano, M.; Norcini, G.; Pellacini, F.; Marchini, F.; Morazzoni, G.; Ferlenga, P.; Pradella, L. Phthalazine PDE4 Inhibitors. Part 2: The Synthesis and Biological Evaluation of 6-Methoxy-1,4-disubstituted Derivatives. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 33–37.

(7) (a) Li, Y.; Linden, A.; Hesse, M. Total Synthesis of the Spermidine Alkaloid 13-Hydroxyisocyclocelabenzine. *Helv. Chim. Acta* **2003**, *86*, 579–591. (b) Hatano, M.; Yamakawa, K.; Kawai, T.; Horibe, T.; Ishihara, K. Enantioselective Cyanosilylation of Ketones with Lithium(I) Dicyanotrimethylsilicate(IV) Catalyzed by a Chiral Lithium(I) Phosphoryl Phenoxide. *Angew. Chem., Int. Ed.* **2016**, *55*, 4021–4025. (c) Hernández, E.; Vélez, J. M.; Vlaar, C. P. Synthesis of 1,4-dihydro-benzo[*d*][1,3]oxazin-2-ones from phthalides via an aminolysis-Hofmann rearrangement protocol. *Tetrahedron Lett.* **2007**, *48*, 8972–8975. (d) Kobayashi, S.; Tamanoi, H.; Hasegawa, Y.; Segawa, Y.; Masuyama, A. Divergent Synthesis of Bioactive Resorcinols Isolated from the Fruiting Bodies of *Hericium erinaceum*: Total Syntheses of Hericenones A, B, and I, Hericenols B–D, and Erinacerins A and B. *J. Org. Chem.* **2014**, *79*, 5227–5238. (e) Hobson, S. J.; Parkin, A.; Marquez, R. Oxidative Rearrangements of Isobenzofurans: Studies toward the Synthesis of the Ajudazols. *Org. Lett.* **2008**, *10*, 2813–2816.

(8) For efficient synthesis of 3,3-disubstituted phthalides through asymmetric catalysis see: (a) Okada, M.; Kaneko, K.; Yamanaka, M.; Shirakawa, S. BINOL-derived bifunctional sulfide catalysts for asymmetric synthesis of 3,3-disubstituted phthalides via bromolactonization. *Org. Biomol. Chem.* **2019**, *17*, 3747–3751. (b) Egami, H.; Asada, J.; Sato, K.; Hashizume, D.; Kawato, Y.; Hamashima, Y. Asymmetric Fluorolactonization with a Bifunctional Hydroxyl Carboxylate Catalyst. *J. Am. Chem. Soc.* **2015**, *137*, 10132–10135. (c) Han, X.; Dong, C.; Zhou, H.-B. C₃-Symmetric Cinchonine-Squaramide-Catalyzed Asymmetric Chlorolactonization of Styrene-

Type Carboxylic Acids with 1,3-Dichloro-5,5-dimethylhydantoin: An Efficient Method to Chiral Isochroman-1-ones. *Adv. Synth. Catal.* **2014**, *356*, 1275–1280. (d) Tanaka, K.; Osaka, T.; Noguchi, K.; Hirano, M. Rhodium-Catalyzed Asymmetric One-Pot Transesterification and [2 + 2 + 2]-Cycloaddition Leading to Enantioenriched 3,3-Disubstituted Phthalides. *Org. Lett.* **2007**, *9*, 1307–1310.

(9) (a) Sicignano, M.; Dentoni Litta, A.; Schettini, R.; De Riccardis, F.; Pierri, G.; Tedesco, C.; Izzo, I.; Della Sala, G. Highly Diastereoselective Crown Ether Catalyzed Arylogous Michael Reaction of 3-Aryl Phthalides. *Org. Lett.* **2017**, *19*, 4383–4386. (b) Sicignano, M.; Schettini, R.; Sica, L.; Pierri, G.; De Riccardis, F.; Izzo, I.; Maity, B.; Minenkov, Y.; Cavallo, L.; Della Sala, G. Unprecedented Diastereoselective Arylogous Michael Addition of Unactivated Phthalides. *Chem. – Eur. J.* **2019**, *25*, 7131–7141. (c) Heisler, T.; Janowski, W. K.; Prager, R. H.; Thompson, M. J. The Chemistry of Phthalidecarboxylic Acid. IV. Michael Additions to Conjugated Ketones. *Aust. J. Chem.* **1989**, *42*, 37–47.

(10) Luo, J.; Wang, H.; Zhong, F.; Kwiatkowski, J.; Xu, L.-W.; Lu, Y. Direct asymmetric Mannich reaction of phthalides: facile access to chiral substituted isoquinolines and isoquinolinones. *Chem. Commun.* **2012**, *48*, 4707–4709.

(11) (a) Luo, J.; Wang, H.; Zhong, F.; Kwiatkowski, J.; Xu, L.-W.; Lu, Y. Highly diastereoselective and enantioselective direct Michael addition of phthalide derivatives to nitroolefins. *Chem. Commun.* **2013**, *49*, 5775–5777. (b) Luo, J.; Jiang, C.; Wang, H.; Xu, L.-W.; Lu, Y. Direct asymmetric Michael addition of phthalide derivatives to chalcones. *Tetrahedron Lett.* **2013**, *54*, 5261–5265. (c) Wang, J.; Li, X.; Cheng, J.-P. Quinine-derived thiourea promoted enantioselective Michael addition reactions of 3-substituted phthalides to maleimides. *Sci. Chin. Chem.* **2019**, *62*, 649–652.

(12) (a) Zhong, F.; Luo, J.; Chen, G.-Y.; Dou, X.; Lu, Y. Highly Enantioselective Regiodivergent Allylic Alkylations of MBH Carbonates with Phthalides. *J. Am. Chem. Soc.* **2012**, *134*, 10222–10227. (b) Hu, Z.-P.; Zhuang, Z.; Liao, W.-W. Asymmetric Synthesis of Dihydronaphthoquinones Containing Adjacent Stereocenters via a Sulfa-Michael Addition Triggered Ring-Expansion Approach. *J. Org. Chem.* **2015**, *80*, 4627–4637. (c) Zhuang, Z.; Hu, Z.-P.; Liao, W.-W. Asymmetric Synthesis of Functionalized Dihydronaphthoquinones Containing Quaternary Carbon Centers via a Metal-Free Catalytic Intramolecular Acylcyanation of Activated Alkenes. *Org. Lett.* **2014**, *16*, 3380–3383.

(13) For quinine-catalyzed reaction of 3-cyanophthalides and MBH carbonates see: Liu, W.; Hu, Z.-P.; Yan, Y.; Liao, W.-W. Highly diastereo- and enantioselective construction of phthalideoxindole hybrids bearing vicinal quaternary chiral centers via an organocatalytic allylic alkylation. *Tetrahedron Lett.* **2018**, *59*, 3132–3135.

(14) For some examples of non-stereoselective introduction of functionalized electrophilic groups at the γ -position of 3-substituted phthalides, see: (a) Kraus, G. A.; Sugimoto, H. Michael Additions of Functionalized Phthalides. *Synth. Commun.* **1977**, *7*, 505–508. (b) Li, T. t.; Walsgrove, T. C. A Convergent Regiospecific Route to the Aklavinone Ring System. *Tetrahedron Lett.* **1981**, *22*, 3741–3744. (c) Chiefari, J.; Janowski, W. K.; Prager, R. H. The Chemistry of Phthalide-3-carboxylic Acid. V. The Reaction of the Acid with Imines. *Aust. J. Chem.* **1989**, *42*, 49–60. (d) Hassan, N. P. S.; Naysmith, B. J.; Sperry, J.; Brimble, M. A. Formal synthesis of nanaomycin D via a Hauser-Kraus annulation using a chiral enone-lactone. *Tetrahedron* **2015**, *71*, 7137–7143. (e) Hu, S.; Teng, L.; Zhang, J.; Meng, J.; Zhuang, Z.; Du, M.; Wang, D.; Liao, W. Synthesis of a Novel Compound and Mediated Neuro-protection in Differentiated PC12 Cells via Mitochondria Related Pathway. *Chem. J. Chin. U.* **2015**, *36*, 2322–2328.

(15) ((a)) *Asymmetric Phase Transfer Catalysis*; Maruoka, K.; Eds.; Wiley-VCH: Weinheim, 2008. (b) Shirakawa, S.; Maruoka, K. Recent Developments in Asymmetric Phase-Transfer Reactions. *Angew. Chem., Int. Ed.* **2013**, *52*, 4312–4348. (c) Ooi, T.; Maruoka, K. Recent Advances in Asymmetric Phase-Transfer Catalysis. *Angew. Chem., Int. Ed.* **2007**, *46*, 4222–4266. (d) Kaneko, S.; Kumatabara, Y.; Shirakawa, S. A new generation of chiral phase-transfer catalysts. *Org.*

Biomol. Chem. **2016**, *14*, 5367–5376. (e) Tan, J.; Yasuda, N. Contemporary Asymmetric Phase Transfer Catalysis: Large-Scale Industrial Applications. *Org. Process Res. Dev.* **2015**, *19*, 1731–1746. (f) Schettini, R.; Sicignano, M.; De Riccardis, F.; Izzo, I.; Della Sala, G. Macrocyclic Hosts in Asymmetric Phase-Transfer Catalyzed Reactions. *Synthesis* **2018**, *50*, 4777–4795. (g) Novacek, J.; Waser, M. Bifunctional Chiral Quaternary Ammonium Salt Catalysts: A Rapidly Emerging Class of Powerful Asymmetric Catalysts. *Eur. J. Org. Chem.* **2013**, 637–648. (h) Qian, D.; Sun, J. Recent Progress in Asymmetric Ion-Pairing Catalysis with Ammonium Salts. *Chem. – Eur. J.* **2019**, *25*, 3740–3751.

(16) For an extensive screening of catalysts and reaction conditions see the [Supporting Information](#).

(17) (a) Lygo, B.; Allbutt, B.; James, R. S. Identification of a highly effective asymmetric phase-transfer catalyst derived from α -methylnaphthylamine. *Tetrahedron Lett.* **2003**, *44*, 5629–5632. (b) Lygo, B.; Allbutt, B.; Beaumont, D. J.; Butt, U.; Gilks, J. A. R. Synthesis and Evaluation of Chiral Dibenzazepinium Halide Phase-Transfer Catalysts. *Synlett* **2009**, 675–680.

(18) CCDC 1978176 (17aa) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

(19) We also examined the reaction with aliphatic alkylating agents, obtaining only complicated mixture of products.

(20) (a) Kamachi, T.; Yoshizawa, K. Enantioselective Alkylation by Binaphthyl Chiral Phase-Transfer Catalysts: A DFT-Based Conformational Analysis. *Org. Lett.* **2014**, *16*, 472–475. (b) Petrova, G. P.; Li, H.-B.; Maruoka, K.; Morokuma, K. Asymmetric Phase-Transfer Catalysis with Homo- and Heterochiral Quaternary Ammonium Salts: A Theoretical Study. *J. Phys. Chem. B* **2014**, *118*, 5154–5167.

(21) Wagner, H.; Burghart, J. Macrocyclische Spermidinalkaloide aus *Maytenus mossambicensis* (KLOTZSCH) BLAKELOCK. *Helv. Chim. Acta* **1982**, *65*, 739–752.

(22) Shipilovskikh, S. A.; Vaganov, V. Y.; Denisova, E. I.; Rubtsov, A. E.; Malkov, A. V. Dehydration of Amides to Nitriles under Conditions of a Catalytic Appel Reaction. *Org. Lett.* **2018**, *20*, 728–731.

(23) The synthesis of another formal precursor of 13-hydroxyisocyclocelabenzine, namely the primary amine corresponding to **18**, has also been reported, although as the (*S*)-enantiomer, potentially leading to the (13*S*)-configured dihydroisoquinolinone moiety (ref 7b).

(24) Liu, Y.; Provencher, B. A.; Bartelson, K. J.; Deng, L. Highly enantioselective asymmetric Darzens reactions with a phase transfer catalyst. *Chem. Sci.* **2011**, *2*, 1301–1304.

(25) Zhang, Y.-H.; Shi, B.-F.; Yu, J.-Q. Palladium(II)-Catalyzed ortho Alkylation of Benzoic Acids with Alkyl Halides. *Angew. Chem., Int. Ed.* **2009**, *48*, 6097–6100.

(26) Tatsugi, J.; Hara, T.; Izawa, Y. Photochemical Behavior of Indane-1,2,3-trione in Degassed Alcoholic Solutions: Formation of 3-Substituted Phthalides. *Chem. Lett.* **1997**, *26*, 177–178.

(27) Ishibashi, M.; Wagner, A.; Mioskowski, C.; Sylvain, C. Preparation of Carboxytetrahydrofuranone Derivatives as Selective Preventives/Remedies for Progressive Lesions after Organ Damage. *Int. Pat. Appl.* WO2001072730A1, 2001.