

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Carbonyl-isocyanide mono-substitution in [Fe2Cp2(CO)4]: A re-visitation

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version: Biancalana L., Ciancaleoni G., Zacchini S., Pampaloni G., Marchetti F. (2020). Carbonyl-isocyanide mono-substitution in [Fe2Cp2(CO)4]: A re-visitation. INORGANICA CHIMICA ACTA, 517, 1-9 [10.1016/j.ica.2020.120181].

Availability: This version is available at: https://hdl.handle.net/11585/793274 since: 2021-01-29

Published:

DOI: http://doi.org/10.1016/j.ica.2020.120181

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:L. Biancalana, G. Ciancaleoni, S. Zacchini, G. Pampaloni, F. Marchetti, "Carbonyl-isocyanide mono-<br/>substitution in [Fe2Cp2(CO)4]: A re-visitation", *Inorg. Chim. Acta*, **2020**, *517*, 120181.Thefinalpublishedversionisavailableonlineat:https://doi.org/10.1016/j.ica.2020.120181

Rights / License: Licenza per Accesso Aperto. Creative Commons Attribuzione - Non commerciale - Non opere derivate 4.0 (CCBYNCND)

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

# Carbonyl-Isocyanide Mono-Substitution in[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>]: a Re-Visitation<sup>†</sup>

Lorenzo Biancalana,<sup>a</sup> Gianluca Ciancaleoni,<sup>a</sup> Stefano Zacchini,<sup>b</sup> Guido Pampaloni<sup>a,\*</sup> and Fabio

Marchetti<sup>a,\*</sup>

<sup>a</sup>University of Pisa, Department of Chemistry and Industrial Chemistry, Via G. Moruzzi 13, I-56124

Pisa, Italy.

<sup>b</sup>University of Bologna, Department of Industrial Chemistry "Toso Montanari", Viale Risorgimento 4,

I-40136 Bologna, Italy.

<sup>†</sup> This work is dedicated to Prof. Maurizio Peruzzini, on the occasion of his 65<sup>th</sup> birthday

\* Authors to whom correspondence should be addressed:

guido.pampaloni@unipi.it, fabio.marchetti1974@unipi.it

#### Abstract

The reactions of  $[Fe_2Cp_2(CO)_4]$  with a series of isocyanides, CNR, were conducted in acetonitrile and afforded, after a thermal treatment, the mono-isocyanide derivatives  $[Fe_2Cp_2(CO)_3(CNR)]$  [R = 1H-indol-5-yl, 1; CH<sub>2</sub>P(O)(OEt)<sub>2</sub>, 2; Cy = C<sub>6</sub>H<sub>11</sub>, 3; 4-C<sub>6</sub>H<sub>4</sub>OMe, 4; Xyl = 2,6-C<sub>6</sub>H<sub>3</sub>Me<sub>2</sub>, 5; Me, 6; 2-naphthyl, 7; Bn = CH<sub>2</sub>Ph, 8]. In order to avoid multiple substitution, the diiron reactant was used in a molar excess with respect to the isocyanide (1.6 equivalents; 1.1 for the synthesis of 8). The products were separated from unreacted  $[Fe_2Cp_2(CO)_4]$  by chromatography or via reversible protonation, and finally isolated in 50-83 % yields. IR and NMR spectroscopy indicate that the isocyanide ligand is bridging coordinated in 2 and 7, terminal in 3 and 5, while in the remaining cases a mixture of terminal- and bridging-CNR isomers is obtained. The molecular structure of 5 was ascertained by X-ray diffraction. In general, the coordination mode of the isocyanide is scarcely influenced by the environment (solvents with different polarities, solid state). Sluggish partial isocyanide migration from terminal to bridging position was recognized for 3 and 5 upon heating in refluxing toluene, a process which was reproduced by DFT calculations.

Keywords: diiron complexes, carbonyl ligand, isocyanide ligands, substitution reaction, DFT calculations

#### Introduction

Isocyanides are ubiquitous ligands in coordination chemistry [<sup>1</sup>], and their incorporation in organometallic structures is often achieved through the replacement of carbon monoxide ligands [<sup>2</sup>]. Cyclopentadienyl iron(II) dicarbonyl dimer, [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (Cp =  $\eta^5$ -C<sub>5</sub>H<sub>5</sub>), is an easily available and robust platform, for which the reactivity with isocyanides aroused the interest of several research groups for more than 20 years in the last century [<sup>3</sup>]. In principle, the resulting complexes may exist as

mixtures of isomers, originated from cis or trans mutual orientation of the Cp ligands and terminal or bridging coordination of the isocyanide ligand(s); this situation is often complicated by the occurrence of interconversion processes in solution [<sup>4</sup>, <sup>5</sup>]. Various monosubstituted derivatives were afforded in ca. 60% yields from the reaction of  $[Fe_2Cp_2(CO)_4]$  with a slight molar excess of a variety of alkylisocyanides in benzene at room temperature, Scheme 1A [<sup>6</sup>]. Similarly, the di-isocyanides CN(CH<sub>2</sub>)<sub>n</sub>NC (n = 2,3,4,6) were used to build dinuclear structures,  $[{Fe_2Cp_2(CO)_3}_2{\mu-CN(CH_2)_nNC}]$  [<sup>7</sup>], and it was found that the terminal coordination of the bidentate isocyanide is favored on increasing the length of the alkyl chain. The synthesis of  $[Fe_2Cp_2(CO)_3(CN^tBu)]$  was alternatively carried out in the presence of a catalytic amount of BEt<sub>3</sub> [5c, <sup>8</sup>]: in this case, a radical chain mechanism was proposed, essentially based on the transfer of an ethyl group to a CO ligand, favoring its extrusion.

A series of di-alkyl-isocyanide complexes were prepared in approximately 50% yields by reacting  $[Fe_2Cp_2(CO)_4]$  with a large excess of the organic reactant in benzene at reflux temperature, Scheme 1B  $[^9, ^{10}]$ .Few tri-isocyanide species were obtained upon prolonged heating of these mixtures and finally isolated in low yield after work-up [10].In order to realize the full substitution of the four carbonyl ligands in  $[Fe_2Cp_2(CO)_4]$ , the reactions with an eight-fold excess of methyl- and ethyl-isocyanides required to be conducted in refluxing xylene, Scheme 1C  $[^{11}]$ ; under these experimental conditions, only the bis-substituted product was obtained from benzyl-isocyanide.

In general, aryl-isocyanides are more reactive towards  $[Fe_2Cp_2(CO)_4]$ , and the isolation of monoderivatives is tricky; interestingly, an IR spectroscopy study on the 1:1 reaction with CNXyl (Xyl = 2,6-C<sub>6</sub>H<sub>3</sub>Me<sub>2</sub>) indicated that  $[Fe_2Cp_2(CO)_3(CNXyl)]$  adds further CNXyl more rapidly than  $[Fe_2Cp_2(CO)_4]$  [<sup>12</sup>].Manning and co-workers described the photolytic reaction of  $[Fe_2Cp_2(CO)_4]$  witha three-fold excess of aryl-isocyanides in THF solution in the light, affording either  $[Fe_2Cp_2(CO)_3(CNR)]$  or  $[Fe_2Cp_2(CO)_2(CNR)_2]$  in 60-70% yields, depending on the steric hindrance of the aryl group [<sup>13</sup>,<sup>14</sup>], Scheme 1D. For instance, the former type of product is almost exclusivelygeneratedfrom2,6dimethylphenyl-isocyanide, while the latter from phenyl-isocyanide (Scheme 1D).The high temperature reactions of  $[Fe_2Cp_2(CO)_4]$  with a four-fold excess of CNAr (Ar = Ph, Xyl) proceed smoothly to give the tetra-isocyanides in over 60% yields, Scheme 1E [12].



**Scheme 1**. Former studies on the substitution reactions of  $[Fe_2Cp_2(CO)_4]$  with isocyanides. References: A [6], B [9,10], C [11], D [13,14], E [12].

The structures of some derivatives were ascertained by single crystal X-ray diffraction. It can be generally observed that the isocyanide moiety, when terminally coordinated to one iron center, typically exhibits a linear C-N-R geometry [5c, 12, <sup>15</sup>]; otherwise, bridging isocyanides substantially deviate from linearity and feature an elongated carbon-nitrogen bond, thus suggesting a substantial imine character which is in alignment with IR data [9,<sup>16</sup>]. As a representative example, a view of the X-ray structure of *cis*-[Fe<sub>2</sub>Cp<sub>2</sub>(CO)( $\mu$ -CO)(CN<sup>i</sup>Pr)( $\mu$ -CN<sup>i</sup>Pr)], containing one terminal and one bridging isopropyl-isocyanide units, is shown in Figure 1 with relevant bonding parameters listed in the caption [10].



**Figure 1**. A) View of the X-ray structure of *cis*-[Fe<sub>2</sub>Cp<sub>2</sub>(CO)( $\mu$ -CO)(CN<sup>i</sup>Pr)( $\mu$ -CN<sup>i</sup>Pr)], from reference [10]; Selected bond lengths (Å) and angles (°): Fe1-Fe2 2.521(7), Fe1-C21.89(3), Fe2-C2 1.88(4), Fe1-C1 2.01(3), Fe2-C2 1.86(3), C2-N1 1.26(3), N1-C5 1.45(4), Fe2-C4 1.76(3), C4-N2 1.22(3), N2-C8 1.55(4), C2-N1-C5 128(3), Fe2-C4-N2 179(3), C4-N2-C8 171(3).

Compounds derived from  $[Fe_2Cp_2(CO)_4]$  by carbon monoxide/isocyanide single replacement are particularly of interest, since the subsequent alkylation of the nitrogen provides the access to cationic aminocarbyne complexes [<sup>17</sup>]. This transformation usually converges to a unique isomeric form, with the aminocarbyne ligand in bridging position and the Cp rings adopting the cis configuration. Such aminocarbyne compounds exhibit a rich chemistry and represent useful starting materials for the preparation of potential anticancer drugs [<sup>18</sup>]. Recently, we described a straightforward and general method to synthesize diiron aminocarbyne compounds (Scheme 2) [17], whereby the preliminary CO/CNR substitution(Scheme 2, step i) is selective upon reaction of a variety of alkyl- and arylisocyanides with an excess (ca. 1.6 equivalents) of  $[Fe_2Cp_2(CO)_4]$  in acetonitrile. The elaborated chromatographic separation of the mono-isocyanide adducts from the unreacted tetracarbonyl species is avoided, and the cationic products are finally isolated in yields up to 95% (Scheme 2, step ii).



**Scheme 2**. Synthesis of cationic diiron  $\mu$ -aminocarbyne complexes from [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] through the preliminary formation of neutral mono-isocyanide adducts. R, reaction temperature: Me, reflux; CH<sub>2</sub>Ph, reflux; 2,6-C<sub>6</sub>H<sub>3</sub>Me<sub>2</sub> (Xyl), RT; 2,6-C<sub>6</sub>H<sub>3</sub>MeCl, RT; 2-naphthyl, RT.

Herein, we present a systematic study focused on the mono-isocyanide adducts [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>(CNR)]: using the optimized synthetic approach (Scheme 2, step i), a series of compounds (including two unprecedented compounds and three isolated for the first time) has been prepared, isolated and characterized by IR and <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy. With the assistance of DFT calculations, the relative stability of the isomeric forms is elucidated and rationalized.

There is a debate on the existence of an iron-iron bond in  $[Fe_2Cp_2(CO)_4]$  [<sup>19</sup>], which might be reasonably extended to  $[Fe_2Cp_2(CO)_3(CNR)]$ ; however, here we will keep the bond in the pictures, basically because we consider this representation more in alignment with the chemistry of the complexes.

#### **Results and discussion**

#### 1) Synthesis and isolation of complexes

The reactions of diethyl isocyanomethylphosphonate and 1H-indol-5-yl isocyanide with ca. 1.6 equivalents of  $[Fe_2Cp_2(CO)_4]$  in acetonitrile, proceeded with selective carbonyl mono-substitution to give the novel compounds 1-2.By means of the same procedure, we prepared the previously reported 3-8 (Scheme 3a), of which4, 5 and 7 were isolated for the first time. Compound 8 was synthesized using only a slight excess of  $[Fe_2Cp_2(CO)_4]$  (1.1 eq), following the previous indication that benzyl isocyanide has a low tendency to poly-substitution  $[10, ^{20}]$ . All the products underwent a thermal treatment (T  $\geq$  90 °C) for at least 3 hours, to ensure complete consumption of the isocyanide, and were obtained in admixture with unreacted  $[Fe_2Cp_2(CO)_4]$ . The separation of1, 2 and 4, containing N/O functions in the isocyanide group, was easily achieved on neutral alumina or Florisil<sup>®</sup> columns(see Experimental for details). On the other hand, chromatographic separation of the remaining, less polar compounds proved

to be more challenging, and only **3** was isolated in low yield. Therefore, we moved towards a different strategy, relying on the reversible protonation of the isocyanide ligand (Scheme 3b)[<sup>21</sup>].

Treatment of  $Et_2O$  solutions of **3-8** with aqueous tetrafluoroboric acid afforded the insoluble aminocarbyne salts  $[Fe_2Cp_2(CO)_3(CNHR)]BF_4$ , which could be separated from  $[Fe_2Cp_2(CO)_4]$  by filtration. Next, triethylamine was employed to regenerate the isocyanide function and the neutral compounds were finally purified by alumina chromatography. Compounds **1-8** were finally obtained as dark red-purple solids which appeared to be rather inert to air/moisture; they were kept under N<sub>2</sub> for long-term storage. Conversely, solutions of **1-8**in any common organic solvent underwent appreciable degradation when exposed to ambient air over few hours, with precipitation of a rusty-brown solid / film over glassware.



**Scheme 3**. (a) Synthesis of diiron bis-cyclopentadienyl isocyanide complexes **1-8** by CO/CNR exchange from  $[Fe_2Cp_2(CO)_4]$ ; (b) Subsequent protonation/deprotonation steps for the isolation of **3-8**.

#### 2) IR and NMR spectroscopy and X-ray diffraction

In principle, four isomers are predictable for the mono-isocyanide adducts **1-8** (see Introduction), and these are shown in Figure 2. According to the mechanism based on NMR studies and proposed by Adams and Cottons for  $[Fe_2Cp_2(CO)_4]$ , cis and trans isomers may interconvert in solution through a bridge-opened structure and rotation around the Fe-Fe bond[<sup>22</sup>]. It was quantified that approximately 10% of  $[Fe_2Cp_2(CO)_4]$  is present in solution in such form displaying all terminal ligands [4]. Adams

and Cotton expanded their concepts to  $[Fe_2Cp_2(CO)_3(CNR)]$  complexes [5c], for which the bridgeopened structure (structure **bo** in Figure 2) gives rise to several conformers; the activation energy for the bridge opening and closing was determined to be 9.4 kcal mol<sup>-1</sup> in the case of**6** in CD<sub>2</sub>Cl<sub>2</sub> solution

[4].



**Figure 2**. Structures of the four possible isomers of  $[Fe_2Cp_2(CO)_3(CNR)]$ , and the bridge-opened intermediate for isomer interconversion in solution.

It was previously observed that the isomeric composition in this class of complexes is variable in different solvents, the more polar solvents usually favoring the more polar isomers (especially cis vs. trans) [5b]. Thus, we collected the IR spectra of **1-8** in MeCN ( $\mu$  = 3.92 D), CH<sub>2</sub>Cl<sub>2</sub>( $\mu$  = 1.60 D) and toluene ( $\mu$  = 0.36 D) solution as well as in the solid state. For each compound, the coordination mode of the isocyanide can be recognized from the infrared bands associated to the CN stretching vibration. The latter falls in within the intervals 2050-2150 and 1690-1720 cm<sup>-1</sup> for terminal and bridging isocyanides, respectively. Note that values related to terminal isocyanides are 10-30 cm<sup>-1</sup> lower than those available for the corresponding non-coordinated isocyanides (Table 1), indicating the occurrence of an appreciable Fe<sup>1</sup> to ligand backdonation in **1-8**. With reference to CH<sub>2</sub>Cl<sub>2</sub> solutions (Table 1), the isocyanide in **2**and**7**manifests a strong tendency to occupy a bridging site, whereas in **3** and **5** the terminal coordination is largely favored. The result obtained for**3** and**5** is coherent with former findings [6, 12]. In the other cases (**1**, **4**, **6** and **8**), a mixture of terminal- and bridging-isocyanide isomers in comparable amounts is inferred. The result concerning**6** is in accordance with the former literature findings (vide infra) [4].

**Table 1**. IR data related to the isocyanide moiety in  $[Fe_2Cp_2(CO)_3(CNR)]$  complexes(CH<sub>2</sub>Cl<sub>2</sub> solution) and noncoordinated CNR molecules.

[Fe <sub>2</sub> Cp <sub>2</sub> (CO) <sub>3</sub> (CNR)]					CNR
R	Cpd. –	Ũ / cm <sup>−1[a]</sup>		Isocyanide	$\tilde{u}(\mathbf{C}=\mathbf{N})/\mathbf{cm}^{-1}[\mathbf{a}]$
		t-CN	μ-ϹΝ	coordination mode <sup>[b]</sup>	
1H-indol-5-yl	1	2106	1703	bridging > terminal	2117(KBr) <sup>23</sup>
CH <sub>2</sub> PO <sub>3</sub> Et <sub>2</sub>	2	-	1690	bridging	2140 <sup>24</sup>
C <sub>6</sub> H <sub>11</sub>	3	2128	-	terminal	2144 <sup>25</sup>
4-C <sub>6</sub> H <sub>4</sub> OMe	4	2102	1698	bridging > terminal	2125 <sup>26</sup>
Xyl	5	2090	-	terminal	2120
Ме	6	2153	1719	terminal > bridging	2166 (gas) <sup>27</sup>
2-naphthyl	7	-	1694	bridging	2124(KBr) <sup>28</sup>
CH <sub>2</sub> Ph	8	2131	1710	bridging > terminal	2152 <sup>29</sup>

[a] **1-8**:  $CH_2Cl_2$  solution; non-coordinated CNR: neat liquid/solid state,unless otherwise specified. [b] According to the relative transmittance of t-CN (**ct** + **tt** isomers in Fig. 2) and  $\mu$ -CN (**cb** + **tb** isomers in Fig. 2) stretching bands.

X-ray quality crystals of **5-ct** were collected from a .... solution, and a view of the structure of this compound in shown in Figure 3, with relevant bonding parameters given in the caption.

**5-ct** adds to the very few [Fe<sub>2</sub>Cp<sub>2</sub>(CNR)(CO)(μ-CO)<sub>2</sub>] complexes structurally characterized [ref: R. D. Adams, F. A.

Cotton, J. M. Troup, Inorg. Chem. 1974, 13, 257; I. L. C. Campbell, F. S. Stephens, J. Chem. Soc. Dalton Trans.

1975, 982]. Its molecular structure and bonding parameters are comparable to those previously reported.

Thus, the Cp ligands adopt a *cis* geometry, and the two  $\mu$ -CO ligands display a marked asymmetry [Fe(1)-C(2)]

1.947(3), Fe(2)-C(2) 1.893(3), Fe(1)-C(3) 1.941(3), Fe(2)-C(3) 1.911(3)] in view of the greater  $\pi$ -acidity of t-CO

compared to t-CNR. The C(11)-N(1)-C(12) [169.6(4)°] group is almost linear as expected for a terminal

isonitrile.



**Figure 3**. Molecular structure of cis-[Fe<sub>2</sub>Cp<sub>2</sub>(CNXyl)(CO)<sub>2</sub>( $\mu$ -CO)], **5a-ct**, with labelling. Displacement ellipsoids are at the 30% probability level. H-atoms have been omitted for clarity. Selected bond distances (Å) and angles (°): Fe(1)-Fe(2) 2.5267(7), Fe(1)-C(1) 1.752(4), Fe(1)-C(2) 1.947(3), Fe(2)-C(2) 1.893(3), Fe(1)-C(3) 1.941(3), Fe(2)-C(3) 1.911(3), Fe(2)-C(11) 1.816(3), C(1)-O(1) 1.142(4), C(2)-O(2) 1.161(4), C(3)-O(3) 1.164(4), C(11)-N(1) 1.161(4), N(1)-C(12) 1.398(4), Fe(1)-C(1)-O(1) 178.0(3), Fe(2)-C(11)-N(1) 177.1(3), C(11)-N(1)-C(12) 169.6(4), Fe(1)-C(2)-Fe(2) 82.29(13), Fe(1)-C(3)-Fe(2) 81.99(13).

In general, the coordination mode adopted by the isocyanide ligand is negligibly affected by the environment, and the IR spectra in the solid-state (Figures S1-S8 in the SI) and in acetonitrile and toluene solutions (Figures S9-S16) are substantially comparable to those recorded in CH<sub>2</sub>Cl<sub>2</sub>. Exceptions are discussed in the following. First, despite the xylyl-isocyanide ligand in **5** is always found terminal, the number and the relative intensity of CO/CN absorptions are variable. Two bands of comparable intensity due to terminal CNXyl (at 2136 and 2082cm<sup>-1</sup>) and other two bands due to bridging CO (at 1775 and 1756 cm<sup>-1</sup>) appear in acetonitrile solution. For sake of comparison, the IR spectrum of **5** in the solid state displays only one CNXyl band (at 2081 cm<sup>-1</sup>) and one CO band (at 1943 cm<sup>-1</sup>), and the same qualitative pattern was found in toluene and dichloromethane. Presumably, both the **tt** and **ct** isomers of **5** are stable in acetonitrile, while the **tt** structure is dominant in less polar solvents and in the solid state (where the **ct** isomer is also present and has been detected by X-ray diffraction, see above). The second exception to the general trend was observed for **4** and **8**: while these complexes exist as mixtures of terminal- and bridging-isocyanide isomers in solution (with prevalence of the latter), the disappearance of the band around 2100 cm<sup>-1</sup>in the solid state indicates the absence of **ct** and **tt** isomers (Figures 2 and 4). This feature seems imputable to solid state effects, on considering that no marked differences were observed in the relative intensities of terminal and bridging isocyanide absorptions on changing the solvent polarity.



**Figure 4**. Comparison of IR spectra (1500-2300 cm<sup>-1</sup>) in CH<sub>2</sub>Cl<sub>2</sub> solution (blue line) and in the solid state (red line) for [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>{CN(4-C<sub>6</sub>H<sub>4</sub>OMe)}], **4**(**a**) and [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>(CNCH<sub>2</sub>Ph)], **8** (**b**).

A previous NMR study assessed the percentage of the isomers of  $6 \text{ at } -26 \text{ °C in } \text{CD}_2\text{Cl}_2 \text{ solution, resulting 77\%}$ (ct), 2% (tt), 16% (cb), 5% (tb) [4]: a strong prevalence is evidenced of cis isomers (*ct* + *cb*, 93%) and terminal coordination of methylisocyanide (79%). In the low temperature <sup>13</sup>C NMR spectrum, the resonances due to terminal and bridging coordination of CNMe are diagnostic, falling at 160 and 250 ppm, respectively. On the other hand, the carbonyl ligands in the same complex resonate at ca. 210 and ca. 280 ppm when occupying terminal and bridging sites.

The NMR spectra of **1-8**were recorded in CDCl<sub>3</sub> at room temperature, and they generally display broadenedsignals indicative of fluxionality (see Figure 2). Overall, the detected <sup>13</sup>C NMR signals for the isocyanide ligands(Table 2) are substantially in alignment with the IR features; in particular, **2** and **7** exhibits one broad signal due to the bridging isocyanide at ca. 260 ppm, while the signal for the isocyanide in **3** and **5** falls at around 170 ppm as expected for terminal coordination. The <sup>13</sup>C NMR chemical shifts of the compounds existing as mixtures of terminal- and bridging-isocyanide isomers (**1**, **4**, **6** and **8**) are not averaged, but consistent with the predominant isocyanide coordination mode as assessed by IR spectroscopy. For instance, the CN resonance falls at *ca*. 250 ppm for **1**, **4**, **8** and *ca*. 160 ppm for **6**. For the latter compound, two methyl resonances are visible (at 47 and 30 ppm), as in the low-temperature spectrum [4].

**Table 2**. <sup>13</sup>C NMR spectral assignments for**ct** and **cb** isomers of  $[Fe_2Cp_2(CO)_3(CNMe)]$  (6) according to the literature<sup>[a]</sup> and comparison of selected <sup>13</sup>C{<sup>1</sup>H}NMR data for complexes **1-8**.<sup>[b]</sup>

280 ppm	Compound	R —	δ / ppm			
<sup>213</sup> ppm C N <sup>30</sup> ppm			μ-CO	t-CO	<u>C</u> NR	Ср
Fe—Fe <sup>159</sup> ppm	<b>1</b> <sup>[c]</sup>	1H-indol-6-yl	274	214	249	89br
86, 87	2	$CH_2PO_3Et_2$	272	213, 210	260	90, 88
0 ppm	3	$C_6H_{11}$	282	217, 213	162	87, 86sh
46 ppm	4	4-C <sub>6</sub> H <sub>4</sub> OMe	273	217br, 211	254	89br, 88
N 211 ppm    248	5	Xyl	278br	215-212br	176br	88
OC C ppm CO	6	Me	281	213, 211	163	87
Fé—Fé	7	2-naphthyl	272	213br, 211	255	90br, 88
C 87, 88 o n.d.	8	CH₂Ph	280br, 273	215-212br, 211	251	88

[a]Spectra recorded in CD<sub>2</sub>Cl<sub>2</sub> at – 26 °C [4];[b]Spectra recorded in CDCl<sub>3</sub>at 298 K unless otherwise specified;[c]CD<sub>3</sub>CN solution.

# 3) DFT calculations

The relative stability of the possible isomers of complexes **1-8** (Figure 2)was evaluated by DFT calculations with reference to dichloromethane medium, and the results are reported in Table 3.

Table 3. Relative energies of the four isomers predictable for complexes 1-8(C-PCM,CH<sub>2</sub>Cl<sub>2</sub>).

Complex	R			$\mathbf{v}_{\mathbf{r}}^{\mathbf{r}} = \mathbf{v}_{\mathbf{r}}^{\mathbf{r}} + \mathbf{v}_{\mathbf$	oc c c c c c c c c c c c c c c c c c c
1	1H-indol-6-yl	0.0	0.4	2.5	2.8
2	$CH_2PO_3Et_2$	0.0	1.3	1.8	3.6
3	C <sub>6</sub> H <sub>11</sub>	0.0	1.1	1.0	1.6
4	4-C <sub>6</sub> H <sub>4</sub> OMe	0.0	0.4	2.8	3.2
5	Xyl	0.0	0.4	0.9	1.3
6	Me	0.0	1.1	0.9	1.6

7	2-naphthyl	0.0	0.5	4.1	4.4
8	Bn	0.0	2.4	2.4	3.0

The **cb** isomer is always the most stable one, although in some cases the energy difference with respect to the other isomers is low (note that differences smaller than 0.5 kcal mol<sup>-1</sup> exceed the accuracy of the DFT methods). For instance, the **cb** and **tb** isomers of **1**, **4**, **5** and **7** are practically isoenergetic. The geometries of the four isomers of **6** are shown in Figure 5, while those ones of **1**-**5** and **7**-**8** are supplied as Supporting Information.



**Figure 5**. DFT-optimized geometries of the four isomers of **6**. Relative energies are indicated in brackets and expressed in kcalmol<sup>-1</sup>.

Overall, the calculations highlight the higher stability provided by the bridging coordination of the isocyanide ligand. This contrasts especially with the experimental evidence that **3** and **5**are obtained almost exclusively as terminal-isocyanide structures through a synthetic procedure including high temperature treatment in a high polar solvent (refluxing acetonitrile,  $T \ge 90^{\circ}$ C). It is noteworthy that additional calculations pointed out that the relative energies of the isomers of **5** do not significantly change in acetonitrile medium with respect to dichloromethane (variations are within 0.1 kcal mol<sup>-1</sup>). When toluene solutions of **3** and **5** were heated at reflux (T ~110 °C), a sluggish, partial conversion into the predicted stable "bridging" products was ascertained by IR spectroscopy, and the new IR patterns were maintained at ambient temperature (see Experimental and figures S41-S42 for details). It is reasonable that, in general, the kinetic product of the CO/CNR substitution features a

terminal isocyanide (structures **ct** and **tt** in Figure 2), on considering that the binding of the terminal CO ligands in  $[Fe_2Cp_2(CO)_4]$  is weaker than that of the bridging ones. It seems that the subsequent bridge closing from **bo**, placing the isocyanide group in bridging position to afford either **cb** or **tb**, isinhibited with encumbered R groups such as cyclohexyl and 2,6-dimethylphenyl.

#### Conclusions

A series of mono-isocyanide complexes obtained by substitution reactions from  $[Fe_2Cp_2(CO)_4]$  has been synthesized using an optimized procedure, isolated and characterized by spectroscopic techniques and by X-ray diffraction in one case. Consistently with previous literature studies, different isomers are generally observed both in the solid state and in solution, where interconversion can take place following the Adams-Cotton mechanism. Although the polarity of the solvent is expected to significantly affect the cis/trans ratio, on the other hand it scarcely influences the coordination mode of the isocyanide ligand.DFT calculations indicate the bridging coordination as the most stable one, and experiments reveal that the exclusive observation of terminal isocyanide structures in some cases (cyclohexyl-isocyanide, xylyl-isocyanide) isrelated to kinetic reasons associated to steric issues.

#### Experimental

#### 1. General experimental details.

[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (99%) was purchased from Strem Chemicals, other reactants and solvents were obtained from Alfa Aesar, Merck, Apollo Scientific or TCI Chemicals and were of the highest purity available. Methyl isocyanide was prepared according to the literature [<sup>30</sup>], while other isocyanides were commercial products stored at low temperature and used as received. Contaminated labware was treated with bleach. The syntheses of diiron compounds were carried out under dry  $N_2$  using standard Schlenk techniques and MeCN distilled from CaH<sub>2</sub>. Chromatography separations were carried out under N₂on neutral alumina (≈ 150 mesh, dry / Brockmann Number I, unless otherwise specified) or Florisil<sup>®</sup> (60-100 mesh) columns. Aqueous HBF<sub>4</sub> nominally 48% w/w was titrated with NaOH (acidity 8.4 mol/L). Solutions of diiron compounds were always prepared and manipulated under N<sub>2</sub>. Solid samples containing the diiron compounds were manipulated with no special precaution over short times but were eventuallystored under N2. NMR spectra were recorded at 25 °C on a Bruker Avance II DRX400 instrument equipped with a BBFO broadband probe. CDCl<sub>3</sub> and CD<sub>3</sub>CN stored in the dark over 3A MS were used for NMR analysis. Chemical shifts (expressed in parts per million) are referenced to the residual solvent peaks [31] (<sup>1</sup>H, <sup>13</sup>C) or to external standards [<sup>32</sup>] (<sup>31</sup>P to 85% H<sub>3</sub>PO<sub>4</sub>). <sup>1</sup>H and <sup>13</sup>C spectra were assigned with the assistance of <sup>1</sup>H{<sup>31</sup>P}, <sup>13</sup>C DEPT 135 and <sup>1</sup>H-<sup>13</sup>C *qs*-HSQC experiments [33].IR spectra of solid samples (650-4000 cm<sup>-1</sup>) were recorded on a Perkin Elmer Spectrum One FT-IR spectrometer, equipped with a UATR sampling accessory. IR spectra of solutions were recorded using a CaF<sub>2</sub> liquid transmission cell (1500-2300 cm<sup>-1</sup>) on a Perkin Elmer Spectrum 100 FT-IR spectrometer. IR bands and <sup>13</sup>C NMR resonances attributed to terminal and bridging CO/CNR ligands are indicated as t-CO/t-CN and  $\mu$ -CO/ $\mu$ -CN, respectively. IR spectra were processed with Spectragryph software [34].

#### 2. Synthesis and characterization of diiron compounds.

*Carbon monoxide/isocyanide exchange (general procedure).* In a Schlenk flaskunder N<sub>2</sub>, the selected isocyanide, or its solution in MeCN (5 mL, for solid compounds), was added dropwise to a suspension of [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (1.6

6

eq for 1-7; 1.1 eq for 8) in anhydrous MeCN. Alkyl isocyanides: the mixture was heated at reflux ( $T \ge 90$  °C) for 8 h, then stirred at room temperature for additional 14 h. Aryl isocyanides: the mixture was stirred at room temperature for 14 h then heated at heated at reflux ( $T \ge 90$  °C) for 3 h. After the conversion was checked by IR,the dark red-brown suspension was dried under vacuum (40 °C). The resulting brown solid is predominantly a mixture of [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] and [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>(CNR)], according to the Fe/isocyanide molar ratio used. Compounds 1-4 were purified from the raw material by column chromatography, as detailed below for each compound.Compounds 3 and 5-8 were purified via the protonation/deprotonation route, as described in the following general procedure.

*Purification by protonation/deprotonation (general procedure)*. In a 100 mL Schlenk flask under N<sub>2</sub>, the brown solid was suspended in Et<sub>2</sub>O (30 mL) under vigorous stirring. Dropwise addition of 48% aqueous HBF<sub>4</sub> (1.6 eq.) resulted in the immediate, massive formation of a bright red solid. The mixture was vigorously stirred for 30' then filtered (G4 porous filter). The solid was thoroughly washed with Et<sub>2</sub>Oto remove [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>]. Next, the solid was moved back into the 100 mL Schlenk tube under N<sub>2</sub> and suspended in Et<sub>2</sub>O (20 mL) under vigorous stirring. Dropwise addition of Et<sub>3</sub>N (1.7 eq) was subsequently followed by a color change to dark red-violet and (partial) dissolution of the solid. The mixture was stirred for 30' then moved on top of an alumina column (h 5 cm, d 2.8 cm). Impurities were eluted with Et<sub>2</sub>O, then a red-violet band containing the title product was eluted with a CH<sub>2</sub>Cl<sub>2</sub>/Et<sub>2</sub>O 1:1  $\nu/\nu$  solution. Volatiles were removed under vacuum (40 °C), affording a dark red-purple solid.

#### [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>{CN(1H-indol-6-yl)}], 1 (Chart 1).



Prepared from [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (2.060 g, 5.82 mmol) and 1H-indol-6-yl isocyanide (530 mg, 3.73 mmol), in MeCN (90 mL). The raw product was dissolved in toluene and moved on top of a Florisil<sup>®</sup> column (h 5.5 cm, d 4.5 cm). A reddish-brown band containing  $[Fe_2Cp_2(CO)_4]$  was eluted with a toluene/CH<sub>2</sub>Cl<sub>2</sub> 10:1.5 v/v mixture then a dark-red band containing the title product was eluted with THF. Volatiles were removed under vacuum (40 °C), the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and gently evaporated under vacuum (RT), affording a dark redpurple solid. Yield: 77 % (with respect to isocyanide). Compound 1 is soluble in THF, CH<sub>2</sub>Cl<sub>2</sub>, Et<sub>2</sub>O, MeCN, poorly soluble in hexane. IR(solid state):  $\tilde{v}/cm^{-1} = 3387w$ -br (NH), 3107w, 2953w-sh, 2922m, 2853m; 2099m, 2077m-sh (t-CN); 1973s, 1933s (t-CO); 1788s-sh, 1736s (µ-CO); 1697s-br, 1661s (µ-CN); 1616s-sh, 1573m, 1507w, 1464m, 1416m, 1377w, 1357w, 1342w, 1313m, 1262w, 1243w, 1214w, 1174w, 1118w, 1089w, 1060m, 1035w, 1014m, 1001m, 941w, 928w, 893w, 865m-sh, 821m, 805m, 760m, 726w, 694w. IR (CH<sub>2</sub>Cl<sub>2</sub>):  $\tilde{v}/cm^{-1} = 1$ 2106m, 2085m-sh (t-CN); 1988s, 1948s (t-CO); 1779s, 1749s (µ-CO); 1703s (µ-CN), 1618w, 1576w, 1466m, 1417w, 1344w, 1321w, 1312w, IR (MeCN):  $\tilde{v}/cm^{-1} = 2104m$ , 2081w-sh (t-CN): 1983s, 1944s (t-CO): 1784m-sh, 1753s ( $\mu$ -CO); 1706m/s ( $\mu$ -CN), 1576w. IR (toluene):  $\tilde{v}$ /cm<sup>-1</sup> = 2102m, 2079w-sh (t-CN); 1988s, 1945s (t-CO); 1792s, 1763s ( $\mu$ -CO); 1708s ( $\mu$ -CN).<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 8.30 (s-br, 1H), 7.50–7.31, 7.24, 7.20–7.08, 6.62-6.43 (m-br, 4H) (C<sub>8</sub>H<sub>5</sub>); 4.72 (s-br, 10H, Cp).<sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 251 (br,  $\mu$ -CN); 213-211 (br, t-CO); 144.2-142.2, 134-133, 128, 125.8-125.2, 120-117 (br.), 111.5, 103.1 (C<sub>8</sub>H<sub>5</sub>); 87.9 (br, Cp).<sup>1</sup>H NMR  $(CD_3CN): \delta/ppm = 9.38 \text{ (s-br, 1H, NH)}; 7.29 \text{ (m-br)}, 6.49 \text{ (s-br)} (5H, C_8H_5); 4.78 \text{ (s-br, 10H, Cp)}.$ <sup>13</sup>C{<sup>1</sup>H} NMR (CD<sub>3</sub>CN): δ/ppm = 273.6 (μ-CO); 249.2 (μ-CN); 213.5 (t-CO); 143.7, 135.4, 134.2, 129.9, 129.2, 127.9, 126.8, 118.3, 112.7, 111.9, 102.8 (C<sub>8</sub>H<sub>6</sub>); 88.9 (br, Cp).

#### [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>{CN(CH<sub>2</sub>PO<sub>3</sub>Et<sub>2</sub>)}], 2 (Chart 2).



Prepared from[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (1.73 g, 4.87 mmol) and diethyl isocyanomethylphosphonate (0.50 mL, 3.12 mmol) in MeCN (90 mL). The raw product was suspended in Et<sub>2</sub>O and moved on top of alumina column (containing 4 % *w/w* water; h 8 cm, d 3.2 cm). A reddish-brown band containing [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] was eluted with Et<sub>2</sub>O, then a raspberry red band containing the title product was eluted with a THF/MeCN 95:5 *v/v* solution. Volatiles were removed under vacuum and the resulting red-violet solid was stored under N<sub>2</sub>.Vield: 910 mg, 58 % (with respect to isocyanide). Compound **2** is soluble in CH<sub>2</sub>Cl<sub>2</sub>, MeOH, MeCN, toluene, poorly soluble in Et<sub>2</sub>O, hexane and insoluble in water. IR (solid state):  $\tilde{v}$ /cm<sup>-1</sup> = 3109w, 3086w, 2986w, 2906w, 2886w; 1975s, 1937s (t-CO); 1771s (µ-CO), 1687s-br (µ-CN), 1480w, 1440w, 1418m, 1405m, 1390w, 1367w, 1359w, 1290w, 1263m-sh, 1238s, 1163m, 1096w, 1050s, 1024s, 980m-sh, 957s, 945s, 892m, 849m-sh, 830s, 797m, 774m, 721m, 671s.IR (CH<sub>2</sub>Cl<sub>2</sub>):  $\tilde{v}$ /cm<sup>-1</sup> = 1992s, 1952m (t-CO); 1789m (µ-CO), 1690m-br (µ-CN).IR (MeCN):  $\tilde{v}$ /cm<sup>-1</sup> = 1987s, 1946w (t-CO); 1788m (µ-CO), 1692m-br (µ-CN).IR (toluene):  $\tilde{v}$ /cm<sup>-1</sup> = 1990s, 1950m (t-CO); 1795m (µ-CO), 1693mbr (µ-CN).<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 4.88, 4.78 (s-br, 10H, Cp); 4.40 (d-br, <sup>2</sup>J<sub>HP</sub> = 13.5 Hz, 2H, NCH<sub>2</sub>), 4.29 (m-br, 4H, OCH<sub>2</sub>), 1.42 (m-br, 6H, CH<sub>3</sub>).<sup>13</sup>C {<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 272.0 (µ-CO); 212.8, 210.4 (t-CO); 260.6–260.2 (m, µ-CN); 89.5, 87.6 (Cp); 62.7 (d, <sup>2</sup>J<sub>CP</sub> = 6 Hz, OCH<sub>2</sub>), 57.5 (d, <sup>1</sup>J<sub>CP</sub> = 162 Hz, NCH<sub>2</sub>), 16.7 (d, <sup>3</sup>J<sub>CP</sub> = 4 Hz, CH<sub>3</sub>).<sup>31</sup>P {<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 22.7.

[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>{CN(C<sub>6</sub>H<sub>11</sub>)}], 3 (Chart 3)[6, <sup>35</sup>].



Prepared from[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (1.0 g, 2.83 mmol) and cyclohexyl isocyanide (0.21 mL, 1.69 mmol) in MeCN (30 mL). Purified by the protonation/deprotonation route. Yield: 625 mg, 82 % (with respect to isocyanide). Alternatively, the raw product was dissolved in toluene and moved on top of a Florisil<sup>\*</sup> column (h 5 cm, d 4.5 cm). A reddish-brown band containing [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] was eluted with a toluene/CH<sub>2</sub>Cl<sub>2</sub> 10:1.5 v/v mixture then a dark red band containing the title product was eluted with Et<sub>2</sub>O. Volatiles were removed under vacuum (40 °C), affording a dark red-purple solid. Yield: 211 mg, 29 % (with respect to isocyanide).Compound **3** is soluble in CH<sub>2</sub>Cl<sub>2</sub>, acetone, MeCN, toluene, poorly soluble in MeOH, insoluble in water. IR (solid state):  $\tilde{v}$ /cm<sup>-1</sup> = 3108w, 2932m, 2856w, 2117s-br (t-CN); 1954s-sh, 1940s-br (t-CO); 1766w-sh, 1726s-br ( $\mu$ -CO); 1650-1550m-br, 1448m, 1432m, 1416m, 1353m, 1316m, 1270w, 1238w, 1150w, 1129w, 1114w, 1061w, 1013m, 997m, 927w, 864m-sh, 824m, 809m, 664m. IR (CH<sub>2</sub>Cl<sub>2</sub>):  $\tilde{v}$ /cm<sup>-1</sup> = 2128s (t-CN), 1985w, 1947s (t-CO), 1780w-sh, 1746s ( $\mu$ -CO). IR (toluene):  $\tilde{v}$ /cm<sup>-1</sup> = 2122s (t-CN), 1988w, 1946s (t-CO), 1784w, 1760s ( $\mu$ -CO). IR (MeCN):  $\tilde{v}$ /cm<sup>-1</sup> = 2125s (t-CN); 1980w, 1943s (t-CO); 1782w-sh, 1749s ( $\mu$ -CO); 1636w-br. <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 4.68, 4.61 (s-br, 10H, Cp); 3.36 (s, 1H, NCH); 1.69–1.52, 1.45–1.30 (m, 10H, CH<sub>2</sub>).<sup>13</sup>C {<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 281.7 ( $\mu$ -CO); 217.3, 212.7 (t-CO); 161.9 (t-CN); 87.2, 86.7sh. (Cp); 55.0 (NCH); 33.0, 25.1, 23.0 (CH<sub>2</sub>).

## [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>{CN(4-C<sub>6</sub>H<sub>4</sub>OMe)}], 4 (Chart 4)[<sup>36</sup>].



Prepared from[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (4.205 g, 11.88 mmol) and 4-methoxyphenyl isocyanide (1.0 g, 7.51 mmol) in MeCN (80 mL). The raw product was dissolved in toluene and moved on top of an alumina column (h 5.5 cm, d 4.5 cm). A reddish-brown band containing [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] was eluted with a toluene/CH<sub>2</sub>Cl<sub>2</sub> 10:1.5  $\nu/\nu$  mixture then a dark red band containing the title product was eluted with Et<sub>2</sub>O. Volatiles were removed under vacuum (40 °C), affording a dark red-purple solid. Yield: 80 % (with respect to isocyanide). Compound **4** is soluble in CH<sub>2</sub>Cl<sub>2</sub>, acetone, MeCN, toluene moderately soluble in Et<sub>2</sub>O, poorly soluble in pentane.IR (solid state):  $\tilde{\nu}$ /cm<sup>-1</sup> = 3109w, 3024w, 2958w, 2924m, 2853w; 2002s-sh, 1966s, 1949s (t-CO); 1774s ( $\mu$ -CO), 1684s-br ( $\mu$ -CN); 1599m-sh, 1575w, 1496s, 1463m, 1437m, 1416m, 1378w, 1356w, 1284m, 1260m, 1238s, 1197m, 1178m, 1161m, 1102m,

1059m, 1026s, 1013m, 859w-sh, 830s, 817s, 806s, 727w-sh, 718s, 668s. IR (CH<sub>2</sub>Cl<sub>2</sub>):  $\tilde{v}/cm^{-1} = 2102m$  (t-CN); 1989s, 1949s (t-CO); 1788s, 1751s (μ-CO); 1698s-br (μ-CN), 1604w, 1575w, 1500s, 1466w, 1442w. IR (MeCN):  $\tilde{v}/cm^{-1} = 2100w$  (t-CN); 1984s, 1945m (t-CO); 1788m, 1754m (μ-CO); 1698s-br (μ-CN). IR (toluene):  $\tilde{v}/cm^{-1} = 2097w$  (t-CN); 1990s, 1947s (t-CO); 1795s, 1764m (μ-CO); 1703s-br (μ-CN).<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta/ppm =$ 7.11 (m-br, 2H), 6.97 (m-br, 2H) (C<sub>6</sub>H<sub>4</sub>); 4.72 (s-br, 10H, Cp), 3.85 (s-br, 3H, OCH<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta/ppm =$ 272.6 (μ-CO); 253.6 (μ-CN); 217br., 211.0 (t-CO); 156.2, 143.5, 121.9, 114.5 (C<sub>6</sub>H<sub>4</sub>); 89.0br., 87.9 (Cp); 55.7 (OCH<sub>3</sub>).

# $[Fe_2Cp_2(CO)_3(CN(2,6-C_6H_3Me_2))], 5 (Chart 5)[^{37}].$



Prepared from[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (9.66 g, 27.3 mmol) and xylyl isocyanide (2.25 g, 17.17 mmol)in MeCN (100 mL). A fraction of the raw material (542 mg) was purified by the protonation/deprotonation route. Yield: 308 mg, 83 % (with respect to isocyanide; re-scaled from the mass fraction of the raw material).Compound **5** is soluble in CH<sub>2</sub>Cl<sub>2</sub>, CHCl<sub>3</sub>, toluene, moderately soluble in Et<sub>2</sub>O, poorly soluble in MeCN.IR (solid state): v/cm<sup>-1</sup> = 2081s (t-CN); 2000w; 1943s (t-CO); 1901w-sh; 1733m, 1739s ( $\mu$ -CO); 1589w, 1470w, 1432w, 1418w, 1382w, 1358w, 1261w, 1186w, 1165w, 1117w, 1092w, 1059w, 1037w, 1012w, 923w, 864w, 819m-sh, 810m, 779m, 739w, 720w, 676w. IR (CH<sub>2</sub>Cl<sub>2</sub>): v/cm<sup>-1</sup> = 2090s-br (t-CN); 1989m, 1950s (t-CO); 1783w-sh, 1752s ( $\mu$ -CO), 1605w, 1589w. IR (MeCN): v/cm<sup>-1</sup> = 2136m, 2082m-br (t-CN); 1989s, 1947m (t-CO); 1775m, 1756m ( $\mu$ -CO), 1711w ( $\mu$ -CN). IR (toluene): v/cm<sup>-1</sup> = 2082s-br (t-CN); 1989m, 1949s (t-CO); 1784w-sh 1764s, 1741w-br ( $\mu$ -CO).<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 6.96 (s, 3H, C<sub>6</sub>H<sub>3</sub>), 4.73 (s, 10H, Cp), 2.26 (s, 6H, CH<sub>3</sub>). <sup>13</sup>C {<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 278 (br.,  $\mu$ -CO); 215-212 (br., t-CO); 176 (br., t-CN); 134.3 (br.), 127.8, 126.8 (br.) (C<sub>6</sub>H<sub>3</sub>Me<sub>2</sub>); 87.6 (Cp), 18.9 (CH<sub>3</sub>).Crystals suitable for X-ray analysis were collected from a ... solution of **5**....

[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>(CNMe)], 6 (Chart 6)[35].



Prepared from[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (25.7 g, 72.6 mmol) andfreshly prepared methyl isocyanide (2.5 mL, 44.4 mmol) in MeCN (150 mL). A fraction of the raw material (510 mg) was purified by the protonation/deprotonation route. Yield: 190 mg, 60 % (with respect to isocyanide; re-scaled from the mass fraction of the raw material). Compound **6** is soluble in CH<sub>2</sub>Cl<sub>2</sub>, toluene, MeCN, less soluble in Et<sub>2</sub>O.IR (solid state):  $\tilde{v}$ /cm<sup>-1</sup> = 2144m (t-CN); 1985w, 1944s (t-CO); 1770m-sh, 1730s-br ( $\mu$ -CO); 1700m-sh ( $\mu$ -CN); 1454w, 1415w, 1355w, 1260w, 1113w, 1062w, 1013w, 1001w, 930w, 876w, 863w, 840m-sh, 820m, 692w, 656w.IR (CH<sub>2</sub>Cl<sub>2</sub>):  $\tilde{v}$ /cm<sup>-1</sup> = 2153m (t-CN); 1988m, 1948s (t-CO); 1778w, 1745s ( $\mu$ -CO), 1719w-sh ( $\mu$ -CN).IR (MeCN):  $\tilde{v}$ /cm<sup>-1</sup> = 2155m (t-CN); 1982s, 1944s (t-CO); 1788w, 1749s ( $\mu$ -CO); 1727w-sh ( $\mu$ -CN).IR (toluene):  $\tilde{v}$ /cm<sup>-1</sup> = 2147m (t-CN); 1989s, 1946s (t-CO); 1791w, 1760s ( $\mu$ -CO); 1731w ( $\mu$ -CN).<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 4.70 (s-br, 10H, Cp); 3.00 (s-br, 3H, CH<sub>3</sub>).<sup>13</sup>C {<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 280.6 ( $\mu$ -CO); 213.0, 210.5 (t-CO); 162.5 (t-CN); 87.3 (Cp); 46.6, 30.5 (CH<sub>3</sub>).

#### [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>{CN(2-C<sub>10</sub>H<sub>7</sub>)}], 7 (Chart 7)[17].



Prepared from[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (455 mg, 1.29 mmol) and 2-naphthyl isocyanide (124 mg, 0.809 mmol) in MeCN (30 mL). Purified by the protonation/deprotonation route. Yield: 194 mg, 50 % (with respect to

isocyanide).Compound **7** is soluble in CH<sub>2</sub>Cl<sub>2</sub>, toluene, MeCN, Et<sub>2</sub>O, hexane. IR (solid state):  $\tilde{v}/cm^{-1} = 3107w$ , 3053w, 2963w, 2853w; 1973s, 1934s (t-CO); 1782s-sh, 1752w (µ-CO), 1679s-br (µ-CN); 1616s, 1589s, 1501m, 1461m, 1432m, 1417m, 1379m, 1357m, 1266w, 1245w, 1206w, 1157m, 1116w, 1060w, 1013m, 999m, 956w, 879w-sh, 854m-sh, 839m-sh, 829m-sh, 814s, 749m, 711w, 679w. IR (CH<sub>2</sub>Cl<sub>2</sub>):  $\tilde{v}/cm^{-1} = 1991s$ , 1951m (t-CO); 1790m (µ-CO); 1694m-br (µ-CN), 1623w, 1593w. IR (MeCN):  $\tilde{v}/cm^{-1} = 1986s$ , 1947m (t-CO); 1790m (µ-CO); 1694m-br (µ-CN), 1623w. IR (toluene):  $\tilde{v}/cm^{-1} = 1991s$ , 1948s (t-CO); 1797s, 1756w (µ-CO); 1740w-sh, 1700s-br (µ-CN); 1623w. <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta/ppm = 7.91$  (d, J = 8.7 Hz), 7.86 (t, J = 7.6 Hz), 7.53–7.47 (m), 7.46–7.40 (m) (7H, C<sub>10</sub>H<sub>7</sub>); 4.75, 4.72 (s-br, 10H, Cp).<sup>13</sup>C {<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta/ppm = 271.9$  (µ-CO); 255.2 (µ-CN); 213 (br.), 211.0 (t-CO); 147.0, 134.5, 130.7, 129.3, 128.0, 127.7, 126.6, 124.8, 122.1, 117.0 (C<sub>10</sub>H<sub>7</sub>); 89.5 (br.), 87.9 (Cp).

#### [Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>3</sub>(CNCH<sub>2</sub>Ph)], 8 (Chart 8)[35].



Prepared from[Fe<sub>2</sub>Cp<sub>2</sub>(CO)<sub>4</sub>] (337 mg, 0.96 mmol) and benzyl isocyanide (0.10 mL, 0.82 mmol) in MeCN (25 mL).Purified by the protonation/deprotonation route. Yield: 241 mg, 66 % (with respect to isocyanide).Compound **8** is soluble in CH<sub>2</sub>Cl<sub>2</sub>, MeCN, toluene, less soluble in Et<sub>2</sub>O. IR (solid state):  $\tilde{v}$ /cm<sup>-1</sup> = 3103w, 3066w, 3032w, 2965w, 2912w, 2850w; 1968s, 1922s (t-CO); 1757s ( $\mu$ -CO); 1712-1699s-br ( $\mu$ -CN); 1494w, 1453w, 1430w, 1418m, 1357w, 1314m, 1262w, 1198w, 1070w, 1059w, 1027w, 1013w, 1005w, 953w, 881w, 867w-sh, 845m-sh, 826s, 749m, 737w, 699s-sh, 684s-br. IR (CH<sub>2</sub>Cl<sub>2</sub>):  $\tilde{v}$ /cm<sup>-1</sup> = 2131m (t-CN); 1989s, 1948s (t-CO); 1784m, 1747s ( $\mu$ -CO); 1710m ( $\mu$ -CN).IR (MeCN):  $\tilde{v}$ /cm<sup>-1</sup> = 2132w (t-CN); 1983s, 1943m (t-CO);

1785m, 1750m ( $\mu$ -CO); 1713m ( $\mu$ -CN).IR (toluene):  $\tilde{\nu}$ /cm<sup>-1</sup> = 2126w (t-CN); 1989s, 1947s (t-CO); 1793m, 1760m ( $\mu$ -CO); 1710m ( $\mu$ -CN). <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm = 7.51 (s-br), 7.37 (s-br), 7.32–7.27 (m), 7.20 (s-br) (5H, C<sub>6</sub>H<sub>5</sub>); 5.18 (s-br, 1H, CH<sub>2</sub>); 4.82, 4.69 (s-br, 10H, Cp); 4.45 (br, 1H, CH<sub>2</sub>). <sup>13</sup>C {<sup>1</sup>H} NMR (CDCl<sub>3</sub>)  $\delta$ /ppm = 280br, 273.3 ( $\mu$ -CO); 251.1 ( $\mu$ -CN); 215-212br, 210.8 (t-CO); 139.9, 128.9, 128.1, 127.1, 126.8 (C<sub>6</sub>H<sub>5</sub>); 87.5 (Cp); 64.9, 49br (CH<sub>2</sub>).

### 3. X-ray crystallography

Crystal data and collection details for **5-ct** are reported in Table 4. Data were recorded on a Bruker APEX II diffractometer equipped with a PHOTON2 detector using Mo–K $\alpha$  radiation. Data were corrected for Lorentz polarization and absorption effects (empirical absorption correction SADABS).<sup>38</sup> Structures were solved by direct methods and refined by full-matrix least-squares based on all data using  $F^2$ .<sup>39</sup> Hydrogen atoms were fixed at calculated positions and refined by a riding model. All non-hydrogen atoms were refined with anisotropic displacement parameters.

	5-ct
Formula	$C_{22}H_{19}Fe_2NO_3$
FW	457.08
Т, К	293(2)
λ, Å	0.71073
Crystal system	Monockinic
Space group	P2 <sub>2</sub> /n
<i>a</i> , Å	8.1610(7)
<i>b,</i> Å	10.7531(10)
<i>c</i> , Å	21.4677(19)
$\beta,^{\circ}$	96.282(3)
Cell Volume, Å <sup>3</sup>	1869.5(3)
Z	4
D <sub>c</sub> , g·cm <sup>-3</sup>	1.624
$\mu$ , mm <sup>-1</sup>	1.575
F(000)	936
Crystal size, mm	0.16×0.14×0.11
θ limits,°	1.909-25.100
Reflections collected	18556
Independent reflections	3312 [ <i>R<sub>int</sub></i> = 0.0491]

Table 4. Crystal data and measurement details for 5-ct.

Data / restraints	3312 / 0 / 255		
/parameters	5512707255		
Goodness on fit on F <sup>2</sup>	1.143		
$R_1 (I > 2\sigma(I))$	0.0432		
wR <sub>2</sub> (all data)	0.0993		
Largest diff. peak and hole, e Å <sup>-3</sup>	0.509 /0.311		

#### 4. Computational studies

All geometries were optimized with ORCA 4.0.1.2, using the B97 functional in conjunction with a triple- $\zeta$  quality basis set (def2-TZVP). The dispersion corrections were introduced using the Grimme D3-parametrized correction and the Becke–Johnson damping to the DFT energy. Most of the structures were confirmed to be local energy minima (no imaginary frequencies), but in some case a small, unavoidable negative frequency relative to the Cp rotation around the M-Cp axis was observed. The solvent was considered through the Continuum-like Polarizable Continuum Model (C-PCM, dichloromethane).

#### 5. Thermal isomerization study

Dark red-brown solutions of **3** and **5** (20 mg) in anhydrous toluene (5 mL) were heated at reflux under a N<sub>2</sub> atmosphere for 3 h while periodically sampled for IR analysis. The appearance of new weak absorption peaks, ascribable to isocyanide-bridging isomers, was observed. Next, solutions were cooled to room temperature and IR spectra were recorded again, showing no variations. Following volatiles removal under vacuum, the solid-state spectra of the brown residues were recorded, displaying a different CO/CNR band pattern with respect to those of the starting materials, consistent with the presence of the isocyanide-bridging isomers. IR data are reported in the following, and the spectra are displayed in Figures S41 and S42. *Compound 3*. IR (toluene): new absorptions at 1996w (t-CO), 1782w-sh ( $\mu$ -CO), 1696w-br ( $\mu$ -CN) cm<sup>-1</sup>. IR (solid state): v/cm<sup>-1</sup> = 2118m (t-CN); 1991m, 1952s-sh, 1940s-br (t-CO), 1772s-sh, 1752s, 1731s ( $\mu$ -CO); 1695m-sh ( $\mu$ -CN). *Compound 5*. IR (toluene):new absorptions at 1995w (t-CO), 1782w-sh ( $\mu$ -CO), 1720w-br ( $\mu$ -CN) cm<sup>-1</sup>. IR (solid state): v/cm<sup>-1</sup> = 2081s (t-CN); 1995w, 1958w-sh, 1932s (t-CO); 1767m-sh, 1755m-sh, 1741s ( $\mu$ -CO); 1706s-sh ( $\mu$ -CN).

#### Acknowledgements

We gratefully thank the University of Pisa (Fondi di Ateneo 2019) for financial support.

#### **Supporting Information Available**

IR and NMR spectra of products; DFT geometries of **1-5** and **7-8**.CCDC reference number XXXXX (**5-ct**) contains the supplementary crystallographic data for the X-ray study reported in this paper. This data can be obtained free of charge at https://www.ccdc.cam.ac.uk/structures/.Cartesian coordinates of the DFT-optimized structures are collected in a separated .xyz file.

#### References

- (a) K. T. Mahmudov, V. Yu. Kukushkin, A. V. Gurbanov, M. A. Kinzhalov, V. P. Boyarskiy, M. F. C. Guedes da Silva, A. J. L. Pombeiro, Coord. Chem. Rev. 384 (2019) 65-89.
- (b) K. V. Luzyanin, A. J. L. Pombeiro in Isocyanide Chemistry: Applications in Synthesis and Material Science, Ed. V. G. Nenajdenko, pp. 531-550, Ed. Wiley.
- (c) A. J. L. Pombeiro, M. F. C. Guedes da Silva, R. A. Michelin, Coord. Chem. Rev. 218 (2001) 43-74.
- (d) J. F. Kronauge, D. J. Mindiola, Organometallics 35 (2016) 3432-3435.
- (e) A. Arroyave, M. Gembicky, A. L. Rheingold, J. S. Figueroa, Inorg. Chem. 59 (2020) 11868-11878.

2 C. Elschenbroich, Organometallics (2006) 3rd Edition, Wiley.

<sup>1</sup> Selected references:

<sup>(</sup>f) D.-A. Rosca, J. Fernandez-Cestau, A. S. Romanov, M. Bochmann, J. Organomet. Chem. 792 (2015) 117-122.

- 3 R. Mazzoni, F. Marchetti, A. Cingolani, V. Zanotti, Inorganics 7 (2019) 25-43; doi:10.3390/inorganics7030025.
- 4 M. A. Guillevic, E. L. Hancox, B. E. Mann, J. Chem. Soc. Dalton Trans. (1992) 1729-1733.
- 5 (a) J. A. S. Howell, P. Mathur, J. Organomet. Chem. 174 (1977) 335-341.
  (b) J. A. S. Howell, A. J. Rowan, J. Chem. Soc. Dalton Trans. (1980) 503-510.
  (c) R. D. Adams, F. A. Cotton, Inorg. Chem. 13 (1974) 257-262.
- J. Bellerby, M. J. Boylan, M. Ennis, A. R. Manning, J. Chem. Soc. Dalton Trans. (1978) 1185-1189.
- 7 J. A. S. Howell, A. J. Rowan, J. Chem. Soc. Dalton Trans. (1981) 297-301.
- 8 S. Nakanishi, Y. Taniki, Y. Otsuji, J. Chem. Soc. Chem. Commun. (1993) 709-710.
- 9 F. A. Cotton, B. A. Frenz, Inorg. Chem. 13 (1974) 253-256.
- 10 M. Ennis, R. Kumar, A. R. Manning, J. A. S. Howell, P. Mathur, A. J. Rowan, F. S. Stephens, J. Chem. Soc. Dalton Trans. (1981) 1251-1259.
- 11 G. Mcnally, P. T. Murray, A. R. Manning, J. Organomet. Chem. 243 (1983) C87-C88.
- 12 N. J. Coville, M. O. Albers, E. Singleton, J. Chem. Soc. Dalton Trans. (1982) 1389-1391.
- 13 A. R. Manning, G. McNally, P. Soye, Inorg. Chim. Acta 180 (1991) 103-110.
- 14 J. A. S. Howell, M. J. Mays, I. D. Hunt, O. S. Mills, J. Organomet. Chem. 128 (1977) C29-C30.
- (a) F. Marchetti, S. Zacchini, V. Zanotti, Organometallics 34 (2015) 3658–3664.
  (b) F. Marchetti, S. Zacchini, V. Zanotti, Organometallics 33 (2014) 3990-3997.
- 16 (a) K. K. Joshi, O. S. Mills, P. L. Pauson, B. W. Shaw, W. H. Stubbs, J. Chem. Soc. Chem. Commun. (1965) 181-182.
  - (b) I. D. Hunt, O. S. Mills, Acta Cryst. B33 (1977) 2432-2435.
  - (c) W. P. Fehlhammer, A. Mayr, W. Kehr, J. Organomet. Chem. 197 (1980), 327-334.
  - (d) M. O. Albers, A. A. Chalmers, G. J. Kruger, D. C. Liles, H. E. Oosthuizen, E. Singleton, N. J. Coville, J. Organomet. Chem. 306 (1986) 385-394.
- 17 G. Agonigi, M. Bortoluzzi, F. Marchetti, G. Pampaloni, S. Zacchini, V. Zanotti, Eur. J. Inorg. Chem. (2018) 960-971.
- (a) F. Marchetti, Eur. J. Inorg. Chem. (2018) 3987-4003.
  (b) G. Agonigi, L. Biancalana, M. G. Lupo, M. Montopoli, N. Ferri, S. Zacchini, F. Binacchi, T. Biver, B. Campanella, G. Pampaloni, V. Zanotti, F. Marchetti, Organometallics 39 (2020) 645-657.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/)

# When citing, please refer to the published version.

(c) D. Rocco, L. K. Batchelor, G. Agonigi, S. Braccini, F. Chiellini, S. Schoch, T. Biver, T. Funaioli, S. Zacchini, L. Biancalana, M. Ruggeri, G. Pampaloni, P. J. Dyson, F. Marchetti, Chem. Eur. J. 25 (2019) 14801-14816.

- 19 J. A. Labinger, Inorg. Chim. Acta 424 (2015) 14-19.
- 20 V. G. Albano, L. Busetto, L. Carlucci, M. C. Cassani M. Monari, V. Zanotti, J. Organomet. Chem. 488 (1995) 133-139.
- 21 S. Willis, A. R. Manning, F. S. Stephens, J. Chem. Soc., Dalton Trans. (1979) 23-27
- 22 L. J. Farrugia, J. Chem. Soc., Dalton Trans. (1997) 1783-1792.
- <sup>23</sup> M. A. Mironov, M. I. Tokareva, M. N. Ivantsova, V. S. Mokrushin, Russ. J. Org. Chem. 40 (2004) 847.853
- <sup>24</sup> U. Schöllkopf, R. Schröder, Tetrahedron Letters 9 (1973) 633-636.
- <sup>25</sup> M. Hanack, Michael, R. Thies, Chem. Ber., 121 (1988) 1225-1230.
- <sup>26</sup> S. Kamijo, T. Jin, Y. Yamamoto, J. Am. Chem. Soc. 123 (2001) 9453-9454.
- <sup>27</sup> R. L. Williams, J. Chem. Phys. 25 (1956) 656-660.
- <sup>28</sup> B. Kim, J. M. Beebe, Y. Jun, F. G. Zhu, J. Am. Chem. Soc.128 (2006) 4970-4971.
- <sup>29</sup> U. Hess, H. Brosig, J. Komenda, Pharmazie 54 (1999) 412-417.
- 30 R. E. Schuster, J. E. Scott, J. Casanova Jr. Methyl Isocyanide, Org. Synth. 46 (1966) 75.
- 31 G. R. Fulmer, A. J. M. Miller, N. H. Sherden, H. E. Gottlieb, A. Nudelman, B. M. Stoltz, J. E. Bercaw, K. I. Goldberg, Organometallics 29 (2010) 2176-2179.
- <sup>32</sup> R. K. Harris, E. D. Becker, S. M. Cabral De Menezes, R. Goodfellow, P. Granger, Pure Appl. Chem. 73 (2001) 1795-1818.
- 33 W. Willker, D. Leibfritz, R. Kerssebaum, W. Bermel, Magn. Reson. Chem. 31 (1993) 287-292.
- <sup>34</sup> F. Menges, "Spectragryph optical spectroscopy software", Version 1.2.9 @ (2018)
   http://www.effemm2.de/spectragryph.
- <sup>35</sup> M. J. Boylan, J. Bellerby, J. Newman and A. R. Manning, J. Organomet. Chem. 47 (1973) C33-C35.
- <sup>36</sup> L. Busetto, F. Marchetti, S. Zacchini, V. Zanotti, Organometallics 25 (2006) 4808-4816.
- <sup>37</sup> V. G. Albano, L. Busetto, M. Monari, V. Zanotti, J. Organomet. Chem. 606 (2000) 163-168.
- 38 G. M. Sheldrick, SADABS-2008/1 Bruker AXS Area Detector Scaling and Absorption Correction, Bruker AXS: Madison, Wisconsin, USA, 2008
- 39 G. M. Sheldrick, Acta Crystallogr. 2015, 71c, 3-8.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.