

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Fatty acid profile, oxidative status, and content of volatile organic compounds in raw and cooked meat of different chicken strains

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Cartoni Mancinelli A., Silletti E., Mattioli S., Dal Bosco A., Sebastiani B., Menchetti L., et al. (2021). Fatty acid profile, oxidative status, and content of volatile organic compounds in raw and cooked meat of different chicken strains. *POULTRY SCIENCE*, 100(2), 1273-1282 [10.1016/j.psj.2020.10.030].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/801250> since: 2021-02-18

*Published:*

DOI: <http://doi.org/10.1016/j.psj.2020.10.030>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

Fatty acid profile, oxidative status, and content of volatile organic compounds in raw and cooked meat of different chicken strains

A. Cartoni Mancinelli, E. Silletti, S. Mattioli, A. Dal Bosco, B. Sebastiani, L. Menchetti, A. Koot, S. van Ruth, C. Castellini

PII: S0032-5791(20)30783-5

DOI: <https://doi.org/10.1016/j.psj.2020.10.030>

Reference: PSJ 776

To appear in: *Poultry Science*

Received Date: 19 March 2020

Revised Date: 8 October 2020

Accepted Date: 20 October 2020

Please cite this article as: Mancinelli A.C., Silletti E., Mattioli S., Bosco A.D., Sebastiani B., Menchetti L., Koot A., van Ruth S. & Castellini C., Fatty acid profile, oxidative status, and content of volatile organic compounds in raw and cooked meat of different chicken strains, *Poultry Science* (2020), doi: <https://doi.org/10.1016/j.psj.2020.10.030>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© YEAR Published by Elsevier Inc. on behalf of Poultry Science Association Inc.



# EFFECT OF GENOTYPE ON RAW AND COOKED POULTRY MEAT

## Fatty acid profile, oxidative status, and content of volatile organic compounds in raw and cooked meat of different chicken strains

A. Cartoni Mancinelli<sup>\*1</sup>, E. Silletti<sup>+</sup>, S. Mattioli<sup>\*</sup>, A. Dal Bosco<sup>\*</sup>, B. Sebastiani<sup>#</sup>, L. Menchetti<sup>§</sup>, A. Koot<sup>+</sup>, S. van Ruth<sup>+</sup> and C. Castellini<sup>\*</sup>

*<sup>\*</sup>Department of Agricultural, Environmental and Food Science, University of Perugia, Borgo XX Giugno 74, 06124, Perugia, Italy.*

*<sup>+</sup>Wageningen Food Safety Research, P.O. Box 230, 6700 AE Wageningen, The Netherlands.*

*<sup>#</sup>Department of Chemistry, Biology and Biotechnology University of Perugia, via Elce di Sotto, 8 - 06123 Perugia, Italy.*

*<sup>§</sup>Department of Veterinary Medicine, University of Perugia, Via San Costanzo 74, 06126, Perugia, Italy.*

Declarations of interest: none

<sup>1</sup>Corresponding author: [acartonimancinelli@gmail.com](mailto:acartonimancinelli@gmail.com)

**ABSTRACT**

Chicken meat is rich in unsaturated fatty acids. Therefore, it is more susceptible to lipid oxidation and production of volatile organic compounds (**VOC**). In this study, we evaluated the fatty acids, antioxidants, and VOC profiles of raw and cooked meat samples derived from four strains of chicken differing in their growth rates, which were as follows: Slow-Growing (**SG**, Leghorn), Medium-Growing (**MG**, Hubbard and Naked Neck) and Fast-Growing (**FG**, Ross). The VOC profile of meat was measured using proton-transfer reaction mass spectrometry (**PTR-MS**). The VOC were identified using PTR-time of flight-MS (**PTR-ToF-MS**). The data were analyzed using both univariate and multivariate models. Twenty main VOC were identified, which were classified into the following chemical categories: aldehydes, alkadienes, alkenes, furans, amides, alcohols, and other compounds. Our results revealed that the chicken genotype and the method of cooking strongly influenced the VOC profile of the meat. Identifying the relationships between these traits allowed us to highlight the trade-off of the main substrates such as n-3 and n-6 Polyunsaturated Fatty Acids (**PUFA**), protective substances (antioxidants) and degradation products (VOC) of the poultry meat produced during cooking. The extent of VOC production and n-3 loss was found to be higher for the SG genotype. Reduction of n-6 was higher in MG whereas small losses in antioxidants and PUFA were observed in the FG genotype, consequently, resulting in the lowest production of VOC.

The SG and MG are genotypes more active from a kinetic point of view respect to the FG ones. For this reason, in the FG genotypes the antioxidants are less involved in the oxidative stress induced by the movement thus, they were available to protect the lipid of the meat during the cooking process. These results suggested that the use of SG and MG genotypes requires a specific dietary protocol (e.i. increasing the antioxidants content) in order to counteract the lipid oxidations in all the phases: *in vivo*, *post-mortem*, and during/after cooking.

**Keywords:** genotype, cooking, volatile organic compound, fatty acid profile, antioxidant.

## INTRODUCTION

Humans consume diets with highly unbalanced lipid profiles and there is a consensus on the need to reduce the consumption of saturated fatty acids (**SFA**) and increase that of polyunsaturated fatty acids (**PUFA**), particularly of the n-3 series.

Fish is the main source of n-3 long-chain PUFA ( $\geq 20$  carbons) such as eicosapentaenoic (**EPA**) and docosahexaenoic acid (**DHA**). However, increased fishing and aquaculture has resulted in issues of sustainability as they depend on the marine trophic chain and use of wild fish as feed for farmed fish, respectively (Tocher et al., 2019).

Thus, it is crucial to find other sustainable food sources containing long chain n-3 (Menchetti. et al., 2018). Consequently, the ability of terrestrial animals to elongate and desaturate  $\alpha$ -linolenic acid (**ALA**), the precursor of EPA and DHA, needs to be carefully investigated.

Although standard poultry meat contains low levels of EPA and DHA, several factors such as sex, feed, rearing systems, and genotypes may affect their content. Particularly, genotype is an important factor affecting the fatty acid composition of poultry meat, which is as follows: usually, a higher percentage of n-3 PUFA is produced by slow-growing (**SG**) chickens compared to the fast-growing (**FG**) ones (Sirri et al., 2010). Castellini et al., (2002) showed that this content could increase in SG birds raised in free-range and organic systems due to high pasture intake. Simultaneously, intake of grass improves their antioxidant responses and prevents PUFA from oxidation (Dal Bosco et al., 2016, Cartoni Mancinelli et al., 2019).

Nevertheless, there is a lack of knowledge on how meat processing (freezing, cooking) can affect its nutritional properties and the development of unpleasant odors and aroma, which are important for consumer acceptability. Volatile organic compounds (**VOC**) (Holm et al., 2013), are some of the molecules responsible for such alteration in the food, the production of which, is influenced by many factors (Leroy et al., 2009). For instance, degradation of nutrients on the surface of cured products, particularly by lactic acid bacteria, can produce unpleasant odours (Holm et al., 2012).

73 However, hygienic conditions and use of low storage temperature, which restrict the growth of  
74 spoilage microorganisms, can prevent excessive generation of VOC (Gill, 2007) in raw meat.  
75 Additionally, oxidation of lipids during cooking is the main factor responsible for VOC production  
76 (Angelo et al., 1987). Warmed-over flavors develop due to the thermal autoxidation of PUFA,  
77 although other components such as proteins or carbohydrates can also contribute to the process. A  
78 high cooking temperature induces the Maillard reaction, which also results in the development of  
79 VOC (Byrne et al., 2001). Therefore, meat is a highly perishable food and its susceptibility to lipid  
80 oxidation depends on the species of livestock, which particularly decreases in the order of fish >  
81 poultry > pork > beef > lamb. This ranking is attributed to the different levels of PUFA and  
82 presence of specific antioxidants, such as tocopherols, in the meat (Rhee et al., 1996).  
83 Since the production of VOC is linked to the contents of PUFA and antioxidants, which vary with  
84 the species and genotype of the livestock (Sirri et al., 2010), this study evaluates the effect of  
85 cooking on lipid oxidation, PUFA, and VOC in different poultry strains. Accordingly, the amount  
86 of VOC produced, fatty acid profile, and content of tocopherols and thiobarbituric acid-reactive  
87 substances (**TBARS**) in raw and cooked meat derived from four strains of chicken with different  
88 growth rates were analyzed.

## MATERIALS AND METHODS

Pure linoleic acid (**LA**) and ALA, linolenic acid methyl ester, and organic solvents were obtained from Sigma-Aldrich (Germany). The methyl ester of stearidonic acid (C18:4n-3) was obtained from Cayman Chem (Ann Arbor, Michigan, USA).

### *Experimental layout and sample preparation*

Poultry with 4 different genotypes and growth rates, namely SG (Leghorn), medium growing (**MG**, Naked Neck and Hubbard Red JA), and FG (Ross 308) were reared at the experimental section of the University of Perugia (Perugia, Italy) according to the EU Regulation 834/07 and Directive 2010/63/EU and transposed as per Regional Directive n.26 on animal welfare for experimental and other scientific purposes.

The animals were reared indoors in equal densities (5 chickens/m<sup>2</sup>) and administered the same standard diets (Table S1).

Since the growth rate for different genetic strains differs significantly, the birds were slaughtered upon attaining the same stages of maturity (about 60% of the adult weight). Leghorn, Hubbard and naked neck, and Ross chickens were reared for 100, 56, and 42 days, respectively (Table S2). All chickens were slaughtered in a commercial slaughtering house.

For each genotype, ten breast samples (five males and females, each) were collected from the carcasses. From every breast sample 3 replicates (3 g of meat each) were analyzed both for raw and cooked meat. From each animal, two aliquots of approximately 100 g of the breast muscle were collected, vacuum-packaged, and frozen at -30 °C. After two weeks of storage, all the samples were thawed at 4 °C for 6 h, of which, half were analyzed immediately, and whereas the other aliquots were cooked by placing them in 250-mL glass bottles and boiling in a water bath at 80 °C for 30 min. With this cooking method the muscle samples reach a temperature of 60 °C ± 1. All samples

were stored at 5° C until completion of the analysis, which was carried out within 12 h from the time of thawing.

### ***Proton-transfer reaction–mass spectrometry (PTR–MS) and PTR–Time-of-Flight (ToF)-MS measurements***

Finely cut raw or cooked meat weighing 3 g was placed in a 250-mL glass bottle and equilibrated in a water bath at 35 °C for 30 min. The samples analyzed on day 1 were randomized across genotypes and replicates.

After temperature equilibration, the bottles were connected to a PTR–MS inlet, which was heated to 60 °C. The headspace flow rate was adjusted to 55 mL/min. A flow of 32 mL/min was directed into the PTR–MS *via* a teflon tubing of diameter 0.25 mm. A constant drift voltage and pressure of 600 V and  $2.19 \pm 0.01$  mbar, respectively, were maintained in the reaction chamber. A quadrupole mass spectrometer was used to analyze the masses of the samples, which were detected as the ion counts per second using a secondary electron multiplier. The ion intensities of the masses were converted to volume mixing ratio (ppb) values, based on the method employed by Lindinger et al.,(1998).

Mass spectral data were collected over a range of 20–160  $m/z$  with a dwell time of 200 ms. Although data of 5 cycles were collected for each measurement, only the average of cycles 2 to 4 was used for analysis.

Prior to analyzing each meat sample, the PTR-MS spectrum of a blank was acquired by sampling the contents from an empty bottle. This spectrum was subtracted from the sample spectra before data analysis (Aprea et al., 2007).

To identify the different molecules in raw and cooked meat, the samples were analyzed using PTR-ToF-MS (Ionicon GmbH, Innsbruck, Austria). Cubes of approximately 3 g were equilibrated at 30 °C for 30 min in a 250-mL screw-cap glass bottle, and the sample from the headspace was directed to the inlet of the high-sensitivity (**HS**) PTR-MS system (Ionicon GmbH) with an airflow rate of 75



mL/min. The temperatures of the inlet and drift chamber were maintained at 60 °C. The HS PTR-MS was operated at a standard  $E/N$  (ratio of the electric field strength across the reaction chamber,  $E$ , to the number density of the buffer gas,  $N$ , within the chamber) of 138 Td (1 Td =  $10^{-17}$  cm<sup>2</sup> V molecule<sup>-1</sup>) and measured in the “mass scan” mode. Thus, a complete mass spectrum in the range of 20–150 atomic mass units (amu) at a mass detection rate of 0.2 s mass<sup>-1</sup> was obtained in 26 s. the samples were analyzed independently in triplicates and seven mass scan cycles were used to measure each replicate. The background measurements were obtained by alternating seven scan cycles of the sample with seven scan cycles of a blank air sample. The individual components in the fraction were identified by directly comparing the mass spectra obtained from the samples with those available in NIST92 and Wiley5 libraries and of pure standards.

#### ***Chemical analysis and Fatty Acid determination***

The proximate composition of raw and cooked meat was determined using the AOAC method (Anastassiades et al., 2003). The fatty acid profile of raw and cooked breast meat was determined by analyzing the lipids extracted from approximately 5 g of the samples in a homogenizer using 20 mL of 2:1 chloroform/methanol (Folch et al., 1957), followed by filtration through a Whatman no. 1 filter paper. The fatty acids were identified in the form of their methyl esters using a Varian CP-3800 Gas Chromatograph (Milano, Italy) and a DB wax capillary column (25 mm ø, 30 m long). The percentages of fatty acids were calculated using the Chrom-Card software. Individual fatty acid methyl esters (FAME) were identified by referencing against the retention time of authentic FAME standards (Sigma–Aldrich, Bellefonte, PA, USA). The relative quantity of each fatty acid present in the meat was calculated using heneicosanoic acid (C21:0; Sigma–Aldrich) as the internal standard and expressed as mg/g of meat. The contents of the major classes of fatty acids were also calculated (SFA, monounsaturated fatty acid [**MUFA**], and PUFA of n-3 and n-6 series).

### ***Tocols content***

The content of tocols ( $\alpha$ -tocopherol [ $\alpha$ -T] and its isoforms  $\beta$ + $\gamma$  [ $\gamma$ -T] and  $\delta$  [ $\delta$ -T];  $\alpha$ ,  $\beta$ + $\gamma$  tocotrienol [ $\beta$ -T<sub>3</sub>,  $\gamma$ -T<sub>3</sub>]) in the meat was quantified using high-performance liquid chromatography (HPLC) (Hewavitharana et al., 2004). A single peak for  $\gamma$ -T was detected intermediate to those of  $\alpha$ -T and  $\delta$ -T on the chromatogram, and all three were quantified.

Briefly, 5 mL of distilled water and 4 mL of ethanol were added to 2 g of meat and vortexed for 10 s. Then, 4 mL of hexane containing butylated hydroxytoluene (BHT 200 mg/L) was added, and the mixture was carefully shaken and centrifuged. An aliquot of the supernatant (3 mL) was dried under a nitrogen stream and dissolved using 300  $\mu$ L of acetonitrile. From this, 50  $\mu$ L was injected into the HPLC system (Perkin Elmer Series 200) equipped with an autosampler system (model AS 950-10, Jasco, Tokyo, Japan) and a Synergy Hydro-RP column (4  $\mu$ m, 4.6  $\times$  100 mm; Phenomenex, Bologna, Italy). The peaks were identified using a Jasco FP-1525 FD detector (excitation and emission wavelengths of 295 and 328 nm, respectively) and quantified through external calibration curves obtained using increasing amounts of pure standards (Sigma–Aldrich, Steinheim, Germany) in ethanol.

### ***TBARS assay***

The extent of muscle lipid oxidation was evaluated using a spectrophotometer adjusted to 532 nm (Shimadzu Corp., UV-2550, Kyoto, Japan) according to the modified method of Ke et al., 1984 which measures the absorbance of TBARS. The oxidation products were quantified as equivalents of malondialdehyde (mg MDA/kg muscle) using a 1,1,3,3-tetraethoxypropane calibration curve.

## Statistical analysis

In addition to analyzing single qualitative traits, the differences in few major traits (n-3 and n-6 PUFA, tocopherols, VOC) between the samples of raw and cooked meat ( $\Delta$ ) were also evaluated. All traits were expressed as dry matter (**DM**) to prevent discrepancies due to changes in the moisture content of the raw and cooked samples.

Two different statistical approaches were employed. Firstly, univariate analysis was conducted using a linear model (Muenchen, 2012; StataCorp, 2015; Wang et al., 2016; StataCorp, 2017) in which, the effects of genotype, processing (raw, cooked), and their interactions were considered. The effect of sex was not found to be significant and, hence, omitted from the analysis. The differences were considered significant when  $P \leq 0.05$ . The statistical significance of values was reported only for major traits for simplification purposes.

Secondly, multivariate analysis was performed using an SPSS Statistics version 23. A set of variables was selected by inspecting the R-matrix and communalities (Righi et al., 2019, Twigg, 2010), and subjected to principal component analysis (**PCA**). We used the Kaiser–Meyer–Olkin and Bartlett tests to verify the adequacy of sampling and associated correlations. Principal components (**PC**) with eigenvalues  $> 1$  were retained and rotated using the varimax method. Only factor loadings with an absolute value greater than 0.5 were discussed (Menchetti. et al., 2018, Pituch and Stevens, 2015). Regression scores were calculated and additional groups for each PC (PC1 and PC2) were determined.

## RESULTS

### *Significance of effects*

The effects of genotype, processing, and their interactions on chicken meat are reported in Table S3. We found that the genotype of the poultry and the meat processing affected the content of majority of the analyzed compounds except butene, butanol, methyl, metacryl amide, and  $\alpha$ -T<sub>3</sub>. The interaction of genotype  $\times$  processing was always found to be significant, except in case of some VOC (butene, butanol, methyl furanone, metacrylic amide, and heptanal), fatty acids (14:0, 22:1*cis*9n-9, 20:5n-3, 22:5n-3, 22:6n-3), tocots ( $\delta$ -T<sub>3</sub>,  $\gamma$ -T<sub>3</sub>), and TBARS.

### *VOC development*

Twenty main VOC were identified and grouped into different chemical categories, which were as follows: aldehydes, alkadiene, alkenes, furans, amides, alcohols, and other compounds (Table 1 and Figure 1). Low amounts of aldehydes, alkadienes, alkenes, furans, amides, and alcohols (2.70, 0.42, 0.07, 0.06, 0.04, 0.02% of total VOC, respectively) were detected in the raw meat whereas other compounds (acetic acid, butyramide, methyl pyrrolidine, propionic ether) constituted 96.51%.

Cooking had strongly impacted VOC production. The content of VOC detected in cooked samples was 5.5 times that of raw meat (Figure 1). Aldehydes were the most important group of compounds detected in cooked meat of all genotypes (64.85% of the total VOC), of which, ethanal was the principal compound. Alkenes (2.44%, mainly 1,4-hexadiene), furans, amides, and alcohols were the components of VOC detected in minor amounts.

We determined that the genotype of the poultry was responsible for the major differences detected in the VOC profiles of their meats. Particularly, the highest increase in VOC (47 times) was

observed in the cooked samples of SG (Leghorn) meat compared to the raw ones. Few changes (3.5 and 3.8 times) were observed in the meats derived from other genotypes such as MG (Hubbard and Naked Neck, respectively), while the lowest content of VOC was detected in meats from FG (3 times).

#### ***Chemical composition and Fatty Acids profile***

Significant differences in the total fat content were observed in both raw and cooked samples of different poultry strains. The highest values for fat content were obtained from the meat samples belonging to the genotype Ross, followed by those of Hubbard and Naked Neck, while those of Leghorn were the lowest (Table S4).

It could be elucidated that cooking had affected the chemical composition of the meat mainly due to a reduction in its water content. However, the percentage of protein and ash was almost the same in raw and cooked samples whereas lipid loss was approximately 13-17% in cooked meat (data calculated from table S4).

Upon analyzing the PUFA profiles of cooked meat (Table 2), it was observed that, although values of total n-6 and n-3 fatty acids were lowest in meat of SG chickens, the loss of n-3 ( $\Delta$ ) was the highest. Contrarily, their amounts were higher in the meats of MG and FG strains, but their “absolute” losses were lower.

### ***Antioxidant content and Oxidative stability***

The TBARS values and the antioxidant content of raw and cooked meat and its variation ( $\Delta$ ) due to the cooking process are reported in Table 3. The Hubbard and Ross chickens showed the highest value of total tocopherols both in raw and cooked meat, while Leghorn had the lowest ones. The Hubbard and Ross birds exhibited the highest decrease in tocopherols (35.96 and 30.15  $\mu\text{g/g DM}$ , respectively) after cooking.

As expected, the TBARS increased in the cooked meat. The cooked Leghorn meat showed the highest increase in TBARS (about 4 times) and the most decrease in tocopherols (about 15 times) compared with the other poultry genotypes.

### ***Principal Component Analysis***

Multivariate analysis allowed us to determine the combined relation between the main traits of cooked meat (VOC,  $\Delta$  n-3, n-6, and content of tocopherols). PCA revealed two components, which accounted for 76.06% of the total variance (Table 4). The highest loadings of VOC and  $\Delta$  n-3 were observed in PC1 while PC2 was mainly influenced by  $\Delta$  n-6 and the content of tocopherols, and explained 43.04% and 33.02% of the total data variance, respectively.

The poultry strains were differentiated based on their PC scores on the factor maps determined through PC1 and PC2, according to their growth rates (Figure 2). The PC scores of SG samples (Leghorn) were concentrated in the fourth quadrant with positive PC1 and negative PC2 scores. The PC1 scores of FG samples (Ross) were negative and mainly located in the first and third quadrants. The scores of meats from Naked Neck and Hubbard, which had similar growth rates, were positive for both PC1 and PC2, since both were located in the second quadrant. Based on their position in the factor map (Figure 2), VOC production and  $\Delta$  n-3 were higher for the SG genotype whereas MG

was characterized by a higher  $\Delta n-6$ . Furthermore, small antioxidant and PUFA losses and the least production of VOC were observed in meats of the FG genotype.

## DISCUSSION

### *Volatile Organic Compounds of meat*

Due to its high ratio of unsaturated/saturated fatty acids and low fat content, poultry meat is considered as a healthy food for humans. It is known that PUFA, due to the presence of double bonds, are susceptible to lipid oxidation, which results in the generation of numerous degradation products such as aldehydes, ketones, alcohols, aliphatic hydrocarbons, acids, and esters (Campo et al., 2003, Menchetti et al. 2020), which are responsible for the development of flavors in meat. However, their concentrations depend mainly on the method of cooking because it affects their degree of degradation. For instance, compared to boiling, more compounds such as pyrazines, pyridines, pyrroles, and thiozoles are generated upon roasting, grilling, and frying chicken meat (Shahidi et al., 2020).

The present study had focused on the response of poultry genotypes with different growth rates to the cooking procedures based on the generation of VOC and content of fatty acids, tocopherols and TBARS. To our knowledge, this is one of the few studies to have explored the complex mechanisms, which regulate the relationship between cooking and the dynamics of PUFA, VOC, and antioxidants in meat of different poultry strains.

It was observed that the number of VOC in raw meat were few, which increased significantly upon cooking. The main VOC detected in cooked meat belonged to the group of aldehydes (Table 1), which are generally associated with the degradation of PUFA thus, confirming that the majority of VOC produced during cooking originate through lipolysis (Tornberg, 2005).

307 Pentanal and hexanal derive from the  $\beta$ -oxidation of fatty acids, mainly  $\alpha$ -linolenic acid (Del Pulgar  
308 et al., 2011). Guerrero-Legarreta et al., (2010) have reported that some aldehydes (hexanal,  
309 pentanal, heptanal, octanal, and nonanal) are responsible for off-odors. This suggests that the meat  
310 from Leghorn chickens was more susceptible to the production of unpleasant odors compared to  
311 that of the other genotypes.

312 The main aldehydes detected in our study (butanal, cyclohexadiene, heptanal, hexanal, nonanal,  
313 norbornane, octanal, and pentanal) are characterized by the presence of saturated chains of four to  
314 nine carbon atoms. These compounds are derived through the oxidation of PUFA and contain more  
315 than 20 carbon atoms (C20:4, C20:5, C22:6) whereas C18:2 and MUFA do not contain sufficiently  
316 long and saturated alkyl chains. However, the alkyl radicals from C20 PUFA may degrade the other  
317 fatty acids (C18:2 and C18:1) as a result of a chain reaction (Tornberg, 2005). Accordingly, our  
318 results demonstrated that the content of C20:4n-6 was lower in the meat of genotypes with a higher  
319 alkanal content (Leghorn), suggesting the manifestation of a specific oxidation mechanism.  
320 Probably, the generation of alkanals was not related to the other C20 PUFA because their content  
321 was less than 0.5 mg/g DM compared to C20:4 ( $\geq 1$  mg/g DM).

322 Despite alcohols being the second most abundant VOC detected in the meat samples, their  
323 incidence in the total VOC content was very scarce. In raw meat, alcohols are derived largely due to  
324 the metabolism of hetero-fermentative lactic acid bacteria. Schuster et al., (2018) have found that  
325 the alcohol levels increase in vacuum-packed lamb meat. However, it has been reported that a  
326 minor percentage of alcohol production also depends on the content of PUFA (Del Pulgar et al.,  
327 2011), thus explaining the presence of alcohols in cooked meat also.



### ***Fat content and Fatty Acid profiles of meat***

It is widely known that cooking largely affects the qualitative characteristics of meat. Part of these changes during cooking result from the degradation of substrates (such as PUFA) and concentration of nutrients due to water loss (Badiani et al., 2002).

Our results have confirmed that cooked meats are enriched with nutrients compared to raw ones. Additionally, cooking reduced also the fat content in meat, which mainly affected the content of PUFA and MUFA while that of SFA (Table 2) remained almost stable. This trend was observed in meat samples of birds belonging to all genotypes with variations in the extent of impact.

As mentioned above, chicken meat is highly susceptible to oxidation because it contains long-chain fatty acids with many double bonds. Consequently, losses, mainly in the content of n-3 PUFA, were observed during cooking. Similarly, here, the effect of cooking differed among meats of different genotypes, which is as follows:  $\Delta$  n-6 PUFA was found to be higher in Hubbard and Naked Neck (MG genotypes) compared to those of Ross (FG) and Leghorn (SG), whereas results contrary to these were observed in SG. It should be noted that, while raw meat of SG was not particularly high in the n-3 content (0.78 mg/g DM), its proportion was the highest (6.3 in LEG vs. 4.2, 3.7, and 3.5 % in HUB, NN and ROSS, respectively).

Cortinas et al., (2005) have suggested that a positive correlation exists between the content of PUFA and extent of lipid oxidation in chicken meat.

The PUFA profile of meat could also depends on rearing system, the presence of outdoor runs allow an additional intake of n-3, some authors (Dal Bosco et al., 2012, Castellini et al., 2016) demonstrated a higher percentage of PUFA, in particular of the n-3 series, in SG than FG when birds are reared free-range (Sirri et al., 2011) for the supplementary intake of vegetables, rich of ALA.

### ***Content of Antioxidants and Oxidative stability***

Although the same diet containing 50 mg/kg tocopheryl acetate was administered to the animals, the content of antioxidants in the raw meat of different genotypes was found to differ. This trend probably depends not only on the fat content of the meat, because tocopherols and retinol are fat-soluble antioxidants, but also on the kinetic behavior of the birds of different genotypes (Mattioli et al., 2017).

The same study has revealed that the antioxidant levels in poultry meat depend on the dietary intake and genetic strains of the birds (Mattioli et al., 2017). As stated previously, SG animals are very active, which affects their oxidative metabolism, resulting in high production of reactive oxygen species (ROS) and extensive consumption of antioxidants by the body.

In extensive rearing systems, this requirement is supported by the intake of natural antioxidants (tocols, vitamin C, and polyphenols) contained in the pastures (Mugnai et al., 2014). In our study, the absence of outdoor runs prevented the intake of additional antioxidants, probably resulting in the low content of tocopherols in SG meat compared to those of the FG and MG genotypes.

Animals of the SG genotype, which demonstrate a superior kinetic behavior and oxidative metabolism, tend to consume more antioxidants enabling them to counteract the oxidative thrust induced by the generation of ROS (Pisoschi and Pop, 2015).

Animals belonging to the MG strains (Hubbard and Naked Neck), being less kinetic, were able to preserve their tissue antioxidant stock better, while the highest content of antioxidant compounds was detected in meats of the FG genotype (Ross), which are recognized as more “static” animals (Dal Bosco et al., 2012).

The process of cooking increased the level of TBARS and decreased that of  $\alpha$ -T because the high temperature of cooking enhances oxidation. Rhee et al. (1996) compared raw and cooked meat of beef, pork, and poultry, and revealed that the TBARS content of cooked chicken drumstick was the highest due to its higher PUFA content. In our study, the lowest level of  $\alpha$ -T was detected in the raw

meat of the Leghorn genotype and, consequently, the highest levels of TBARS were detected after cooking it compared to that of the other genotypes. Selim et al., (2013) demonstrated the protective effect of a diet supplemented with vitamin E against PUFA autoxidation, which neutralized free radicals in both blood plasma and the muscles. Similarly, Castellini et al., (1998) demonstrated that the supplementation of  $\alpha$ -tocopheryl acetate could increase the contents of n-3 fatty acid in both raw and cooked rabbit meat. In agreement, Nardoia et al., (2017) detected lower levels of TBARS in the breast meat of chicken supplemented with vitamin E, grape skin, and grape pomace.

### ***Principal Component Analysis***

In addition to the trend obtained for single traits such as the content of total fat, fatty acids, antioxidants, TBARS, and VOC in poultry meat, we analyzed the specific relationships among these traits. Previous studies have investigated the effect of cooking on different attributes of meat quality. However, only a handful have evaluated the trade-off of substrates (fatty acids), protective substances (antioxidants), and oxidative and degradation products (TBARS, VOC) in the transition of poultry meat from a raw to cooked state (see Fig. 2).

It has been previously reported that raw meat is characterized by the presence of some antioxidants (mainly tocopherols) and PUFA (n-3 and n-6), which, during cooking, are oxidized easily, consequently producing radicals. Majorly, ROS and reactive nitrogen species (NOS) are the main radicals generated in this process, which, due to the presence of an unpaired electron, trigger an oxidative chain reaction with other compounds (proteins) or PUFA themselves. The effect of these radicals is modulated by the presence of antioxidants (which are primarily fat-soluble vitamins such as tocopherols) on the muscle enabling it to reduce this pro-oxidant action. If the concentration of tocopherols is not sufficient to counteract this oxidative process, the content of PUFA decreases due to VOC generation. In our trial, even the presence of PUFA series affected the amount of VOC produced during cooking, the results of which, are as follows: VOC generation was higher when n-3 PUFA

404 were used as substrates instead of n-6 due to a higher degree of unsaturation in the former  
405 (Scislowski et al. 2005). This confirmed that n-3 PUFA, due to the presence of many double bonds,  
406 were more susceptible to oxidation compared to their n-6 counterparts.

407 A multivariate approach permitted us to highlight the pattern of the aforementioned variables and  
408 distinguish the effect of cooking on meat from poultry with different genotypes. Tocols (PC2) were  
409 the most stable in meat of FG chickens, with the lowest variation in both n-6 and n-3, and having a  
410 low content of VOC. Contrarily, SG meat, which was characterized by low levels of antioxidants  
411 and the highest n-3 PUFA content, produced more VOC (PC1) compared to the others. The  
412 genotypes for MG, which were characterized by the presence of the highest amount of n-6 PUFA  
413 compared to n-3 PUFA, resulted in a lower production of VOC, which was consistent with this  
414 assumption.

415 In figure 3, a schematic representation of the mechanisms regulating the relationship between  
416 PUFA, VOC, and antioxidants in poultry meat during cooking has been reported.

417 Based on this perspective, more researches should focus on the stability of the PUFA profile during  
418 cooking. Our findings have indicated that these compounds undergo significant degradation in the  
419 meat after cooking, producing VOC and other substances (TBARS) with potential concerns for  
420 human health. Accordingly, greater attention should be paid to protect against PUFA, particularly in  
421 case of meats with high nutraceutical values.

## CONCLUSIONS

VOC such as alkanes, alkenes, aldehydes, ketones, alcohols, represent the principal products of the lipid oxidation and consequently are the main responsible of the reduction of the meat quality. A drastic increase in the content of VOC (aldehydes and alkanals) was observed in cooked chicken meat derived mainly due to the oxidation of PUFA, which was significantly correlated with the genotype of the organism. Maximum changes in the VOC content due to cooking were observed in the meat of breeds which could be defined as “more vulnerable”, with greater susceptibility to PUFA oxidation and associated with a lower content of antioxidants (Leghorn), resulting in a remarkable loss of n-3. Contrarily, losses of n-3 PUFA from the cooked meat of the FG genotypes were lower. Moreover, due to their lower activity, dietary antioxidants were rendered available for protecting the lipids in their meats.

Nowadays, increasing attention towards animal welfare has promoted the use of SG and MG genotypes, particularly in alternative rearing systems (organic, free-range) because these strains are more adaptable with a better welfare status in outdoor conditions. This adaptation is strictly related to the kinetic behavior of chickens. Hence, the relationship between activity (oxidative burst) and the nutritional quality of meat (PUFA profile) needs to be better defined. Our results have suggested that the use of these genotypes (MG and SG) requires a specific dietary protocol to balance the oxidative status of chickens *in vivo*, post-mortem, and, possibly during/after cooking. Accordingly, additional studies are needed to understand the mechanism for protecting n-3 PUFA from oxidation. For example, using dietary antioxidants (natural as per organic systems or synthetic, for the conventional) and/or finding more appropriate cooking methods to preserve the meat quality.

445 **Acknowledgement** Authors wish to thank Mr Giovanni Migni, for his contribute in animal  
446 handling.

447 **Funding** Partly funded by PRIN2017 – Prot. 2017S229WC, A.C.M. acknowledges the Erasmus+  
448 program for financing her student fellowship at the Wageningen University and Research.

449

Journal Pre-proof

## REFERENCE

450

- 451 Anastassiades, M., Lehotay, S. J., Štajnbaher, D., & Schenck, F. J. (2003). Fast and easy  
 452 multiresidue method employing acetonitrile extraction/partitioning and “dispersive solid-phase  
 453 extraction” for the determination of pesticide residues in produce. *Journal of AOAC*  
 454 *international*, 86(2), 412-431.
- 455 Angelo, A. S., Vercellotti, J. R., Legendre, M. G., VinnelT, C. H., Kuan, J. W., James Jr, C., &  
 456 Dupuy, H. P. (1987). Chemical and instrumental analyses of warmed-over flavor in  
 457 beef. *Journal of food science*, 52(5), 1163-1168.
- 458 Aprea, E., Biasioli, F., Märk, T. D., & Gasperi, F. (2007). PTR-MS study of esters in water and  
 459 water/ethanol solutions: Fragmentation patterns and partition coefficients. *International Journal*  
 460 *of Mass Spectrometry*, 262(1-2), 114-121.
- 461 Badiani, A., Stipa, S., Bitossi, F., Gatta, P. P., Vignola, G., & Chizzolini, R. (2002). Lipid  
 462 composition, retention and oxidation in fresh and completely trimmed beef muscles as affected  
 463 by common culinary practices. *Meat Science*, 60(2), 169-186.
- 464 Byrne, D. V., Bredie, W. L. P., Bak, L. S., Bertelsen, G., Martens, H., & Martens, M. (2001).  
 465 Sensory and chemical analysis of cooked porcine meat patties in relation to warmed-over flavour  
 466 and pre-slaughter stress. *Meat Science*, 59(3), 229-249.
- 467 Campo, M. M., Nute, G. R., Wood, J. D., Elmore, S. J., Mottram, D. S., & Enser, M. (2003).  
 468 Modelling the effect of fatty acids in odour development of cooked meat in vitro: part I—sensory  
 469 perception. *Meat science*, 63(3), 367-375.
- 470 Castellini, C., Dal Bosco, A., Bernardini, M., & Cyril, H. W. (1998). Effect of dietary vitamin E on  
 471 the oxidative stability of raw and cooked rabbit meat. *Meat Science*, 50(2), 153-161.
- 472 Castellini, C., Mugnai, C. A. N. D., & Dal Bosco, A. (2002). Effect of organic production system  
 473 on broiler carcass and meat quality. *Meat science*, 60(3), 219-225.
- 474 Castellini, C., Mugnai, C., Moscati, L., Mattioli, S., Guarino Amato, M., Cartoni Mancinelli, A., &  
 475 Dal Bosco, A. (2016). Adaptation to organic rearing system of eight different chicken genotypes:  
 476 behaviour, welfare and performance. *Italian Journal of Animal Science*, 15(1), 37-46.
- 477 Cortinas, L., Barroeta, A., Villaverde, C., Galobart, J., Guardiola, F., & Baucells, M. D. (2005).  
 478 Influence of the dietary polyunsaturation level on chicken meat quality: Lipid oxidation. *Poultry*  
 479 *Science*, 84(1), 48-55.
- 480 Dal Bosco, A., Mugnai, C., Mattioli, S., Rosati, A., Ruggeri, S., Ranucci, D., & Castellini, C.  
 481 (2016). Transfer of bioactive compounds from pasture to meat in organic free-range  
 482 chickens. *Poultry science*, 95(10), 2464-2471.
- 483 Dal Bosco, A., Mugnai, C., Ruggeri, S., Mattioli, S., & Castellini, C. (2012). Fatty acid composition  
 484 of meat and estimated indices of lipid metabolism in different poultry genotypes reared under  
 485 organic system. *Poultry Science*, 91(8), 2039-2045.

- 486 Del Pulgar, J. S., Soukoulis, C., Biasioli, F., Cappellin, L., García, C., Gasperi, F., Granitto, P.,  
487 Märk, T.D., Piasentier, E., and Schuhfried, E. (2011). Rapid characterization of dry cured ham  
488 produced following different PDOs by proton transfer reaction time of flight mass spectrometry  
489 (PTR-ToF-MS). *Talanta*, 85(1), 386-393.
- 490 Folch, J., Lees, M., & Stanley, G. S. (1957). A simple method for the isolation and purification of  
491 total lipides from animal tissues. *Journal of biological chemistry*, 226(1), 497-509.
- 492 Gill, C. O. (2007). Microbiological conditions of meats from large game animals and birds. *Meat*  
493 *science*, 77(2), 149-160.
- 494 Guerrero-Legarreta, I., & Hui, Y. H. (2010). Processed poultry products: a primer. *Handbook of*  
495 *poultry science and technology*, 2, 3-11.
- 496 Hewavitharana, A. K., Lanari, M. C., & Becu, C. (2004). Simultaneous determination of vitamin E  
497 homologs in chicken meat by liquid chromatography with fluorescence detection. *Journal of*  
498 *Chromatography A*, 1025(2), 313-317.
- 499 Holm, E. S., Adamsen, A. P. S., Feilberg, A., Schäfer, A., Løkke, M. M., & Petersen, M. A. (2013).  
500 Quality changes during storage of cooked and sliced meat products measured with PTR-MS and  
501 HS-GC-MS. *Meat science*, 95(2), 302-310.
- 502 Holm, E. S., Schäfer, A., Skov, T., Koch, A. G., & Petersen, M. A. (2012). Identification of  
503 chemical markers for the sensory shelf-life of saveloy. *Meat science*, 90(2), 314-322.
- 504 Leroy, F., Vasilopoulos, C., Van Hemelryck, S., Falony, G., & De Vuyst, L. (2009). Volatile  
505 analysis of spoiled, artisan-type, modified-atmosphere-packaged cooked ham stored under  
506 different temperatures. *Food Microbiology*, 26(1), 94-102.
- 507 Lindinger, W., & Jordan, A. (1998). Proton-transfer-reaction mass spectrometry (PTR-MS): on-line  
508 monitoring of volatile organic compounds at pptv levels. *Chemical Society Reviews*, 27(5), 347-  
509 375.
- 510 Mattioli, S., Dal Bosco, A., Ruggeri, S., Martino, M., Moscati, L., Pesca, C., & Castellini, C.  
511 (2017). Adaptive response to exercise of fast-growing and slow-growing chicken strains: Blood  
512 oxidative status and non-enzymatic antioxidant defense. *Poultry science*, 96(11), 4096-4102.
- 513 Menchetti, L., Canali, C., Castellini, C., Boiti, C., & Brecchia, G. (2018). The different effects of  
514 linseed and fish oil supplemented diets on insulin sensitivity of rabbit does during  
515 pregnancy. *Research in veterinary science*, 118, 126-133.
- 516 Muenchen, R. A. (2012). The popularity of data analysis software. URL <http://r4stats.com/popularity>.  
517
- 518 Mugnai, C., Sossidou, E. N., Dal Bosco, A., Ruggeri, S., Mattioli, S., & Castellini, C. (2014). The  
519 effects of husbandry system on the grass intake and egg nutritive characteristics of laying  
520 hens. *Journal of the Science of Food and Agriculture*, 94(3), 459-467.



- 521 Nardoia, M., Ruiz-Capillas, C., Herrero, A. M., Pintado, T., Jiménez Colmenero, F., Chamorro, S.,  
522 & Brenes, A. (2017). Effect of added grape seed and skin on chicken thigh patties during chilled  
523 storage.  
524 *International Journal of Food and Nutritional Science* 4 (1): 67-73
- 525 Pisoschi, A. M., & Pop, A. (2015). The role of antioxidants in the chemistry of oxidative stress: A  
526 review. *European journal of medicinal chemistry*, 97, 55-74.
- 527 Pituch, K. A., & Stevens, J. P. (2015). *Applied multivariate statistics for the social sciences:*  
528 *Analyses with SAS and IBM's SPSS*. Routledge.
- 529 Rhee, K. S., Anderson, L. M., & Sams, A. R. (1996). Lipid oxidation potential of beef, chicken, and  
530 pork. *Journal of Food Science*, 61(1), 8-12.
- 531 Righi, C., Menchetti, L., Orlandi, R., Moscati, L., Mancini, S., & Diverio, S. (2019). Welfare  
532 Assessment in Shelter Dogs by Using Physiological and Immunological  
533 Parameters. *Animals*, 9(6), 340.
- 534 Schuster, L., Franke, C., Silcock, P., Beauchamp, J., & Bremer, P. J. (2018). Development of a  
535 novel sample reuse approach to measure the impact of lean meat, bone and adipose tissue on the  
536 development of volatiles in vacuum-packed chilled lamb stored at 2° C for 15 days. *Meat*  
537 *science*, 145, 31-39.
- 538 Selim, N. A., Nada, S. A., Abdel-Salam, A. F., & Youssef, S. F. (2013). Evaluation of some natural  
539 antioxidant sources in broiler diets: 2-effect on chemical and microbiological quality of chilled  
540 and frozen broiler meat. *International Journal of Poultry Science*, 12(10), 572.
- 541 Shahidi, F., & Oh, W. Y. (2020). Lipid-derived flavor and off-flavor of traditional and functional  
542 foods: an overview. *Journal of Food Bioactives*, 10.
- 543 Sirri, F., Castellini, C., Bianchi, M., Petracci, M., Meluzzi, A., & Franchini, A. (2011). Effect of  
544 fast-, medium-and slow-growing strains on meat quality of chickens reared under the organic  
545 farming method. *Animal: an International Journal of Animal Bioscience*, 5(2), 312.
- 546 Sirri, F., Castellini, C., Roncarati, A., Franchini, A., & Meluzzi, A. (2010). Effect of feeding and  
547 genotype on the lipid profile of organic chicken meat. *European journal of lipid science and*  
548 *technology*, 112(9), 994-1002.
- 549 StataCorp, L. L. C. (2017). *Stata statistical software: Release 15* (2017). College Station, TX:  
550 StataCorp LP.  
551
- 552 StataCorp. 2015. *Stata Statistical Software: Release 14*. College Station, TX: StataCorp LP. 2015.
- 553 Tocher, D. R., Betancor, M. B., Sprague, M., Olsen, R. E., & Napier, J. A. (2019). Omega-3 long-  
554 chain polyunsaturated fatty acids, EPA and DHA: bridging the gap between supply and  
555 demand. *Nutrients*, 11(1), 89.
- 556 Tornberg, E. V. A. (2005). Effects of heat on meat proteins–Implications on structure and quality of  
557 meat products. *Meat science*, 70(3), 493-508.
- 558 Twigg, M. W. (2010). *Discovering Statistics using SPSS*. 3rd Edn A. Field 259× 190 mm. Pp. 856.  
559 Illustrated. 2009. SAGE Publications: London. ISBN: 97811847879073. *British Journal of*

560 Surgery, 97(6), 967-967.

561 Wang, Y., Yoon, B. J., & Qian, X. (2016, March). Co-segmentation of multiple images through  
562 random walk on graphs. In 2016 IEEE International Conference on Acoustics, Speech and Signal  
563 Processing(ICASSP) (pp.1811-1815).IEEE.

Journal Pre-proof

## TABLES

Table 1. Influence of genetic strain and processing (raw vs. cooked) on VOC composition (ppb) of chicken meat

Compounds	RAW				COOKED				Pooled SE <sup>1</sup>
Genotypes <sup>2</sup>	LEG	HUB	NN	ROSS	LEG	HUB	NN	ROSS	
Aldehydes									
2-Butenal	0.9	1.9	1.0	1.0	10.5	6.5	6.9	4.2	2.1
3-Ciano-propanal	0.3	0.3	0.4	0.3	38.9	13.2	17.3	9.0	4.8
Butanal	8.1	20.4	5.5	45.6	64.1	79.2	63.9	92.5	25.8
Butene	9.7	8.3	8.2	11.7	27.3	24.7	37.8	32.6	300.2
Cyclohexadiene	0.2	0.7	0.2	0.4	3.9	1.5	1.6	1.1	0.6
Ethanal	24.2	13.1	18.4	9.0	7736.2	5359.2	5613.1	3090.8	1493.9
Heptanal	0.1	0.2	0.2	0.4	2.0	1.2	1.3	0.9	0.6
Hexanal	0.3	0.4	0.4	0.1	46.8	12.7	22.2	9.0	11.9
Nonanal	0.2	0.3	0.5	0.2	14.3	3.3	5.0	1.9	3.0
Norbornane	0.2	0.3	0.3	0.2	9.5	6.1	8.0	5.1	4.3
Octanal	0.4	0.4	0.4	0.4	2.8	1.4	1.7	1.1	0.7
Pentanal	0.7	1.4	1.7	1.5	19.9	10.5	13.2	9.5	4.3
Xylenes	0.4	0.9	0.4	0.3	9.5	10.5	8.4	9.9	5.7
Alkadiene									
1,2-Butadiene	3.9	7.8	16	2.1	557.3	122.3	222.8	73.2	81.8
Alkenes									
1,4-Hexadiene	0.7	1.3	1.7	1.1	504.7	114.7	195.5	71.8	71.5
Furanas									
Methyl furanone	0.8	0.8	0.9	1.0	12.5	12.5	10.7	10.5	4.0
2-Pentyl Furan	0.2	0.2	0.1	0.2	1.7	1.1	1.4	1.0	0.8
Ammides									
Butyramide	0.1	0.1	0.1	0.1	1.1	0.1	1.1	1.5	0.3
Methachrylic amide	0.2	0.2	0.3	0.2	3.8	4.3	3.3	3.3	1.3

## Alcohols

Butanol	0.5	1.3	0.2	0.5	0.01	23.8	1.5	2.7	72.2
$\Sigma$ Aldehydes	45.7	48.6	37.6	71.1	7985.7	5530.0	5800.4	3267.6	693.7
$\Sigma$ Alkadiene	3.9	7.8	16	2.1	557.3	122.3	222.8	73.2	81.8
$\Sigma$ Alkenes	0.7	1.3	1.7	1.1	504.7	114.7	195.5	71.8	71.5
$\Sigma$ Furans	1	1	1	1.2	14.2	13.6	12.1	11.5	3.9
$\Sigma$ Amides	0.3	0.3	0.4	0.3	4.9	4.4	4.4	4.8	1.1
$\Sigma$ Alcohols	0.5	1.3	0.2	0.5	0.01	23.8	1.5	2.7	72.2
Other compounds	192.0	2417.0	2494.0	1695.0	2421.	2967.0	3462.0	1958.3	1120.2
Total VOC <sup>3</sup>	244.1	2477.3	2550.9	1771.3	11487.8	8775.8	9698.7	5389.9	3210.0

567

568 Abbreviations:<sup>1</sup>SE, standard error;569 <sup>2</sup> HUB, Hubbard; NN, Naked Neck; LEG, Leghorn; ROSS, Ross 308;570 <sup>3</sup>VOC, Volatile organic compounds;

Table 2. Influence of genetic strains and processing (raw vs. cooked) on the fatty acid profiles (mg/g DM) of poultry meat

Compounds <sup>1</sup>	RAW				COOKED				Pooled SE <sup>3</sup>
Genotypes <sup>2</sup>	LEG	HUB	NN	ROSS	LEG	HUB	NN	ROSS	
14:0	0.17	0.90	0.52	0.35	0.12	0.24	0.22	0.31	0.10
16:0	2.45	5.43	4.45	6.89	2.36	6.27	5.00	6.81	1.62
18:0	1.55	2.8	2.35	3.31	1.80	2.46	1.91	2.18	1.15
Σ SFA	4.17 <sup>a</sup>	9.13 <sup>c</sup>	7.32 <sup>b</sup>	10.55 <sup>c</sup>	4.28 <sup>a</sup>	8.97 <sup>bc</sup>	7.13 <sup>b</sup>	9.30 <sup>c</sup>	0.25
16:1cis9	0.85	2.21	1.94	2.86	0.58	1.61	1.32	2.3	0.20
18:1cis9 n-9	2.62	6.42	4.94	7.73	2.36	5.36	4.17	6.83	1.35
22:1cis9	0.15	0.31	0.34	0.30	0.17	0.35	0.35	0.37	0.01
Σ MUFA	3.73 <sup>a</sup>	9.14 <sup>bc</sup>	7.46 <sup>b</sup>	10.92 <sup>c</sup>	3.38 <sup>a</sup>	7.83 <sup>b</sup>	6.28 <sup>b</sup>	9.89 <sup>c</sup>	0.20
18:2n-6 (LA)	2.5	5.58	4.68	5.72	2.17	4.18	3.36	5.20	5.58
20:4n-6 (AA)	1.1	1.72	1.32	1.08	0.91	1.20	1.07	0.89	1.72
Σ n-6 PUFA	3.60 <sup>a</sup>	7.30 <sup>b</sup>	6.00 <sup>b</sup>	6.80 <sup>b</sup>	3.08 <sup>a</sup>	5.38 <sup>b</sup>	4.43 <sup>ab</sup>	6.09 <sup>b</sup>	1.38
Δ n-6 PUFA					0.52 <sup>a</sup>	1.92 <sup>b</sup>	1.57 <sup>b</sup>	0.71 <sup>a</sup>	0.10
18:3n-3 (ALA)	0.35	0.57	0.36	0.40	0.15	0.32	0.27	0.30	0.57
20:5n-3 (EPA)	0.22	0.35	0.28	0.39	0.15	0.30	0.22	0.34	0.35
22:5n-3 (DPA)	0.11	0.11	0.09	0.16	0.08	0.08	0.07	0.13	0.11
22:6n-3 (DHA)	0.10	0.11	0.09	0.08	0.06	0.09	0.07	0.05	0.11
Σ n-3 PUFA	0.78 <sup>a</sup>	1.14 <sup>b</sup>	0.82 <sup>a</sup>	1.03 <sup>b</sup>	0.44 <sup>a</sup>	0.79 <sup>b</sup>	0.63 <sup>ab</sup>	0.82 <sup>b</sup>	0.13
Δ n-3 PUFA					0.43 <sup>b</sup>	0.30 <sup>ab</sup>	0.23 <sup>a</sup>	0.20 <sup>a</sup>	0.08

Abbreviations:<sup>1</sup>SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; AA, arachidonic acid; LA, linolenic acid; ALA, α-linolenic acid; EPA, eicosapentaenoic acid; DPA, docosapentaenoic acid; DHA, docosahexaenoic acid; LCP, long chain PUFA;

<sup>2</sup>HUB, Hubbard; NN, Naked Neck 1; LEG: Leghorn; ROSS: Ross 308;

<sup>3</sup>SE: standard error;

<sup>a, b, c</sup> on the same row indicate the same type of processing (raw, cooked), with means of  $P < 0.05$ ;

Table 3. Influence of genetic strains and processing (raw vs. cooked) on the content of antioxidants ( $\mu\text{g/g DM}$ ) and oxidative stability ( $\text{mg/kg DM}$ ) of poultry meat

Compounds <sup>1</sup>	RAW				COOKED				Pooled SE <sup>3</sup>
Genotypes <sup>2</sup>	LEG	HUB	NN	ROSS	LEG	HUB	NN	ROSS	
$\alpha$ -T	10.73	34.46	16.01	32.98	0.98	4.22	2.93	3.35	1.73
$\delta$ -T	1.09	0.50	0.63	0.99	0.03	0.23	0.17	0.16	0.02
$\gamma$ -T	0.13	0.37	0.17	0.41	0.01	0.05	0.02	0.06	0.04
$\alpha$ -T <sub>3</sub>	5.28	6.03	8.65	6.07	0.24	1.23	0.74	0.52	0.65
$\delta$ -T <sub>3</sub>	1.13	0.17	1.38	0.87	0.30	0.03	0.17	0.20	0.14
$\gamma$ -T <sub>3</sub>	1.00	0.35	0.85	0.89	0.27	0.03	0.12	0.18	0.09
TBARS	0.34 <sup>a</sup>	0.70 <sup>b</sup>	0.54 <sup>a</sup>	1.12 <sup>c</sup>	1.28 <sup>c</sup>	0.95 <sup>b</sup>	0.65 <sup>a</sup>	1.55 <sup>c</sup>	0.14
$\Sigma$ -T	19.50 <sup>a</sup>	41.76 <sup>c</sup>	28.22 <sup>b</sup>	42.19 <sup>c</sup>	1.28 <sup>a</sup>	5.80 <sup>b</sup>	4.20 <sup>b</sup>	12.04 <sup>c</sup>	1.82
$\Delta$ tocols (raw-cooked)					18.22 <sup>a</sup>	35.96 <sup>c</sup>	24.02 <sup>b</sup>	30.15 <sup>c</sup>	1.11

Abbreviations: <sup>1</sup> $\alpha$ -T,  $\alpha$ -tocopherol;  $\delta$ -T,  $\delta$ -tocopherol;  $\gamma$ -T,  $\gamma$ -tocopherol;  $\alpha$ -T<sub>3</sub>,  $\alpha$ -tocotrienol;  $\delta$ -T<sub>3</sub>,  $\delta$ -tocotrienol;  $\gamma$ -T<sub>3</sub>,  $\gamma$ -tocotrienol; TBARS, Substances Reactive to Thiobarbituric acid;

<sup>2</sup>HUB, Hubbard; NN: Naked Neck 1; LEG: Leghorn; ROSS: Ross 308;

<sup>3</sup> SE, standard error;

<sup>a, b, c</sup> on the same row indicates the same type of processing (raw, cooked), with means of  $P < 0.05$ ;

590 Table 4. Principal components loadings, eigenvalue and variance


	PC1	PC2
VOC	0.71	-0.11
$\Delta$ n-3	0.59	0.05
$\Delta$ n-6	0.31	0.64
$\Delta$ tocols	0.20	0.75
Eigenvalue	1.72	1.32
% variance explained	43.04	33.02
Cumulative variance explained	76.06	

591  
592 Abbreviations: VOC, Volatile organic compounds;

593 Loadings  $\geq 0.50$  or  $\leq -0.50$  are presented in bold;

## FIGURES


Figure 1. Main VOC classes (%) in raw (a) and cooked (b) chicken meat



Abbreviations: VOC, Volatile Organic Compound;




Figure 2. Principal component analysis plot depicting the loadings and variables



Abbreviations: HUB, Hubbard; NN, Naked Neck; LEG, Leghorn; ROSS, Ross 308;

Figure 3. Scheme of PUFA, VOC and antioxidant dynamics in poultry meat during cooking process



Abbreviations: VOC, volatile organic compound; PUFA, polyunsaturated fatty acid; NOS, reactive nitrogen species; ROS, reactive oxygen species;

## SUPPLEMENTARY MATERIAL

Table S1. Formulation and chemical analysis of chicken feed

			Starter	Grower	Finisher
	<b>Ingredients</b>				
	Corn	%	53.92	55.95	53.11
	soybean meal 48%	%	30.23	24.67	15.69
	Extruded corn flour	%	5.08	8.90	11.45
	wheat	%	5.00	5.00	15.00
	Dicalcium phosphate	%	1.71	1.58	1.21
	Calcium carbonate	%	1.23	1.16	1.29
	Corn gluten 70	%	1.00	1.00	/
	Soybean oil	%	0.62	0.54	1.15
	Vitamin supplement	%	0.40	0.40	0.40
	salt	%	0.20	0.18	0.23
	Mineral supplement	%	0.16	0.16	0.11
	Sodium bicarbonate	%	0.15	0.15	0.15
	<b>Chemical composition</b>				
	Dry Matter (DM)	%	87.80	87.89	88
	Crude protein	% DM	24.01	22.16	18.41
	Lipids	% DM	3.99	3.98	4.55
	Crude fiber	% DM	3.48	3.58	3.60
	Ashes	% DM	6.92	6.43	5.78
	Metabolizable energy	kcal/kg	3245	3242	3295
	Vitamin A	U.I.	11385	11377	11364
	Vitamin E	Mg	36.43	36.41	36.37

647 Table S2. Productive performance of the four genotypes (slaughtered at about 60% adult weight)

648				
	Genotypes	slaughter age (d)	slaughter weight (kg)	daily gain (g)
649	Leghorn	100	1.3	13
650	Hubbard	56	2.0	35
651	Naked Neck	56	2.4	42
652	Ross	42	3.0	71
653				

654 Table S3. Significance of the effects on volatile compounds (A) total lipids and fatty acids (B) and  
 655 oxidative parameters (C)

A) Volatile Organic Compounds <sup>1</sup>	Genotype - G	Processing – P (raw vs. cooked)	GxP
Ethanal	***	***	***
Butadiene	***	***	***
Butene	ns	*	ns
Butenal	***	***	***
Butanal	***	***	**
Butanol	ns	ns	ns
Cyclohexadiene	***	***	***
Hexadiene	***	***	***
Cyano-propanal	***	***	***
Methyl furanone	ns	***	ns
Metacrylic amide	ns	***	ns
Pentanal	***	***	***
Butyramide	***	***	***
Norbomane	**	***	**
Hexanal	***	***	***
Xylenes	***	***	***
Heptanal	***	***	ns
Octanal	***	***	***
2-Pentyl furan	***	***	***
Nonanal	**	***	***
Σ Aldheydes	***	***	***
Σ Alchool	**	***	**
Σ Alkanales	***	***	***
Σ Ammine	***	***	***
Others	***	***	ns
Σ VOC	***	***	***

656 Abbreviations: <sup>1</sup>VOC: volatile organic compounds; \*\*\* P < 0.001; \*\* P < 0.01; P < 0.05. ns: not significant.

B) Fatty Acids <sup>2</sup>	Genotype – G	Processing – P (raw vs. cooked)	GxP
16:0	***	***	***
18:0	***	**	***
ΣSFA	**	**	***
16:1cis9	***	***	***
18:1cis9 n-9	***	***	***
22:1cis9	***	***	ns
ΣMUFA	***	***	***
20:4n-6 (AA)	***	***	ns
18:2n-6 (LA)	***	**	***
Σn-6	***	***	***
18:3n-3 (ALA)	***	***	***
20:5n-3 (EPA)	***	***	ns
22:5n-3 (DPA)	***	***	ns
22:6n-3 (DHA)	***	**	ns
Σn-3	***	**	***
n-3/n-6	***	***	***
ΣPUFA	***	***	***
ΣLCP	***	***	**

657

658

659

660

661

662

<sup>2</sup>SFA, saturated fatty acids; MUFA; monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; AA, arachidonic acid; LA, linolenic acid; ALA, α-linolenic acid; EPA, eicosapentaenoic acid; DPA, docosapentaenoic acid; DHA, docosahexaenoic acid; LCP, long chain polyunsaturated fatty acids;

\*\*\* P < 0.001; \*\* P < 0.01; P < 0.05. ns: not significant;

C) Oxidative parameters <sup>3</sup>	Genotype - G	Processing – P (raw vs. cooked)	GxP
$\alpha$ -T	***	***	***
$\delta$ -T	***	***	**
$\gamma$ -T	***	***	*
$\alpha$ -T <sub>3</sub>	ns	***	*
$\delta$ -T <sub>3</sub>	**	***	ns
$\gamma$ -T <sub>3</sub>	***	***	ns
TBARs	**	**	ns

663

664 Abbreviations:<sup>3</sup> $\alpha$ -T,  $\alpha$ -tocopherol;  $\gamma$ -T,  $\gamma$ -tocopherol;  $\delta$ -T,  $\delta$ -tocopherol;  $\alpha$ -T<sub>3</sub>,  $\alpha$ -tocotrienol;  $\gamma$ -T<sub>3</sub>,  $\gamma$ -tocotrienol;  $\delta$ -T<sub>3</sub>,  $\delta$ -tocotrienol; TBARs,  
665 substances reactive to the thiobarbituric acid;

666 \*\*\* P < 0.001; \*\* P < 0.01; P < 0.05. ns: not significant.

667

668 Table S4. Proximate composition of raw and cooked samples in the different poultry genotypes

Raw		LEG	HUB	NN	ROSS	Pooled SE
DM	%	23.85	24.23	23.92	24.20	1.82
Protein	g/100 g DM	93.92	92.28	92.39	92.31	8.23
Ash	“	4.61	4.62	5.02	4.17	0.32
Lipids	“	1.47 <sup>a</sup>	3.10 <sup>b</sup>	2.59 <sup>b</sup>	3.51 <sup>b</sup>	0.10
Cooked						
DM	%	33.71	34.82	35.18	34.30	3.20
Protein	g/100 g DM	93.89	92.48	92.61	92.71	9.02
Ash	“	4.89	4.88	5.12	4.37	0.40
Lipid	“	1.22 <sup>a</sup>	2.64 <sup>b</sup>	2.27 <sup>b</sup>	2.9 <sup>b</sup>	0.12

669

670 Abbreviations: HUB, Hubbard; NN, Naked Neck; LEG, Leghorn; ROSS, Ross 308;

671 a..b on the same row:  $P \leq 0.05$ .

672



# TABLES

Table 1. Influence of genetic strain and processing (raw vs. cooked) on VOC composition (ppb) of chicken meat

Compounds	RAW				COOKED				Pooled SE <sup>1</sup>
Genotypes <sup>2</sup>	LEG	HUB	NN	ROSS	LEG	HUB	NN	ROSS	
Aldehydes									
2-Butenal	0.9	1.9	1.0	1.0	10.5	6.5	6.9	4.2	2.1
3-Ciano-propanal	0.3	0.3	0.4	0.3	38.9	13.2	17.3	9.0	4.8
Butanal	8.1	20.4	5.5	45.6	64.1	79.2	63.9	92.5	25.8
Butene	9.7	8.3	8.2	11.7	27.3	24.7	37.8	32.6	300.2
Cyclohexadiene	0.2	0.7	0.2	0.4	3.9	1.5	1.6	1.1	0.6
Ethanal	24.2	13.1	18.4	9.0	7736.2	5359.2	5613.1	3090.8	1493.9
Heptanal	0.1	0.2	0.2	0.4	2.0	1.2	1.3	0.9	0.6
Hexanal	0.3	0.4	0.4	0.1	46.8	12.7	22.2	9.0	11.9
Nonanal	0.2	0.3	0.5	0.2	14.3	3.3	5.0	1.9	3.0
Norbornane	0.2	0.3	0.3	0.2	9.5	6.1	8.0	5.1	4.3
Octanal	0.4	0.4	0.4	0.4	2.8	1.4	1.7	1.1	0.7
Pentanal	0.7	1.4	1.7	1.5	19.9	10.5	13.2	9.5	4.3
Xylenes	0.4	0.9	0.4	0.3	9.5	10.5	8.4	9.9	5.7
Alkadiene									
1,2-Butadiene	3.9	7.8	16	2.1	557.3	122.3	222.8	73.2	81.8
Alkenes									
1,4-Hexadiene	0.7	1.3	1.7	1.1	504.7	114.7	195.5	71.8	71.5
Furanas									
Methyl furanone	0.8	0.8	0.9	1.0	12.5	12.5	10.7	10.5	4.0
2-Pentyl Furan	0.2	0.2	0.1	0.2	1.7	1.1	1.4	1.0	0.8
Ammides									
Butyramide	0.1	0.1	0.1	0.1	1.1	0.1	1.1	1.5	0.3
Methachrylic amide	0.2	0.2	0.3	0.2	3.8	4.3	3.3	3.3	1.3

## Alcohols

Butanol	0.5	1.3	0.2	0.5	0.01	23.8	1.5	2.7	72.2
$\Sigma$ Aldehydes	45.7	48.6	37.6	71.1	7985.7	5530.0	5800.4	3267.6	693.7
$\Sigma$ Alkadiene	3.9	7.8	16	2.1	557.3	122.3	222.8	73.2	81.8
$\Sigma$ Alkenes	0.7	1.3	1.7	1.1	504.7	114.7	195.5	71.8	71.5
$\Sigma$ Furans	1	1	1	1.2	14.2	13.6	12.1	11.5	3.9
$\Sigma$ Ammides	0.3	0.3	0.4	0.3	4.9	4.4	4.4	4.8	1.1
$\Sigma$ Alcohols	0.5	1.3	0.2	0.5	0.01	23.8	1.5	2.7	72.2
Other compounds	192.0	2417.0	2494.0	1695.0	2421.	2967.0	3462.0	1958.3	1120.2
Total VOC <sup>3</sup>	244.1	2477.3	2550.9	1771.3	11487.8	8775.8	9698.7	5389.9	3210.0

4

5 Abbreviations:<sup>1</sup>SE, standard error;6 <sup>2</sup>HUB, Hubbard; NN, Naked Neck; LEG, Leghorn; ROSS, Ross 308;7 <sup>3</sup>VOC, Volatile organic compounds;

8 Table 2. Influence of genetic strains and processing (raw vs. cooked) on the fatty acid profiles (mg/g  
9 DM) of poultry meat

Compounds <sup>1</sup>	RAW				COOKED				Pooled SE <sup>3</sup>
Genotypes <sup>2</sup>	LEG	HUB	NN	ROSS	LEG	HUB	NN	ROSS	
14:0	0.17	0.90	0.52	0.35	0.12	0.24	0.22	0.31	0.10
16:0	2.45	5.43	4.45	6.89	2.36	6.27	5.00	6.81	1.62
18:0	1.55	2.8	2.35	3.31	1.80	2.46	1.91	2.18	1.15
Σ SFA	4.17 <sup>a</sup>	9.13 <sup>c</sup>	7.32 <sup>b</sup>	10.55 <sup>c</sup>	4.28 <sup>a</sup>	8.97 <sup>bc</sup>	7.13 <sup>b</sup>	9.30 <sup>c</sup>	0.25
16:1cis9	0.85	2.21	1.94	2.86	0.58	1.61	1.32	2.3	0.20
18:1cis9 n-9	2.62	6.42	4.94	7.73	2.36	5.36	4.17	6.83	1.35
22:1cis9	0.15	0.31	0.34	0.30	0.17	0.35	0.35	0.37	0.01
Σ MUFA	3.73 <sup>a</sup>	9.14 <sup>bc</sup>	7.46 <sup>b</sup>	10.92 <sup>c</sup>	3.38 <sup>a</sup>	7.83 <sup>b</sup>	6.28 <sup>b</sup>	9.89 <sup>c</sup>	0.20
18:2n-6 (LA)	2.5	5.58	4.68	5.72	2.17	4.18	3.36	5.20	5.58
20:4n-6 (AA)	1.1	1.72	1.32	1.08	0.91	1.20	1.07	0.89	1.72
Σ n-6 PUFA	3.60 <sup>a</sup>	7.30 <sup>b</sup>	6.00 <sup>b</sup>	6.80 <sup>b</sup>	3.08 <sup>a</sup>	5.38 <sup>b</sup>	4.43 <sup>ab</sup>	6.09 <sup>b</sup>	1.38
Δ n-6 PUFA					0.52 <sup>a</sup>	1.92 <sup>b</sup>	1.57 <sup>b</sup>	0.71 <sup>a</sup>	0.10
18:3n-3 (ALA)	0.35	0.57	0.36	0.40	0.15	0.32	0.27	0.30	0.57
20:5n-3 (EPA)	0.22	0.35	0.28	0.39	0.15	0.30	0.22	0.34	0.35
22:5n-3 (DPA)	0.11	0.11	0.09	0.16	0.08	0.08	0.07	0.13	0.11
22:6n-3 (DHA)	0.10	0.11	0.09	0.08	0.06	0.09	0.07	0.05	0.11
Σ n-3 PUFA	0.78 <sup>a</sup>	1.14 <sup>b</sup>	0.82 <sup>a</sup>	1.03 <sup>b</sup>	0.44 <sup>a</sup>	0.79 <sup>b</sup>	0.63 <sup>ab</sup>	0.82 <sup>b</sup>	0.13
Δ n-3 PUFA					0.43 <sup>b</sup>	0.30 <sup>ab</sup>	0.23 <sup>a</sup>	0.20 <sup>a</sup>	0.08

10 Abbreviations:<sup>1</sup>SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; AA, arachidonic acid; LA,  
11 linolenic acid; ALA, α-linolenic acid; EPA, eicosapentaenoic acid; DPA, docosapentaenoic acid; DHA, docosahexaenoic acid; LCP, long chain  
12 PUFA;

13 <sup>2</sup>HUB, Hubbard; NN, Naked Neck 1; LEG: Leghorn; ROSS: Ross 308;

14 <sup>3</sup>SE: standard error;

15 <sup>a, b, c</sup> on the same row indicate the same type of processing (raw, cooked), with means of  $P < 0.05$ ;

16  
17

Table 3. Influence of genetic strains and processing (raw vs. cooked) on the content of antioxidants ( $\mu\text{g/g DM}$ ) and oxidative stability ( $\text{mg/kg DM}$ ) of poultry meat

Compounds <sup>1</sup>	RAW				COOKED				Pooled SE <sup>3</sup>
Genotypes <sup>2</sup>	LEG	HUB	NN	ROSS	LEG	HUB	NN	ROSS	
$\alpha$ -T	10.73	34.46	16.01	32.98	0.98	4.22	2.93	3.35	1.73
$\delta$ -T	1.09	0.50	0.63	0.99	0.03	0.23	0.17	0.16	0.02
$\gamma$ -T	0.13	0.37	0.17	0.41	0.01	0.05	0.02	0.06	0.04
$\alpha$ -T <sub>3</sub>	5.28	6.03	8.65	6.07	0.24	1.23	0.74	0.52	0.65
$\delta$ -T <sub>3</sub>	1.13	0.17	1.38	0.87	0.30	0.03	0.17	0.20	0.14
$\gamma$ -T <sub>3</sub>	1.00	0.35	0.85	0.89	0.27	0.03	0.12	0.18	0.09
TBARS	0.34 <sup>a</sup>	0.70 <sup>b</sup>	0.54 <sup>a</sup>	1.12 <sup>c</sup>	1.28 <sup>c</sup>	0.95 <sup>b</sup>	0.65 <sup>a</sup>	1.55 <sup>c</sup>	0.14
$\Sigma$ -T	19.50 <sup>a</sup>	41.76 <sup>c</sup>	28.22 <sup>b</sup>	42.19 <sup>c</sup>	1.28 <sup>a</sup>	5.80 <sup>b</sup>	4.20 <sup>b</sup>	12.04 <sup>c</sup>	1.82
$\Delta$ tocols (raw-cooked)					18.22 <sup>a</sup>	35.96 <sup>c</sup>	24.02 <sup>b</sup>	30.15 <sup>c</sup>	1.11

Abbreviations: <sup>1</sup> $\alpha$ -T,  $\alpha$ -tocopherol;  $\delta$ -T,  $\delta$ -tocopherol;  $\gamma$ -T,  $\gamma$ -tocopherol;  $\alpha$ -T<sub>3</sub>,  $\alpha$ -tocotrienol;  $\delta$ -T<sub>3</sub>,  $\delta$ -tocotrienol;  $\gamma$ -T<sub>3</sub>,  $\gamma$ -tocotrienol; TBARS, Substances Reactive to Thiobarbituric acid;

<sup>2</sup>HUB, Hubbard; NN: Naked Neck 1; LEG: Leghorn; ROSS: Ross 308;

<sup>3</sup> SE, standard error;

<sup>a, b, c</sup> on the same row indicates the same type of processing (raw, cooked), with means of  $P < 0.05$ ;


Table 4. Principal components loadings, eigenvalue and variance

	PC1	PC2
VOC	0.71	-0.11
$\Delta$ n-3	0.59	0.05
$\Delta$ n-6	0.31	0.64
$\Delta$ tocols	0.20	0.75
Eigenvalue	1.72	1.32
% variance explained	43.04	33.02
Cumulative variance explained	76.06	

Abbreviations: VOC, Volatile organic compounds;


Loadings  $\geq 0.50$  or  $\leq -0.50$  are presented in bold;

Figure 1. Main VOC classes (%) in raw (a) and cooked (b) chicken meat




Abbreviations: VOC, Volatile Organic Compound;

Figure 2. Principal component analysis plot depicting the loadings and variables



Abbreviations: HUB, Hubbard; NN, Naked Neck; LEG, Leghorn; ROSS, Ross 308;

Figure 3. Scheme of PUFA, VOC and antioxidant dynamics in poultry meat during cooking process



Abbreviations: VOC, volatile organic compound; PUFA, polyunsaturated fatty acid; NOS, reactive nitrogen species; ROS, reactive oxygen species;