

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Feeding European sea bass with increasing dietary fibre levels: Impact on growth, blood biochemistry, gut histology, gut evacuation

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Bonvini, E., Bonaldo, A., Parma, L., Mandrioli, L., Sirri, R., Grandi, M., et al. (2018). Feeding European sea bass with increasing dietary fibre levels: Impact on growth, blood biochemistry, gut histology, gut evacuation. AQUACULTURE, 494, 1-9 [10.1016/j.aquaculture.2018.05.017].

Availability: This version is available at: https://hdl.handle.net/11585/672940 since: 2021-02-25

Published:

DOI: http://doi.org/10.1016/j.aquaculture.2018.05.017

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

1	
2	
3	This is the final peer-reviewed accepted manuscript of:
4 5	Feeding European sea bass with increasing dietary fibre levels: impact on growth, blood biochemistry, gut histology, gut evacuation
6 7	The final published version is available online at: <u>https://doi.org/10.1016/j.aquaculture.2018.05.017</u>
8	
9	
10	Rights / License:
11 12 13	The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>) When citing, please refer to the published version.

25	Feeding European sea bass with increasing dietary fibre levels: impact on growth,
26	blood biochemistry, gut histology, gut evacuation
27	
28	Erika Bonvini <sup>a</sup> , Alessio Bonaldo <sup>a</sup> , Luca Parma <sup>a</sup> *, Luciana Mandrioli <sup>a</sup> , Rubina Sirri <sup>a</sup> ,
29	Monica Grandi <sup>a</sup> , Ramon Fontanillas <sup>c</sup> , Cinzia Viroli <sup>b</sup> , Pier Paolo Gatta <sup>a</sup>
30	
31	
32	<sup>a</sup> Department of Veterinary Medical Sciences, University of Bologna, Via Tolara di Sopra
33	50, 40064 Ozzano Emilia, Bologna, Italy
34	<sup>b</sup> Department of Statistical Sciences "Paolo Fortunati", University of Bologna, Via delle
35	Belle Arti 41, 40126 Bologna, Italy
36	<sup>c</sup> Skretting Aquaculture Research Centre, Sjøhagen 3, 4016 Stavanger, Norway
37	
38	
39	*Corresponding author: Luca Parma, Department of Veterinary Medical Sciences,
40	University of Bologna, Viale Vespucci 2, 47042 Cesenatico, FC, Italy. Tel.: +39 0547
41	338931; <i>Fax</i> : +39 0547 338941
42	E-mail address: <u>luca.parma@unibo.it</u> (L. Parma)
43	
44	
45	
46	
47	
48	

49 Abstract

50

51 Changing trends in fish feed formulation, with progressively higher inclusion levels of 52 plant ingredients, are invariably introducing more fibre despite the fact that this 53 component cannot be utilized by most fish. The effects of increasing insoluble dietary 54 fibre level on growth, nutrient utilization, blood parameters and gut health in European 55 sea bass (*Dicentrarchus labrax* L.) were studied over a period of 117 days. Moreover, 56 investigation on digesta transit time through gastrointestinal evacuation pattern and 57 digesta characteristics (moisture of digesta) were studied. Five iso-proteic diets were 58 formulated to contain increasing insoluble fibre levels, neutral detergent fibre, NDF (7.2, 59 8.9, 11.5, 13.1 and 15.5%) derived by the inclusion of sunflower hulls and soybean hulls. 60 No significant differences due to fibre inclusion levels were observed in final body 61 weight, specific growth rate, feed intake, feed conversion rate, protein and lipid 62 efficiency. No significant differences in serum total protein, glucose, triglycerides, 63 alkaline phosphatase and inorganic phosphorous were found. All the histological sections 64 showed normal intestinal architecture, and inflammatory and/or degenerative changes 65 were not present in any histological section from all subjects examined. The investigation 66 into gastrointestinal evacuation pattern revealed no significant differences between 67 treatments, however higher dietary fibre levels seem to increase the time required to 68 empty the stomach while the time required to empty 90% of the hindgut content was 69 similar in all the treatments: around 46-47 h. No differences were found between diets in 70 the moisture content of digesta along the digestive tracts. We can conclude that the 71 different insoluble fibre levels tested in this trial have no effects on overall performances 72 and feed efficiency in European sea bass. Results from blood biochemistry profile and

histology confirm good nutritional and health status of fish under all feeding treatments.
The inclusion of fibre had no influence on digesta transit time. In formulation of feed for
the on-growing of European sea bass insoluble fibre derived from sunflower hulls and
soybean hulls can be included at a level of up to 15.5 %.

77

78 Keywords: European sea bass, fibre, growth, blood biochemistry, gut histology, gut
79 evacuation

80

### 81 **1.Introduction**

82

83 Changing trends in fish feed formulation, with progressively higher inclusion levels of plant ingredients, are invariably introducing more fibre in aquafeeds. Cellulose and other 84 85 fibrous carbohydrates are found in the structural components of plants and are indigestible 86 to monogastric (simple-stomach) animals, such as fish. Fibre can be divided into soluble 87 and insoluble (according to their extractability in a neutral buffer solution) that may have 88 different antinutritive effects in endothermic animals (Dalsgaard et al., 2016). Soluble 89 fibres in mammals, poultry and some fish species tend to increase digesta viscosity and retard absorption of nutrients (Krogdahl et al., 2005; Amirkolaie et al., 2005), while 90 91 insoluble fibres appear to act largely as physical bulking agents (Bach Knudsen 2001) 92 and tend to increase digesta transit, resulting in reduced absorption time (Krogdahl et al., 93 2005). Regarding insoluble dietary fibres, cellulose seems to be a relatively inert dietary 94 component that may reduce the dry matter digestibility in Atlantic salmon (Salmo salar), 95 Atlantic cod (Gadus morhua L.), Nile tilapia (Oreochromis niloticus L.) and rainbow 96 trout (Oncorhynchus mykiss) (Amirkolaie et al. 2005; Hansen & Storebakken 2007;

97 Kraugerud et al. 2007; Glencross 2009; Glencross et al. 2012; Lekva et al. 2010). 98 Increasing dietary levels of cellulose have also negatively affected lipid digestibility in 99 cod (Lekva et al. 2010) and energy and protein digestibility in rainbow trout (Glencross 100 2009; Glencross et al. 2012). Moreover, as reported by Altan and Korkut (2011), low dietary concentrations of dietary fibre (3-5 %) may have a beneficial effect on fish 101 102 growth, but high dietary fibre (>8 %), on the contrary, may decrease dry matter 103 digestibility of the diet and reduce the availability of other nutrients. Some discrepancies 104 between values of maximum dietary fibre level have been reported for fish: less than 7 % 105 (Altan and Korkut, 2011); less than 8 % (Eusebio et al., 2004); as low as possible and not 106 exceeding 10 % (NRC, 2011). Information about the optimal percentage of fibre inclusion 107 in practical feed formulation for the on-growing of the European sea bass (Dicentrarchus 108 labrax L.) is scarce but Kousoulaki et al. (2015) reported that the fibre content in 109 commercial feeds for this species can range between 1.5-3.2 %. The aim of the present 110 study was to assess the effects of increasing dietary insoluble fibre level derived from 111 sunflower hulls and soybean hulls on growth, nutrient utilisation, blood parameters, gut 112 health, gastrointestinal evacuation pattern and digesta characteristics in European sea bass 113 over 117 days, in order to identify the maximum amount of dietary fibre inclusion without 114 negatively affecting zootechnical performance and health.

115

#### 116 **2. Materials and methods**

117

118 2.1 Experimental diets

Five iso-proteic diets were formulated to contain increasing insoluble fibre levels
(Neutral detergent fibre, NDF 7.2, 8.9, 11.5, 13.1 and 15.5%; F7.2, F8.9, F11.5, F13.1.

121 and F15.5, respectively). Diets were formulated with fishmeal and with a mixture of 122 vegetable ingredients currently used in aquafeed (Parma et al., 2016; Bonvini et al., 2017). 123 The fibre content was increased by increasing levels of a combination of sunflower hulls 124 and soybean hulls to provide same proportion of fibre from each ingredient. Sunflower 125 hulls and soybean hulls were chosen because they are one of the most used and available 126 raw material for feed production. Lipid levels were slightly increased at increasing fibre 127 content in order to compensate for the loss of available energy due to the higher fibre 128 content. The diets were produced by extrusion process by Skretting Aquaculture Research 129 Centre, Stavanger, Norway. The diameter of the pellet was 4 mm. Ingredients and 130 proximate composition of the experimental diets are presented in Table 1.

131

#### 132 2.2 Fish and feeding trial

133

134 The experiment was carried out at the Laboratory of Aquaculture, Department of 135 Veterinary Medical Sciences of the University of Bologna, Cesenatico, Italy. European 136 sea bass juveniles were obtained from Maricoltura Mattinatese (Mattinata, Foggia, Italy). 137 At the beginning of the trial, 60 fish (initial average weight:  $69.4 \pm 2.3$  g) per tank were 138 randomly distributed into fifteen 900 L square tanks with a conical base. Each diet was 139 administered to triplicate groups, assigned in a completely random manner, over 117 140 days. Tanks were provided with natural seawater and connected to a closed recirculating 141 system (overall water volume: 18 m<sup>3</sup>). The rearing system consisted of a mechanical sand 142 filter (PTK 1200, Astralpool, Barcelona, Spain), ultraviolet lights (PE 25mJ/cm<sup>2</sup>: 32m<sup>3</sup> h<sup>-</sup> <sup>1</sup>, Blaufish, Barcelona, Spain) and a biofilter (PTK 1200, Astralpool, Barcelona, Spain). 143 144 The water exchange rate within each tank was 100% every hour, while the overall water 145 renewal amount in the system was 5 % daily. During the trial, the temperature was kept 146 constant at  $22 \pm 1.0$  °C and the photoperiod was maintained at 12 h light and 12 h dark 147 through artificial light. The oxygen level was kept constant  $(8.0 \pm 1.0 \text{ mg L}^{-1})$  by a liquid 148 oxygen system regulated by a software programme (B&G Sinergia snc, Chioggia, Italy). 149 Ammonia (total ammonia nitrogen  $\leq 0.1 \text{ mg L}^{-1}$ ), nitrite ( $\leq 0.2 \text{ mg L}^{-1}$ ) and salinity (25 g L<sup>-1</sup>) were daily monitored spectrophotometrically (Spectroquant Nova 60, Merck, Lab 150 151 business, Darmstadt, Germany). Sodium bicarbonate was added on a daily basis to keep 152 pH constant at 7.8–8.0. Feed was provided to apparent satiation by oversupplying the feed 153 by automatic feeders by approximately 10% of the daily ration, twice a day for six days 154 a week, while one meal was supplied on Sundays, as reported by Mongile et al. (2014). 155 Each meal lasted 1 hour, after which the uneaten pellets of each tank were gathered, dried 156 overnight at 105°C, and their weight was deducted for overall calculation.

157

158 2.3 Sampling

159

160 At the beginning and at the end of the experiment, all the fish in each tank were anaesthetised by 2-phenoxyethanol at 300 mg L<sup>-1</sup> and individually weighed. Specific 161 growth rate (SGR), voluntary feed intake (VFI) and feed conversion rate (FCR) were 162 163 calculated. The proximate composition of the carcasses was determined at the beginning 164 of the trial on a pooled sample of 10 fish and on a pooled sample of 5 fish per tank at the 165 end of the trial. Protein efficiency rate (PER), gross protein efficiency (GPE) and gross 166 lipid efficiency (GLE) were calculated. Furthermore, at the end of the trial, wet weight, 167 viscera, liver and mesenteric fat weight were individually recorded for 10 fish per tank to 168 determine viscerosomatic index (VSI), hepatosomatic index (HSI) and mesenteric fat 169 index (MFI). At the end of the trial, five fish per tank (15 fish per dietary treatment) were 170 sampled for intestine histology examination. At the end of the trial, the fish left were kept 171 in the same rearing and feeding conditions for three more days and then were sampled to 172 perform blood analyses of serum total protein (TP), triglycerides (TRIG), glucose (GLU), 173 alkaline phosphatase (ALP) and inorganic phosphorus (P). Blood from 4 fish per tank 174 was collected 5 h postprandial from the caudal vein. Samples were then centrifuged (3000 175 g for 10 min at 4°C), serum aliquots were stored at 4°C and analysed during the same day 176 according to Bonvini et al. (2017). All experimental procedures were evaluated and 177 approved by the Ethical-Scientific Committee for Animal Experimentation of the 178 University of Bologna, in accordance with European directive 2010/63/UE on the 179 protection of animals used for scientific purposes.

180

## 181 2.4 Calculations

182

### 183 The formulae employed were as follows:

184 Specific growth rate (SGR) (% day<sup>-1</sup>) =  $100 * (\ln FBW - \ln IBW) / days$  (where FBW and 185 IBW represent the final and the initial body weights). Voluntary Feed Intake (VFI) (g 186 feed fish<sup>-1</sup>) = g feed ingested / fish number. Feed conversion ratio (FCR) = feed intake / 187 weight gain. Viscerosomatic index (VSI) (%) = 100 \* (viscera weight / body weight). 188 Hepatosomatic index (HSI) (%) = 100 \* (liver weight / body weight). Mesenteric fat index 189 (MFI) (%) = 100 \* (mesenteric fat weight / body weight). Protein efficiency rate (PER) 190 = (FBW - IBW) / protein intake. Gross protein efficiency (GPE) (%) = 100 \* [(% final 191 body protein \* FBW) - (% initial body protein \* IBW)] / total protein intake fish. Gross lipid efficiency (GLE) (%) = 100 \* [(% final body lipid \* FBW) - (% initial body lipid \*
IBW)] / total lipid intake fish.

194

195 2.5 Histology

196

197 After euthanasia of 5 fish per tank, the gut was removed and the intestine was divided 198 into two segments (midgut and hindgut). From each segment, a 5 mm-long piece was 199 sectioned and fixed in 10% buffered formalin. Samples were processed for routine 200 histology to obtain a transversal section, which was stained with haematoxylin and eosin 201 (H&E). Sections were evaluated blind under a light microscope (Nikon Eclipse 80i, 202 Nikon Corporation, Japan) to verify the preservation of the normal intestinal architecture. 203 In particular, the histological investigation was focused on the main cell constituents of 204 the mucosal layer (goblet cells, supranuclear absorption vacuoles in the enterocytes), 205 capillary within the intestinal folds, lymphoplasmacytic cells within lamina propria 206 (GALT-like tissue). Moreover, any possible degenerative and diet adaptive induced 207 changes were taken into consideration. Photographs of the sections were made with a 208 Nikon Eclipse 80i microscope (Nikon, Tokyo, Japan) equipped with a Nikon Digital Sight 209 SD-MS camera and the Nikon software NIS-Elements. Adobe Photoshop CS3 Extended 210 was used for the final photographic preparation without altering the original integrity of 211 the pictures.

212

213 2.6 Gastrointestinal evacuation, time and digesta characteristics

215 Following the feeding trial, sampling for gastrointestinal evacuation pattern and 216 digesta characteristics was conducted. We adopted the approach used in the studies by 217 Adamidou et al. (2009) and Nikolopoulou et al. (2011). Fish were kept fasting for 72 h 218 before being fed to make sure that the gastrointestinal tract was empty. Four fish per tank 219 were sacrificed at 1, 4, 10, 16, 28 and 48 h after feeding a single meal to satiation. Each 220 sampled fish was weighed, then the abdominal cavity was opened and the digestive tract 221 carefully removed and separated into three parts: stomach, midgut and hindgut. Midgut 222 was defined as the section from the pyloric sphincter to the ileorectal valve and hindgut 223 from the ileorectal valve to the anus. Stomach and intestinal contents from the above 224 sections were collected in pre-weighed dishes, weighed for each fish separately, dried and 225 reweighed. The measured weights were used for the calculations described below. The 226 geometric means of stomach and intestinal dry digesta content divided by the fish weight 227 were regressed against time separately for each diet, in order to fit to a model for 228 calculating gastric evacuation rate (GER) and gastric evacuation time (GET) for the 229 stomach, midgut and hindgut filling time (MFT, HFT) and midgut and hindgut evacuation 230 time (MET, HET) for midgut and hindgut.

231

232 2.7 Stomach evacuation pattern calculation

233

In the case of stomach, GER was calculated according to the formula described by Elliott (1972) and adapted by Nikolopoulou et al. (2011) The geometric mean of the stomach dry digesta content divided by the fish weight was regressed against time separately for each diet. GER is estimated as the value of r of the regression model:

238  $W_t = A e^{-rt}$ 

239 which is equivalent to the semi-logarithmic model:

 $lnW_t = lnA-rt$ 

241 where Wt is the geometric mean weight of stomach dry matter digesta at time t, A is an

242 intercept estimated from the model regression and r is the rate of gastric evacuation. GET

243 50%, GET 75% and GET 90% is the evacuation time (expressed in hours) required to

empty 50%, 75% and 90% of the stomach. It was computed through:

245 GET *p*%=[ln100-ln(100-p)]/r

where *p* is the digestible organic matter to be evacuated from the stomach.

247

248 2.8 Midgut and Hindgut Filling/Evacuation pattern calculations

249

In the case of midgut and hindgut the best model is the quadratic regression, represented as a parabola. It was not possible to calculate GER, because the trend of digesta in midgut and hindgut was not linear. Points from each curve were estimated to determine the time of maximum midgut and hindgut filling and evacuation time. The MFT and HFT, (maximum filling time) was calculated as the vertex of the parabola. In these cases, the equations used were:

256  $W_t = A + r_1 t + r_2 t^2$ 

257 where  $r_1$  is the coefficient of the linear part of the model and  $r_2$  is the parameter of the

- 258 quadratic part of the model.
- 259 The vertex of the parabola:

260  $[-(r_1/2r_2), (4Ar_2 - r_1^2)/4r_2]$ 

and MFT and HFT:

262  $-(r_1/2r_2)$ 

263	MET and HET 50%, 75% and 90% is the evacuation time (expressed in hours) required
264	to empty 50%, 75% and 90% of midgut and hindgut and was computed through:
265	$(1-p)^*[(4Ar_2 - r_1^2)/4r_2] = A + r_1t + r_2t^2.$
266	MET and HET 50% , 75% and 90% is the solution with respect to t of the equation:
267	A-(1-p)*[(4Ar <sub>2</sub> - $r_1^2$ )/4 $r_2$ ]+ $r_1t$ + $r_2t^2$ =0
268	by Newton Raphson optimization where p=0.50, 0.75, 0.90.
269	
270	2.9 Digesta moisture determination
271	
272	The weights of stomach and intestine contents, at each sampling time, were used to
273	calculate the moisture of digesta as percentage of digesta weight in each gastrointestinal
274	segment, according to Nikolopoulou et al. (2011).
275	
276	2.10 Analytical methods
277	
278	Diets and whole body were analysed for proximate composition. Moisture content was
279	obtained by weight loss after drying samples in a stove at 105 $^{\circ}$ C until a constant weight
280	was achieved. Crude protein was determined as total nitrogen (N) by using the Kjeldahl
281	method and multiplying N by 6.25. Total lipids were determined according to Bligh and
282	Dyer's (1959) extraction method. Ash content was estimated by incineration to a constant
283	weight in a muffle oven at 450 °C. NDF, acid-detergent fibre (ADF) and lignin were
284	determined according to Van Soest and Robertson, (1981). Cellulose content was

286 NDF-ADF content. Gross energy was determined by a calorimetric bomb (Adiabatic

285

determined based on the ADF-lignin. Hemicellulose content was determined based on

Calorimetric Bomb Parr 1261; PARR Instrument, IL, U.S.A.). Serum glucose, alkaline
phosphatase, inorganic phosphorus, triglycerides and total protein were measured using
colorimetric methods (Total protein OSR6232, Triglyceride OSR61118, Glucose
OSR6121, Alkaline phosphatase OSR6004, Inorganic phosphorus OSR6122; Beckman
Coulter, Brea, CA, USA) on an automated analyser (AU 400, Beckman Coulter, Brea
CA, USA).

293

294 2.11 Statistical analysis

295

296 All data are presented as mean ± standard deviation (SD). Tank was used as the 297 experimental unit for analyzing growth and performance, a pool of five sampled fish was 298 considered the experimental unit for analyzing carcass composition, whereas individual 299 fish was used for analyzing VSI, HSI, MFI, and blood parameters. Data of growth 300 performance, VSI, HSI, MFI, nutritional indices and blood parameters were analysed by 301 a one-way ANOVA and in case of significance ( $P \le 0.05$ ) Tukey's post hoc test was 302 performed. The normality and/or homogeneity of variance assumptions were validated 303 for all data preceding ANOVA. Moisture of digesta was analysed by two-way ANOVA 304 using diet and time as independent factors. Statistical analysis was performed using the 305 software R version 3.1.0 (Revolution analytics, Palo Alto, CA, USA). The differences 306 among treatments were considered significant at  $P \leq 0.05$ .

307

308 **3. Results** 



311

312 Growth performances are summarised in Table 2. No significant differences due to 313 fibre inclusion levels were observed after the 117 days in terms of growth performance 314 (final body weight and SGR), feed intake (VFI) and feed utilisation (FCR). Data on 315 biometric indices, body composition and nutritional indices are shown in Table 3. VSI 316 and HSI values significantly decreased with increasing fibre dietary levels, while no 317 significant differences were found in MFI. Regarding whole body protein content, fish 318 fed diet F7.2 showed higher value in comparison to those fed F13.1, while no significant 319 differences were found among the lipid, ash and moisture content of fish. No significant 320 differences among treatments were found in PER, GPE and GLE. Serum TP, GLU, TRIG, 321 ALP and P levels are shown in Table 4. No significant differences among treatments were 322 found. Data on P were significant for ANOVA (P = 0.0469) but no differences among 323 treatments were detected by the multiple comparison test.

324

325 *3.2 Histology* 

326

All the histological sections showed normal intestinal architecture. No differences were found in the mucosal layer (goblet cells, supranuclear absorption vacuoles in the enterocytes), capillarity network within the intestinal folds, lymphoplasmacytic cells within the *lamina propria*. Moreover, inflammatory and/or degenerative changes were not present in the histological sections taken from the examined subjects (Fig. 1).

332

333 *3.3 Gut evacuation rate and time* 

335 Data on GER, GET, MFT, HFT, MET and HET are presented in Table 5. Dry weight 336 of the stomach content decreased with time as shown in Fig. 2. No significant differences 337 were found among treatments for stomach, midgut and hindgut however diet F7.2 338 required 22 h to evacuate 90% of the initial digesta content, while diet F13.1 and F15.5 339 required 38 h and 35 h respectively. The data extracted for the midgut and for the hindgut 340 evacuation fitted a quadratic model (Fig. 3 a, b). According to the equations, the time 341 required to empty the whole gastrointestinal tract was estimated to be around 46-47 h for 342 fish fed under all treatments. Data on moisture of digesta for each gastrointestinal tracts 343 at 1, 4, 10, 16, 28, and 48 h post prandial are shown in Table 6. No differences were found 344 between treatments, but moisture in the stomach changed with time increasing from 1 to 345 4 h postprandial and then remained constant until 28 h postprandial.

346

#### 347 **4. Discussion**

348

349 To our knowledge, this is the first study designed to specifically evaluate the effects of 350 graded levels of dietary insoluble fibre derived from sunflower hulls and soybean hulls 351 on performance, health and digesta transit time of European sea bass using formulations 352 at practical level. Crude fibre which is basically cellulose, insoluble hemicellulose and 353 lignin, does not have any value in the nutrition of carnivorous fish and according to Altan 354 and Korkut (2011) should be restricted to less than 7 % in fish diet. The results of our 355 study demonstrated that neutral detergent fibre (hemicellulose, cellulose and lignin) up to 356 level of 15.5 % had no effects on growth performance, feed intake and protein/lipid 357 utilisation. The performances registered in this trial are similar to those found in other 358 studies on European sea bass (Tulli et al., 2010; Guerreiro et al., 2015; Bonvini et al.,

359 2018). Hansen and Hemre (2013) reported that plant protein can contain soluble fibres 360 and antinutrients which can interfere with nutrient digestibility. Soluble fibres increase 361 the viscosity of gut content, which potentially can reduce digestible enzyme activities and 362 negatively affect nutrient digestion and absorption (Leenhouwers et al., 2006). Recently, Adamidou et al. (2011) reported that an inclusion of fibre up to a level of 5 g 100 g<sup>-1</sup> did 363 364 not affect growth performance or nutrient digestibility in sharpsnout seabream (Diplodus 365 *puntazzo*). Bou et al. (2014) have demonstrated that using diets with the inclusion of fibre 366 from cellulose up to 18 % at the expense of carbohydrate did not affect growth 367 performance in gilthead sea bream (Sparus aurata). Both sharpsnout and gilthead sea 368 bream belong to the family of Sparidae, which are considered omnivorous fish and more 369 predisposed to digest plant-based ingredients then carnivorous species, such as European 370 sea bass (Stone, 2003). In general, dietary fibre did not affect whole body composition 371 except for the protein content which was slightly lower in F13.1 compared to F7.2 but 372 without resulting in lower protein efficiency.

VSI and HSI values increased in diets with the lowest fibre inclusion. No evidence is reported in the literature about any influence of fibre levels on somatic indices. One possible explanation of the trend in HSI, found in our study, can be related to the decrease in level of wheat in diets at the increasing of fibre. Previous studies on European sea bass and other species reported that increasing NFE, wheat and starch level in the diets can increase HSI values probably due to an enhancement in glycogen liver deposition (Bonaldo et al., 2008, 2010; Bonvini et al., 2017; Bou et al., 2014).

Information on the nutritional status and health in fish species can be achieved through the study of blood metabolites (Bonaldo et al., 2015; Bonvini et al., 2015). To our knowledge, no studies have assessed blood parameters in response to fibre inclusion in

383 diets in marine species. Among the blood parameters, TP, TRIG, GLU, ALP and P seem 384 to have potential as predicative diagnostic tools for evaluation of European sea bass 385 nutritional status (Peres et al., 2014). In this trial, no differences were found among 386 treatments in all parameters analysed. TP level is usually very stable in well-nourished 387 animals but decreases under fasting conditions (Peres et al., 2014). In this study, plasma 388 TP averaged between 5.6 and 5.1 g dl<sup>-1</sup> and these values are within the ranges reported 389 for European sea bass under good nutritional status (Peres et al., 2014). TRIG levels are 390 in agreement with previously reported values for this species (Bonvini et al., 2017), but 391 higher than those reported by Peres et al. (2014), a difference which can be related to the 392 different lipid level and formulation of the diet used, in comparison to our study. GLU 393 levels are in agreement with previously reported values for this species (Peres et al., 394 2014). Adamidou et al. (2009), reported that GLU levels in European sea bass serum were 395 also affected by the type of carbohydrate ingested, with wheat starch showing the most 396 rapid increase and decrease in serum glucose compared to fish fed pea and chickpea diets, 397 while faba bean starch resulted in a delay in serum glucose peak and a lower range of 398 glucose values. ALP levels are higher in comparison to Peres et al. (2014). Alkaline 399 phosphatase is involved in the absorption and transport of lipid and carbohydrates from 400 the intestine, and intestinal activities are positively correlated with food ingestion and 401 growth rate (Lemieux et al., 1999). Finally, inorganic phosphorus levels are in agreement 402 with previously reported values for this species (Peres et al., 2014). Plasma electrolytes 403 (univalent and bivalent) are considered to be valuable indicators of secondary stress and 404 osmoregulation ability in fish (Roque et al., 2010). Plasma phosphorus has been identified 405 as a good indicator of stress (i.e. starvation and stocking density) and pathological 406 situations (Roque et al., 2010; Peres et al., 2014).

407 Besides the study of blood metabolites, histological analysis of the digestive system is 408 considered a valid tool for evaluating the gut health status. No inflammatory and/or 409 degenerative changes were recorded in any of the histological sections of the gut. The 410 examination revealed no alteration in the mucosal layer (goblet cells and supranuclear 411 absorption vacuoles in the enterocytes), no vascular changes within the intestinal folds 412 and no lymphoplasmacytic infiltration within the lamina propria. Few references are 413 available in literature on the effects of fibre on fish intestine. For example, Olsen et al. 414 (2007) reported in Atlantic cod an increase in cellularity of the lamina propria and a 415 modest goblet cell hypertrophy and hyperplasia especially in fish fed a diet with 100% of 416 plant protein mixture. These results were ascribed to a significant amount of fibres in 417 plant-based diets. In our study these morphological changes were not observed in any of 418 the diet treatments confirming a good tolerance of European sea bass intestine to a fibre 419 content up to 15.5%. Leigh et al. (2017) reported in zebrafish (Danio rerio) changes in 420 the epithelial surface area and overall gut length as a response to fibre intake level but 421 without any inflammatory or degenerative effects.

422 Dietary fibre can affect gastrointestinal transit time of feed (Zhou et al., 2004). The 423 investigation on gastrointestinal evacuation rate/time in this study reported no significant 424 differences between treatments, however higher fibre dietary levels led to an increased 425 time required to empty the stomach. A marked difference in term of mean values was 426 infact observed between diets F7.2, F13.1, F15.5 which required 22, 38 and 35 h, 427 respectively to empty 90% of the stomach content. In addition slower evacuation was also 428 observed in F13.1 and F15.5 compared to F7.2 to empty 50% and 75% of the stomach 429 content. In the present trial if we consider that the inclusion of high dietary fibre level did 430 not exert a negative effect on performance and feed utilization, a possible explanation 431 may be related to a reduced gastric evacuation at high fibre level which might have432 improved feed digestion and absorption.

433 On the other hand, the evacuation time for the hindgut was very similar in all the 434 treatments, around 46-47 h to empty 90% of hindgut content. There is little published 435 information regarding the effects of fibre on gastric evacuation in marine carnivorous 436 species. When plant-based ingredients are included in the diet, gastric evacuation time is 437 higher, increasing more with legumes than with cereals (Adamidou et al., 2009). 438 Moreover, Venou et al. (2003) reported that differences in ingredient processing can also 439 modify gastric evacuation time, which is higher for extruded cereals than for raw cereals 440 in gilthead sea bream. In the same species, García-Meilán et al. (2014) found that 441 differences in diet composition, such as high lipid levels or high starch content, may be 442 involved in the differential transit rate and Fountoulaki et al. (2005) found that a low 443 transit rate was related to a high lipid content. Since in the present work diets were 444 designed at a practical level in order to identify the maximum amount of dietary fibre 445 inclusion without negatively affecting zootechnical performance and health, we cannot 446 totally exclude the effects that the differences in the lipid content could have exerted on 447 the gastric evacuation pattern. Moisture of digesta in the gastrointestinal tract showed no 448 differences between treatments. Moisture of stomach increased with time from 1 to 4 h 449 after feeding and remained unchanged until 28 h post prandial. The water required for 450 feed moisturisation originates from feed water, initial water absorption of pellets, 451 drinking and stomach secretions (Kristiansen and Rankin, 2001). In this context, dietary 452 fibre behaves within the gastrointestinal tract as a polymer matrix with variable 453 physicochemical properties including water-holding capacity (Kay, 1982).

454

455 **5. Conclusion** 

456

457 In conclusion, the different fibre levels derived from sunflower hulls and soybean hulls 458 tested in this trial had no effects on overall performances and feed efficiency in European 459 sea bass. Results from blood biochemistry profile and gut histology confirm a good 460 nutritional and health status of fish under all feeding treatments. The inclusion of fibres 461 derived from sunflower and soybean hulls had no influence also on digesta transit time 462 and digesta moisture even if higher dietary fibre levels seem to increase the time required 463 to empty the stomach. According to the results it seems feasible to include insoluble fibre 464 up to 15.5% in feed formulation for the on-growing of European sea bass without 465 negatively affecting zootechnical performance and health.

466

#### 467 Acknowledgments

The authors would like to thank Paola Parazza and Marina Silvi for technical assistance during laboratory analyses. The author would like to thank Gillian Forlivesi Heywood for English language editing. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The authors declare no conflict of interest.

473

#### 474 **References**

475

Adamidou, S., Nengas, I., Alexis, M., Fountoulaki, E., Nikolopoulou, D., Campbell, P.,
Karacostas, I., Rigos, G., Bell, G.J., Jauncey, K., 2009. Apparent nutrient
digestibility and gastrointestinal evacuation time in European seabass

- 479 (*Dicentrarchus labrax*) fed diets containing different levels of legumes.
  480 Aquaculture 289, 106-112.
- Adamidou, S., Rigos, G., Mente, E., Nengas, I., Foundoulaki, E., 2011. The effects of
  dietary lipid and fibre levels on digestibility of diet and on the growth performance
  of sharpsnout seabream (*Diplodus puntazzo*). Mediterr. Mar. Sci. 12, 401-412.
- Altan, O., Korkut, A.Y., 2011. Appearent Digestibility of Plant Protein Based Diets by
  European Sea Bass *Dicentrarchus labrax* L., 1758. Turk. J. Fish. Aquat. Sci. 11,
  87-92.
- 487 Amirkolaie, A.K., Leenhouwers, J.I., Verreth, J.A.J., Schrama, J.W., 2005. Type of
  488 dietary fibre (soluble versus insoluble) influences digestion, faeces characteristics
  489 and faecal waste production in Nile tilapia (*Oreochromis niloticus* L.). Aquacult.
  490 Res., 36, 1157–1166.
- 491 Bach Knudsen, K.E., 2001. The nutritional significance of "dietary fibre" analysis. Anim.
  492 Feed Sci. Technol. 90, 3–20.
- Bligh, E.G., Dyer, W.J., 1959. A rapid method of total lipid extraction and purification.
  Can. J. Biochem. Phys. 37, 911–917.
- Bonaldo, A., Roem, A.J., Fagioli, P., Pecchini, A., Cipollini, I., Gatta, P.P., 2008.
  Influence of dietary levels of soybean meal on the performance and gut histology
  of gilthead sea bream (*Sparus aurata* L.) and European sea bass (*Dicentrarchus labrax* L.). Aquacult. Res. 39, 970-978.
- 499 Bonaldo, A., Isani, G., Fontanillas, R., Parma, L., Grilli, E., Gatta, P.P., 2010. Growth
- 500 and feed utilization of gilthead sea bream (Sparus aurata, L.) fed to satiation and
- 501 restrictively at increasing dietary energy levels. Aquacult. Int. 18, 909-919.

502	Bonaldo, A., Di Marco, P., Petocchi, T., Marino, G., Parma, L., Fontanillas, R., Koppe,
503	W., Mongile, F., Finoia, M.G., Gatta, P.P., 2015. Feeding turbot juveniles Psetta
504	maxima L. with increasing dietary plant protein levels affects growth performance
505	and fish welfare. Aquacult. Nutr. 21, 401-413.
506	Bonvini, E., Parma, L., Mandrioli, L., Sirri, R., Brachelente, C., Mongile, F., Gatta, P.P.,
507	Bonaldo, A., 2015. Feeding common sole (Solea solea) juveniles with increasing
508	dietary lipid levels affects growth, feed utilization and gut health. Aquaculture
509	449, 87-93.
510	Bonvini, E., Bonaldo, A., Mandrioli, L., Sirri, R., Dondi, F., Bianco, C., Fontanillas, R.,
511	Mongile, F., Gatta, P.P., Parma, L., 2017. Effects of feeding low fishmeal diets
512	with increasing soybean meal levels on growth, gut histology and plasma
513	biochemistry of sea bass. Animal 1-8. doi:10.1017/S1751731117002683.
514	Bonvini, E., Parma, L., Badiani, A., Fontanillas, R., Gatta, P.P., Sirri, F., Nannoni, E.,
515	Bonaldo, A., 2018. Integrated study on production performance and quality traits
516	of European sea bass (Dicentrarchus labrax) fed high plant protein diets,
517	Aquaculture, 484, 126-132.
518	Bou, M., Todorčević, M., Fontanillas, R., Capilla, E., Gutiérrez, J., Navarro, I., 2014.
519	Adipose tissue and liver metabolic responses to different levels of dietary
520	carbohydrates in gilthead sea bream (Sparus aurata). Comp. Biochem. Physiol. A
521	175, 72–81.
522	Dalsgaard, J., Bach Knudsen, K.E., Verlhac, V., Ekmann, K.S., Pedersen, P.B., 2016.
523	Supplementing enzymes to extruded, soybean-based diet improves breakdown of
524	non-starch polysaccharides in rainbow trout (Oncorhynchus mykiss). Aquac. Nutr.
525	22, 419–426.

- 526 Elliott, J.M., 1972. Rates of gastric evacuation in brown trout, *Salmo trutta* L. Freshw.
  527 Biol. 2, 1-18.
- Eusebio, P.S., Coloso, R.M., Mamauag, R.E.P., 2004. Apparent digestibility of selected
  ingredients in diets for juvenile grouper, *Epinephelus coiodes* (Hamilton).
  Aquacult. Res. 35, 1261-1269.
- Fountoulaki, E., Alexis, M.N., Nengas, I., Venou, B., 2005. Effect of diet composition on
  nutrient digestibility and digestive enzyme levels of gilthead sea bream (*Sparus aurata* L.). Aquacult. Res. 26, 1243–1251.
- García-Meilán, I., Ordóñez-Grande, B., Gallardo, M.A., 2014. Meal timing affects
  protein-sparing effect by carbohydrates in sea bream: Effects on digestive and
  absorptive processes. Aquaculture 434, 121-128.
- Glencross, B., 2009. The influence of soluble and insoluble lupin non-starch
  polysaccharides on the digestibility of diets fed to rainbow trout (*Oncorhynchus mykiss*). Aquaculture, 294, 256–261.
- Glencross, B., Rutherford, N., Bourne, N., 2012. The influence of various starch and nonstarch polysaccharides on the digestibility of dies fed to rainbow trout
  (*Oncorhynchus mykiss*). Aquaculture, 356–357, 141–146.
- Guerreiro, I., Oliva-Teles, A., Enes, P., 2015. Improved glucose and lipid metabolism in
  European sea bass (*Dicentrarchus labrax*) fed short-chain fructooligosaccharides
  and xylooligosaccharides. Aquaculture 441, 57–63.
- 546 Hansen, J.O., Storebakken, T., 2007. Effects of dietary cellulose level on pellet quality
- and nutrient digestibilities in rainbow trout (Oncorhynchus mykiss). Aquaculture,
  272, 458–465.

- Hansen, A.C., Hemre, G.I., 2013. Effect of replacing fish meal and oil with plant
  resources in on-growing diets for Atlantic cod (*Gadus morhua* L.). Aquacult.
  Nutr. 19, 641-650.
- 552 Kay, R.M., 1982. Dietary fiber. J. Lipid Res. 23, 221-242.
- Kousoulaki, K., Sæther, B.S., Albrektsen, S., Noble, C., 2015. Review on European sea
  bass (*Dicentrarchus labrax*, Linnaeus, 1758) nutrition and feed management: a
  practical guide for optimizing feed formulation and farming protocols. Aquacult.
  Nutr. 21, 129-151.
- Kraugerud, O.F., Penn, M., Storebakken, T., Refstie, S., Krogdahl, A., Svihus, B., 2007.
  Nutrient digestibilities and gut function in Atlantic salmon (*Salmo salar*) fed diets
- 559 with cellulose or non-starch polysaccharides from soy. Aquaculture, 273, 96–107.
- Kristiansen, H.R., Rankin, J.C., 2001. Discrimination between endogenous and
  exogenous water sources in juvenile rainbow trout fed extruded dry feed. Aquat.
  Living Resour. 14, 359–366.
- Krogdahl, A., Hemre, G.I., Mommsen, T.P., 2005. Carbohydrates in fish nutrition;
  digestion and absorption in postlarval stages. Aquacult. Nutr. 11, 103-122.
- 565 Leenhouwers, K.I., Adjei-Boateng, D., Verreth, J.A.J., Schrama, J.W., 2006. Digesta
- viscosity, nutrient digestibility and organ weights in African catfish (*Clarias gariepinus*) fed diets supplemented with different levels of a soluble non-starch
  polysaccharide. Aquacult. Nutr. 12, 111–116.
- 569 Leigh, S.C., Nguyen-Phuc, B.Q., German, D.P., 2017. The effects of protein and fiber
- 570 content on gut structure and function in zebrafish (*Danio rerio*). J Comp Physiol
- 571 B. doi: 10.1007/s00360-017-1122-5.

572	Lekva, A., Hansen, A.C., Rosenlund, G., Karlsen, O., Hemre, G.I., 2010. Energy dilution
573	with alpha-cellulose in diets for Atlantic cod (Gadus morhua L.) juveniles -
574	effects on growth, feed intake, liver size and digestibility of nutrients.
575	Aquaculture, 300, 169–175.

- 576 Lemieux, H., Blier, P., Dutil, J.D., 1999. Do digestive enzymes set a physiological limit
  577 on growth rate and food conversion efficiency in the Atlantic cod (*Gadus*578 *morhua*)? Fish Physiol. Biochem. 20, 293-303.
- Mongile, F., Bonaldo, A., Fontanillas, R., Mariani, L., Badiani, A., Bonvini, E., Parma,
  L., 2014. Effects of dietary lipid level on growth and feed utilisation of gilthead
  seabream (*Sparus aurata* L.) reared at Mediterranean summer temperature. Ital. J.
  Anim. Sci. 13, 30-35.
- Nikolopoulou, D., Moutou, K.A., Fountoulaki, E., Venou, B., Adamidou, S., Alexis,
  M.N., 2011. Patterns of gastric evacuation, digesta characteristics and pH changes
  along the gastrointestinal tract of gilthead sea bream (*Sparus aurata* L.) and
  European sea bass (*Dicentrarchus labrax* L.). Comp. Biochem. Physiol. A 158,
  406-414.
- Nutrient requirements of fish and shrimp (NRC) 2011. National Academic Press,
  Washington DC, USA.
- Olsen, R.E., Hansen, A.C., Rosenlund, G., Hemre, G.I., Mayhew, T.W., Knudsen, D.L.,
  Eroldogan, O.T., Myklebust, R., Karlsen, Ø., 2007. Total replacement of fish meal
  with plant proteins in diets for Atlantic cod (*Gadus morhua* L.) II: Health aspects.
  Aquaculture 272, 612–624.
- Parma, L., Candela, M., Soverini, M., Turroni, S., Consolandi, C., Brigidi, P., Mandrioli,
  L., Sirri, R., Fontanillas, R., Gatta, P.P., Bonaldo, A., 2016. Next-generation

- sequencing characterization of the gut bacterial community of gilthead sea bream
  (*Sparus aurata*, L.) fed low fishmeal based diets with increasing soybean meal
  levels. Anim. Feed Sci. Technol. 222, 204-216.
- Peres, H., Santos, S., Oliva-Teles, A., 2014. Blood chemistry profile as indicator of
  nutritional status in European seabass (*Dicentrarchus labrax*). Fish Physiol.
  Biochem. 40, 1339-1347.
- Roque, A., Yildiz, H.Y., Carazo, I., Duncan, N., 2010. Physiological stress responses of
  seabass (*Dicentrarchus labrax*) to hydrogen peroxide (H2O2) exposure.
  Aquaculture 304, 104-107.
- 605 Stone, D.A.J., 2003. Dietary carbohydrate utilization by fish. Rev. Fish. Sci. 11, 337-369.

Tulli, F., Messina, M., Calligaris, M., Tibaldi, E., 2010. Response of European sea bass

- 607 (*Dicentrarchus labrax*) to graded levels of methionine (total sulfur amino acids)
  608 in soya protein-based semi-purified diets. Br. J. Nutr. 104, 664-673.
- Van Soest, P.J., Robertson, J.B., 1981. The detergent system of analysis and its
  application to human foods. Chapter 8. In: James, W.P.T., Theander, O. (Eds.),
  The Analysis of Dietary Fibre in Food. Marcel Dekker Inc., New York, p. 123.
- 612 Venou, B., Alexis, M.N., Fountoulaki, E., Nemges, I., Apostolopoulou, M. Castritsi-
- 613 Cathariou, I., 2003. Effect of extrusion of wheat and corn on gilthead sea bream
- 614 (*Sparus aurata*) growth, nutrient utilization efficiency, rates of gastric evacuation
- and digestive enzyme activities. Aquaculture 225, 207-223.

- 616 Zhou, Q.C., Tan, B.P., Mai, K.S., Liu, Y.J., 2004. Apparent digestibility of selected feed
- 617 ingredients for juvenile cobia *Rachycentron canadum*. Aquaculture 241, 441-451.

	F7.2	F8.9	F11.5	F13.1	F15.5
Ingredient, % of the diet					
Sunflower Hulls	0.0	1.5	3.2	4.8	6.4
Soybean Hulls	0.0	2	4.2	6.4	8.6
Fish meal	20	20	20	20	20
Soybean concentrate	13	13	13	13	13
Wheat	28.4	23.3	17.9	12.4	6.9
Corn gluten	8	8	8	8	8
Wheat gluten	15.2	15.6	16.1	16.6	17
Fish oil	7.5	8	8.6	9.2	9.8
Rapeseed oil	7.5	8	8.6	9.2	9.8
Vit/Min premix <sup>1</sup>	0.5	0.5	0.5	0.5	0.5
Proximate composition, % on	a wet weigh	nt basis			
Moisture	7.28	7.77	7.71	7.49	7.74
Protein	43.6	43.2	43.5	43.8	43.7
Lipid	22.5	24.3	24.8	25.4	27.1
Ash	4.42	4.47	4.53	4.63	4.73
$NDF^2$	7.20	8.93	11.46	13.14	15.46
$ADF^3$	2.79	4.45	5.99	7.11	9.26
Hemicellulose	4.41	4.48	5.48	6.03	6.20
Cellulose	2.33	3.37	4.73	5.73	7.07
Lignin	0.46	1.08	1.26	1.38	2.20
Gross energy MJ kg <sup>-1</sup>	21.5	22.1	22.1	22.5	23.0

**Table 1.** Ingredients and proximate composition of the experimental diets

<sup>1</sup> Vitamin and mineral premix; Skretting, Stavanger, Norway (fulfilling recommendations for marine fish species given by NRC, 2011) <sup>2</sup>Neutral detergent fibre <sup>3</sup>Acid detergent fibre

Table 2. Growth performance and feed intake of European sea bass fed experimental diets over 117 days.

uuys.						
	Experimental	l diets				
	F7.2	F8.9	F11.5	F13.1	F15.5	P value
IBW (g)	$69.5 \pm 2.1$	$68.1\pm4.0$	$68.7 \pm 1.1$	$69.7 \pm 1.3$	$71.2\pm2.5$	0.5944
FBW (g)	$215\pm11.7$	$204 \pm 14.9$	$215\pm15.9$	$223\pm6.3$	$219 \pm 15.8$	0.5369
SGR (% day <sup>-1</sup> )	$0.97\pm0.03$	$0.94\pm0.02$	$0.97\pm0.07$	$0.99\pm0.01$	$0.96\pm0.04$	0.5136
VFI (g feed fish <sup>-1</sup> )	$237\pm7.7$	$219\pm7.7$	$230 \pm 11.6$	$232\pm8.1$	$237 \pm 13.4$	0.2759
FCR	$1.63\pm0.06$	$1.43\pm0.24$	$1.58\pm0.11$	$1.51\pm0.03$	$1.60\pm0.06$	0.3773
Survival %	$100\ \pm 0.0$	$100\pm0.0$	$99.4 \pm 1.0$	$100\pm0.0$	$100\pm0.0$	0.4516

Data are given as the mean  $(n=3) \pm SD$ . In each line, different superscript letters indicate significant differences among treatments (*P*  $\leq$  0.05).

IBW = Initial body weight. FBW = Final body weight.

SGR = Specific growth rate (% day<sup>-1</sup>) =  $100 * (\ln FBW- \ln IBW) / days$ . VFI = Voluntary Feed Intake (g feed/fish) = g feed ingested / number of fish FCR = Feed conversion rate = feed intake / weight gain.

625

626

the experimental diets									
	Experimental diets								
	F7.2	F8.9	F11.5	F13.1	F15.5	P value			
Biometric ind	dices								
VSI	$12.1^{\text{b}}\pm1.5$	$11.4^{ab}\pm1.8$	$11.3^{ab}\pm2.5$	$11.1^{ab} \pm 1.4$	$10.5^{\rm a}\pm1.9$	0.0303			
HSI	$3.2^{\rm c}\pm0.5$	$3.1^{\rm c}\pm0.5$	$2.7^{b}\pm0.6$	$2.6^{ab}\pm0.5$	$2.3^{a}\pm0.5$	< 0.0001			
MFI	$6.7\pm1.5$	$6.1 \pm 1.8$	$6.4\pm2.5$	$6.3\pm1.5$	$5.8 \pm 1.6$	0.4604			
Whole body	composition, 9	6							
Protein	$17.1^{\text{b}}\pm0.1$	$16.4^{ab}\pm0.3$	$16.6^{ab}\pm0.3$	$16.2^{\rm a}\pm0.5$	$16.3^{ab}\pm0.3$	0.0465			
Lipid	$17.6\pm1.3$	$18.6 \pm 1.4$	$17.6\pm2.3$	$17.8 \pm 1.9$	$19.0\pm1.1$	0.7498			
Ash	$2.8\pm0.21$	$2.7\pm0.03$	$2.7\pm0.07$	$2.7\pm0.04$	$2.8\pm0.15$	0.7554			
Moisture	$61.6\pm0.8$	$61.2\pm0.4$	$60.9\pm0.5$	$60.3\pm0.5$	$60.0\pm1.9$	0.3485			
Nutritional ir	ndices								
PER	$1.41\pm0.05$	$1.43\pm0.08$	$1.46\pm0.11$	$1.51\pm0.03$	$1.43\pm0.05$	0.4484			
GPE	$24.4\pm0.97$	$23.5 \pm 1.78$	$24.2 \pm 1.24$	$24.2\pm0.62$	$23.2\pm0.65$	0.6601			
GLE	$56.4\pm7.1$	$57.2\pm8.2$	$52.9 \pm 11.7$	$54.2\pm6.3$	$52.7\pm4.1$	0.9343			

Table 3. Biometric indices	, body composition	and nutritional	indices of	European s	ea bass fed
the experimental diets				-	

Data are given as the mean (n=3; n=30 for VSI, HSI, MFI)  $\pm$  SD. In each line, different superscript letters indicate significant differences among treatments ( $P \le 0.05$ ). VSI = Viscerosomatic index (%) = 100\*(viscera weight/FBW). HSI = Hepatosomatic index (%) = 100\*(liver weight/FBW).

MFI = Mesenteric fat index (%) = 100\*(mesenteric fat weight/FBW).PER = Protein efficiency ratio = ((FBW-IBW)/protein intake).

GPE = Gross protein efficiency = 100\*[(%final body protein\*FBW) - (%initial body protein\*IBW)]/total protein intake fish.

GLE = Gross lipid efficiency = 100\*[(%final body lipid\*FBW) - (%initial body lipid\*IBW)]/total lipid intake fish.

	Experiment	al diets				
	F7.2	F8.9	F11.5	F13.1	F15.5	P value
TP (g dL <sup>-1</sup> )	$5.5\pm0.40$	$5.6\pm0.98$	$5.3\pm0.70$	$5.5\pm0.63$	$5.1\pm1.16$	0.5586
TRIG (mg dL <sup>-1</sup> )	$1884\pm818$	$2023\pm600$	$1769\pm788$	$1790\pm715$	$1797\pm670$	0.9064
GLU (mg dL <sup>-1</sup> )	$98\pm55.1$	$90\pm30.1$	$77 \pm 17.3$	$89\pm26.0$	$111\pm33.8$	0.2544
ALP (U L <sup>-1</sup> )	$235\pm58.9$	$252\pm59.1$	$227\pm56.7$	$207\pm49.7$	$210\pm51.9$	0.2300
$P (mg dL^{-1})$	$8.4\pm0.85$	$9.4 \pm 1.21$	$8.5\pm0.72$	$8.4 \pm 1.01$	$8.9 \pm 1.00$	0.0469

Table 4. Blood biochemistry of European sea bass fed the experimental diets

Each value is mean from 12 samples  $\pm$  SD. Data on P were significant for ANOVA (P = 0.0469) but no differences among treatments were detected by the multiple comparison test.

TP = total protein TRIG = triglycerides

GLU = glucose

ALP = alkaline phosphatase

P = inorganic phosphorus



	Experimental diets							
	F7.2	F8.9	F11.5	F13.1	F15.5			
Stomach								
GER	0.1031	0.0945	0.0899	0.0607	0.0657			
GET 50% (h)	6.72	7.33	7.71	11.43	10.55			
GET 75% (h)	13.44	14.66	15.41	22.86	21.09			
GET 90% (h)	22.32	24.36	25.60	37.96	35.04			
$\mathbb{R}^2$	0.82	0.92	0.92	0.98	0.77			
Midgut								
MFT (h)	12.99	21.01	12.52	19.45	19.99			
MET 50% (h)	35.63	39.50	35.39	38.65	38.88			
MET 75% (h)	40.72	43.65	35.39	42.97	43.12			
MET 90% (h)	43.36	45.81	43.20	45.21	45.33			
$\mathbb{R}^2$	0.33	0.55	0.33	0.43	0.53			
Hindgut								
HFT (h)	25.71	23.28	23.71	25.66	25.34			
HET 50% (h)	41.60	41.00	40.07	41.54	41.20			
HET 75% (h)	45.71	44.99	43.74	45.11	44.76			
HET 90% (h)	47.03	47.06	45.66	46.97	46.62			
$\mathbb{R}^2$	0.71	0.42	0.65	0.70	0.60			

**Table 5.** Gastric evacuation rate (GER), gastric evacuation time (GET, expressed in hours), midgut filling and evacuation time (MFT, MET, h) and hindgut filling and evacuation time (HFT, HET, h) of European sea bass fed the experimental diets

No significant differences among treatments and within each gastrointestinal tract were detected  $P \le 0.05$ 

140) allel	leeding								
Stomach	T1	T4	T10	T16	T28	T48	Time	Diet	Time*Diet
F7.2	$63.0\pm2.5^{\rm a}$	73.9 ±4.3 <sup>ab</sup>	$77.9 \pm 5.0^{b}$	$82.9 \pm 2.6^{b}$	77.1±1.2 <sup>b</sup>				
F8.9	$58.9 \pm 11.5^{a}$	72.9±3.4 <sup>b</sup>	$77.2 \pm 4.4^{b}$	$80.4{\pm}1.4^{b}$	$80.0 \pm 4.9^{b}$				
F11.5	$60.4 \pm 6.8^{a}$	69.3±9.3 <sup>b</sup>	$79.8 {\pm} 5.0^{\rm b}$	$80.5 \pm 2.9^{b}$	$80.8 \pm 6.1^{b}$		0.0001	0.43	0.115
F13.1	$59.4 \pm 4.2^{a}$	$72.3 \pm 3.0^{b}$	$75.9 \pm 4.1^{b}$	$77.1 \pm 5.1^{b}$	$78.8{\pm}8.8^{\mathrm{b}}$				
F15.5	59.6±3.0 <sup>a</sup>	$74.6 \pm 5.2^{b}$	$80.5 \pm 2.2^{b}$	$78.4 \pm 2.2^{b}$	$75.1 \pm 6.3^{b}$				
Midgut									
F7.2		86.1±2.4	83.0±7.1	$85.4{\pm}1.6$	87.5±1.5				
F8.9		$87.0 \pm 2.0$	85.3±1.1	$85.9 \pm 2.2$	85.3±2.8				
F11.5		85.4±1.6	$85.4 \pm 2.6$	$83.2 \pm 6.8$	85.5±4.7		0.608	0.647	0.769
F13.1		84.5±5.7	85.5±1.4	$84.2 \pm 3.9$	$84.2 \pm 6.1$				
F15.5		85.1±2.5	85.1±2.0	85.1±3.8	$84.9 \pm 2.6$				
Hindgut									
F7.2		80.2±9.1	$80.4{\pm}10.1$	82.5±2.3	$79.5 \pm 5.9$	$80.4 \pm 6.7$			
F8.9		$80.8 \pm 3.4$	$81.9 \pm 1.9$	$78.8 \pm 10.4$	81.6±5.3	$77.8 \pm 5.1$			
F11.5		77.1±8.7	$80.8 \pm 7.2$	$80.3 \pm 8.6$	$79.0 \pm 7.2$	71.3±7.3	0.140	0.087	0.936
F13.1		$84.0 \pm 5.0$	83.0±2.2	79.7±11.9	82.1±4.5	$80.6 \pm 3.4$			
F15.5		81.2±5.8	83.0±2.4	81.3±6.0	81.6±5.7	75.9±11.1			

**Table 6.** Moisture, % of digesta obtained for each gastrointestinal tract of European sea bass from 1 to 48 hours (T1, T48) after feeding

Values within a row with different superscripts differ significantly at P < 0.05. Statistical significance by two-way ANOVA using time and diet as independent factors.

650 Figure captions

651

652 Figure 1. Histology of the midgut (a, b) and hindgut (c, d) of European sea bass fed diets 653 F7.2 (a, c) and F15.5 (b, d). All the histological sections showed normal architecture of 654 the mucosal layer (regular columnar epithelium with polarized and basally located 655 nuclei), submucosal (loose connective tissue rich in capillary network) and muscular layer 656 (H&E, bar=100 µm). 657 658 Figure 2. Exponential curves showing stomach evacuation of g digesta dry matter % body 659 weight (DM %BW) over the 48 h sampling period of European sea bass fed the 660 experimental diets. No significant differences among treatments were detected  $P \le 0.05$ . 661 662 Figure 3. Midgut (a) and hindgut (b) quadratic curves describing the evacuation of g 663 digesta dry matter % body weight (DM %BW) over the 48 h sampling period of European 664 sea bass fed the experimental diets. No significant differences among treatments were 665 detected  $P \leq 0.05$ .

666

## 668 Figure 1















