



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Feeding European sea bass with increasing dietary fibre levels: Impact on growth, blood biochemistry, gut histology, gut evacuation

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Bonvini, E., Bonaldo, A., Parma, L., Mandrioli, L., Sirri, R., Grandi, M., et al. (2018). Feeding European sea bass with increasing dietary fibre levels: Impact on growth, blood biochemistry, gut histology, gut evacuation. *AQUACULTURE*, 494, 1-9 [10.1016/j.aquaculture.2018.05.017].

Availability:

This version is available at: <https://hdl.handle.net/11585/672940> since: 2021-02-25

Published:

DOI: <http://doi.org/10.1016/j.aquaculture.2018.05.017>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

This is the final peer-reviewed accepted manuscript of:

Feeding European sea bass with increasing dietary fibre levels: impact on growth, blood biochemistry, gut histology, gut evacuation

The final published version is available online at:
<https://doi.org/10.1016/j.aquaculture.2018.05.017>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

25 **Feeding European sea bass with increasing dietary fibre levels: impact on growth,**
26 **blood biochemistry, gut histology, gut evacuation**

27

28 Erika Bonvini^a, Alessio Bonaldo^a, Luca Parma^{a*}, Luciana Mandrioli^a, Rubina Sirri^a,
29 Monica Grandi^a, Ramon Fontanillas^c, Cinzia Viroli^b, Pier Paolo Gatta^a

30

31

32 ^aDepartment of Veterinary Medical Sciences, University of Bologna, Via Tolara di Sopra
33 50, 40064 Ozzano Emilia, Bologna, Italy

34 ^bDepartment of Statistical Sciences “Paolo Fortunati”, University of Bologna, Via delle
35 Belle Arti 41, 40126 Bologna, Italy

36 ^cSkretting Aquaculture Research Centre, Sjøhagen 3, 4016 Stavanger, Norway

37

38

39 **Corresponding author:* Luca Parma, Department of Veterinary Medical Sciences,
40 University of Bologna, Viale Vespucci 2, 47042 Cesenatico, FC, Italy. *Tel.:* +39 0547
41 338931; *Fax:* +39 0547 338941

42 E-mail address: luca.parma@unibo.it (L. Parma)

43

44

45

46

47

48

49 **Abstract**

50

51 Changing trends in fish feed formulation, with progressively higher inclusion levels of
52 plant ingredients, are invariably introducing more fibre despite the fact that this
53 component cannot be utilized by most fish. The effects of increasing insoluble dietary
54 fibre level on growth, nutrient utilization, blood parameters and gut health in European
55 sea bass (*Dicentrarchus labrax* L.) were studied over a period of 117 days. Moreover,
56 investigation on digesta transit time through gastrointestinal evacuation pattern and
57 digesta characteristics (moisture of digesta) were studied. Five iso-proteic diets were
58 formulated to contain increasing insoluble fibre levels, neutral detergent fibre, NDF (7.2,
59 8.9, 11.5, 13.1 and 15.5%) derived by the inclusion of sunflower hulls and soybean hulls.
60 No significant differences due to fibre inclusion levels were observed in final body
61 weight, specific growth rate, feed intake, feed conversion rate, protein and lipid
62 efficiency. No significant differences in serum total protein, glucose, triglycerides,
63 alkaline phosphatase and inorganic phosphorous were found. All the histological sections
64 showed normal intestinal architecture, and inflammatory and/or degenerative changes
65 were not present in any histological section from all subjects examined. The investigation
66 into gastrointestinal evacuation pattern revealed no significant differences between
67 treatments, however higher dietary fibre levels seem to increase the time required to
68 empty the stomach while the time required to empty 90% of the hindgut content was
69 similar in all the treatments: around 46-47 h. No differences were found between diets in
70 the moisture content of digesta along the digestive tracts. We can conclude that the
71 different insoluble fibre levels tested in this trial have no effects on overall performances
72 and feed efficiency in European sea bass. Results from blood biochemistry profile and

73 histology confirm good nutritional and health status of fish under all feeding treatments.
74 The inclusion of fibre had no influence on digesta transit time. In formulation of feed for
75 the on-growing of European sea bass insoluble fibre derived from sunflower hulls and
76 soybean hulls can be included at a level of up to 15.5 %.

77

78 **Keywords:** European sea bass, fibre, growth, blood biochemistry, gut histology, gut
79 evacuation

80

81 **1.Introduction**

82

83 Changing trends in fish feed formulation, with progressively higher inclusion levels of
84 plant ingredients, are invariably introducing more fibre in aquafeeds. Cellulose and other
85 fibrous carbohydrates are found in the structural components of plants and are indigestible
86 to monogastric (simple-stomach) animals, such as fish. Fibre can be divided into soluble
87 and insoluble (according to their extractability in a neutral buffer solution) that may have
88 different antinutritive effects in endothermic animals (Dalsgaard et al., 2016). Soluble
89 fibres in mammals, poultry and some fish species tend to increase digesta viscosity and
90 retard absorption of nutrients (Krogdahl et al., 2005; Amirkolaie et al., 2005), while
91 insoluble fibres appear to act largely as physical bulking agents (Bach Knudsen 2001)
92 and tend to increase digesta transit, resulting in reduced absorption time (Krogdahl et al.,
93 2005). Regarding insoluble dietary fibres, cellulose seems to be a relatively inert dietary
94 component that may reduce the dry matter digestibility in Atlantic salmon (*Salmo salar*),
95 Atlantic cod (*Gadus morhua* L.), Nile tilapia (*Oreochromis niloticus* L.) and rainbow
96 trout (*Oncorhynchus mykiss*) (Amirkolaie et al. 2005; Hansen & Storebakken 2007;

97 Kraugerud et al. 2007; Glencross 2009; Glencross et al. 2012; Lekva et al. 2010).
98 Increasing dietary levels of cellulose have also negatively affected lipid digestibility in
99 cod (Lekva et al. 2010) and energy and protein digestibility in rainbow trout (Glencross
100 2009; Glencross et al. 2012). Moreover, as reported by Altan and Korkut (2011), low
101 dietary concentrations of dietary fibre (3–5 %) may have a beneficial effect on fish
102 growth, but high dietary fibre (>8 %), on the contrary, may decrease dry matter
103 digestibility of the diet and reduce the availability of other nutrients. Some discrepancies
104 between values of maximum dietary fibre level have been reported for fish: less than 7 %
105 (Altan and Korkut, 2011); less than 8 % (Eusebio et al., 2004); as low as possible and not
106 exceeding 10 % (NRC, 2011). Information about the optimal percentage of fibre inclusion
107 in practical feed formulation for the on-growing of the European sea bass (*Dicentrarchus*
108 *labrax* L.) is scarce but Kousoulaki et al. (2015) reported that the fibre content in
109 commercial feeds for this species can range between 1.5-3.2 %. The aim of the present
110 study was to assess the effects of increasing dietary insoluble fibre level derived from
111 sunflower hulls and soybean hulls on growth, nutrient utilisation, blood parameters, gut
112 health, gastrointestinal evacuation pattern and digesta characteristics in European sea bass
113 over 117 days, in order to identify the maximum amount of dietary fibre inclusion without
114 negatively affecting zootechnical performance and health.

115

116 **2. Materials and methods**

117

118 *2.1 Experimental diets*

119 Five iso-proteic diets were formulated to contain increasing insoluble fibre levels
120 (Neutral detergent fibre, NDF 7.2, 8.9, 11.5, 13.1 and 15.5%; F7.2, F8.9, F11.5, F13.1.

121 and F15.5, respectively). Diets were formulated with fishmeal and with a mixture of
122 vegetable ingredients currently used in aquafeed (Parma et al., 2016; Bonvini et al., 2017).
123 The fibre content was increased by increasing levels of a combination of sunflower hulls
124 and soybean hulls to provide same proportion of fibre from each ingredient. Sunflower
125 hulls and soybean hulls were chosen because they are one of the most used and available
126 raw material for feed production. Lipid levels were slightly increased at increasing fibre
127 content in order to compensate for the loss of available energy due to the higher fibre
128 content. The diets were produced by extrusion process by Skretting Aquaculture Research
129 Centre, Stavanger, Norway. The diameter of the pellet was 4 mm. Ingredients and
130 proximate composition of the experimental diets are presented in Table 1.

131

132 *2.2 Fish and feeding trial*

133

134 The experiment was carried out at the Laboratory of Aquaculture, Department of
135 Veterinary Medical Sciences of the University of Bologna, Cesenatico, Italy. European
136 sea bass juveniles were obtained from Maricoltura Mattinatese (Mattinata, Foggia, Italy).
137 At the beginning of the trial, 60 fish (initial average weight: 69.4 ± 2.3 g) per tank were
138 randomly distributed into fifteen 900 L square tanks with a conical base. Each diet was
139 administered to triplicate groups, assigned in a completely random manner, over 117
140 days. Tanks were provided with natural seawater and connected to a closed recirculating
141 system (overall water volume: 18 m^3). The rearing system consisted of a mechanical sand
142 filter (PTK 1200, Astralpool, Barcelona, Spain), ultraviolet lights ($PE 25 \text{ mJ/cm}^2: 32 \text{ m}^3 \text{ h}^{-1}$,
143 ¹, Blaufish, Barcelona, Spain) and a biofilter (PTK 1200, Astralpool, Barcelona, Spain).
144 The water exchange rate within each tank was 100% every hour, while the overall water

145 renewal amount in the system was 5 % daily. During the trial, the temperature was kept
146 constant at 22 ± 1.0 °C and the photoperiod was maintained at 12 h light and 12 h dark
147 through artificial light. The oxygen level was kept constant (8.0 ± 1.0 mg L⁻¹) by a liquid
148 oxygen system regulated by a software programme (B&G Sinergia snc, Chioggia, Italy).
149 Ammonia (total ammonia nitrogen ≤ 0.1 mg L⁻¹), nitrite (≤ 0.2 mg L⁻¹) and salinity (25 g
150 L⁻¹) were daily monitored spectrophotometrically (Spectroquant Nova 60, Merck, Lab
151 business, Darmstadt, Germany). Sodium bicarbonate was added on a daily basis to keep
152 pH constant at 7.8–8.0. Feed was provided to apparent satiation by oversupplying the feed
153 by automatic feeders by approximately 10% of the daily ration, twice a day for six days
154 a week, while one meal was supplied on Sundays, as reported by Mongile et al. (2014).
155 Each meal lasted 1 hour, after which the uneaten pellets of each tank were gathered, dried
156 overnight at 105°C, and their weight was deducted for overall calculation.

157

158 *2.3 Sampling*

159

160 At the beginning and at the end of the experiment, all the fish in each tank were
161 anaesthetised by 2-phenoxyethanol at 300 mg L⁻¹ and individually weighed. Specific
162 growth rate (SGR), voluntary feed intake (VFI) and feed conversion rate (FCR) were
163 calculated. The proximate composition of the carcasses was determined at the beginning
164 of the trial on a pooled sample of 10 fish and on a pooled sample of 5 fish per tank at the
165 end of the trial. Protein efficiency rate (PER), gross protein efficiency (GPE) and gross
166 lipid efficiency (GLE) were calculated. Furthermore, at the end of the trial, wet weight,
167 viscera, liver and mesenteric fat weight were individually recorded for 10 fish per tank to
168 determine viscerosomatic index (VSI), hepatosomatic index (HSI) and mesenteric fat

169 index (MFI). At the end of the trial, five fish per tank (15 fish per dietary treatment) were
170 sampled for intestine histology examination. At the end of the trial, the fish left were kept
171 in the same rearing and feeding conditions for three more days and then were sampled to
172 perform blood analyses of serum total protein (TP), triglycerides (TRIG), glucose (GLU),
173 alkaline phosphatase (ALP) and inorganic phosphorus (P). Blood from 4 fish per tank
174 was collected 5 h postprandial from the caudal vein. Samples were then centrifuged (3000
175 g for 10 min at 4°C), serum aliquots were stored at 4°C and analysed during the same day
176 according to Bonvini et al. (2017). All experimental procedures were evaluated and
177 approved by the Ethical-Scientific Committee for Animal Experimentation of the
178 University of Bologna, in accordance with European directive 2010/63/UE on the
179 protection of animals used for scientific purposes.

180

181 *2.4 Calculations*

182

183 The formulae employed were as follows:

184 Specific growth rate (SGR) (% day⁻¹) = 100 * (ln FBW - ln IBW) / days (where FBW and
185 IBW represent the final and the initial body weights). Voluntary Feed Intake (VFI) (g
186 feed fish⁻¹) = g feed ingested / fish number. Feed conversion ratio (FCR) = feed intake /
187 weight gain. Viscerosomatic index (VSI) (%) = 100 * (viscera weight / body weight).
188 Hepatosomatic index (HSI) (%) = 100 * (liver weight / body weight). Mesenteric fat index
189 (MFI) (%) = 100 * (mesenteric fat weight / body weight). Protein efficiency rate (PER)
190 = (FBW - IBW) / protein intake. Gross protein efficiency (GPE) (%) = 100 * [(% final
191 body protein * FBW) - (% initial body protein * IBW)] / total protein intake fish. Gross

192 lipid efficiency (GLE) (%) = 100 * [(% final body lipid * FBW) - (% initial body lipid *
193 IBW)] / total lipid intake fish.

194

195 *2.5 Histology*

196

197 After euthanasia of 5 fish per tank, the gut was removed and the intestine was divided
198 into two segments (midgut and hindgut). From each segment, a 5 mm-long piece was
199 sectioned and fixed in 10% buffered formalin. Samples were processed for routine
200 histology to obtain a transversal section, which was stained with haematoxylin and eosin
201 (H&E). Sections were evaluated blind under a light microscope (Nikon Eclipse 80i,
202 Nikon Corporation, Japan) to verify the preservation of the normal intestinal architecture.
203 In particular, the histological investigation was focused on the main cell constituents of
204 the mucosal layer (goblet cells, supranuclear absorption vacuoles in the enterocytes),
205 capillary within the intestinal folds, lymphoplasmacytic cells within lamina propria
206 (GALT-like tissue). Moreover, any possible degenerative and diet adaptive induced
207 changes were taken into consideration. Photographs of the sections were made with a
208 Nikon Eclipse 80i microscope (Nikon, Tokyo, Japan) equipped with a Nikon Digital Sight
209 SD-MS camera and the Nikon software NIS-Elements. Adobe Photoshop CS3 Extended
210 was used for the final photographic preparation without altering the original integrity of
211 the pictures.

212

213 *2.6 Gastrointestinal evacuation, time and digesta characteristics*

214

215 Following the feeding trial, sampling for gastrointestinal evacuation pattern and
216 digesta characteristics was conducted. We adopted the approach used in the studies by
217 Adamidou et al. (2009) and Nikolopoulou et al. (2011). Fish were kept fasting for 72 h
218 before being fed to make sure that the gastrointestinal tract was empty. Four fish per tank
219 were sacrificed at 1, 4, 10, 16, 28 and 48 h after feeding a single meal to satiation. Each
220 sampled fish was weighed, then the abdominal cavity was opened and the digestive tract
221 carefully removed and separated into three parts: stomach, midgut and hindgut. Midgut
222 was defined as the section from the pyloric sphincter to the ileorectal valve and hindgut
223 from the ileorectal valve to the anus. Stomach and intestinal contents from the above
224 sections were collected in pre-weighed dishes, weighed for each fish separately, dried and
225 reweighed. The measured weights were used for the calculations described below. The
226 geometric means of stomach and intestinal dry digesta content divided by the fish weight
227 were regressed against time separately for each diet, in order to fit to a model for
228 calculating gastric evacuation rate (GER) and gastric evacuation time (GET) for the
229 stomach, midgut and hindgut filling time (MFT, HFT) and midgut and hindgut evacuation
230 time (MET, HET) for midgut and hindgut.

231

232 *2.7 Stomach evacuation pattern calculation*

233

234 In the case of stomach, GER was calculated according to the formula described by
235 Elliott (1972) and adapted by Nikolopoulou et al. (2011) The geometric mean of the
236 stomach dry digesta content divided by the fish weight was regressed against time
237 separately for each diet. GER is estimated as the value of r of the regression model:

238 $W_t = A e^{-rt}$

239 which is equivalent to the semi-logarithmic model:

$$240 \ln W_t = \ln A - rt$$

241 where W_t is the geometric mean weight of stomach dry matter digesta at time t , A is an
242 intercept estimated from the model regression and r is the rate of gastric evacuation. GET
243 50%, GET 75% and GET 90% is the evacuation time (expressed in hours) required to
244 empty 50%, 75% and 90% of the stomach. It was computed through:

$$245 \text{GET } p\% = [\ln 100 - \ln(100 - p)] / r$$

246 where p is the digestible organic matter to be evacuated from the stomach.

247

248 *2.8 Midgut and Hindgut Filling/Evacuation pattern calculations*

249

250 In the case of midgut and hindgut the best model is the quadratic regression,
251 represented as a parabola. It was not possible to calculate GER, because the trend of
252 digesta in midgut and hindgut was not linear. Points from each curve were estimated to
253 determine the time of maximum midgut and hindgut filling and evacuation time. The
254 MFT and HFT, (maximum filling time) was calculated as the vertex of the parabola. In
255 these cases, the equations used were:

$$256 W_t = A + r_1 t + r_2 t^2$$

257 where r_1 is the coefficient of the linear part of the model and r_2 is the parameter of the
258 quadratic part of the model.

259 The vertex of the parabola:

$$260 [-(r_1/2r_2), (4Ar_2 - r_1^2)/4r_2]$$

261 and MFT and HFT:

$$262 -(r_1/2r_2)$$

263 MET and HET 50%, 75% and 90% is the evacuation time (expressed in hours) required
264 to empty 50%, 75% and 90% of midgut and hindgut and was computed through:

$$265 (1-p)*[(4Ar_2 - r_1^2)/4r_2]=A+ r_1t+r_2t^2.$$

266 MET and HET 50% , 75% and 90% is the solution with respect to t of the equation:

$$267 A-(1-p)*[(4Ar_2 - r_1^2)/4r_2]+ r_1t+r_2t^2=0$$

268 by Newton Raphson optimization where p=0.50, 0.75, 0.90.

269

270 *2.9 Digesta moisture determination*

271

272 The weights of stomach and intestine contents, at each sampling time, were used to
273 calculate the moisture of digesta as percentage of digesta weight in each gastrointestinal
274 segment, according to Nikolopoulou et al. (2011).

275

276 *2.10 Analytical methods*

277

278 Diets and whole body were analysed for proximate composition. Moisture content was
279 obtained by weight loss after drying samples in a stove at 105 °C until a constant weight
280 was achieved. Crude protein was determined as total nitrogen (N) by using the Kjeldahl
281 method and multiplying N by 6.25. Total lipids were determined according to Bligh and
282 Dyer's (1959) extraction method. Ash content was estimated by incineration to a constant
283 weight in a muffle oven at 450 °C. NDF, acid-detergent fibre (ADF) and lignin were
284 determined according to Van Soest and Robertson, (1981). Cellulose content was
285 determined based on the ADF–lignin. Hemicellulose content was determined based on
286 NDF–ADF content. Gross energy was determined by a calorimetric bomb (Adiabatic

287 Calorimetric Bomb Parr 1261; PARR Instrument, IL, U.S.A.). Serum glucose, alkaline
288 phosphatase, inorganic phosphorus, triglycerides and total protein were measured using
289 colorimetric methods (Total protein OSR6232, Triglyceride OSR61118, Glucose
290 OSR6121, Alkaline phosphatase OSR6004, Inorganic phosphorus OSR6122; Beckman
291 Coulter, Brea, CA, USA) on an automated analyser (AU 400, Beckman Coulter, Brea
292 CA, USA).

293

294 *2.11 Statistical analysis*

295

296 All data are presented as mean \pm standard deviation (SD). Tank was used as the
297 experimental unit for analyzing growth and performance, a pool of five sampled fish was
298 considered the experimental unit for analyzing carcass composition, whereas individual
299 fish was used for analyzing VSI, HSI, MFI, and blood parameters. Data of growth
300 performance, VSI, HSI, MFI, nutritional indices and blood parameters were analysed by
301 a one-way ANOVA and in case of significance ($P \leq 0.05$) Tukey's post hoc test was
302 performed. The normality and/or homogeneity of variance assumptions were validated
303 for all data preceding ANOVA. Moisture of digesta was analysed by two-way ANOVA
304 using diet and time as independent factors. Statistical analysis was performed using the
305 software R version 3.1.0 (Revolution analytics, Palo Alto, CA, USA). The differences
306 among treatments were considered significant at $P \leq 0.05$.

307

308 **3. Results**

309

310 *3.1 Growth and blood biochemistry*

311

312 Growth performances are summarised in Table 2. No significant differences due to
313 fibre inclusion levels were observed after the 117 days in terms of growth performance
314 (final body weight and SGR), feed intake (VFI) and feed utilisation (FCR). Data on
315 biometric indices, body composition and nutritional indices are shown in Table 3. VSI
316 and HSI values significantly decreased with increasing fibre dietary levels, while no
317 significant differences were found in MFI. Regarding whole body protein content, fish
318 fed diet F7.2 showed higher value in comparison to those fed F13.1, while no significant
319 differences were found among the lipid, ash and moisture content of fish. No significant
320 differences among treatments were found in PER, GPE and GLE. Serum TP, GLU, TRIG,
321 ALP and P levels are shown in Table 4. No significant differences among treatments were
322 found. Data on P were significant for ANOVA ($P = 0.0469$) but no differences among
323 treatments were detected by the multiple comparison test.

324

325 *3.2 Histology*

326

327 All the histological sections showed normal intestinal architecture. No differences
328 were found in the mucosal layer (goblet cells, supranuclear absorption vacuoles in the
329 enterocytes), capillarity network within the intestinal folds, lymphoplasmacytic cells
330 within the *lamina propria*. Moreover, inflammatory and/or degenerative changes were
331 not present in the histological sections taken from the examined subjects (Fig. 1).

332

333 *3.3 Gut evacuation rate and time*

334

335 Data on GER, GET, MFT, HFT, MET and HET are presented in Table 5. Dry weight
336 of the stomach content decreased with time as shown in Fig. 2. No significant differences
337 were found among treatments for stomach, midgut and hindgut however diet F7.2
338 required 22 h to evacuate 90% of the initial digesta content, while diet F13.1 and F15.5
339 required 38 h and 35 h respectively. The data extracted for the midgut and for the hindgut
340 evacuation fitted a quadratic model (Fig. 3 a, b). According to the equations, the time
341 required to empty the whole gastrointestinal tract was estimated to be around 46-47 h for
342 fish fed under all treatments. Data on moisture of digesta for each gastrointestinal tracts
343 at 1, 4, 10, 16, 28, and 48 h post prandial are shown in Table 6. No differences were found
344 between treatments, but moisture in the stomach changed with time increasing from 1 to
345 4 h postprandial and then remained constant until 28 h postprandial.

346

347 **4. Discussion**

348

349 To our knowledge, this is the first study designed to specifically evaluate the effects of
350 graded levels of dietary insoluble fibre derived from sunflower hulls and soybean hulls
351 on performance, health and digesta transit time of European sea bass using formulations
352 at practical level. Crude fibre which is basically cellulose, insoluble hemicellulose and
353 lignin, does not have any value in the nutrition of carnivorous fish and according to Altan
354 and Korkut (2011) should be restricted to less than 7 % in fish diet. The results of our
355 study demonstrated that neutral detergent fibre (hemicellulose, cellulose and lignin) up to
356 level of 15.5 % had no effects on growth performance, feed intake and protein/lipid
357 utilisation. The performances registered in this trial are similar to those found in other
358 studies on European sea bass (Tulli et al., 2010; Guerreiro et al., 2015; Bonvini et al.,

2018). Hansen and Hemre (2013) reported that plant protein can contain soluble fibres and antinutrients which can interfere with nutrient digestibility. Soluble fibres increase the viscosity of gut content, which potentially can reduce digestible enzyme activities and negatively affect nutrient digestion and absorption (Leenhouwers et al., 2006). Recently, Adamidou et al. (2011) reported that an inclusion of fibre up to a level of 5 g 100 g⁻¹ did not affect growth performance or nutrient digestibility in sharpsnout seabream (*Diplodus puntazzo*). Bou et al. (2014) have demonstrated that using diets with the inclusion of fibre from cellulose up to 18 % at the expense of carbohydrate did not affect growth performance in gilthead sea bream (*Sparus aurata*). Both sharpsnout and gilthead sea bream belong to the family of Sparidae, which are considered omnivorous fish and more predisposed to digest plant-based ingredients than carnivorous species, such as European sea bass (Stone, 2003). In general, dietary fibre did not affect whole body composition except for the protein content which was slightly lower in F13.1 compared to F7.2 but without resulting in lower protein efficiency.

VSI and HSI values increased in diets with the lowest fibre inclusion. No evidence is reported in the literature about any influence of fibre levels on somatic indices. One possible explanation of the trend in HSI, found in our study, can be related to the decrease in level of wheat in diets at the increasing of fibre. Previous studies on European sea bass and other species reported that increasing NFE, wheat and starch level in the diets can increase HSI values probably due to an enhancement in glycogen liver deposition (Bonaldo et al., 2008, 2010; Bonvini et al., 2017; Bou et al., 2014).

Information on the nutritional status and health in fish species can be achieved through the study of blood metabolites (Bonaldo et al., 2015; Bonvini et al., 2015). To our knowledge, no studies have assessed blood parameters in response to fibre inclusion in

383 diets in marine species. Among the blood parameters, TP, TRIG, GLU, ALP and P seem
384 to have potential as predicative diagnostic tools for evaluation of European sea bass
385 nutritional status (Peres et al., 2014). In this trial, no differences were found among
386 treatments in all parameters analysed. TP level is usually very stable in well-nourished
387 animals but decreases under fasting conditions (Peres et al., 2014). In this study, plasma
388 TP averaged between 5.6 and 5.1 g dl⁻¹ and these values are within the ranges reported
389 for European sea bass under good nutritional status (Peres et al., 2014). TRIG levels are
390 in agreement with previously reported values for this species (Bonvini et al., 2017), but
391 higher than those reported by Peres et al. (2014), a difference which can be related to the
392 different lipid level and formulation of the diet used, in comparison to our study. GLU
393 levels are in agreement with previously reported values for this species (Peres et al.,
394 2014). Adamidou et al. (2009), reported that GLU levels in European sea bass serum were
395 also affected by the type of carbohydrate ingested, with wheat starch showing the most
396 rapid increase and decrease in serum glucose compared to fish fed pea and chickpea diets,
397 while faba bean starch resulted in a delay in serum glucose peak and a lower range of
398 glucose values. ALP levels are higher in comparison to Peres et al. (2014). Alkaline
399 phosphatase is involved in the absorption and transport of lipid and carbohydrates from
400 the intestine, and intestinal activities are positively correlated with food ingestion and
401 growth rate (Lemieux et al., 1999). Finally, inorganic phosphorus levels are in agreement
402 with previously reported values for this species (Peres et al., 2014). Plasma electrolytes
403 (univalent and bivalent) are considered to be valuable indicators of secondary stress and
404 osmoregulation ability in fish (Roque et al., 2010). Plasma phosphorus has been identified
405 as a good indicator of stress (i.e. starvation and stocking density) and pathological
406 situations (Roque et al., 2010; Peres et al., 2014).

407 Besides the study of blood metabolites, histological analysis of the digestive system is
408 considered a valid tool for evaluating the gut health status. No inflammatory and/or
409 degenerative changes were recorded in any of the histological sections of the gut. The
410 examination revealed no alteration in the mucosal layer (goblet cells and supranuclear
411 absorption vacuoles in the enterocytes), no vascular changes within the intestinal folds
412 and no lymphoplasmacytic infiltration within the *lamina propria*. Few references are
413 available in literature on the effects of fibre on fish intestine. For example, Olsen et al.
414 (2007) reported in Atlantic cod an increase in cellularity of the *lamina propria* and a
415 modest goblet cell hypertrophy and hyperplasia especially in fish fed a diet with 100% of
416 plant protein mixture. These results were ascribed to a significant amount of fibres in
417 plant-based diets. In our study these morphological changes were not observed in any of
418 the diet treatments confirming a good tolerance of European sea bass intestine to a fibre
419 content up to 15.5%. Leigh et al. (2017) reported in zebrafish (*Danio rerio*) changes in
420 the epithelial surface area and overall gut length as a response to fibre intake level but
421 without any inflammatory or degenerative effects.

422 Dietary fibre can affect gastrointestinal transit time of feed (Zhou et al., 2004). The
423 investigation on gastrointestinal evacuation rate/time in this study reported no significant
424 differences between treatments, however higher fibre dietary levels led to an increased
425 time required to empty the stomach. A marked difference in term of mean values was
426 infact observed between diets F7.2, F13.1, F15.5 which required 22, 38 and 35 h,
427 respectively to empty 90% of the stomach content. In addition slower evacuation was also
428 observed in F13.1 and F15.5 compared to F7.2 to empty 50% and 75% of the stomach
429 content. In the present trial if we consider that the inclusion of high dietary fibre level did
430 not exert a negative effect on performance and feed utilization, a possible explanation

431 may be related to a reduced gastric evacuation at high fibre level which might have
432 improved feed digestion and absorption.

433 On the other hand, the evacuation time for the hindgut was very similar in all the
434 treatments, around 46-47 h to empty 90% of hindgut content. There is little published
435 information regarding the effects of fibre on gastric evacuation in marine carnivorous
436 species. When plant-based ingredients are included in the diet, gastric evacuation time is
437 higher, increasing more with legumes than with cereals (Adamidou et al., 2009).
438 Moreover, Venou et al. (2003) reported that differences in ingredient processing can also
439 modify gastric evacuation time, which is higher for extruded cereals than for raw cereals
440 in gilthead sea bream. In the same species, García-Meilán et al. (2014) found that
441 differences in diet composition, such as high lipid levels or high starch content, may be
442 involved in the differential transit rate and Fountoulaki et al. (2005) found that a low
443 transit rate was related to a high lipid content. Since in the present work diets were
444 designed at a practical level in order to identify the maximum amount of dietary fibre
445 inclusion without negatively affecting zootechnical performance and health, we cannot
446 totally exclude the effects that the differences in the lipid content could have exerted on
447 the gastric evacuation pattern. Moisture of digesta in the gastrointestinal tract showed no
448 differences between treatments. Moisture of stomach increased with time from 1 to 4 h
449 after feeding and remained unchanged until 28 h post prandial. The water required for
450 feed moisturisation originates from feed water, initial water absorption of pellets,
451 drinking and stomach secretions (Kristiansen and Rankin, 2001). In this context, dietary
452 fibre behaves within the gastrointestinal tract as a polymer matrix with variable
453 physicochemical properties including water-holding capacity (Kay, 1982).

454

455 **5. Conclusion**

456

457 In conclusion, the different fibre levels derived from sunflower hulls and soybean hulls
458 tested in this trial had no effects on overall performances and feed efficiency in European
459 sea bass. Results from blood biochemistry profile and gut histology confirm a good
460 nutritional and health status of fish under all feeding treatments. The inclusion of fibres
461 derived from sunflower and soybean hulls had no influence also on digesta transit time
462 and digesta moisture even if higher dietary fibre levels seem to increase the time required
463 to empty the stomach. According to the results it seems feasible to include insoluble fibre
464 up to 15.5% in feed formulation for the on-growing of European sea bass without
465 negatively affecting zootechnical performance and health.

466

467 **Acknowledgments**

468 The authors would like to thank Paola Parazza and Marina Silvi for technical
469 assistance during laboratory analyses. The author would like to thank Gillian Forlivesi
470 Heywood for English language editing. This research did not receive any specific grant
471 from funding agencies in the public, commercial, or not-for-profit sectors. The authors
472 declare no conflict of interest.

473

474 **References**

475

476 Adamidou, S., Nengas, I., Alexis, M., Fountoulaki, E., Nikolopoulou, D., Campbell, P.,
477 Karacostas, I., Rigos, G., Bell, G.J., Jauncey, K., 2009. Apparent nutrient
478 digestibility and gastrointestinal evacuation time in European seabass

479 (*Dicentrarchus labrax*) fed diets containing different levels of legumes.
480 Aquaculture 289, 106-112.

481 Adamidou, S., Rigos, G., Mente, E., Nengas, I., Foundoulaki, E., 2011. The effects of
482 dietary lipid and fibre levels on digestibility of diet and on the growth performance
483 of sharpsnout seabream (*Diplodus puntazzo*). Mediterr. Mar. Sci. 12, 401-412.

484 Altan, O., Korkut, A.Y., 2011. Apparent Digestibility of Plant Protein Based Diets by
485 European Sea Bass *Dicentrarchus labrax* L., 1758. Turk. J. Fish. Aquat. Sci. 11,
486 87-92.

487 Amirkolaie, A.K., Leenhouders, J.I., Verreth, J.A.J., Schrama, J.W., 2005. Type of
488 dietary fibre (soluble versus insoluble) influences digestion, faeces characteristics
489 and faecal waste production in Nile tilapia (*Oreochromis niloticus* L.). Aquacult.
490 Res., 36, 1157–1166.

491 Bach Knudsen, K.E., 2001. The nutritional significance of “dietary fibre” analysis. Anim.
492 Feed Sci. Technol. 90, 3–20.

493 Bligh, E.G., Dyer, W.J., 1959. A rapid method of total lipid extraction and purification.
494 Can. J. Biochem. Phys. 37, 911–917.

495 Bonaldo, A., Roem, A.J., Fagioli, P., Pecchini, A., Cipollini, I., Gatta, P.P., 2008.
496 Influence of dietary levels of soybean meal on the performance and gut histology
497 of gilthead sea bream (*Sparus aurata* L.) and European sea bass (*Dicentrarchus*
498 *labrax* L.). Aquacult. Res. 39, 970-978.

499 Bonaldo, A., Isani, G., Fontanillas, R., Parma, L., Grilli, E., Gatta, P.P., 2010. Growth
500 and feed utilization of gilthead sea bream (*Sparus aurata*, L.) fed to satiation and
501 restrictively at increasing dietary energy levels. Aquacult. Int. 18, 909-919.

502 Bonaldo, A., Di Marco, P., Petocchi, T., Marino, G., Parma, L., Fontanillas, R., Koppe,
503 W., Mongile, F., Finoia, M.G., Gatta, P.P., 2015. Feeding turbot juveniles *Psetta*
504 *maxima* L. with increasing dietary plant protein levels affects growth performance
505 and fish welfare. *Aquacult. Nutr.* 21, 401-413.

506 Bonvini, E., Parma, L., Mandrioli, L., Sirri, R., Brachelente, C., Mongile, F., Gatta, P.P.,
507 Bonaldo, A., 2015. Feeding common sole (*Solea solea*) juveniles with increasing
508 dietary lipid levels affects growth, feed utilization and gut health. *Aquaculture*
509 449, 87-93.

510 Bonvini, E., Bonaldo, A., Mandrioli, L., Sirri, R., Dondi, F., Bianco, C., Fontanillas, R.,
511 Mongile, F., Gatta, P.P., Parma, L., 2017. Effects of feeding low fishmeal diets
512 with increasing soybean meal levels on growth, gut histology and plasma
513 biochemistry of sea bass. *Animal* 1–8. doi:10.1017/S1751731117002683.

514 Bonvini, E., Parma, L., Badiani, A., Fontanillas, R., Gatta, P.P., Sirri, F., Nannoni, E.,
515 Bonaldo, A., 2018. Integrated study on production performance and quality traits
516 of European sea bass (*Dicentrarchus labrax*) fed high plant protein diets,
517 *Aquaculture*, 484, 126-132.

518 Bou, M., Todorčević, M., Fontanillas, R., Capilla, E., Gutiérrez, J., Navarro, I., 2014.
519 Adipose tissue and liver metabolic responses to different levels of dietary
520 carbohydrates in gilthead sea bream (*Sparus aurata*). *Comp. Biochem. Physiol. A*
521 175, 72–81.

522 Dalsgaard, J., Bach Knudsen, K.E., Verlhac, V., Ekmann, K.S., Pedersen, P.B., 2016.
523 Supplementing enzymes to extruded, soybean-based diet improves breakdown of
524 non-starch polysaccharides in rainbow trout (*Oncorhynchus mykiss*). *Aquac. Nutr.*
525 22, 419–426.

526 Elliott, J.M., 1972. Rates of gastric evacuation in brown trout, *Salmo trutta* L. Freshw.
527 Biol. 2, 1-18.

528 Eusebio, P.S., Coloso, R.M., Mamauag, R.E.P., 2004. Apparent digestibility of selected
529 ingredients in diets for juvenile grouper, *Epinephelus coiodes* (Hamilton).
530 Aquacult. Res. 35, 1261-1269.

531 Fountoulaki, E., Alexis, M.N., Nengas, I., Venou, B., 2005. Effect of diet composition on
532 nutrient digestibility and digestive enzyme levels of gilthead sea bream (*Sparus*
533 *aurata* L.). Aquacult. Res. 26, 1243–1251.

534 García-Meilán, I., Ordóñez-Grande, B., Gallardo, M.A., 2014. Meal timing affects
535 protein-sparing effect by carbohydrates in sea bream: Effects on digestive and
536 absorptive processes. Aquaculture 434, 121-128.

537 Glencross, B., 2009. The influence of soluble and insoluble lupin non-starch
538 polysaccharides on the digestibility of diets fed to rainbow trout (*Oncorhynchus*
539 *mykiss*). Aquaculture, 294, 256–261.

540 Glencross, B., Rutherford, N., Bourne, N., 2012. The influence of various starch and non-
541 starch polysaccharides on the digestibility of diets fed to rainbow trout
542 (*Oncorhynchus mykiss*). Aquaculture, 356–357, 141–146.

543 Guerreiro, I., Oliva-Teles, A., Enes, P., 2015. Improved glucose and lipid metabolism in
544 European sea bass (*Dicentrarchus labrax*) fed short-chain fructooligosaccharides
545 and xylooligosaccharides. Aquaculture 441, 57–63.

546 Hansen, J.O., Storebakken, T., 2007. Effects of dietary cellulose level on pellet quality
547 and nutrient digestibilities in rainbow trout (*Oncorhynchus mykiss*). Aquaculture,
548 272, 458–465.

549 Hansen, A.C., Hemre, G.I., 2013. Effect of replacing fish meal and oil with plant
550 resources in on-growing diets for Atlantic cod (*Gadus morhua* L.). *Aquacult.*
551 *Nutr.* 19, 641-650.

552 Kay, R.M., 1982. Dietary fiber. *J. Lipid Res.* 23, 221-242.

553 Kousoulaki, K., Sæther, B.S., Albrektsen, S., Noble, C., 2015. Review on European sea
554 bass (*Dicentrarchus labrax*, Linnaeus, 1758) nutrition and feed management: a
555 practical guide for optimizing feed formulation and farming protocols. *Aquacult.*
556 *Nutr.* 21, 129-151.

557 Kraugerud, O.F., Penn, M., Storebakken, T., Refstie, S., Krogdahl, A., Svihus, B., 2007.
558 Nutrient digestibilities and gut function in Atlantic salmon (*Salmo salar*) fed diets
559 with cellulose or non-starch polysaccharides from soy. *Aquaculture*, 273, 96–107.

560 Kristiansen, H.R., Rankin, J.C., 2001. Discrimination between endogenous and
561 exogenous water sources in juvenile rainbow trout fed extruded dry feed. *Aquat.*
562 *Living Resour.* 14, 359–366.

563 Krogdahl, A., Hemre, G.I., Mommsen, T.P., 2005. Carbohydrates in fish nutrition;
564 digestion and absorption in postlarval stages. *Aquacult. Nutr.* 11, 103-122.

565 Leenhouwers, K.I., Adjei-Boateng, D., Verreth, J.A.J., Schrama, J.W., 2006. Digesta
566 viscosity, nutrient digestibility and organ weights in African catfish (*Clarias*
567 *gariiepinus*) fed diets supplemented with different levels of a soluble non-starch
568 polysaccharide. *Aquacult. Nutr.* 12, 111–116.

569 Leigh, S.C., Nguyen-Phuc, B.Q., German, D.P., 2017. The effects of protein and fiber
570 content on gut structure and function in zebrafish (*Danio rerio*). *J Comp Physiol*
571 *B.* doi: 10.1007/s00360-017-1122-5.

572 Lekva, A., Hansen, A.C., Rosenlund, G., Karlsen, O., Hemre, G.I., 2010. Energy dilution
573 with alpha-cellulose in diets for Atlantic cod (*Gadus morhua* L.) juveniles –
574 effects on growth, feed intake, liver size and digestibility of nutrients.
575 *Aquaculture*, 300, 169–175.

576 Lemieux, H., Blier, P., Dutil, J.D., 1999. Do digestive enzymes set a physiological limit
577 on growth rate and food conversion efficiency in the Atlantic cod (*Gadus*
578 *morhua*)? *Fish Physiol. Biochem.* 20, 293-303.

579 Mongile, F., Bonaldo, A., Fontanillas, R., Mariani, L., Badiani, A., Bonvini, E., Parma,
580 L., 2014. Effects of dietary lipid level on growth and feed utilisation of gilthead
581 seabream (*Sparus aurata* L.) reared at Mediterranean summer temperature. *Ital. J.*
582 *Anim. Sci.* 13, 30-35.

583 Nikolopoulou, D., Moutou, K.A., Fountoulaki, E., Venou, B., Adamidou, S., Alexis,
584 M.N., 2011. Patterns of gastric evacuation, digesta characteristics and pH changes
585 along the gastrointestinal tract of gilthead sea bream (*Sparus aurata* L.) and
586 European sea bass (*Dicentrarchus labrax* L.). *Comp. Biochem. Physiol. A* 158,
587 406-414.

588 Nutrient requirements of fish and shrimp (NRC) 2011. National Academic Press,
589 Washington DC, USA.

590 Olsen, R.E., Hansen, A.C., Rosenlund, G., Hemre, G.I., Mayhew, T.W., Knudsen, D.L.,
591 Eroldogan, O.T., Myklebust, R., Karlsen, Ø., 2007. Total replacement of fish meal
592 with plant proteins in diets for Atlantic cod (*Gadus morhua* L.) II: Health aspects.
593 *Aquaculture* 272, 612–624.

594 Parma, L., Candela, M., Soverini, M., Turrone, S., Consolandi, C., Brigidi, P., Mandrioli,
595 L., Sirri, R., Fontanillas, R., Gatta, P.P., Bonaldo, A., 2016. Next-generation

596 sequencing characterization of the gut bacterial community of gilthead sea bream
597 (*Sparus aurata*, L.) fed low fishmeal based diets with increasing soybean meal
598 levels. Anim. Feed Sci. Technol. 222, 204-216.

599 Peres, H., Santos, S., Oliva-Teles, A., 2014. Blood chemistry profile as indicator of
600 nutritional status in European seabass (*Dicentrarchus labrax*). Fish Physiol.
601 Biochem. 40, 1339-1347.

602 Roque, A., Yildiz, H.Y., Carazo, I., Duncan, N., 2010. Physiological stress responses of
603 seabass (*Dicentrarchus labrax*) to hydrogen peroxide (H₂O₂) exposure.
604 Aquaculture 304, 104-107.

605 Stone, D.A.J., 2003. Dietary carbohydrate utilization by fish. Rev. Fish. Sci. 11, 337-369.

606 Tulli, F., Messina, M., Calligaris, M., Tibaldi, E., 2010. Response of European sea bass
607 (*Dicentrarchus labrax*) to graded levels of methionine (total sulfur amino acids)
608 in soya protein-based semi-purified diets. Br. J. Nutr. 104, 664-673.

609 Van Soest, P.J., Robertson, J.B., 1981. The detergent system of analysis and its
610 application to human foods. Chapter 8. In: James, W.P.T., Theander, O. (Eds.),
611 The Analysis of Dietary Fibre in Food. Marcel Dekker Inc., New York, p. 123.

612 Venou, B., Alexis, M.N., Fountoulaki, E., Nemes, I., Apostolopoulou, M. Castritsi-
613 Cathariou, I., 2003. Effect of extrusion of wheat and corn on gilthead sea bream
614 (*Sparus aurata*) growth, nutrient utilization efficiency, rates of gastric evacuation
615 and digestive enzyme activities. Aquaculture 225, 207-223.

616 Zhou, Q.C., Tan, B.P., Mai, K.S., Liu, Y.J., 2004. Apparent digestibility of selected feed
617 ingredients for juvenile cobia *Rachycentron canadum*. Aquaculture 241, 441-451.

Table 1. Ingredients and proximate composition of the experimental diets

	F7.2	F8.9	F11.5	F13.1	F15.5
<i>Ingredient, % of the diet</i>					
Sunflower Hulls	0.0	1.5	3.2	4.8	6.4
Soybean Hulls	0.0	2	4.2	6.4	8.6
Fish meal	20	20	20	20	20
Soybean concentrate	13	13	13	13	13
Wheat	28.4	23.3	17.9	12.4	6.9
Corn gluten	8	8	8	8	8
Wheat gluten	15.2	15.6	16.1	16.6	17
Fish oil	7.5	8	8.6	9.2	9.8
Rapeseed oil	7.5	8	8.6	9.2	9.8
Vit/Min premix ¹	0.5	0.5	0.5	0.5	0.5
<i>Proximate composition, % on a wet weight basis</i>					
Moisture	7.28	7.77	7.71	7.49	7.74
Protein	43.6	43.2	43.5	43.8	43.7
Lipid	22.5	24.3	24.8	25.4	27.1
Ash	4.42	4.47	4.53	4.63	4.73
NDF ²	7.20	8.93	11.46	13.14	15.46
ADF ³	2.79	4.45	5.99	7.11	9.26
Hemicellulose	4.41	4.48	5.48	6.03	6.20
Cellulose	2.33	3.37	4.73	5.73	7.07
Lignin	0.46	1.08	1.26	1.38	2.20
Gross energy MJ kg ⁻¹	21.5	22.1	22.1	22.5	23.0

¹ Vitamin and mineral premix; Skretting, Stavanger, Norway (fulfilling recommendations for marine fish species given by NRC, 2011)

²Neutral detergent fibre

³Acid detergent fibre

618

619

620

621

622

623

624

Table 2. Growth performance and feed intake of European sea bass fed experimental diets over 117 days.

	Experimental diets					<i>P</i> value
	F7.2	F8.9	F11.5	F13.1	F15.5	
IBW (g)	69.5 ± 2.1	68.1 ± 4.0	68.7 ± 1.1	69.7 ± 1.3	71.2 ± 2.5	0.5944
FBW (g)	215 ± 11.7	204 ± 14.9	215 ± 15.9	223 ± 6.3	219 ± 15.8	0.5369
SGR (% day ⁻¹)	0.97 ± 0.03	0.94 ± 0.02	0.97 ± 0.07	0.99 ± 0.01	0.96 ± 0.04	0.5136
VFI (g feed fish ⁻¹)	237 ± 7.7	219 ± 7.7	230 ± 11.6	232 ± 8.1	237 ± 13.4	0.2759
FCR	1.63 ± 0.06	1.43 ± 0.24	1.58 ± 0.11	1.51 ± 0.03	1.60 ± 0.06	0.3773
Survival %	100 ± 0.0	100 ± 0.0	99.4 ± 1.0	100 ± 0.0	100 ± 0.0	0.4516

Data are given as the mean (n=3) ± SD. In each line, different superscript letters indicate significant differences among treatments ($P \leq 0.05$).

IBW = Initial body weight.

FBW = Final body weight.

SGR = Specific growth rate (% day⁻¹) = 100 * (ln FBW - ln IBW) / days.

VFI = Voluntary Feed Intake (g feed/fish) = g feed ingested / number of fish

FCR = Feed conversion rate = feed intake / weight gain.

625

626

627

Table 3. Biometric indices, body composition and nutritional indices of European sea bass fed the experimental diets

	Experimental diets					<i>P</i> value
	F7.2	F8.9	F11.5	F13.1	F15.5	
Biometric indices						
VSI	12.1 ^b ± 1.5	11.4 ^{ab} ± 1.8	11.3 ^{ab} ± 2.5	11.1 ^{ab} ± 1.4	10.5 ^a ± 1.9	0.0303
HSI	3.2 ^c ± 0.5	3.1 ^c ± 0.5	2.7 ^b ± 0.6	2.6 ^{ab} ± 0.5	2.3 ^a ± 0.5	<0.0001
MFI	6.7 ± 1.5	6.1 ± 1.8	6.4 ± 2.5	6.3 ± 1.5	5.8 ± 1.6	0.4604
Whole body composition, %						
Protein	17.1 ^b ± 0.1	16.4 ^{ab} ± 0.3	16.6 ^{ab} ± 0.3	16.2 ^a ± 0.5	16.3 ^{ab} ± 0.3	0.0465
Lipid	17.6 ± 1.3	18.6 ± 1.4	17.6 ± 2.3	17.8 ± 1.9	19.0 ± 1.1	0.7498
Ash	2.8 ± 0.21	2.7 ± 0.03	2.7 ± 0.07	2.7 ± 0.04	2.8 ± 0.15	0.7554
Moisture	61.6 ± 0.8	61.2 ± 0.4	60.9 ± 0.5	60.3 ± 0.5	60.0 ± 1.9	0.3485
Nutritional indices						
PER	1.41 ± 0.05	1.43 ± 0.08	1.46 ± 0.11	1.51 ± 0.03	1.43 ± 0.05	0.4484
GPE	24.4 ± 0.97	23.5 ± 1.78	24.2 ± 1.24	24.2 ± 0.62	23.2 ± 0.65	0.6601
GLE	56.4 ± 7.1	57.2 ± 8.2	52.9 ± 11.7	54.2 ± 6.3	52.7 ± 4.1	0.9343

Data are given as the mean (n=3; n=30 for VSI, HSI, MFI) ± SD. In each line, different superscript letters indicate significant differences among treatments ($P \leq 0.05$).

VSI = Viscerosomatic index (%) = 100*(viscera weight/FBW).

HSI = Hepatosomatic index (%) = 100*(liver weight/FBW).

MFI = Mesenteric fat index (%) = 100*(mesenteric fat weight/FBW).

PER = Protein efficiency ratio = ((FBW-IBW)/protein intake).

GPE = Gross protein efficiency = 100*[(%final body protein*FBW) - (%initial body protein*IBW)]/total protein intake fish.

GLE = Gross lipid efficiency = 100*[(%final body lipid*FBW) - (%initial body lipid*IBW)]/total lipid intake fish.

628

629

630

631

632

633

634

635

636

Table 4. Blood biochemistry of European sea bass fed the experimental diets

	Experimental diets					<i>P</i> value
	F7.2	F8.9	F11.5	F13.1	F15.5	
TP (g dL ⁻¹)	5.5 ± 0.40	5.6 ± 0.98	5.3 ± 0.70	5.5 ± 0.63	5.1 ± 1.16	0.5586
TRIG (mg dL ⁻¹)	1884 ± 818	2023 ± 600	1769 ± 788	1790 ± 715	1797 ± 670	0.9064
GLU (mg dL ⁻¹)	98 ± 55.1	90 ± 30.1	77 ± 17.3	89 ± 26.0	111 ± 33.8	0.2544
ALP (U L ⁻¹)	235 ± 58.9	252 ± 59.1	227 ± 56.7	207 ± 49.7	210 ± 51.9	0.2300
P (mg dL ⁻¹)	8.4 ± 0.85	9.4 ± 1.21	8.5 ± 0.72	8.4 ± 1.01	8.9 ± 1.00	0.0469

Each value is mean from 12 samples ± SD. Data on P were significant for ANOVA ($P = 0.0469$) but no differences among treatments were detected by the multiple comparison test.

TP = total protein

TRIG = triglycerides

GLU = glucose

ALP = alkaline phosphatase

P = inorganic phosphorus

Table 5. Gastric evacuation rate (GER), gastric evacuation time (GET, expressed in hours), midgut filling and evacuation time (MFT, MET, h) and hindgut filling and evacuation time (HFT, HET, h) of European sea bass fed the experimental diets

	Experimental diets				
	F7.2	F8.9	F11.5	F13.1	F15.5
<i>Stomach</i>					
GER	0.1031	0.0945	0.0899	0.0607	0.0657
GET 50% (h)	6.72	7.33	7.71	11.43	10.55
GET 75% (h)	13.44	14.66	15.41	22.86	21.09
GET 90% (h)	22.32	24.36	25.60	37.96	35.04
R ²	0.82	0.92	0.92	0.98	0.77
<i>Midgut</i>					
MFT (h)	12.99	21.01	12.52	19.45	19.99
MET 50% (h)	35.63	39.50	35.39	38.65	38.88
MET 75% (h)	40.72	43.65	35.39	42.97	43.12
MET 90% (h)	43.36	45.81	43.20	45.21	45.33
R ²	0.33	0.55	0.33	0.43	0.53
<i>Hindgut</i>					
HFT (h)	25.71	23.28	23.71	25.66	25.34
HET 50% (h)	41.60	41.00	40.07	41.54	41.20
HET 75% (h)	45.71	44.99	43.74	45.11	44.76
HET 90% (h)	47.03	47.06	45.66	46.97	46.62
R ²	0.71	0.42	0.65	0.70	0.60

No significant differences among treatments and within each gastrointestinal tract were detected $P \leq 0.05$

638

639

640

641

642

643

644

645

646

Table 6. Moisture, % of digesta obtained for each gastrointestinal tract of European sea bass from 1 to 48 hours (T1, T48) after feeding

<i>Stomach</i>	T1	T4	T10	T16	T28	T48	Time	Diet	Time*Diet
F7.2	63.0 ± 2.5 ^a	73.9 ± 4.3 ^{ab}	77.9 ± 5.0 ^b	82.9 ± 2.6 ^b	77.1 ± 1.2 ^b				
F8.9	58.9 ± 11.5 ^a	72.9 ± 3.4 ^b	77.2 ± 4.4 ^b	80.4 ± 1.4 ^b	80.0 ± 4.9 ^b				
F11.5	60.4 ± 6.8 ^a	69.3 ± 9.3 ^b	79.8 ± 5.0 ^b	80.5 ± 2.9 ^b	80.8 ± 6.1 ^b		0.0001	0.43	0.115
F13.1	59.4 ± 4.2 ^a	72.3 ± 3.0 ^b	75.9 ± 4.1 ^b	77.1 ± 5.1 ^b	78.8 ± 8.8 ^b				
F15.5	59.6 ± 3.0 ^a	74.6 ± 5.2 ^b	80.5 ± 2.2 ^b	78.4 ± 2.2 ^b	75.1 ± 6.3 ^b				
<i>Midgut</i>									
F7.2		86.1 ± 2.4	83.0 ± 7.1	85.4 ± 1.6	87.5 ± 1.5				
F8.9		87.0 ± 2.0	85.3 ± 1.1	85.9 ± 2.2	85.3 ± 2.8				
F11.5		85.4 ± 1.6	85.4 ± 2.6	83.2 ± 6.8	85.5 ± 4.7		0.608	0.647	0.769
F13.1		84.5 ± 5.7	85.5 ± 1.4	84.2 ± 3.9	84.2 ± 6.1				
F15.5		85.1 ± 2.5	85.1 ± 2.0	85.1 ± 3.8	84.9 ± 2.6				
<i>Hindgut</i>									
F7.2		80.2 ± 9.1	80.4 ± 10.1	82.5 ± 2.3	79.5 ± 5.9	80.4 ± 6.7			
F8.9		80.8 ± 3.4	81.9 ± 1.9	78.8 ± 10.4	81.6 ± 5.3	77.8 ± 5.1			
F11.5		77.1 ± 8.7	80.8 ± 7.2	80.3 ± 8.6	79.0 ± 7.2	71.3 ± 7.3	0.140	0.087	0.936
F13.1		84.0 ± 5.0	83.0 ± 2.2	79.7 ± 11.9	82.1 ± 4.5	80.6 ± 3.4			
F15.5		81.2 ± 5.8	83.0 ± 2.4	81.3 ± 6.0	81.6 ± 5.7	75.9 ± 11.1			

Values within a row with different superscripts differ significantly at $P < 0.05$. Statistical significance by two-way ANOVA using time and diet as independent factors.

650 **Figure captions**

651

652 **Figure 1.** Histology of the midgut (a, b) and hindgut (c, d) of European sea bass fed diets
653 F7.2 (a, c) and F15.5 (b, d). All the histological sections showed normal architecture of
654 the mucosal layer (regular columnar epithelium with polarized and basally located
655 nuclei), submucosal (loose connective tissue rich in capillary network) and muscular layer
656 (H&E, bar=100 μ m).

657

658 **Figure 2.** Exponential curves showing stomach evacuation of g digesta dry matter % body
659 weight (DM %BW) over the 48 h sampling period of European sea bass fed the
660 experimental diets. No significant differences among treatments were detected $P \leq 0.05$.


661

662 **Figure 3.** Midgut (a) and hindgut (b) quadratic curves describing the evacuation of g
663 digesta dry matter % body weight (DM %BW) over the 48 h sampling period of European
664 sea bass fed the experimental diets. No significant differences among treatments were
665 detected $P \leq 0.05$.

666

667

668 Figure 1




669

670


671

672 Figure 2
673
674
675



676
677
678
679

680 Figure 3



681

682

683

684

685

686

687


688

689

690

691

692



693