

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Generalized integration operators on hardy spaces

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Generalized integration operators on hardy spaces / Chalmoukis N.. - In: PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY. - ISSN 0002-9939. - ELETTRONICO. - 148:8(2020), pp. 3325-3337. [10.1090/proc/15016]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/802868> since: 2021-03-14

*Published:*

DOI: <http://doi.org/10.1090/proc/15016>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

**Chalmoukis, N. (2020). Generalized integration operators on hardy spaces. Proceedings of the American Mathematical Society, 148(8), 3325-3337.**

The final published version is available online at: <http://dx.doi.org/10.1090/proc/15016>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

# GENERALIZED INTEGRATION OPERATORS ON HARDY SPACES

NIKOLAOS CHALMOUKIS

ABSTRACT. We introduce a natural generalization of a well studied integration operator acting on the family of Hardy spaces in the unit disc. We study the boundedness and compactness properties of the operator and finally we use these results to give simple proofs of a result of Rättyä and another result by Cohn.

## 1. INTRODUCTION

Let  $H^p$  be the Hardy space of analytic function in the unit disc, i.e. functions  $f$  holomorphic in  $\mathbb{D}$  such that

$$\|f\|_p^p := \frac{1}{2\pi} \sup_{0 < r < 1} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta < \infty.$$

In this article  $p$  will be a positive exponent. There is a variety of linear operators acting between Hardy spaces which have been studied. In particular in [1], motivated by some earlier work of Pommerenke in [2] and Berkson and Porta in [3] Aleman and Siskakis introduced the integral operator  $T_g$ , depending on a symbol  $g$  analytic in the unit disc, which generalized the classical Cesàro averaging operator on  $H^2$ , defined by

$$T_g f(z) = \int_0^z f(t)g'(t)dt.$$

The operator has been studied extensively, both for its intrinsic interest and also for its applications. In particular, such an operator seems to have been introduced by Pommerenke in [2], in order to give a slick proof of the analytic John-Nirenberg inequality. Later on it became also evident the connection to factorization theorems for derivatives of functions in the Hardy space (see [4]), and therefore to previous work of Aleksandrov and Peller on Foguel-Hankel operators in [5]. More precisely suppose we consider the bilinear operator  $T(f, g) := T_g f$ ,  $T : X \times Y \rightarrow Z$ , where  $X, Y, Z$  are Banach spaces of analytic functions on the unit disc. Then the question whether or not, for every  $h \in Z$  there exists a factorization of the form  $h' = fg'$ , where  $f \in X, g \in Y$  and  $\|f\|_X + \|g\|_Y \lesssim \|h\|_Z$ , can be translated into a question about the openness of the bilinear operator  $T$ . Weak factorization theorems can be also approached in this way (see [4]).

The first attempt to characterize the boundedness properties of  $T_g$  was done in [1] by Aleman and Siskakis where they provided necessary and sufficient conditions for the boundedness of  $T_g$  on  $H^p$  for  $p \geq 1$ , while in [6] the result was extended for  $p > 0$ . More recently, a characterization of its spectrum was found in [7].

---

2010 *Mathematics Subject Classification*. Primary: 30H10. Secondary: 30H35, 47G10.

The first author is supported by the fellowship INDAM-DP-COFUND-2015 "INDAM Doctoral Programme in Mathematics and/or Applications Cofunded by Marie Skłodowska-Curie Actions", Grant 713485.

In the present work we study a generalization of the aforementioned operator. Let us introduce the notation  $I$  for the integration operator, i.e.

$$I f(z) := \int_0^z f(t) dt.$$

Fix now an analytic symbol  $g$  and a sequence of coefficients  $a \in \mathbb{C}^{n-1}$ . We shall define the generalized integration operator  $T_{g,a}$  by

$$T_{g,a} f(z) = I^n(fg^{(n)} + a_1 f' g^{(n-1)} \dots a_{n-1} f^{(n-1)} g')(z).$$

The question which naturally arises is, for which symbols  $g$  this operator defines a bounded linear operator between two Hardy spaces. In this direction we prove the following theorems.

**Theorem 1.1.** *Let  $0 < p < \infty$  and  $a \in \mathbb{C}^{n-1}$ . Then,  $T_{g,a}$  is bounded from  $H^p$  to itself if and only if  $g \in BMOA$ . Moreover,  $T_{g,a}$  is compact if and only if  $g \in VMOA$ .*

In the next theorem we use the notation  $\Lambda_\alpha$  to denote the analytic Lipschitz space. That is the space of holomorphic functions  $f$  in the unit disc such that

$$|f'(z)| = \mathcal{O}((1 - |z|)^{1-\alpha}).$$

The little Lipschitz space  $\lambda_\alpha$  corresponds to the "little  $o$ " condition in the above definition.

**Theorem 1.2.** *Let  $0 < p < q < \infty$  and  $a \in \mathbb{C}^{n-1}$ . Define also  $\alpha = \frac{1}{p} - \frac{1}{q} > 0$  and  $k = \max\{l : a_l \neq 0\}$  or  $k = 0$  if  $a = 0$ . Then if  $l < \alpha \leq l + 1 \leq n - k$  for some  $l \in \mathbb{N}$ ,  $T_{g,a}$  is bounded from  $H^p$  to  $H^q$  if and only if  $g^{(l)} \in \Lambda_{\alpha-l}$ . If  $\alpha > n - k$  and  $T_{g,a}$  is bounded from  $H^p$  to  $H^q$  then  $T_{g,a}$  is the zero operator.*

**Theorem 1.3.** *Let  $0 < q < p < \infty$  and  $a \in \mathbb{C}^{n-1}$ . If  $g \in H^s$ , where  $\frac{1}{s} = \frac{1}{q} - \frac{1}{p}$  then  $T_{g,a}$  is bounded from  $H^p$  to  $H^q$ . In the special case that  $n = 2$  and  $a = 0$ , if  $T_{g,a} : H^p \rightarrow H^q$  is bounded, then  $g \in H^s$ .*

It is natural to state the following conjecture.

**Conjecture 1.4.** *Is it true that if  $T_{g,a} : H^p \rightarrow H^q, 0 < q < p < \infty$  is bounded then  $g$  must be in  $H^s, \frac{1}{s} = \frac{1}{q} - \frac{1}{p}$ ?*

Theorems 1.1 and 1.3 show that the behaviour, regarding boundedness, of the operator  $T_{g,a}$ , in the known cases, is essentially the same with this of  $T_g$  when  $p \geq q$ . In the case that  $p < q$  the operator exhibits a different behaviour because its boundedness depends on its last non zero term. It is interesting to notice that in all known cases, if  $T_{g,a}$  is bounded, every term comprising the operator is forced to be bounded as well. In other words there is no cancellation between the terms.

In the light of the manifold applications that the generalized Cesàro operator has in function theory of the Hardy space, it is natural to investigate potential applications of the generalized integration operator. Not surprisingly, using  $T_{g,a}$  one can derive a result which generalizes the analytic John-Nirenberg inequality and a second theorem which generalizes factorization of derivatives in the Hardy space. Both results exist already in the literature but our method allows for a unified and simpler approach.

The first one is of Rättyä [8] on solutions of complex linear differential equations. To formulate it we shall use the notation  $\|\cdot\|_*$  for the semi-norm of the  $BMOA$  space, i.e.

$$\|f\|_* := \sup_{a \in \mathbb{D}} \|f \circ \varphi_a - f(a)\|_{H^2},$$

where  $\varphi_a$  is the biholomorphic involution of the unit disc which maps 0 to  $a$ .

**Theorem 1.5.** *Let  $n \in \mathbb{N}, 0 < p < \infty, f_0 \in H^p, G \in BMOA$  and  $g_i \in \mathcal{B}, 1 \leq i < n$ . There exists a constant  $A > 0$  depending on  $p$  such that if  $\|G\|_*, \|g_i\|_{\mathcal{B}} < A$ , every solution of the non homogeneous linear differential equation*

$$G^{(n)}f + g_1^{(n-1)}f' + \cdots + g_{n-1}'f^{(n-1)} + f^{(n)} = f_0^{(n)}$$

*is in  $H^p$ . If  $G \in VMOA$  and  $g_i \in \mathcal{B}_0$ , the same result holds without the restriction in the norm of  $g_i$  and  $G$ .*

*Proof.* To see why this is a Corollary of Theorem 1.1 define the operator

$$Df = \Gamma^n(fG^{(n)} + f'g_1^{(n-1)} + \cdots + f^{(n-1)}g_{n-1}')_0$$

which by Theorem 1.1 is bounded on  $H^p$ , as a sum of  $n-1$  bounded operators. By an application of the closed graph theorem, there exists a constant  $C$  depending only on  $p$  such that

$$\|D\|_p \leq C(\|G\|_* + \|g_1\|_{\mathcal{B}} + \cdots + \|g_{n-1}\|_{\mathcal{B}}).$$

If  $\|G\|_*, \|g_i\|_{\mathcal{B}} < 1/(nC)$ , then by triangular inequality  $\|D\| < 1$ , and hence since the spectrum of a bounded operator is contained in the disc centred at 0 with radius the norm of the operator, it follows that  $-1$  is in the resolvent set of  $D$  on  $H^p$ . Let now  $f$  any solution of the above differential equation. There exists  $F \in H^p$  such that  $F - f_0$  is a polynomial of degree less than  $n$  and  $F^{(i)}(0) = f^{(i)}(0), 0 \leq i < n$ . Then for some  $\tilde{f} \in H^p, D\tilde{f} + \tilde{f} = F$ . But  $\tilde{f}$  and  $f$  satisfy the same differential equation with the same initial conditions, hence by the uniqueness of solution of the initial value problem  $f = \tilde{f} \in H^p$ .

If  $G \in VMOA$  and  $g_i \in \mathcal{B}_0$ ,  $D$  is compact. We prove that in this case its spectrum is the singleton  $\{0\}$ , thus the result follows by the previous argument. Since the operator is compact it suffices to prove that there are no non zero eigenvalues. In fact if  $\lambda \in \mathbb{C} \setminus \{0\}$ , the equation  $Df = \lambda f$  is equivalent to the initial value problem

$$fG^{(n)} + f'g_1^{(n-1)} + \cdots + f^{(n-1)}g_{n-1}' = \lambda f^{(n)}, \quad f(0) = \cdots = f^{(n-1)}(0) = 0,$$

which again by the uniqueness of solutions implies that  $f \equiv 0$ . □

The second result which follows from our analysis is a factorization theorem for derivatives of functions in the Hardy space. For  $n = 1$  the result can be found in [4], and it was later generalized by Cohn for derivatives of any order in [9].

**Theorem 1.6.** *Let  $f \in H^p$ , then there exist  $F \in H^p$  and  $G_n \in BMOA, n \in \mathbb{N}$  such that  $f^{(n)} = FG_n^{(n)}$ .*

The paper is organized as follows. In Section 2 we give some preliminary results in Hardy spaces which we shall use repeatedly later. In Section 3 we give the proofs of the main theorems.

**Notation.** As it is customary we shall use the notation  $A \lesssim B$ , when there exists some constant  $C > 0$  independent of the parameters on which  $A, B$  depend such that  $A \leq CB$ . If  $A \lesssim B$  and  $B \lesssim A$  we write  $A \approx B$ .

**Acknowledgements.** Most of the present work was done while the author was a MSc student at Lund University under the supervision of Professor Alexandru Aleman. I would like to express my gratitude to him for introducing me to the problems discussed in this article, and for his guidance and help.

## 2. PRELIMINARY RESULTS

First we will look at some variations on the Hardy-Stein identity. To do so we shall introduce the Stolz angle  $\Gamma_\sigma(e^{i\theta})$  with vertex at  $e^{i\theta}$  and aperture  $\sigma$ . That is, the interior of the convex hull of the point  $e^{i\theta}$  and the disc  $D(0, \sigma)$ . Then we can define the so called square function or Lusin area function  $S_f$ .

$$S_f(\zeta) = \left( \int_{\Gamma_\sigma(\zeta)} |f'|^2 dA \right)^{1/2}, \zeta \in \mathbb{T}.$$

A well known result of Fefferman and Stein (see [10]) states that if  $0 < p < \infty$  there exist constants  $C_1, C_2 > 0$ , depending only on  $\sigma$  and  $p$ , such that

$$C_1 \|f\|_p^p \leq |f(0)|^p + \int_{\mathbb{T}} S_f^p dm \leq C_2 \|f\|_p^p,$$

for any  $f$  analytic in  $\mathbb{D}$ . Another related function is the Paley-Littlewood  $G$ -function, which is defined by

$$G(f)(\zeta) = \left( \int_0^1 |f'(r\zeta)|^2 (1-r) dr \right)^{1/2}, \quad \zeta \in \mathbb{T}.$$

The Paley-Littlewood  $G$ -function enjoys the same property as the Lusin area function, i.e. there exist  $C_1, C_2 > 0$ , depending only on  $\sigma$  and  $p$ , such that

$$(1) \quad C_1 \|f\|_p^p \leq |f(0)|^p + \int_{\mathbb{T}} G^p(f) dm \leq C_2 \|f\|_p^p,$$

for any  $f$  analytic in  $\mathbb{D}$ . For more information about these functions the reader is referred to [10].

For our purposes, we need a version of the Paley-Littlewood  $G$ -function, involving only the  $n$ -th derivative of  $f$ . We start with a lemma.

**Lemma 2.1.** *Let  $f$  be an analytic function in the unit disc, and  $n \geq 1$  then for  $z \in \mathbb{D}$*

$$|f^{(n)}(z)|^2 \leq \frac{n!(n-1)!2^{2n}}{(1-|z|)^{2n}} \int_{D(z, \frac{1-|z|}{2})} |f'(\zeta)|^2 dA(\zeta).$$

*Proof.* The expression of the Dirichlet integral of  $f$  in terms of its Taylor coefficients is

$$\int_{\mathbb{D}} |f'(\zeta)|^2 dA(\zeta) = \sum_{k \geq 0} k |a_k|^2.$$

Using this we have,

$$(2) \quad |f^{(n)}(0)|^2 = (n!)^2 |a_n|^2 \leq n!(n-1)! \sum_{k \geq 0} k |a_k|^2 = n!(n-1)! \int_{\mathbb{D}} |f'(\zeta)|^2 dA(\zeta).$$

Now let  $z \in \mathbb{D}$  fixed and set  $r = \frac{1-|z|}{2}$ . Applying (2) to the function  $f_z(\zeta) = f(z + r\zeta)$ ,  $\zeta \in \mathbb{D}$ , and using a change of variables:

$$\begin{aligned} r^{2n} |f^{(n)}(z)|^2 &\leq n!(n-1)! \int_{\mathbb{D}} r^2 |f'(z + r\zeta)|^2 dA(\zeta) \\ &= n!(n-1)! \int_{D(z, \frac{1-|z|}{2})} |f'(\zeta)|^2 dA(\zeta), \end{aligned}$$

which gives the desired inequality.  $\square$

Now, let  $f$  analytic in  $\mathbb{D}$ . We define the Paley-Littlewood  $G_k$ -function of order  $k$  to be

$$(3) \quad G_k(f)(\zeta) = \left( \int_0^1 |f^{(k)}(r\zeta)|^2 (1-r)^{2k-1} dr \right)^{1/2}.$$

**Proposition 2.2.** *Let  $p > 0$  and  $f$  analytic in  $\mathbb{D}$  with  $f^{(i)}(0) = 0$ ,  $0 \leq i < k$  then, for any  $k \in \mathbb{Z}_+$ , there exist constants  $C_1, C_2$  depending only on  $p$  and  $k$  such that*

$$C_1 \|f\|_{H^p} \leq \|G_k(f)\|_{L^p} \leq C_2 \|f\|_{H^p}.$$

*Proof.* We prove the left inequality for  $k = 2$ , the general case follows by induction. First, let  $f$  be analytic in an open set containing the closure of  $\mathbb{D}$ . Then for  $t \in \mathbb{R}$  fixed

$$\begin{aligned} G_1^2(f)(e^{it}) &= \int_0^1 |f'(re^{it})|^2 (1-r) dr = \frac{1}{2} \int_0^1 \frac{\partial |f'(re^{it})|^2}{\partial r} (1-r)^2 dr \\ &= \int_0^1 \operatorname{Re}(e^{it} f''(re^{it}) \bar{f}'(re^{it})) (1-r)^2 dr \\ &\leq \int_0^1 |f''(re^{it})| |f'(re^{it})| (1-r)^2 dr \\ &\leq G_1(f)(e^{it}) G_2(f)(e^{it}), \end{aligned}$$

by Cauchy-Schwarz. Dividing through by  $G_1(f)(e^{it})$  we have the result for  $f$  analytic in a larger disc. Now if  $f$  is an arbitrary analytic function, fix  $0 < \rho < 1$  and consider the dilations  $f_\rho(z) = f(\rho z)$ . Then,

$$\begin{aligned} G_1(f_\rho)(t) &\leq \rho^4 \int_0^1 |f''(\rho e^{it})|^2 (1-r)^3 dr \\ &\leq \rho \int_0^\rho |f''(ue^{it})|^2 (1-u)^3 du. \end{aligned}$$

Then by taking  $\liminf$  in both sides as  $\rho \rightarrow 1^-$  and applying Fatou's lemma on the left and monotone convergence on the right we conclude that  $G_1(f) \leq G_2(f)$ . The desired inequality then follows by (1).

To prove the right inequality, we will use Lemma 2.1. First note that for any  $z = re^{it} \in \mathbb{D}$ ,  $D(re^{it}, \frac{1-r}{2}) \subset \Gamma_{\frac{1}{2}}(e^{it})$ , which, together with Lemma 2.1, justifies the following calculation

$$\begin{aligned} G_k^2(f)(e^{it}) &= \int_0^1 |f^{(k)}(re^{it})|^2 (1-r)^{2k-1} dr \\ &\leq C_k \int_0^1 \int_{D(re^{it}, \frac{1-r}{2})} |f'(\zeta)|^2 dA(\zeta) (1-r)^{-1} dr \\ &= C_k \int_{\Gamma_{\frac{1}{2}}(e^{it})} |f'(\zeta)|^2 \int_0^1 \chi_{D(r, \frac{1-r}{2})}(\zeta) (1-r)^{-1} dr dA(\zeta). \end{aligned}$$

It is routine to check that if  $|\zeta| < \frac{3r-1}{2}$  or  $|\zeta| > \frac{1+r}{2}$  then  $\zeta \notin D(r, \frac{1-r}{2})$ . Hence

$$\int_0^1 \chi_{D(r, \frac{1-r}{2})}(\zeta) (1-r)^{-1} dr \leq \int_{\frac{2|\zeta|-1}{3}}^{\frac{2|\zeta|+1}{3}} (1-r)^{-1} dr = \log 3$$

Therefore we have proven that  $G_k^2(f)(e^{it}) \leq S_f^2(e^{it})$  and the estimate follows by Fefferman-Stein's theorem.  $\square$

We conclude this section with a lemma of linear algebra. For  $\gamma > 0$  we use the notation  $(\gamma)_0 = 1, (\gamma)_k = \gamma(\gamma+1)\cdots(\gamma+k-1)$  for  $k \geq 1$ .

**Lemma 2.3.** *Suppose that  $f_0, f_2, \dots, f_{n-1}$  are complex valued functions on the unit disc (not necessarily analytic), such that for any  $\gamma \in \mathbb{R}$  sufficiently large there exists  $C_\gamma > 0$  such that*

$$(4) \quad \left| \sum_{k=0}^{n-1} f_k(z) (\gamma)_k \right| \leq C_\gamma, z \in \mathbb{D}.$$

Then all  $f_k$  are bounded.

*Proof.* Choose distinct  $\gamma_0, \gamma_2, \dots, \gamma_{n-1}$  sufficiently large. It is a tedious but standard calculation that

$$\det \begin{bmatrix} (\gamma_0)_1 & (\gamma_0)_2 & \cdots & (\gamma_0)_{n-1} \\ (\gamma_1)_1 & (\gamma_1)_2 & \cdots & (\gamma_1)_{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ (\gamma_{n-1})_1 & (\gamma_{n-1})_2 & \cdots & (\gamma_{n-1})_{n-1} \end{bmatrix} = \prod_{0 \leq i < j < n} (\gamma_j - \gamma_i) \neq 0.$$

In other words, the vectors  $\Gamma_i = ((\gamma_i)_0, \dots, (\gamma_i)_{n-1}), k = 0, 1, \dots, n-1$  form a basis of  $\mathbb{R}^n$ . Therefore for a fixed  $k, 0 \leq k < n$ , there exist  $r_0(k), \dots, r_{n-1}(k) \in \mathbb{R}$ , such that

$$\sum_{i=0}^{n-1} r_i(k) \Gamma_i = e_k,$$

where  $e_k$  is the  $k$ -th vector of the standard basis of  $\mathbb{R}^n$ . In other words,

$$\sum_{i=0}^{n-1} r_i(k) (\gamma_i)_j = \delta_{kj}, \quad 0 \leq k, j < n,$$



where  $\delta$  is the Kronecker delta. Therefore,

$$\begin{aligned} f_k(z) &= \sum_{j=0}^{n-1} f_j(z) \left( \sum_{i=0}^{n-1} r_i (\gamma_i)_j \right) \\ &= \sum_{i=0}^{n-1} r_i \sum_{j=0}^{n-1} f_j(z) (\gamma_i)_j. \end{aligned}$$

Hence, by the our assumptions,

$$|f_k(z)| \leq \sum_{i=0}^{n-1} |r_i| C_{\gamma_i}.$$

□

### 3. PROOFS OF THE MAIN THEOREMS

In order to understand the behaviour of the operator  $T_{g,a}$  it will be useful to consider the operators  $T_g^{n,k}$  defined for an analytic function  $g$  and natural numbers  $n, k$  such that  $0 \leq k < n$ , by the formula

$$T_g^{n,k} f = I^n(f^{(k)} g^{(n-k)}).$$

#### 3.1. Boundedness.

**Proof of Theorem 1.1.** The main step to prove the sufficiency part of Theorem 1.1, is the following proposition.

**Proposition 3.1.** *Let  $n, k \in \mathbb{Z}_+$ ,  $k < n$ , and  $g \in \mathcal{B}$ . Then the operator  $T_g^{n,k} : H^p \rightarrow H^p$  is bounded.*

*Proof.* Let  $f$  be an analytic function in an open set containing the closure of the unit disc. We can assume without loss of generality that  $f^{(i)}(0) = 0$ ,  $0 \leq i < k$  because it is readily checked that  $T_g^{n,k}$  maps the set of polynomials in  $H^p$ . Then we have that

$$\begin{aligned} G_n(T_g^{n,k} f)(e^{it}) &= \left( \int_0^1 |f^{(k)}(re^{it})|^2 |g^{(n-k)}(re^{it})|^2 (1-r)^{2n-1} dr \right)^{1/2} \\ &\lesssim \|g\|_{\mathcal{B}} \left( \int_0^1 |f^{(k)}(re^{it})|^2 (1-r)^{2k-1} dr \right)^{1/2} \\ &\lesssim \|g\|_{\mathcal{B}} G_k(f)(e^{it}). \end{aligned}$$

The result follows immediately from Proposition 2.2, and the fact that polynomials are dense in  $H^p$ . □

To prove sufficiency in Theorem 1.1, notice that

$$(5) \quad T_g = T_g^{n,0} + \sum_{k=1}^{n-1} \binom{n-1}{k} T_g^{n,k}.$$

Therefore if  $g \in BMOA \subset \mathcal{B}$ , by Proposition 3.1 and the fact that  $T_g$  is bounded when  $g \in BMOA$  (see [6]),  $T_g^{n,0}$  is bounded as well. Hence  $T_{g,a}$  is bounded as well.

We suppose that  $T_{g,a} : H^p \rightarrow H^p$  is bounded. It suffices to prove that in that case  $g \in \mathcal{B}$ . this together with Proposition 3.1 and equation (5) imply that  $T_g$  is bounded, and therefore  $g \in BMOA$ , because the result is known to hold for  $n = 1$  (see [6]).

For  $\lambda \in \mathbb{D}, \gamma > 1/p$  set

$$f_{\lambda,\gamma}(z) = \frac{(1 - |\lambda|^2)^{\gamma-1/p}}{(1 - \bar{\lambda}z)^\gamma}.$$

Hence,

$$(6) \quad f_{\lambda,\gamma}^{(k)}(\lambda) = \frac{(\gamma)_k \bar{\lambda}^k}{(1 - |\lambda|^2)^{k+1/p}}, \quad k \in \mathbb{N},$$

where  $(\gamma)_0 = 1, (\gamma)_k = \gamma(\gamma+1) \cdots (\gamma+k-1)$ . Also there exists a positive constant  $C_\gamma$  such that  $\|f_{\lambda,\gamma}\|_p \leq C_\gamma, \lambda \in \mathbb{D}$ . Then the growth estimate for  $H^p$  functions gives

$$\begin{aligned} \frac{C_\gamma}{(1 - |\lambda|^2)^{n+1/p}} &\geq |(T_{g,a} f_{\lambda,\gamma})^{(n)}(\lambda)| \\ &= \left| \sum_{k=0}^{n-1} \frac{a_k \bar{\lambda}^k (\gamma)_k}{(1 - |\lambda|^2)^{k+1/p}} g^{(n-k)}(\lambda) \right|. \end{aligned}$$

Or, rearranging the inequality,

$$(7) \quad \left| \sum_{k=0}^{n-1} g^{(n-k)}(\lambda) (1 - |\lambda|^2)^{(n-k)} a_k \bar{\lambda}^k (\gamma)_k \right| \leq C_\gamma.$$

By applying Lemma 2.3 we can infer that  $\sup_{\lambda \in \mathbb{D}} |g^{(n)}(\lambda)| (1 - |\lambda|^2)^n < \infty$ , i.e.  $g \in \mathcal{B}$ .

At this point we are able to give the promised simple proof of Theorem 1.6. We shall need the next lemma which is of some interesting on its own right.

**Lemma 3.2.** *If  $g$  is in  $BMOA$  the operator  $T_g^{n,n-1}$  is a bounded linear operator from  $\mathcal{B}$  to  $BMOA$ .*

*Proof.* Let  $f \in \mathcal{B}$  and set  $F = T_g^{n,n-1} f$ . Then for an arbitrary  $h \in H^2$

$$T_{F,0} h = I^n (f^{(n-1)} g' h) = T_f^{n,1} T_g h.$$

But  $T_g h \in H^2$  since  $g \in BMOA$  and by Proposition 3.1,  $T_f^{n,1}$  is bounded on  $H^2$ . Hence  $T_{F,0}$  is bounded on  $H^2$  as well. By the necessity part of Theorem 1.1 we get that  $F \in BMOA$ .  $\square$

*Proof of Theorem 1.6.* Let  $f \in H^p$ , and define  $F$  by

$$F(z) = \left( \int_{\mathbb{T}} \frac{\zeta + z}{\zeta - z} |f(\zeta)|^{p/2} dm(\zeta) \right)^{2/p}.$$

Since  $F$  has positive real part,  $\log F \in BMOA$ . Now we proceed by induction. The statement is trivial for  $n = 0$ , hence suppose that it is true for  $n - 1$ ,

$$\begin{aligned} f^{(n)} &= (F G_{n-1}^{(n-1)})' \\ &= F((\log F)' G_{n-1}^{(n-1)} + G_{n-1}^{(n)}) \\ &= F(T_{\log F}^{n,n-1} G_{n-1} + G_{n-1})^{(n)}. \end{aligned}$$

But  $\log F \in BMOA \subset \mathcal{B}$ , hence, by the previous lemma  $T_{\log F}^{n,n-1} G_{n-1} \in BMOA$ .  $\square$

**Proof of Theorem 1.2.** As before we prove a seemingly stronger statement.

**Proposition 3.3.** *Let  $0 < p < q < \infty, n \in \mathbb{N}$  and  $0 \leq k < n$  fixed. Then set  $\alpha = \frac{1}{p} - \frac{1}{q}$ . If  $l < \alpha \leq l+1 \leq n-k$  for some  $l \in \mathbb{N}$  and  $g^{(l)} \in \Lambda_{\alpha-l}$ , then  $T_g^{n,k}$  is bounded from  $H^p$  to  $H^q$ .*

*Proof.* For  $n = 1$  the operator is just  $T_g$ , and the statement of the theorem reduces to the known result about  $T_g$  (see [6]). Suppose now that the statement is true for some  $n \geq 1$  and proceed by induction. We distinguish two cases. If  $n-k < \alpha \leq n+1-k$ , take an arbitrary  $f \in H^p$  and write  $f^{(k)} = FG_k^{(k)}$  for some  $F \in H^p$  and  $G_k \in BMOA$ . The following identity is true,

$$T_g^{n+1,k}f = I^{(n-k)}T_g^{k+1,1}T_{g^{(n-k)}}F.$$

By the assumption that  $g^{(n-k)} \in \Lambda_{\alpha-n+k}$  and by the case  $n = 1$  we conclude that  $T_{g^{(n-k)}}$  is bounded from  $H^p$  to  $H^{p'}$ , where  $\frac{1}{p'} = \frac{1}{q} + n - k$ . It follows, using Proposition 3.1 and the boundedness properties of  $I$  that in this case  $T_{g,a} : H^p \rightarrow H^q$  is bounded. Suppose now that  $\alpha \leq n - k$ . Assuming without loss of generality that  $g$  has sufficient zero multiplicity at the origin. Integrating by parts we get

$$T_g^{n+1,k}f = T_g^{n,k}f - T_g^{n+1,k+1}f.$$

The first term on the right hand side is bounded by the induction hypothesis. To prove the boundedness of the second term, factorize  $f^{(k+1)}$  as in Theorem 1.6. Then,

$$T_g^{n+1,k+1}f = T_{G_k}^{n+1,n-k}T_g^{n-k,0}F.$$

The result follows by the induction hypothesis and Theorem 1.1 applied to the operator  $T_{G_k}^{n+1,n-k}$ .  $\square$

The proof of necessity in Theorem 1.2 is similar to that of the necessity part of Theorem 1.1. Consider again the family of test functions  $f_{\lambda,\gamma}$ . As before,

$$\left| \sum_{k=0}^{n-1} \frac{a_k \bar{\lambda}^k (\gamma)_k}{(1-|\lambda|^2)^k} g^{(n-k)}(\lambda) \right| \leq \frac{C_\gamma}{(1-|\lambda|^2)^{\frac{1}{q} - \frac{1}{p} + n}},$$

by applying again Lemma 2.3 we can separate the previous condition to the following.

$$(8) \quad |g^{(n-k)}(\lambda)| \lesssim \frac{1}{(1-|\lambda|^2)^{\frac{1}{q} - \frac{1}{p} + n - k}}, \quad \lambda \in \mathbb{D},$$

where  $k = \max\{l : a_l \neq 0\}$ . Since  $a_k \neq 0$  it follows immediately that  $g^{(l)} \in \Lambda_{\alpha-l}$  if  $\alpha \leq n - k$  and  $g^{(n-k)} = 0$  if  $\alpha > n - k$ .

**Proof Theorem 1.3.** The proof of sufficiency in Theorem 1.3 follows the same pattern. Again we will prove by induction on  $n$  that for  $0 \leq k < n$ ,  $T_g^{n,k}$  is bounded from  $H^p$  to  $H^q$ . As usual write  $f^{(k)} = FG_k^{(k)}$  as in Theorem 1.6 and use the same recursive formula, i.e.,

$$T_g^{n+1,k}f = T_g^{n,k}f - T_{G_k}^{n+1,n-k}T_g^{n-k,0}F.$$

By induction hypothesis  $T_g^{n,k}f, T_g^{n-k,0}F \in H^q$ . Furthermore,  $G_k \in BMOA$  thus by Theorem 1.1  $T_{G_k}^{n+1,n-k}T_g^{n-k,0}F \in H^q$ .

The proof of necessity in Theorem 1.3 is more involved and is based on the following lemma.

**Lemma 3.4.** *Suppose that  $n = 2, a \in \mathbb{C}$ . For every  $p > q > 0$  there exists a positive constant  $C = C(q) > q$  such that if  $p > C(q)$  and  $T_{g,a} : H^p \rightarrow H^q$  then  $g \in H^s, \frac{1}{s} = \frac{1}{q} - \frac{1}{p}$ .*

*Proof.* Assume that  $g(0) = g'(0) = 0$ , multiple integrations by parts allow us to rewrite the formula expressing  $T_{g,a}$  as

$$(9) \quad T_{g,a}f = fg + (1-a) \int \int f''g + (a-2) \int f'g$$

$$(10) \quad = fg + (1-a) \int \int \left(\frac{f''}{f} - \frac{f'^2}{f^2}\right)gf + (1-a) \int \int \frac{f'^2}{f^2}gf + (a-2) \int \frac{f'}{f}gf,$$

for any  $f \in H^p$ , where we used the convention

$$\int f = \int_0^z f(t)dt.$$

Now let  $\varepsilon > 0, 0 < r < 1$ . By  $g_r$  we denote the dilation of  $g$ . Define  $G = \mathcal{H}\left(\frac{|g_r|^\alpha}{(1+\varepsilon|g_r|)^\alpha}\right)^\beta$ , where  $\alpha$  and  $\beta$  are positive constants to be specified later and  $\mathcal{H}$  denotes the Herglotz transform, i.e.

$$G(z) = \left( \int_{\mathbb{T}} \frac{|g(r\zeta)|^\alpha}{(1+\varepsilon|g(r\zeta)|)^\alpha} \frac{\zeta+z}{\zeta-z} dm(\zeta) \right)^\beta.$$

$G^{\frac{1}{\beta}}$  has positive real part, hence  $\|\log G\|_* \lesssim \beta$ , that follows by the definition of the  $BMOA$  norm and Littlewood's subordination principle. Because, for  $a \in \mathbb{D}$  the function  $\log \frac{G^{\frac{1}{\beta}} \circ \varphi_a}{G^{\frac{1}{\beta}}(a)}$  is subordinate to  $\log \frac{1+z}{1-z}$ . Hence,

$$\|\log G\|_* = \beta \sup_{a \in \mathbb{D}} \left\| \log \frac{G^{\frac{1}{\beta}} \circ \varphi_a}{G^{\frac{1}{\beta}}(a)} \right\|_{H^2} \leq \left\| \log \frac{1+z}{1-z} \right\|_{H^2} < +\infty.$$

Also  $G \in BMOA$ , since the Herglotz transform maps  $C^\infty(\overline{\mathbb{D}}) \cap Hol(\mathbb{D})$  into bounded analytic functions. If we assume that  $\beta p > 1$  then by M. Riesz's theorem [11, Theorem 4.1]

$$(11) \quad \|G\|_p^p = \int_{\mathbb{T}} \mathcal{H}\left(\frac{|g_r|^\alpha}{(1+\varepsilon|g_r|)^\alpha}\right)^{\beta p} dm$$

$$(12) \quad \lesssim \int_{\mathbb{T}} \left(\frac{|g_r|}{1+\varepsilon|g_r|}\right)^{\alpha\beta p} dm.$$

The last estimate we need for  $G$  is that  $|G| \geq (\operatorname{Re} G^{1/\beta})^\beta = \left(\frac{|g_r|}{1+\varepsilon|g_r|}\right)^{\alpha\beta}$  on  $\mathbb{T}$ . Consequently,

$$(13) \quad \|Gg\|_q^q = \int_{\mathbb{T}} |Gg|^q dm \geq \int_{\mathbb{T}} \frac{|g_r|^{\alpha\beta q} |g|^q}{(1+\varepsilon|g_r|)^{\alpha\beta q}} dm.$$

With these preliminaries we are going to estimate  $\|T_{g,a}G\|_q$ . First we estimate separately the last three terms in the right hand side of (9).

$$(14) \quad \left\| \int \int (\log G)'' Gg \right\|_q \leq C_1(q) \|\log G\|_* \|Gg\|_q$$

$$(15) \quad \leq C_2(q) \beta \|Gg\|_q,$$

by the sufficiency part of Theorem 1.1. Similarly,

$$\begin{aligned} \left\| \int \int \frac{G'^2}{G^2} gG \right\|_q &= \left\| \int \int (\log G)' \left( \int (\log G)' gG \right)' \right\|_q \\ &\leq C_3(q) \beta^2 \|gG\|_q. \end{aligned}$$

$$\left\| \int \frac{G'}{G} gG \right\|_q \leq C_4(q) \beta \|gG\|_q.$$

Then (9) and the boundedness of  $T_{g,a}$  imply that

$$\|Gg\|_q (1 - C_5(q)\beta - C_6(q)\beta^2) \lesssim \|T_{g,a}\|_{p,q} \|G\|_p.$$

Furthermore, if  $\beta C_5(q) + \beta^2 C_6(q) < 1$  (or equivalently  $\beta < C(q)$  where  $C(q)$  is a continuous function of  $C_5, C_6$ ) we arrive at

$$\|Gg\|_q \leq C_{q,\beta} \|T_{g,a}\|_{p,q} \|G\|_p,$$

which together with estimates (12) and (13) give

$$\left( \int_{\mathbb{T}} \frac{|g_r|^{\alpha\beta q} |g|^q}{(1 + \varepsilon |g_r|)^{\alpha\beta q}} dm \right)^{1/q} \leq \tilde{C}_{q,\beta} \|T_{g,a}\|_{p,q} \left( \int_{\mathbb{T}} \left( \frac{|g_r|}{1 + \varepsilon |g_r|} \right)^{\alpha\beta p} \right)^{1/p}.$$

Letting  $r \rightarrow 1^-$  by Fatou's Lemma and Bounded Convergence Theorem we arrive at

$$\left( \int_{\mathbb{T}} \frac{|g|^{(\alpha\beta+1)q}}{(1 + \varepsilon |g|)^{\alpha\beta q}} dm \right)^{1/q} \leq \tilde{C}_{q,\beta} \|T_{g,a}\|_{p,q} \left( \int_{\mathbb{T}} \left( \frac{|g|}{1 + \varepsilon |g|} \right)^{\alpha\beta p} \right)^{1/p}.$$

Now, choose  $\alpha$  such that  $(\alpha\beta+1)q = \alpha\beta p$  and note that in this case  $(\alpha\beta+1)q = s = \alpha\beta p$ . Hence

$$\left( \int_{\mathbb{T}} \left( \frac{|g|}{1 + \varepsilon |g|} \right)^s dm \right)^{1/q} \leq \left( \int_{\mathbb{T}} \frac{|g|^s}{(1 + \varepsilon |g|)^{\alpha\beta q}} dm \right)^{1/q} \leq \widetilde{C}_{q,\beta} \|T_{g,a}\|_{p,q} \left( \int_{\mathbb{T}} \left( \frac{|g|}{1 + \varepsilon |g|} \right)^s \right)^{1/p}.$$

Finally letting  $\varepsilon \rightarrow 1^-$  Fatou's lemma then gives  $g \in H^s$ .  $\square$

We can now proceed to the proof of the necessity part in Theorem 1.3. Let  $T_{g,0} : H^p \rightarrow H^q, p > q$  be bounded. And set  $\frac{1}{s} = \frac{1}{q} - \frac{1}{p}$ . Note that by the Riesz-Thorin Theorem (see for example [12, Remark 2.2.5]) we can choose the constant  $C = C(q)$  in the previous lemma to stay bounded if  $0 < \varepsilon \leq q \leq 1/\varepsilon$  for some  $\varepsilon > 0$ . Therefore

$$C_0 = \sup_{\frac{1}{s} \leq \frac{1}{q} \leq \frac{1}{s} + \frac{1}{p}} C(q) < \infty.$$

The proof requires an iterative application of the previous lemma. Pick a natural number  $n$  such that  $np > C_0$  and define  $p' = np$ . Then if  $f_i \in H^{p'}$

$$T_{g,0}(f_1 f_2 \cdots f_n) = T_{T_{g,0}(f_2 f_3 \cdots f_n), 0}(f_1).$$

Keeping  $f_2, f_3, \dots, f_n$  fixed and applying the previous lemma to the operator  $T_{T_{g,0}(f_2 f_3 \cdots f_n), 0}$  we have that  $T_{g,0}(f_2 f_3 \cdots f_n) \in H^{q_1}$ ,  $\frac{1}{q_1} = \frac{1}{q} - \frac{1}{p'}$ . After  $n-2$  more applications of the Lemma 3.4 we arrive at  $T_{g,0} f_n \in H^{q_n}$ , with  $\frac{1}{q_n} = \frac{1}{q} - \frac{n-1}{p'}$ . We can still apply the lemma a last time since  $\frac{1}{s} \leq \frac{1}{q_{n-1}} \leq \frac{1}{s} + \frac{1}{p}$ . We thus get  $g \in H^{q_n}$ ,  $\frac{1}{q_n} = \frac{1}{q} - \frac{n}{p'} = \frac{1}{s}$ .

**3.2. Compactness.** Now, we are going to prove the compactness part of Theorem 1.1. The proof is similar to the one for boundedness, therefore, first we prove the result corresponding to Proposition 3.1 for the operators  $T_g^{n,k}$  when  $g$  is in the little Bloch space.

**Proposition 3.5.** *Let  $0 < p < \infty$  and  $g \in \mathcal{B}_0$ . Then for  $n > 1, 1 < k < n$  the operator  $T_g^{n,k}$  is a compact operator from  $H^p$  to itself. If  $g \in VMOA$ ,  $T_g^{n,0}$  is compact from  $H^p$  to itself.*

*Proof.* By integration by parts one can write  $T_g^{n,k}$  in the following form

$$T_g^{n,k} f = c_1 I^{(k)} M_{g^{(k)}} f + c_2 I^{(k+1)} M_{g^{(k+1)}} f + \cdots + c_{n-k} I^n M_{g^{(n)}} f.$$

Where  $M_\phi$  is the multiplication operator defined by

$$M_\phi f(z) = f(z)\phi(z).$$

If  $g$  is polynomial,  $M_{g^{(i)}}$ , for  $k \leq i \leq n$  is bounded on  $H^p$ . It is a well known fact (see [6]) that  $I$  is compact on  $H^p$ , hence if  $g$  is polynomial  $T_g^{n,k}$  is a compact operator. Suppose now that  $g \in \mathcal{B}_0$ . Then there exists a sequence of polynomials  $g_n$  converging in the Bloch norm to  $g$ . Therefore by an application of the open mapping theorem,

$$\|T_g^{n,k} - T_{g_n}^{n,k}\|_p \leq C_p \|g_n - g\|_{\mathcal{B}},$$

i.e.  $T_g^{n,k}$  is compact as the norm limit of compact operators.

The proof of the second part is identical, since  $VMOA$  is the closure of the set of polynomials in  $BMOA$ .  $\square$

By Proposition 3.5 and the fact that  $VMOA \subset \mathcal{B}_0$ , it follows immediately that  $T_{g,a}$  is compact if  $g \in VMOA$ .

To show necessity it suffices to show that  $g \in \mathcal{B}_0$ , because then by Proposition 3.5 and formula (5) it follows that  $T_g$  is compact and therefore  $g \in VMOA$ . Let  $f_{\lambda,\gamma}$  be defined by (6). For fixed  $\gamma$ ,  $f_{\lambda,\gamma}$  converges to zero on compact sets as  $|\lambda| \rightarrow 1^-$ . The compactness of  $T_{g,a}$  then implies that  $T_{g,a} f_{\lambda,\gamma}$  converges strongly to 0.

Using the same estimates as in the proof of necessity of Theorem 1.1, we get the following inequality.

$$\left| \sum_{k=0}^{n-1} g^{(n-k)}(\lambda) (1 - |\lambda|^2)^{(n-k)} a_k \bar{\lambda}^k (\gamma)_k \right| \leq \|T_{g,a} f_{\lambda,\gamma}\|_p.$$

Again, by applying Lemma 2.3 we can separate the above estimate.

$$|g^{(n)}(\lambda)| (1 - |\lambda|^2)^n \lesssim \sum_{i=1}^n c_i \|T_{g,a} f_{\lambda,\gamma_i}\|_p,$$

for some positive constants  $c_i$ . This last inequality gives the desired result, since the right part converges to zero as  $|\lambda| \rightarrow 1^-$ .

## REFERENCES

- [1] A. Aleman and A. Siskakis, “An integral operator on  $H^p$ ,” *Complex Var., Theory Appl.*, vol. 28, no. 2, pp. 149–158, 1995.
- [2] C. Pommerenke, “Schlichte funktionen und analytische funktionen von beschränkter mittlerer oszillation,” *Comment. Math. Helv.*, vol. 52, pp. 591–602, 1977.
- [3] E. Berkson and H. Porta, “Semigroups of analytic functions and composition operators,” *Michigan Math. J.*, vol. 25, no. 1, pp. 101–115, 1978.
- [4] A. Aleman, “A class of integral operators on spaces of analytic functions,” *Proceedings of the Winter School in Operator Theory and Complex Analysis, Univ. Malaga*, pp. 3–30, 2007. Málaga.
- [5] A. B. Alexandrov and V. V. Peller, “Hankel operators and similarity to a contraction,” *Internat. Math. Res. Notices*, no. 6, pp. 263–275, 1996.
- [6] A. Aleman and J. A. Cima, “An integral operator on  $H^p$  and Hardy’s inequality,” *J. Anal. Math.*, vol. 85, no. 1, pp. 157–176, 2001.
- [7] A. Aleman and J. A. Peláez, “Spectra of integration operators and weighted square functions,” *J. Anal. Math.*, vol. 61, no. 2, pp. 775–793, 2012.
- [8] J. Rättyä, “Linear differential equations with solutions in Hardy spaces,” *Complex Var. Elliptic Equ.*, vol. 52, no. 9, pp. 785–795, 2007.
- [9] W. S. Cohn and I. E. Verbitsky, “Factorization of tent spaces and Hankel operators,” *J. Funct. Anal.*, vol. 175, no. 2, pp. 308 – 329, 2000.
- [10] C. Fefferman and E. M. Stein, “ $H^p$  spaces of several variables,” *Acta Math.*, vol. 129, pp. 137–193, 1972.
- [11] P. L. Duren, *Theory of  $H^p$  spaces*. Courier Corporation, 2000.
- [12] K. Zhu, *Operator Theory in Function Spaces*. Mathematical surveys and monographs, M. Dekker, 1990.

N. CHALMOUKIS, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, 40127 BOLOGNA, ITALY  
 Email address: nikolaos.chalmoukis2@unibo.it