

# May Measurement Month 2019: results of blood pressure screening from 47 countries

Neil R. Poulter<sup>1\*</sup>, Claudio Borghi<sup>2</sup>, Albertino Damasceno<sup>3</sup>, Tazeen H. Jafar<sup>4,5</sup>, Nadia Khan<sup>6</sup>, Yoshihiro Kokubo<sup>7</sup>, Peter M. Nilsson<sup>8</sup>, Dorairaj Prabhakaran<sup>9</sup>, Markus P. Schlaich<sup>10</sup>, Aletta E. Schutte<sup>11</sup>, George S. Stergiou<sup>12</sup>, Thomas Unger<sup>13</sup>, and Thomas Beaney<sup>1,14</sup>

<sup>1</sup>Imperial Clinical Trials Unit, Imperial College London, 68 Wood Lane, London W12 7RH, UK

<sup>2</sup>Department of Medical and Surgical Sciences, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Via Massarenti 9, Bologna, Italy

<sup>3</sup>Faculty of Medicine, Eduardo Mondlane University, Maputo, Mozambique

<sup>4</sup>Health Services and Systems Research, DUKE-NUS Medical School, Singapore, Singapore

<sup>5</sup>Duke Global Health Institute, Duke University, Durham, NC, USA

<sup>6</sup>Department of Medicine, Center for Health Evaluation and Outcomes Sciences, University of British Columbia, Vancouver, Canada

<sup>7</sup>Department of Preventive Cardiology, National Cerebral and Cardiovascular Centre, Suita, Japan

<sup>8</sup>Department of Clinical Sciences, Lund University, Skane University Hospital, Malmo, Sweden

<sup>9</sup>Public Health Foundation of India, Plot 47, Sector 44, Haryana, India

<sup>10</sup>Dobney Hypertension Centre, School of Medicine, Royal Perth Hospital Unit, University of Western Australia, Level 3, MRF Building, 50 Rear Street, Perth WA 6000, Australia

<sup>11</sup>School of Population Health, University of New South Wales; The George Institute for Global Health, Sydney, NSW 2052, Australia

<sup>12</sup>Hypertension Center STRIDE-7, National and Kapodistrian University of Athens, School of Medicine, Third Department of Medicine, Sotiria Hospital, 152 Mesogion Avenue, Athens 11527, Greece

<sup>13</sup>CARIM School for Cardiovascular Diseases, Maastricht University P.O. Box 616,6200 Maastricht, The Netherlands and

<sup>14</sup>Department of Primary Care and Public Health, St Dunstan's Road, Imperial College London, London, W6 8RP, UK

## Background

There continues to be an inexorable rise in the death toll due to raised blood pressure (BP) which remains the biggest single contributor to global death and the global burden of disease.<sup>1</sup> It is estimated that in 2019 about 19% of all deaths (10.8 million) were due to raised BP, having risen from 9.4 million deaths in 2014.<sup>2</sup>

Hypertension causes over 50% of cases of heart disease, stroke, and heart failure<sup>3</sup> and it is estimated that about 10% of global healthcare spending arises from raised BP and its complications.<sup>4</sup> Moreover, hypertension-mediated organ damage increases risk of severe infections from COVID-19, including risk of death.<sup>5</sup> For any of these reasons, it is critical to prevent and, failing that, identify and manage raised

BP that appears to differentially affect the most vulnerable groups in society.

Given that BP is easy and inexpensive to measure and that several relatively inexpensive and effective drug classes are available to control hypertension, it is remarkable that a large proportion of people with hypertension remain unaware of their condition and that only a small proportion get their BPs controlled even to the now conservative target of <140/90 mmHg.<sup>6,7</sup>

This is the background that stimulated the launch of May Measurement Month (MMM) which began as an initiative of the International Society of Hypertension (ISH) in 2016.<sup>8</sup> The primary aim of MMM is to raise awareness of the importance of BP measurement at the individual and population level both nationally and globally. The first screening campaign took place in 2017 (MMM17), screening over 1.2 million adults worldwide,<sup>9</sup> followed by increasingly successful campaigns in 2018 (MMM18)<sup>10</sup> and 2019 (MMM19),<sup>11</sup>

\*Corresponding author. Tel: +44 (0)207 594 3445, Email: n.poulter@imperial.ac.uk

screening over 1.5 million each. Sadly, due to the COVID-19 pandemic, the 2020 campaign had to be deferred.

This supplement presents the data from the 47 individual countries each of which screened at least 2500 adults during MMM19 to complement the two previous supplements reporting national data from the highest screening countries in MMM17<sup>12</sup> and MMM18.<sup>13</sup>

## MMM19 summary

Details of the methods and results of MMM19 have been published previously.<sup>10</sup> In summary, the MMM19 campaign was a cross-sectional opportunistic survey of the BP levels of adults (aged  $\geq 18$  years) who volunteered to be screened. Screening sites were set up in a wide range of places from clinical settings such as hospitals and pharmacies to public spaces, such as supermarkets. Three sitting BP readings were obtained on each screenee using standardized methods and the mean of the last two was used in the analyses.

Those screenees found to have BP levels in the hypertensive range (systolic BP  $\geq 140$  mmHg and/or diastolic BP  $\geq 90$  mmHg) were given advice on diet and lifestyle to help reduce their BP and locally tailored advice to facilitate further follow-up of their raised BP. Prior to BP measurement, a brief questionnaire was administered collecting data on demographic, medical, social, and lifestyle variables.

Data were collected from 1 508 130 screenees from 92 countries in 2019.<sup>10</sup> Almost one third (32%) of screenees had never had their BP measured previously and 34% of all screenees were found to be hypertensive (systolic BP  $\geq 140$  mmHg and/or diastolic BP  $\geq 90$  mmHg, or taking antihypertensive medication). Of those with hypertension, 59% were aware of their condition and 55% were receiving antihypertensive treatment. Of those on treatment, 58% were controlled to  $<140/90$  mmHg and 29% to  $<130/80$  mmHg and only 50% were taking two or more antihypertensive agents. Overall, only 32% of hypertensive screenees were controlled to  $<140/90$  mmHg and 350 825 (23%) of all screenees had untreated or inadequately treated hypertension.

## From global to national data from MMM19

For many countries, the data collected during the MMM campaigns reflect the largest BP screening ever to take place in their country.<sup>8</sup> In order to make these data available and bring focus to the national level, we asked all 51 countries who had screened at least 2500 adults in MMM19 to produce their individual national report to collate in this *European Heart Journal Supplement* issue. Although to some extent arbitrary, the 2500 cut-off point was set to ensure a database of sufficient size to generate reasonably valid results from the analyses carried out.

A summary of the key results of the 47 countries who accepted the invitation to generate their national reports and of the other four countries is shown in *Table 1*.

The protocol for MMM19 was common to all participating countries and so the methods for each country are essentially the same. However, data from previously available

BP screening in each country vary as did the logistics and the sources of the convenience samples screened. These details potentially impact significantly on the interpretation of the results obtained in each country and the observed differences among them.

## Challenges for MMM19

Similar challenges to those which pertained in MMM18 persisted in 2019. Namely—acquiring ethical approval (where it was needed) and the distribution of BP measuring devices (once again kindly donated by OMRON Healthcare) caused delays and financial difficulties in some countries.

The quality of data collection in terms of missing data and the speed of transfer of the data to allow central analyses were both improved compared with 2018 which in turn improved on 2017.

Nevertheless, despite improvements to the bespoke MMM app, its use was limited to only 15.8% of participants, albeit increased from 12.4% in 2018. Consequently, central data cleaning remained a large, time-consuming task and we were only able to lock the database and initiate analyses in October 2019—4 months after screening for MMM19 had in theory ended.

## Methodological issues

The completeness of data collection varied across sites, with some sites systematically not collecting certain variables. Key demographic variables such as age (99.0%) and sex (99.6%) were well-documented, but others, particularly questions newly added in 2019, such as number of antihypertensive medications (86.0%) were less consistently recorded. Analyses of associations between BP and any variables for which recorded responses were limited in number at the national level were not carried out. Although the protocol advised three BP measurements in each participant, this was not possible in 25% of cases for various reasons, including logistical and time pressures, or participant preference. Our findings from MMM17, MMM18, and MMM19 showed significant declines in BP levels on average across the first to second, and to third readings, with the mean of the second and third resulting in the lowest proportion identified as hypertensive.<sup>9–11</sup> In order to provide comparative readings for those with only one or two readings, multiple imputation using chained equations was used, based on at least one BP measurement and other available participant characteristics, running on the global dataset as described previously.<sup>11</sup> Consideration was given to running imputations individually for each country; however, for many countries, the relatively small numbers involved, made an averaging over the global dataset more suitable. Furthermore, there was a priori no strong rationale for differential variation in subsequent BP measurements across countries.

Although efforts have been made to ensure consistency between individuals based on the number of BP readings, comparisons between countries should be interpreted cautiously. Participants were screened opportunistically, and samples at screening site and national level are not

**Table 1** Summary statistics for 51 countries with at least 2500 participants from May Measurement Month 2019

| Country                      | Total participants | Proportion of all participants with hypertension | Proportion of hypertensives aware | Proportion of hypertensives on medication | Proportion of those on medication with controlled BP | Proportion of all hypertensives controlled |
|------------------------------|--------------------|--|-----------------------------------|---|--|--|
| India                        | 362 708            | 29.4%  | 43.8%                             | 42.0%                                     | 55.5%  | 23.3%                                      |
| China                        | 238 387            | 27.8%  | 51.5%                             | 48.4%                                     | 60.2%  | 29.1%                                      |
| Argentina                    | 94 523             | 52.5%  | 81.1%                             | 77.7%                                     | 59.2%  | 46.0%                                      |
| Philippines                  | 89 941             | 53.3%  | 65.0%                             | 62.8%                                     | 61.1%  | 38.4%                                      |
| Nepal                        | 74 205             | 27.5%  | 46.3%                             | 37.5%                                     | 54.3%  | 20.3%                                      |
| Colombia                     | 48 324             | 27.9%  | 63.7%                             | 60.0%                                     | 64.0%  | 38.4%                                      |
| Mexico                       | 39 700             | 25.5%  | 43.8%                             | 41.7%                                     | 66.8%  | 27.8%                                      |
| Kenya                        | 33 992             | 26.1%  | 34.5%                             | 31.5%                                     | 59.7%  | 18.8%                                      |
| United Arab Emirates         | 32 152             | 23.9%  | 54.5%                             | 49.6%                                     | 59.7%  | 29.6%                                      |
| Cameroon                     | 30 187             | 20.8%  | 29.9%                             | 24.0%                                     | 46.7%  | 11.2%                                      |
| Democratic Republic of Congo | 29 857             | 25.5%  | 33.1%                             | 23.2%                                     | 51.5%  | 11.9%                                      |
| Vietnam                      | 25 887             | 33.8%  | 69.8%                             | 65.5%                                     | 51.2%  | 33.5%                                      |
| Saudi Arabia                 | 25 023             | 29.2%  | 60.8%                             | 60.8%                                     | 64.6%  | 39.3%                                      |
| Bangladesh                   | 24 941             | 28.0%  | 76.3%                             | 71.6%                                     | 64.2%  | 46.0%                                      |
| Taiwan                       | 24 851             | 49.7%  | 84.7%                             | 82.1%                                     | 72.0%  | 59.2%                                      |
| Venezuela                    | 24 672             | 48.9%  | 86.2%                             | 82.6%                                     | 64.5%  | 53.3%                                      |
| Albania                      | 19 154             | 38.6%  | 64.7%                             | 62.0%                                     | 48.3%  | 29.9%                                      |
| Cabo Verde                   | 17 627             | 34.0%  | 68.4%                             | 51.3%                                     | 46.5%  | 23.8%                                      |
| Ecuador                      | 15 885             | 41.9%  | 65.0%                             | 64.2%                                     | 76.5%  | 49.1%                                      |
| Brazil                       | 13 476             | 50.9%  | 68.8%                             | 65.3%                                     | 55.2%  | 36.1%                                      |
| Georgia                      | 13 267             | 64.1%  | 85.4%                             | 85.0%                                     | 34.8%  | 29.6%                                      |
| Tunisia                      | 11 271             | 38.1%  | 72.5%                             | 67.5%                                     | 56.6%  | 38.2%                                      |
| Italy                        | 10 182             | 31.1%  | 62.1%                             | 16.2%                                     | 45.8%  | 7.4%                                       |
| Republic of Korea            | 9975               | 47.6%  | 76.2%                             | 74.0%                                     | 68.2%  | 50.5%                                      |
| Armenia                      | 9818               | 41.6%  | 72.8%                             | 65.4%                                     | 46.5%  | 30.4%                                      |
| Malawi                       | 9723               | 26.3%  | 17.4%                             | 15.2%                                     | 51.9%  | 7.9%                                       |
| United Kingdom and Ireland   | 9233               | 33.4%  | 33.5%                             | 29.7%                                     | 38.2%  | 11.4%                                      |
| Zambia                       | 9232               | 30.7%  | 42.6%                             | 27.6%                                     | 35.0%  | 9.7%                                       |
| Libya                        | 8686               | 42.8%  | 64.8%                             | 59.2%                                     | 48.2%  | 28.5%                                      |
| Mauritius                    | 8262               | 29.4%  | 64.7%                             | 60.8%                                     | 57.3%  | 34.8%                                      |
| Angola                       | 7112               | 38.6%  | 59.8%                             | 50.6%                                     | 41.2%  | 20.8%                                      |
| Ghana                        | 7102               | 25.9%  | 36.5%                             | 30.0%                                     | 46.1%  | 13.8%                                      |
| Poland                       | 7072               | 55.4%  | 83.0%                             | 80.4%                                     | 58.2%  | 46.7%                                      |
| Lebanon                      | 7019               | 36.6%  | 64.1%                             | 62.3%                                     | 62.6%  | 39.0%                                      |
| Pakistan                     | 6919               | 52.1%  | 56.2%                             | 49.5%                                     | 40.0%  | 19.8%                                      |
| Chile                        | 6876               | 35.4%  | 65.9%                             | 60.1%                                     | 57.2%  | 34.4%                                      |
| Mongolia                     | 6522               | 32.5%  | 62.2%                             | 50.1%                                     | 54.1%  | 27.1%                                      |
| Greece                       | 5814               | 41.6%  | 78.7%                             | 73.1%                                     | 66.1%  | 48.3%                                      |
| Botswana                     | 5459               | 32.1%  | 44.8%                             | 41.5%                                     | 47.0%  | 19.5%                                      |
| Russia                       | 5447               | 30.6%  | 72.8%                             | 63.0%                                     | 46.8%  | 29.5%                                      |
| Slovenia                     | 4974               | 61.1%  | 76.4%                             | 68.0%                                     | 46.2%  | 31.4%                                      |
| South Africa                 | 4727               | 31.9%  | 42.5%                             | 36.1%                                     | 51.5%  | 18.6%                                      |
| Spain                        | 4433               | 42.5%  | 77.2%                             | 71.1%                                     | 64.9%  | 46.1%                                      |
| Paraguay                     | 4301               | 41.4%  | 70.0%                             | 65.3%                                     | 44.6%  | 29.1%                                      |
| Nigeria                      | 3646               | 39.2%  | 62.9%                             | 55.4%                                     | 46.8%  | 25.9%                                      |
| Benin                        | 3637               | 37.5%  | 64.5%                             | 43.9%                                     | 34.9%  | 15.3%                                      |
| Republic of the Congo        | 3157               | 33.5%  | 42.6%                             | 37.3%                                     | 62.4%  | 23.3%                                      |
| Malaysia                     | 3062               | 18.7%  | 63.2%                             | 57.2%                                     | 70.3%  | 40.2%                                      |
| Australia                    | 2877               | 31.3%  | 50.5%                             | 40.6%                                     | 54.3%  | 22.0%                                      |
| Hungary                      | 2766               | 46.5%  | 74.4%                             | 69.8%                                     | 53.2%  | 37.1%                                      |
| Jamaica                      | 2550               | 41.4%  | 69.9%                             | 62.5%                                     | 44.4%  | 27.8%                                      |

expected to be representative at the population level. An understanding of the local context of screening is required, in terms of target populations and sites of screening, which the national papers in this supplement aim to provide.

## Limitations of MMM19

Although MMM19 was just larger in terms of the number of countries included (92) and the total number of screenees (1 508 130) a few countries including Sudan and Cote D'Ivoire that had made large contributions in previous years, were unable to take part due to civil upheavals or financial constraints.

Once again, we were limited in the scope of data collection by the very reasonable request of local investigators to restrict the time and complexity of the interaction spent with each screenee. Consequently, our database is limited in terms of the number of variables evaluated and blood, urine and more complex anthropometry are beyond the capacity of the personnel and available budget.

By design, as discussed, MMM does not attempt to collect nationally representative samples from each country in which it takes place. Nevertheless, analyses in progress show significant associations between national stroke mortality and various measures of national BP management and control seen in MMM.<sup>14</sup>

Although effective intervention on raised BP was not the primary aim of MMM, the detection in the three campaigns to date, of almost one million adults with either untreated or inadequately treated hypertension begs the question of whether MMM improves the health outcomes of those detected. Due to the cross-sectional design, collection of follow-up data was beyond the scope of the campaign and we do not know whether participants instigated lifestyle changes, modified health-seeking behaviours, or were started on or had any anti-hypertensive medication increased. However, a study of a cohort of older adults in China reported that an approach very similar to that used in MMM to detect raised BP was associated with a significant and important reduction in systolic BP two years after screening.<sup>15</sup> Nevertheless, an MMM pilot study is in discussion to evaluate the efficacy of a collaboration with the RESOLVE to Save Lives programme<sup>16</sup> to facilitate more direct links with effective drug treatment and thereby to ensure improved BP levels.

## Strengths of and prospects for the MMM campaign

In MMM17, 34 of the 80 countries included, reported that MMM was the largest BP screening to have taken place in their country.<sup>9</sup> Since then, coverage has increased to 92 countries and the number of screenees included has risen from over 1.2 million in MMM17 to over 1.5 million in MMM19. It is not easy to produce hard data to confirm whether the primary aim of MMM—to raise awareness of BP measurement—has been achieved. However, significant numbers of participants (almost half a million in 2019 alone) had never had a BP measured previously, while across all three campaigns almost one million adults have

been found with untreated or inadequately treated hypertension. This suggests that at the individual level, at least in these groups, awareness has been improved. We are further persuaded by the extent of television, radio, media, and social media coverage around the world that, at the population level, awareness has also been raised, although this is difficult to quantify.

Due to the COVID-19 pandemic, MMM20 had to be deferred. Despite persisting uncertainties surrounding this pandemic, we expect that MMM21 will take place, but it seems that 'May' is unlikely to be the central month of the campaign. Indeed, the window for data collection will be extended to anytime between May and November 2021, depending on local pandemic conditions and the critical acquisition of local ethical clearance.

Because all the MMM investigators and volunteers measuring BPs around the world do so pro bono and the campaign receives many charitable donations centrally, and at the national level, the MMM campaign remains an inexpensive but very effective means of detecting large numbers who require additional help with their raised BP—the most putative of cardiovascular risk factors. In addition to demonstrating associations between measures of BP detection, management and control at a national level and national stroke mortality, analyses are in progress to evaluate the potential utility of MMM-based data for characterizing nationally representative BP data.

Together these two sets of analyses will allow MMM to address one of its four key objectives—to use MMM data to motivate governments to improve BP screening facilities and management. Meanwhile, annual MMM surveys will continue to provide an inexpensive and hopefully temporary substitute for systematic screening in many countries worldwide.

## Acknowledgements

Our sincere thanks to Judith Bunn (MMM Project Manager 2017-2019) for administering MMM19 and to Harsha McArdle (MMM Project Manager 2020-current) and Ranjit Rayat (Editing Assistant) for their superb dedicated efforts towards this national supplement project. We would also like to thank the officers of the International Society of Hypertension for their background support, OMRON Healthcare and Servier Pharmaceuticals Co for their continued generous contributions. Most importantly our thanks go out to the thousands of volunteers for their support in making the MMM campaign a huge success.

## Data availability

Data are not publicly available but are available with permission from the MMM Management Board, on request through the MMM website: [maymeasure.org](http://maymeasure.org).

## Funding

Most of the financial burden of MMM within each country falls on local resources and as ever we are indebted to those who have provided the necessary financial and

logistical support within each country and to all those who have volunteered their time to the project. We shall continue to seek further sources of funding to supplement the generous annual donation made by Servier Pharmaceuticals Co to at least offset a greater proportion of the local costs of running MMM. We must also thank OMRON Healthcare for their continued support with the supply of BP measuring devices.

**Conflict of interest:** N.R.P. was supported by the National Institute for Health Research Senior Investigator Awards, Biomedical Research Centre funding, and the British Heart Foundation Research Centre Excellence Award. He has received financial support from several pharmaceutical companies which manufacture BP-lowering agents, for consultancy fees (Servier), research projects and staff (Servier, Pfizer) and for arranging and speaking at educational meetings (AstraZeneca, Lri Therapharma, Napi, Servier, Sanofi, Eva Pharma and Pfizer). He holds no stocks and shares in any such companies. C.B., A.D., N.K., Y.K., P.M.N., D.P., T.U., and T.B. report no conflicts of interest. T.H.J. is supported by the National Medical Research Council, Singapore and has received consulting fees from RESOLVE to Save Lives. M.P.S. is supported by an NHMRC Research Fellowship and has received consulting fees and/or travel and research support from Medtronic, Abott, Novartis, Servier, Pfizer and Boehringer-Ingelheim. A.E.S. has received lecture and/or travel and research support from Abott, Novartis, Servier, OMRON Healthcare and Takeda. G.S.S. has received lecture and consultancy fees from OMRON Healthcare.

## References

- GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018; **392**:1923.
- GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; **396**: 1223-1249.
- Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L, Alexander L, Estep K, Hassen Abate K, Akinyemiju TF, Ali R, Alvis-Guzman N, Azzopardi P, Banerjee A, Barnighausen T, Basu A, Bekele T, Bennett DA, Biadgilign S, Catalá-López F, Feigin VL, Fernandes JC, Fischer F, Gebru AA, Gona P, Gupta R, Hankey GJ, Jonas JB, Judd SE, Khang Y-H, Khosravi A, Kim YJ, Kimokoti RW, Kokubo Y, Kolte D, Lopez A, Lotufo PA, Malekzadeh R, Melaku YA, Mensah GA, Misganaw A, Mokdad AH, Moran AE, Nawaz H, Neal B, Ngalesoni FN, Ohkubo T, Pourmalek F, Rafay A, Rai RK, Rojas-Rueda D, Sampson UK, Santos IS, Sawhney M, Schutte AE, Sepanlou SG, Shifa GT, Shiuie I, Tedla BA, Thrift AG, Tonelli M, Truelsen T, Tsilimparis N, Ukwaja KN, Uthman OA, Vasankari T, Venketasubramanian N, Vlassov VV, Vos T, Westerman R, Yan LL, Yano Y, Yonemoto N, Zaki MES, Murray CJL. Global burden of hypertension and systolic blood pressure of at least 110 to 115 mm Hg, 1990-2015. *JAMA* 2017; **317**:165-182.
- Gaziano TA, Bitton A, Anand S, Weinstein MC. The global cost of non-optimal blood pressure. *J Hypertens* 2009; **27**:1472-1477.
- Yang J, Zheng Y, Gou X, Pu K, Chen Z, Guo Q, Ji R, Wang H, Wang Y, Zhou Y. Prevalence of comorbidities and its effect in patients with CARS-CoV-2: a systematic review and meta-analysis. *Int J Infect Dis* 2020; **94**:91-95.
- NCD Risk Factor Collaboration (NCD-RisC). Long-term and recent trends in hypertension awareness, treatment, and control in 12 high-income countries: an analysis of 123 nationally representative surveys. *Lancet* 2019; **394**:639-651.
- Geldsetzer P, Manne-Goehler J, Marcus M-E, Ebert C, Zhumadilov Z, Wesseh CS, Tsabedze L, Supiyev A, Sturua L, Bahendeka SK, Sibai AM, Quesnel-Crooks S, Norov B, Mwangi KJ, Mwalim O, Wong-McClure R, Mayige MT, Martins JS, Lunet N, Labadarios D, Karki KB, Kagaruki GB, Jorgensen JMA, Hwalla NC, Houinato D, Houehanou C, Msaidié M, Guwatudde D, Gurung MS, Gathecha G, Dorobantu M, Damasceno A, Bovet P, Bicaba BW, Aryal KK, Andall-Brereton G, Agoudavi K, Stokes A, Davies JI, Barnighausen T, Atun R, Vollmer S, Jaacks LM. The state of hypertension care in 44 low-income and middle-income countries: a cross-sectional study of nationally representative individual-level data from 1.1 million adults. *Lancet* 2019; **394**:652-662.
- Poulter NR, Lackland DT. May Measurement Month: a global blood pressure screening campaign. *Lancet* 2017; **389**:1678-1680.
- Beaney T, Schutte AE, Tomaszewski M, Ariti C, Burrell LM, Castillo RR, Charchar FJ, Damasceno A, Kruger R, Lackland DT, Nilsson PM, Prabhakaran D, Ramirez AJ, Schlaich MP, Wang J, Weber MA, Poulter NR; MMM Investigators. May Measurement Month 2017: an analysis of blood pressure screening results worldwide. *Lancet Glob Health* 2018; **6**:e736-e743.
- Beaney T, Burrell LM, Castillo RR, Charchar FJ, Cro S, Damasceno A, Kruger R, Nilsson PM, Prabhakaran D, Ramirez AJ, Schlaich MP, Schutte AE, Tomaszewski M, Touyz R, Wang JG, Weber MA, Poulter NR; MMM Investigators. May Measurement Month 2018: a pragmatic global screening campaign to raise awareness of blood pressure by the International Society of Hypertension. *Eur Heart J* 2019; **40**: 2006-2017.
- Beaney T, Schutte AE, Stergiou GS, Borghi C, Burger D, Charchar F, Cro S, Diaz A, Damasceno A, Espeche W, Jose AP, Khan N, Kokubo Y, Maheshwari A, Marin MJ, More A, Neupane D, Nilsson P, Patil M, Prabhakaran D, Ramirez A, Rodriguez P, Schlaich M, Steckelings UM, Tomaszewski M, Unger T, Wainford R, Wang JG, Williams B, Poulter NR; on behalf of MMM Investigators. May Measurement Month 2019: the Global Blood Pressure Screening Campaign of the International Society of Hypertension. *Hypertension* 2020; **76**:333-341.
- Poulter NR, Borghi C, Castillo RR, Charchar FJ, Ramirez AJ, Schlaich MP, Beaney T. May Measurement Month 2017: results of 39 national blood pressure screening programmes. *Eur Heart J Suppl* 2019; **21**(Suppl D):D1-D132.
- Poulter NR, Borghi C, Burger D, Castillo RR, Damasceno A, Ito S, Jose AP, Kruger R, Morgan T, Nilsson PM, Schlaich MP, Schutte AE, Stergiou G, Unger T, Wainford RD, Beaney T. May Measurement Month 2018: results of blood pressure screening from 41 countries. *Eur Heart J Suppl* 2020; **22**(Suppl H):H1-H141.
- Lin, Q Ye, T Beaney, T Poulter, NR. The associations between hypertension and stroke from May Measurement Month and between national hypertension parameters with stroke mortality from the Global Burden of Disease. ESH/ISH 2021 Conference Abstract.
- Chen S, Sudharsanan N, Huang F, Liu Y, Geldsetzer P, Barnighausen T. Impact of community based screening for hypertension on blood pressure after two years: regression discontinuity analysis in a national cohort of older adults in China. *BMJ* 2019; **366**:14064.
- Frieden TR, Varghese CV, Kishore SP, Campbell NRC, Moran AE, Padwal R, Jaffe MG. Scaling up effective treatment of hypertension—a pathfinder for universal health coverage. *J Clin Hypertens* 2019; **21**:1442-1449.